The effect of inter-group competition on intra-group affiliation in primates

B. Majolo^{1*}, A. de Bortoli Vizioli^{2,3}, J. Lehmann⁴

¹School of Psychology, University of Lincoln, Lincoln, U.K. Email: bmajolo@lincoln.ac.uk

²Istituto di Scienze e Tecnologie della Cognizione, Consiglio Nazionale delle Ricerche,
Rome, Italy.

³Dipartimento di Psicologia dello Sviluppo e della Socializzazione, Università di Padova,
Padua, Italy. Email: aurora.debortolivizioli@gmail.com

⁴Centre for Research in Evolutionary Anthropology, University of Roehampton, London,
U.K. Email: J.Lehmann@roehampton.ac.uk

*Corresponding author:

Bonaventura Majolo, School of Psychology, University of Lincoln, Brayford Pool, Lincoln,
LN6 7TS, U.K. Phone: +44-01522-837015, fax: +44-01522-886026, email:
bmajolo@lincoln.ac.uk

Word count: 5,514

The effect of inter-group competition on intra-group affiliation in primates

Word count: 5,975

Researchers from various disciplines have hypothesised a positive correlation between the level of inter-group contest competition (IGCC) and the evolution of behavioural traits, such as co-operation, altruism and friendship, which promote intra-group affiliation. Empirical support for this hypothesis is, however, scarce and mainly available from humans. We tested if the level of IGCC affects intra-group affiliation (i.e. intra-group grooming exchange) among male and female non-human primates. To quantify intra-group affiliation, we used social network measures and a grooming index. Our measure of IGCC combined frequency of inter-group encounters and proportion of aggressive encounters and was calculated separately for males and females. We ran our analyses on 27 wild groups of primates belonging to 15 species (13 Cercopithecinae, 1 Colobinae and 1 Cebinae). Our analyses reveal a clear pattern of correlated evolution between grooming network density and inter-individual variation in the number of grooming partners on the one hand, and the intensity of IGCC on the other in females, but not males. Thus, our data suggest that the exact nature of the relationship between IGCC and intra-group affiliation is sex-specific. These results may be explained by the differential costs and benefits males and females experience during aggressive inter-group confrontations and by sex-specific differences in intra-group affiliation.

Keywords: affiliation, collective action problem, competition, grooming, monkeys

Inter-group contest competition (IGCC), where some/all members of one group aggressively attempt to access and monopolise valuable resources (e.g. food or water) at the expense of another group, is a phenomenon observed in many group-living species (e.g. Rubenstein 1986; Kitchen and Beehner 2007; Radford 2008a; Crofoot and Wrangham 2010). The outcome of an aggressive inter-group encounter can affect individual fitness (e.g. Radford and Fawcett, 2014), especially when ecological conditions elicit frequent aggressive interactions between groups (e.g. due to habitat saturation or food scarcity) that bear high cost/benefits payoffs (e.g. Mitani et al. 2010). Aggressive inter-group encounters represent a typical example of collective action problems (Nunn and Lewis 2001; Willems et al. 2013; Willems and van Schaik, 2015): groups where a larger proportion of animals take part in the aggressive confrontation are expected to have the best chances to win a fight (other things being equal; e.g. group size, sex ratio) and thus gain access to the resources at stake. However, because aggressive inter-group encounters also bear potential high costs for individuals (e.g. injuries or death; Mitani et al. 2010), the risk of free-riding is high: some individuals could gain the benefits of their group winning a fight with another group, without incurring any potential cost, by not taking an active role in the aggressive confrontation (Nunn and Lewis 2001; Crofoot and Gilby 2012; Willems et al. 2013; Willems and van Schaik, 2015).

Researchers have hypothesised that strong intra-group affiliation should reduce the probability that animals in a group free-ride during collective action problems (Alexander and Borgia 1978; Choi and Bowles 2007; Puurtinen and Mappes 2009). Thus, individuals from groups where intra-group affiliation is strong should out-perform groups with weak intra-group affiliation during inter-group confrontations, because everybody is taking part in the interaction. These groups should then gain overall fitness benefits. Under this scenario, individuals in populations/species that face a high level of IGCC should become

progressively more affiliative towards their group members over evolutionary time, thereby increasing the overall intra-group affiliation of the group. Furthermore, individuals in populations/species where intra-group affiliation is strong should also be more likely to take part in collective actions (such as protecting a food source from other groups) than individuals in populations/species where intra-group affiliation is weak (Conradt and List 2009; Miller et al. 2013). Following this logic, researchers from a range of different disciplines, including evolutionary biology (Reeve and Hölldobler 2007), behavioural ecology (Wrangham 1980; Sterck et al. 1997), psychology (West et al. 2006; van Vugt and Park 2009) and anthropology (Haas 1990; Choi and Bowles 2007), have predicted that a high level of IGCC should favour the evolution of behavioural traits which promote intra-group affiliation, such as social bonding and parochialism (Alexander and Borgia 1978; Reeve and Hölldobler 2007). However, different mechanisms and factors have been proposed to be involved, depending on the discipline and/or study taxon. Genetic similarity among group members (e.g. due to limited dispersal; Reeve and Hölldobler 2007), environmental conditions (Sterck et al. 1997), and the evolution of social norms (e.g. rules on fairness and punishment of free-riders; e.g. Gintis et al. 2015) can all affect the relationship between inter-group competition and intra-group affiliation over evolutionary time.

Several mathematical models, that support the hypothesis that the level of IGCC has a positive effect on the evolution of intra-group affiliation, have been developed (e.g. Reeve and Hölldobler 2007; Bowles 2009). However, empirical tests of this hypothesis are sparse and have mostly focused on the proximate link between IGCC and intra-group affiliation. There is consistent experimental evidence in humans showing that co-operation increases when groups are competing with one another compared to situations without inter-group competition; this effect is independent from kin relationships amongst group members (Erev et al. 1993; West et al. 2006; Tan and Bolle 2007; Puurtinen and Mappes 2009). This

evidence has led some authors to propose that the extensive period of warfare that our ancestors faced in the Pleistocene and Holocene has driven the evolution of group-beneficial behaviours, such as in-/out-group categorisation in humans (Choi and Bowles 2007; Bowles 2009).

For non-human species, data on the proximate link between IGCC and intra-group affiliation are relatively scarce (for a review: Grueter 2013), although the same principles should apply to all taxonomic groups (see e.g. Wrangham 1980; Reeve and Hölldobler 2007). A few studies have focused on post inter-group encounter allo-preening and allo-grooming, as a measure of intra-group affiliation (Radford 2008a), as these two behaviours have important social functions in many mammals and birds (e.g. they promote agonistic support and food tolerance; Dunbar 1991; Ventura et al. 2006; Radford 2008a). In the co-operatively breeding green woodhoopoe (*Phoeniculus purpureus*) the frequency and duration of allo-preening between group members increases following a conflict with another group (Radford 2008a, b, 2011). Conversely, in captive tufted capuchin monkeys (*Cebus apella*) visual interactions between groups did not result in increased grooming exchange within the group (Polizzi di Sorrentino et al. 2012). Another measure of affiliation that has been used is spatial proximity between group members. In chimpanzees (*Pan troglodytes*) and spider monkeys (*Ateles geoffroyi*) individuals were found to stay closer together when feeding in the home range of a neighbouring group (i.e. an area where the risk of a conflict with another group is high) than when feeding in their home range (Aureli et al. 2006; Mitani et al. 2010). However, all of these studies have tested the immediate effects of inter-group encounters on intra-group affiliation. It remains unclear if such effects are only short-lived and thus temporary (e.g. an emotional response to the stress of the encounter; Radford 2008a, 2011; Polizzi di Sorrentino 2012), if such an increase in affiliative behaviour occurs primarily between already closely bonded partners and/or those who have actively participated to the

inter-group encounter (i.e. affiliation is partner specific), or if such behavioural responses to inter-group encounters would indeed lead to an overall more general increase in intra-group affiliation over evolutionary time.

To our knowledge, only two studies (Cheney 1992; Grueter 2013) have attempted to test the generality of the hypothesised relationship between IGCC and intra-group affiliation across a number of different species. Cheney (1992) used allo-grooming (hereafter grooming) distribution (averaged across individuals of the same group) as a measure of group affiliation in non-human primates. She predicted that animals would increase their effort to distribute their grooming more evenly to their potential grooming partners, instead of mostly grooming with fewer individuals, the higher the level of IGCC they faced. Grueter (2013) also investigated non-human primates and used the average grooming time per species as a measure of intra-group affiliation. These two studies hypothesised that the proximate effect of IGCC on intra-group affiliation would extend beyond the first few minutes after an inter-group encounter. Groups facing higher IGCC would have overall stronger intra-group affiliation than groups facing a lower level of IGCC, irrespective of the timing and occurrence of each inter-group encounter (Wrangham 1980; Sterck et al. 1997). Contrary to what was predicted, IGCC was not found to have a significant effect on either grooming distribution among group members (Cheney 1992) or on total grooming time (Grueter 2013). However, these two studies could not effectively tackle some important issues. Cheney's (1992) study was restricted to female primates and did not control for the phylogenetic relationship between species. Grueter (2013) used total grooming time to measure group cohesion, without taking into account how animals distributed their grooming among their group members. Moreover, Grueter (2013) used a proxy of IGCC (i.e. home range overlap) that could equally measure inter-group scramble competition (Wrangham 1980; Sterck et al. 1997). These factors could potentially explain the discrepancy between the human and the

non-human primate data.

The mixed results available on this topic so far make it difficult to determine if IGCC affects the evolution of group-beneficial behaviours (Choi and Bowles 2007; Bowles 2009) in non-human animals. Our aim was to analyse the occurrence of correlated evolution between IGCC and intra-group affiliation across primate social groups, using a phylogenetically controlled analysis and a variety of different measures of intra-group affiliation. We measured the level of IGCC by multiplying the frequency of inter-group encounters with the proportion of aggressive encounters (independently calculated for males and females). Intra-group affiliation, using grooming distribution, was measured by three different variables: (a) the evenness of grooming, assessing the distribution of grooming effort across potential grooming partners (following Cheney 1992), (b) the inter-individual variation in the number of grooming partners, and (c) grooming network density. We predicted that species that experience more IGCC, will also exhibit more intra-group affiliation, that is, animals would increase their effort to distribute their grooming more evenly to their group companions, inter-individual variation in grooming partner numbers would be low, and more individuals would be involved in grooming interactions (high network density). Due to the scarcity of data available on a wide range of primate taxa our dataset is mostly composed of Cercopithecinae (See supplementary table 1), i.e. of species where grooming is the main affiliative behaviour and where females are the philopatric sex. Although not ideal, this allowed us to some extent to control for species-specific differences in affiliative behaviours. In addition, we ran the analyses separately for male and female primates, because intra-group same-sex affiliation can differ strongly between non-human primate species (Wrangham 1980; Sterck et al. 1997; Crofoot and Wrangham 2010). Moreover, the two sexes differ in their aggressive participation in inter-group encounters (Majolo et al. 2005; Kitchen and Beehner 2007; Crofoot and Wrangham 2010) and might have markedly different cost-benefit

ratios of IGCC. We did not consider grooming interactions between males and females, as a different set of hypotheses (e.g. hired-gun hypothesis; Rubenstein 1986) than the ones tested here focus on hetero-sexual grooming.

METHODS

Data collection

We used the ISI Web of Knowledge (© Thomson Reuters) to review the primatological literature published between 1st January 1950 and 1st June 2011. We selected papers that contained detailed data on grooming interactions between adult and sub-adult group members (e.g. a matrix of grooming exchange) and on IGCC on one or more social groups. Data on juveniles and/or infants were not included in the analyses. When these data were not available in a single paper, we combined different published studies containing the relevant data collected on the same social group within the same overall time period (see Supplementary material for further details).

Our literature review showed that published studies providing detailed data on grooming and IGCC on a single social group are scarce. Therefore, we contacted colleagues working on primates in the field, asking if they were willing to share with us the following data for each of their study groups: study species, dates when data were collected, duration of the study (in months) and location, group size (i.e. number of adults, sub-adults, juveniles and infants in the group), whether the group was provisioned (i.e. the group relied extensively on human-provisioned food) or not, and frequency of intra-group contest food competition (i.e. number of aggressive interactions over food / hour). We also requested a matrix containing the proportion or frequency of grooming interactions among the adult and sub-adult males and/or females of the study group. Finally, we requested the following data on IGCC: frequency of inter-group encounters (i.e. number of inter-group encounters / day) for the

study group and proportion of inter-group encounters in which at least one adult male and/or female from the study group was aggressive towards the other group (i.e. N aggressive encounters for males/females divided by total encounters observed). We asked colleagues to give us data for this latter variable separately for males and females. Moreover, we asked colleagues to provide us with their definition of inter-group encounter (or we extracted this information from the selected literature). The minimum distance between two groups, for an inter-group encounter to occur, ranged between 15 and 100 meters across the studies included in our dataset, with an average distance of approximately 50 metres. Aggression towards another group was defined as an animal displaying threatening facial expressions or giving a threat call towards, chasing or biting one or more animals from the other group.

Data analysis

The analyses presented here were based on 27 study groups from 15 species, 13 Cercopithecinae, 1 Colobinae and 1 Cebinae (males: 11 species, 19 study groups; females: 15 species, 25 study groups; electronic supplementary table S1). We quantified the level of IGCC, our independent variable, by multiplying the frequency of inter-group encounters (N of encounters/day) by the proportion of these encounters that were aggressive (Willems & van Schaik 2015). The level of IGCC had different values for analyses done on males and females, as we calculated the proportion of aggressive encounters separately for males and females (based on their involvement in them).

Matrices of grooming exchanges between group members were used to obtain three measures of intra-group affiliation (our dependent variables). First, following Cheney (1992), we used the Shannon-Wiener index (SWI; Wilson and Bossert 1971) to measure how an individual distributes grooming across all the potential grooming partners (Supplementary material for further details). The higher the value of SWI the more animals have evenly

distributed grooming relationships (i.e. they attempt to groom all of the available partners about equally). In addition to the SWI, we calculated two commonly used social network metrics to quantify intra-group affiliation (Borgatti et al. 2006; Wey et al. 2008). Because of the differences between studies in the measures of grooming (e.g. percentage of grooming duration or simply frequency) we only calculated network measures for binary networks (i.e. the presence/absence of a grooming relationship) and not on bond strength. Our two network measures of intra-group affiliation were standard deviation of the out degree and density (Borgatti et al. 2006). Standard deviation of the out degree (SDOD) measures the inter-individual variation in the number of partners groomed. A low SDOD value indicates that all group members groom a similar number of partners while a high SDOD indicates that the number of grooming partners is skewed between group members. Density measures the number of existing grooming relationships in relation to all possible grooming relationships. All network metrics were calculated using UCInet 6 (Borgatti et al. 2006). The three dependent variables were not significantly correlated to one another (Spearman pair-wise correlations; all $P > 0.28$), suggesting that our indices of intra-group affiliation measured different aspects of grooming distribution.

We controlled for observation effort (i.e. study period length, measured in months) and for four biologically relevant variables, which may affect the level of IGCC and/or overall intra-group affiliation (electronic supplementary methods for further details): provisioning (binary variable: yes/no), group size (number of animals in the social group, including juveniles and infants), intra-group contest competition (aggressive events/hour) and predation risk (categorical variable: low, medium or high risk). We also controlled for additional variables that could bias our results, including a ‘data quality’ variable (see electronic supplementary methods for details).

Statistical analyses

We ran phylogenetically controlled generalized least square regression models (PGLS) in R 2.14.0 (R Development Core Team 2011), using the CAPER 0.4 package (Orme et al. 2012). The phylogenetic tree for the analyses was obtained from the 10ktree primate phylogeny version 2 (Arnold et al. 2010). We derived lambda via maximum likelihood estimation as provided in CAPER. All the analyses were run considering data on each social group as a single data point, as IGCC and intra-group affiliation are not likely to solely be species-specific traits and can vary significantly across populations/groups (for a similar approach see Majolo et al. 2012). To be able to control for phylogeny we added these populations into the phylogenetic tree (via Mesquite) using very small branch lengths of 0.000001 my (Majolo et al. 2012). This value was chosen to reflect the fact that these are in fact not different species. Because this is an arbitrary value which might have affected our results, we re-ran the analyses using, respectively, 0.00001 my and 0.0001 my as branch lengths for populations of the same species. These two additional sets of analyses (not shown here for the sake of brevity) yielded the same results as the ones obtained using 0.000001 my as a branch length, suggesting that the chosen branch length did not affect the results.

Continuous variables were tested for normality using Kolmogorov-Smirnov test. Three variables (level of IGCC, study period, and intra-group contest competition) were found to significantly deviate from normality. Level of IGCC and intra-group contest competition were ln-transformed; study period was square root-transformed. These transformations solved the normality problem.

Altogether we ran three models for each sex, with each of these models containing our independent variable (level of IGCC) and the four additional control variables, namely study period length, group size and intra-group contest competition as covariates, and provisioning (2 levels) and predation risk (3 levels) as confounding factors, both of which

were dummy-coded. In addition, we ran the same analyses separately on the two variables that we used to construct our measure of IGCC (i.e. the frequency of inter-group encounters per day and the proportion of aggressive inter-group encounters for males/females). The results for these models were representative of our overall findings. Therefore, we discuss briefly these results below but we present these analyses in the supplementary material (electronic supplementary tables S2-S5).

RESULTS

Females

A phylogenetic signal was detected for the standard deviation of the out degree (SDOD; $\lambda = 0.90$) and for density ($\lambda = 0.76$), but not for the Shannon-Wiener index (SWI; Table 1). The level of IGCC was not significantly related to SWI (Table 1). Conversely, and supporting our predictions, females showed significantly less variation in their number of grooming partners (i.e. lower SDOD; Figure 1) and had significantly denser grooming networks (Figure 2) when levels of IGCC were higher (Table 2). Among the control variables, we found that group size significantly affected each of the three dependent variables: larger groups had less evenly distributed grooming, less variation in their number of grooming partners, and less dense grooming networks than smaller groups (Table 1). In addition, the level of intra-group contest competition was positively and significantly linked to the variation in the number of grooming partners (Table 2). Finally, the variation in the number of partners groomed by a particular individual was higher in groups/species facing high predation risk than, respectively, medium or low predation risk. Results of the analyses on the two variables used to calculate IGCC showed a significant effect of, respectively, frequency of inter-group encounters per day on grooming density and the proportion of aggressive encounters on SDOD (supplementary tables S2 and S3).

Males

A phylogenetic signal was detected in the model for SWI ($\lambda = 1$) whereas λ was equal to zero in the models on SDOD and density (Table 2). Contrary to our predictions, the level of IGCC was not significantly affected by any of our intra-group affiliation measures (Table 2). For the control variables, we found that provisioning as well as group size were affected by SWI, i.e. grooming was more evenly distributed in non-provisioned and in larger groups. The other control variables did not have a significant effect on our measures of intra-group affiliation. Supporting our results, the two variables used to calculate IGCC had no effect on within-group affiliation among males (supplementary tables S4 and S5).

DISCUSSION

In contrast to previous reports (Cheney, 1992; Grueter, 2013), our study is the first to provide partial support for the hypothesis of correlated evolution between IGCC and intra-group affiliation and suggests that the exact nature of this effect is sex specific. In males, the level of IGCC had no significant effect on grooming distribution. In females, the level of IGCC had a positive and significant effect on the variation in the number of grooming partners and on grooming network density, but not on SWI. The fact that the frequency of inter-group encounters per day and the proportion of aggressive encounters had different effects on within-group affiliation among females suggest that different aspects of IGCC affects grooming distribution differently. Our findings on female primates, therefore, corroborate hypotheses from socio-ecological models (Alexander and Borgia 1978; Wrangham 1980; Sterck et al., 1997) and evolutionary biology (Reeve and Hölldobler 2007), and are in line with what has been found, and expected to occur in humans (Haas 1990; West et al. 2006;

Choi and Bowles 2007; van Vugt and Park 2009). Because the data available for our analysis were mostly from Old World monkeys, we do not know if these findings also extend to New World Monkeys. In addition, many species of New World monkeys are known to show very little grooming (Dunbar 1991) and intra-group affiliation would have to be measured differently in those species; however, we would expect to find a similar relationship between IGCC and intra-group affiliation using appropriate affiliation measures.

Sex differences

By maintaining less varied and denser networks of intra-group affiliation, female primates may be better able to cope with high levels of IGCC. Other things being equal (e.g. group size) females in strongly affiliated groups might more effectively out-compete other groups and access/monopolise resources (e.g. shelter or water) that are essential to maximise their reproductive outputs. The fact that we found support for our prediction only in females but not males is interesting. As males usually take a more active role in aggressive inter-group encounters than females (Majolo et al. 2005; Kitchen and Beehner 2007) and have been reported to have strong social bonds even in species with male dispersal (Silk 1994; Mitani 2009; Schülke et al. 2010; Young et al 2014), a positive effect of IGCC on intra-group affiliation was predicted to occur among males. The lack of support for this prediction in males may be explained by two main factors. Firstly, the benefits and costs of aggressively taking part in inter-group encounters differ between the two sexes. The benefits males can gain from winning an encounter with another group are likely to be highly skewed in favour of the dominant male, whereas in females benefits are expected to be more evenly distributed (Cheney 1992; Kitchen and Beehner 2007; Crofoot and Wrangham 2010). This might explain why overall levels of same-sex affiliation are not linked to IGCC in males. By out-competing another group, every female can potentially gain benefits like access to food and reduced risk

of immigration by infanticidal males, although these benefits could be somewhat greater for dominant and/or lactating females (Sterck et al. 1997). Conversely, the main benefits that males can gain during/following aggressive confrontations with other groups are opportunities for extra-group copulations and/or for emigration, and defence of receptive females in their group (Kitchen and Beehner 2007; Crofoot and Wrangham 2010). These benefits are likely to not be shared equally by group males as, for example dominant males often monopolize mating opportunities (e.g. Majolo et al. 2012). Indeed, it has been shown that aggressive participation in inter-group encounters in males largely depends on the age and dominance status of the animals (Majolo et al. 2005; Kitchen and Beehner 2007). Secondly, as all species in our sample are female philopatric, social relationships among dispersing males may be much more variable and opportunistic than those in females, and be based on other behaviours (e.g. agonistic support) than grooming (Schülke et al. 2012), which for lack of data could not be included into our analysis. IGCC may be affected by male social relationships but this might not be reflected in the grooming network. Thus, taken together, socio-ecological factors and the different cost/benefits of participation in inter-group confrontations for the two sexes may result in different evolutionary pressure for a correlated evolution between IGCC and intra-group affiliation in males and females.

Confounding variables

Our analysis of the control variables showed that group size was significantly linked to intra-group affiliation in four of our six models, although the effects of group size on our dependent variables were small in terms of the actual model coefficients for all but one (i.e. SDOD in males) variable. It is well known that group size affects the response of animals to collective action problems as well as the outcome of inter-group encounters (Crofoot and Gilby, 2012; Willems and van Schaik, 2015). Group size also affects individual behaviour

and has been shown to have fitness consequences for individual animals (Majolo et al., 2008). Therefore, group size is expected to modulate the relationship between IGCC and intra-group affiliation, which is supported by our results. Similarly, provisioning, the level of intra-group contest competition and predation risk affected some of our measures of intra-group affiliation. All three variables have also been shown to affect inter-individual distances and activity budgets (e.g. Wrangham, 1980; Sterck et al., 1997), which may explain why they were significant in our models. Since all of the species in our dataset are female philopatric, the relationship between grooming and IGCC may also be affected by kinship, that is, females in species/populations facing a higher level of IGCC might have larger matriline, which in turn may result in larger and denser grooming networks. In our dataset, we could not control for kin-relationships (as kinship data were not available for most of the study groups in the dataset) but this would be an interesting hypothesis to test in future studies. By including group size into the models, however, we aimed to control for these effects in our analyses.

From non-human primate to humans

When comparing the relationship between IGCC and intra-group affiliation in humans and non-human animals (Crofoot and Wrangham, 2010; Gintis et al. 2015), three considerations have to be made. First, unlike many non-human primates, in human hunter-gatherer societies both sexes can disperse and establish long-distance and long-lasting social bonds and cooperative relationships (e.g. Apicella et al. 2012). As such, differences in social structure and inter-individual relatedness in humans societies compared to non-human primates could affect correlated evolution between IGCC and intra-group affiliation. Second, inter-group conflicts in non-human animals and human warfare may be two behaviourally different phenomena (Crofoot and Wrangham 2010). Human warfare in small scale societies, despite

lacking the complex structure, strategic approach and forced recruitment of soldiers typical of military confrontations, is often characterized by social norms (e.g. participation in warfare as a sign of adulthood) and punishment rules (e.g. social isolation or reduced power) that favour individual contribution to aggressive encounters (Haas 1990). These factors should act on limiting the occurrence of free-riding and the risk posed by collective action problems in humans compared to non-human primates (Nunn and Lewis 2001; Crofoot and Gilby 2012; Willems et al. 2013; Willems and van Schaik, 2015). Similarly, the different risk posed by collective action problems can explain why the relationship between IGCC and intra-group affiliation appears to be stronger in cooperative breeding birds (Radford 2008a, b, 2011) than in the primate species included in our study. In cooperatively breeding species, group members are often genetically related, share resources with one another (Radford and Fawcett, 2014), and groups are generally small; all conditions that reduce the risk of collective action problems (Willems and van Schaik, 2015) and favour genetic group selection (Reeve and Hölldobler, 2007). Conversely, our dataset was mostly composed of species living in large groups where the benefits and costs of intense IGCC are likely to be skewed across group members. Third, inter-group conflicts in non-human animals and human warfare may bear different consequences for individual fitness. The risk and occurrence of deaths resulting from inter-group aggressive confrontations are thought to have been high during the course of human evolution (Haas 1990; Choi and Bowles 2007; but see Fry and Söderberg 2013), especially since the use of projectile weapons (Sahle et al 2013). Conversely, the risk of deaths resulting from inter-group aggressive confrontations is likely to be lower in non-human vertebrates than in humans, although deaths due to inter-group confrontations have been reported (e.g. lions and anthropoid primates: Packer et al. 1990; Mitani et al. 2010) and estimating the probability of their occurrence is extremely difficult. If such differences in the cost of inter-group aggressive confrontations are indeed relevant, there

might have been stronger evolutionary pressure for an adaptive link between IGCC and intra-group affiliation in the human lineage than in non-human primates or in other phylogenetic taxa.

In conclusion, our study provides partial support for the hypothesis of correlated evolution between IGCC and intra-group affiliation, similar to what has been shown in humans and in cooperatively breeding birds. The fact that a link between IGCC and intra-group affiliation was found in females but not in males, suggests different cost/benefit ratios of inter-group encounters for the sexes. The generality of our findings to non-human primate species not included in this study, and to other taxa, is limited by the scarcity and patchiness of data on this topic. When data become available, further studies will be needed to test the effect of factors that we could not analyse here, such as social organization, breeding system, relatedness and participation patterns in inter-group encounters. In addition, data on more species are required and different measures, especially measures involving bond strength, for intra-group affiliation should be explored. Despite the limitations of our study, our analyses suggest for the first time a relationship between IGCC and intra-group affiliation in non-human female primates.

REFERENCES

- Alexander RD, Borgia G. 1978. Group selection, altruism and the levels of organisation of life. *Ann Rev Ecol Sys* 9: 449–474.
- Apicella CL, Marlowe FW, Fowler JH, Christakis NA. 2012. Social networks and cooperation in hunter-gatherers. *Nature* 481: 497-501.
- Arnold C, Matthews LJ, Nunn CL. 2010. The 10kTrees Website: A new online resource for primate phylogeny. *Evol Anthropol* 19:114-118.
- Aureli F, Schaffner CM, Verpooten J, Slater K, Ramos-Fernandez G. 2006. Raiding parties of male spider monkeys: insights into human warfare? *Am J Phys Anthropol* 131:486-497.
- Borgatti SP, Everett MG, Freeman L. C. 2006. UCInet for Windows: software for social network analysis. Harvard, Massachusetts: Analytic Technologies.
- Bowles S. 2009. Did warfare among ancestral hunter-gatherer groups affect the evolution of human social behaviors? *Science* 324: 1293-1298.
- Brown M, Crofoot M. 2013. Social and spatial relations between primate groups. In: Sterling E, Bynum E, Blair M, editors. *Primate ecology and conservation*. 1st ed. Oxford, UK: Oxford University Press. p. 151-176.
- Cheney DL. 1992. Intragroup cohesion and intergroup hostility: the relationship between grooming distributions and intergroup competition among female primates. *Behav Ecol* 3: 334-345.
- Choi JK, Bowles S. 2007. The coevolution of parochial altruism and war. *Science* 318: 636–640.
- Conradt L, List C. 2009. Group decisions in humans and animals: a survey. *Phil Trans R Soc B* 364: 719-742.
- Crofoot MC, Gilby IC. 2012. Cheating monkeys undermine group strength in enemy territory. *Proc Natl Acad Sci* 109: 501-505.

- Crofoot MC, Wrangham R. W. 2010. Intergroup aggression in primates and humans: the case for a unified theory. In: Kappeler PM, Silk J, editors. Mind the gap: tracing the origins of human universals. 1st ed. New York: Springer. p. 171-195
- Dunbar RIM. 1991. Functional significance of social grooming in primates. *Folia Primatol* 57:121-131.
- Erev I, Bornstein G, Galili R. 1993. Constructive intergroup competition as a solution to the free rider problem in the workplace. *J Exp Soc Psychol* 29:463-478.
- Fry DP, Söderberg P. 2013. Lethal aggression in mobile forager bands and implications for the origins of war. *Science* 341: 270-273.
- Gintis, H., van Schaik, C., & Boehm, C. 2015. Zoon politikon: the evolutionary origins of human political system. *Current Anthropology* 3: 327-353.
- Grueter CC. 2013. No effect of inter-group conflict on within-group harmony in non-human primates. *Commun Integr Biol* 6:e26801.
- Haas J. 1990. The anthropology of war. Cambridge, UK: Cambridge University Press.
- Kitchen DM, Beehner JC. 2007. Factors affecting individual participation in group-level aggression among non-human primates. *Behaviour* 144: 1551–1581.
- Majolo B, Ventura R, Koyama NF. 2005. Sex, rank and age differences in the Japanese macaque (*Macaca fuscata yakui*) participation in inter-group encounters. *Ethology* 111:455-468.
- Majolo B, de Bortoli Vizioli A. & Schino G (2008). Costs and benefits of group living in primates: group size effects on behaviour and demography. *Animal Behaviour*, 76: 1235-1247.
- Majolo B, Lehmann J, de Bortoli Vizioli A, Schino G. 2012. Fitness-related benefits of dominance in primates. *Am J Phys Anthropol* 147: 652-660.
- Miller N, Garnier S, Hartnett AT, Couzin ID. 2013. Both information and social cohesion

- determine collective decisions in animal groups. *Proc Natl Acad Sci* 110: 5263-5268.
- Mitani JC. 2009. Male chimpanzees form enduring and equitable social bonds. *Anim Behav* 77:633-640.
- Mitani JC, Watts DP, Amsler SJ. 2010. Lethal intergroup aggression leads to territorial expansion in wild chimpanzees. *Curr Biol* 20: R507-508.
- Nunn CL, Lewis RJ. 2001. Cooperation and collective action in animal behavior. In: Noë r, van Hooff JARAM, Hammerstein P, editors. *Economics in nature*. 1st ed. Cambridge, UK: Cambridge University Press. p. 42-66
- Orme CDL, Freckleton RP, Thomas GH, Petzoldt T, Fritz SA, Isaac NJB. 2012. CAPER: comparative analyses of phylogenetics and evolution in R. Package version 0.5. Available at <http://cran.r-project.org/web/packages/caper/index.html>. Accessed May 30th, 2013.
- Packer C, Scheel D, Pusey AE. 1990. Why lions form groups: food is not enough. *Am Nat* 136: 1-19.
- Polizzi di Sorrentino E, Schino G, Massaro L, Visalberghi E, Aureli F. 2012 Between-group hostility affects within-group interactions in tufted capuchin monkeys. *Anim Behav* 83: 445-451.
- Puurtilinen M, Mappes T. 2009. Between-group competition and human cooperation. *Proc R Soc B* 267: 355-360.
- R Development Core Team 2011. A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>. ISBN 3-900051-07-0.
- Radford AN. 2008a. Duration and outcome of intergroup conflict influences intragroup affiliative behaviour. *Proc R Soc B* 275: 2787-2791.
- Radford AN. 2008b. Type of threat influences postconflict allopreening in a social bird. *Curr*

Biol 18: R114-115.

Radford AN. 2011. Preparing for battle? Potential intergroup conflict promotes current intragroup affiliation. *Biol Lett* 7: 26-29.

Radford AN, Fawcett TW. 2014. Conflict between groups promotes later defense of a critical resource in a cooperatively breeding bird. *Curr Biol* 24: 2935-2939.

Reeve HK, Hölldobler B. 2007. The emergence of a superorganism through intergroup competition. *Proc Natl Acad Sci* 104: 9736 - 9740.

Rubenstein DI. 1986: Ecology and sociality in horses and zebras. In: Rubenstein DI, Wrangham RW, editors. *Ecological determinants of social evolution*. 1st ed. Princeton: Princeton University Press. p. 282-302

Schülke O, Bhagavatula J, Vigilant L, Ostner J. 2010. Social bonds enhance reproductive success in male macaques. *Curr Biol* 220: 2207-2210.

Silk J. 1994. Social relationships of male bonnet macaques: male bonding in a matrilineal society. *Behaviour* 130:271-291.

Sahle Y, Hutchings WK, Braun DR, Sealy JC, Morgan LE, Negash A & Atnafu B. 2013. Earliest stone-tipped projectiles from the Ethiopian rift date to >279,000 years ago. *PLoS ONE* 8(11): e78092. doi:10.1371/journal.pone.0078092

Sterck EHM, Watts DP, van Schaik CP. 1997. The evolution of female social relationships in nonhuman primates. *Behav Ecol Sociobiol* 41: 291-309.

Tan JHW, Bolle F. 2007. Team competition and the public goods games. *Econom Lett* 96: 133-139.

van Vugt M, Park JH. 2009. Guns, germs, and sex: how evolution shaped our intergroup psychology. *Soc Personal Psychol Compass* 3: 927-938.

Ventura R, Majolo B, Koyama NF, Hardie S, Schino G. 2006. Reciprocation and interchange in wild Japanese macaques: Grooming, co-feeding and agonistic support. *Am J*

Primatol 68: 1138-1149.

West SA, Gardner A, Shuker DM, Reynolds T, Burton-Chellow M, Sykes EM, Guinness MA,

Griffin AS. 2006. Cooperation and the scale of competition in humans. *Curr Biol* 16: 1103–1106.

Wey T, Blumstein DT, Shen W, Jordán F. 2008. Social network analysis of animal behaviour: a promising tool for the study of sociality. *Anim Behav* 75: 333-344.

Willems EP, Hellriegel B, van Schaik CP. 2013. The collective action problem in primate territory economics. *Proc R Soc B* 280:20130081.

Willems EP, van Schaik CP. (2015). Collective action and the intensity of between-group competition in nonhuman primates. *Behav Ecol* 26: 625-631.

Wilson EO, Bossert WH. 1971. *A primer in population biology*. Boston, Massachusetts: Sinauer Associates.

Wrangham RW. 1980. An ecological model of female-bonded primate groups. *Behaviour* 75:262-300.

Young C, Majolo B, Heistermann M, Schülke O, Ostner J (2014). Responses to social and environmental stress are attenuated by strong male bonds in wild macaques. *Proc Natl Acad Sci* 201411450.

Table 1: PGLS results on the effect of the level of IGCC and of the control variables on the three measures of intra-group affiliation among females (n=25; significant results in bold).

Independent variables	Model No. - Dependent variables								
	Model #1 - SWI			Model #2 - SDOD			Model #3 - Density		
	Lambda = 0			Lambda = 0.90			Lambda = 0.76		
	Estimate <i>SE</i>	$\pm$ <i>t</i>	<i>P</i>	Estimate <i>SE</i>	$\pm$ <i>t</i>	<i>P</i>	Estimate <i>SE</i>	$\pm$ <i>t</i>	<i>P</i>
Level of IGCC	0.12±0.39	0.31	0.76	-14.83±3.99	-3.72	<0.01	0.32±0.15	2.15	<0.05
Study period	0.14±0.08	1.62	0.12	0.03±0.98	0.04	0.97	0.06±0.04	1.62	0.12
Provisioning	0.06±0.26	0.25	0.80	-3.94±3.13	-1.26	0.23	-0.07±0.11	-0.63	0.54
Group size	-0.01±0.00	-3.01	<0.01	-0.05±0.02	-2.46	<0.05	-0.00±0.00	-4.36	<0.001
Level of intra-group contest competition	-0.11±0.21	-0.52	0.61	6.16±2.83	2.17	<0.05	-0.04±0.10	-0.42	0.68
Predation risk									
High vs. low	0.18±0.42	0.43	0.67	-18.37±5.03	-3.65	<0.01	-0.18±0.18	-0.99	0.34
High vs. medium	0.17±0.44	0.38	0.71	-14.78±5.48	-2.70	<0.05	-0.26±0.19	-1.35	0.20
Intercept	1.44±0.53	2.74	<0.05	24.83±9.41	2.64	<0.05	0.82±0.29	2.80	<0.05

Table 2: PGLS results on the effect of the level of IGCC and of the control variables on the three measures of intra-group affiliation among males (n=19; significant results in bold).

Independent variables	Model No. - Dependent variables										
	Model #1 - SWI				Model #2 - SDOD			Model #3 - Density			
	Lambda = 1				Lambda = 0			Lambda = 0			
	Estimate <i>SE</i>	±	<i>t</i>	<i>P</i>	Estimate ± <i>SE</i>	<i>t</i>	<i>P</i>	Estimate <i>SE</i>	±	<i>t</i>	<i>P</i>
Level of IGCC	-3.90±2.16		-1.81	0.10	151.43±145.84	1.04	0.32	-0.05±0.22		-0.24	0.81
Study period	4.33±2.15		2.01	0.07	- 107.12±131.31	-0.82	0.44	-0.20±0.20		-0.99	0.35
Provisioning	-2.87±0.87		-3.31	<0.01	12.92±105.03	0.12	0.91	-0.25±0.16		-1.58	0.14
Group size	0.09±0.02		5.14	<0.001	-2.92±2.70	-1.08	0.30	-0.01±0.00		-1.88	0.09
Level of intra-group contest competition	0.66±0.77		0.85	0.41	1.06±81.87	0.01	0.90	0.21±0.12		1.71	0.12
Predation risk											
High vs. low	-0.76±0.61		-1.25	0.24	- 406.93±201.30	-2.02	0.07	-0.11±0.31		-0.37	0.72
High vs. medium	-1.94±1.41		-1.37	0.20	- 411.62±207.39	-1.99	0.07	-0.13±0.32		-0.42	0.70
Intercept	-0.88±7.54		-0.12	0.91	723.55±281.98	2.57	<0.05	0.93±0.43		2.17	<0.05

Figure 1: Scatter plots showing the relationship between the level of IGCC (ln-transformed), on the y axis, and the standard deviation of out degree (SDOD) among female primates.

Figure 2: Scatter plots showing the relationship between the level of IGCC (ln-transformed) and the density of grooming networks among female primates.

Figure 1

Figure 2