**Predictably harsh environment is associated with reduced cognitive flexibility
in wild food-caching mountain chickadees**

Rebecca Croston¹, Carrie L. Branch¹, Angela M. Pitera¹, Dovid Y. Kozlovsky¹, Eli S.

Bridge², Thomas L. Parchman¹, Vladimir V. Pravosudov^{1,*}

Cognition is one of the mechanisms underlying behavioral flexibility, but flexibility of cognition itself may vary as a result of trade-offs between the ability to learn new information and the ability to retain old memories. How and when cognitive flexibility is constrained by this trade-off remains poorly understood. We investigated cognitive flexibility in wild food-caching mountain chickadees in the field at different elevations experiencing different levels of environmental harshness, using a spatial learning and memory reversal paradigm. There were no significant differences in sampling strategies between elevations, but high elevation chickadees performed worse than low elevation chickadees on the reversal task, indicating lower cognitive flexibility. Compared to the initial learning task, low elevation chickadees improved their performance during the reversal task, while high elevation chickadees performed worse. High elevation birds inspected previously rewarding locations more frequently than other locations, suggesting that reduced cognitive flexibility is associated with proactive interference. Considering that high elevation chickadees cache more food and are likely more dependent on these caches than their conspecifics from low elevation, and that chickadees from both elevations use similar sampling strategies, our findings suggest that stronger memories of more caches might interfere with acquisition and retention of new memories. Overall, our results suggest that predictably harsh environments might favor stronger memories at the expense of decreased cognitive flexibility, which is likely driven by increased proactive interference.

Keywords: behavioral flexibility, cognitive flexibility, spatial memory, food caching, chickadee, interference, reversal learning.

INTRODUCTION

Behavioral flexibility has been considered critical for success in highly variable and changing environments (Lefebvre et al. 2004; Bond et al. 2007; Wright et al. 2010). Cognition, defined as the acquisition, processing, storage, and use of information (Dukas 2004; Shettleworth 2010), is one of the main mechanisms allowing for behavioral flexibility (Lefebvre et al. 2004; Dukas 2004). For example, learning can allow animals to rapidly alter their behavior in response to changing environments (Dukas 2004; Shettleworth 2010). Hence, we would expect more complex, predictably unpredictable environments to favor enhanced learning abilities, resulting in the evolution of learning and behavioral flexibility (Lefebvre et al. 2004; Pravosudov & Roth 2013). At the same time, if the environment is completely unpredictable, learning abilities should not be favored because all predictors would be unreliable environmental signals (e.g. Dunlap & Stevens 2009).

Cognition, however, can vary in its degree of flexibility, and learning new information may come as a trade-off with previously learned information in the form of proactive and retroactive interference (Shettleworth 2010; Anderson & Neely 1996). Specifically, previously formed memories might interfere with the ability to form new memories (e.g. proactive interference) and new memories might interfere with recollection of old memories (e.g. retroactive interference; Anderson & Neely 1996). The magnitude of proactive interference has been suggested to depend on the strength (e.g. longevity, persistence) of previous memories; i.e..

stronger memories should cause more interference (Jacoby et al. 2001; Kuhl & Wagner 2009). Proactive and retroactive interference appear to represent a trade-off in which animals might either 1) initially learn new information well, but learn subsequent similar information less well (proactive interference), or 2) learn new information well, but in doing so, weaken or eliminate old memories (retroactive interference). While the evolution of better cognitive abilities concerns both proactive and retroactive interference, most evolutionary behavioral ecologists have focused on what should promote cognitive flexibility (Lefebvre et al. 2004; Bond et al. 2007), often overlooking the potential cognitive trade-offs associated with cognitive flexibility, and the question of when reduced cognitive flexibility might be favored.

Food-caching animals present an interesting opportunity to address the trade-offs associated with cognitive flexibility. It has been hypothesized that food-caching species have evolved specialized spatial memory along with an enlarged hippocampus used in cache recovery (Krebs et al. 1989; Sherry et al. 1989; Pravosudov & Roth 2013). These animals cache large amounts of food items during late summer and autumn and subsequently recover these caches, relying at least in part on spatial memory (Krebs et al. 1989; Sherry et al. 1989), throughout the winter (Pravosudov 2006). There is variation in the degree of caching specialization both within and among species, with more specialized species having higher reliance on food caches and exhibiting more accurate spatial memory and a larger hippocampus (Garamszegi & Eens 2004; Lucas et al. 2004; Pravosudov & Roth 2013). Within species, individuals from populations experiencing harsher winter

climates with higher demands for food caches have also been shown to cache more food, have more accurate spatial memory, larger hippocampi, higher total number of hippocampal neurons, and faster adult hippocampal neurogenesis (Pravosudov & Clayton 2002; Roth & Pravosudov 2009; Roth et al. 2011; Freas et al. 2012). There is some indirect evidence that differences in spatial memory and the hippocampus between some of these populations have been shaped by natural selection (Roth et al. 2012; Freas et al. 2013; Pravosudov & Roth 2013).

Many food-caching animals store and retain large amounts of spatial information about numerous cache locations during a relatively short-term burst of caching, usually in autumn. Preserving these memories is integral to the use of this information, and should be of highest priority. It therefore might be predicted that food-caching animals should be better at initial memory acquisition and retention (which could include the period of creating long-term caches in autumn), at the expense of being able to learn new information equally well, due to proactive interference associated with stronger (e.g. longer lasting) previous memories. In other words, selection for adaptive specialization of memory could be expected to be associated with reduced cognitive flexibility allowing for better retention of older memories. Initially the opposite was hypothesized; that food-caching species should remember specific caching locations while non-caching species should remember all locations visited, which should produce more proactive interference in non-caching species [Clayton & Krebs 1994; Hampton et al. 1998]. However, some available evidence shows higher proactive interference in specialized food cachers (Bond et al. 2007; Hampton et al. 1998; Lewis & Kamil 2006, but see Urhan & Brodin 2015).

Here, we aimed to test whether the degree of caching specialization, resulting from differences in winter climate severity, might be associated with differences in cognitive flexibility in mountain chickadees (*Poecile gambeli*) from high and low elevations. Chickadees living at higher elevations experience predictably harsher and longer winters compared to birds at lower elevations (Freas et al. 2012; Croston et al. 2016). Based on such differences, it was hypothesized that chickadees at high elevation should be more dependent on food caches for survival (Pravosudov & Clayton 2002). Indeed, birds at high elevation cache more food, perform better on spatial memory tasks (both acquisition and retention), have larger hippocampi, and have more hippocampal neurons with higher adult hippocampal neurogenesis rates (Freas et al. 2012, 2013). Moreover, differences in food caching rates and spatial memory were confirmed in the same free-living population (Croston et al. 2016).

In this study, we used a spatial learning and memory reversal task to test for potential differences in cognitive flexibility. Reversal learning is a common method to test for cognitive flexibility in which an individual learns an initial task (e.g. location of a food reward) and once the task is acquired, the rewarding target is changed so that the individual has to learn and remember a new task, such as a new location (Anderson & Neely 1996; Strang & Sherry 2014; Bebus et al. 2016).

Differences in the ability to learn and remember in reversal tasks are interpreted as differences in cognitive flexibility; more flexible individuals are expected to perform better, while less flexible individuals are expected to perseverate and make more errors, potentially due to proactive interference (Lewis and Kamil 2006; Bond et al. 2007). By comparing learning and memory performance on the initial learning task

and the reversal task, and analyzing errors in the reversal task we can provide evidence for proactive interference. Proactive interference would be characterized by worse performance during the reversal task (versus during the initial learning task) associated specifically with visits to the previously rewarded locations (Anderson & Neely 1996; Jacoby et al. 2001).

Variation in sampling strategies could potentially result in disparate performance on a learning and memory task. For instance, if a population or species tends to have a ‘win-stay’ strategy (e.g. repeatedly visiting a rewarding location) they may have more difficulty learning a reversal task than individuals with a ‘win-shift’ strategy (e.g. always going to a new location after visiting a rewarding location), which, in turn should have more difficulty learning an initial learning task with the same rewarding location (Olton & Schlosberg 1978; Sulikowski & Burke 2012). Thus we assessed sampling strategies of chickadees at high and low elevations during the shaping/pre-training period, when all feeders were accessible and a visit to any feeder resulted in reinforcement.

We carried out our study using wild chickadees fitted with unique PIT-tags and ‘smart’ RFID-based feeders arranged in a spatial array directly in the field (Croston et al. 2016). If birds from high elevations are indeed more specialized in food caching and exhibit higher levels of dependence on caches than birds from low elevations, it might be expected that they are also less cognitively flexible.

MATERIALS AND METHODS

Study Subjects and Site

We tested spatial memory performance using wild mountain chickadees (herein called chickadees) fitted with unique PIT-tag identifiers (Croston et al. 2016). The initial spatial learning task involved 167 chickadees (90 from high elevation and 77 from low elevation), and the subsequent spatial reversal task involved 143 of these birds (74 from high elevation and 69 from low elevation). While most of the birds tested in this study were new to this season, there were also chickadees that were tested one year ago (Croston et al. 2016): 27 individuals at high and 9 individuals at low elevation during the initial learning task and 22 individuals at high and 7 individuals at low during the reversal task. These birds were always assigned target feeders different from the last year assignments to avoid any carryover effects. It must be noted that after testing last year, all feeders were kept open and then again all feeders were open at the beginning of testing for this study, minimizing any carryover effects.

Mountain chickadees were trapped using mist-nets at 17 feeder sites and at nests throughout Sagehen Experimental Forest, in the Sierra Nevada, northern California, U.S.A. Birds were trapped at 'low' (ca. 1900 m) and 'high' (ca. 2400 m) elevation sites, following our previous work (Croston et al. 2016), and fitted with unique color band combinations, including one band with a passive integrated transponder (PIT) tag (IB Technology, Leicestershire, U.K.; <http://www.buyrfid.co.uk/store/index.php?route=product/categoryandpath=42>) attached.

Sagehen Experimental Forest is a primarily mixed-conifer forest, which varies predictably in ambient temperature and snow cover across elevations. At

higher elevations, temperatures are lower, snow cover is deeper and longer-lasting throughout the year, and climate is generally harsher than at lower elevations (Freas et al. 2012; Croston et al. 2016).

Spatial memory testing took place at spatial arrays constructed at four separate locations within Sagehen Experimental Forest – two each at high and low elevations following our previous study (Croston et al. 2016). Within elevation, arrays were separated by ca. 1.2 km. No birds moved between arrays during testing.

'Smart' Feeders and Spatial Arrays

We tested initial spatial memory performance and spatial memory reversal using four spatial arrays, each consisting of eight RFID-programmable “smart” feeders mounted on a square frame (122 x 122 cm) with two feeders on each side. Arrays were suspended from trees approximately 2.5-3 m above ground level. Feeders were designed and manufactured specifically for use within this system based on design specifications by E.S.B. and V.V.P., and the RFID design in Bridge and Bonter (2011). Feeder functionality is described in detail in Croston et al. (2016), however, the external feeder mechanism was redesigned for this study (described below).

Briefly, RFID circuitry and data logging were implemented on a circuit board similar to that described in Bridge & Bonter (2011), and a second small board was added that featured a BD6222HFP-TR motor controller (ROHM Semiconductor, Kyoto, Japan). The primary circuit board integrated an RFID module (UB22270, Atmel Corporation, San Jose, CA, USA), microprocessor (PIC16F688, Microchip, Inc., Chandler, AZ, USA), memory module (24LC512, Microchip, Inc.), real-time clock

207 (DS1307, Maxim Integrated Products, Sunnyvale, CA, USA), an SD card (for
208 removable storage), and external device support for integration with the feeder
209 motor and antenna (see Bridge & Bonter 2011 for details). The circuitry controlled a
210 geared motor that powered a rack and pinion system mounted against a door, which
211 moved up and down to restrict and allow access to black-oil sunflower seeds. The
212 top of the feeder door opening was fitted with either a small safety brush (one array
213 at high elevation and one at low elevation) or a small safety door (one array at high
214 elevation and one at low elevation), which restricted access to seed above the closed
215 door, without creating a hard stopper surface that could trap birds when the door
216 was closing. Both of these safety features were identical among the feeders within
217 each array and balanced between elevations, and birds would only interact with
218 these features in very rare cases where they failed to move away from the closing
219 door, as they were located above the feeder opening. Considering the above, we do
220 not believe that these features affected behavior, or differences in behavior, at any
221 arrays. These differences were not planned, but were implemented because of
222 technical issues..

223 Birds accessed seed by landing on a wooden perch, which we embedded with
224 an antenna (approximately 32 x 45 mm and 1.35 mH) made of coiled 30 AWG
225 magnet wire and encased in weatherproof PlastiDip (<http://www.plastidip.com>).
226 We built and tested each antenna in our lab and actual inductance ranged between
227 1.345 and 1.355 mH (an inductance value of 1.350 mH provides optimal RFID
228 detection; <http://animalmigration.org/RFID/RFIDbuild.htm>). Feeders cycled on an
229 automatic timer, and were active from 0600 hours to 1930 hours. When active, they

polled for tags for 0.5 s and then shifted into a low-power mode for 0.5 s. Feeders were powered by lithium-ion batteries (12V, 6800 mAh), which usually last for more than seven days. During testing, we conservatively replaced batteries every second day, and no feeders were found inoperable due to power supply. The exteriors of the feeders were constructed from PVC sheeting and were close to identical in appearance.

The feeder system allowed three operational feed modes, which controlled birds' access to sunflower seeds stored within the feeder hopper. In "open" mode, the access door remains open, allowing any bird to access seed, while recording the RFID, date, and time of each visit by a PIT-tagged chickadees. In "feed all" mode, the access door rested in closed position, but would open for any PIT-tagged chickadee recorded landing on the perch. In "target" mode, the access door would only open for birds whose unique RFID codes were programmed manually into the microprocessor memory – these birds were assigned to that "target" feeder. In each mode, the reader recorded each visit to the feeder perch, regardless of whether the feeder door opened. We collected data from all feeders every second day, and at each data collection we verified feeder parameters and performance with two control PIT-tags.

Memory testing - Initial

Birds were first presented with feeders in "open" mode for approximately 10 weeks, beginning August 28th, 2015. Feeders were then switched to 'feed all' mode on November 11th, 2015, allowing chickadees 19 days to acclimate to the feeder

mechanism and door movement. We switched feeders to ‘target’ mode on November 30th, 2015, triggering the beginning of memory testing. Due to technical issues preventing many individuals from accessing food, we restarted two arrays (one each at high and low elevation), and returned them to “open” mode a second time on December 5th, 2015. This restarted the sequence of acclimating chickadees to the mechanism by allowing access to seed at all feeders within those arrays. Testing and reversal sequences at those arrays were offset from the others by one week. Overall, the initial spatial learning task took place from November 30th until December 18th in two arrays (one from each elevation) and from December 11th until December 30th in the other two arrays.

During the initial learning phase, chickadees were only able to obtain food from a single, randomly assigned ‘target’ feeder, and learning performance was examined relative to that feeder. Chickadees which were either known to have previously visited a given array, or were initially trapped at that same location, were assigned a target feeder within that array. Under this paradigm, following another chickadee to their target feeder is rarely rewarded, and social learning is discouraged. For the birds that tested in the previous year (see Croston et al. 2016), we also avoided assigning targets in the same location within the array as had been rewarded a year ago. It is important to add that immediately following testing last year, all feeders were returned to the open position for more than two months, and food was visible and available again at all feeders. During this study, all feeders were also maintained in ‘open’ position for a long time prior to target assignments.

We recorded all instances of any individual landing at any feeder (a “visit”), and subsequently compiled both the number of visits to non-target feeders (“visit errors”) per trial per individual, and the number of non-target feeders visited (“location errors”) per trial per individual as in our previous study (Croston et al. 2016). We used the number of location errors (number of incorrect feeders visited prior to visiting the rewarding feeder) as a measure of learning (Croston et al. 2016). A trial was defined as beginning either with an individual’s first visit to any feeder within the array in a given day, or with an individual’s first visit subsequent to visiting its target feeder. A trial was defined as ending either with the end of the day (in the case of no visits to the target feeder), or with a visit to the target feeder (Croston et al. 2016).

Memory testing – Reversal

In order to test for chickadees’ ability to learn the location of a new target feeder following learning their initial target, we assigned each bird that participated in the initial testing stage a new target feeder (“reversal target”) immediately following the initial learning task. Reversal testing took place from December 18th until December 30th in the early two arrays and from December 30th until January 15th, 2016 in the latter two arrays. Reversal targets were assigned semi-randomly; we avoided assigning reversal targets adjacent to initial targets for the same individual, as adjacent feeders are sometimes visited as a byproduct of either flight path, or an already-occupied feeder perch. We also avoided assigning the same reversal targets to sets of chickadees which had been assigned the same initial learning target, so

that those individuals could not learn the new location by following the same individuals that were assigned their same previous target feeder. We recorded all instances of any individual landing at any feeder, and compiled visit errors and location errors as during initial learning. Custom Perl scripts were used to process data downloaded from the smart feeders into data files containing information on each individual's feeder activity each day and summary data on the number of errors during each trial.

Foraging strategies

To test whether differences in performance on spatial memory tests could reflect the use of different foraging strategies in exploring the arrays, we examined visit data from the first eight days of the pre-testing period, when all feeders were programmed to provide food to any PIT-tagged chickadee landing on any of the perches. For example, if chickadees forage using a 'win-stay' strategy, they would be expected to continue visiting only one feeder following their first food reward, and to not visit any other feeders even though they all provide rewards (e.g. Olton & Schlossberg 1978). If, on the other hand, chickadees use a 'win-shift' strategy, they would be expected to visit all feeders in the array equally as they should shift to a different feeder after visiting the rewarding feeder (Olton & Schlossberg 1978). We ranked all eight feeders in each array by the frequency of visits (from most visited to the least visited) by each individual, so that visit frequencies can be directly compared across individuals. We must add that this comparison is not limited to

testing for just ‘win-stay’ and ‘win-shift’ strategies and should reveal if chickadees from different elevations sample the feeders differently.

Statistical Analyses

For initial and reversal learning, we compared performance across trials for the entire testing window (separately for both tasks) for each bird, grouped by elevation, using repeated measures generalized linear mixed models (GLMMs) fitted by maximum likelihood with Satterthwaite approximations using the lme4 package in R, as in Croston et al. (2016). The number of location errors per individual per trial was treated as the response variable, with PIT-tag ID as a random variable accounting for unbalanced repeated measures, and elevation (high versus low) and trial number (unique to each individual’s testing sequence) as fixed effects. PIT-tag ID was implicitly nested within elevation in this model and trial number was nested within PIT-tag ID, as this was a repeated measure unique to each individual. Trial number accounted for time across testing because number and timing of trials varied across individuals due to the voluntary nature of testing. For all subsequent analyses, we used mean number of errors per trial averaged across multiple trials, which should provide a more accurate measure accounting for potential differences in timing among different trials.

In our analyses, learning is described by decreasing numbers of location errors per trial, given the assumption that chickadees which have learned their rewarding target feeder will visit only that feeder and will avoid visiting the other un-rewarding feeders. We excluded data from days 17-19 of initial testing due to an

343 anomalous dramatic increase in the number of visit and location errors on these
344 days at all four arrays, which is likely attributable to major shifts in weather
345 associated with a winter storm.

346 We next estimated the mean number of location errors per trial per
347 individual over the entire testing period (separately for the initial learning period
348 and for the reversal learning period), as well as during the first 5, 10 and 20 trials as
349 mean estimates of learning and used them as dependent variables in all subsequent
350 analyses. Comparing the number of errors between the first 5, 10 and 20 trials
351 provides a measure of learning within the first 20 trials; a reduction in the number
352 of errors between the first 5, 10 and 20 trials will indicate learning as individuals
353 are getting better with experience.

354 In order to investigate potential memory mechanisms (e.g. interference)
355 underlying differences in reversal performance between elevations, we examined
356 whether chickadees continued visiting the previously rewarded target (i.e. initial
357 targets) even after these were no longer rewarded (during the reversal phase). We
358 calculated the ratio of visits to the initial targets relative to the total number of visits
359 to all other feeders across the entire reversal test. Errors in the reversal
360 performance associated specifically with visits to the initial target feeder would
361 suggest continuing memory for these targets, which would be consistent with
362 proactive interference.

363 To compare sampling strategies, we calculated the frequency of visits to each
364 feeder during the pre-testing period when all feeders provided food, and ranked
365 feeders by the frequency of visits for each individual. We then directly compared

proportions of visits to each feeder ranking across individuals using a repeated-measures general linear model (GLM) where relative visit frequency for each ranked feeder within an array is treated as the within-subject repeated measure and elevation as the between-subject variable.

Ethical Note

This research adhered to the guidelines of the Institutional Animal Care and Use Committee of the University of Nevada – Reno (Protocol #00603), as well as all local and federal guidelines (California Department of Fish and Wildlife Permit D-0011776516-4). This is a field study, so animals were only individually banded and immediately released. We did not detect any detrimental effects of tagging and collecting blood samples.

RESULTS

Initial spatial learning task

Across the entire initial learning testing window and all trials, chickadees showed significant improvement in performance as measured by reduction in location errors (GLMM, trial number $t = -3.99$; $P < 0.001$), but there were no significant differences between elevations ($t = -1.00$; $P = 0.315$, interaction $P = 0.83$). The mean number of location errors per trial over the testing period was also not significantly different between chickadees from high versus low elevation (GLM, $F_{1,159} = 2.0$, $P = 0.16$; Fig. 1).

388 **FIGURE 1.** *Mean number of location errors per individual per trial across each*
389 *testing period (either initial learning or reversal) at high versus low elevations during*
390 *the initial and the reversal spatial learning tasks. Error bars represent Standard*
391 *Errors.*

392

393

394 Learning was evident during the first five trials, as the mean number of
395 location errors per individual per trial was much lower than would be expected by
396 chance (chance performance is four location errors). Chickadees from both
397 elevations improved significantly during the first 20 trials (repeated-measures
398 ANOVA with the mean number of location errors during the first 5, 10 and 20 trials
399 as a repeated measure; number of trials: $F_{2,318} = 52.04$, $P < 0.001$). There were no
400 significant differences between elevations ($F_{1,159} = 0.25$, $P = 0.61$) and the

interaction between elevation and the number of trials was also not significant ($F_{2,318} = 0.28, P = 0.76$; Fig. 2).

FIGURE 2. Mean number of location errors per individual across the first 5, 10 and 20 trials at high versus low elevations during the initial and reversal spatial learning tasks. Error bars represent Standard Errors.

Reversal spatial learning task

As in the initial testing, chickadees showed significant improvement in memory performance across the entire reversal testing period, as measured by location

errors (GLMM, trial number $t = -3.75$, $P < 0.001$), with no significant differences between elevations ($t = -1.81$, $P = 0.07$), and the interaction between the number of location errors and elevation also not being significant ($t = 1.15$, $P = 0.25$).

When we averaged the number of location errors per trial across the entire testing period, the mean number of location errors per trial was significantly higher in chickadees from high elevation (GLM, $F_{1,141} = 4.70$, $P = 0.03$; Fig. 1), with chickadees from lower elevation showing more accurate spatial learning performance during the reversal task.

When we directly compared performance on the initial learning and on the reversal learning task, there were significant effects of elevation (repeated-measures GLM, $F_{1,140} = 4.47$, $P = 0.036$) and testing stage (initial vs reversal; $F_{1,140} = 6.20$, $P = 0.01$), but the interaction was not significantly different ($F_{1,140} = 1.09$, $P = 0.29$; Fig. 1). Pairwise planned comparisons revealed that high elevation chickadees performed significantly worse than both low elevation chickadees on the reversal task, ($P = 0.02$) and their own performance on the initial learning task ($P = 0.01$). There were no significant differences between high and low elevation chickadees at the initial learning task ($P = 0.28$) and no significant differences between the initial learning and the reversal task in low elevation chickadees ($P = 0.31$).

There were small but significant differences in the mean number of birds present at each array during each individual's testing (the mean number of birds present at each day of testing for each bird across their unique testing window, as in Croston et al. 2016) between elevations (Initial learning: high elevation – 35.1, SE = 0.8, low elevation – 32.4, SE = 0.8, $F_{1,159} = 5.69$, $P = 0.02$; Reversal learning: high

elevation – 36.6, SE = 0.7, low elevation – 33.2, SE = 0.7, $F_{1,141} = 10.76$, $P = 0.001$). To control for potential effects of social environment, we included the mean number of birds present per day for each individual bird's testing window as a covariate. . The effect of the number of birds present during testing on learning performance was significant ($F_{1,139} = 12.2$, $P < 0.001$), but despite this, the effect of elevation ($F_{1,139} = 8.4$, $P = 0.004$). and testing phase (Initial vs reversal; $F_{1,139} = 14.3$, $P < 0.001$) were significant. The interaction between elevation and reversal was not significant ($F_{1,139} = 3.2$, $P = 0.07$). Pairwise planned comparisons revealed the same patterns as when social environment is not included: high elevation birds performed worse than low elevation birds during the reversal ($P = 0.01$) and worse compared to the initial learning task ($P = 0.009$). There were no differences in performance between high and low elevation chickadees in the initial learning task ($P = 0.26$) and there was no significant difference in performance between the initial and the reversal tasks in low elevation chickadees ($P = 0.29$).

As during the initial learning task, learning during the reversal was already evident during the first five trials, as the mean number of location errors per individual per trial was lower than would be expected if chickadees were searching randomly (Fig. 2). Learning performance improved significantly during the first 20 trials (repeated-measures ANOVA with the mean number of location errors during the first 5, 10 and 20 trials as a repeated measure; number of trials: $F_{2,282} = 64.06$, $F < 0.001$), but chickadees from high elevation performed significantly worse than birds from low elevation as indicated by the higher mean number of location errors per trial (elevation effect: $F_{1,141} = 8.12$, $P = 0.005$), while the interaction between

elevation and the sampling period (e.g. first 5, 10 or 20 trials) was not significant ($F_{2,282} = 2.70, P = 0.07$; Fig. 2).

Compared to low elevation chickadees, chickadees from high elevation visited their initial target location significantly more often relative to their total number of visits to all other feeders across the entire reversal task (GLM, Elevation: $F_{1,140} = 13.83, P < 0.001$; total number of visits to all other feeders as a covariate: $F_{1,140} = 25.58, P < 0.001$; Fig. 3A). Furthermore, the mean number of location errors per trial over the entire reversal testing was significantly and positively associated with the relative frequency of visits to the initial target feeder during the entire reversal task (regression, $b = 0.30, t_{140} = 3.73, P < 0.001$; elevation: $b = -0.16, t_{140} = -1.97, P = 0.05$; adjusted $R^2 = 0.11$; Fig. 3B). Individuals that visited the previously rewarded locations more frequently also made significantly more location errors during the reversal task. When the relative frequency of visits to the initial targets during the entire reversal trial was added as a covariate to the repeated-measures GLM comparing the mean number of location errors per trial during the first 5, 10 and 20 trials, it was also statistically significant ($F_{1,140} = 6.49, P = 0.01$), and the differences between elevations ($F_{1,140} = 7.27, P = 0.007$) and between sampling periods (first 5, 10 and 20 trials; $F_{2,280} = 56.17, P < 0.001$) remained statistically significant.

FIGURE 3. A. *Total number of visits to the initial target location across the entire reversal testing period relative to the total number of visits to all other feeders (least square means from the GLM with the total number of visits to the initial target*

across the entire reversal period as a dependent variable and the total number of visits to all other feeders across the entire reversal period as a covariate). B: The relationship between the mean number of location errors per individual per trial across during the entire reversal testing period and the ratio of the number of visits to the previously rewarded location to the total number of visits to all other feeders. Error bars represent Standard Errors.

A.

B.

Compared to performance on the initial learning task, chickadees from high elevation performed significantly worse, on average, during the reversal as indicated by the significantly higher mean number of location errors per individual per trial over the entire reversal testing (pairwise planned comparisons, $P = 0.01$; repeated-measures ANOVA, elevation: $F_{1,140} = 4.47$, $P = 0.04$, initial vs reversal: $F_{1,140} = 6.21$, $P = 0.01$, interaction: $F_{1,140} = 1.09$, $P = 0.29$; Fig. 1). Low elevation chickadees' performance did not differ from reversal to initial learning (pairwise planned comparisons, $P = 0.31$).

Likewise, when we compared the initial and the reversal learning performance during the first 5, 10 and 20 trials, chickadees from high elevation made significantly more location errors during the reversal task (pairwise planned

comparisons, $P < 0.03$ for 5, 10 and 20 first trials; repeated-measures ANOVA, elevation: $F_{1,140} = 3.34$, $P = 0.07$; initial vs reversal: $F_{1,140} = 0.07$, $P = 0.78$; interaction with elevation: $F_{1,140} = 5.13$, $P = 0.02$; number of trials: $F_{2,280} = 107.14$, $P < 0.001$; interaction with elevation: $F_{2,280} = 0.71$, $P = 0.49$). In this analysis, however, low elevation chickadees, performed significantly better on the reversal task during the first 5, 10 and 20 trials (pairwise planned comparisons, $P < 0.006$ for all comparisons, Fig. 2) than during the initial learning task.

When the total number of birds present during testing was included as a covariate, it did not have a significant effect on initial task performance (repeated-measures GLM, $F_{1,139} = 0.26$, $P = 0.61$), or elevation ($F_{1,139} = 3.59$, $P = 0.06$) but there was a significant interaction between elevation and testing phase ($F_{1,139} = 6.33$, $P = 0.01$). The pairwise comparisons showed the same patterns as in the analyses where social environment (number of birds present at the array) was not included.

Across both elevations, individuals that performed better during the initial learning task also performed better on the reversal learning task, albeit the positive slope of regression was significantly less than one (regression on the mean location errors during the entire testing period: $b = 0.33$, $SE = 0.08$, $t_{139} = 4.17$, $P < 0.001$; elevation: $b = -0.12$, $t_{139} = -1.54$, $P = 0.13$; Fig. 4). These results suggest that an improvement in one task is associated with only incremental improvement in the other task.

FIGURE 4. *The relationship between the mean number of location errors per individual per trial during the initial learning period and the mean number of location*

errors per bird per trial during the reversal learning period. Error bars represent Standard Errors.

Learning performance and the number of trials

While there were no significant differences between low and high elevation in the mean number of trials completed during the initial learning period (high elevation mean = 1238.7 trials per individual, SE=110.6, n = 90; low elevation mean = 1117.9 trials per individual, SE = 96.8, n = 77; GLM, $F_{1,165} = 0.65$, $P = 0.42$), chickadees from high elevation completed significantly more trials during the reversal learning task (high elevation mean = 1466.9 trials per individual, SE = 192.9, n = 74; low elevation mean = 1002.1 trials per individual, SE = 79.1, n = 69; GLM, $F_{1,141} = 4.72$, $P = 0.031$).

The total number of trials completed during the initial learning testing period was significantly negatively associated with spatial memory performance (mean number of location errors) as early as the first five trials (regression, number of trials: $b = -0.23$, $t_{158} = -3.01$, $P = 0.003$, elevation: $b = 0.03$, $t_{158} = 0.41$, $p = 0.68$, adjusted $R^2 = 0.04$), as well as after 10 trials ($b = -0.30$, $t_{158} = -3.94$, $P < 0.001$, elevation: $b = 0.01$, $t_{158} = 0.08$, $P = 0.94$, adjusted $R^2 = 0.08$) and 20 trials ($b = -0.37$, $t_{158} = -5.03$, $P < 0.001$, elevation: $b = 0.01$, $t_{158} = 0.05$, $P = 0.96$, adjusted $R^2 = 0.13$; Fig. 5A). Memory performance during just the first 5, 10 or 20 trials is a significant predictor of how many trials that individual will complete, and chickadees that go on to complete more trials show better memory from the very onset of testing (Fig. 5A).

FIGURE 5. *The relationship between the mean number of location errors per individual per trial during the first 20 trials and the total number of trials for that individual during the initial learning testing (A) and during the reversal learning task (B).*

565 A.

566

567

568

569

570

571

572

573

574 B.

The same significance patterns were present during the reversal learning; individuals that completed more trials showed significantly better performance (as indicated by fewer location errors) as early as the first five trials (regression, number of trials: $b = -0.21$, $t_{140} = -2.57$, $P = 0.01$, elevation: $b = -0.28$, $t_{140} = -3.50$, $P < 0.001$, adjusted $R^2 = 0.09$), 10 (number of trials: $b = -0.34$, $t_{140} = -4.28$, $P < 0.001$, elevation: $b = -0.27$, $t_{140} = -3.45$, $P < 0.001$, adjusted $R^2 = 0.14$) and 20 (number of trials: $b = -0.38$, $t_{140} = -4.95$, $P < 0.001$, elevation: $b = -0.27$, $t_{140} = -3.55$, $P < 0.001$, adjusted $R^2 = 0.17$; Fig. 5B) trials. Individuals that ultimately completed more trials made fewer location errors during the first 5-20 trials (Fig. 5B).

Sampling strategies

During the first eight days of the pre-testing period, when all feeders in the array were programmed to provide food to any PIT-tagged individual, each chickadee visited the array on average 731.2 times (SE=43.3, n = 78) at low elevation and 554.3 times (SE=44.5, n = 74) at high elevation. At both elevations, individuals primarily visited just one feeder (ca. 40% of all visits, Fig. 6), with the five least visited feeders combined accounting for only approximately 25% of feeder visits (Fig. 6). The differences between the proportion of visits to different feeders within an array were highly significant (repeated-measures GLM with feeder rank as a within-subject variable and elevation as between-subject variable; feeders – $F_{7,1050} = 353, P < 0.0001$) while there were no significant differences between chickadees from high and low elevations ($F_{1,150} = 0.00, P = 0.57$) and the interaction between the elevation and the feeders was also not statistically significant ($F_{7,1050} = 0.00, P = 0.97$; Fig. 6). These data suggest that chickadees at both elevations use similar sampling strategy.

FIGURE 6. *Mean proportion (and Standard Errors) of visits to every feeder in the array during the first 8 days of the pre-testing period, when all feeders opened for all PIT-tagged birds. For each individual, the feeders are ranked from the most (1) to the least (8) visited.*

DISCUSSION

Chickadees from harsher, high elevations performed significantly worse on the spatial learning and memory reversal task than did chickadees from milder, low elevations. Performance during the initial learning task did not differ. Moreover, compared to their own initial learning and memory performance, chickadees from high elevation performed worse on the reversal task, while chickadees from low elevation performed better. During the reversal task, high elevation chickadees visited their previously rewarded sites relatively and significantly more often than low elevation chickadees, and individuals that visited their previously rewarded sites relatively more often also made more overall location errors. At the same time, chickadees from both elevations appeared to use the same sampling strategy, as

indicated by visiting all feeders according to the same frequency patterns at both elevations during the pretesting period, and heavily favoring just one rewarding feeder despite all feeders providing food.

Taken together, these results (1) demonstrate significant variation in behavioral flexibility in spatial learning and memory across elevations and (2) suggest that such variation is consistent with differences in proactive interference as indicated by errors associated specifically with the previously rewarded location. Given exactly the same design of spatial arrays, the fact that low elevation chickadees learned and remembered the new spatial location within the same array equally well as the initial location and that birds from both elevations learned and remembered the initial location equally well, proactive memory interference is the most likely explanation for detected differences. High elevation chickadees appear to exhibit more proactive interference, which might negatively affect their ability to learn and remember new locations in a reversal-learning task within the same testing paradigm. Low elevation chickadees, on the other hand, did not appear to experience proactive interference, as their performance improved on the reversal relative to the initial learning task. Overall, our data suggest that high elevation chickadees are less cognitively flexible than low elevation chickadees, which could potentially be associated with stronger (e.g. longer lasting) memories for the initially learned spatial information.

The ability to successfully learn and remember spatial locations has been linked to different foraging strategies, and, more specifically, to ‘win-stay’ and ‘win-shift’ strategies (e.g. Olton & Schlosberg 1978; Shettleworth 2010). Animals

predisposed to the 'win-shift' sampling strategy are predicted to shift location following each reward, leading to poor performance on spatial memory tasks with a single location. Animals predisposed to the 'win-stay' sampling strategy are expected to visit the same rewarding location repeatedly, which may lead to worse performance in task reversal, as this would require sufficient exploration to find their new rewarding location (Olton & Schlossberg 1978; Shettleworth 2010). If chickadees from high and low elevations use different sampling strategies, this might have contributed to the differences in reversal performance that we detected. Our data, however, indicate that chickadees at high and low elevations follow similar patterns in feeder visits, and chickadees at both high and low elevations repeatedly visit the same single feeder even when all feeders in an array provided food. Given that chickadees from high and low elevations visit rewarding feeders in similar patterns, and that chickadees do sample all of the feeders within the array, differences in proactive interference appear to be most consistent with our results.

Hypothetically, it is possible that the differences between elevations were due to differences in other cognitive abilities, such as the ability to innovate, or problem solving. If birds are more innovative and better at problem solving, they might be able to learn the reversal task faster than less innovative conspecifics. Our previous data, however, demonstrate that high elevation birds are better at problem solving than low elevation birds, and there are no significant differences in neophobia between high and low elevation chickadees (Kozlovsky et al. 2015). Yet here, high elevation birds performed less accurately on the reversal learning and memory task.

681 Our findings are especially interesting in light of earlier hypotheses
682 suggesting that specialized food-caching species should not experience proactive
683 interference (Clayton & Krebs 1994; Hampton et al. 1998; Hampton & Shettleworth
684 1996). The initial argument (Clayton & Krebs 1994) was that non-caching species
685 might remember both rewarded and non-rewarded locations equally well, resulting
686 in higher memory load and hence higher proactive interference, while food-caching
687 species were suggested to form strong memories specifically of rewarded locations,
688 resulting in less proactive interference. Direct tests of this hypothesis found
689 equivocal results (Hampton & Shettleworth 1996; Hampton et al. 1998; Urhan &
690 Brodin 2015). For example, Hampton et al. (1998) reported more proactive
691 interference in a food-caching species, which was associated with better memory of
692 previous locations, just as in our study. In addition, when non-caching species made
693 errors they were equally likely to be associated with both familiar and novel
694 locations. Highly specialized Clark's nutcrackers (*Nucifraga columbiana*) have also
695 been shown to exhibit proactive interference (Lewis & Kamil 2006). Urhan and
696 Brodin (2015) failed to detect proactive interference in marsh tits (*Poecile*
697 *palustris*). In their paradigm birds recovered their food caches at each trial and did
698 not need to remember specific rewarding locations in subsequent trials.
699 In our experiment, however, birds were trained that the target location always
700 provides food over hundreds of trials, and so when the rewarding location was
701 changed, they were unable to unlearn the initial target quickly even after it stopped
702 providing food. That our birds' errors were due to persistent visits to the previously
703 rewarded location, and that low elevation chickadees performed equally well during

the reversal, suggests proactive interference (e.g. Anderson & Neely 1996; Hampton et al. 1998; Jacoby et al. 2001).

It was impossible for us to use a proper control group of birds which would only be used in the reversal test for the first time (e.g. Jacoby et al. 2001), and compared to the birds that participated in both initial learning and reversal. Our low elevation chickadees, however, performed equally well in both initial and reversal tasks using exactly the same spatial arrays as high elevation birds. In addition, the same feeders that rewarded some birds during the initial learning were used for other birds during the reversal, so all feeders were used simultaneously.

Considering that there are also no differences in sampling strategy or neophobia (Kozlovsky et al. 2015), differences in performance on the reversal, but not on the initial learning and memory indicate that reversal performance differences were not the result of an uncontrolled variable only present at one elevation or during one testing phase.

In our study system, chickadees from high elevations appear to be more reliant on food caches, as indicated by higher food caching rates both in the lab (Freas et al. 2012) and in the field (Croston et al. 2016), better spatial memory acquisition (Freas et al. 2012; Croston et al. 2016) and retention (Freas et al. 2012), as well as by significantly larger hippocampi, a larger total number of hippocampal neurons and higher adult hippocampal neurogenesis rates (Freas et al. 2012, 2013). Our study suggests that chickadees with better spatial memory and a larger hippocampus are less cognitively flexible, perhaps due to associated differences in proactive interference. If higher reliance on memory associated with larger memory

loads (e.g. more caches) is associated with higher costs of proactive interference, then cognitive flexibility might represent a trade-off between stronger, longer-lasting memories and the ability to quickly learn and remember new information while forgetting previously learned, potentially no longer useful information. For example, experimental impairment of initial spatial learning performance was associated with enhanced subsequent spatial learning reversal performance in lab rats (Walsh et al. 2011), supporting the idea of an initial vs. reversal memory trade-off. Additional support for this idea comes from studies showing a negative correlation between performance on initial and reversal tasks; individuals that perform better during initial learning, perform worse during reversal and vice versa (Bebus et al. 2016).

Adult hippocampal neurogenesis has been suggested to combat effects of proactive interference via mediating forgetting (Epp et al. 2016). Yet in our system chickadees from high elevation were previously shown to have significantly higher hippocampal neurogenesis rates (Freas et al. 2012), but in this study they seemed to exhibit stronger proactive interference. Because chickadees at high elevations cache more food and likely have a higher memory load, faster neurogenesis might assist with minimizing proactive interference and mediate forgetting of retrieved caches. Forgetting locations of retrieved caches while retaining memories of untapped caches might be a more beneficial strategy than retaining memories of all cache locations in order to avoid re-visiting them (Bond et al. 2007). If that were the case, high elevation chickadees would initially form strong memories of cache locations in autumn and once caching stopped and recovery began, they would only need to

keep track of remaining caches while gradually forgetting locations of recovered caches. This process may be facilitated by higher hippocampal neurogenesis rates, albeit this might not be enough to completely eliminate proactive interference.

Chickadees, which are more reliant on food caches in harsher environments, might be better at initial spatial learning and memory of their cache locations at the expense of reduced cognitive flexibility. Considering that (a) food caching is seasonal, (b) cache locations rarely change (only in the case of re-caching), and (c) caches are recovered throughout winter, and having accurate, long lasting memories of these numerous cache locations should provide greater survival benefits than the ability to constantly learn and remember new, previously unused spatial locations, if such ability might compromise older memories (e.g. retroactive interference). At the same time, once caches have been retrieved, instead of remembering these locations as emptied (Bond et al. 2007), forgetting them might be a simpler mechanism preventing animals from revisiting these previously depleted sites.

Considering that these birds cache thousands of food items during autumn, an important question is whether proactive interference has a negative effect on remembering numerous cache locations during caching. Spreading caches is a behavior that has been suggested to function in minimizing pilferage risk (Waite & Reeve 1993), but since proactive interference appears to be most pronounced in cases where new information is extremely similar to old information (Anderson & Neely 1996), cache spreading may also function to mitigate effects of proactive interference during caching itself. For example, when Clark's nutcrackers (*Nucifraga columbiana*) were tested on a reversal learning task with new spatial cues they

performed significantly better than on a similar task where reversal trials were conducted with the same cues present as during the initial learning (Lewis et al. 2013). Chickadees from lower elevations associated with milder environments do not cache as many food items and appear to rely less on food caches (Freas et al. 2012), therefore, being more cognitively flexible might be more advantageous even if it might compromise previous memories.

It has been suggested that better learning ability is associated with behavioral flexibility as indicated by higher innovation rates (Lefebvre et al. 2004). Considering that behavioral flexibility is related specifically to reversal learning (i.e. ability to quickly learn a new rule or set of information) and that better reversal learning might come at the cost of forgetting previously learned information, it is important to differentiate between initial and reversal learning, as they might be negatively associated (Bebus et al. 2016). Better performance in reversal learning and memory tasks is frequently associated with less proactive interference and more forgetting (Anderson & Neely 1996), therefore, it appears to be a good measure of cognitive flexibility. Interestingly, in our mountain chickadee system, low elevation chickadees appear to be more flexible, at least in the spatial context, yet they were worse at novel problem solving compared to high elevation chickadees (Kozlovsky et al. 2015). We do not know, however, if flexibility in spatial learning extends to flexibility in all cognitive arenas. Nevertheless, the association between learning, reversal learning, and innovation might be more complex than presented in Lefebvre et al. 2004) and cognitive flexibility may not necessarily extend to novel problem solving ability.

796 It is also important to consider potential differences in timing ‘between trials’
797 (inter-trial intervals) or retention intervals, as longer inter-trial intervals might
798 require better memory retention. During the initial learning stage, there were no
799 significant elevation-related differences in the number of trials completed over
800 exactly the same testing time period, suggesting that there were no significant
801 differences in the inter-trial interval times on average. During the reversal stage,
802 high elevation chickadees completed more trials, suggesting shorter inter-trial
803 intervals on average. Despite such shorter inter-trial intervals, high elevation birds
804 performed less accurately, consistent with proactive interference. Considering that
805 individuals completed hundreds and sometimes thousands of trials over the course
806 of testing, and that we average performance data over many trials, our estimates
807 likely reflect real differences in spatial memory performance rather than artifacts of
808 differences in trial timing and memory retention. Given hundreds of trials, animals
809 should successfully learn the association with a single continuously rewarding
810 location. The fact that the mean performance over just the first 5, 10 or 20 trials
811 highly correlated with the mean performance over hundreds of trials suggests that
812 our measures are accurate. In addition, when the total number of trials was included
813 as a covariate to account for potential differences in the ‘between trials’ interval, the
814 outcomes of all analyses remained essentially the same.

815 In contrast with both of our previous field and lab studies that demonstrated
816 reliance on spatial rather than non-spatial cues (Freas et al. 2012; Croston et al.
817 2016), we detected no significant elevation-related differences in spatial memory
818 during the initial learning task. Winter conditions during data collection for the

819 present study were notably more severe than those during data collection for our
820 earlier work. It is unclear how these harsher winter conditions might directly affect
821 memory performance in wild chickadees and, unfortunately, we cannot directly
822 compare performance between these two studies because of significant changes in
823 feeder appearance and design.

824 It is possible that memory performance might change between years and
825 depend on some differences in developmental condition, that our testing paradigm
826 is too simple to consistently detect elevation-related differences, or that other
827 aspects of memory drive these differences. For example, we previously showed that
828 high elevation chickadees have better memory retention under laboratory
829 conditions than do low elevation chickadees (Freas et al. 2012), although we have
830 not directly tested this in the field. It remains unknown whether memory capacity
831 differs between chickadees from high versus low elevations. It is possible that the
832 observed differences in retention length may actually reflect differences in memory
833 capacity such that low elevation chickadees have lower memory capacity and “write
834 over” older memories earlier than do high elevation chickadees given equal
835 numbers of cache locations.

836 Our previous study also linked caching intensity with spatial memory
837 performance, showing a positive association between the total number of trials
838 completed by an individual, and performance on the spatial memory task (Croston
839 et al. 2016). This study has revealed the same pattern. Memory performance during
840 the first 5, 10 and 20 trials in individuals from both elevations were significant
841 predictors of the total number of trials completed during both the initial and

reversal phases of testing. These data suggest that better memory performance in individuals that completed more trials cannot be explained by more experience, as it was already evident during the first five trials.

Overall, our study demonstrated significant differences in cognitive flexibility and suggested that such differences are associated with proactive interference in food-caching birds that have specialized spatial memory. It is possible that these differences are a direct product of natural selection, or alternatively, they might be a byproduct of potential differences in memory load and retention. If selection favors increased memory retention, it might come with costs of memory interference. Further research is needed to understand the origin and maintenance of these differences. Our results, combined with our previous studies of memory in this system showing more accurate and longer lasting spatial memory in chickadees from high elevations (Freas et al. 2012; Croston et al. 2016), suggest that animals might face a trade-off between keeping strong long-term memories at the expense of being less flexible due to proactive interference and being more cognitively flexible, likely at the expense of forgetting previously acquired information. The relative importance of previously learned information should determine which side of this trade-off provides higher fitness benefits under different environmental constraints. For highly specialized food cachers, remembering locations of previously made caches should be the highest priority and hence these individuals might be expected to be less cognitively. Further studies should provide direct tests of proactive interference as a potential mechanism underlying differences in reversal learning and cognitive flexibility.

865

866 AUTHORS' CONTRIBUTIONS

867 VP designed the experiment, participated in field data collection and analyses and in
868 writing the manuscript; RC participated in designing the experiment, collected field
869 data and participated in analyses and in writing the manuscript; CLB, AMP and DYK
870 participated in field data collection and in writing the manuscript; EB provided
871 expertise with designing and running RFID-based feeders and participated in
872 writing the manuscript; TP wrote computer code necessary for processing and
873 organizing raw data from RFID-based feeders and participated in writing the
874 manuscript. All authors gave final approval for publication.

875

876 ACKNOWLEDGEMENTS

877 This study was supported by the National Science Foundation, grant IOS-1351295,
878 to VVP. Initial development of the low-cost RFID units was assisted by funding from
879 the National Science Foundation (awards IOS-0919528 and DBI-1458402 to ESB).
880 Critical and constructive criticisms from Anders Brodin, an anonymous reviewer
881 and the Editor significantly improved the manuscript.

882

883 REFERENCES

884 Anderson, M. C., Neely, J. H. (1996). *Interference and inhibition in memory retrieval*.
885 In: "Memory. Handbook of Perception and Cognition", E.L.Bjork & R.A. Bjork
886 (Eds.), pp. 237-313, San Diego, CA, Academic Press.

887 Bebus, S. E., Small, T. W., Jones, B. C., Elderbrock, E. K., Schoech, S. J. (2016).
888 Associative learning is inversely related to reversal learning and varies with
889 nestling corticosterone exposure. *Animal Behaviour*, 111, 251-260.

890 Bond, A. B., Kamil, A. C., Balda, R. P. (2007). Serial reversal learning and the
891 evolution of behavioral flexibility in three species of North American Corvids
892 (*Gymnorhinus cyanocephalus*, *Nucifraga columbiana*, *Aphelocoma californica*).
893 *Journal of Comparative Psychology*, 121, 372-379.

894 Bridge, E. S., Bonter, D. N. (2011). A low-cost radio frequency identification device
895 for ornithological research. *Journal of Field Ornithology*, 82, 52-59.

896 Clayton, N. S., Krebs, J. R. (1994). One-trial associative memory: comparison of food-
897 storing and nonstoring species of birds. *Animal Learning and Behavior*, 22,
898 366-372.

899 Croston, R., Branch, C. L., Kozlovsky, D. Y., Roth, T. C., LaDage, L. D., Freas, C. A.,
900 Pravosudov, V. V. (2015). Potential mechanisms driving population variation
901 in spatial memory and the hippocampus in food-caching chickadees.
902 *Integrative and Comparative Biology*, 55, 354-371.

903 Croston, R., Kozlovsky, D. Y., Branch, C. L., Parchman, T. L., Bridge, E. S., Pravosudov,
904 V. V. (2016). Individual variation in spatial memory performance in wild
905 mountain chickadees from different elevations. *Animal Behaviour*, 111, 225-
906 234.

907 Dukas, R. (2004). Evolutionary biology of animal cognition. *Annual Reviews in*
908 *Ecology, Evolution and Systematics*, 347-374.

909 Dunlap, A. S., Stephens, D. W. (2009). Components of change in the evolution of
 910 learning and unlearned preference. *Proceedings of the Royal Society B*, 276,
 911 3201-3208.

912 Epp, J. R., Mera, R. S., Kohler, S., Jesselyn, S. A., Frankland, P. W. (2016).
 913 Neurogenesis-mediated forgetting minimizes proactive interference. *Nature*
 914 *Communication* 7,10838.

915 Freas, C. A., Bingman, K., LaDage, L. D., Pravosudov, V. V. (2013). Untangling
 916 elevation-related differences in the hippocampus in food-caching mountain
 917 chickadees: the effect of a uniform captive environment. *Brain Behavior and*
 918 *Evolution*, 82, 199-209.

919 Freas, C. A., LaDage, L. D., Roth, T. C., Pravosudov, V. V. (2012). Elevation-related
 920 differences in memory and the hippocampus in mountain chickadees, *Poecile*
 921 *gambeli*. *Animal Behaviour*, 84, 121–127.

922 Garamszegi, L., Eens, M. (2004). The evolution of hippocampus volume and brain
 923 size in relation to food hoarding in birds. *Ecology Letters*, 7, 1216-1224.

924 Hampton, R. R., Shettleworth, S. J., Westwood, R. P. (1998). Proactive interference,
 925 recency, and associative strength: comparisons of black-capped chickadees
 926 and dark-eyed juncos. *Animal Learning and Behavior*, 26, 475-485.

927 Hampton, R. R., Shettleworth, S. J. (1996). Hippocampus and memory in a food-
 928 storing and in a nonstoring bird species. *Behavioral Neuroscience*, 110, 946-
 929 964.

930 Jacoby, L. L., Hay, J. F., Debner, J. A. (2001) Proactive interference, accessibility bias,
 931 and process dissociations: valid subjective reports of memory. *Journal of*
 932 *Experimental Psychology*, 27, 686-700.

933 Kozlovsky, D. Y., Branch, C. L., Pravosudov, V. V. (2015). Problem solving ability and
 934 response to novelty in mountain chickadees (*Poecile gambeli*) from different
 935 elevations. *Behavioral Ecology and Sociobiology*, 69, 635-643.

936 Krebs, J. R., Sherry, D. F., Healy, S. D., Perry, V. H., Vaccarino, A. L. (1989).
 937 Hippocampal specialization of food-storing birds. *Proceedings of the National*
 938 *Academy of Sciences USA*, 86, 1388–1392.

939 Kuhl, B. A., Wagner, A. D. (2009). *Forgetting and retrieval*. In: “Handbook of
 940 Neuroscience for the Behavioral Sciences”, G. G. Berntson & J. T. Cacioppo,
 941 Eds., John Wiley & Sons.

942 Lefebvre, L., Reader, S. M., Sol, D. (2004). Brains, innovations and evolution in birds
 943 ad primates. *Brain Behavior and Evolution*, 63, 233-246.

944 Lewis, J. L., Kamil, A. C. (2006). Interference effects in the memory for serially
 945 presented locations in Clark’s nutcrackers, *Nucifraga columbiana*. *Journal of*
 946 *Experimental Psychology: Animal Behavior Processes*, 32, 407-418.

947 Lewis, J. L., Kamil, A. C., Webbink, K. E. (2013) Changing room cues reduces the
 948 effects on proactive interference in Clark’s nutcrackers, *Nucifraga*
 949 *columbiana*. *Journal of Experimental Psychology*, 39, 187-192.

950 Lucas, J. R., Brodin, A., de Kort, S. R., Clayton, N. S. (2004). Does hippocampal volume
 951 size correlate with the degree of caching specialization? *Proceedings of the*
 952 *Royal Society B*, 271, 2423-2429

953 Olton, D. S., Schlosberg, P. (1978). Food-searching strategies in young rats: win-shift
 954 predominates over win-stay. *Journal of Comparative and Physiological*
 955 *Psychology*, 92: 609-618.

956 Pravosudov, V. V., Clayton, N. S. (2002). A test of the adaptive specialization
 957 hypothesis: population differences in caching, memory, and the hippocampus
 958 in black-capped chickadees (*Poecile atricapilla*). *Behavioral Neuroscience*,
 959 116, 515–522.

960 Pravosudov, V. V., Roth, T. C. (2013). Cognitive Ecology of Food Hoarding: The
 961 Evolution of Spatial Memory and the Hippocampus. *Annual Reviews in*
 962 *Ecology, Evolution and Systematics*, 44, 173–193.

963 Pravosudov, V. V. (2006). On seasonality of food caching behavior in parids: do we
 964 know the whole story? *Animal Behaviour*, 71, 1455-1460.

965 R Core Team. 2014 A language and environment for statistical computing. R
 966 Foundation for Statistical Computing, Vienna, Austria. URL: [http://www.R-](http://www.R-project.org/)
 967 [project.org/](http://www.R-project.org/)

968 Roth, T.C., LaDage, L. D., Freas, C. A., Pravosudov, V. V. (2012). Variation in memory
 969 and the hippocampus across populations from different climates: a common
 970 garden approach. *Proceedings of the Royal Society B*, 279, 402-410.

971 Roth, T. C., Pravosudov, V. V. (2009). Hippocampal volume and neuron numbers
 972 increase along a gradient of environmental harshness – a large-scale
 973 comparison. *Proceedings of the Royal Society B*, 276, 401-405.

974 Roth, T. C., LaDage, L. D., Freas, C. A., Pravosudov, V.V. (2011). Variation in memory
 975 and the hippocampus across populations from different climates: a common
 976 garden approach. *Proceedings of the Royal Society B*, 279, 402-410.

977 Sherry, D. F., Vaccarino, A. L., Buckenham, K., Herz, R. S. (1989). The hippocampal
 978 complex of food-storing birds. *Brain, Behavior and Evolution*, 34, 308-317.

979 Shettleworth, S. J. (2010). *Cognition, evolution, and behavior*, 2nd edition. Oxford
 980 University Press, New York.

981 Sulikowski, D., Burke, D. (2012). Win shifting in nectarivorous birds: selective
 982 inhibition of the learned win-stay response. *Animal Behaviour*, 83, 519-524.

983 Strang, C. G., Sherry, D. F. (2014). Serial reversal learning in bumblebees (*Bombus*
 984 *impatiens*). *Animal Cognition*, 17, 723-734.

985 Urhan, A. U., Brodin, A. (2015). No evidence for memory interference across sessions
 986 in food hoarding tits *Poecile palustris* under laboratory conditions. *Animal*
 987 *Cognition*, 18, 649-656.

988 Waite, T. A., Reeve, J. D. (1993). Food storage in gray jays: source type and cache
 989 dispersion. *Ethology*, 93: 326-336.

990 Walsh, C. M., Booth, V., Poe, G. R. (2011). Spatial and reversal learning in the Morris
 991 water maze are largely resistant to six hours of REM sleep deprivation
 992 following training. *Learning and Memory*, 18, 422-434.

993 Wright, T. F., Eberhard, J. R., Hobson, E. A., Avery, M. L., Russello, M. A. (2010).
 994 Behavioral flexibility and species invasions: the adaptive flexibility
 995 hypothesis. *Ethology, Ecology and Evolution*, 22, 393-404.

996

ACKNOWLEDGEMENTS

This study was supported by the National Science Foundation, grant IOS-1351295, to VVP. Initial development of the low-cost RFID units was assisted by funding from the National Science Foundation (awards IOS-0919528 and DBI-1458402 to ESB). Critical and constructive criticisms from Anders Brodin, an anonymous reviewer and the Editor significantly improved the manuscript.

Figure 1 rev

Figure 2 rev

Figure 3a rev

Figure 3b rev

Figure 4 rev

Figure 5a rev

Figure 5b rev

Figure 6 rev