WDR5 positively regulates p53 stability by inhibiting p53 ubiquitination

Qingqing Xie¹, Zengpeng, Li¹, Jianming Chen²

1 School of Life Sciences, Xiamen University, Xiamen 361102, Fujian, China

2 State Key Laboratory Breeding Base of Marine Genetic Resources, Third Institute of Oceanography, State Oceanic Administration, Xiamen 361005, Fujian, China

Corresponding authors:

Jianming Chen: Third Institute of Oceanography, State Oceanic Administration, Xiamen 361005, Fujian, China. Tel/Fax: +86592-2195337, E-mail: chenjm@xmu.edu.cn

Abstract:

WD40 repeat protein WDR5 is a core component of the Set/MLL histone methyltransferase complex which catalyzes histone H3 Lys4 trimethylation and activates gene transcription in human cells. WDR5 promotes Set/MLL complex assembly and mediates the complex binding to Lys4-dimethylated histone H3 tail. Most earlier studies report that WDR5 exerts profound effects on various cellular and organismal processes mainly through epigenetic regulation of gene transcription. However, the functions of WDR5 in lung cancer remain largely unknown. Here, we report that WDR5 positively regulates p53 stability by inhibiting p53 ubiquitination in human lung cancer A549 cells. Overexpression of WDR5 dramatically increases p53 protein levels and its half-life in A549 cells, while depletion of WDR5 with WDR5-specific siRNAs significantly decreases p53 protein levels. We also observe that WDR5 is required for p53 induction in response to cisplatin treatment. Mechanistically, WDR5 colocalizes with p53 and inhibits p53 ubiquitination, resulting in p53 stabilization. Consequently, overexpression of WDR5 induces G1 phase arrest in A549 cells, and knocking down WDR5 by siRNAs reduces the population at G1 phase. Furthermore, p53 expression levels is at least in part determined by the p53 positive regulator WDR5 in some cancer cells. Taken together, these data suggest that WDR5 is directly involved in p53 signaling pathway. Our studies provide a new insight into WDR5 functions in A549 cells.

Keywords: WDR5; p53; ubiquitination; A549 cells

Introduction

WD40 repeat protein WDR5 is a core component of Set/MLL complex which catalyzes histone H3 Lys4 trimethylation (H3K4me3) in human cells, along with other common subunits Ash2L, RbBP5, and DPY30 [1,2]. WDR5 specifically interacts with the K4-methylated histone H3 to mediate K4 trimethylation and is required for global H3K4me3 [2,3]. As a key component of Set/MLL complex, WDR5 binds an arginine-bearing motif in MLL, promoting the complex assembly [1]. Disrupting the WDR5-MLL interaction by inhibitors or mutating the MLL WDR5 interaction motif severely inhibits the histone methyltransferase activity [4,5,6,7]. WDR5 represents a potential coactivator of transcription factors by associating transcription factors and promoting histone methylation, resulting in transcription activation [8,9,10,11]. In response to p53 induction, WDR5 along with Rbbp5 and Ash2L increasingly occupies the promoters of all the p53 target genes, promoting p53 target genes expression [8]. WDR5 is found to associate with OCT4 and activates transcription of key self-renewal genes, leading to embryonic stem cell self-renewal [9]. WDR5 is also reported to interact with MYC and is needed for MYC recruitment to chromatin to drive tumorigenesis [12,13,14]. WDR5 binds Cbx8 and together regulate Notch gene expression by maintaining H3K4me3 in breast cancer cells [10]. In addition, post-transcriptional modifications are involved in the

regulation of WDR5. PAS domain containing protein kinase (Pask) phosphorylates WDR5 at ser49, facilitating the transition to the trimethylated state of the Myog promoter, leading to transcription activation and differentiation [11].

Recent study reveals that WDR5 cooperates with other chromatin modifiers, such as HDAC3, to coordinately regulate gene transcription. WDR5 is induced in response to hypoxia and cooperates with HDAC3 to activate mesenchymal gene expression [15]. H3K4me3 is also involved in DNA replication and transcription processing [16,17]. WDR5 localizes at replication origins and is directly involved in DNA replication by regulating H3 K4 methylation [16]. WDR5 along with Ash2L and RbBP5 are also involved in pre-mRNA processing of immediate early genes without affecting the levels of H3K4me3 at the promoter region [17]. More recently, functions of WDR5 independent of the Set/MLL complex and H3K4me3 are reported. WDR5 is recruited to the midbody and promotes abscission [18]. In *C.elegans*, WDR5 homologs are required for oogenesis by promoting TRA-1-mediated transcription repression, independent of histone methylation [19]. In addition, WDR5 is essential for assembling distinct complexes, including NSL complex and VISA complex [20,21]. WDR5 interacts with NSL1 and promotes NSL complex assembling, which contains histone acetyltransferase MOF [20]. In a virus infection dependent manner, WDR5 translocates to the mitochondria and is essential for the VISA complex assembly, leading to IRF3 and NFκB activation [21].

In this study, we report that WDR5 regulates p53 stability in A549 cells. We show that WDR5 colocalizes with p53 and inhibits p53 ubiquitination, leading to p53 stabilization.

WDR5 is required for p53 induction in response to cisplatin treatments and regulates G1 phase arrest in A549 cells. Interestingly, we found that the expression of WDR5 is positively correlated with p53 expression in some cancer cell lines, indicating that the levels of p53 in cancer cells is at least in part regulated by WDR5.

Materials and Methods

Plasmids

Full-length cDNA encoding human WDR5 was amplified by PCR using cDNA generated from HEK293 cells. Then WDR5 cDNA was subcloned into the pCMV-myc vector for transfection or pLVX-IRES-ZsGreen 1 vector (Clontech) for lentivirus infection. The plasmid constructions were verified by sequencing (Beijing Genomics Institute). Primers are as follows: WDR5-forward: ATGGCGACGGAGGAGAAG; WDR5-reverse: TTAGCAGTCACTCTTCCACAGTT.

Cell culture, transfection, and reagents

HEK293 and A549 cells were cultured in Dulbecco's modified eagle medium (Gibco), supplemented with 10% fetal bovine serum (FBS) (Gibco) and 1% penicillin–streptomycin (100 U/ml penicillin and 100 mg/ml streptomycin) at 37°C in a humidified incubator containing 5% CO₂. Transient transfections were carried out with Lipofectamine 3000 (Thermo Fisher Scientific) according to the manufacture's instruction. Cisplatin, cycloheximide, MG132 and puromycin were purchased from Sigma-Aldrich.

Western blotting analysis

Cells were harvested and lysed in a lysis buffer (150 mM NaCl, 20mM Tris-HCl, pH 7.4, 1 mM EDTA, 1% Triton, 1 mM β -glycerolphosphate, 1 mM sodium orthovanadate, 1 mg/ml leupeptin, 1 mM phenylmethylsulfonylfluoride) for 45 minutes on ice. Equal amounts of protein lysates were boiled and separated on 12% SDS-polyacrylamide gels. The samples were transferred to Immobilon PVDF membranes (Millipore) and blocked with 5% skim milk for 1 h. Then the membranes were probed with primary antibodies: anti-p53 (DO-1) and anti-myc (9E10) monoclonal antibody (Santa Cruze); anti- β -actin monoclonal antibody (Sigma-Aldrich); anti-WDR5 (ab56919) monoclonal antibody and anti-bax (E63) (ab32503) polyclonal antibody (abcam). After extensive washing, blots were incubated with horseradish peroxidase-conjugated goat anti-mouse and anti-rabbit antibody (Amersham Pharmacia Biotech) for 1 h, and then the chemiluminescence reaction was performed.

Quantitative real-time RT-PCR

The simple Total RNA Kit (Tiangen Biotech, Beijing, China) was used to extract total RNA from A549 cells and revers transcription was performed using First Strand cDNA Synthesis kit (Toyobo, Osaka, Japan). The cDNA samples were then analyzed by quantitative real-time PCR with SYBR Green PCR mix (Tiangen Biotech, Beijing, China). Primers are as follows, p53 forward: GCTTTCCACGACGGTGAC; p53 reverse: GCTCGACGCTAGGATCTGAC; p21 forward: GCAGACCAGCATGACAGATTTC; p21 reverse: GCGGATTAGGGCTTCCTCTT; puma forward: GTGCCCTCGGCAGTGTCC;

puma reverse: GTACTGTGCGTTGAGGTC; bax forward: CACCAGCTCTGAGCAGATC;
bax reverse: GCTGCCACTCGGAAAAAG; mdm2 forward: GTGCTGTAACCACTCAC;
mdm2 reverse: GCTCTTTCACAGAGAAGC; WDR5 forward:
TCTCCCCGAACGGCAAATAC; WDR5 reverse: TGTGGCCAGTGACGTCTTC.

Cell cycle analysis

A549-control and A549-WDR5 cells were treated with 20 μ M cisplatin for 24 hours, and then cells were harvested and fixed in ice-cold 70% ethanol at 4°C overnight. After being washed with PBS, cells were incubated with 200 μ g/ml DNase-free RNase A at 37°C for 20 minutes, and then incubated with 50 μ g/ml PI for another 30 minutes in the dark. And finally, 2×10^4 cells were counted and analyzed each sample by a Flow Cytometer (BD C6).

Immunofluorescence

A549 cells were plated on glass coverslips and transfected with pCMV-myc-WDR5. 24 hours after transfection, cells were fixed in 4% paraformaldehyde and permeabilized in 0.2% Triton X-100. And then cells were stained in a standard protocol. Images were acquired with Zeiss LSM 800 laser-scanning confocal microscope and ZEN software.

RNA interference

The nucleotide sequences for siRNAs to human WDR5 are as follows, siWDR5-1:
UCUGGAACCUUCAGACGAATT; siWDR5-2: GGUCACAAGCUGGGAAUAUTT;

RNAiMax (Invitrogen) was used to transfect A549 cells with siRNAs according to the manufacture's instruction.

Ubiquitination of p53 in vivo

HEK293 cells were transfected with expression vectors of HA-ub and myc-WDR5, as indicated. 24 hours later, cells were treated with 10 μ M MG132 for 3 hours before being collected. Then cells were lysed with RIPA buffer (0.5% Nonidet P-40, 0.2% SDS, 0.5% sodium deoxycolate, 1 mM PMSF, 2 μ g/ml aprotinin, 10 μ g/ml leupeptin). Ubiquitinated p53 was immunoprecipitated with anti p53 antibody followed by Western blot analysis with anti-HA antibody (against HA-ubiquitin).

Chip assay

A total of 1×10^7 cells were taken for chip assay using SimpleChIP Plus Enzymatic Chromatin IP kit (Magnetic Beads) (Cell Signaling Technology) according to the manufacture's instruction. Cells were cross-linked in 1% formaldehyde in DMEM medium for 10 min and then 0.215 M Glycine was added for 5 min to stop the cross-linking. Then nuclei were prepared and digested by 1 μ l micrococcal nuclease for 20 min at 37°C. Nuclear pellet was then sonicated for 3 min on ice. 10 μ g of digested and cross-linked chromatin were incubated with H3K4me3 (ab1012) (abcam) antibody or rabbit IgG overnight at 37°C, then 30 μ l protein G magnetic beads were added to IP reaction and incubated for 2 hours. Then magnetic beads were washed with low salt buffer 3 times and high salt buffer once. Chromatin were eluted at 65°C for 30 min and reversal of cross-links were performed in NaCl and Protein K at 65°C overnight. Purified DNA were subjected to

a real-time quantitative PCR analysis. Primers are as follows, bax promoter, forward: TAATCCCAGCGCTTTGGAAG, reverse: TGCAGAGACCTGGATCTAGCAA; mdm2 promoter, forward: GGTTGACTCAGCTTTTCCTCTTG, reverse: GGAAAATGCATGGTTTAAATAGCC.

Results

WDR5 regulates p53 stability

By serendipity, we observed that silencing WDR5 with siRNAs in human lung A549 cells affects cell proliferation rate. To determine whether WDR5 could regulate p53, the key regulator of cell proliferation, we transfected increasing amounts of WDR5 into A549 cells. As figure 1A shows, overexpression of WDR5 led to a dose-dependent increase in endogenous p53 levels. Then we infected A549 cells with retroviruses encoding WDR5. Again, the endogenous p53 level was increased upon WDR5 expression with no effect on p53 mRNA levels (Figure 1BC). We also observed increased mRNA levels of p53 target genes, p21, bax, mdm2 and puma (Figure 1C). Since WDR5 is a core component of Set/MLL complex which catalyzes H3K4me3 and activates gene transcription in human cells. We examined whether WDR5 overexpression would activate H3K4me3 on the promoters of p53 target genes bax and mdm2. However, as Figure 1D shows, overexpression of WDR5 was not sufficient to promote H3K4me3 on bax and mdm2 promoters. These data suggest that WDR5 stabilizes p53 and promotes p53 target genes expression.

WDR5 is required for p53 induction in response to cisplatin treatment

Then we determined whether WDR5 affected p53 accumulation in response to DNA

damage stresses, such as cisplatin treatment. As shown in Figure 2A, p53 induction in A549 cells stably expression WDR5 was increased than that of A549 control cells in response to cisplatin treatment. To confirm the role of WDR5 in regulating p53 levels, we depleted WDR5 with WDR5-specific siRNAs, and we found that downregulation of WDR5 decreased p53 levels (Fig 2B). In agreement with the results of WDR5 overexpression assays, downregulation of WDR5 attenuated p53 accumulation in response to cisplatin treatments. These results indicate that WDR5 is required for p53 induction in response to cisplatin treatment.

WDR5 colocalizes with p53 and inhibits p53 ubiquitination

To further confirm that WDR5 regulated p53 stability, we performed half-life assay by treating A549 control cells or cells stably expressing WDR5 with cycloheximide (CHX). As Figure 3A shows, half-life of p53 was ~20 minutes in control A549 cells. However, overexpression of WDR5 prolonged p53 half-life to ~40 minutes. The stability of p53 is tightly controlled by ubiquitination and proteasomal-dependent degradation through E3 ubiquitin ligase, such as mdm2 and ARF-BP1/Mule. It is likely that WDR5 functions to inhibit the ubiquitination of p53, resulting in the stabilization of p53. Indeed, as shown in Figure 3B, overexpression of HA-ub significantly induced the ubiquitination of p53, while co-expression of WDR5 effectively decreased the ubiquitination of p53, indicating that WDR5 negatively regulates p53 ubiquitination in cells. Furthermore, we found that WDR5 and p53 colocalized very well in the nucleus and cytoplasm (Figure 3C). These results suggest that WDR5 colocalizes with p53 and inhibits p53 ubiquitination.

WDR5 induces G1 cell cycle arrest in A549 cells

As a tumor suppressor, p53 inhibits cell growth by inducing cell cycle arrest, apoptosis and senescence. Since our data showed WDR5 affects p53's stability, it's possible that WDR5 regulates p53-dependent functions, including cell cycle regulation. As shown in Figure 4A, overexpression of WDR5 induced G1 phase accumulation in A549 cells. In contrast, inactivation of WDR5 by WDR5 RNAi reduced the cell population at G1 phase (Figure 4B). WDR5 induced G1 cell cycle arrest in A549 cells probably through the induction of p21 (Fig 1C). p53 is a crucial tumor suppressor which is frequently mutated or dysregulated in human cancers. We hypothesized that WDR5 may correlate with p53 dysregulation in cancers. To test the hypothesis, we tested the expression of WDR5 and p53 in several cancer cell lines. Indeed, we found that the expression levels of WDR5 is dramatically decreased in p53 wild-type A549 and H460 lung cancer cells with lower p53 expression, when comparing with p53-mutant cell lines, including DU145 prostate cancer cells and AsPC-1 pancreatic cancer cells with higher p53 expression levels (Fig 4C). Interestingly, the protein levels of WDR5 is also lower in PC3 prostate cancer cells with p53 not detected (Fig 4C). These results suggest that p53 protein levels in cancer cells is at least in part regulated by WDR5 and the function of WDR5 may be highly cell-context dependent.

Discussion

As a core component of Set/MLL complex, WDR5 mainly functions to mediate histone H3 K4 trimethylation in human cells. WDR5 represents a potential transcription coactivator

through mediating H3K4me3, promoting gene transcription [9,10,11,13,14]. In addition, H3K4me3 not only participates in gene transcription regulation, recent studies show that WDR5 and Set/MLL complex are also directly involved in DNA replication, chromosomal polyploidy and pre-mRNA processing [16,17]. Furthermore, WDR5 serves as a binding platform for assembling distinct complexes, including MOF-containing NSL complex and virus-induced VISA complex [20,21].

In this work, we provide strong evidence that WDR5 positively regulates p53 stability. We show that overexpression of WDR5 increases p53 protein levels without influencing the p53 mRNA levels. We also observe that overexpression of WDR5 prolongs the half-life of p53. Moreover, WDR5 is required for p53 induction in response to DNA damage stress caused by cisplatin treatment. In agreement with overexpression results, depletion of WDR5 with siRNAs decreases p53 protein levels and p53 induction in response to cisplatin treatment. Mechanistically, we show that WDR5 colocalizes with p53 and inhibits p53 ubiquitination. Overexpression of WDR5 causes G1 phase accumulation in A549 cells, which is possibly caused by the increasing of p21 levels, and depletion of WDR5 releases the G1 phase arrest.

It is reported that WDR5 occupancy is enhanced at all the p53 target promoters in response to p53 stabilization [8]. However, we demonstrate that overexpression of WDR5 alone was not enough to promote H3K4me3 at p53 target genes, suggesting that WDR5 increases p53 stability without influencing the H3K4me3 status of promoters of p53 target genes. WDR5 strongly inhibits p53 ubiquitination, however, the underlying mechanism through which WDR5 inhibits p53 ubiquitination is unknown. Given that WDR5

significantly colocalizes with p53 and WD40 propeller mediates protein-protein interactions, it is possible that WDR5 inhibits p53 ubiquitination through binding to and inhibiting the activities of E3 ubiquitin ligases toward p53, such as mdm2. Further studies are needed to test the hypothesis in A549 cells.

Many early studies have shown that WDR5 is overexpressed in various cancers, including bladder cancer, prostate cancer, breast cancer and leukemia, promoting cell proliferation and self-renewal by mediating H3K4me3 [22,23,24,25]. In addition, WDR5 forms a complex with N-Myc at the mdm2 promoter, leading to mdm2 transcription activation and p53 protein decreasing in neuroblastoma cells [26]. Early studies strongly suggest that WDR5 is a potential oncogene. However, the function of WDR5 in lung cancer remains largely unknown. In this study, we demonstrate that WDR5 increases p53 stability by inhibiting p53 ubiquitination in human lung cancer A549 cells. One possibility is that the functions of WDR5 in cancers are more complicated than earlier imagined, and WDR5 may have different functions in different cancers. Indeed, we provide evidence that the expression of WDR5 is significantly different in various cancers (Fig. 4C). WDR5 is highly expressed in AsPC-1 pancreatic cancer cells and DU145 human prostate cancer cells. While the expression of WDR5 is lower in A549, H460 lung cancer cells, and PC3 prostate cancer cells (Fig. 4C). Interestingly, the expression of p53 is also higher in AsPC-1 and DU145 cells comparing with A549 and H460 cells. It is possible that extremely low expression of p53 in A549 and H460 cells is at least in part caused by low expression of p53 positive regulator WDR5, higher expression of mutant p53 is connected with higher expression of WDR5 in p53-mutant AsPC-1 and DU145 cells. Importantly, we observed

that the cell lines used in earlier researches on WDR5 functions are mostly p53-mutant, including UM-UC-3, T24, THP-1 and U937 cells [22,24]. Overexpression of WDR5 in these cells may result in the stabilization of mutant p53, leading to cell proliferation. Thus, functions of WDR5 in cancer cells appears to be highly cell-context dependent. WDR5 is lowly expressed in p53 wild-type A549 cells, leading to p53 ubiquitination and degradation, and consequently promoting cancer cells proliferation. While WDR5 is overexpressed in some cancer cells with mutant p53, leading to mutant p53 stabilization and cancer cell proliferation.

In conclusion, we found that WDR5 up-regulates p53 protein by inhibiting p53 ubiquitination in A549 cells. Our work reveals a novel role of WDR5 in p53 signaling pathway and suggests a molecular mechanism of p53 induction induced by anticancer drug cisplatin, which may lead to the development of new strategies for lung cancer therapy.

Acknowledgements

This work was supported by National Science Foundation grant 31271567, China Postdoctoral Science Foundation 2013M541853 and 2014T70606 and The Scientific Research Foundation of Third Institute of Oceanography, SOA 2014007. This work was also supported by National Nature Science Foundation of China (41606160) and The Scientific Research Foundation of Third Institute of Oceanography, SOA (No. 2015011).

References

- [1] R.C. Trievel, A. Shilatifard, WDR5, a complexed protein, *Nat Struct Mol Biol* 16 (2009) 678-680.
- [2] J. Wysocka, T. Swigut, T.A. Milne, Y. Dou, X. Zhang, A.L. Burlingame, R.G. Roeder, A.H. Brivanlou, C.D. Allis, WDR5 associates with histone H3 methylated at K4 and is essential for H3 K4 methylation and vertebrate development, *Cell* 121 (2005) 859-872.
- [3] J.F. Couture, E. Collazo, R.C. Trievel, Molecular recognition of histone H3 by the WD40 protein WDR5, *Nat Struct Mol Biol* 13 (2006) 698-703.
- [4] N.L. Alicea-Velazquez, S.A. Shinsky, D.M. Loh, J.H. Lee, D.G. Skalnik, M.S. Cosgrove, Targeted Disruption of the Interaction between WD-40 Repeat Protein 5 (WDR5) and Mixed Lineage Leukemia (MLL)/SET1 Family Proteins Specifically Inhibits MLL1 and SETd1A Methyltransferase Complexes, *J Biol Chem* 291 (2016) 22357-22372.
- [5] D.D. Li, Z.H. Wang, W.L. Chen, Y.Y. Xie, Q.D. You, X.K. Guo, Structure-based design of ester compounds to inhibit MLL complex catalytic activity by targeting mixed lineage leukemia 1 (MLL1)-WDR5 interaction, *Bioorg Med Chem* 24 (2016) 6109-6118.
- [6] F. Grebien, M. Vedadi, M. Getlik, R. Giambruno, A. Grover, R. Avellino, A. Skucha, S. Vittori, E. Kuznetsova, D. Smil, D. Barsyte-Lovejoy, F. Li, G. Poda, M. Schapira, H. Wu, A. Dong, G. Senisterra, A. Stukalov, K.V. Huber, A. Schonegger, R. Marcellus, M. Bilban, C. Bock, P.J. Brown, J. Zuber, K.L. Bennett, R. Al-Awar, R. Delwel, C. Nerlov, C.H. Arrowsmith, G. Superti-Furga, Pharmacological targeting of the Wdr5-MLL interaction in C/EBPalpha N-terminal leukemia, *Nat Chem Biol* 11 (2015) 571-578.
- [7] D.D. Li, W.L. Chen, Z.H. Wang, Y.Y. Xie, X.L. Xu, Z.Y. Jiang, X.J. Zhang, Q.D. You, X.K.

- Guo, High-affinity small molecular blockers of mixed lineage leukemia 1 (MLL1)-WDR5 interaction inhibit MLL1 complex H3K4 methyltransferase activity, *Eur J Med Chem* 124 (2016) 480-489.
- [8] S.K. Mungamuri, S. Wang, J.J. Manfredi, W. Gu, S.A. Aaronson, Ash2L enables P53-dependent apoptosis by favoring stable transcription pre-initiation complex formation on its pro-apoptotic target promoters, *Oncogene* 34 (2015) 2461-2470.
- [9] Y.S. Ang, S.Y. Tsai, D.F. Lee, J. Monk, J. Su, K. Ratnakumar, J. Ding, Y. Ge, H. Darr, B. Chang, J. Wang, M. Rendl, E. Bernstein, C. Schaniel, I.R. Lemischka, Wdr5 mediates self-renewal and reprogramming via the embryonic stem cell core transcriptional network, *Cell* 145 (2011) 183-197.
- [10] C.Y. Chung, Z. Sun, G. Mullokandov, A. Bosch, Z.A. Qadeer, E. Cihan, Z. Rapp, R. Parsons, J.A. Aguirre-Ghiso, E.F. Farias, B.D. Brown, A. Gaspar-Maia, E. Bernstein, Cbx8 Acts Non-canonically with Wdr5 to Promote Mammary Tumorigenesis, *Cell Rep* 16 (2016) 472-486.
- [11] C.K. Kikani, X. Wu, L. Paul, H. Sabic, Z. Shen, A. Shakya, A. Keefe, C. Villanueva, G. Kardon, B. Graves, D. Tantin, J. Rutter, Pask integrates hormonal signaling with histone modification via Wdr5 phosphorylation to drive myogenesis, *Elife* 5 (2016).
- [12] L.R. Thomas, Q. Wang, B.C. Grieb, J. Phan, A.M. Foshage, Q. Sun, E.T. Olejniczak, T. Clark, S. Dey, S. Lorey, B. Alicie, G.C. Howard, B. Cawthon, K.C. Ess, C.M. Eischen, Z. Zhao, S.W. Fesik, W.P. Tansey, Interaction with WDR5 promotes target gene recognition and tumorigenesis by MYC, *Mol Cell* 58 (2015) 440-452.
- [13] A. Carugo, G. Genovese, S. Seth, L. Nezi, J.L. Rose, D. Bossi, A. Cicalese, P.K. Shah, A.

Viale, P.F. Pettazzoni, K.C. Akdemir, C.A. Bristow, F.S. Robinson, J. Tepper, N. Sanchez, S. Gupta, M.R. Estecio, V. Giuliani, G.I. Dellino, L. Riva, W. Yao, M.E. Di Francesco, T. Green, C. D'Alesio, D. Corti, Y. Kang, P. Jones, H. Wang, J.B. Fleming, A. Maitra, P.G. Pelicci, L. Chin, R.A. DePinho, L. Lanfrancone, T.P. Heffernan, G.F. Draetta, In Vivo Functional Platform Targeting Patient-Derived Xenografts Identifies WDR5-Myc Association as a Critical Determinant of Pancreatic Cancer, *Cell Rep* 16 (2016) 133-147.

[14] L.R. Thomas, A.M. Foshage, A.M. Weissmiller, W.P. Tansey, The MYC-WDR5 Nexus and Cancer, *Cancer Res* 75 (2015) 4012-4015.

[15] M.Z. Wu, Y.P. Tsai, M.H. Yang, C.H. Huang, S.Y. Chang, C.C. Chang, S.C. Teng, K.J. Wu, Interplay between HDAC3 and WDR5 is essential for hypoxia-induced epithelial-mesenchymal transition, *Mol Cell* 43 (2011) 811-822.

[16] F. Lu, X. Wu, F. Yin, C. Chia-Fang Lee, M. Yu, I.S. Mihaylov, J. Yu, H. Sun, H. Zhang, Regulation of DNA replication and chromosomal polyploidy by the MLL-WDR5-RBBP5 methyltransferases, *Biol Open* 5 (2016) 1449-1460.

[17] P.L. Teoh, A.D. Sharrocks, WDR5, ASH2L, and RBBP5 control the efficiency of FOS transcript processing, *Cell Mol Biol Lett* 19 (2014) 215-232.

[18] J.K. Bailey, A.T. Fields, K. Cheng, A. Lee, E. Wagenaar, R. Lagrois, B. Schmidt, B. Xia, D. Ma, WD repeat-containing protein 5 (WDR5) localizes to the midbody and regulates abscission, *J Biol Chem* 290 (2015) 22447.

[19] T. Li, W.G. Kelly, A role for WDR5 in TRA-1/Gli mediated transcriptional control of the sperm/oocyte switch in *C. elegans*, *Nucleic Acids Res* 42 (2014) 5567-5581.

- [20] J. Dias, N. Van Nguyen, P. Georgiev, A. Gaub, J. Brettschneider, S. Cusack, J. Kadlec, A. Akhtar, Structural analysis of the KANSL1/WDR5/KANSL2 complex reveals that WDR5 is required for efficient assembly and chromatin targeting of the NSL complex, *Genes Dev* 28 (2014) 929-942.
- [21] Y.Y. Wang, L.J. Liu, B. Zhong, T.T. Liu, Y. Li, Y. Yang, Y. Ran, S. Li, P. Tien, H.B. Shu, WDR5 is essential for assembly of the VISA-associated signaling complex and virus-triggered IRF3 and NF-kappaB activation, *Proc Natl Acad Sci U S A* 107 (2010) 815-820.
- [22] X. Chen, W. Xie, P. Gu, Q. Cai, B. Wang, Y. Xie, W. Dong, W. He, G. Zhong, T. Lin, J. Huang, Upregulated WDR5 promotes proliferation, self-renewal and chemoresistance in bladder cancer via mediating H3K4 trimethylation, *Sci Rep* 5 (2015) 8293.
- [23] X. Dai, W. Guo, C. Zhan, X. Liu, Z. Bai, Y. Yang, WDR5 Expression Is Prognostic of Breast Cancer Outcome, *PLoS One* 10 (2015) e0124964.
- [24] Z. Ge, E.J. Song, Y.I. Kawasaki, J. Li, S. Dovat, C. Song, WDR5 high expression and its effect on tumorigenesis in leukemia, *Oncotarget* 7 (2016) 37740-37754.
- [25] J.Y. Kim, T. Banerjee, A. Vinckevicius, Q. Luo, J.B. Parker, M.R. Baker, I. Radhakrishnan, J.J. Wei, G.D. Barish, D. Chakravarti, A role for WDR5 in integrating threonine 11 phosphorylation to lysine 4 methylation on histone H3 during androgen signaling and in prostate cancer, *Mol Cell* 54 (2014) 613-625.
- [26] Y. Sun, J.L. Bell, D. Carter, S. Gherardi, R.C. Poulos, G. Milazzo, J.W. Wong, R. Al-Awar, A.E. Tee, P.Y. Liu, B. Liu, B. Atmadibrata, M. Wong, T. Trahair, Q. Zhao, J.M. Shohet, Y. Haupt, J.H. Schulte, P.J. Brown, C.H. Arrowsmith, M. Vedadi, K.L. MacKenzie, S.

Huttelmaier, G. Perini, G.M. Marshall, A. Braithwaite, T. Liu, WDR5 Supports an N-Myc Transcriptional Complex That Drives a Protumorigenic Gene Expression Signature in Neuroblastoma, *Cancer Res* 75 (2015) 5143-5154.

Figure legends

Figure 1. WDR5 stabilizes p53. (A) A549 cells were transfected with increasing amounts of WDR5. 24 hours later, proteins were extracted and subjected to Western blot. (B) A549 cells infected with lentivirus encoding WDR5 were lysed and lysates were blotted with the indicated antibodies. (C) mRNA extracted from A549 control and A549-WDR5 cells were subjected to real-time quantitative analysis. (D) Chip analysis showing H3K4me3 occupancy on bax and mdm2 promoters in A549-control and A549-WDR5 cells. IP, immunoprecipitation.

Figure 2. WDR5 is required for p53 induction in response to cisplatin treatments. (A) A549-control and A549-WDR5 cells were treated with 20 μ M cisplatin in the indicated times and cells were lysed and the lysates were subjected to western blot analysis with the indicated antibodies. (B) A549 cells were transfected with non-specific control siRNAs or WDR5-specific siRNAs, 36 hours later, proteins were extracted and subjected to western blot analysis with indicated antibodies. (C) A549 cells were transfected with siRNAs, 36 hours later, cells were treated with 20 μ M cisplatin for the indicated times and then, cells were lysed and subjected to western blot analysis with the indicated antibodies.

Figure 3. WDR5 colocalizes with p53 and inhibits p53 ubiquitination. (A) A549-control

and A549-WDR5 cells were treated with 50 µg/ml cycloheximide for the indicated times. Cells were harvested and subjected to Western blot analysis with the indicated antibodies. And quantification of the p53 protein levels relative to β-actin was shown. (B) WDR5 inhibits p53 ubiquitination. HEK293 cells were transfected with the indicated plasmids. 24 hours after transfection, cells were harvested and lysed. The lysates were immunoprecipitated with anti-p53 (DO-1) antibody and immunoblotted with anti-HA antibody. (C) Subcellular localization of WDR5 and p53. HEK293 cells were transfected with myc-WDR5, 24 hours later, cells were fixed and stained as indicated. The nuclei were stained with DAPI.

Figure 4. WDR5 affects cell cycle distribution in A549 cells. (A) A549-control and A549-WDR5 cells were harvested and stained with propidium iodide followed by FACS analysis. (B) A549 cells were transfected with WDR5-specific siRNAs, 36 hours later, cells were collected and cell cycle distribution was determined by FACS. (C) Western blot analysis of WDR5 and p53 protein expression in two p53 wild type cell lines (A549 and H460) and three p53 mutant cell lines (AsPC-1, DU145 and PC3).

Figures

Figure 1

Figure 2

Figure 3

Figure 4

graphical abstract