1 For the *Perspectives* section of *Biological Conservation*

2 **When natural habitat fails to enhance biological** 3 **pest control – five hypotheses**

4

5 Teja Tschamntke¹ (ttschar@gwdg.de), Daniel S. Karp² (dkarp@ucdavis.edu),
6 Rebecca Chaplin-Kramer³ (bchaplin@stanford.edu), Péter Batáry¹
7 (p.batary@gwdg.de), Fabrice DeClerck⁴ (f.declerck@cgiar.org), Claudio Gratton⁵
8 (cgratton@wisc.edu), Lauren Hunt⁶ (lhunt1@ssync.org), Anthony Ives⁷
9 (arives@wisc.edu), Mattias Jonsson⁸ (mattias.jonsson@slu.se), Ashley Larsen⁹
10 (aelarsen@berkeley.edu), Emily A. Martin¹⁰ (emily.martin@uni-wuerzburg.de),
11 Alejandra Martínez-Salinas¹¹ (amartinez@catie.ac.cr), Timothy D. Meehan¹²
12 (tmeehan@neoninc.org), Megan O'Rourke¹³ (megorust@vt.edu), Katja Poveda¹⁴
13 (kap235@cornell.edu), Jay A. Rosenheim¹⁵ (JARosenheim@ucdavis.edu), Adrien
14 Rusch¹⁶ (adrien.rusch@bordeaux.inra.fr), Nancy Schellhorn¹⁷
15 (Nancy.Schellhorn@csiro.au), Thomas C. Wanger^{1,8} (tomcwanger@gmail.com),
16 Stephen Wratten¹⁸ (Steve.Wratten@lincoln.ac.nz), Wei Zhang¹⁹ (w.zhang@cgiar.org)

17

18 ¹ Agroecology, Department of Crop Sciences, University of Göttingen,
19 Grisebachstrasse 6, D-37077 Göttingen, Germany

20 ² Institute for Resources, Environment, and Sustainability; University of British
21 Columbia, Vancouver, British Columbia, Canada

22 ³ Natural Capital Project, Woods Institute for the Environment, Stanford University,
23 USA

24 ⁴ Biodiversity International, CGIAR, Landscape Restoration and Management,
25 Montpellier, France

⁵ University of Wisconsin, Department of Entomology, Madison, WI (USA)

⁶ Department of Entomology, University of Maryland, 4112 Plant Sciences Building,
College Park, MD 20742-4454, USA

⁷ University of Wisconsin, Department of Zoology, Madison, WI (USA)

⁸ Swedish University of Agricultural Sciences, Department of Ecology, PO Box 7044,
SE-750 07 Uppsala, Sweden

⁹ Environmental Science, Policy, & Management, University of California, Berkeley,
94720

¹⁰ Department of Animal Ecology and Tropical Biology, Biocenter Am Hubland,
University of Würzburg, 97074 Würzburg, Germany

¹¹ Tropical Agricultural Research and Higher Education Center (CATIE), Turrialba,
Costa Rica; Department of Fish and Wildlife Sciences, University of Idaho, Idaho,
USA

¹² National Ecological Observatory Network, Boulder, Colorado, USA

¹³ Virginia Tech, Department of Horticulture, Blacksburg, VA (USA)

¹⁴ Cornell University, Department of Entomology, Ithaca, NY (USA)

¹⁵ Department of Entomology and Nematology, University of California, Davis, CA
(USA)

¹⁶ INRA, ISVV, Univ. Bordeaux, Bordeaux Sciences Agro, UMR SAVE, F-33883
Villenave d'Ornon

¹⁷ CSIRO, Brisbane, QLD, 4001, Australia

¹⁸ Bio-Protection Research Centre, Burns, 514, Cnr Springs and Ellesmere Junction
Roads, P O Box 85084, Lincoln University, Lincoln 7647, New Zealand

¹⁹ International Food Policy Research Institute, Washington DC, United States

Word count: Main text: 6563; References: 3871, abstract: 212

1 **Abstract**

2 Ecologists and farmers often have contrasting perceptions about the value of natural
3 habitat in agricultural production landscapes, which so far has been little
4 acknowledged in ecology and conservation. Ecologists and conservationists often
5 appreciate the contribution of natural habitat to biodiversity and potential ecosystem
6 services such as biological pest control, whereas many farmers see habitat remnants
7 as a waste of cropland or source of pests. While natural habitat has been shown to
8 increase pest control in many systems, we here identify five hypotheses for when and
9 why natural habitat can fail to support biological pest control, and illustrate each with
10 case studies from the literature: (1) pest populations have no effective natural
11 enemies in the region, (2) natural habitat is a greater source of pests than natural
12 enemies, (3) crops provide more resources for natural enemies than does natural
13 habitat, (4) natural habitat is insufficient in amount, proximity, composition, or
14 configuration to provide large enough enemy populations needed for pest control,
15 and (5) agricultural practices counteract enemy establishment and biocontrol
16 provided by natural habitat. In conclusion, we show that the relative importance of
17 natural habitat for biocontrol can vary dramatically depending on type of crop, pest,
18 predator, land management, and landscape structure. This variation needs to be
19 considered when designing measures aimed at enhancing biocontrol services
20 through restoring or maintaining natural habitat.

21

22 Key words: pest regulation, natural enemies, agricultural management, ecosystem
23 services, spillover, landscape structure, predators, parasitoids

1 Introduction

2 We are facing unprecedented declines in global biodiversity and associated
3 ecosystem services largely due to enduring losses in natural habitat. As agriculture
4 now occupies 38% of Earth's terrestrial area (Foley *et al.* 2011), remnants of natural
5 habitat in human-dominated landscapes deserve increasing attention for
6 conservation. In fact, small patches are the dominant form of natural habitat on Earth
7 (Haddad *et al.* 2015). However, ecologists and farmers often have contrasting
8 perceptions about the value of remaining natural habitat in agricultural landscapes
9 (defined here as the combination of natural or "semi-natural" non-crop habitats such
10 as cropland boundaries, fallows, grasslands, woodlands, wetlands, and forests).
11 Farmers often view natural habitat remnants as a waste of potential cropland,
12 barriers for mechanization or a source of pests and diseases, and thereby, as costs
13 or lost economic opportunity. In contrast, proponents of maintaining or even restoring
14 natural habitat make two arguments. First, keeping natural habitat in agricultural
15 landscapes promotes conservation of wild biodiversity and, second, natural habitat
16 provides important ecosystem services including pest control (Landis *et al.* 2000;
17 Bianchi *et al.* 2006; Karp *et al.* 2013; Shackelford *et al.* 2013; Milligan *et al.* 2016),
18 but also soil conservation (Mäder *et al.* 2002), nutrient retention (Raudsepp-Hearne
19 *et al.* 2010), crop pollination (Klein *et al.* 2003; Carvalheiro *et al.* 2010), and cultural
20 services (van Zanten *et al.* 2014; Riechers *et al.* 2015).

21 Natural habitat heterogeneity at multiple spatial and temporal scales, not just
22 the amount of natural habitat, is a major determinant of biodiversity in agriculture
23 (Benton *et al.* 2003; Schellhorn *et al.* 2015; but see Batáry *et al.* 2011).
24 Heterogeneous landscapes with a diversity of, often intermingled, habitat types
25 generally increase biodiversity and the services that flow from them (Tscharntke *et al.*
26 2005a). Therefore, combining agricultural land use with natural habitat fragments in

mosaic landscapes can be beneficial for biodiversity conservation, increasing environmental benefits, ecosystem services, and human wellbeing (Perfecto and Vandermeer 2010; Tscharntke *et al.* 2012a). However, prioritizing management for biodiversity might limit management options and priorities for provisioning ecosystem services such as crop production (Kleijn *et al.* 2011; Macfadyen *et al.* 2012).

As shown in several recent reviews, natural enemy populations are on average higher and pest pressure can be lower in complex, heterogeneous landscapes versus simple, homogeneous landscapes, leading to enhanced pest suppression and lowered crop injury (Bianchi *et al.* 2006; Tscharntke *et al.* 2007; Chaplin-Kramer *et al.* 2011a; Blitzer *et al.* 2012; Rusch *et al.* 2016). For example, it has been shown that landscapes with large amounts of natural habitat exhibit higher parasitism rates and lower oilseed rape damage by pollen beetles in Germany (Thies and Tscharntke 1999) and higher biocontrol of cereal aphids across Europe (Thies *et al.* 2011; Rusch *et al.* 2013b).

Despite the strong general evidence of the benefits of natural habitat to sustaining or restoring biological pest control in agricultural landscapes, variability is high and there is also scattered evidence for the reverse; that is, natural habitat can have no, or even negative, effects on pest control. In this perspective paper, we identify several conditions under which we should not expect natural habitats to benefit natural biological control of crop pests. We present five, non-mutually exclusive hypotheses (Table 1) for the failure of natural habitats to support biological pest control and illustrate each with published evidence, selecting examples from across geographic regions and taxa:

- (1) Pest populations have no effective natural enemies in the region,
- (2) Natural habitat is a greater source of pests than natural enemies,
- (3) Crops provide more resources for natural enemies than does natural habitat,

(4) Natural habitat is insufficient in amount, proximity, composition or configuration to provide large enough enemy populations for pest control,

(5) Agricultural practices counteract enemy establishment and biocontrol provided by natural habitat.

We provide evidence to support these five hypotheses and derive recommendations for how to manage natural habitat or cropland at different spatial and temporal scales for improving biological control and pest suppression in agricultural landscapes.

(1) Pest populations have no effective natural enemies in the region

For some pests, population dynamics in agricultural landscapes may not be controlled by natural enemies, regardless of the availability of natural habitat at the regional scale. Instead, pest density and outbreaks may be driven by other factors such as abiotic conditions, crop susceptibility, agricultural practices, crop area, or intraguild predation of higher trophic levels.

A classic example of massive pest outbreaks, and probably one of the most damaging pests in the world, is the biblical plague, the outbreak of the migratory locust *Locusta migratoria* (Lomer *et al.* 2001). The factors determining phase polyphenism (the solitary and the gregarious phase) and migratory dynamics in grasshoppers and locusts are complex and variable, but they are dominated by abiotic factors such as rainfall and temperature, and not natural enemies (Lomer *et al.* 2001). Nevertheless, the clearing of natural habitat has increased dry season survival, prompting locust outbreaks in new locations (Farrow 1974).

In addition to the migratory locust, there are many other examples of major pests that do not have effective natural enemy communities. A well-studied example is the Colorado potato beetle, *Leptinotarsa decemlineata*. Even in its natural range in northern Mexico on wild *Solanum*, high early-season fecundity makes defoliation of

host plants likely. Many natural enemies become abundant only late in the season. In potato fields in Canada, Washington, and Wisconsin, no effective natural enemies are present (Hough-Goldstein *et al.* 1993). Potato yield in Colombia has been shown to be negatively affected by higher percentage of land cropped with potatoes due to the response of another specialized pest, the Guatemalan potato moth, while natural enemies had no effect on crop damage or yield (Poveda *et al.* 2012). Corn rootworms (*Diabrotica virgifera* and *D. barberi*) also inflict significant economic damage yet few natural enemy species have been identified (Levine and Oloumi-Sadeghi 1991, but see Lundgren and Fergen 2011) and their response to land-use is thought to be driven by the area of corn in the landscape (O'Rourke *et al.* 2011).

Invasive species usually arrive to a new area without their co-evolved predators, parasites, and pathogens and, by escaping from these mortality agents, often increase and spread rapidly in the new environment (but see Hawkins *et al.* 1997). The practice of classical biological control consists of locating natural enemies in the pest's native range and evaluating the results of their importation, quarantine, testing, and release in the new environment (Messing and Wright 2006) in an effort to re-establish biological control of populations. The globally invasive silverleaf whitefly, *Bemisia tabaci*, is a prime example; several native species of *Bemisia* and their natural enemies are present in Australia, and although 12 of these native natural enemies attacked the invasive *B. tabaci*, control never exceeded 5% (DeBarro and Coombs 2009). In the absence of classical biological control, however, there may simply be no effective natural enemies to control pests, regardless of the amount and/or configuration of the surrounding natural habitat. However, in the long run, generalist natural enemies may exploit new prey resources, causing steadily increased pest mortality rates (Symondson *et al.* 2002, Strong *et al.* 1974).

1 In biological control with parasitoids, the number of parasitoid species can be
2 related to host mortality rates (Tylianakis *et al.* 2006), and hosts with few enemies
3 may not be effectively controlled (but in many other cases, single agents are more
4 effective than multiple agents, Denoth *et al.* 2002). The size of the parasitoid complex
5 differs with the hosts' feeding niche and is, for example, highest in leaf miners and
6 lowest in root feeders (Hawkins 1994): leaf miners suffer on average from 6
7 parasitoid species and roughly 65% parasitism, whereas root feeders suffer from 2
8 parasitoid species with 17% parasitism (Hawkins 1994). When the percentage of
9 parasitism drops below a threshold value of 32-36%, a success in classical biological
10 control has never been found (Hawkins and Cornell 1994). Hence, the feeding niche
11 of the pest may be a reasonable predictor of biocontrol potential.

12 Intraguild predation may also constrain the role of natural enemies for pest
13 suppression. A study in South Korean landscapes found that the effects of natural
14 habitats on pest control were ambiguous (Martin *et al.* 2013). An enclosure
15 experiment showed that density and damage by cabbage butterfly larvae increased
16 with increasing amounts of semi-natural habitat. There was also higher pest control
17 by wasps and syrphids in these areas. However, insectivorous birds appeared to
18 consume not only the cabbage pest caterpillars, but also the wasps that control them,
19 thereby releasing the pests from biocontrol. Since birds benefited from natural
20 habitat, ultimately pest damage was highest and yields lowest in the areas with
21 highest cover of natural habitat (Martin *et al.* 2013). In contrast, Karp *et al.* (2014)
22 found that insectivorous birds mainly consumed herbivorous arthropods, while bats
23 also consumed predatory arthropods. Therefore, birds increased coffee yields and
24 prevented leaf damage, while bats did not. In a DNA study Traugott *et al.* (2012)
25 showed that parasitoids can be major prey of generalist predatory beetles. Davey *et*
26 *al.* (2013) also reported high levels of intraguild predation in wheat fields, between

the very abundant carabid beetle *Pterostichus melanarius* and several juvenile spiders (mostly Linyphiidae). The various and often unpredictable occurrences of intraguild predation or mesopredator suppression in diverse enemy communities may be one reason why effects of natural enemies are variable (as also shown by Tscharntke 1992, 1997) and natural habitat fails to enhance control.

Finally, a positive effect of natural habitats on natural enemies may be present only periodically. For example, Meisner *et al.* (2014) observed apparent pea aphid-parasitoid cycles at the regional scale with a period of roughly 2/3rds of the year, leading to peaks in biocontrol once (mid-summer) or twice (spring and fall) a year in alternating years; any effects of landscape composition on control by parasitoids would thus necessarily be episodic as well. Similarly, after the introduction of soybean aphids into the Midwest USA, regional aphid outbreaks appeared to alternate between years along with variable biocontrol effects by the ladybeetle *Harmonia axyridis* (Bahlai *et al.* 2015), while landscape composition may have been only periodically important. Indeed, the pattern of alternating years of high and low abundance has often been described in aphid pests (Sequeira and Dixon 1997). Within years, a common pattern for soybean aphids (Mueller *et al.* 2010) and other pests is the initial low abundance of natural enemies early in the growing season, followed by an increase in natural enemies as crops mature and pest populations build, so the effects of landscape composition are expected to be seen most strongly later in the season (Costamagna *et al.* 2015, but see Östman *et al.* 2001).

These examples illustrate that pest species may not always be controlled effectively by their natural enemies, in which cases other management practices need to be taken into account for pest suppression. Management approaches may be based on bottom-up control (Scherber *et al.* 2010), e.g. by increasing within-field diversity (Letourneau *et al.* 2011).

(2) Natural habitat is a greater source of pests than natural enemies

Natural habitats can provide a suitable environment for a large number of pest and natural enemy species at several key stages of their life-cycle (Landis *et al.* 2000). It is generally assumed that these habitats host a larger proportion of beneficial or neutral species than detrimental ones (Denys and Tscharntke 2002), and that natural enemies are highly mobile (Schellhorn *et al.* 2014). For example, Keller and Häni (2000) reported that about 9 out of 10 natural enemy species require natural habitats during their life cycle, whereas only 5 out of 10 pest species require natural habitats. However, in some cases, natural habitats can be a major source of pests but not of natural enemies (Blitzer *et al.* 2012).

Natural enemies may have large populations in natural habitats but will not disperse, while weeds and pests from natural habitat can spill over into cropland (Blitzer *et al.* 2012). This may be true for natural enemies in forested habitats that avoid non-shaded open land (e.g., tropical insectivorous bird species) (Tscharntke *et al.* 2008). Similarly, there are many predatory arthropods that rarely venture outside forests. Fischer *et al.* (2013) found that many carabid species remain in forest edges and may immigrate to adjacent hedges, but avoid colonizing neighbouring cropland. Gaines and Gratton (2010) showed that abundant herbivorous carabids in field margins do not venture out into potato crop fields, resulting in lower predation rates on weed seeds. In California, colonization of cotton fields by a major pest, the plant bug *Lygus hesperus*, is lowered in cotton fields surrounded by large areas of cotton (Carrière *et al.* 2006, 2012), whereas uncultivated areas in the surroundings can significantly elevate *Lygus* densities because they provide major overwintering sites for the pests while natural biocontrol is weak (Sivakoff *et al.* 2013).

1 In other cases, natural enemies from natural habitat may colonize cropland,
 2 but simply not as well as pests do. In western Kenya, grasslands surrounding 'push-
 3 pull' maize provided habitat for both stem borers and their parasitoids (Midega *et al.*
 4 2014), but the net effect was an increase in stem borer density in landscapes with
 5 more grassland. Similarly, pollen beetle densities and damage in oilseed rape are
 6 higher in fields surrounded by semi-natural habitats (and especially woodland) in
 7 northwestern France (Rusch *et al.* 2011, 2012, 2013a), despite high levels of
 8 biological control by parasitoids in these complex landscapes. Pollen beetles and
 9 their parasitoids both directly benefited from semi-natural habitats, and parasitism
 10 rates of pollen beetles increased linearly with landscape complexity (i.e. landscapes
 11 with high amount of semi-natural habitats). However, this strong top-down effect was
 12 compensated over time by pollen beetle population dispersal from simple landscapes
 13 (i.e. landscapes with little natural habitat left), where they were not controlled, to
 14 complex landscapes, where they moved for overwintering. So, in the end, biocontrol
 15 of pollen beetles does not seem to be effective in complex landscapes from one year
 16 to the next, even if very high parasitism rates (>80%) are observed (Rusch *et al.*
 17 2011, 2012, 2013b).

18 Counterbalancing effects of landscape complexity have also been found for
 19 cereal aphids in Germany (Thies *et al.* 2005). In complex landscapes with high
 20 amounts of semi-natural habitat, aphid mortality by parasitoids (the main biocontrol
 21 agents) was high, but spring colonization by winged aphids was also high. Hence, in
 22 the beginning of the season, pest density was highest in complex landscapes, but
 23 towards the end of the season, densities were similar (due to higher enemy impact in
 24 complex landscapes) along the landscape complexity gradient.

25 Natural habitat can also serve as an important reservoir for weeds as well as
 26 crop diseases (Wisler and Norris 2005). The annual life cycle of most crops means

1 that perennial wild hosts are usually needed as a major overwintering resource for
2 pathogens (Power and Mitchell 2004). A variety of viruses, bacteria and pathogenic
3 fungi are known to spillover from weeds in overwintering sites and wild habitats to
4 crops (Power and Mitchell 2004; Blitzer *et al.* 2012). The older crop-protection
5 literature provides much evidence for natural habitat as a source and reservoir of
6 pests (>70 arthropod families) (Norris and Kogan 2000), although some of these
7 insects or pathogens can also suppress weeds and may thus be considered
8 beneficial (Hatcher and Paul 2001).

9 Indeed, natural habitat may simultaneously provide both benefits and threats
10 to nearby crops. Corbett and Rosenheim (1996) showed that natural riparian habitat
11 includes plants that are key overwintering sites for egg parasitoids (genus *Anagrus*),
12 which are major biocontrol agents of leafhopper pests attacking cultivated grapes
13 (see also Murphy *et al.* 1996). However, the very same riparian community also
14 includes host plants that harbour the bacterial plant pathogen *Xylella fastidiosa* as
15 well as the insect vectors of this pathogen. As a result, the incidence of bacterial
16 infection in grapevines is most severe near riparian plant communities (Daugherty *et*
17 *al.* 2012). The net effect is that some grape growers choose to completely destroy
18 riparian habitats because the ecosystem disservices (pathogens that kill vines)
19 outweigh the ecosystem services provided by the parasitoids.

20 There is also the possibility that certain elements of natural habitat benefit
21 pests relative to natural enemies, especially when non-native plants invade natural
22 habitats. Remnant native habitat in agricultural landscapes can vary from near
23 pristine to degraded (McIntyre and Hobbs 1999), containing a high proportion of
24 invasive plants. This has been shown in Australia where non-native weeds in native
25 remnants and pastures can host natural enemies, but they often host far more pests,

1 whereas the native plants rarely host pests of crops, but do support their enemies
2 (Schellhorn *et al.* 2010; Parry *et al.* 2015).

3 In the grasslands of California, which are similarly dominated by invasive
4 weeds, non-native weeds in the mustard family allow cabbage aphids to become far
5 more toxic to their natural enemies than when feeding on crops, and thus to build up
6 populations in virtually enemy-free space before spilling back into nearby cole crops
7 in much greater numbers than their enemies can control (Chaplin-Kramer *et al.*
8 2011b). In the Midwestern United States, the dominant winter hosts of soybean aphid
9 are European and glossy buckthorn, non-native shrubs that have invaded the
10 woodlands of the Great Lakes states (Heimpel *et al.* 2010). In Wisconsin, higher
11 amounts of forest in the landscape are associated with increased soybean aphid
12 populations (Stack-Whitney *et al.* unpubl. data). On the other hand, Bahlai *et al.*
13 (2010) found that biocontrol of soybean aphid improves in areas with more natural
14 habitat. Therefore, depending on the abundance of particular plant species
15 (buckthorn), natural habitat may play both positive and negative roles, though the
16 overall effect of natural habitat on biocontrol in that system is often negative.

17 Similarly, plant species providing floral resource subsidies can have different
18 effects on herbivores and their natural enemies (Campbell *et al.* 2012). Using three
19 different herbivore-parasitoid systems, Lavandero *et al.* (2006) showed differential
20 responses of parasitoid fitness and herbivory to nectar sources. Some flowering
21 plants enhance both herbivore and parasitoid fitness whereas other plants, such as
22 buckwheat and phacelia, only enhanced parasitoid fitness. This study nicely
23 illustrates how plant species composition can affect the potential of non-crop habitat
24 to deliver biological control.

25 Segoli and Rosenheim (2012) used a simulation model to understand how
26 pest densities are affected by crop field size and, therefore, the amount of edge

habitat. They argued that the relative dispersal abilities of pests and natural enemies from edges to crop, their likelihood of overwintering in the crop field, and the pest's rate of reproduction in the field determine the outcome of the interaction. This provides some theoretical explanation for why natural habitat appears to benefit pests more than enemies in certain cases, such as the examples shown here.

The studies in this section illustrate how natural habitat can be a major source of pests, if the plant composition in these habitats benefits pests more than natural enemies or if natural enemies remain within natural habitats and contribute little to pest suppression in cropland. This may be due to the enemy species' niche and associated habitat requirements (e.g. preference of forested over non-forested habitat). Furthermore, even if natural habitat is a source of natural enemies, the concurrent enhancement of pests can result in a net positive effect of natural habitats on pest populations. Weeds, pathogens, and arthropod pests may also invade certain cropland, and in countries with high pressure of invasive plants, this is often particularly important. When pests benefit from natural habitat remnants, farmers are understandably reluctant to create or maintain crop boundary habitat. More detailed research is needed to tease apart the differential effects of specific plant composition or other elements of natural habitat on pests and natural enemies.

(3) Crops provide more important resources for natural enemies than do natural habitats

Natural habitat may not always be a panacea for natural enemies. In some cases, natural habitat has relatively low productivity and thus supports relatively small natural enemy populations compared to cropland. This applies particularly to landscapes with poor soils, summer dryness (Mediterranean or continental climate),

1 or arid conditions. In such cases, natural enemies may be more influenced by the
2 composition of surrounding cropland than natural areas.

3 For example, Gardiner *et al.* (2009) found that some species of ladybeetles in
4 landscapes of Midwestern USA are more abundant when cropland increases.
5 *Coleomegilla maculata*, in particular, is a generalist natural enemy more commonly
6 found in corn fields than grasslands, as its diet can consist entirely of corn pollen
7 (Werling *et al.* 2011),

8 Costamagna *et al.* (2015) showed that melon aphid suppression on sentinel
9 rockmelon plants in subtropical Australia was positively correlated with the amount of
10 lucerne pasture within 1.5 km around fields, but neutral or negatively associated with
11 the amount of natural or semi-natural vegetation. Further, the number of predators
12 captured was highest in landscapes with larger areas of crops and lucerne within 0.5
13 km. This is likely because in sub-tropical and Mediterranean regions of Australia,
14 many crops are irrigated, whereas the natural and semi-natural habitats experience
15 long periods of droughts and become relatively unproductive and inhospitable. This is
16 in contrast to temperate and cold regions where natural habitats and forests maintain
17 their productivity during the growing season and provide overwintering sites for
18 dormant natural enemies (Rand et al. 2006).

19 A recent study in temperate vineyard landscapes found decreasing activity-
20 density of ground beetles with increasing landscape complexity, while species
21 richness and evenness remained unchanged along the landscape gradient (Rusch et
22 al., 2016). This negative effect of semi-natural habitats on ground beetle activity-
23 density is attributable to the fact that grass cover is maintained throughout the year in
24 the majority of vineyards in this region, providing overwintering sites and resources
25 throughout the year. This was especially true for the the dominant ground beetle
26 species found in these landscapes, which are known to prefer open-habitats and to

1 overwinter in fields or field margins. This study illustrates that in some cases crops
 2 can provide more resources for natural enemies than do semi-natural habitats,
 3 thereby revealing negative relationships between the amount of semi-natural habitat
 4 in the landscape and the abundance of natural enemies within crops.

5 If natural habitat supports little primary and secondary production (e.g., prey
 6 resources, Dreyer and Gratton 2014), natural enemy communities may be better
 7 supported over the course of a year by providing a continuous flow of crop-based
 8 resources over space and time rather than by maintaining nearby natural habitat
 9 (Schellhorn *et al.* 2015). This could occur through a mosaic of irrigated crops that,
 10 over time, provide complementary resources, helping enemy populations to avoid
 11 spatial and temporal bottlenecks. The potential benefits of maintaining
 12 heterogeneous agricultural landscapes to favour natural enemies and biological
 13 control has been argued for in rice crops (Way & Heong 1994). Evidence suggests
 14 that maintaining asynchronous harvesting and planting of crops will likely favour
 15 natural enemies over pests (Sawada *et al.* 1992), because asynchronous cropping
 16 provides mobile specialist natural enemies with a continuous supply of resources
 17 (Ives & Settle 1997). However, asynchronous harvesting may also supply continuous
 18 sources of food for herbivorous pests.

19 Ensuring continuous availability of food for natural enemies could benefit
 20 biological control through the strategic planting and maintenance of crops with
 21 differing phenologies that create a temporal mosaic of cropland in the landscape, but
 22 temporal and spatial contribution of crop configuration and composition has so far
 23 been little explored.

24
 25 **(4) Natural habitat is insufficient in amount, proximity, composition or**
 26 **configuration to provide enough enemy populations for pest control**

To enhance pest control sufficiently, natural habitats must be both large enough and near enough to farm fields to allow a substantial increase in within-field enemy abundance (Holland *et al.* 2016). Field margin strips, for example, may often be so small that the enemy populations they support are insufficient to be effective for pest control (Segoli and Rosenheim 2012, Dreyer and Gratton 2014). For example, control of pollen beetles by parasitoids was highest in landscapes with large amounts of semi-natural habitat (>20%), lower in simple landscapes with only one fallow field adjacent to the oilseed rape field (>1 ha), and lowest when only a narrow field margin strip (3 m) was present adjacent to the crop (Thies and Tscharntke 1999).

Even within crop fields, the distance between natural habitat patches may be too large to facilitate biological control. Lys *et al.* (1994) subdivided a large winter cereal field to have a series of five 1.5-m-wide weed-strips with 12, 24 or 36 m between the strips. They found that the activity density of ground beetles was highest near the weed strips, such that only very narrow rows between strips had enhanced biological control. Similar effects have been found for parasitoid populations and parasitism rates at different distances from field margins and floral strips (Tylianakis *et al.* 2004; Lavandero *et al.* 2005). Henri *et al.* (2015) found that natural enemies of African fruit crop pests decline and pests increase with distance to natural habitat, and that biocontrol was greatly limited by enemy dispersal. Specialist enemies in particular appear to be less effective colonizers of distant cropland. Chaplin-Kramer *et al.* (2011a) found that specialist enemies respond to prey resources at smaller spatial scales than generalist enemies.

In South Korea, Martin *et al.* (2016) found that the amount of semi-natural habitat had no influence on many natural enemies and even had a negative effect on parasitoids, but the configuration of patches (increasing edge density) and landscape compositional heterogeneity had a positive effect. Configurational and compositional

diversity has been shown to be important to natural enemies in several studies (e.g. Hendrickx *et al.* 2007; Holzschuh *et al.* 2010). Since natural enemies often need to move between crop and non-crop resources (Tscharntke *et al.* 2005b, 2012b), large amounts of habitat may be less critical than easy access between neighbouring habitats and cropland (Martin *et al.* 2016). This finding has been shown theoretically in spatially-explicit simulation models where the spatial arrangement of source habitats for natural enemies of agricultural pests had profound effects on their potential to colonize crops and suppress pest populations (Bianchi *et al.* 2010). Perović *et al.* (2010) demonstrated that spatial configuration metrics had higher explanatory power to explain the within-crop density of some predator species than composition metrics in Australian cotton landscapes; hence, manipulating the connectivity of woodland in the landscape might be a management option to increase density of these predators.

These examples indicate that patches of natural habitat can support biocontrol agents, but if the available habitat area is too small or isolated, biocontrol services will not be provided (e.g., Bianchi and van der Werf 2003). A threshold level of 20% natural habitat at the landscape scale has been suggested as the minimum amount of natural habitat necessary to support biocontrol services, while the necessary level of habitat connectivity is unknown (Tscharntke *et al.* 2007a). Shifting our research focus from local to landscape configuration effects on functional biodiversity will be critical to developing solutions for future management of biological control (Tscharntke *et al.* 2012b).

(5) Agricultural practices counteract enemy establishment and biocontrol provided by natural habitat

Intensive agricultural practices can disrupt the biological control delivered by natural enemy populations from surrounding natural habitats and decrease the effectiveness of 'conservation biological control' (van Driesche *et al.* 2009). Vice versa, pesticide applications in the other fields of the surrounding landscape may disrupt biocontrol in the focal field (Bianchi *et al.* 2013). Broad-spectrum pesticide spraying, deep ploughing, planting high-yield and highly susceptible crop varieties as well as little crop diversification may all negatively affect natural enemies, even if surrounding natural habitats are present (Tscharntke *et al.* 2005; Poveda *et al.* 2008; Letourneau *et al.* 2011; Iverson *et al.* 2014). In a study of pests and parasitoids on *Brassica* (kale) in New Zealand (Jonsson *et al.* 2012), low parasitism rates in intensively managed landscapes were attributed to frequent insecticide applications in surrounding crops rather than a lack of resources in non-crop habitats. Thus, the landscape level management of agricultural land appeared to be more important for biological pest management than availability of natural habitat.

Similarly, in a highly replicated study across eight European regions, Geiger *et al.* (2011) found that wheat yield was negatively related to the density of natural enemies such as carabid beetles and positively related to the survival of aphids and their abundance. This appeared to be due to the high level of pesticide applications in intensively managed high-yielding crop landscapes. On organic farms without pesticide applications, aphid predation increases (Winqvist *et al.* 2011) and aphid density decreases (Thies *et al.* 2011) compared to conventional farms. In China, reducing insecticides also conserved aphid enemies, which subsequently regulated cotton aphid populations. Lu *et al.* (2012) found a marked increase in the abundance of three types of generalist arthropod predators (ladybeetles, lacewings and spiders) and a decreased abundance of aphid pests associated with widespread adoption of Bt (*Bacillus thuringiensis*) cotton and reduced insecticide sprays. However, Lu *et al.*

(2010) suggested that the increase in herbivorous mirid bug populations is due to the regional adoption of transgenic Bt cotton for the control of lepidopteran pests. This is because insecticide use, which had controlled mirid bugs, has also dropped, and there is little control of mirid bugs by natural enemies (Lu and Wu 2008).

In Indonesia's rice production, insecticides were initially subsidised by the government to control a major stem-boring rice pest. Increased pesticide use caused the uprising of the brown planthopper (*Nilaparvata lugens*) by initially releasing it from its natural predators, and subsequently selecting for those individuals that could feed on high performance rice varieties. The brown planthopper developed into a disastrous pest, causing a loss of rice that could have fed two million people for one year and an economic loss of US\$100 millions (Settle et al. 1996). Solutions to this problem should include the removal of insecticide subsidies and introduction of an integrated pest management program (Settle et al. 1996). Bats are natural biocontrol agents of the white backed planthopper, one of the major rice pests in Asia, but are affected by pesticides in ricefields, while roosting boxes in the landscape may boost the bat population and provide additional biocontrol services (Wanger et al. 2014).

Moreover, vegetation management and soil tillage can affect generalist predators and predation rates, whether natural habitat is surrounding the field or not (Perfecto 1990; Rusch et al. 2010). Thorbek and Bilde (2004), for instance, showed direct and indirect effects of soil cultivation and grass cutting on generalist predators such as spiders. Soil tillage has also been found to strongly affect parasitoid populations. Several studies have shown that the abundances of pollen beetle parasitoids overwintering in crop soil were reduced both by soil tillage during parasitoid overwintering and by insecticide treatment at emergence (Nilsson et al. 2010; Rusch et al. 2011).

In addition to counteracting pest predation, management practices may also contribute to high pest densities that no longer can be effectively controlled by natural enemies. For example, in coffee systems, the primary cause of coffee berry borer outbreaks is farmers' failure to prevent borers from overwintering in fallen berries or berries left on coffee plants after harvest (Avelino *et al.* 2012). Similarly, neglecting to remove pathogen-infested cacao pods increases infection levels in the following harvest (Tscharntke *et al.* 2011b).

These illustrative examples make it clear that agricultural practices can interfere with effective biological control. Landscape management is not a panacea, as other practices can strongly influence biocontrol. Important practices may include reducing applications of agrochemicals to maintain and promote natural enemies as well as pest specific measures to prevent the build-up of pest populations.

Conclusions and outlook

The claim that increasing the amount of natural habitat in a landscape is generally a benefit for biological pest control is fairly common, but there appear to be many exceptions to this rule. Surprisingly few studies have shown higher yield of farmers following conservation biological control, driving pests below economically important thresholds through local and landscape management (Settle *et al.* 1996; Cook *et al.* 2007; Landis *et al.* 2008, Maas *et al.* 2013, Pywell *et al.* 2015, Tschumi *et al.* 2016).

Natural habitat often provides biological control, but there are also many unexplained patterns. In this critical review, we aim to identify the conditions under which we do not observe the generally expected biocontrol. We provide evidence for the concern of many farmers that natural habitat sometimes does not help and can

1 even be a source for pests, and we identify five conditions under which natural
2 habitat would not provide effective biological pest control.

3 If natural enemies do not effectively control a particular pest species, or if
4 natural enemies do not disperse to adjacent cropland, then other methods of pest
5 control, such as improved plant resistance, crop rotation, or crop diversification, need
6 to be implemented. If natural habitat is a major source of pests, then natural habitats
7 should be placed at distances to the crop field larger than the dispersal distances of
8 pests. If natural habitats are degraded and contain weeds that are a major source of
9 pests, then weed management is needed. If neighbouring crops provide more
10 important resources for natural enemies than natural habitat, then reconfiguring
11 landscapes into a mosaic of small crop fields that provide continuous resources
12 across seasons may help. If the amount of natural habitat in agricultural landscapes
13 is too small to support economically meaningful biological control, then increasing
14 habitat amount could be worthwhile (if opportunity costs are not considerably higher
15 than the socio-economic benefits of enhanced ecosystem services). Alternatively,
16 targeted measures on small parcels of land may be possible as a means to secure
17 the continuity of resources throughout the life cycle of biological control agents and
18 release limitations to population growth (Schellhorn et al. 2015). Finally, because
19 local agricultural practices in intensively managed farmland (e.g., high levels of
20 pesticide spraying) may counteract pest control derived from natural habitat, farmers
21 may need to eliminate broad-spectrum insecticides and replace them with target-
22 specific products less harmful to biological control agents while working towards
23 judicious spraying to realize benefits.

24 Although landscape-scale habitat composition and configuration is often key
25 for managing local processes such as biological pest control, agri-environment
26 schemes and eco-labelling are usually aimed at the field and farm level (Tschamntke

1 *et al.* 2011a, 2015). Hence, agri-environment programmes should be broadened to
 2 the landscape scale to motivate farmers to adopt schemes beyond their farm (Gabriel
 3 *et al.* 2010). Similarly, certification programmes could help to maintain biodiversity
 4 and ecosystem services at larger spatial scales, if certification is applied across
 5 several neighbouring farms in one region (McKenzie *et al.* 2013; Tscharncke *et al.*
 6 2015). Moreover, participatory approaches, including the development, research, and
 7 implementation of regional programmes for conservation biological control, might
 8 help to involve farmers and to enhance the acceptance and effectiveness of
 9 landscape-wide management of biocontrol and other ecosystem services (Westphal
 10 *et al.* 2015). Farm cooperatives for sharing machinery, crop certification, or water
 11 table management are already well-established examples of large-scale programs
 12 that affect land-use decisions beyond the farm level.

13 In the absence of public financial support, enhancing biological pest control
 14 services by natural habitat is widely viewed as only being successful if there are
 15 economic benefits for farmers, which requires information on a number of critical
 16 ecological and economic conditions. Depending on the expectations of stakeholders
 17 (e.g., scientists and farmers), the conditions that must be met can vary. For example,
 18 if managed natural habitats are seen as part of preventative pest management,
 19 enhancing stability and resilience of cropland, then the ecological and economic
 20 evidence can be: reduced and infrequent pest outbreaks, pest populations staying
 21 below economic injury levels for longer, reduced need for insecticide use, hence
 22 reduced cost, and less occurrence of crop damage and loss. However, if
 23 conservation biological control serviced by natural habitat is seen as the primary pest
 24 management tool, then the burden of proof is far more stringent. First, natural habitat
 25 must support large populations of natural enemies. Second, these natural enemies
 26 must spill over into cropland. Third, natural enemies must significantly suppress pest

1 populations. Fourth, this suppression must lead to reduced crop damage. Fifth,
2 reduced crop damage must increase crop yield or value, or reduce input costs due to
3 insecticide use, and this should increase farmers' profits. Finally, overall increased
4 profit from diversified croplands should outweigh the opportunity cost of setting aside
5 non-crop habitat. The first five steps are supporting ecosystem services, or
6 intermediate contributions to marketed products (Zhang and Swinton 2012), whereas
7 for many farmers, only the final outcome counts: the decreased input cost, or the
8 increased crop yield, quality, and profit (Mace *et al.* 2012; Wielgoss *et al.* 2014; Wong
9 *et al.* 2015; although benefits beyond economic profits, such as public health,
10 reduced environmental pollution and cultural services, need also be considered).

11 Beyond direct economic benefits to farmers, stability of yields, risk reduction,
12 food safety, and an enabling institutional environment for coordination are further
13 important parameters that need to be considered when evaluating the viability of
14 habitat-based biological control. For example, farmers' risk preferences play an
15 important role in agricultural production decisions (Feder 1980; Just and Zilberman
16 1983). Uncertainty in the effectiveness of conservation biological control could hinder
17 adoption, as farmers may choose the (apparently) less risky method of insecticide
18 spray, even if biological pest control is more profitable. Regarding food safety, fear
19 that wildlife might spread foodborne diseases has caused many California produce
20 growers to remove non-crop vegetation (perceived as wildlife habitat) and adopt a
21 view that sanitized, hospital-like conditions are critical for responsible produce
22 management, even though there is no evidence that non-crop vegetation
23 compromises food safety (Karp *et al.* 2015a, b). Still, the global preponderance of
24 organic farming provides an intriguing example of how reliance on biological pest
25 control can be viable. On the one hand, effective biological control could help
26 maintain farm profitability. On the other hand, viability could be supported by price

premium paid by consumers who perceive the benefits of reduced human health risk from insecticides (Sexton *et al.* 2007) and environmental damage (soil, water, air pollution, e.g. due to nitrogen losses, Sutton and Grinsven 2011). Without accounting for the full costs of insecticides on human health and the environment, insecticide price alone may not send a strong enough market signal to promote widespread adoption of alternative pest control methods related to conservation biological control (Meehan *et al.* 2011).

In conclusion, there are many examples in the literature where the amount and configuration of natural and semi-natural habitat in agricultural landscapes helps to enhance pest control and reduces probabilities of pest outbreaks. Our synthesis adds to this literature by showing that the relative importance of natural habitat for supporting natural enemies and pest control can vary dramatically with the type of crop, insects, habitat, management regime, and landscape type considered. We suggest that smart management on local scales (pesticide avoidance, implementing habitat patches, replacing invasive with native flora) and on landscape scales (increasing habitat availability and crop diversity as well as configurational diversity by smaller cropland patches) can help improve the chances that biological control will help us meet future food demand while preserving the environment and ecosystems. These types of management actions, if carefully executed, could also have positive consequences for other ecosystem services, such as cultural services, pollination, soil conservation, nutrient retention, and climate regulation, which need to be acknowledged in a holistic management approach.

Acknowledgements

We are grateful for many helpful comments by three anonymous reviewers. Author sequence follows the “sequence-determines-credit” (from TT to RCK) and the

“equal-contribution” norm (from PB to WZ) (see Tscharntke et al., 2007b). This work benefited from support from the National Socio-Environmental Synthesis Center (SESYNC) - NSF award DBI-1052875 for the project “Evidence and Decision-Support Tools for Controlling Agricultural Pests with Conservation Interventions” organized by Daniel Karp and Becky Chaplin-Kramer. TT acknowledges support by the DFG-CRC 990 EForTS and the DFG-FOR 2432, TT and PB by the Biodiversa project FarmLand, DK by a Killam Postdoctoral Fellowship, and MJ by the Biodiversa project APPEAL. CG was supported in part by the US DOE Great Lakes Bioenergy Research Center (DOE BER Office of Science DE-FC02-07ER64494), the US DOE OBP Office of Energy and Renewable Energy (DE-AC05-76RL01830), and by the USDA NIFA Agriculture and Food Research Initiative (2011-67009-30022). AR acknowledges support by the Era-net project MULTAGRI (EU/FP7), NAS by the Julius Career Award and the Cotton Research & Development Corporation, FDC and WZ by CGIAR’s Water, Land and Ecosystems (WLE) research program, AL by the EPA Science to Achieve Results Fellowship (FP 91762601), and EAM by the EU/FP7 project LIBERATION (311781).

References

- Avelino, J., Romero-Gurdian, A., Cruz-Cuellar, H.F., Declerck, F.A.J. 2012. Landscape context and scale differentially impact coffee leaf rust, coffee berry borer, and coffee root-knot nematodes. *Ecol. Appl.* 22, 584-596.
- Bahlai, C.A., Sikkema, S., Hallett, R.H., Newman, J., Schaafsma, A.W. 2010. Modeling distribution and abundance of soybean aphid in soybean fields using measurements from the surrounding landscape. *Env. Entomol.* 39, 50–56.
- Bahlai, C. A., van der Werf, W., O'Neal, M., Hemerik, L., Landis, D.A. 2015. Shifts in dynamic regime of an invasive lady beetle are linked to the invasion and insecticidal management of its prey. *Ecol. Appl.* 25, 1807-1818.

- 1 Batáry, P., Fischer, J., Báldi, A., Crist, T.O., Tscharntke, T. 2011. Does habitat
2 heterogeneity increase farmland biodiversity? *Front. Ecol. Environ.* 9, 152–153.
- 3 Baur, B. 2014. Dispersal-limited species – A challenge for ecological restoration.
4 *Basic Appl. Ecol.* 15, 559–564.
- 5 Benton, T.G., Vickery, J.A., Wilson, J.D. 2003. Farmland biodiversity: is habitat
6 heterogeneity the key? *Trends Ecol. Evol.* 18, 182–88.
- 7 Bianchi, F.J.J.A., Booij, C.J.H., Tscharntke, T. 2006. Sustainable pest regulation in
8 agricultural landscapes: a review on landscape composition, biodiversity and
9 natural pest control. *Proc. Roy. Soc. B* 273, 1715-1727.
- 10 Bianchi, F.J.J.A., Ives, A.R., Schellhorn, N.A. 2013. Interactions between
11 conventional and organic farming for biocontrol services across the landscape.
12 *Ecol. Appl.* 23, 1531-1543.
- 13 Bianchi, F.J.J.A., Schellhorn, N.A., Buckley, Y.M., Possingham, H.P. 2010. Spatial
14 variability in ecosystem services: simple rules for predator-mediated pest
15 suppression. *Ecol. Appl.* 20, 2322-2333.
- 16 Bianchi, F.J.J.A., van der Werf, W. 2003. The effect of the area and configuration of
17 hibernation sites on the control of aphids by *Coccinella septempunctata*
18 (Coleoptera: Coccinellidae) in agricultural landscapes: a simulation study. *Env.*
19 *Entomol.* 32, 1290-1304.
- 20 Blitzer, E.J., Dormann, C.F., Holzschuh, A., Klein, A.M., Rand, T.A., Tscharntke, T.
21 2012. Spillover of functionally important organisms between managed and natural
22 habitats. *Agric. Ecosyst. Environ.* 146, 34–43.
- 23 Campbell, A.J., Biesmeijer, J.C., Varma, V., Wäckers, F.L. 2012. Realising multiple
24 ecosystem services based on the response of three beneficial insect groups to
25 floral traits and trait diversity. *Basic Appl. Ecol.*, 13, 363-370.
- 26 Carrière, Y., Ellsworth, P.C., Dutilleul, P., Ellers-Kirk, C., Barkley, V., Antilla, L. 2006.
27 A GIS-based approach for areawide pest management: the scales of *Lygus*
28 *hesperus* movements to cotton from alfalfa, weeds, and cotton. *Entomol. Exp.*
29 *Appl.* 118, 203–210.
- 30 Carrière, Y., Goodell, P.B., Ellers-Kirk, C., Larocque, G., Dutilleul, P. et al. 2012.
31 Effects of local and landscape factors on population dynamics of a cotton pest.
32 *PLoS ONE* 7, e39862.

- 1 Carvalheiro, L.G., Seymour, C.L., Veldtman, R., Nicolson, S.W. 2010. Pollination
2 services decline with distance from natural habitat even in biodiversity-rich areas
3 J. Appl. Ecol. 47, 810–820.
- 4 Chaplin-Kramer, R., Kliebenstein, D.J., Chiem, A., Morrill, E., Mills, N.J., Kremen, C.
5 2011b. Chemically-mediated tritrophic interactions: opposing effects of
6 glucosinolates on a specialist herbivore and its predators. J. Appl. Ecol. 48, 880–
7 887.
- 8 Chaplin-Kramer, R., O'Rourke, M.E., Blitzer, E.J., Kremen, C. 2011a. A
9 meta-analysis of crop pest and natural enemy response to landscape complexity
10 Ecol. Letters 14, 922-932.
- 11 Corbett, A., Rosenheim, J.A. 1996. Impact of a natural enemy overwintering refuge
12 and its interaction with the surrounding landscape. Ecol. Entomol. 21,155-164
- 13 Cook, S.M., Khan, Z.R., Pickett, S.A. 2007. The use of push–pull strategies in
14 integrated pest management. Annu. Rev. Entomol. 52, 375–400.
- 15 Costamagna, A.C., Venables, W.N., Schellhorn, N.A. 2015. Landscape-scale pest
16 suppression is mediated by timing of predator arrival. Ecol. Appl. 25,1114-1130.
- 17 Daugherty M.P., Gruber, B.R., Almeida, R.P.P. et al. 2012. Testing the Efficacy of
18 Barrier Plantings for Limiting Sharpshooter Spread. American J. Enol. Vitic. 63,
19 139-143.
- 20 Davey, J.S., Vaughan, I.P., Andrew King, R., Bell, J.R., Bohan, D.A., Bruford, M. W.,
21 Symondson, W.O. 2013. Intraguild predation in winter wheat: prey choice by a
22 common epigeal carabid consuming spiders. J. Appl. Ecol., 50, 271-279.
- 23 De Barro, P.J., Coombs, M.T. 2009. Post-release evaluation of *Eretmocerus hayati*
24 Zolnerowich and Rose in Australia. Bull. Ent. Res. 99,193-206.
- 25 Denoth, M., Frid, L., Myers, J.H. 2002. Multiple agents in biological control: improving
26 the odds?. Biol. Control 24, 20-30.
- 27 Denys, C., Tscharntke, T. 2002. Plant-insect communities and predator-prey ratios in
28 field margin strips, adjacent crop fields, and fallows. Oecologia 130, 315-324.
- 29 Dreyer, J., Gratton, C. 2014. Habitat linkages in conservation biological control:
30 Lessons from the land–water interface. Biol. Control 75, 68–76.
- 31 Farrow, R.A. 1974. Comparative plague dynamics of tropic *Locusta* (Orthoptera,
32 Acrididae). Bull. Ent. Res. 64, 401-411.
- 33 Feder, G. 1980. Farm size, risk aversion and the adoption of new technology under
34 uncertainty. Oxford Economic Papers 32, 263-283.

- 1 Fischer, C., Schlinkert, H., Ludwig, M., Holzschuh, A., Gallé, R., Tschardtke, T.,
2 Batáry, P. 2013. The impact of hedge-forest connectivity and microhabitat
3 conditions on spider and carabid beetle assemblages in agricultural landscapes. *J.*
4 *Insect Conserv.* 17, 1027-1038.
- 5 Foley, J.A., Ramankutty, N., Brauman, K.A., Cassidy, E.S., Gerber, J.S., Johnston,
6 M., *et al.* 2011. Solutions for a cultivated planet. *Nature* 478, 337-342.
- 7 Gabriel, D., Sait, S.M., Hodgson, J.A., Schmutz, U., Kunin, W.E., Benton T.G. 2010.
8 Scale matters: the impact of organic farming on biodiversity at different spatial
9 scales. *Ecol. Letters* 13, 858-869.
- 10 Gaines, H.R., Gratton, C. 2010. Seed predation increases with ground beetle
11 diversity in a Wisconsin (USA) potato agroecosystem. *Agr. Ecosyst. Environ.* 137,
12 329–336.
- 13 Gardiner, M.M., Landis, D.A., Gratton, C., Schmidt, N., O’Neal, M., Mueller, E.,
14 Chacon, J., Heimpel, G.E., DiFonzo, C.D. 2009. Landscape composition
15 influences patterns of native and exotic lady beetle abundance. *Divers. Distrib.* 15,
16 554–564.
- 17 Geiger, F., de Snoo, G.R., Berendse, F., *et al.* 2010. Persistent negative effects of
18 pesticides on biodiversity and biological control potential on European farmland.
19 *Basic Appl. Ecol.* 11, 97–105.
- 20 Haddad, N.M., Brudvig, L.A., Clobert, J., Davies, K.F., Gonzalez, A., Holt, R.D., *et al.*
21 2015. Habitat fragmentation and its lasting impact on Earth’s ecosystems. *Science*
22 *Advances* 1, e1500052, DOI: 10.1126/sciadv.1500052.
- 23 Hatcher, P., Paul, N.D. 2001. Plant pathogen herbivore interactions and their effects
24 on weeds. In: *Biotic Interactions in Plant-Pathogen Associations*, pp. 193–225.
- 25 Hawkins, B.A. 1994. *Pattern and Process in Host-Parasitoid Interactions*. Cambridge
26 University Press, Cambridge.
- 27 Hawkins, B.A., Cornell, H.V. 1994. Maximum parasitism rates and successful
28 biological control. *Science* 266, 1886.
- 29 Hawkins, B. A., H. V. Cornell, and M. E. Hochberg. 1997. Predators, parasitoids, and
30 pathogens as mortality agents in phytophagous insect populations. *Ecology*
31 78:2145-2152.
- 32 Heimpel, G.E., Frelich, L.E., Landis, D.A., Hopper, K.R., Hoelmer, K.A., Sezen, Z., *et*
33 *al.* 2010. European buckthorn and Asian soybean aphid as components of an
34 extensive invasional meltdown in North America. *Biol. Invas.* 12, 2913-2931.

- 1 Hendrickx, F., Maelfait, J., van Wingerden, W., Schweiger, O., Speelmans, M.,
2 Aviron, S., et al. (2007). How landscape structure, land-use intensity and habitat
3 diversity affect components of total arthropod diversity in agricultural landscapes.
4 J. Appl. Ecol. 44, 340–351.
- 5 Henri, D., Jones, O., Tsiattalos, A., Thébault, E., Seymour, C. L., & Veen, F. J. 2015.
6 Natural vegetation benefits synergistic control of the three main insect and
7 pathogen pests of a fruit crop in southern Africa. J. Appl. Ecol. (online).
- 8 Holland, J.M., Bianchi, F.J., Entling, M.H., Moonen, A.C., Smith, B.M., Jeanneret, P.
9 2016. Structure, function and management of semi-natural habitats for
10 conservation biological control: a review of European studies. Pest Manag. Sci.
11 72, 1638-1651.
- 12 Holzschuh, A., Steffan-Dewenter, I., Tscharnkte, T. 2010. How do landscape
13 composition and configuration, organic farming and fallow strips affect the diversity
14 of bees, wasps and their parasitoids? J. Anim. Ecol. 79, 491–500.
- 15 Hough-Goldstein, J.A., Heimpel, G.E., Bechmann, H.E., Mason, C.E. 1993.
16 Arthropod natural enemies of the Colorado potato beetle. Crop Protection 12,
17 324–334.
- 18 Iverson, A.L., Marín, L.E., Ennis, K.K., Gonthier, D.J., Connor-Barrie, B.T., Remfert,
19 J.L., et al. 2014. REVIEW: Do polycultures promote win-wins or trade-offs in
20 agricultural ecosystem services? A meta-analysis. J. Appl. Ecol., 51, 1593-1602.
- 21 Ives, A.R., Settle, W.H. 1997. Metapopulation dynamics and pest control in
22 agricultural systems. Am. Natur. 149, 220-246.
- 23 Jonsson, M., Buckley, H.L., Case, B.S., Wratten, S.D., Hale, R.J., Didham, R.K.
24 2012. Agricultural intensification drives landscape-context effects on host–
25 parasitoid interactions in agroecosystems. J. Appl. Ecol. 49: 706-714.
- 26 Just, Zilberman 1983. Estimation of multicrop production functions. American J.
27 Agric. Econ. 65, 770-780.
- 28 Karp, D.S., Baur, P., Atwill, E.R., Master, K.D.E., Gennet, S., Iles, A., et al. (2015a).
29 The unintended ecological and social impacts of food safety regulations in
30 California's Central Coast region. *Bioscience*, 65, 1173–1183.
- 31 Karp, D.S., Chase, D.M., Figueroa Sandí, R., Chaumont, N., Ehrlich, P.R., Hadly,
32 E.A., Daily, G.C. 2013. Forest bolsters bird abundance, pest control and coffee
33 yield. Ecol. Letters 16, 1339-1347.

- 1 Karp, D.S., Daily, G.C. 2014. Cascading effects of insectivorous birds and bats in
2 tropical coffee plantations. *Ecology* 95, 1065-1074.
- 3 Karp, D.S., Gennet, S., Kilonzo, C., Partyka, M., Chaumont, N., Atwill, E.R., Kremen,
4 C. 2015b. Comanaging fresh produce for nature conservation and food safety.
5 *Proc. Natl. Acad. Sci. USA* 112, 11126-11131.
- 6 Keller, S., Häni, F. 2000. Ansprüche von Nützlingen und Schädlingen an den
7 Lebensraum. In: Streifenförmige ökologische Ausgleichsflächen in der
8 Kulturlandschaft: Ackerkrautstreifen, Buntbrache, Feldränder. Verlag
9 Agrarökologie, Bern, 199-217.
- 10 Kleijn, D., Rundlöf, M., Scheper, J., Smith, H.G., Tscharntke, T. 2011. Does
11 conservation on farmland contribute to halting the biodiversity decline? *Trends*
12 *Ecol. Evol.* 26, 474-481.
- 13 Klein, A.-M., Steffan-Dewenter, I., Tscharntke, T. 2003. Pollination of *Coffea*
14 *canephora* in relation to local and regional agroforestry management. *J. Appl.*
15 *Ecol.* 40, 837-845.
- 16 Klein, A.-M., Vaissière, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A.,
17 Kremen, C., Tscharntke, T. 2007. Importance of pollinators in changing
18 landscapes for world crops. *Proc. Roy. Soc. B* 274, 303-313.
- 19 Landis, D.A., Wratten, S.D., Gurr, G.M. 2000. Habitat management to conserve
20 natural enemies of arthropod pests in agriculture. *Ann. Rev. Entomol.* 45, 175–
21 201.
- 22 Landis, D.A., Gardiner, M.M., van der Werf, W. Swinton, S.M. 2008. Increasing corn
23 for biofuel production reduces biocontrol services in agricultural landscapes. *PNAS*
24 105, 20552-20557.
- 25 Lavandero, B., Wratten, S.D., Didham, R.K., Gurr, G. 2006. Increasing floral diversity
26 for selective enhancement of biological control agents: a double-edged sword?
27 *Basic Appl. Ecol.*, 7, 236-243.
- 28 Letourneau, D.K., Armbrrecht, I., Rivera, B.S., Lerma, J.M., Carmona, E.J., Daza,
29 M.C., et al. 2011. Does plant diversity benefit agroecosystems? A synthetic
30 review. *Ecol. Appl.* 21, 9–21.
- 31 Levine, E., Oloumi-Sadeghi, H. 1991. "Management of Diabroticite Rootworms in
32 Corn." *Ann. Rev. Entomol.* 36, 229-255.
- 33 Lomer, C., Bateman, R., Johnson, D., Langewald, J., Thomas, M. 2001. Biological
34 control of locusts and grasshoppers. *Ann. Rev. Entomol.* 1, 667–702

- 1 Lu, Y., Wu, K., Jiang, Y., Guo, Y., Desneux, N. 2012.. Widespread adoption of Bt
2 cotton and insecticide decrease promotes biocontrol services. *Nature* 487, 362–
3 365.
- 4 Lu, Y., Wu, K., Jiang, Y., Xia, B., Li, B., Feng, H., Wyckhuys, K.A.G., Guo, Y. 2010.
5 “Mirid Bug Outbreaks in Multiple Crops Correlated with Wide-Scale Adoption of Bt
6 Cotton in China.” *Science* 328, 1151-1154.
- 7 Lu, Y.H., Wu, K.M. 2008. *Biology and control of cotton mirids*. Golden Shield Press,
8 Beijing, China.
- 9 Lundgren, J.G. 2009. *Relationships of natural enemies and non-prey foods* (Vol. 7).
10 Springer Science & Business Media.
- 11 Lundgren, J.G., Fergen, J.K. 2011. Enhancing predation of a subterranean insect
12 pest: a conservation benefit of winter vegetation in agroecosystems. *Appl. Soil*
13 *Ecol.* 51, 9-16.
- 14 Lys, J.A., Zimmermann, M., Nentwig, W. 1994. Increase in activity density and
15 species number of carabid beetles in cereals as a result of strip-management.
16 *Entomol. Exp. Appl.* 73, 1-9.
- 17 Maas, B., Clough, Y., Tscharntke, T. 2013. Bats and birds increase crop yield in
18 tropical agroforestry landscapes. *Ecol. Letters* 16, 1480-1487.
- 19 Mace, G.M., Norris, K., Fitter, A.H. 2012. Biodiversity and ecosystem services: a
20 multilayered relationship. *Trends Ecol. Evol.* 27, 19–26.
- 21 Macfadyen, S., Cunningham, S.A., Costamagna, A.C., Schellhorn, N.A. 2012.
22 Managing ecosystem services and biodiversity conservation in agricultural
23 landscapes: are the solutions the same? *J. Appl. Ecol.* 49, 690–694
- 24 Mäder, P., Fliessbach, A., Dubois, D., Gunst, L., Fried, P., Niggli, U. 2002. Soil
25 fertility and biodiversity in organic farming. *Science* 296, 1694-1697.
- 26 Martin, E.A., Seo, B., Park, C.-R., Reineking, B., Steffan-Dewenter, I. 2016. Scale-
27 dependent effects of landscape composition and configuration on natural enemy
28 diversity, crop herbivory and yields. *Ecol. Appl.* 26, 448-462.
- 29 Martin, E.A., Reineking, B., Seo, B., Steffan-Dewenter, I. 2013. Natural enemy
30 interactions constrain pest control in complex agricultural landscapes. *Proc. Nat.*
31 *Acad. Sci. USA* 110, 5534-5539.
- 32 McIntyre, S., Hobbs, R. 1999. A framework for conceptualizing human effects on
33 landscapes and its relevance to management and research models. *Conserv. Biol.*
34 13, 1282-1292.

- 1 McKenzie, A.J., Emery, S.B., Franks, J.R., Whittingham, M.J. 2013. Landscape-scale
2 conservation: collaborative agri-environment schemes could benefit both
3 biodiversity and ecosystem services, but will farmers be willing to participate? J.
4 Appl. Ecol. 50, 1274–1280.
- 5 Meehan, T.D., Werling, B.P., Landis, D.A., Gratton, C. 2011. Agricultural landscape
6 simplification and insecticide use in the Midwestern U.S. PNAS 108, 11500-11505.
- 7 Meisner, M.H., Harmon, J.P., Ives, A.R. 2014. Temperature effects on long-term
8 population dynamics in a parasitoid-host system. Ecol. Monogr. 84, 457-476.
- 9 Messing, R.H., Wright, M.G. 2006. Biological control of invasive species: solution or
10 pollution? Front. Ecol. Environ. 4, 132-140.
- 11 Midega, C.A.O., Jonsson, M., Khan Z.R., Ekbom, B. 2014. Effects of landscape
12 complexity and habitat management on stemborer colonization, parasitism and
13 damage to maize. Agric. Ecosyst. Environ. 188, 289–293.
- 14 Milligan, M.C., Johnson, M.D., Garfinkel, M., Smith, C.J. Njoroge, P. 2016.
15 Quantifying pest control services by birds and ants in Kenyan coffee farms. Biol.
16 Conserv. 194, 58-65.
- 17 Mueller, E.E., Frost, K.E., Esker, P.D., Gratton, C. 2010. Seasonal phenology of
18 *Aphis glycines* (Hemiptera: Aphididae) and other aphid species in cultivated bean
19 and noncrop habitats in Wisconsin. J. Econ. Entomol. 103, 1670-1681.
- 20 Murphy, B.C., Rosenheim, J.A., Granett, J. 1996. Habitat diversification for improving
21 biological control: abundance of *Anagrus epos* (Hymenoptera: Mymaridae) in
22 grape vineyards. Environ. Entomol. 25, 495-504.
- 23 Nilsson, C. 2010. Impact of soil tillage on parasitoids of oilseed rape pests. In: Ed.
24 Williams I.H., Biocontrol-based Integrated Management of Oilseed Rape Pests.
25 Springer. DOI: 10.1007/978-90-481-3983-5_11
- 26 Norris, R., Kogan, M. 2000. Interactions between weeds, arthropod pests, and
27 their natural enemies in managed ecosystems. Weed Science 48, 94–158.
- 28 O'Rourke, M.E., Rienzo-Stack, K., Power, A.G. 2011. A multi-scale, landscape
29 approach to predicting insect populations in agroecosystems. Ecol. Appl., 21,
30 1782-1791.
- 31 Östman, Ö., Ekbom, B., Bengtsson, J. 2001. Landscape heterogeneity and farming
32 practice influence biological control. Basic Appl. Ecol. 2, 365–371.
- 33 Parry, H.R., Macfadyen, S., Hopkinson, J.E., Bianchi, F.J.J.A., Zalucki, M.P., Bourne,
34 A., Schellhorn, N.A. 2015. Plant composition modulates arthropod pest and

- predator abundance: Evidence for culling exotics and planting natives. *Basic Appl. Ecol.* 16, 531-543.
- Perfecto, I. 1990. Indirect and Direct Effects in a Tropical Agroecosystem: The Maize-Pest-Ant System in Nicaragua. *Ecology* 71, 2125-2134.
- Perfecto, I., Vandermeer, J. 2010. The agricultural matrix as an alternative to the land-sparing/agricultural intensification model: facing the food and biodiversity crises. *Proc. Nat. Acad. Science* 107, 5786–5791.
- Perović, D.J., Gurr, G.M., Raman, A., Nicol, H.I. 2010. Effect of landscape composition and arrangement on biological control agents in a simplified agricultural system: a cost–distance approach. *Biol. Contr.* 52, 263-270.
- Poveda, K., Gómez, M.I., Martínez, E. 2008. Diversification practices: their effect on pest regulation and production. *Rev. Colomb. Entomol.* 34, 131-144.
- Poveda, K., Martínez, E., Kersch-Becker, M.F., Bonilla, M.A., Tschardtke, T. 2012. Landscape simplification and altitude affect biodiversity, herbivory and Andean potato yield. *J. Appl. Ecol.* 49, 513–522.
- Power, A.G., Mitchell, C.E. 2004. Pathogen Spillover in Disease Epidemics. *The Am. Nat.* 164, S79–S89. doi:10.1086/424610
- Pywell, R.F., Heard, M.S., Woodcock, B.A., Hinsley, S., Ridding, L., Nowakowski, M., Bullock, J.M. 2015. Wildlife-friendly farming increases crop yield: evidence for ecological intensification. *Proc. R. Soc. B* 282, 2015-1740.
- Rand, T.A., Tylianakis, J.M., Tschardtke, T. 2006. Spillover edge effects: the dispersal of agriculturally subsidized insect natural enemies into adjacent natural habitats. *Ecol. Letters* 9, 603-614.
- Raudsepp-Hearne, C., Peterson, G.D., Bennett, E.M. 2010. Ecosystem service bundles for analyzing tradeoffs in diverse landscapes. *Proc. Nat. Acad. Sci.* 107, 5242-5247.
- Riechers, M., Barkmann, J., Tschardtke, T. 2016. Perceptions of cultural ecosystem services from urban green. *Ecosyst. Serv.* 17: 33-39.
- Rusch, A., Valantin-Morison, M., Sarthou, J.P., Roger-Estrade, J. 2010. Biological Control of Insect Pests in Agroecosystems: Effects of Crop Management, Farming Systems, and Seminal Habitats at the Landscape Scale: A Review. *Adv. Agron.* 109, 219.

- 1 Rusch, A., Chaplin-Kramer, R., Gardiner, M.M., Hawro, V., Holland, J., Landis, D., et
2 al. 2016. Agricultural landscape simplification reduces natural pest control: A
3 quantitative synthesis. *Agric. Ecosyst. Environ.* 221, 198-204.
- 4 Rusch, A., Valantin-Morison, M., Roger-Estrade, J., Sarthou, J.P. 2012. Using
5 landscape indicators to predict high pest infestations and successful natural pest
6 control at the regional scale. *Landsc. Urban Plann.* 105, 62–73.
- 7 Rusch, A., Valantin-Morison, M., Sarthou, J.P., Roger-Estrade, J. 2011. Multi-scale
8 effects of landscape complexity and crop management on pollen beetle parasitism
9 rate. *Landsc. Ecol.*, DOI: 10.1007/s10980-011-9573-7
- 10 Rusch, A., Bommarco, R., Chiverton, P., Öberg, S., Wallin, H., Wikteliuss, S., Ekbom,
11 B. 2013a. Response of ground beetle (Coleoptera, Carabidae) communities to
12 changes in agricultural policies in Sweden over two decades. *Agric. Ecosyst.*
13 *Environ.* 176, 63-69.
- 14 Rusch, A., Bommarco, R., Jonsson, M., Smith, H.G., Ekbom, B. 2013b. Flow and
15 stability of natural pest control services depend on landscape complexity and crop
16 rotation in the landscape. *J. Appl. Ecol.* 50, 345-354.
- 17 Rusch, A., Binet, D., Delbac, L., Thiéry, D., 2016. Local and landscape effects of
18 agricultural intensification on Carabid community structure and weed seed
19 predation in a perennial cropping system. *Landscape Ecol* 1–12.
- 20 Sawada, H., Subroto, S.W.G., Mustaghfirin, G., Wuaya, E.S. 1992. Immigration,
21 population development and outbreaks of brown planthopper, *Nilaparvata lugens*
22 (Stal) under different rice cultivation patterns in central Java, Indonesia. *Technical*
23 *Bulletin -- Food and Fertilizer Technology Center* 130, 9-13.
- 24 Schellhorn, N.A., Bianchi, F.J.J.A., Hsu, C.L. 2014. Movement of entomophagous
25 arthropods in agricultural landscapes: links to pest suppression. *Ann. Rev.*
26 *Entomol.* 59, 559-581.
- 27 Schellhorn, N.A., Gagic, V., Bommarco, R. 2015. Time will tell: resource continuity
28 bolsters ecosystem services. *Trends Ecol. Evol.* 30, 524–530.
- 29 Schellhorn, N.A., Glatz, R.V., Wood, G.M. 2010. The risk of exotic and native plants
30 as hosts for four pest thrips (Thysanoptera: Thripinae). *Bull. Ent. Res.* 100, 501-
31 510.
- 32 Scherber, C., Eisenhauer, N., Weisser, W.W., Schmid, B., Voigt, W., Fischer, M., et
33 al. 2010. Bottom-up effects of plant diversity on multitrophic interactions in a
34 biodiversity experiment. *Nature* 468, 553-556.

- 1 Segoli, M., Rosenheim, J.A. 2012. Should increasing the field size of monocultural
2 crops be expected to exacerbate pest damage? *Agr. Ecosyst. Environ.* 150, 38-44.
- 3 Sequeira, R., Dixon, A.F.G. 1997. Population dynamics of tree-dwelling aphids: the
4 importance of seasonality and time scale. *Ecology* 78, 2603-2610.
- 5 Settle, W.H., Ariawan, H., Astuti, E.T., Cahyana, W., Hakim, A.L., Hindayana, D.,
6 Lestari, A.S. 1996. Managing tropical rice pests through conservation of generalist
7 natural enemies and alternative prey. *Ecology* 1975-1988.
- 8 Sexton, S.E., Lei, Z., Zilberman, D. 2007. The economics of pesticides and pest
9 control. *Intern. Rev. Environm. Res. Econom.* 1, 271-326.
- 10 Settle, W.H., Ariawan, H., Tri Asuti, E., Cahyana, W., Hakim, A.L., Hindayana, D., Sri
11 Lestari, A., Sartano, P. 1996. Managing tropical rice pests through conservation of
12 generalist natural enemies and alternative prey. *Ecology* 77, 1975–1988.
- 13 Sequeira, R., Dixon, A.F.G. 1997. Population dynamics of tree-dwelling aphids: the
14 importance of seasonality and time scale. *Ecology* 78, 2603-2610.
- 15 Shackelford, G., Steward, P.R., Benton, T.G., Kunin, W.E., Potts, S.G., Biesmeijer,
16 J.C., Sait, S.M. 2013. Comparison of pollinators and natural enemies: a
17 meta-analysis of landscape and local effects on abundance and richness in crops.
18 *Biol. Rev.* 88, 1002-1021.
- 19 Sivakoff, F.J., Rosenheim, J.A., Dutilleul, P., Carrière, Y. 2013. Influence of the
20 surrounding landscape on crop colonization by an insect pest. *Entomol. Exp. Appl.*
21 149, 11-21.
- 22 Strong, D. R. (1974). Rapid asymptotic species accumulation in phytophagous insect
23 communities: the pests of cacao. *Science*, 185(4156), 1064-1066.
- 24 Sutton, M.A., van Grinsven, H. 2011. *The European Nitrogen Assessment*.
25 Cambridge University Press, Cambridge.
- 26 Symondson, W. O. C., Sunderland, K. D., & Greenstone, M. H. (2002). Can
27 generalist predators be effective biocontrol agents? 1. Annual review of
28 entomology, 47(1), 561-594.
- 29 Thies, C., Roschewitz, I., Tscharnkte, T. 2005. The landscape context of cereal
30 aphid-parasitoid interactions. *Proc. Roy. Soc. B* 272, 203-210.
- 31 Thies, C., Tscharnkte, T. 1999. Landscape structure and biological control in
32 agroecosystems. *Science* 285, 893–95

- 1 Thies, C., Haenke, S., Scherber, C., Bengtsson, J., Bommarco, R., Clement, L.W., et
2 al. 2011. The relationship between agricultural intensification and biological control
3 – experimental tests across Europe. *Ecol. Appl.* 21, 2187–2196.
- 4 Thorbek, P., Bilde, T. 2004. Reduced numbers of generalist arthropod predators after
5 crop management. *J. Appl. Ecol.* 41, 526-538.
- 6 Traugott, M. Bell, J.R. Raso, L. Sint, D. Symondson, W.O.C. 2012. Generalist
7 predators disrupt parasitoid aphid control by direct and coincidental intraguild
8 predation. *Bull. Entomol. Res.* 102, 239 – 247.
- 9 Tscharntke, T. 1992. Cascade effects among four trophic levels: Bird predation on
10 galls affects density-dependent parasitism. *Ecology* 73, 1689-1698.
- 11 Tscharntke, T. 1997. Vertebrate effects on plant-invertebrate food webs. In: Gange
12 A.C., Brown V.K. (eds): *Multitrophic interactions in terrestrial systems*. Blackwell
13 Science Ltd. (36th Symposium of The British Ecological Society), 277-297.
- 14 Tscharntke, T., Batáry, P., Dormann, C.F. 2011a Set-aside management: How do
15 succession, sowing patterns and landscape context affect biodiversity? *Agric.*
16 *Ecosyst. Environ.* 143, 37– 44.
- 17 Tscharntke, T., Bommarco, R., Clough, Y., Crist, T.O., Kleijn, D., Rand, T.A., et al.
18 2007a. Conservation biological control and enemy diversity on a landscape scale.
19 *Biol. Control* 43, 294-309.
- 20 Tscharntke, T., Clough, Y., Bhagwat, S.A., Buchori, D., Faust, H., Hertel, D., et al.
21 2011b. Multifunctional shade-tree management in tropical agroforestry landscapes
22 – a review. *J. Appl. Ecol.* 48, 619-629.
- 23 Tscharntke, T., Clough, Y., Wanger, T.C., Jackson, L., Motzke, I., Perfecto, I., et al.
24 2012a. Global food security, biodiversity conservation and the future of agricultural
25 intensification. *Biol. Conserv.* 151, 53-59.
- 26 Tscharntke, T., Hochberg, M.E., Rand, T.A., Resh, V.H., Krauss, J. 2007b. Author
27 sequence and credit for contributions in multiauthored publications. *PLoS Biology*
28 5, 13–14.
- 29 Tscharntke, T., Klein, A.M., Kruess, A., Steffan-Dewenter, I., Thies, C. 2005a.
30 Landscape perspectives on agricultural intensification and biodiversity—
31 ecosystem service management. *Ecol. Lett.* 8, 857–74
- 32 Tscharntke, T., Milder, J.C., Schroth, G., Clough, Y., DeClerck, F., Waldron, A., Rice,
33 R., Ghazoul, J. 2015. Conserving biodiversity through certification of tropical
34 agroforestry crops at local and landscape scales. *Conserv. Letters* 8, 14-23.

- 1 Tscharntke, T., Rand, T., Bianchi, F. 2005b. The landscape context of trophic
2 interactions: insect spillover across the crop-noncrop interface. *Ann. Zool. Fenn.*,
3 42, 421–432.
- 4 Tscharntke, T., Sekercioglu, C.H., Dietsch, T.V., Sodhi, N.S., Hoehn, P., Tylianakis,
5 J.M. 2008. Landscape constraints on functional diversity of birds and insects in
6 tropical agroecosystems. *Ecology* 89, 944-951.
- 7 Tscharntke, T., Tylianakis, J.M., Rand, T.A., et al. 2012b. Landscape moderation of
8 biodiversity patterns and processes – eight hypotheses. *Biol. Rev.* 87, 661-685.
- 9 Tschumi, M., Albrecht, M., Bärtschi, C., Collatz, J., Entling, M.H., Jacot, K., 2016.
10 Perennial, species-rich wildflower strips enhance pest control and crop yield.
11 *Agriculture, Ecosystems & Environment* 220, 97–103.
- 12 Tylianakis, J.M., Didham, R.K., Wratten, S.D. 2004. Improved fitness of aphid
13 parasitoids receiving resource subsidies. *Ecology* 85, 658–666.
- 14 Tylianakis, J.M., Tscharntke, T., Klein, A.-M. 2006. Diversity, ecosystem function, and
15 stability of parasitoid-host interactions across a tropical gradient of habitat
16 gradient. *Ecology* 87, 3047-3057.
- 17 van Driesche, R., Hoddle, M., Center, T. 2009. *Control of Pests and Weeds by*
18 *Natural Enemies: An Introduction to Biological Control*. John Wiley & Sons.
- 19 van Zanten, B.T., Verburg, P.H., Koetse, M.J., van Beukering, P.J. 2014. Preferences
20 for European agrarian landscapes: A meta-analysis of case studies. *Landsc.*
21 *Urban Plann.* 132, 89-101.
- 22 Wanger, T.C., Darras, K., Bumrungsri, S., Tscharntke, T., Klein, A. M. 2014. Bat pest
23 control contributes to food security in Thailand. *Biol. Conserv.* 171, 220-223.
- 24 Way, M.J., Heong, K.L. 1994. The role of biodiversity in the dynamics and
25 management of insect pests of tropical irrigated rice - a review. *Bull. Entomol. Res.*
26 84, 567-587.
- 27 Werling, B.P., Meehan, T.D., Gratton, C., Landis, D.A. 2011. Influence of habitat and
28 landscape perennality on insect natural enemies in three candidate biofuel crops.
29 *Biol. Control* 59, 304-312.
- 30 Westphal, C., Vidal, S., Horgan, F.G., Gurr G.M., Escalada M., Chien H.V. et al.
31 2015. Promoting multiple ecosystem services with flower strips and participatory
32 approaches in rice production landscapes. *Basic Appl. Ecol.*
33 <http://dx.doi.org/10.1016/j.baae.2015.10.004>.

- Wielgoss, A., Tscharntke, T., Rumed, A., Fiala, B., Seidel, H., Shahabuddin, S., Clough, Y. 2014. Interaction complexity matters: disentangling services and disservices of ant communities driving yield in tropical agroecosystems. *Proc. R. Soc. B* 281, 20132144; <http://dx.doi.org/10.1098/rspb.2013.2144>.
- Winqvist, C., Bengtsson, J., Aavik, T., Berendse, F., Clement, L.W., Eggers, S., et al. (2011). Mixed effects of organic farming and landscape complexity on farmland biodiversity and biological control potential across Europe. *J. Appl. Ecol.* 48, 570-579.
- Wisler, G.C., Norris, R.F. 2005. Interactions between weeds and cultivated plants as related to management of plant pathogens. *Weed Science* 53, 914-917.
- Wong, C.P., Jiang, B., Kinzig, A.P., Lee, K.N., Ouyang, Z. 2015. Linking ecosystem characteristics to final ecosystem services for public policy. *Ecol. Letters* 18, 108–118
- Zhang, W., Swinton, S.M. 2012. Optimal control of soybean aphid in the presence of natural enemies and the implied value of their ecosystem services. *J. Environ. Manage.* 96, 7-16.

Figure Legends

Figure 1: Five hypotheses for when and why natural habitat can fail to support biological pest control: (a) Pest populations have no effective natural enemies in the region, (b) natural habitat is a greater source of pests than natural enemies, (c) crops provide more important resources for natural enemies than does natural habitat, (d) natural habitat is insufficient in amount, proximity, composition, or configuration to provide enemy populations needed for pest control, and (e) agricultural practices counteract enemy establishment and biocontrol provided by natural habitat.

Table 1: Natural habitat and biological pest control: Five hypotheses, with explanations and references, for when natural habitat does not enhance biocontrol

Name of hypothesis	Explanation	References
(1) Pest populations have no effective natural enemies in the region	Pest density may be driven by factors other than biocontrol, such as environmental conditions, crop susceptibility, agricultural practices, crop area, or intraguild predation of higher trophic levels.	Hough-Goldstein <i>et al.</i> 1983; Martin <i>et al.</i> 2013; Karp <i>et al.</i> 2014; Meisner <i>et al.</i> 2014; O'Rourke <i>et al.</i> , 2011; Poveda <i>et al.</i> 2008
(2) Natural habitat is a greater source of pests than natural enemies	Natural habitats can provide a suitable environment for a large number of pest species at several key stages of their life-cycle, and/or natural enemies may not disperse from natural habitat.	Blitzer <i>et al.</i> 2011; Wisler & Norris 2005; Power & Mitchell 2004; Carrière <i>et al.</i> 2012; Parry <i>et al.</i> 2015; Rusch <i>et al.</i> 2013b; Midega <i>et al.</i> 2014
(3) Crops provide more important resources for natural enemies than does natural habitat	Natural habitat may not always be a panacea for natural enemies (e.g., because of low productivity), which may be more influenced by the surrounding cropland than natural areas.	Rand <i>et al.</i> 2006; Gardiner <i>et al.</i> 2009; Blitzer <i>et al.</i> 2011; Costamagna <i>et al.</i> 2015; Schellhorn <i>et al.</i> 2015
(4) Natural habitat is insufficient in amount, proximity, composition or configuration to provide large enough enemy populations for pest control	To enhance pest control effectively, natural habitats must be both large enough and proximate enough to farm fields to facilitate a substantial increase in within-field enemy abundance.	Segoli & Rosenheim 2012; Dreyer & Gratton 2014; Thies & Tschardtke 1999; Tschardtke <i>et al.</i> 2007
(5) Agricultural practices counteract natural enemy establishment and biocontrol provided by natural habitat	Pesticide spraying, deep ploughing, planting highly susceptible crop varieties and little crop diversity may all negatively affect natural enemies and support pests, even if surrounding natural habitats are present.	Iverson <i>et al.</i> 2014; Geiger <i>et al.</i> 2011; Rusch <i>et al.</i> 2011; Letourneau <i>et al.</i> 2011; Jonsson <i>et al.</i> 2012

Figure 1: Five hypotheses for when and why natural habitat can fail to support biological pest control: (a) Pest populations have no effective natural enemies in the region, (b) natural habitat is a more important source of pests than natural enemies, (c) crops provide more important resources for natural enemies than does natural habitat, (d) natural habitat is insufficient in amount, proximity, composition, or configuration to provide enemy populations needed for pest control, and (e) agricultural practices counteract enemy establishment and biocontrol provided by natural habitat.