1 Title: Identifying overlap between humpback whale foraging grounds and the Antarctic krill fishery.

2 Authors: Ben G. Weinstein¹, Michael Double², Nick Gales², David W. Johnston³, & Ari S. Friedlaender¹

3 ¹Department of Fisheries and Wildlife, Marine Mammal Institute, Oregon State University, 2030 Marine
4 Science Drive, Newport, OR 97365, USA

5 ²Australian Antarctic Division, 203 Channel Highway, Kingston, Tasmania, 7050, Australia

6 ³Duke University Marine Laboratory, 135 Duke Marine Lab Road, Beaufort, NC, 28516, USA

7 Keywords: Cetaceans, Bayesian movement models, Gerlache Strait, CCAMLR, Fisheries Management

8 ABSTRACT

9 The Antarctic krill fishery is the largest in the southern ocean, but currently operates without fine-scale
10 information on whale movement and behavior. Using a multi-year dataset of satellite-tagged whales, as
11 well as information on krill catch levels, we analyzed the spatial distribution of whales and fisheries
12 effort within the small-scale management units defined by the Convention for the Conservation of
13 Antarctic Marine Living Resources (CCAMLR). Using a Bayesian movement model to partition whale
14 movement into traveling and area-restricted search states, we found that both whale behavior and krill
15 catch effort were spatially clustered, with distinct hotspots of the whale activity in the Gerlache and
16 southern Bransfield Straits. These areas align with increases in krill fishing effort, and present potential
17 areas of current and future conflict. We recommend that the Antarctic West and Bransfield Strait West
18 management units merit particular attention when setting fine-scale catch limits and, more broadly,
19 consideration as critical areas for krill predator foraging.

20 INTRODUCTION

21 Minimizing overlap between fishing effort and predators is a persistent challenge in managing marine
22 fisheries. To create ecologically informed guidelines, managers need detailed information on

occurrence, abundance and behavior of predator species. This is especially difficult in polar regions where information can be scarce and observation windows are curtailed due to extreme climate. High-resolution tracking data provides a bridge between species locations and geographically dependent behaviors (Hays et al., 2016). By incorporating behavior into conservation action, we gain greater insight into the threats animals face, and increase the likelihood that management plans achieve positive conservation goals (Ellison et al., 2012; Trathan et al., 2015).

The Antarctic krill (*Euphausia superba*) is the keystone species of the Antarctic ecosystem, providing the primary food source for a diverse group of predators including fish, penguins, seals, and whales (Hill et al., 2006). Antarctic krill form dense swarms, sometimes in excess of 10,000 individuals per cubic meter, extending 200-300m in depth and several kilometers wide (Tarling et al. 2009, Nowacek et al. 2011, Espinasse et al. 2012). The Antarctic Krill fishery is the largest in the Southern Ocean with a reported total catch of 293,815 metric tonnes (mt) in 2015. This catch has fluctuated greatly, with a peak of nearly 500,000 mt in the 1980s, to a low of less than 100,000 mt in the early 2000s, to a rapid increase to 300,000 mt since 2009 (Nicol et al., 2012). This rise is due to the expansion of the fishery by new nations, as well as an extension of fishing effort into austral autumn and winter months (CCAMLR, 2015).

The krill fishery is managed by the Commission for the Conservation of Marine Living Resources (CCAMLR). CCAMLR was formed in 1982 as part of the Antarctic treaty system with the goal of 'providing maintenance of the ecological relationships between harvested, dependent and related populations of Antarctic marine resources' (Constable et al., 2000). The CCAMLR management guidelines require that the krill fishery not interfere with the population growth of Antarctic krill predators (Kawaguchi et al., 2006). However, the management of the krill fishery has not assessed the needs and behavior of baleen whales, which are the largest krill predators in the Antarctic.

The krill fishery historically included large expanses of Antarctic waters, but is currently concentrated in the South Atlantic (CCAMLR Area 48). Records from 2015 report 146,191 mt taken from the Antarctic Peninsula and South Shetlands Islands (Subarea 48.1), 72,455 mt from the South Orkneys Islands (Subarea 48.2), and 75,169 mt from South Georgia Island (Subarea 48.3) (CCAMLR 2015). The current precautionary annual catch limit of 5.61 million mt is shared among these subareas, and is substantially more than the total annual catch of the current fishery (Nicol et al., 2012). The current limit is described as ‘precautionary’, meaning that the level would have low risk of long-term adverse effects on the krill stock and the surrounding ecosystem (Nicol and Foster, 2016).

Our study focuses solely on Subarea 48.1 (Fig. 1), an area of increasing importance to the fishery. In 1993, CCAMLR introduced ‘trigger levels’ for each subarea, which could not be exceeded until CCAMLR had distributed the precautionary catch limit at finer spatial scales. In 2009, CCAMLR adopted an interim measure to distribute the trigger level across subareas, with a limit of 155,000 mt for subarea 48.1. This level has been reached four times since 2010, with each event leading to the closure of the subarea for the remainder of the season (Nicol and Foster, 2016). CCAMLR has attempted to manage the spatial distribution of fishing effort by agreeing upon small-scale management units (SSMUs; Fig. 1). However, to date, CCAMLR has failed to agree on a method to distribute the total precautionary catch limit between SSMUs (Hewitt et al., 2004). Our aim is to determine the overlap between the krill fishery and areas of whale foraging, in order to recommend whether specific small-scale management units should have restrictions in fishing effort or duration.

Since the cessation of commercial whaling in the Southern Hemisphere in the late 20th century, humpback whales (*Megaptera novaeangliae*) have recovered to become the most numerous whale species in the region (Clapham et al., 1999; Herr et al., 2016; Matsuoka et al., 2006). Previous work has shown that humpback distribution is related to the distribution and abundance of krill (Friedlaender et al., 2006; Nowacek et al., 2011). At the fine-scale, the foraging behavior of humpback whales is affected

by the depth and density of krill patches, due to the energetic demands of whale foraging and life-history (Friedlaender et al., 2013; Tyson et al., 2016). Humpbacks whales may therefore be vulnerable to disturbance from the krill fishery due to their reliance on krill as a primary food source (Nicol et al., 2008). Current information on the foraging behavior of Antarctic humpback whales comes largely from short-term tagging efforts (e.g. Friedlaender et al., 2013; Johnston et al., 2012), and from ship-based surveys (Hedley et al., 2001; Herr et al., 2016; Santora et al., 2010). We used a large multi-year dataset (>40,000 Argos locations) of tagged humpbacks to assess the space use and behavior within CCAMLR small-scale management units. Our goals are to 1) describe the areas of humpback presence in reference to the CCAMLR small-scale management units, 2) partition movement into traveling and area-restricted search states, and 3) compare the areas of whale behavior with the distribution of krill fishery activity along the Antarctic Peninsula.

METHODS

Satellite Tagging and Tracking

We deployed Wildlife Computers (Redmond, WA, USA) SPOT5 Platform Transmitting Terminals (PTTs) in 2012, 2013, 2015, and 2016 (Table S1). Each tag is contained in a sterilized housing designed to penetrate the whale's skin and blubber up to 290mm, and is anchored in the tissue beneath the blubber with stainless steel barbs, with the transmitting antenna remaining free outside of the animal. Tags were deployed by experienced researchers from a Zodiac Mark V or a Solas ridged-hulled inflatable boat with a 40hp 4-stroke engine using an ARTS Whale Tagging PLT compressed air system. Whales were approached at idle speed from oblique angles so as not to cross over the flukes. No dependent calves were tagged, all whales were presumed to be adults based on their size. Tags were deployed from a range of 3-10 meters and placed near the dorsal fin, which contains the thickest blubber layer, and also provides the greatest height to transmit positional information via the exposed antenna.

All tags transmitted positional information via the Argos satellite system and were activated via a salt-water switch. In 2012, tags transmitted only during the hours 00:00-04:00 and 12:00-16:00 GMT. All subsequent tags were set without duty cycling and attempted to transmit data on each surfacing. Our raw data included 46,421 observations for 42 humpback whales from 2012-2016. We filtered observations without location data (n=103), duplicate timestamps (n=186), locations on land (n=1800), and implausible speed between consecutive locations (n=2552) to create a filtered set of observations. The duty cycle from the 2012 tags made it difficult to use tracks with large gaps and they were removed (n=6). From this initial dataset, we kept tracks that had points within 6 hour intervals, and discarded tracks less than 24 hours (Fig. S1).

CCAMLR Units Krill Fishery Data

Commercial krill catch data were provided with permission by CCAMLR for Subareas 48.1 - 48.4, aggregated into 0.5° latitude and 1.0° longitude grid cells. CCAMLR's small-scale management units range in size from 16,000 km² to 440,000 km² covering the northern section of the Antarctic Peninsula to south of Anvers Island (Fig. 1). Krill catches in kg were used from 1980-2015 and summed across all CCAMLR member nations. Catch is reported on a haul-by-haul basis, on either monthly or 5-day intervals, depending on the amount reported. The haul-by-haul data requires locations for start and end fishing positions. For analysis, the total amount of krill collected in each georeferenced grid cell was aggregated across seasons to represent the catch intensity in the waters surrounding the western side of the Antarctic Peninsula.

Hierarchical State-Space Models

To associate spatial patterns of animal movement with predicted behavior states, we used a hierarchical state-space model. State-space movement models combine a process model that estimates movement parameters and an observation model that accounts for the spatial uncertainty using Markov Chain

Monte Carlo (MCMC). We used a dynamic correlated random walk model following Jonsen *et al.* (2005) and Bestley *et al.* (2013), in which each movement stems from either a ‘traveling’ or ‘area-restricted search’ state (Bestley *et al.*, 2013; Beyer *et al.*, 2013; Jonsen *et al.*, 2005). The traveling state is defined as long straight tracks with small turning angles. Area-restricted search is defined as short step lengths with large turning angles. We prefer area-restricted search over ‘foraging’, since we recognize that whales will not be exclusively feeding during the defined state. Humpbacks display a threshold response to prey availability, likely due to the physiological effort required for foraging (Goldbogen *et al.*, 2008). This creates a characteristic spatial signature of whale foraging movements that may last up to several days (Dalla Rosa *et al.*, 2008).

Our discrete time observation model assigns each Argos location to a behavioral state based on the movement characteristics among consecutive Argos locations. We chose to model movement on a six hour time step, with individuals assumed to move in a straight line within in each time step. This assumption allows us to balance between the tractability of model-fitting and the temporal autocorrelation of whale behavior. The six hour time step is a short enough that we believe it is unlikely that we would miss transitions between foraging and traveling state, but coarse enough that we can fit the model using the data available. Given the strongly stereotyped nature of whale movements and the coarse categorization of the CCAMLR management units, we do not believe addition fine-scale variation in Argos error will influence our analysis. Our movement process model states that the geographic location of individual (*i*) at time (*t*) along a track (*g*) is multivariate normally distributed with a mean (*d*) and a variance (*σ*). The mean position (*d*) is a first order markov process, meaning that it depends on the previous step ($Y_{i,g,t} - Y_{i,g,t-1}$), plus the movement step at time *t*. This movement is a function of the degree of autocorrelation in step length (*γ*) and the turning angle (*T*). Step lengths and turning angles are considered to come from two behavioral states, ‘traveling’ and ‘area-restricted search’. The predicted behavioral state (*S*) is a multinomial draw with a probability of being in the traveling state

140 $(\phi_{i,g,t})$ or in the feeding state $(1 - \phi_{i,g,t})$. The probability for each behavioral state at time t is a
 141 function of the parameter $\alpha_{s_{i,g,t-1}}$, which is the estimated temporal autocorrelation in behavioral states.
 142 The variance in the argos observation estimate (τ_{argos}) was fixed for Argos error following the results
 143 reported in Jonsen et al. (2005), Bestley et al. (2013), (2015) and Jonsen (2016).

144 Process Model

$$Y_{i,g,t+1} \sim \text{Multivariate Normal}(d_{i,g,t}, \sigma)$$

$$d_{i,g,t} = Y_{i,g,t} + \gamma_{s_{i,g,t}} * T_{i,g,t} * (Y_{i,g,t} - Y_{i,g,t-1})$$

$$T_{i,g,t} = \begin{pmatrix} \cos(\theta_{s_{i,g,t}}) & -\sin(\theta_{s_{i,g,t}}) \\ \sin(\theta_{s_{i,g,t}}) & \cos(\theta_{s_{i,g,t}}) \end{pmatrix}$$

$$S_{i,g,t} \sim \text{Multinomial}(\phi_{i,g,t} \quad 1 - \phi_{i,g,t})$$

$$\text{logit}(\phi_{i,g,t}) = \alpha_{s_{i,g,t-1}}$$

145 Observation Model

$$w_{i,g,t,u} \sim \text{Multivariate Normal}(\hat{z}_{i,g,t,u}, \tau_{argos})$$

$$\hat{z}_{i,g,t,u} = (1 - j_{i,g,t,u}) * Y_{i,g,t-1} + j_{i,g,t,u} * Y_{i,g,t}$$

146 We ran our Bayesian model in Jags (Plummer, 2003) for 2 chains, each with 40,000 runs, with a burn-in
 147 of 38,000 draws. To reduce correlation among chains, we thinned the posterior by saving every 4th draw,
 148 yielding 500 values for the posterior distribution of each parameter.

149 To associate specific geographic areas with whale activity, we overlaid our study area with a 100
 150 by 100 grid of equal area cell ($\sim 23 \text{ km}^2$ in polar stereographic projection). We then calculated the total
 151 time each animal spent in each cell by taking the difference between the first and last timestamp of
 152 consecutive observations. The cell size was intentionally coarse, such that whales could only rarely pass

through a cell in a single observation point. These single observations were ignored, since they did not represent stationery occupancy of a cell. By adding the individual time rasters together, we created a map of the total time in each cell based on traveling behavior, area-restricted search, and both behaviors combined. To assess whether our results had been influenced by the location of tag placement, we performed the same analysis after trimming the first 7 days of data off each tag deployment. Whales can travel >100km in this time frame, and we use these data to compare to the map of whale behavior from the full dataset.

Do whales occupy SSMUs non-randomly?

Given that the SSMUs cover almost the entire study area, we must determine whether the pattern of spatial use by whales differs from random. Given the number of observation hours, and the differences in size of each of the SSMU, which units are occupied more or less than expected? We used parametric randomizations to create a null distribution of whale occupancy days based on an equal probability of presence in any cell. Using a 100 X 100 grid to cover the study area, for each of the 4818 cells ($cell_{x,y}$) within the SSMUs, the null expectation for the number of whale days is:

$$Whale\ days_{x,y} \sim Multinomial(k, \rho)$$

where k is the total number of whale days (3129 days) and $\rho = (1/4818)$ (i.e equal likelihood among all cells). We then summed the total number of whale days for each SSMU. We repeated this randomization 5000 times to generate a null distribution. We then compared our true number of whale days per SSMU to the null distribution of whale days per SSMU to determine if it was more, less or not-differing form random space use. If the observed value was outside the lower 5th or upper 95th of the null distribution, we considered it different from random.

Do whales behave non-randomly inside SSMUs?

To determine whether whale behavior differed from random within each SSMU, we used a null model to generate the expected proportion of area-restricted search given the total number of observations. Given that the model has to be in either traveling in area-restricted search state, the null expectation is a 50/50 split among behaviors. As the observation time decreases, there may be insufficient power to differentiate our results from random behavior. We evaluated this using the following null model:

$$\text{Proportion Area restricted search}_{x,y} \sim \text{Binomial}(n, k)$$

Where n = the number of whale observation days within each SSMU and k is 0.5 (i.e. equal probability of foraging or traveling). We drew 1000 draws from this null model to determine the expected deviation equal split given our sampling effort. If the observed value was outside the lower 5th or upper 95th of the null distribution, we considered our result different from random.

RESULTS

Our analyzed data contained 29,346 observations from 33 individuals from 2013-2016. On average individuals had 11.95 tracks, with an average track duration of 80.9 hours. The average number of observations per individual within our study area was 1187 (min = 31, max=2731). Average deployment for each individual within our study area was 46.06 days (min = 1.75, max = 145.5). Average time between observations was one hour and five minutes. Average distance between points was 824 meters, although this does not include estimate Argos error ellipses, and should be seen as approximate measure. We witnessed 11 whales begin their migration northward from Antarctic waters (defined as north of -60 latitude), ranging from March 22nd to July 17th (mean = May 25th). These migration events were assumed to relate to other life history functions (e.g. mating, calving) and were not included within this analysis.

Our behavioral model partitioned animal movement resulting from traveling and area-restricted search states. The traveling state was associated with high movement autocorrelation (mean $\gamma_1 = 0.91$ (0.88, 0.93)), and the area-restricted state with low movement autocorrelation (mean $\gamma_1 = 0.05$ (0.03, 0.06)). The mean probability of switching from traveling to area-restricted search was 9.9% (7.2%, 12.3%). The mean probability of switching from area-restricted search to traveling was 4.7% (3.4%, 6.2%). Transition probabilities must add to one, therefore the mean probability of remaining in the traveling state was 90.1% (87.7%, 92.8%), and the probability of remaining in the area-restricted search state was 95.3% (93.8%, 96.6%).

Overlaying the predicted behavioral state on our observed tracks showed strong concentrations of activity on the south side of Anvers Island, in the Gerlache Strait and along nearshore waters in Marguerite Bay (Fig. 3). Area-restricted search in the Bransfield Strait was restricted to the continental coastline, and away from the deep canyon south of Livingston Island. With the exception of one major feeding event on North side of Livingston Island in the late austral fall, no bouts of area-restricted search were seen north of the South Shetland Islands. We found no evidence of area-restricted search as whales crossed the Drake Passage and continued north towards tropical breeding grounds.

Partitioning our results by the CCAMLR small-scale management units (SSMUs), the majority of whale activity occurred in the Antarctic Peninsula West and Bransfield Strait West SSMUs (Table 1). This activity was disproportionately in the area-restricted search state. Only in the Bransfield Strait East and Drake Passage East units was there greater time spent in the traveling state. We recorded a total of 47 whale days in the area-restricted search state outside of the current SSMUs for the krill fishery, with the majority of this activity occurring southwest of Antarctic Pelagic Area SSMU boundary towards Marguerite Bay.

Aggregating the krill catch effort by SSMU showed uneven spatial catch by the krill fishery (Table 1). The Bransfield Strait West and the Drake Passage West units had the most catch between 1980 and 2009. Since 2009, the Drake Passage West unit has been used less frequently, but remains the 2nd most used unit by area. In contrast, the catch in the Bransfield Strait West unit increased dramatically during this period. Smaller increases were seen in Antarctic Peninsula West and Bransfield Strait East units. The temporal trend in krill fishery distribution is moving towards nearshore areas, with a decrease in the reported catch from the Elephant Island unit, and the lack of reported catch in the Antarctic East unit since 2009.

We asked whether whales used SSMUs non-randomly, given the differences in size of the management units. Compared to a random model of distribution, whales used the Bransfield Strait West, Drake Passage West, and Antarctic Peninsula West units more frequently than expected (Fig. 5a). The greatest number of observation days was within the Antarctic Peninsula West management unit. While several of the management units had less whale days than expected by random chance, we refrained from overly interpreting this finding (see discussion).

Our second randomization asked whether whales behaved non-randomly, given their presence in each of the SSMU (Fig. 5b). We found we had sufficient sampling to differentiate the observed behavior from random in the Drake Passage West, Antarctic Peninsula West, and Bransfield Strait West units. In all three units, we found a greater proportion of the area-restricted search state than expected. For the remaining units where we had whale observations, there was insufficient sampling to differentiate the observed allocation of behaviors from random.

DISCUSSION

Understanding the spatial distribution of krill predator foraging is imperative to manage the Antarctic Krill fishery. Humpback whales are the largest krill predator commonly found in the waters near the

West Antarctic Peninsula. The unknown effects of a warming climate on krill stock (Atkinson et al., 2004), combined with the slowly recovering cetacean community, suggest that precautionary management of krill fishery is required (Constable et al., 2000). We found that while humpbacks traverse the length of the entire western side of the Antarctic Peninsula, foraging areas are highly spatially clustered with respect to the CCAMLR small-scale management units (SSMUs). Whales used the Antarctic Peninsula West, Bransfield Strait West, and Drake Passage West SSMUs more often than expected. Within these units, whales spent more time in the area-restricted search state. Similarly, the krill fishery catch effort was spatially uneven, and has increasingly moved south and toward nearshore areas (Nicol and Foster, 2016). This finer scale information on humpback presence suggests that the uniform krill catch limit for all SSMUs in the Antarctic Peninsula Subarea 48.1 is overly broad and ignores important variation in humpback distribution and critical foraging areas. Resolving the disparity between the preference for specific areas by humpback whales and the uniform nature of the krill catch limits is critical in minimizing the potential conflict between whales and the fishery.

Our analysis identified key areas for humpback whale foraging, as represented by the area-restricted search state. Foremost, the Gerlache strait, in the Antarctic Peninsula West SSMU, had the greatest number of whale days in the area-restricted search state. While fishing effort is comparatively low in this SSMU, it is critical that CCAMLR proactively minimize conflict with foraging whales as the fishery shifts farther south, especially given the potential seasonal implications for whale life history. During the austral fall, krill that are distributed along the continental shelf move inshore towards areas that will be soon covered by sea-ice (Brierley, 2002). As humpbacks follow krill inshore, they congregate in greater densities and have reduced foraging space (Curtice et al., 2015; Nowacek et al., 2011). The shift towards krill fishing later in the austral fall (CCAMLR, 2015) coincides with a lack of ice-free areas outside the Gerlache and Bransfield straits during this period (Friedlaender et al., 2011), highlighting the potential for negative impacts from the fishery on concentrated whale aggregations. Given the relatively

low use of the Antarctic Peninsula West SSMU by the krill fishery, we advocate for a closure of this area, especially during the austral fall, as a preventative measure to preempt conflict between the fishery and marine predators.

In contrast to the Antarctic Peninsula West Unit, the adjoining Bransfield Strait West unit is the most used SSMU by the krill fishery. We found significant presence of humpbacks in this unit, especially in the nearshore areas south of Trinity Island. The spatial overlap between whale presence and concentrated fishing activities has negative implications for whale mortality due to the potential for ship-strikes and long-term effects on resource competition between the fishery and prey base. Given the dominance of this unit in the overall fisheries catch, it is unlikely that member nations will agree to large-scale catch limits. Nearby anecdotal observations suggest that whales move closer to shore in the austral fall (Nowacek et al., 2011), and further work must focus on the seasonal patterns of whale movement to determine whether distance to shore restrictions would reduce the potential overlap between fishing vessels and whales.

To differentiate observed patterns of whale distribution and behavior from random expectation, we used randomization tests to account for the different sizes of the SSMUs and our overall sampling effort. One remaining question is whether tagging location could have influenced our results. We believe that this effect is likely to be small. Rather than stay near tagging locations, whales traveled widely, covering the full breadth of the Antarctic Peninsula throughout the summer. This means that the importance of the initial location of tagging quickly diminished over time. Re-analyzing our map of area-restricted search days after removing the first week of observations showed the same spatial pattern as our full dataset (Fig. S2). Ship-board surveys along the north end of the Antarctic Peninsula have previously highlighted deep water areas in the Bransfield Strait (Santora et al., 2010; Santora and Veit, 2013), and Dalla Rosa (2008) documented two cases of resightings of tagged humpbacks returning to the Gerlache Strait, underlining the importance of this area. While our dataset is larger than most

287 cetacean movement studies, it would be inappropriate to conflate the lack of presence with inferred
288 absence. We focus on where whales were observed, and our results should not be interpreted as
289 dismissing the conservation potential of other areas within the study region.

290 The distribution of humpback whales may provide insights into the recovery of other krill
291 foraging species. Humpbacks travel widely throughout the peninsula, and are not tied to breeding
292 colonies or haul-out locations that limit the mobility of other krill predators, such as penguins and seals.
293 The presence of humpback whales may therefore be a strong indicator of krill abundance, and our
294 distribution patterns may represent important foraging areas for other krill predators along the West
295 Antarctic Peninsula. For cetaceans, the SSMUs with the highest krill catch rates also overlap with areas
296 that historically had high numbers of fin and blue whales (Kemp and Bennett 1932). These species yet to
297 recover from historic whaling, when at least 11,927 fin and 6,628 blue whales were killed in this region
298 during the 1920's (Kemp and Bennett 1932). Current distribution information is sparse, but observations
299 of fin whales, which are few, are largely restricted to the Bransfield Strait East unit (e.g. Herr *et al.* 2016).
300 Observations of blue whales are limited in the continental shelf waters of the West Antarctic Peninsula
301 (Friedlaender unpublished data). As these animals repatriate the region, precaution must be taken to
302 ensure that populations recover unhindered from diminished prey availability due to commercial
303 harvest. We still lack information on the consumption rates and energetic demands of baleen whales
304 (Goldbogen et al., 2015; Hazen et al., 2009), and a loss of food base has been shown to have negative
305 demographic effects in cetaceans (Leaper et al., 2006), as well as other Antarctic marine wildlife (Croxall
306 et al., 1999).

307 We highlight the concentrated space use of humpback whales in the Antarctic West, and
308 Bransfield Strait West CCAMLR small-scale management units. To avoid conflict with the krill fishery,
309 additional management action is needed to refine the spatial scale of catch limits, and provide
310 ecologically informed guidelines for near-shore restrictions. Given the potential influence of global

climate change to allow both whales and ships to stay longer in the Antarctic, regulations must also consider the temporal timing of krill movement and habitat availability in reducing competition for ice-free areas during the austral fall.

ACKNOWLEDGEMENTS

This research was supported by research grants from the Antarctic Wildlife Research Fund, International Whaling Commission, Southern Ocean Research Partnership and National Science Foundation ANT-0823101, 1250208, and 1440435. Research was conducted under NMFS Permits 14907, 14809, and 14856, ACA Permits 2009-013 and 2015-011, Duke University IACUC A049-122-02 and Oregon State University ACUP 4513. We would like to thank the crews of the *R/V Point Sur* and *ARSV Laurence M Gould* for their support, and numerous colleagues for their constructive comments and feedback during the development of this project including Rodolfo Werner, Doug Nowacek, Andy Read, and Caroline Casey. We are grateful for the field contributions of John Durban, Bob Pitman, Reny Tyson, Matthew Bowers, Ladd Irvine, Heather Foley, Erin Pickett, and numerous marine technicians. We would also like to thank members of CCAMLR for providing data from the krill fishery.

LITERATURE CITED

- Atkinson, A., Siegel, V., Pakhomov, E.A., Rothery, P., 2004. Long-term decline in krill stock and increase in salps within the Southern Ocean. *Nature* 432, 100–103. doi:10.1038/nature02950.1.
- Bestley, S., Jonsen, I.D., Hindell, M.A., Harcourt, R.G., Gales, N.J., 2015. Taking animal tracking to new depths: Synthesizing horizontal - vertical movement relationships for four marine predators. *Ecology* 96, 417–427. doi:10.1890/14-0469.1
- Bestley, S., Jonsen, I.D., Hindell, M. a, Guinet, C., Charrassin, J.-B., 2013. Integrative modelling of animal movement: incorporating in situ habitat and behavioural information for a migratory marine predator. *Proc. R. Soc. B Biol. Sci.* 280, 2012–2262. doi:10.1098/rspb.2012.2262

334 Beyer, H.L., Morales, J.M., Murray, D., Fortin, M.J., 2013. The effectiveness of Bayesian state-space
 335 models for estimating behavioural states from movement paths. *Methods Ecol. Evol.* 4, 433–441.
 336 doi:10.1111/2041-210X.12026

337 Brierley, A.S., 2002. Antarctic krill under sea ice: elevated abundance in a narrow band just south of ice
 338 edge. *Science* (80-.). 295, 1890–1892. doi:10.1126/science.1068574

339 CCAMLR, 2015. Krill Fishery Report. *Comm. Conserv. Antarct. Mar. Living Resour.* 1–35.
 340 doi:https://www.ccamlr.org/en/system/files/00%20KRI48%202015_1.pdf

341 Clapham, P.J., Young, S.B., Brownell Jr., R.L., 1999. Baleen whales: conservation issues and the status of
 342 the most endangered populations. *Mamm. Rev.* 29, 37–60. doi:10.1046/j.1365-2907.1999.00035.x

343 Constable, A.J., de la Mare, W.K., Agnew, D.J., Everson, I., Miller, D., 2000. Managing fisheries to
 344 conserve the Antarctic marine ecosystem: practical implementation of the Convention on the
 345 Conservation of Antarctic Marine Living Resources (CCAMLR). *ICES J. Mar. Sci.* 57, 778–791.
 346 doi:10.1006/jmsc.2000.0725

347 Croxall, J.P., Reid, K., Prince, P.A., 1999. Diet, provisioning and productivity responses of marine
 348 predators to differences in availability of Antarctic krill. *Mar. Ecol. Prog. Ser.* 177, 115–131.
 349 doi:10.3354/meps177115

350 Curtice, C., Johnston, D.W., Ducklow, H.W., Gales, N.J., Halpin, P.N., Friedlaender, A.S., 2015. Modeling
 351 the spatial and temporal dynamics of foraging movements of humpback whales (*Megaptera*
 352 *novaeangliae*) in the Western Antarctic Peninsula. *Mov. Ecol.* 3, 13. doi:10.1186/s40462-015-0041-
 353 x

354 Dalla Rosa, L., Secchi, E.R., Maia, Y.G., Zerbini, A.N., Heide-Jørgensen, M.P., 2008. Movements of
 355 satellite-monitored humpback whales on their feeding ground along the Antarctic Peninsula. *Polar*

356 Biol. 31, 771–781. doi:10.1007/s00300-008-0415-2

357 Ellison, W.T., Southall, B.L., Clark, C.W., Frankel, A.S., 2012. A New Context-Based Approach to Assess
358 Marine Mammal Behavioral Responses to Anthropogenic Sounds. *Conserv. Biol.* 26, 21–28.
359 doi:10.1111/j.1523-1739.2011.01803.x

360 Friedlaender, A.S., Halpin, P.N., Qian, S.S., Lawson, G.L., Wiebe, P.H., Thiele, D., Read, A.J., 2006. Whale
361 distribution in relation to prey abundance and oceanographic processes in shelf waters of the
362 Western Antarctic Peninsula. *Mar. Ecol. Prog. Ser.* 317, 297–310. doi:10.3354/meps317297

363 Friedlaender, A.S., Johnston, D.W., Fraser, W.R., Burns, J., Patrick N., H., Costa, D.P., 2011. Ecological
364 niche modeling of sympatric krill predators around Marguerite Bay, Western Antarctic Peninsula.
365 *Deep. Res. Part II Top. Stud. Oceanogr.* 58, 1729–1740. doi:10.1016/j.dsr2.2010.11.018

366 Friedlaender, A.S., Tyson, R.B., Stimpert, A.K., Read, A.J., Nowacek, D.P., 2013. Extreme diel variation in
367 the feeding behavior of humpback whales along the western Antarctic Peninsula during autumn.
368 *Mar. Ecol. Prog. Ser.* 494, 281–289. doi:10.3354/meps10541

369 Goldbogen, J.A., Calambokidis, J., Croll, D.A., Harvey, J.T., Newton, K.M., Oleson, E.M., Schorr, G.,
370 Shadwick, R.E., 2008. Foraging behavior of humpback whales: kinematic and respiratory patterns
371 suggest a high cost for a lunge. *J. Exp. Biol.* 211, 3712–3719. doi:10.1242/jeb.023366

372 Goldbogen, J.A., Hazen, E.L., Friedlaender, A.S., Calambokidis, J., Deruiter, S.L., Stimpert, A.K., Southall,
373 B.L., 2015. Prey density and distribution drive the three-dimensional foraging strategies of the
374 largest filter feeder. *Funct. Ecol.* 1–11. doi:10.1111/1365-2435.12395

375 Hays, G.C., Ferreira, L.C., Sequeira, A.M.M., Meekan, M.G., Duarte, C.M., Bailey, H., Bailleul, F., Bowen,
376 W.D., Caley, M.J., Costa, D.P., Eguíluz, V.M., Fossette, S., Friedlaender, A.S., Gales, N., Gleiss, A.C.,
377 Gunn, J., Harcourt, R., Hazen, E.L., Heithaus, M.R., Heupel, M., Holland, K., Horning, M., Jonsen, I.,

378 Kooyman, G.L., Lowe, C.G., Madsen, P.T., Marsh, H., Phillips, R.A., Righton, D., Ropert-Coudert, Y.,
 379 Sato, K., Shaffer, S.A., Simpfendorfer, C.A., Sims, D.W., Skomal, G., Takahashi, A., Trathan, P.N.,
 380 Wikelski, M., Womble, J.N., Thums, M., 2016. Key Questions in Marine Megafauna Movement
 381 Ecology. *Trends Ecol. Evol.* 31, 463–475. doi:10.1016/j.tree.2016.02.015

382 Hazen, E.L., Friedlaender, A.S., Thompson, M.A., Ware, C.R., Weinrich, M.T., Halpin, P.N., Wiley, D.N.,
 383 2009. Fine-scale prey aggregations and foraging ecology of humpback whales *Megaptera*
 384 *novaeangliae*. *Mar. Ecol. Prog. Ser.* 395, 75–89. doi:10.3354/meps08108

385 Hedley, S., Reilly, S., Borberg, J., Holland, R., Hewitt, R., Watkins, J., Naganobu, M., Sushin, V., 2001.
 386 Modelling whale distribution: a preliminary analysis of data collected on the CCAMLR-IWC Krill
 387 Synoptic Survey, 2000. *Rep. Int. Whal. Comm.* 1–38.

388 Herr, H., Viquerat, S., Siegel, V., Kock, K.H., Dorschel, B., Huneke, W.G.C., Bracher, A., Schröder, M., Gutt,
 389 J., 2016. Horizontal niche partitioning of humpback and fin whales around the West Antarctic
 390 Peninsula: evidence from a concurrent whale and krill survey. *Polar Biol.* 39, 799–818.
 391 doi:10.1007/s00300-016-1927-9

392 Hewitt, R.P., Watters, G., Trathan, P.N., Croxall, J.P., Goebel, M.E., Ramm, D., Reid, K., Trivelpiece, W.Z.,
 393 Watkins, J.L., 2004. Options for allocating the precautionary catch limit of krill among small-scale
 394 management units in the Scotia Sea. *CCAMLR Sci.* 11, 81–97.

395 Hill, S.L., Murphy, E.J., Reid, K., Trathan, P.N., Constable, A.J., 2006. Modelling Southern Ocean
 396 ecosystems: krill, the food-web, and the impacts of harvesting. *Biol. Rev.* 81, 581–608.
 397 doi:10.1017/S1464793106007123

398 Johnston, D.W., Friedlaender, A.S., Read, A.J., Nowacek, D.P., 2012. Initial density estimates of
 399 humpback whales *Megaptera novaeangliae* in the inshore waters of the western Antarctic

400 Peninsula during the late autumn. *Endanger. Species Res.* 18, 63–71. doi:10.3354/esr00395

401 Jonsen, I., 2016. Joint estimation over multiple individuals improves behavioural state inference from
 402 animal movement data. *Sci. Rep.* 6, 20625. doi:10.1038/srep20625

403 Jonsen, I.D., Flemming, J., Myers, R., 2005. Robust State-Space Modeling of Animal Movement Data.
 404 *Ecology* 86, 2874–2880.

405 Kawaguchi, S., Nicol, S., Taki, K., Naganobu, M., 2006. Fishing ground selection in the Antarctic krill
 406 fishery: Trends in patterns across years, seasons and nations. *CCAMLR Sci.* 13, 117–141.

407 Kemp, S., and A.G. Bennett. 1932. On the distribution and movements of whales on the South Georgia
 408 and South Shetland whaling grounds. *Discovery Reports*, VI, 165-190.

409 Leaper, R., Cooke, J., Trathan, P., Reid, K., Rowntree, V., Payne, R., 2006. Global climate drives southern
 410 right whale (*Eubalaena australis*) population dynamics. *Biol. Lett.* 2, 289–292.
 411 doi:10.1098/rsbl.2005.0431

412 Matsuoka, K., Hakamada, T., Kiwada, H., Murase, H., Nishiwaki, S., 2006. Distributions and standardized
 413 abundance estimates for humpback , fin and blue whales in the Antarctic Areas IIIE, IV, V and VIW
 414 (35°E -145°W), south of 60°S. *Reports Int. Whal. Comm. SC/D06/J7*, 1–37.

415 Nicol, S., Foster, J., 2016. The Fishery for Antarctic Krill: Its Current Status and Management Regime, in:
 416 Siegel, V. (Ed.), *Biology and Ecology of Antarctic Krill, Advances in Polar Ecology*. Springer
 417 International Publishing, Cham, pp. 387–421. doi:10.1007/978-3-319-29279-3

418 Nicol, S., Foster, J., Kawaguchi, S., 2012. The fishery for Antarctic krill - recent developments. *Fish Fish.*
 419 13, 30–40. doi:10.1111/j.1467-2979.2011.00406.x

420 Nicol, S., Worby, A., Leaper, R., 2008. Changes in the Antarctic sea ice ecosystem: Potential effects on

421 krill and baleen whales. *Mar. Freshw. Res.* 59, 361–382. doi:10.1071/MF07161

422 Nowacek, D.P., Friedlaender, A.S., Halpin, P.N., Hazen, E.L., Johnston, D.W., Read, A.J., Espinasse, B.,
423 Zhou, M., Zhu, Y., 2011. Super-aggregations of krill and humpback whales in Wilhelmina bay,
424 Antarctic Peninsula. *PLoS One* 6, 2–6. doi:10.1371/journal.pone.0019173

425 Plummer, M., 2003. JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling.
426 *Proc. 3rd Int. Work. Distrib. Stat. Comput.* 1–10. doi:ISSN 1609-395X

427 Santora, J.A., Reiss, C.S., Loeb, V.J., Veit, R.R., 2010. Spatial association between hotspots of baleen
428 whales and demographic patterns of Antarctic krill *Euphausia superba* suggests size-dependent
429 predation. *Mar. Ecol. Prog. Ser.* 405, 255–269. doi:10.3354/meps08513

430 Santora, J.A., Veit, R.R., 2013. Spatio-temporal persistence of top predator hotspots near the Antarctic
431 Peninsula. *Mar. Ecol. Prog. Ser.* 487, 287–304. doi:10.3354/meps10350

432 Tarling, G.A., Klevjer, T., Fielding, S., Watkins, J., Atkinson, A., Murphy, E., Korb, R., Whitehouse, M.,
433 Leaper, R., 2009. Variability and predictability of Antarctic krill swarm structure. *Deep. Res. Part I*
434 *Oceanogr. Res. Pap.* 56, 1994–2012. doi:10.1016/j.dsr.2009.07.004

435 Trathan, P.N., García-Borboroglu, P., Boersma, D., Bost, C.-A., Crawford, R.J.M., Crossin, G.T., Cuthbert,
436 R.J., Dann, P., Davis, L.S., De La Puente, S., Ellenberg, U., Lynch, H.J., Mattern, T., Pütz, K., Seddon,
437 P.J., Trivelpiece, W., Wienecke, B., 2015. Pollution, habitat loss, fishing, and climate change as
438 critical threats to penguins. *Conserv. Biol.* 29, 31–41. doi:10.1111/cobi.12349

439 Tyson, R.B., Friedlaender, A.S., Nowacek, D.P., 2016. Does optimal foraging theory predict the foraging
440 performance of a large air-breathing marine predator? *Anim. Behav.* 116, 223–235.
441 doi:10.1016/j.anbehav.2016.03.034

443 TABLES

444 Table 1. Number of whale hours in each the CCAMLR small-scale management units (SSMUs) within
 445 Subarea 48.1. The duration of time was calculated by the time difference between the last observation
 446 and the first observation for each movement track for each animal. The sum value across animals is
 447 shown. We also included ‘outside proposed units’, which largely consists of observations west of
 448 Marguerite Bay (see Fig. 3). Ordered by total time in area-restricted search. Total krill catch is shown
 449 between 1980-2009 and 2009-2015, as well as the catch density per year, to account for difference in

Unit	Whale Traveling (Days)	Whale Area-restricted Search (Days)	Krill Catch (Metric Tons/Year)		Catch Density (Metric Tons/ km ² /Year)	
			1980-2009	2009-2015	1980-2009	2009-2015
Antarctic Peninsula West	222.17	430.78	332	9335	0.01	0.25
Bransfield Strait West	61.33	193.68	13,486	55,688	0.06	2.47
Antarctic Peninsula Pelagic Area	122.82	134.54	2,082	141	>0.01	>0.01
Drake Passage West	41.42	63.64	17,774	10,132	1.11	0.63
Outside Proposed Units	42.15	47.78	-	-	-	-
Bransfield Strait East	5.19	4.29	1,357	11,044	0.05	0.40
Drake Passage East	6.53	5.32	6,923	4,882	0.42	0.30
Elephant Island	0.00	0.00	10,917	620	0.30	0.02
Antarctic Peninsula East	0.00	0.00	156	-	0.08	-

450 SSMU area.

451 FIGURES

452 Figure 1. CCAMLR small-scale management units for the West Antarctic Peninsula krill fishery (CCAMLR

453 Sub-unit 48.1). For brevity “Antarctic Peninsula” is abbreviated AP.

454

455 Figure 2. Total krill catch between 1980-2009 and 2009-2015 in the West Antarctic Peninsula. The krill
456 fishery has moved inshore during recent years. The eight CCAMLR small-scale management units for
457 region 48.1 are shown in black lines.

458

459 Figure 3. Predicted behaviors for each filtered whale track based on the state-space model. If the mean
460 probability of switching to the area-restricted state (ϕ_{ARS}) was greater than 0.5, the observation was
461 labeled as area-restricted search. The eight CCAMLR small-scale management units for region 48.1 are
462 shown in black lines.

463 Figure 4. Total whale days for each behavior state. Using our state-space model to estimate the
464 movement process we divided locations as belonging to traveling or area restricted search states. We
465 then calculated the total time each whale spent in each state at each grid cell. Combining the results for
466 all individuals we mapped the distribution of behavioral states across the waters surrounding the West
467 Antarctic Peninsula.

Figure 5. Randomization tests for the occurrence and behavior within CCAMLR small-scale management units (SSMUs): **A)** Do whales use SSMUs non-randomly? We compared the observed duration of whale presence (in days) for each SSMU with a null expectation based on equal probability of occurrence in all cells. Black dots are the observed total days in each of the SSMUs. Grey bars are the lower 5th and upper 95th of the null distribution. Points to the right of the grey bars indicate that whales spent more time in a SSMU than expected. Points to the left of the grey bars indicate whales spent less time in a SSMU than expected at random. **B)**: Do whales behave non-randomly within SSMUs? Using our movement model, we calculated the percentage of time in area-restricted search in each the SSMUs and compared it to a null distribution of foraging given the total number of observations. Black points are the observed proportion, grey bars are the lower 5th and upper 95th of the null distribution. Black points to the right of the grey bar indicate a greater proportion of area-restricted search than expected given the level of sampling within a SSMU.