1
2
3
4
5
6
7
8
9
10 **Applications and limitations of U-Th disequilibria systematics for determining ages of**
11 **carbonate alteration minerals in peridotite**

12 Evelyn M. Mervine^{a,*}, Kenneth W. W. Sims^b, Susan E. Humphris^c, and Peter B. Kelemen^d

13 Prepared for submission to *Chemical Geology*
14

15 ^aMIT/WHOI Joint Program, Geology and Geophysics Department, Woods Hole Oceanographic
16 Institution, 266 Woods Hole Road, Woods Hole, MA 02543, USA, emervine@whoi.edu

17
18 ^bGeology and Geophysics Department, University of Wyoming, 1000 East University Avenue,
19 Laramie, Wyoming 82071, USA, ksims7@uwyo.edu

20
21 ^cGeology and Geophysics Department, Woods Hole Oceanographic Institution, 266 Woods Hole
22 Road, Woods Hole, MA 02543, USA, shumphris@whoi.edu

23
24 ^dLamont-Doherty Earth Observatory, 61 Route 9W, Palisades, NY 10964, USA,
25 peterk@ldeo.columbia.edu
26

27
28
29
30
31 *Corresponding author. Present address: De Beers Marine, DBM Gardens, Golf Park 2,
32 Raapenberg Road, Pinelands, 7405, Cape Town, South Africa, Tel: +27 (0)21 658 3224.
33 Evelyn.Mervine@debeersgroup.com.

Abstract

^{238}U – ^{234}U – ^{230}Th dating was conducted on carbonate alteration minerals in the peridotite layer of the Samail Ophiolite, Sultanate of Oman, in order to assess the applicability of U-series dating techniques to these types of Quaternary terrestrial carbonates and also to further constrain natural rates of carbonation of the peridotite. Due to their low U concentrations and relatively high Th/U ratios, Samail carbonates are challenging to date with the ^{230}Th technique because of the sensitivity of ages to corrections for initial ^{230}Th . Uncorrected ^{230}Th ages for Ca-rich travertines are consistently older than previously obtained ^{14}C ages. However, geologically reasonable initial ^{230}Th corrections bring the two sets of ages into concordance. This age concordance suggests that the travertines are generally closed systems, adding a level of credence to the reliability of previously obtained ^{14}C ages. In contrast, uncorrected ^{230}Th ages for Mg-rich carbonate veins are generally younger than previously obtained ^{14}C ages. These young ages are interpreted in terms of remobilization of hexavalent U, which is subsequently deposited as tetravalent U by reduced serpentinization fluids. Two Mg-rich carbonate veins sampled at a roadcut have near-equilibrium ($^{230}\text{Th}/^{238}\text{U}$) and ($^{234}\text{U}/^{238}\text{U}$) values, which indicate that these veins are >375,000 years in age, consistent with their “ ^{14}C dead” (>50,000 yr BP) ages. The variable young and old ages for these Mg-rich carbonate veins indicate that carbonation of the peridotite layer of the Samail Ophiolite is an ongoing process and that there have been multiple generations of subsurface carbonate vein formation. Overall, this study provides insights into some of the challenges associated with applying U-series dating methods to Quaternary terrestrial carbonates, in particular carbonate alteration minerals in peridotites, and highlights some areas where there is room for improvement, such as obtaining better constraints on the isotopic composition of admixed detritus, and also some advantages, such as the ability to identify open system behavior not apparent from ^{14}C dating and stable C and O isotopic analysis alone.

Keywords: carbonate; peridotite; uranium; thorium; U-series dating; ophiolite

1. Introduction

Determining timescales of the formation and preservation of carbonate alteration minerals in mantle peridotite is essential to better understand the role of this potentially important sink in the global carbon cycle and also to evaluate the feasibility of using artificially enhanced, *in situ* formation of carbonates in peridotite as a method for mitigating the buildup of anthropogenic CO₂ emissions in the atmosphere (e.g. Seifritz, 1990; Lackner et al., 1995; Lackner, 2002; Kelemen and Matter, 2008; Matter and Kelemen, 2009; Kelemen et al., 2011). While natural carbonation of peridotite is commonly observed subaerially and on the seafloor (e.g. Trommsdorff and Evans, 1977; Trommsdorff et al., 1980; Ferry, 1995; Surour and Arafa, 1997; Kelley et al., 2001; Fröh-Green et al., 2003; Ludwig et al., 2006, 2011; Kelemen and Matter, 2008; Matter and Kelemen, 2009; Power et al., 2009; Kelemen et al., 2011; Pronost et al., 2011; Beinlich and Austrheim, 2012; Harrison et al., 2013; Chavagnac et al., 2013a, b; Mervine et al., 2014), the natural rate of peridotite carbonation and therefore the rate of CO₂ uptake via this alteration mechanism is poorly known (e.g. Wilson et al., 2006, 2009a, 2009b; Kelemen and Matter, 2008; Kelemen et al., 2011; Mervine et al., 2014). In part, this is because carbonate alteration minerals in peridotite are challenging to date. Several studies (e.g. Kelemen and Matter, 2008; Wilson et al., 2009b; Kelemen et al., 2011; Mervine et al., 2014) have employed ¹⁴C dating, but this dating technique has a practical limit of only ~50,000 years. Use of the ²³⁸U-²³⁴U and ²³⁴U-²³⁰Th disequilibria dating techniques (which have practical dating limits of ~1 million years and ~375,000 years, respectively) and the U-Pb dating technique (which is suitable for dating on timescales of millions of years) to investigate longer timescales of carbonation is limited by the generally low (ppb level) U concentrations of peridotites and their associated carbonate alteration minerals (e.g. Hanghøj et al., 2010; Bodinier and Godard, 2014). Carbonates formed in peridotites at the Lost City Hydrothermal Field, which is located off-axis

of the mid-Atlantic Ridge, have been dated using U-Th disequilibria techniques (Ludwig et al., 2011). However, these carbonates have high U concentrations (ppm level) due to contribution of U from seawater (Ludwig et al., 2011). Historically, little work has been done to apply the U-Pb dating technique to carbonates although recently there has been some development in this field, with a focus on dating of speleothem samples with high U and low common Pb concentrations (e.g. Rasbury and Cole, 2009; Woodhead and Pickering, 2012; Hellstrom and Pickering, 2015).

Furthermore, while some carbonates, particularly corals and speleothems, are routinely dated with the U-Th disequilibria methods (e.g. Edwards et al., 2003), dating of terrestrial carbonates with these techniques can be challenging due to the significant presence of admixed detritus. Unlike most corals and speleothems, terrestrial carbonates are usually affected by the presence of initial ^{230}Th and, to a lesser extent, by ^{234}U and ^{238}U contributed by detrital material (e.g. Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991; Kaufman, 1993). This detritus is often fine-grained and intergrown with the carbonate, making physical separation very difficult (e.g. Ku and Liang, 1984; Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991). Various approaches have been attempted over the years to account for detrital ^{230}Th , ^{234}U , and ^{238}U in terrestrial carbonates, including light acid leaching, separating the carbonate material from the admixed detritus, and pseudoisochron methods for determining the composition of incorporated detritus (e.g. Osmond et al., 1970; Schwarcz and Latham, 1989; Przybylowicz et al., 1991; Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991). However, these techniques have had limited success. In most cases, employing total sample dissolutions with corrections for detrital inputs is the most robust way to obtain U-Th disequilibria ages for terrestrial carbonates (e.g. Kaufman, 1993; Ludwig and Titterton, 1994; Edwards et al., 2003). However, terrestrial carbonates with low U concentrations and high Th/U ratios, such as carbonate alteration products formed in

peridotites, can be challenging to date with U-Th disequilibria methods since the ages are highly sensitive to detrital corrections.

In this study we examine the applications and limitations of using U-Th disequilibria methods to date carbonate alteration minerals that formed in peridotites of the Samail Ophiolite, Sultanate of Oman (Figure 1). The Samail Ophiolite is one of the largest and best-exposed ophiolites in the world (Glennie et al., 1973, 1974; Coleman, 1977, 1981; Lippard et al., 1986; Nicolas et al., 2000) and is thus an excellent location for investigating natural rates of carbonate formation in peridotite. The goal of this study was to determine ^{238}U - ^{234}U and ^{234}U - ^{230}Th ages and age limits for Samail carbonates in order to further assess the range of ages of carbonate alteration minerals in the peridotite layer of the Samail Ophiolite. This study builds upon previous ^{14}C dating and stable C and O isotope analyses of Samail carbonates, which are presented in Mervine et al. (2014) and also in Clark and Fontes (1990), Clark et al. (1992), Kelemen and Matter (2008), and Kelemen et al. (2011). Because these Quaternary terrestrial carbonates contain significant admixed fine-grained detritus, we undertook methods to: (1.) attempt to separate the aluminosilicate detritus from the carbonate or (2.) accurately correct for the effect the detritus is having on the ^{238}U - ^{234}U - ^{230}Th isotope systematics of the carbonate minerals. Comparing the ^{238}U - ^{234}U and ^{234}U - ^{230}Th ages and age limits determined for the Samail carbonates with previously published ^{14}C ages on the same samples (Mervine et al., 2014) enables us to more carefully scrutinize the reliability of ages obtained with both dating systems and to further constrain timescales of natural carbonation of Samail Ophiolite peridotites.

2. Background

2.1 Terrestrial carbonates in the Samail Ophiolite

The Samail Ophiolite ranges from approximately 50 to 100 km in width and extends for greater than 500 km (e.g. Lippard et al., 1986; Nicolas et al., 2000). The ophiolite consists of several uplifted, thrust-bounded blocks (nappes) of oceanic crust and mantle that were obducted at ~80 to 95 Ma on top of autochthonous shelf carbonate rocks (the Hajar Supergroup) and parautochthonous continental slope carbonate rocks (the Sumeini Group), which rest on top of pre-Permian crystalline basement (Coleman, 1981; Lippard et al., 1986; Searle and Cox, 1999, 2002). These nappes consist of ~4 to 7 km of crustal rocks (layered gabbros, sheeted dikes, and volcanics, including pillow lavas) and ~8 to 12 km of upper mantle, primarily harzburgite (Glennie et al., 1973, 1974; Coleman, 1981; Lippard et al., 1986; Nicolas et al., 2000). Abundant carbonate veins as well as calcite-rich travertines are found throughout the peridotite layer of the ophiolite (e.g. Neal and Stanger, 1984, 1985; Clark and Fontes, 1990; Clark et al., 1992; Kelemen and Matter, 2008; Kelemen et al., 2011; Chavagnac et al., 2013 a, b; Mervine et al., 2014).

This study focuses on two types of carbonate alteration products that form in the peridotite layer of the Samail Ophiolite: (1.) travertine precipitated from high pH springs (Clark and Fontes, 1990; Clark et al., 1992; Kelemen and Matter, 2008; Kelemen et al., 2011; Paukert et al., 2012; Chavagnac et al., 2013a, 2013b; Mervine et al., 2014) and (2.) carbonate veins that form *in situ* in partially hydrated (serpentinized) peridotite (Kelemen and Matter, 2009; Kelemen et al., 2011; Streit et al., 2012; Mervine et al., 2014). These carbonates are thought to form as a result of low temperature alteration of peridotite through interaction with meteoric water (Figure 2) (e.g. Barnes et al., 1967, 1978; Barnes and O'Neil, 1969, 1971; Neal and Stanger, 1985; Bruni et al., 2002; Cipolli et al., 2004; Kelemen et al., 2011). When meteoric water weathers partially

serpentinized peridotite, $\text{Mg}^{2+}\text{-HCO}_3^-$ rich waters (known as “Type I” waters) are formed. As these waters percolate deeper into peridotite bedrock where they are no longer in equilibrium with the atmosphere, they precipitate Mg-rich carbonates, serpentines, and clays. As a result of this precipitation and other reactions in the subsurface, the waters transform into $\text{Ca}^{2+}\text{-OH}^-$ waters (known as “Type II” waters) that have very high pH (10 to 12) and low Mg, C, and oxygen fugacity (Eh approximately -200 mV) (Neal and Stanger, 1983, 1984, 1985; Clark and Fontes, 1990; Kelemen and Matter, 2008; Kelemen et al., 2011; Paukert et al., 2012; Chavagnac et al., 2013a, b). When Type II waters return to the surface as hyperalkaline springs, they rapidly react with atmospheric CO_2 to precipitate calcite-rich travertines (e.g. Clark and Fontes, 1990; Clark et al., 1992; Kelemen and Matter, 2008; Paukert et al., 2012; Chavagnac et al., 2013a, b; Mervine et al., 2014).

The calcite-rich travertines (Figure 3, Panels A to C) are precipitated from the hyperalkaline springs found throughout the Samail Ophiolite, predominantly in the peridotite layer and concentrated along the basal thrust of the ophiolite as well as along the crust-mantle boundary (Neal and Stanger, 1984, 1985; Clark and Fontes, 1990; Clark et al., 1992; Dewandel et al., 2003, 2004, 2005; Kelemen and Matter, 2008; Kelemen et al., 2011; Paukert et al., 2012; Chavagnac et al., 2013a, b). At these springs, thick travertine deposits (up to several meters, on average ~1 to 2 m) form on top of peridotite, which is highly fractured with abundant carbonate-serpentine veins. Turquoise-blue alkaline pools lined with white carbonate precipitate are a feature of many hyperalkaline springs. Browner, more weathered (and generally older) travertine deposits are often found upslope and to the side of the actively precipitating travertine deposits (Kelemen and Matter, 2008; Mervine et al., 2014).

The carbonate veins (Figure 3, Panels D to H) form *in situ* in the peridotite and, to a lesser extent, in the gabbro and basalt layers of the ophiolite (Kelemen and Matter, 2008; Kelemen et al., 2011; Mervine et al., 2014). Carbonate veins adjacent to surface travertine deposits are Ca-rich and similar in composition to the travertines. Carbonate veins sampled far from travertines are generally Mg-rich and presumably formed in the deeper subsurface during the transformation of Type I waters to Type II waters. The Mg-rich carbonate veins are predominately composed of magnesite and dolomite, range in size from a few millimeters to several meters thick, and can extend for hundreds of meters (Kelemen and Matter, 2008; Kelemen et al., 2011; Mervine et al., 2014). The Mg-rich carbonate veins are often intergrown with serpentine veins.

2.2 Previous dating of carbonate alteration minerals in peridotite in the Samail Ophiolite

Carbonate alteration minerals formed in the peridotite layer of the Samail Ophiolite have been previously dated by ^{14}C , with ages reported in Clark and Fontes (1990), Clark et al. (1992), Keleman and Matter (2008), Kelemen et al. (2011), and Mervine et al. (2014). Clark and Fontes (1990) and Clark et al. (1992) dated ~50 travertines and travertine veins sampled in the vicinity of four hyperalkaline springs near the town of Nizwa. They found that these travertines and travertine veins varied in age from >modern (post-1950) to ~35,000 yr BP, with one sample “ ^{14}C dead” (older than the ~50,000 year limit of ^{14}C dating). Kelemen and Matter (2008) and Kelemen et al. (2011) dated ~60 carbonates, both travertines and Mg-rich carbonate veins, from numerous locations throughout the southern ~2/3 of the ophiolite. Similar to the previous studies in the Nizwa area, they found that the travertines varied in age from >modern to ~40,000 yr BP. They also found that the Mg-rich carbonate veins had an average ^{14}C age of ~26,000 yr BP, an

important observation since previously many of these veins were believed to be tens of millions of years old. Only two of their samples were ^{14}C dead. Mervine et al. (2014) obtained ^{14}C ages for an additional ~100 Samail carbonates, including several layered travertine terrace sequences. Evaluating all available ^{14}C data, Mervine et al. (2014) concluded that the travertines range in age from >modern to >45,000 yr BP in age and that between ~30,000 and ~45,000 yr BP the travertine deposition rate was ~0.1 to 0.3 mm/yr for the four travertine sequences analyzed. Furthermore, they found that new travertine ^{14}C ages filled in previously observed “gaps” in the Clark and Fontes (1990) dataset, implying that travertine deposition has been fairly continuous and not necessarily controlled by climatic variations, as was previously hypothesized by Clark and Fontes (1990). Notably, Mervine et al. (2014) observed that a significant proportion of Mg-rich carbonate veins sampled at three roadcuts were ^{14}C dead. A location-weighted average indicates that ~40% of Mg-rich carbonate veins sampled at roadcuts are ^{14}C dead. In contrast, no Mg-rich carbonate veins sampled at outcrops (i.e. on the natural weathering surface of the peridotite) are ^{14}C dead. Mervine et al. (2014) concluded that there have been multiple generations of Mg-rich carbonate vein formation and that this vein formation continues to the present day. Furthermore, they speculated that some of the apparent ^{14}C ages for Mg-rich carbonate veins could be affected by open system behavior, such as dissolution and re-precipitation. Overall, the ^{14}C data indicate that carbonation of the peridotite layer of the Samail Ophiolite is a recent and on-going process and not exclusively an ancient one.

3. Methods

3.1 Sample descriptions

Sample descriptions are given in Table 1. Carbonates were collected from four locations (Figure 1) of travertine deposition (Qafeefah Travertine, Misht Travertine, Wadi Uqaybah Travertine, and Wadi Sudari Travertine), from three roadcuts exposing carbonate veins in peridotite (Qafeefah Roadcut, Fanja Roadcut, and Al-Wuqbah Roadcut), and from one campsite located on the natural peridotite weathering surface and containing abundant carbonate veins exposed in outcrop (Wadi Sudari Campsite). All carbonates analyzed for U and Th were also dated with ^{14}C and analyzed for $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, the results of which have been previously reported in Mervine et al. (2014). Three water samples were collected from hyperalkaline pools at Wadi Sudari Travertine, Misht Travertine, and Wadi Dima (near Qafeefah, see Figure 1) during January 2009 by Jürg Matter (University of Southampton). The water samples (500 mL) were filtered and acidified with ultrapure HNO_3 and were later analyzed for U and Th concentrations.

Three types of surficial travertines were analyzed for U and Th: (1.) recently-formed travertine precipitates collected from the bottoms of hyperalkaline pools (two samples), (2.) travertine terraces (four samples), and (3.) travertine “pseudospeleothems” (two samples), which are speleothem-like formations that have developed in overhangs, such as underneath travertine terraces. Note that these features are not true speleothems because they were not formed within caves. Also note that travertine terrace sample OM09-109C-MS is part of a younger travertine terrace forming between weathered, older travertine terraces; these travertines were distinguished as “channel-filling travertines” in Mervine et al. (2014). In addition, U and Th measurements were obtained for one Ca-rich travertine vein.

Mg-rich carbonate veins (13 samples) were also analyzed for U and Th. For the purposes of this study, these veins have been divided into “outcrop veins” (sampled from the natural peridotite weathering surface) and “roadcut veins” (sampled at roadcuts). Both types of veins are interpreted to have formed in the deeper subsurface; however, the roadcut veins are generally less weathered than the outcrop veins, which are often partially eroded. U and Th concentration measurements were also obtained for two serpentine mineral fractions that were separated out of intergrown carbonate-serpentine veins sampled at roadcuts.

Finally, U and Th measurements were obtained for two sediment samples, which are a potential source of detritus in the carbonates. These sediment samples were collected from small wadis or valleys adjacent to the carbonates sampled at Fanja Roadcut and Wadi Uqaybah Travertine. There is no significant soil formation at any of the carbonate sampling locations, so these sediments consist of loose detritus that contains fragments of peridotite and carbonate as well as aeolian particles.

3.2 Geochemical analyses

Carbonates selected for U and Th analysis were crushed using a jaw crusher and then purified through magnetic separation and hand picking. Many of the carbonate veins consist of multiple minerals (e.g. magnesite, calcite, dolomite, serpentine, brucite). Whenever possible, monomineralic (or nearly so) separates were prepared, and their purity was evaluated by powder X-ray diffraction (XRD) analysis at Woods Hole Oceanographic Institution using a Philips Analytical PW1830 XRD machine. Mineral identification was conducted using the International Center for Diffraction Data Powder Diffraction File. While large fragments of peridotite and serpentine were easily removed through magnetic separation and hand picking, fine-grained

aluminosilicate material incorporated into the travertines and, to a lesser extent, into the subsurface carbonate veins was impossible to remove through physical separation. Note that the two recently formed travertine precipitates (OM09-8COPS-MS and OM09-10COPS-MS) were not purified (aside from removing large, obvious peridotite clasts) due to their fine-grained nature.

For travertines, approximately 1 to 2 g of material was required for analysis. For Mg-rich carbonate veins, approximately 20 to 30 g of material was required because of their extremely low U and Th concentrations. For select Mg-rich carbonate veins as well as the two serpentine mineral separates, 1 to 2 g of material was first analyzed for U and Th concentrations in order to identify samples that were most promising for isotopic analysis (i.e. had the highest U concentrations). For the sediments, 2 to 10 g of a bulk powder was analyzed for each sample.

Carbonates selected for ^{230}Th analyses were completely dissolved using HNO_3 , HF , HClO_4 , and H_3BO_3 . To investigate the behavior of samples during dissolution and also the nature of the incorporated aluminosilicate detritus, separate weak HNO_3 leaches were carried out for select samples. The aluminosilicate residues left behind by these weak HNO_3 leaches were also dissolved and analyzed separately. A progressive dissolution experiment was carried out on one travertine sample (OM09-107C-MS). The dissolution experiments are discussed in detail in the Supplementary Information. In brief, the dissolution experiments indicate that both elemental and isotopic fractionation of U and Th occurs during partial dissolution of Samail carbonates. This fractionation is most significant for the surface travertines, which incorporate higher amounts of aluminosilicate detritus than the subsurface carbonate veins. The observed fractionation as a result of partial dissolutions reinforces the need to employ a total dissolution

approach for ^{230}Th dating of terrestrial carbonates that contain significant admixed detritus (e.g. Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991).

After dissolution, U and Th were purified using two columns (see details in Ball et al., 2008 and Sims et al., 2008a). The first was a nitric anion column that removes U and Th from the rock matrix. The second was a hydrochloric anion column that separates U from Th. For large samples, U and Th were first co-precipitated with Fe hydroxide. For the Fe hydroxide precipitation, 3.5 mg of an ultrapure Fe ICP standard manufactured by Ricca Chemical was added to each sample (see Edwards et al., 1986 for discussion of the Fe-hydroxide precipitation step).

U and Th concentrations were measured by isotope dilution ICP-MS using ^{229}Th and ^{233}U spikes. The isotope dilution measurements were carried out on a Thermo Element2 mass spectrometer at Woods Hole Oceanographic Institution following procedures detailed in Sims et al. (2008a). U and Th isotopes were measured on chromatographically separated U and Th aliquots on separate days by MC-ICP-MS on the ThermoFinnegan NEPTUNE at Woods Hole Oceanographic Institution using techniques described in Ball et al. (2008) and Sims et al. (2008a, 2008b). When measuring Th isotopes, ^{230}Th is in the center SEM using the RPQ high abundance sensitivity filter, and ^{232}Th is measured on an up-mass Faraday. Th isotope mass bias and SEM – Faraday yield was corrected for by sample standard bracketing using a $^{230}\text{Th}/^{232}\text{Th}$ atom ratio of $11.380 \pm 0.170 \times 10^{-6}$ for IRMM 35 (see Sims et al., 2008b for details). For U isotopic measurements, ^{234}U was measured on the center SEM, and ^{235}U and ^{238}U were measured on up-mass Faradays. U isotopes were corrected for mass bias using both an internal correction and assuming the samples $^{235}\text{U}/^{238}\text{U} = 0.0072527$ (Cheng et al, 2000) and an external correction using NBL U010 for a mass bias and SEM–Faraday yield using the Richter and Goldberg (2003)

values $^{235}\text{U}/^{238}\text{U} = 0.0101382$ and $^{234}\text{U}/^{238}\text{U} = 5.4483 \times 10^{-5}$ (see Ball et al., 2008 for details). For a matrix matched quality assurance standard, the carbonate standard (JCp-1) was analyzed for U and Th concentrations and isotopes (Tables 2 and 3). In addition, a large compilation of U and Th concentration and isotopic ratio measurements for five synthetic Th standards (IRMM 35, IRMM36, UCSC ThA, WUN, OU ThU) and six rock standards (UCSC TML, Icelandic ATHO, USGS BCR2, USGS W2, USGS BHVO-2, and LV18) that were analyzed at the same time of this Oman study can be found in Sims et al. (2008a), which represents a concerted international effort of six labs (WHOI; UCSC; LLNL; UI Urbana; GEOMOC, AU; Bristol, UK) to cross calibrate these U-series standards. The same instrumentation, methods, and ^{229}Th and ^{233}U spikes were used in both this study and in Sims et al. (2008a). Total procedural blanks, including the acid used for sample introduction, are typically less than <4 pg (1×10^{10} atoms) for ^{238}U and <3 pg (7×10^9 atoms) for ^{232}Th . Blank levels for ^{234}U and ^{230}Th are un-measurable but are approximately 10^{-4} of ^{238}U and 10^{-5} of ^{232}Th , respectively.

4. Results

4.1 Mineralogy

The results of X-ray diffraction analysis of powdered samples are presented in Table 1. Minerals were identified as “major” or “minor” but were not further quantified.

The mineralogy of the Samail carbonates is discussed in detail in Mervine et al. (2014). In brief, the travertine samples consist predominately of calcite with trace quantities of unidentified clay minerals. One recently formed travertine precipitate (OM09-8COPS-MS) also contains minor aragonite and possibly hydromagnesite. The other recently formed travertine precipitate (OM09-12S-MS) contains trace quartz, possibly from aeolian sand incorporated into

the sample. The Mg-rich subsurface carbonate veins consist primarily of magnesite or dolomite except for one sample (OM10-84C-MG), which consists primarily of brucite.

The two sediment samples have distinct mineral compositions. The first sample collected near Fanja Roadcut (OM09-8S-MS) consists primarily of lizardite with minor quartz, dolomite, magnesite, and calcite. The second sample collected near Wadi Uqaybah Travertine (OM09-12S-MS) consists primarily of quartz and calcite with minor chrysotile, dolomite, enstatite, and possibly aragonite. The difference in mineralogy reflects the different detrital inputs from the nearby environment and also from aeolian sources at these two locations.

4.2 U and Th concentrations

Samail carbonates have U and Th concentrations ranging over several orders of magnitude from <1 ppb to ~1 ppm (Table 2 and Figure 4). The travertine terraces and wadi sediments have the highest U and Th concentrations, ranging from ~0.1 to ~1 ppm for both U and Th. The Mg-rich carbonate veins have U and Th concentrations ranging from <1 ppb to ~10 ppb, three or four orders of magnitude lower than most of the travertine U and Th concentrations. Sample OM09-63C-MS, an Mg-rich carbonate vein, has U and Th concentrations comparable to the total procedural blank and thus below detection limit. The two serpentine mineral separates (from serpentine-carbonate veins) have very low (<1 ppb) U and Th concentrations comparable with those of the Mg-rich carbonate veins. The three hyperalkaline (500 mL) water samples analyzed by isotope dilution were well below the instrumental detection limits for U and Th, which are ~1 ppt. Reproducibility of U and Th concentration measurements for JCp-1 (n = 4) is better than the propagated analytical error of the individual measurements. The reproducibility of the travertine samples and Mg carbonate samples is also quite good with the exception of OM09-

47C-MS, which is likely due to this sample's low U and Th concentrations and potential for heterogeneity from fine-grained admixed material.

4.3 U and Th isotopes

U and Th isotopic data are presented in Table 3. Note that in the tables, and throughout this paper, parentheses indicate activities. Due to low U and Th concentrations, only a subset of the samples analyzed for U and Th concentrations could be analyzed for U and Th isotopes.

Many of the Samail carbonates are out of equilibrium with respect to ($^{230}\text{Th}/^{238}\text{U}$) (Figure 5). Samples with ($^{230}\text{Th}/^{238}\text{U}$) = 1 plot on the equiline, indicating isotopic equilibrium. Four of the Mg-rich carbonate outcrop veins, two of the travertine terraces, and the travertine vein have ($^{230}\text{Th}/^{238}\text{U}$) < 1 and thus have ^{238}U excesses. One travertine terrace sample, the two travertine pseudospeleothems, the two recently-formed travertine precipitates, one of the Mg-rich carbonate outcrop vein samples, four Mg-rich carbonate roadcut vein samples, and the two sediment samples fall very close to the equiline.

Many of the Samail carbonates are also out of equilibrium with respect to ($^{234}\text{U}/^{238}\text{U}$) (Figure 6). The travertine terrace samples, recently formed travertine precipitates, and travertine pseudospeleothems all have ^{234}U excesses with ($^{234}\text{U}/^{238}\text{U}$) values ranging from 1.118 to 1.255. However, the travertine vein sample plots on the other side of the equiline with a ($^{234}\text{U}/^{238}\text{U}$) value of 0.832, which may indicate that the hyperalkaline Type II fluids from which this sample was precipitated reacted with material that has low $^{234}\text{U}/^{238}\text{U}$ due to prior water-rock interaction. The Mg-rich carbonate outcrop vein samples have a fairly narrow range of ($^{234}\text{U}/^{238}\text{U}$) values of 1.187 to 1.237. The Mg-rich carbonate roadcut vein samples have lower ($^{234}\text{U}/^{238}\text{U}$) values than the outcrop vein samples, ranging from 0.998 to 1.123. Two of the roadcut vein samples have

near-equilibrium ($^{234}\text{U}/^{238}\text{U}$) values of 0.998 (Sample OM09-58C-MS) and 1.010 (Sample OM09-57C-MS). Sediment sample OM09-8S-MS plots just above the equiline with ($^{234}\text{U}/^{238}\text{U}$) = 1.032 while sediment sample OM09-12S-MS, which contains significant calcite that likely derives from nearby travertines, has higher ($^{234}\text{U}/^{238}\text{U}$) = 1.102, which approaches the values observed in the surface travertines.

There is a large range of ($^{230}\text{Th}/^{232}\text{Th}$) values observed in the Samail carbonates and sediments (Figure 7). Most samples have ($^{230}\text{Th}/^{232}\text{Th}$) values that fall in the range of ~0 to 6. However, the Mg-rich carbonate roadcut vein sample OM09-57C-MS has ($^{230}\text{Th}/^{232}\text{Th}$) = 30.515.

5. Discussion

5.1 Systematics of U and Th in Samail carbonates and sediments

5.1.1 Origin of variable U and Th concentrations

The travertine samples, particularly the surface travertine samples, have the highest U and Th concentrations (Figure 4). The high U and Th concentrations of the travertines are likely a result of incorporation of a significant amount of aluminosilicate detritus. There are multiple lines of evidence that support this interpretation. First, while the higher U concentrations of the travertines could be explained by U enrichment in fluids due to low water/rock ratios, the Th concentrations can only be explained by the incorporation of detritus since Th is insoluble under almost all conditions and is believed to be insoluble in both the Type I and the Type II waters flowing through Samail peridotite (e.g. Langmuir and Herman, 1980; Porcelli and Swarzenski, 2003). Second, light acid leaches of travertine samples (see Supplementary Information) leave behind significant aluminosilicate detritus, which has higher U and Th concentrations than the

carbonate material dissolved during the light acid leach. In contrast, most of the Mg-rich carbonate vein samples do not leave behind significant aluminosilicate residues after light acid leaches. Third, the U and Th concentrations of the travertine vein sample (OM09-84C-MS), which formed in the subsurface and likely incorporated less detritus, are lower than the other travertines. Finally, the travertine samples have U and Th concentrations that are comparable to the wadi sediments, which are likely sources of detritus.

Since the travertine samples incorporate significant detritus, constraining the composition of this component is essential to correct for detrital Th in the ^{230}Th age calculations. While fragments of serpentine and altered peridotite are observed in the travertines, these cannot be the only sources of detritus in the travertine samples because the travertine terrace samples have U and Th concentrations that are much higher than those measured in serpentine and partially serpentinized peridotites (Figure 4; peridotite data from Hanghøj et al., 2010). However, the two wadi sediments have U and Th concentrations that are comparable with those of the travertine terrace samples. The wadi sediments contain serpentine and enstatite, which likely originate from local peridotite bedrock (Table 1). They also contain significant modal quartz, which is most likely aeolian in origin, and significant carbonate minerals, such as calcite, dolomite, and magnesite. The magnesite likely originates from local magnesite veins in peridotite. However, the calcite and dolomite could originate from local carbonate alteration products in peridotite, such as weathered travertines, and from non-local carbonates transported by wind. For example, aeolian carbonate could originate from the Cretaceous Natih formation, which has U concentrations up to ~9 ppm (Figure 4; Natih carbonate data from Fleitmann et al., 2003). The two pseudospeleothems also have lower U and Th concentrations than the travertine terrace samples. This is likely because the formation beneath overhangs and the fine-grained texture of

these pseudospeleothems reduces the amount of detritus that they incorporate. The low U concentrations of the Mg-rich carbonate vein samples reflect the low U concentrations of the peridotites (e.g. Hanghøj et al., 2010) that are altering to form these carbonates, and low U concentrations in the reacting fluids (Figure 4).

5.1.2 Systematics of U and Th isotopes

The Samail carbonates and sediments either have ^{238}U excesses or fall on the equiline of the $(^{230}\text{Th}/^{232}\text{Th})$ versus $(^{238}\text{U}/^{232}\text{Th})$ plot (Figure 5). During initial precipitation from fluids, Samail carbonates should have ^{238}U excesses. Over time, as the ^{238}U decay chain returns to equilibrium, the isotopic compositions of the carbonates will move toward the equiline. After approximately 375,000 years (5 times the half-life of ^{230}Th , which equates to an uncertainty of 3% in $(^{230}\text{Th}/^{238}\text{U})$), the isotopic compositions of the carbonate samples will plot on the equiline. Thus, one interpretation is that the carbonate samples which plot on or close to the equiline (five travertines, one Mg-rich carbonate outcrop vein sample, and four Mg-rich carbonate roadcut veins sample) are 375,000 years or older in age. This interpretation may be valid for the carbonate veins, which incorporate lower amounts of aluminosilicate detritus, but may not be valid for the surface travertines.

As discussed previously, the travertine samples likely contain significant aluminosilicate detritus, which explains their elevated U and Th concentrations relative to the subsurface carbonates. Therefore, travertine samples that plot near or on the $(^{230}\text{Th}/^{232}\text{Th})$ versus $(^{238}\text{U}/^{232}\text{Th})$ equiline could either: (1.) be ~375,000 years or older in age or (2.) have incorporated a significant quantity of equilibrium (or near-equilibrium) detritus that overwhelms the isotopic signature of the pure travertine. The second explanation is favored since the two

recently formed travertine precipitates, which have >modern (post-1950) ^{14}C ages (Mervine et al., 2014), fall on the equiline. These precipitates are known to have formed very recently, so old ^{230}Th ages for these samples can be rejected. Therefore, these samples must have incorporated significant near-equilibrium detritus, such as wadi sediments, two samples of which (OM09-8S-MS and OM09-12S-MS) fall close to the equiline. The isotopic compositions of the other travertine samples are also likely affected by the incorporation of aluminosilicate detritus. The fact that the two wadi sediment samples plot close to the equiline is not unexpected. Most aluminosilicate detritus (see the data compilation in the Supplementary Information) falls on the equiline or has slight ^{230}Th excesses on a ($^{230}\text{Th}/^{232}\text{Th}$) versus ($^{238}\text{U}/^{232}\text{Th}$) plot. Aluminosilicate detritus rarely has ^{238}U excesses (e.g. Szabo and Rosholt, 1982; Porcelli and Swarzenski, 2003). This is because rocks generally weather under oxidizing conditions present at and near the Earth's surface, where U is soluble and will be leached from rocks (e.g. Porcelli and Swarzenski, 2003).

5.2 Approaches and limitations for determining ^{230}Th ages for Samail carbonates

5.2.1 ^{230}Th dating of terrestrial carbonates

^{230}Th dating of carbonates measures the ingrowth of ^{230}Th from the decay of ^{238}U and ^{234}U incorporated into the carbonate matrix during formation. Since Th is generally highly insoluble relative to U, carbonate that forms (either through inorganic precipitation or biological processes) from marine and terrestrial waters generally contains relatively high U concentrations and low Th concentrations. This creates disequilibria in the ^{238}U decay chain, which gradually returns to equilibrium as ^{238}U (half-life: $\sim 4.47 \times 10^9$ years) and ^{234}U (half-life: $\sim 2.45 \times 10^5$ years) decay to ^{230}Th (half-life: $\sim 7.56 \times 10^4$ years).

²³⁰Th dating of carbonates relies on two primary assumptions: (1.) there was no initial ²³⁰Th incorporated when the carbonate formed and (2.) there has been no loss or addition of U and Th since carbonate formation. In practice, these two assumptions are valid only for unrecrystallized, pure carbonates such as some corals and speleothems (e.g. Edwards et al., 2003; Scholz and Hoffmann, 2008). For many carbonates, these assumptions are not valid and corrections must be employed in the dating technique. If initial ²³⁰Th is present in a carbonate sample, then it may be possible to correct for this initial ²³⁰Th (e.g. Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991; Kaufman, 1993; Edwards et al., 2003). Carbonates affected by open system behavior after formation, such as recrystallized corals and lacustrine carbonate with added hydrogenous U or Th, are challenging to date unless the open system behavior can be estimated using models (e.g. Fontes et al., 1992; Thompson et al., 2003; Villemant and Feullet, 2003; Haase-Schramm et al., 2004).

Terrestrial carbonates are usually affected by the presence of initial ²³⁰Th and, to a lesser extent, by ²³⁴U and ²³⁸U contributed by detrital material (e.g. Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991; Kaufman, 1993). Detritus incorporated into terrestrial carbonates is often fine-grained and intergrown with the carbonate, making physical separation very difficult (e.g. Ku and Liang, 1984; Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991). Various approaches have been attempted over the years to account for detrital ²³⁰Th, ²³⁴U, and ²³⁸U in terrestrial carbonates to enable dating of these carbonates. Early studies (e.g. Harmon et al., 1977) attempted to separate the carbonate material from the admixed detritus by employing light acid leaches. However, numerous subsequent studies (e.g. Schwarcz and Latham, 1989; Przybylowicz et al., 1991; Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991) have demonstrated that fractionation of U and Th concentrations and shifts in isotopic ratios can occur during incomplete acid leaching.

Fractionation of U and Th concentrations and isotopes was also observed during partial dissolution experiments conducted on our Samail carbonate samples (see Supplementary Information). A number of studies have investigated and employed pseudoisochron methods for determining the composition of incorporated detritus (e.g. Osmond et al., 1970; Schwarcz and Latham, 1989; Przybylowicz et al., 1991; Bischoff and Fitzpatrick, 1991; Luo and Ku, 1991). However, pseudoisochron methods are limited by the requirement of a set of coeval samples with variable detritus/carbonate ratios and thus are not suitable for dating the Samail carbonates analyzed in this study since the Samail travertines and Mg-rich carbonate veins are known to have a range of ages based on previous ^{14}C dating (Mervine et al., 2014).

For most terrestrial carbonates, employing total sample dissolutions with corrections for detrital ^{230}Th (and sometimes also ^{234}U and ^{238}U) is the most robust way to determine ^{230}Th ages for the carbonates (e.g. Ludwig and Titterton, 1994; Edwards et al., 2003; Ludwig et al., 2011). The simplest and most commonly employed detrital correction approach uses ^{232}Th as an indicator isotope for the ^{230}Th correction. In this method, all ^{232}Th is assumed to originate from detritus, and the measured ($^{232}\text{Th}/^{238}\text{U}$) of the terrestrial carbonate is used to weight the ^{230}Th correction. The ($^{230}\text{Th}/^{232}\text{Th}$) composition of the detrital material is rarely known explicitly and is generally the largest source of uncertainty in corrected ^{230}Th ages. Often, a Bulk Silicate Earth (BSE) value for ($^{230}\text{Th}/^{232}\text{Th}$) of 0.814 (calculated assuming equilibrium with ($^{238}\text{U}/^{232}\text{Th}$) = ($^{230}\text{Th}/^{232}\text{Th}$)) is assumed for the detritus (e.g. Ludwig and Titterton, 1994; Ludwig et al., 2011). Although the required assumptions are not necessarily geologically sound, assuming a BSE ($^{230}\text{Th}/^{232}\text{Th}$) value works for older carbonates with high U and low ^{232}Th contents because the resulting correction is relatively insignificant, such that the assumed ($^{230}\text{Th}/^{232}\text{Th}$) value has a small effect on the calculated age (e.g. Bischoff and Fitzpatrick, 1991; Kaufman, 1993; Ludwig

and Titterton, 1994). However, for young carbonates and for carbonates with relatively low U concentrations and high ^{232}Th concentrations (such as the Samail carbonates), the assumed ($^{232}\text{Th}/^{230}\text{Th}$) value for the detrital material has a large effect on the calculated age, and therefore the ($^{232}\text{Th}/^{230}\text{Th}$) value must be known explicitly in order to reliably determine corrected ^{230}Th ages.

We have employed total dissolutions with corrections for initial ^{230}Th in an attempt to determine ^{230}Th ages for Samail carbonates. We have not employed corrections for detrital ^{234}U and ^{238}U because the initial ^{230}Th correction has the largest influence on the age corrections.. Because Samail carbonate ^{230}Th ages are so sensitive to initial ^{230}Th corrections and because there is not an obvious single detrital ($^{230}\text{Th}/^{232}\text{Th}$) value to use for the corrections, a sensitivity analysis approach has been employed to evaluate the influence of detrital ^{230}Th on the calculated ^{230}Th ages of the Samail carbonates.

To evaluate a reasonable range of detrital ($^{230}\text{Th}/^{232}\text{Th}$) values to use for the initial ^{230}Th correction for the Samail carbonate ^{230}Th ages, we compiled 300 literature values for ($^{230}\text{Th}/^{232}\text{Th}$) measured in various types (soil, saprolite, alluvium, colluvium, till, and volcanic ash) of aluminosilicate detritus (see Supplementary Information). These data were compiled from Rosholt et al. (1966), Hansen and Stout (1968), Hansen (1970), Rosholt et al. (1985), Mathieu et al. (1995), Dequincey et al. (2002), Krishnaswami et al. (2004), Dosseto et al. (2008), Pelt et al. (2008), and Ma et al. (2010). This compilation has ($^{230}\text{Th}/^{232}\text{Th}$) values ranging from 0.02 to 4.61 (average: 0.77, 2σ standard deviation: 0.50). While the average ($^{230}\text{Th}/^{232}\text{Th}$) of the compiled detritus values is fairly close to the equilibrium BSE value of 0.814, aluminosilicate detritus clearly has a wide range of ($^{230}\text{Th}/^{232}\text{Th}$) values. Our results show that the ($^{230}\text{Th}/^{232}\text{Th}$) value of the detritus incorporated into the Samail carbonates may be higher than the equilibrium

BSE value. The recently formed travertine precipitates OM09-8COPS-MS and OM09-10COPS-MS have ($^{230}\text{Th}/^{232}\text{Th}$) values of 1.376 and 1.416, respectively. These two precipitates are essentially zero age with respect to the half-life of ^{230}Th (based on ^{14}C measurements and field relations, see Mervine et al., 2014). While the ($^{230}\text{Th}/^{232}\text{Th}$) value of detritus incorporated in Samail carbonates is clearly spatially and temporally variable, it is important to note that the ($^{230}\text{Th}/^{232}\text{Th}$) values measured in the recently-formed precipitates (likely formed a matter of days or weeks before collection and therefore should have minimal ingrown ^{230}Th) are similar and provide a possible correction value to use for older travertine samples. ($^{230}\text{Th}/^{232}\text{Th}$) values measured in wadi sediments also provide possible values to use for initial ^{230}Th corrections for the Samail carbonate ^{230}Th ages. One of the wadi sediments has a ($^{230}\text{Th}/^{232}\text{Th}$) value of 2.093, somewhat similar to the values measured in the two recently-formed travertine precipitates. However, wadi sediment OM09-8S-MS, which consists primarily of serpentine, has a higher ($^{230}\text{Th}/^{232}\text{Th}$) value of 5.215. Clearly, there is significant variability in the measured ($^{230}\text{Th}/^{232}\text{Th}$) values of detritus that may have been incorporated into the Samail carbonates.

5.2.2 ^{230}Th dating of Samail travertines

Uncorrected ^{230}Th ages for Samail travertines are presented in Table 3. For all travertines analyzed, the uncorrected ^{230}Th ages are significantly older than the corresponding ^{14}C ages (Mervine et al., 2014). The low U concentrations and relatively high U/Th ratios found in the Samail travertines makes obtaining detritus corrected ^{230}Th ages highly challenging. The travertines contain significant detrital Th, and the values of corrected ^{230}Th ages are thus highly sensitive to the selected value of initial ($^{230}\text{Th}/^{232}\text{Th}$) employed in the corrections. Varying the assumed initial ($^{230}\text{Th}/^{232}\text{Th}$) over the range observed in wadi sediments, BSE, and our

564 compilation of aluminosilicate detritus (see Supplementary Information) changes the corrected
565 travertine ^{230}Th ages by tens of thousands of years. As an example, Figure 8 displays the effect
566 of the selected initial ($^{230}\text{Th}/^{232}\text{Th}$) value on the corrected ^{230}Th age for travertine terraces sample
567 OM09-106C-MS-A. Similar figures for all travertine samples are presented in the Supplementary
568 Information.

569 For all travertine samples, geologically plausible detrital ($^{230}\text{Th}/^{232}\text{Th}$) values bring the
570 corrected ^{230}Th ages into concordance with the ^{14}C ages presented in Mervine et al. (2014). For
571 example, an assumed detrital ($^{230}\text{Th}/^{232}\text{Th}$) value of ~ 1.7 produces a corrected ^{230}Th or sample
572 OM09-106C-MS-A that is concordant with the ^{14}C age of $30,023 \pm 235$ yr BP reported in
573 Mervine et al. (2014). Unfortunately, no single detrital ($^{230}\text{Th}/^{232}\text{Th}$) value consistently corrects
574 the travertine ^{230}Th ages to the corresponding ^{14}C ages. The different ($^{230}\text{Th}/^{232}\text{Th}$) corrections
575 required for each travertine are likely a result of the observed heterogeneous nature of the
576 detritus. Therefore, although geologically plausible ($^{230}\text{Th}/^{232}\text{Th}$) values can be selected for each
577 travertine to bring the corrected ^{230}Th age into concordance with the ^{14}C age, the lack of a
578 uniform ($^{230}\text{Th}/^{232}\text{Th}$) correction value precludes obtaining precise ages for Samail travertines
579 using ^{230}Th dating alone.

581 5.2.3 ^{230}Th dating of Samail Mg-rich carbonate veins

582 Uncorrected ^{230}Th ages for Mg-rich carbonate vein samples are presented in Table 3.
583 Four out of the five Mg-rich carbonate outcrop vein samples have uncorrected ^{230}Th ages that are
584 younger than the corresponding ^{14}C ages (Mervine et al., 2014). The exception is sample OM10-
585 84C-MG, which has a ^{230}Th age older than its ^{14}C age. While not particularly high, the ~ 2 ppb
586 Th concentration of sample OM10-84C-MG is significant relative to its low U concentration of

~1 ppb. Therefore, sample OM10-84C-MG likely contained significant initial ^{230}Th relative to its U concentration. Furthermore, the Th/U ratio (1.995) of OM10-84C-MG is significantly higher than the other carbonate outcrop veins, which have Th/U ratios of ~0.2 to ~0.3.

Since the uncorrected ^{230}Th ages of the Mg-rich carbonate outcrop veins are generally younger than the corresponding ^{14}C ages, corrections for initial ^{230}Th from detritus cannot be employed to make the two sets of ages concordant since initial ^{230}Th corrections would make the ^{230}Th ages even younger. Figure 9 illustrates the effect of a range of initial ^{230}Th corrections for Mg-rich carbonate outcrop vein OM10-13C-MG. Similar figures for all of the Mg-rich carbonate outcrop vein samples are presented in the Supplementary Information. Possibly, a detrital correction employing an unusual detritus with ^{238}U enrichment, such as a highly altered peridotite, could be employed to make the ^{230}Th ages older.

There are two possible end member explanations for the discrepancies between the ^{230}Th and ^{14}C ages for the Mg-rich carbonate outcrop vein samples. The first explanation is that the ^{230}Th ages are correct but the ^{14}C ages are too old because of the incorporation of recycled, ^{14}C dead carbon. The second explanation is that the ^{230}Th ages are too young because of ^{238}U addition from U-bearing fluids. Essentially, hexavalent U, which is highly soluble, is being carried down from the surface by the oxidized Type I waters; this U is subsequently deposited as tetravalent U by the Type II reducing serpentinization fluids. We also note that if the carbonate outcrop veins were modified by U addition from fluid, then their ^{14}C ages may have also been partially or fully re-set. Therefore, because of the possible effects of incorporated old carbon with no ^{14}C and/or open system addition of young carbon with high ^{14}C , the ^{14}C ages of altered Mg-rich carbonate veins should be interpreted with caution. The ^{14}C ages could be affected by both types of end member open system behavior, including multiple generations of open system

behavior. The uncorrected ^{230}Th ages for the Mg-rich carbonate roadcut vein samples are significantly older than the uncorrected ^{230}Th ages for the Mg-rich carbonate outcrop vein samples. The roadcut vein samples have ($^{230}\text{Th}/^{238}\text{U}$) values approaching equilibrium. Mg-rich carbonate roadcut vein samples OM09-35C-MS, OM09-55C-MS, and OM09-57C-MS have ($^{230}\text{Th}/^{238}\text{U}$) values of 1.068, 0.956, and 0.984, respectively. Furthermore, Mg-rich roadcut vein samples OM09-57C-MS, and OM09-58C-MS have near-equilibrium ($^{234}\text{U}/^{238}\text{U}$) values of 1.010, and 0.998, respectively. These four Mg-rich carbonate roadcut vein samples are ^{14}C dead, suggesting that their ages are >50,000 yr BP (Mervine et al., 2014). Thus, the older ^{230}Th ages suggested by the near-equilibrium ($^{230}\text{Th}/^{238}\text{U}$) and ($^{234}\text{U}/^{238}\text{U}$) values are consistent with the ^{14}C data. As shown in Figure 10, applying a wide range of ($^{230}\text{Th}/^{232}\text{Th}$) detrital values for an initial ^{230}Th correction does not affect the ^{230}Th age of sample OM09-57C-MS significantly. Unfortunately, Th isotopes could not be measured in Sample OM09-58C-MS due to its low Th concentration. Given the long half-life of ^{234}U ($t_{1/2} = 245.3$ kyr), the near-equilibrium ($^{234}\text{U}/^{238}\text{U}$) values for these two veins suggest that these samples are several hundred thousand years old. An alternate explanation would be that these samples started out with ($^{234}\text{U}/^{238}\text{U}$) in equilibrium. However, we consider this an unlikely initial condition given: (1.) the extent of ($^{234}\text{U}/^{238}\text{U}$) disequilibria observed in the other Mg-carbonate samples (Figure 6); (2.) that most crustal waters (seemingly including those forming these veins, see Figure 6) are enriched in ^{234}U relative to ^{238}U , because of well-documented alpha recoil and lattice damage effects (Porcelli and Swarzenski, 2003); and (3.) these inferred old ages are consistent with other isotopic characteristics which also indicate that they are old, namely equilibrium ($^{230}\text{Th}/^{238}\text{U}$) and ^{14}C dead radiocarbon measurements.

Mg-rich carbonate roadcut vein sample OM10-52C-MG has a ($^{230}\text{Th}/^{238}\text{U}$) value of 1.542, which is a value that cannot be reached through closed-system ^{238}U decay and ^{230}Th ingrowth. Therefore, no ^{230}Th age can be determined for this sample. The high ($^{230}\text{Th}/^{238}\text{U}$) value of this sample is likely explained by high detrital ^{230}Th since it has a very high Th/U ratio of 5.6.

5.3 Implications for natural rates of carbonate formation in Samail peridotite

The determination of precise ^{230}Th ages for Samail carbonates is limited by large and uncertain initial ^{230}Th corrections. Nevertheless, the U and Th systematics and the age limits from the ^{230}Th dating attempts permit further refinement of previous estimates of natural rates of carbon sequestration in Samail carbonates (Kelemen and Matter, 2008; Kelemen et al., 2011; Mervine et al., 2014).

The fact that geologically reasonable initial ^{230}Th corrections bring ^{230}Th and ^{14}C ages for travertines into concordance further supports prior inferences that ^{14}C dating of travertines is generally reliable and these ages can be interpreted as formation ages. This places additional confidence on the estimates, based on ^{14}C dating, that travertine precipitation rates are ~ 0.1 to ~ 0.3 mm / yr and that travertines in the Samail Ophiolite are sequestering a maximum of ~ 1 to 3×10^6 kg CO_2 / yr (Mervine et al., 2014).

U and Th systematics and ^{230}Th ages for the Mg-rich carbonate vein samples also provide further evidence that there are multiple generations of Mg-rich carbonate veins in the peridotite subsurface. The fact that the Mg-rich carbonate outcrop veins have ^{230}Th ages younger than the corresponding ^{14}C ages suggests that these carbonate veins may have been altered by U-bearing fluids within the last 375,000 years. The lack of concordance between ^{230}Th and ^{14}C ages for the Mg-rich carbonate outcrop veins also suggests that these carbonate veins may have been affected

by open system behavior, most likely dissolution and re-precipitation of previous generations of carbonate veins. Such open system behavior may have perturbed both the ^{14}C and ^{230}Th ages for Mg-rich carbonate veins.

The near-equilibrium ($^{230}\text{Th}/^{238}\text{U}$) and ($^{234}\text{U}/^{238}\text{U}$) values measured in the less weathered, ^{14}C dead, Mg-rich carbonate roadcut veins suggest that, at certain exposures such as Fanja Roadcut, there may be much older (possibly older than 1 million years, based on the equilibrium $^{234}\text{U}/^{238}\text{U}$ values) generations of Mg-rich carbonate veins preserved at these localities. Therefore, carbonate vein formation in the peridotite layer of the Samail Ophiolite has likely been ongoing for at least the past 50,000 years and possibly for a much longer time. Hence, sensitivity analysis, such as that presented in Mervine et al. (2014), that considers the impact of multiple generations of carbonate vein formation on the estimated rate of CO_2 sequestration in subsurface carbonate veins is essential since there is no single “average” carbonate vein age. The sensitivity analysis presented in Mervine et al. (2014) indicates that ongoing Mg-rich carbonate vein formation in the peridotite layer of the Samail Ophiolite sequesters on the order of 10^7 kg of CO_2 per year, consistent with less well-constrained, previous estimates (Kelemen and Matter 2008; Kelemen et al. 2011).

5.4 Implications for U-Th disequilibria dating of terrestrial Quaternary carbonates

The results of this study indicate that U-Th disequilibria dating of Quaternary carbonate alteration minerals formed in ophiolite peridotites is highly challenging, due to low U concentrations, high Th/U ratios, and the open system nature of some of the carbonates. This study concurs with previous investigations (e.g. Schwarcz and Latham, 1989; Przybylowicz et al., 1991; Kaufman, 1993) that partial dissolution approaches are not recommended for U-Th

disequilibria dating of terrestrial carbonates, due to fractionation of U and Th concentrations and isotopic values. Unfortunately, U-Th disequilibria dating approaches employing total dissolutions with detrital corrections (e.g. Kaufman, 1993; Ludwig et al., 2003) were also challenging to implement. The travertine samples analyzed in this study contained high amounts of detritus, which appears to be heterogeneous in nature with no single ($^{230}\text{Th}/^{232}\text{Th}$) correction value that can be employed to determine reliable ages that are comparable with previously obtained ^{14}C ages. The very low U concentrations of the subsurface carbonate vein samples required the dissolution of large amounts of carbonate material. In addition, although the subsurface carbonate vein samples contained significantly less detritus than the travertines, the very low U concentrations of these samples means that the age calculations remain highly sensitive to detrital corrections. Furthermore, combined analysis of U-Th disequilibria systematics and ^{14}C dating indicates that some of the subsurface carbonate veins have been affected by open system behavior. In this case, neither the U-Th disequilibria nor the ^{14}C ages can be considered reliable. However, investigation of U-Th systematics was a useful tool for identifying this open system behavior, even though U-Th disequilibria ages could not be determined.

The determination of precise U-Th disequilibria ages for carbonate alteration minerals formed in peridotites with low U concentrations is likely to remain highly challenging. Thus, other dating techniques, such as (U-Th)-He dating of magnetite, an alteration product formed in association with serpentine minerals and also carbonate (e.g. Hernandez Goldstein et al., 2014), may have to be explored to better understand timescales of natural peridotite alteration. However, U-Th disequilibria dating techniques should remain useful for dating carbonate

alteration products in seafloor peridotites since these carbonates commonly have higher U concentrations due to U contributions from seawater (e.g. Ludwig et al., 2011).

U-Th disequilibria dating techniques, as well as U-Pb dating techniques, could also potentially prove useful for dating carbonate alteration products that form in other types of igneous rocks that have higher U concentrations. For example, these techniques could potentially be used to determine ages of kimberlite carbonates. Although kimberlites, like peridotites, are ultramafic rocks comprised largely of olivine (as a primary mineral and as macrocrysts) and pyroxene (as xenocrysts), they have significantly higher U concentrations (ppm level) than peridotites since they are formed through low degrees of mantle melting (e.g. Mitchell, 1986). Potentially, the U-Th disequilibria and U-Pb dating techniques could help kimberlite geologists better understand which carbonates are primary (i.e. magmatic) and which are secondary (i.e. formed through alteration processes). However, some work will be required to better understand the distribution of U and Th in kimberlite minerals. For example, U will most likely be concentrated in minerals such as apatite and perovskite (e.g. Mitchell, 1986), and magmatic carbonates could potentially contain significant Th. Although challenges remain and these must be properly understood and accounted for, U-Th disequilibria dating nevertheless remains an important tool to consider for the dating of several types of terrestrial carbonates, including some types of carbonate alteration minerals.

6. Conclusions

This study highlights the difficulties of using ^{238}U - ^{234}U - ^{230}Th disequilibria measurements to determine ages for carbonate alteration minerals formed in peridotite. Our efforts to use ^{238}U -

^{234}U - ^{230}Th measurements to date both travertines and Mg-rich carbonate veins in the peridotite layer of the Samail Ophiolite show:

(1.) Samail carbonates have extremely low U and Th concentrations, which reflect the low U and Th concentrations found in Samail peridotites and associated groundwaters. The higher U and Th concentrations observed in surface travertine terraces relative to subsurface carbonate veins likely result from the higher amounts of aluminosilicate detritus incorporated into these surface samples. The low U concentrations and relatively high Th/U ratios found in Samail travertines make obtaining ^{230}Th ages highly challenging.

(2.) The ($^{238}\text{U}/^{232}\text{Th}$) versus ($^{230}\text{Th}/^{232}\text{Th}$) pseudoisochron method cannot be rigorously used to date the Samail carbonate alteration minerals. This method requires coeval samples, and the Samail travertines and Mg-rich carbonate veins have a large range of ^{14}C ages (Mervine et al., 2014) relative to the half-life of ^{230}Th . The travertine ages in particular can vary by thousands of years over short distances (mm to cm scale).

(3.) Partial dissolution experiments to separate the fine-grained, inter-grown aluminosilicate detritus from the carbonate show that even light acid leaching significantly perturbs U/Th and ($^{230}\text{Th}/^{232}\text{Th}$) of the dissolved carbonates. These isotopic and elemental perturbations are from partial leaching of the fine-grained silicates. Because this leached out silicate contribution cannot be explicitly quantified, partial dissolution is not a viable method for dating the carbonate.

(4.) For the travertines, because of the issues outlined in conclusions 1, 2 and 3 above, the most viable method for obtaining reliable ^{238}U - ^{234}U - ^{230}Th ages is total dissolution and correcting for the ^{230}Th contributed by the detritus. Because these travertines contain significant initial ^{230}Th from detritus, the values of the ages obtained through correction for ^{230}Th from detritus are highly sensitive to the selected value of initial ($^{230}\text{Th}/^{232}\text{Th}$) used in the correction. For all travertine samples, varying the initial ($^{230}\text{Th}/^{232}\text{Th}$) used in the correction changes the ages of the travertines by tens of thousands of years. However, geologically plausible initial ($^{230}\text{Th}/^{232}\text{Th}$) values correct the travertine ^{230}Th ages to the corresponding ^{14}C ages (Mervine et al., 2014). This age concordance suggests that the travertines are closed systems, adding a level of credence to the reliability of the ^{14}C ages (Mervine et al., 2014) and even the U-series dates (albeit with large uncertainties) obtained in this study.

(5.) Most Mg-rich carbonate veins from outcrops on the natural peridotite weathering surface have uncorrected ^{230}Th ages that are younger than the corresponding ^{14}C ages. The lack of concordance between ^{230}Th and ^{14}C ages for the Mg-rich carbonate outcrop veins indicates that these carbonate veins have been affected by open system behavior, most likely dissolution and re-precipitation of previous generations of carbonate veins and old carbon. Alternatively, the continuous addition of ^{238}U from fluids would make the ^{230}Th ages too young. If the carbonate outcrop veins were modified by U addition from fluid, then their ^{14}C ages may have also been partially or

fully re-set and thus the ^{14}C ages of altered Mg-rich carbonate veins should be interpreted with caution.

(6.) Some ^{14}C dead, Mg-rich carbonate veins from roadcuts have ($^{230}\text{Th}/^{238}\text{U}$) and ($^{234}\text{U}/^{238}\text{U}$) values approaching equilibrium and are likely $\gg 375,000$ years in age.

The variable (young and old) ages for the Mg-rich carbonate veins support the conclusions of ^{14}C dating (Clark and Fontes, 1990; Clark et al., 1992; Kelemen and Matter, 2008; Kelemen et al., 2011; Mervine et al., 2014) that carbonation of the peridotite layer of the Samail Ophiolite is an ongoing process and that there are multiple generations of subsurface carbonate vein formation that must be considered when estimating natural rates of CO_2 sequestration in the peridotite layer of the Samail Ophiolite.

(7.) Overall, this study has shown that using ^{238}U - ^{234}U - ^{230}Th disequilibria measurements to determine ages for carbonate alteration minerals formed in peridotite is challenging due to the generally low U concentrations and high Th/U ratios of the carbonates. In particular, further development is required to better account for the isotopic signature of admixed aluminosilicate detritus. Nevertheless, this study has shown that investigation of U-Th systematics is a useful tool for identifying open system behavior in some carbonate alteration minerals, even when precise U-Th disequilibria ages cannot be determined. In addition, U-Th disequilibria dating techniques remain useful for dating carbonate alteration products in seafloor peridotites since these carbonates commonly have higher U concentrations due to U contributions from seawater (e.g. Ludwig et al., 2011). U-Th disequilibria dating techniques could also

potentially prove useful for dating carbonate alteration products that form in other types of igneous rocks that have higher U concentrations, such as basalts and kimberlites.

7. Acknowledgments

Ali Al-Rajhi, Ministry of Commerce and Industry, Sultanate of Oman, and Sobhi Nasir, Sultan Qaboos University, are thanked for logistical support during field seasons. The Oman field teams are thanked for assistance with fieldwork and for scientific discussions. Jerzy Blusztajn is thanked for mentorship and assistance in the laboratory, and Vera Pavel is thanked for assistance with MATLAB programming. Jürg Matter is thanked for providing the water samples. This research was funded by National Science Foundation grants EAR-1049281 and EAR-1049905, the Deep Ocean Exploration Institute at Woods Hole Oceanographic Institution, the Academic Programs Office at Woods Hole Oceanographic Institution, the Mellon Foundation, a Research Initiative in Science and Engineering Grant from Columbia University to Peter Kelemen and Jürg Matter, and Peter Kelemen's Arthur D. Storke Chair at Columbia University. Lucy McGee and an anonymous reviewer are thanked for comments which improved the paper. David Hilton is thanked for editorial handling.

8. References

- Ball, L., Sims, K. W. W., Schwieters, J., 2008. Measurement of $^{234}\text{U}/^{238}\text{U}$ and $^{230}\text{Th}/^{232}\text{Th}$ in volcanic rocks using the Neptune MC-ICP-MS. *Journal of Analytical Atomic Spectroscopy* 23: 173-180.
- Barnes, I., LaMarche, V. C., Himmelberg, G., 1967. Geochemical evidence of present day serpentinization. *Science* 156, 830-832.

- Barnes, I., O'Neil, J. R., 1969. The relationship between fluids in some fresh Alpine type ultramafics and possible modern serpentinization, Western United States. Geological Society of America Bulletin 80, 1947-1960.
- Barnes, I., O'Neil, J. R., 1971. Calcium-magnesium carbonate solid solutions from Holocene conglomerate cements and travertines in the Coast Range of California. Geochimica et Cosmochimica Acta 35, 699-718.
- Barnes, I., O'Neil, J. R., Trescases, J. J., 1978. Present-day serpentinization in New Caledonia, Oman, and Yugoslavia. Geochimica et Cosmochimica Acta 42, 144-145.
- Beinlich, A., Austrheim, H., 2012. *In situ* sequestration of atmospheric CO₂ at low temperature and surface cracking of serpentinized peridotite in mine shafts. Chemical Geology 332-333, 32-44.
- Bischoff, J. L., Fitzpatrick, J. A., 1991. U-series dating of impure carbonates: An isochron technique using total-sample dissolution. Geochimica et Cosmochimica Acta 55, 543-554.
- Bodinier, J.-L., Godard, M., 2014. Orogenic, Ophiolitic, and Abyssal Peridotites. In: Holland, H., Turekian, K. (Eds.) Treatise on Geochemistry, 2nd Edition, 103-166.
- Brennecke, G. A., Borg, L. E., Hutcheon, I. D., Sharp, M. A., Anbar, A. D., 2010. Natural variations in uranium isotope ratios of uranium ore concentrates: Understanding the ²³⁸U/²³⁵U fractionation mechanism. Earth and Planetary Science Letters 291, 228-233.
- Bruni, J., Canepa, M., Chiodini, G., Cioni, R., Cipolli, F., Longinelli, A., Marini, L., Ottonello, G., Zuccolini, M. V., 2002. Irreversible water-rock mass transfer accompanying the generation of the neutral, Mg-HCO₃ and high-pH, Ca-OH spring waters of the Genova province, Italy. Applied Geochemistry 17, 455-474.
- Charlou, J. L., Fouquet, Y., Bougalt, H., Donval, J. P., Etoubleau, J., Jean-Baptiste, P., Dapoigny, A., Apriou, P., Rona, P. A., 1998. Intense CH₄ plumes generated by serpentinization of ultramafic rocks at the intersection of the 15°20'N fracture zone and the Mid-Atlantic Ridge. Geochimica et Cosmochimica Acta 62, 2323-2333.
- Chavagnac, V., Ceuleneer, G., Monnin, C., Lansac, B., Hoareau, G., Boulart, C., 2013a. Mineralogical assemblages forming at hyperalkaline warm springs hosted on ultramafic rocks: A case study of Oman and Ligurian ophiolites. Geochemistry Geophysics Geosystems 14, 2474-2495.
- Chavagnac, V., Monnin, C., Ceuleneer, G., Boulart, C., Hoareau, G., 2013b. Characterization of hyperalkaline fluids produced by low-temperature serpentinization of mantle peridotites in the Oman and Ligurian ophiolites. Geochemistry Geophysics Geosystems 14, 2496-2522.
- Cheng, H., Edwards, R. L., Hoff, J., Gallup, C. D., Richards, D. A., and Asmerom, Y., 2000. The half-lives of uranium-234 and thorium-230. Chemical Geology 169, 17-33.

- Cipolli, F., Gambardella, B., Marini, L., Otonello, G., Zuccolini, M. V., 2004. Geochemistry of high-pH waters from serpentinites of the Gruppo di Voltri (Genova, Italy) and reaction path modeling of CO₂ sequestration in serpentine aquifers. *Applied Geochemistry* 19, 787-802.
- Clark, I. D., Fontes, J.-C., 1990. Paleoclimate reconstruction in northern Oman based on carbonate from hyperalkaline groundwaters. *Quaternary Research* 33, 320-336.
- Clark, I. D., Fontes, J.-C., Fritz, P., 1992. Stable isotope disequilibria in travertine from high pH waters: Laboratory investigations and field observations from Oman. *Geochimica et Cosmochimica Acta* 56, 2041-2050.
- Coleman R. G., 1977. *Ophiolites*. Springer-Verlag, New York.
- Coleman, R. G., 1981. Tectonic setting for ophiolite obduction in Oman. *Journal of Geophysical Research* 86, 2497-2508.
- Dequincey, O., Chabaux, F., Clauer, N., Sigmarsson, O., Liewig, N., Leprun, J.-C. 2002. Chemical mobilizations in laterites: Evidence from trace elements and ²³⁸U-²³⁴U-²³⁰Th disequilibria. *Geochimica et Cosmochimica Acta* 66, 1197-1210.
- Dewandel, B., Lachassagne, P., Bakalowicz, M., Weng, P., Al-Malki, A., 2003. Evaluation of aquifer thickness by analyzing recession hydrographs: Application to the Oman ophiolite hard-rock aquifer. *Journal of Hydrology* 274, 248-269.
- Dewandel, B., Lachassagne, P., Qatan, A., 2004. Spatial measurements of stream baseflow, a relevant method for aquifer characterization and permeability evaluation: Application to a hard rock aquifer, Oman ophiolite. *Hydrological Processes* 18, 3391-3400.
- Dewandel, B., Lachassagne, P., Boudier, F., Al-Hattali, S., Ladouche, B., Pinault, J.-L., Al-Suleimani, Z., 2005. A conceptual and hydrogeological model of ophiolite hard-rock aquifers in Oman based on a multiscale and multidisciplinary approach. *Hydrogeology Journal* 13, 708-726.
- Dooley, J. R., Tatsumoto, M., Rosholt, J. N., 1964. Radioactive disequilibrium studies of roll features, Shirley Basin, Wyoming. *Economic Geology* 59, 586-595.
- Dooley, J. R., Granger, H. C., Rosholt, J. N., 1965. Uranium-234 fractionation in sandstone-type uranium deposits of the Ambrosia Lake District, New Mexico. *Economic Geology* 61, 1362-1382.
- Dossetto, A., Turner, S.P., Chappell, J. 2008. The evolution of weathering profiles through time: New insights from uranium-series isotopes. *Earth and Planetary Science Letters* 274, 359-371.
- Edwards, L. R., Chen, J. H., Wasserburg, G. J., 1986. ²³⁸U-²³⁴U-²³⁰Th-²³²Th systematics and the precise measurement of time over the past 500,000 years. *Earth and Planetary Science Letters* 81, 175-192.

- Edwards, L. R., Gallup, C., Cheng, H., 2003. Uranium-series dating of marine and lacustrine carbonates. In: Bourdon, B., Henderson, G. M., Lundstrom, C. C., Turner, S. P. (Ed.), *Uranium-Series Geochemistry. Reviews in Mineralogy and Geochemistry* 52, 363-406.
- Ferry, J. M., 1995. Fluid-flow during contact-metamorphism of ophicarbonate rocks in the Bergell-Aureole, Val-Malenco, Italian Alps. *Journal of Petrology* 36, 1039-1053.
- Fleitmann, D., Burns, S. J., Neff, U., Mangini, A., Matter, A., 2003. Changing moisture sources over the last 350,000 years in northern Oman from fluid inclusion evidence in speleothems. *Quaternary Research* 60, 223-232.
- Fontes, J.-C., Andrews, J. N., Causse, C., Gibert, E., 1992. A comparison of radiocarbon and U/Th ages on continental carbonates. *Radiocarbon* 34, 602-610.
- Früh-Green, G. L., Kelley, D. S., Bernasconi, S. M., Karson, J. A., Ludwig, K. A., Butterfield, D. A., Boschi, C., Proskurowski, G. 2003. 30,000 years of hydrothermal activity at the Lost City vent field. *Science* 301, 495-498.
- Gascoyne, M., 1992. Geochemistry of the actinides and their daughters. In: Ivanovich, M., Harmon, R. S. (Eds.), *Uranium-series Disequilibria*. Oxford, Clarendon Press, pp. 34-61.
- Ghaleb, B., Hillaire-Marcel, C., Causse, C., Gariépy, C., Vallières, S., 1990. Fractionation and recycling of U and Th isotopes in a semi-arid endoreic depression of central Syria. *Geochimica et Cosmochimica Acta* 54, 1025-1035.
- Glennie, K. W., Bouef, M. G. A., Hughes-Clarke, M. W., Moody-Stuart, M., Pilaar, W. F. H., Reinhart, B. M., 1973. Late Cretaceous nappes in the Oman mountains and their geological evolution. *American Association of Petroleum Geologists Bulletin* 57, 5-27.
- Glennie, K. W., Bouef, M. G. A., Hughes-Clarke, M. W., Moody-Stuart, M., Pilaar, W. F. H., Reinhart, B. M. 1974. *Geology of the Oman Mountains. Part I: Text. Part II: Tables and Illustrations. Part III: Enclosures. Verhandelingen van het Koninklijk Nederlands Geologisch Mijnbouwkundig Genootschap* 31, pp. 423.
- Govindaraju, K., 1994. Compilation of working values and sample description for 383 geostandards. *Geostandards Newsletter* 18 Special Issue, pp. 158.
- Haase-Schramm, A., Goldstein, S. L., Stein, M., 2004. U-Th dating of Lake Lisan (late Pleistocene Dead Sea) aragonite and implications for glacial East Mediterranean climate change. *Geochimica et Cosmochimica Acta* 68, 985-1005.
- Haggerty, J. A., 1991. Evidence from fluid seeps atop submarine seamounts in the Mariana forearc: clues for emplacement of the seamounts and their relationship to forearc tectonics. *Marine Geology* 102, 293-309.

955 Hanhøj, K., Kelemen, P. B., Hassler, D., Godard, M., 2010. Composition and genesis of depleted
956 mantle peridotites from the Wadi Tayin massif, Oman Ophiolite; Major and trace element
957 geochemistry, and Os isotope and PGE systematics. *Journal of Petrology* 51, 201-227.
958

959 Hansen, R.O. and Stout, P.R. 1968. Isotopic distributions of uranium and thorium in soils. *Soil*
960 *Science* 105, 44-50.

961 Hansen, R.O. 1970. Radioactivity of a California terrace soil. *Soil Science* 110, 31-36.
962

963 Harmon, S. R., Ford, D. C., Schwarcz, H. P., 1977. Interglacial chronology of the Rocky and
964 Mackenzie Mountains based upon ^{230}Th - ^{234}U dating of calcite speleothems. *Canadian Journal of*
965 *Earth Sciences* 14, 2543-2552.
966

967 Harrison, A. L., Power, I. M., Dipple G. M., 2013. Accelerated carbonation of brucite in mine
968 tailings for carbon sequestration. *Environmental Science and Technology* 47, 126-134.
969

970 Hellstrom, J., Pickering, R., 2015. Recent advances and future prospects of the U-Th and U-Pb
971 chronometers applicable to archaeology. *Journal of Archaeological Science* 56: 32-40.
972

973 Hernandez Goldstein, E., Stockli, D., Ketcham, R., Seman, S., 2014. (U-Th)/He dating of
974 magnetites in serpentinites (Abstract). Goldschmidt Geochemistry Conference, Sacramento,
975 California.
976

977 Jaffey A. H., Flynn K. F., Glendenin L. E., Bentley W. C., Essling, A. M., 1971. Precision
978 measurement of half-lives and specific activities of ^{235}U and ^{238}U . *Physical Review C* 4, 1889-
979 1906.

980 Kaufman, A., 1993. An evaluation of several methods for determining $^{230}\text{Th}/\text{U}$ ages in impure
981 carbonates. *Geochimica et Cosmochimica Acta* 57, 2303-2317.
982

983 Kelemen, P. B., Matter, J. M., 2008. *In situ* carbonation of peridotite for CO_2 storage.
984 *Proceedings of the National Academy of Sciences* 105, 17295-17300.
985

986 Kelemen, P. B., Matter, J. M., Streit, E. E., Rudge, J. F., Curry, W. B., Blusztajn, J., 2011. Rates
987 and mechanisms of mineral carbonation in peridotite: Natural processes and recipes for
988 enhanced, *in situ* CO_2 capture and storage. *Annual Review of Earth and Planetary Sciences* 39,
989 545-576.
990

991 Kelley, D. S., Karson, J. A., Blackman, D. K., Früh-Green, G. L., Butterfield, D. A., Lilley, M.
992 D., Olson, E. J., Schrenk, M. O., Roe, K. K., Lebon, G.T., Rivizzigo, P., 2001. An off-axis
993 hydrothermal vent field near the Mid-Atlantic Ridge at 30 degrees N. *Nature* 412, 145-149.
994

995 Kelley, D. S., Karson, J. A., Früh-Green, G. L., Yoerger, D. R., Shank, T. M., Butterfield, D. A.,
996 Hayes, J. M., Schrenk, M. O., Olson, E. J., Proskurowski, G., Jakuba, M., Bradley, A., Larson,
997 B., Ludwig, K., Glickson, D., Buckman, K., Bradley, A. S., Brazelton, W. J., Roe, K., Elend, M.
998 J., Delacour, A., Bernasconi, S. M., Lilley, M. D., Baross, J. A., Summons, R. E., Sylvia, S. P.,

2005. A serpentinite-hosted ecosystem: The Lost City hydrothermal field. *Science* 307, 1428-1434.
- Krishnaswami, S., Williams, G.A., Graustein, W.C., and Turekian, K.K. 2004. The effect of weathering regime on uranium decay series and osmium in two soil profiles. *Geochemical Journal* 38, 651-660.
- Ku, T.-L., Liang, Z.-C., 1984. The dating of impure carbonates with decay-series isotopes. *Nuclear Instruments and Methods in Physics Research* 223, 563-571.
- Lackner, K. S., Wendt, C. H., Butt, D. P., Joyce, E. L., Sharp, D. H., 1995. Carbon dioxide disposal in carbonate minerals. *Energy* 20, 1153-1170.
- Lackner, K. S., 2002. Carbonate chemistry for sequestering fossil carbon. *Annual Review of Energy and the Environment* 27, 193-232.
- Langmuir, D., Herman, J. S., 1980. The mobility of thorium in natural waters at low temperatures. *Geochimica et Cosmochimica Acta* 44, 1753-1766.
- Le Roux, L. J., Glendenin, L. E., 1963. Half-life of ^{232}Th , *Proc. Natl. Meet. Nucl. Energy*, 83-94.
- Lippard, S. J., Shelton, A. W., Gass, I. G., 1986 *The Ophiolite of Northern Oman*. The Geological Society, Memoir No. 11. Blackwell Scientific Publications, London.
- Lively, R. S., Harmon, R. S., Levinson, A. A., Bland, C. J., 1979. Disequilibrium in the ^{238}U series in samples from Yeelirrie, Western Australia. *Journal of Geochemical Exploration* 12, 57-65.
- Ludwig, K. R., Titterton, D. M., 1994. Calculation of $^{230}\text{Th}/\text{U}$ isochrons, ages, and errors. *Geochimica et Cosmochimica Acta* 58: 5031-5042.
- Ludwig, K. A., Kelley, D. S., Butterfield, D. A., Nelson, B. K., Fröh-Green, G. L., 2006. Formation and evolution of carbonate chimneys at the Lost City hydrothermal field. *Geochimica et Cosmochimica Acta* 70, 3625-2645.
- Ludwig, K. A., Shen, C.-C., Kelley, D. S., Cheng, H., Edwards, L. R., 2011. U-Th systematics and ^{230}Th ages of carbonate chimneys at the Lost City hydrothermal field. *Geochimica et Cosmochimica Acta* 75, 1869-1888.
- Luo, Y., Ku, T.-L., 1991. U-series isochron dating: A generalized method employing total sample dissolution. *Geochimica et Cosmochimica Acta* 55, 555-564.
- Ma, L., Chabaux, F., Pelt, E., Blaes, E., Jin, L., and Brantley, S. 2010. Regolith production rates calculated with uranium-series isotopes at Susquehanna/Shale Hills Critical Zone Observatory. *Earth and Planetary Science Letters* 297, 211-225.

1043 Mathieu, D., Bernat, M., and Nahon, D. 1995. Short-lived U and Th isotope distribution in a
 1044 tropical laterite derived from granite (Pitinga river basin, Amazonia, Brazil): Application to
 1045 assessment of weathering rate. *Earth and Planetary Science Letters* 136, 703-714.

1046 Matter, J. M. and Kelemen, P. B., 2009. Permanent storage of carbon dioxide in geological
 1047 reservoirs by mineral carbonation. *Nature Geoscience* 2, 837-841.

1048

1049 Mervine, E. M., Humphris, S. E., Sims, K. W. W., Kelemen, P. K., Jenkins, W. J., 2014.
 1050 Carbonation rates of peridotite in the Samail Ophiolite, Sultanate of Oman, constrained through
 1051 ¹⁴C dating and stable isotopes. *Geochimica et Cosmochimica Acta* 126, 371-397.

1052

1053 Mitchell, R. H., 1986. *Kimberlites: Mineralogy, Geochemistry, and Petrology*. Plenum Press,
 1054 New York.

1055

1056 Neal, C., Stanger, G., 1983. Hydrogen generation from mantle source rocks in Oman. *Earth and*
 1057 *Planetary Science Letters* 66, 315-320.

1058

1059 Neal, C., Stanger, G., 1984. Calcium and magnesium-hydroxide precipitation from alkaline
 1060 groundwaters in Oman, and their significance to the process of serpentinization. *Mineralogical*
 1061 *Magazine* 48, 237-241.

1062

1063 Neal, C., Stanger, G., 1985. Past and present serpentinization of ultramafic rocks: An example
 1064 from the Semail ophiolite nappe of northern Oman. In: Drewer, J. I. (Ed.), *The Chemistry of*
 1065 *Weathering*. D. Reidel Publishing Company, Holland, pp. 249-275.

1066

1067 Nicolas, A., Boudier, F., Ildefonse, V., Ball, E., 2000. Accretion of Oman and United Arab
 1068 Emirates ophiolite—Discussion of a new structural map. *Marine Geophysical Researches* 21,
 1069 147-149.

1070

1071 Osmond, J. K., May, J. P., Tanner, W. F., 1970. Age of the Cape Kennedy Barrier-and-Lagoon
 1072 Complex. *Journal of Geophysical Research* 75, 469-479.

1073

1074 Paukert, A. N., Matter, J. M., Kelemen, P. B., Shock, E. L., Havig, J. R. 2012. Reaction path
 1075 modeling of enhanced *in situ* CO₂ mineralization for carbon sequestration in the peridotite of the
 1076 Samail Ophiolite, Sultanate of Oman. *Chemical Geology* 330-331, 86-100.

1077

1078 Pelt, E., Chabaux, F., Innocent, C., Navarre-Sitchler, A.K., Sak, P.B., and Brantley, S.L. 103
 1079 2008. Uranium-thorium chronometry of weathering rinds: Rock alteration rate 104 and
 1080 paleoisotopic record of weathering fluids. *Earth and Planetary Science Letters* 276, 98-105.

1081

1082 Peretti, A., Dubessy, J., Mullis, J., Frost, B. R., Tromsdorff, V., 1992. Highly reducing
 1083 conditions during alpine metamorphism of the Malenco peridotite (Sondrio, northern Italy)
 1084 indicated by mineral paragenesis and H₂ in fluid inclusions. *Contributions to Mineralogy and*
 1085 *Petrology* 112, 329-340.

- Porcelli, D., Swarzeski, P. W., 2003. The behavior of U- and Th-series nuclides in groundwater. In: Bourdon, B., Henderson, G. M., Lundstrom, C. C., Turner, S. P. (Ed.), Uranium-Series Geochemistry. Reviews in Mineralogy and Geochemistry 52, 363-406.
- Power I. M., Wilson S. A., Thom J. M., Dipple G. M., Gabites J. E. and Southam G., 2009. The hydromagnesite playas of Atlin, British Columbia, Canada: A biogeochemical model for CO₂ sequestration. Chemical Geology 260, 286-300.
- Pronost, J., Beaudoin, G., Tremblay, J., Larachi, F., Duchesne, J., Hébert, R., Constantin, M., 2011. Carbon sequestration kinetic and storage capacity of ultramafic mining waste. Environmental Science and Technology 45, 9413-9420.
- Przybylowicz, W., Schwarcz, H. P., Latham, A. G., 1991. Dirty Calcites 2. Uranium series dating of artificial calcite-detritus mixtures. Chemical Geology 86, 161-178.
- Rasbury, E. T., Cole, J. M., 2009. Directly dating geologic events: U-Pb dating of carbonates. Reviews of Geophysics 47: 1-27.
- Robinson, C. S., Rosholt, J. N., 1961. Uranium migration and geochemistry of uranium deposits in sandstone above, at, and below the water table. Part II: Relationship of uranium migration dates, geology, and chemistry of the ore deposits. Economic Geology 56, 1404-1420.
- Rosholt, J. N., 1961. Uranium migration and geochemistry of uranium deposits in sandstone at, above, and below the water table. Part I: Calculation of apparent dates of uranium migration in deposits above and at the water table. Economic Geology 56, 1392-1403.
- Rosholt, J. N., Harshman, E. N., Shields, W. R., Garner, E. L., 1964. Isotopic fractionation of uranium related to roll features in sandstone, Shirley Basin, Wyoming. Economic Geology 59, 570-585.
- Rosholt, J.N., Doe, B.R., and Tatsumoto, M. 1966. Evolution of the isotopic composition of uranium and thorium in soil profiles. Geological Society of America Bulletin 77, 987-1004.
- Rosholt, J.N., Bush, C.A., Shroba, R.R., Pierce, K.L., and Richmond, G.M. 1985. Uranium-trend dating and calibrations for Quaternary sediments. Open-File Report 85-299, U.S. Geological Survey, Lakewood, Colorado, pp. 48
- Scholz, D., Hoffmann, D., 2008. ²³⁰Th/U-dating of fossil corals and speleothems. Quaternary Science Journal 57, 52-76.
- Schwarcz, H. P., Lathan, A. G., 1989. Dirty Calcites 1. Uranium-series dating of contaminated calcite using leachates alone. Chemical Geology 80, 35-42.
- Searle, M. P., Cox, J., 1999. Tectonic setting, origin, and obduction of the Oman ophiolite. Geological Society of America Bulletin 111, 104-122.

- Searle, M. P., Cox, J., 2002. Subduction zone metamorphism during formation and emplacement of the Semail ophiolite in the Oman Mountains. *Geological Magazine* 139, 241-255.
- Seifritz W., 1990. CO₂ disposal by means of silicates. *Nature* 345, 486.
- Sims, K. W. W., Hart, S. R., Reagen, M. K., Blusztajn, J., Staudigel, H., Sohn, R. A., Layne, G. D., Ball, L. A., Andrews, J., 2008a. ²³⁸U-²³⁰Th-²²⁶Ra-²¹⁰Pb-²¹⁰Po, ²³²Th-²²⁸Ra, and ²³⁵U-²³¹Pa constraints on the ages and petrogenesis of Vailulu'u and Malumalu Lavas, Samoa. *Geochemistry, Geophysics, Geosystems* 9, 1-30.
- Sims, K. W. W., Gill, J. B., Dosseto, A., Hoffman, D. L., Lundstrom, C. C., Williams, R. W., Ball, L., Tollstrup, D., Turner, S., Prytulak, J., Glessner, J. J. G., Standish, J. J., Elliot, T., 2008b. An inter-laboratory assessment of the thorium isotopic composition of synthetic and rock reference materials. *Geostandards and Geoanalytical Research* 32, 65-91.
- Streit, E., Kelemen, P., Eiler, J., 2012 Coexisting serpentine and quartz from carbonate-bearing serpentinized peridotite in the Semail Ophiolite, Oman. *Contributions to Mineralogy and Petrology* 164, 821-837.
- Surour, A. A., Arafa, E. H., 1997. Ophicarbonates: Calichified serpentinites from Gebel Mohagara, Wadi Ghadir area, Eastern Desert, Egypt. *Journal of African Earth Sciences* 24, 315-324.
- Szabo, B. J., Rosholt, J. N., 1982. Surficial continental sediments. In: Ivanovich, M., Harmon, R. S. (Eds.), *Uranium Series Disequilibrium: Applications to Environmental Problems*. London, Claredon Press, pp. 257-260.
- Thompson, W. G., Spiegelman, M. W., Goldstein, S. L., Speed, R. C., 2003. An open system model for U-series age determinations of fossil corals. *Earth and Planetary Science Letters* 210, 365-381.
- Trommsdorff, V., Evans, B. W., 1977. Antigorite-ophicarbonates: Contact metamorphism in Valmalenco, Italy. *Contributions to Mineralogy and Petrology* 62, 301-312.
- Trommsdorff, V., Evans, B. W., Pfeifer, H. R., 1980. Ophicarbonate rocks: Metamorphic reactions and a possible origin. *Archives des Sciences Genève* 33, 361-364.
- Villemant, B., Feuillet, N., 2003. Dating open systems by the ²³⁸U-²³⁴U-²³⁰Th method: application to Quaternary reef terraces. *Earth and Planetary Science Letters* 210, 105-118.
- Wilson, S. A., Raudsepp, M., Dipple, G. M., 2006. Verifying and quantifying carbon fixation in minerals from serpentine-rich mine tailings using the Rietveld method with X-ray powder diffraction data. *American Mineralogist* 91, 1331-1341.
- Wilson, S. A., Raudsepp, M., Dipple, G. M., 2009a. Quantifying carbon fixation in trace minerals from processed kimberlite: A comparative study of quantitative methods using X-ray

powder diffraction data with applications to the Diavik Diamond Mine, Northwest Territories, Canada. *Applied Geochemistry* 24, 95-112.

Wilson, S. A., Dipple, G. M., Power, I. M., Thom, J. M., Anderson, R. G., Raudsepp, M., Gabites, J. E., Southam, G., 2009b. Carbon dioxide fixation within mine wastes of ultramafic hosted ore deposits: Examples from the Clinton Creek and Cassiar chrysotile deposits, Canada. *Economic Geology* 104, 95-112.

Woodhead, J., Pickering, R., 2012. Beyond 500 ka: Progress and prospects in the U-Pb chronology of speleothems, and their application to studies in paleoclimate, human evolution, biodiversity, and tectonics. *Chemical Geology* 322-323: 290-299.

List of tables

Table 1: Sample Locations and Descriptions

Table 2: U and Th Concentrations. U and Th concentrations were measured by isotope dilution using a Thermo Fisher Element 2 high resolution sector-field ICP-MS at WHOI. Mass fractionation was corrected by sample-standard bracketing with NBL-112A, using a linear interpolation of $^{235}\text{U}/^{238}\text{U}$ measurements of the standard. Errors include uncertainties in spike calibrations, in run measurement error, and uncertainty in standards against which instrument measurements were calibrated.

Table 3: U and Th Isotopic Values Isotopic compositions were measured with a Thermo Fisher Neptune MC-ICP-MS at WHOI. ^{232}Th and ^{238}U were measured on a Faraday cup and ^{230}Th on an axial, discrete dynode ion counter with a repelling potential quadrupole (RPQ). Abundance sensitivity over 2 AMU at 85% transmission was ~50 ppb. An exponential correction was used to correct for the tailing of ^{232}Th on ^{230}Th . Use of UCSC Th 'A' for Th-bracketing results in inter-day reproducibility of 0.4-1.5% (2σ). $^{236}\text{U}/^{238}\text{U}$ ratios of U010 were measured to correct for mass bias drift and to calibrate SEM yield. Standard-sample-standard bracketing was used for uranium analyses.

() denotes activity. Activity ratios calculated using: $\lambda_{230} = 9.158 \times 10^{-6} \text{ yr}^{-1}$ (Cheng et al., 2000); $\lambda_{232} = 4.948 \times 10^{-11} \text{ yr}^{-1}$ (Le Roux and Glendenin, 1963); $\lambda_{234} = 2.8263 \times 10^{-6} \text{ yr}^{-1}$ (Cheng et al., 2000); and, $\lambda_{238} = 1.551 \times 10^{-10} \text{ yr}^{-1}$; (Jaffey et al., 1971).

Errors for activity ratios include uncertainties in spike calibrations for ^{232}Th and ^{238}U concentrations, in run measurement error for $^{230}\text{Th}/^{232}\text{Th}$ and uncertainties in the standards against which instrument measurements were calibrated. Errors do not include uncertainties in λ_{238} (0.07%), λ_{232} (0.5%), λ_{230} (0.3%) or λ_{234} (0.1%).

List of figures

Figure 1: The eight field locations (blue circles) where carbonate and peridotite samples were collected in the Samail Ophiolite during 2009 and 2010 field seasons. Map made using ArcGIS, Version 10.1.

Figure 2: Schematic illustrating the behavior of CO₂, U, and Th in Type I and Type II waters associated with alteration of partially-serpentinized peridotite. Figure modified from Neal and Stanger (1985).

Figure 3: Representative images of carbonates and altered peridotites in the Samail Ophiolite. A: Recently-formed travertine precipitate in a hyperalkaline streambed at Qafeefah Travertine. B: A turquoise hyperalkaline pool and recently-formed travertine precipitate at Wadi Sudari Travertine. Note the hammer for scale. C: Small hills of altered peridotite (background) adjacent to travertine terraces at Misht Travertine. D: Calcite-rich travertine veins exposed underneath a ~1 to 2 m thick travertine terrace deposit at Wadi Sudari Travertine. The veins can be seen in the overhang, next to the person. E: A white magnesite vein with botryoidal texture exposed on the natural peridotite weathering surface near Fanja Roadcut. Note the hammer for scale. F: Dolomite-serpentine veins exposed on the surface of Fanja Roadcut. Note the marker for scale. G: Very highly altered peridotite exposed underneath a travertine terrace at Misht Travertine. H: Highly altered peridotite exposed at Misht Travertine. Note the hammer for scale.

Figure 4: Plot of U and Th concentrations for Samail travertines (orange circles for travertine terraces, pink circles for recently-formed travertine precipitates, orange triangles for travertine pseudospeleothems, and an orange square for the travertine vein), Mg-rich carbonate veins (blue circles for Mg-rich carbonate outcrop veins and blue squares for Mg-rich carbonate roadcut veins), serpentine mineral separates from carbonate-serpentine veins (purple circles), and wadi sediments (brown triangles). Note that both axes have a log scale. Shown for reference are U and Th concentrations for harzburgites (green circles) and dunites (green squares) from Wadi Tayin, a massif in the southern part of the Samail Ophiolite (data from Hanghøj et al., 2010); for carbonates formed at Lost City, a hydrothermal site located in peridotite ~15 km off-axis of the Mid-Atlantic Ridge (data from Ludwig et al., 2011); and for speleothems from Hoti Cave, which is located in Cretaceous age Natih limestone in northern Oman (data from Fleitmann et al., 2003).

Figure 5: Plot of (²³⁰Th/²³²Th) versus (²³⁸U/²³²Th) for Samail travertines, Mg-rich carbonate veins, highly altered peridotites, and wadi sediments using the same symbol scheme as Figure 4.

Figure 6: Plot of (²³⁴U/²³⁸U) versus U concentration for Samail travertines, Mg-rich carbonate veins, and wadi sediments using the same symbol scheme as Figure 4. The dashed line indicates equilibrium for (²³⁴U/²³⁸U).

Figure 7: Plot of (²³⁰Th/²³²Th) versus Th concentration for Samail travertines, Mg-rich carbonate veins, and wadi sediments using the same symbol scheme as Figure 4.

Figure 8: Sensitivity analysis figure showing the effect of the selected detritus (²³⁰Th/²³²Th) correction value on the corrected ²³⁰Th age. This figure is for sample OM09-106C-MS-A, a travertine terrace sample from Wadi Uqaybah. The blue line indicates the corrected ²³⁰Th age calculated from the detritus value on the x-axis. Note that the uncorrected ²³⁰Th age plots at x = 0. For comparison, the calibrated ¹⁴C age (from Mervine et al., 2014) is plotted as the red line. The gray box indicates the 2σ range of (²³⁰Th/²³²Th) values for aluminosilicate detritus (a literature compilation of ~300 values, see Supplementary Information). The vertical lines

indicate possible values to use for the detritus ($^{230}\text{Th}/^{232}\text{Th}$) correction value: the average of the aluminosilicate detritus literature compilation (yellow line), equilibrium Bulk Silicate Earth (pink line), wadi sediments (green lines), and recently-formed travertine precipitates (black lines). The choice of the ($^{230}\text{Th}/^{232}\text{Th}$) correction value has a very significant impact on the determined ^{230}Th age, which ranges from ~50,000 years to zero depending on the correction value selected. However, a ($^{230}\text{Th}/^{232}\text{Th}$) correction value of approximately 1.7 brings the corrected ^{230}Th age into concordance with the ^{14}C age.

Figure 9: Sensitivity analysis figure showing the effect of the selected detritus ($^{230}\text{Th}/^{232}\text{Th}$) correction value on the corrected ^{230}Th age. This figure is for sample OM10-13C-MG, an Mg-rich carbonate outcrop vein from Fanja Roadcut. The blue line indicates the corrected ^{230}Th age calculated from the detritus value on the x-axis. Note that the uncorrected ^{230}Th age plots at x = 0. For comparison, the calibrated ^{14}C age (from Mervine et al., 2014) is plotted as the red line. The gray box indicates the 2σ range of ($^{230}\text{Th}/^{232}\text{Th}$) values for aluminosilicate detritus (a literature compilation of ~300 values, see Supplementary Information). The vertical lines indicate possible values to use for the detritus ($^{230}\text{Th}/^{232}\text{Th}$) correction value: the average of the aluminosilicate detritus literature compilation (yellow line), equilibrium Bulk Silicate Earth (pink line), wadi sediments (green lines), and recently-formed travertine precipitates (black lines). The choice of the ($^{230}\text{Th}/^{232}\text{Th}$) correction value has a very significant impact on the determined ^{230}Th age, which ranges from ~30,000 years to zero depending on the correction value selected. Since the uncorrected ^{230}Th age is younger than the ^{14}C age, no ($^{230}\text{Th}/^{232}\text{Th}$) correction value can bring the two ages into concordance.

Figure 10: Sensitivity analysis figure showing the effect of the selected detritus ($^{230}\text{Th}/^{232}\text{Th}$) correction value on the corrected ^{230}Th age. This figure is for sample OM09-57C-MS, an Mg-rich carbonate roadcut vein from Fanja Roadcut. The blue line indicates the corrected ^{230}Th age calculated from the detritus value on the x-axis. Note that the uncorrected ^{230}Th age plots at x = 0. For comparison, the calibrated ^{14}C age (from Mervine et al., 2014) is plotted as the red line. The gray box indicates the 2σ range of ($^{230}\text{Th}/^{232}\text{Th}$) values for aluminosilicate detritus (a literature compilation of ~300 values, see Supplementary Information). The vertical lines indicate possible values to use for the detritus ($^{230}\text{Th}/^{232}\text{Th}$) correction value: average of aluminosilicate detritus literature compilation (yellow line), equilibrium Bulk Silicate Earth (pink line), wadi sediments (green lines), and recently-formed travertine precipitates (black lines). The choice of the ($^{230}\text{Th}/^{232}\text{Th}$) correction value has a minimal impact on the determined ^{230}Th age, which remains greater than 350,000 years. This is consistent with the ^{14}C dead (>50,000 yr BP) age limit of this sample.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10

Table 1

Sample Name:	Location:	UTM-Easting	UTM-Northing	Description:	Major Minerals (XRD):	Trace Minerals (XRD):
<i>Travertines:</i>						
OM09-76C-MS-A (Top)	Wadi Sudari Travertine	0443115	2650257	Layered pseudospeleothem (drop)	calcite	none
OM09-76C-MS-C (Bottom)	Wadi Sudari Travertine	0443115	2650257	Layered pseudospeleothem (drop)	calcite	unidentified clay
OM09-8COPS-MS	Wadi Sudari Travertine	0443118	2650078	Recently-formed travertine precipitate	calcite	aragonite, hydromagnesite?, unidentified clay
OM09-84C-MS	Wadi Sudari Travertine	0443082	2650304	Travertine vein	calcite, brucite	hydromagnesite?, unidentified clay
OM09-106C-MS-A (Top)	Wadi Uqaybah Travertine	0426245	2633924	Travertine terrace	calcite	none
OM09-106C-MS-B (Bottom)	Wadi Uqaybah Travertine	0426245	2633924	Travertine terrace	calcite	unidentified clay
OM09-107C-MS	Wadi Uqaybah Travertine	0426309	2633950	Travertine terrace	calcite	unidentified clay
OM09-109C-MS	Wadi Uqaybah Travertine	0426208	2633925	Travertine terrace	calcite	none
OM09-10COPS-MS	Wadi Uqaybah Travertine	0426183	2633965	Recently-formed travertine precipitate	calcite	quartz, unidentified clay
<i>Mg-Rich Carbonate Outcrop Veins:</i>						
OM09-47C-MS	Fanja Roadcut	0609304	2597565	Carbonate outcrop vein	magnesite	unidentified clay
OM10-13C-MG	Fanja Roadcut	0609451	2597416	Carbonate outcrop vein	magnesite	dolomite, calcite
OM10-53C-MG	Al-Wuqbah Roadcut	0440574	2643360	Carbonate outcrop vein	magnesite	none
OM09-91C-MS	Wadi Sudari Campsite	0446151	2647471	Carbonate outcrop vein	magnesite	calcite
OM10-82C-MG	Wadi Sudari Campsite	0445905	2647602	Carbonate outcrop vein	magnesite	dolomite, calcite, unidentified clay
OM10-84C-MG	Wadi Sudari Campsite	0445991	2647684	Carbonate outcrop vein	brucite, calcite	magnesite, dolomite, unidentified clay
<i>Mg-Rich Carbonate Roadcut Veins:</i>						
OM09-35C-MS	Qafeefah Roadcut	0647852	2537682	Carbonate/serpentine	magnesite	chrysotile, unidentified clay

				roadcut vein		
OM09-36C-MS	Qafeefah Roadcut	0647818	2537657	Carbonate roadcut vein	magnesite	calcite, dolomite
OM09-55C-MS	Fanja Roadcut	0609351	2597507	Carbonate/serpentine roadcut vein	dolomite	aragonite?, unidentified clay
OM09-57C-MS	Fanja Roadcut	0609356	2597501	Carbonate/serpentine roadcut vein	dolomite	aragonite?, calcite, unidentified clay
OM09-58C-MS	Fanja Roadcut	0609370	2597506	Carbonate roadcut vein	dolomite, periclase?	aragonite?, unidentified clay
OM09-63C-MS	Fanja Roadcut	0609470	2597481	Carbonate roadcut vein	magnesite	unidentified clay
OM10-52C-MG	Al-Wuqbah Roadcut	0440396	2643047	Carbonate roadcut vein	magnesite	halite?, unidenitified clay
<i>Serpentines:</i>						
OM10-27C-MG-SERP	Qafeefah Roadcut	0647791	2537644	Carbonate/serpentine roadcut vein	lizardite	none
OM10-54C-MG-SERP	Al-Wuqbah Roadcut	0440832	2643931	Carbonate/serpentine roadcut vein	lizardite	none
<i>Sediments:</i>						
OM09-8S-MS	Fanja Roadcut	0609493	2597536	Sediment collected from wadi floor	lizardite	quartz, dolomite, magnesite, calcite
OM09-12S-MS	Wadi Uqaybah Travertine	0426173	2633970	Sediment collected from wadi floor	quartz, calcite	chrysotile, dolomite, enstatite, aragonite?
<i>Carbonate Standard:</i>						
JCp-1	Ryukyu Islands, Japan			<i>Porites</i> coral	--	--

Table 2

Sample Name:	U (ppm):	2σ (%):	Th (ppm):	2σ (%):	Th/U:	2σ (%):
<i>Travertines:</i>						
OM09-76C-MS-A (Top)	0.0447	3.14	0.0673	1.45	1.5070	3.46
OM09-76C-MS-C (Bottom)	0.0195	1.81	0.0472	1.83	2.4239	2.58
OM09-8COPS-MS	0.0596	1.95	0.1561	1.30	2.6202	2.35
OM09-84C-MS	0.0161	1.42	0.0041	1.39	0.2532	1.99
OM09-106C-MS-A (Top)	0.8275	5.42	0.2917	1.96	0.3525	5.77
OM09-106C-MS-B (Bottom)	0.3434	1.57	0.1507	2.20	0.4389	2.70
	0.3456	1.88	0.1507	1.25	0.4361	2.26
	0.3445	1.73	0.1507	1.73	0.4375	2.48
OM09-107C-MS	0.6016	1.52	0.4250	1.25	0.7065	1.96
OM09-109C-MS	0.0687	1.61	0.1072	1.98	1.5607	2.55
OM09-10COPS-MS	0.2693	3.06	0.3455	4.20	1.2829	5.19
	0.1732	1.56	0.3560	1.52	2.0551	2.18
	0.2213	2.31	0.3507	2.86	1.5852	1.24
<i>Mg-Rich Carbonate Outcrop Veins:</i>						
OM09-47C-MS	0.0030	2.62	0.0007	2.10	0.2317	3.36
	0.0007	7.05	0.0003	3.44	0.3547	7.85
	0.0019	4.84	0.0005	2.77	0.2554	5.60
OM10-13C-MG	0.0010	1.65	0.0002	4.49	0.1683	4.78
OM10-53C-MG	0.0032	2.72	0.0006	2.48	0.1819	3.69
OM09-91C-MS	0.0114	8.72	0.0030	12.68	0.2626	15.39
OM10-82C-MG	0.0009	3.96	0.0003	2.45	0.2903	4.66
OM10-84C-MG	0.0010	1.74	0.0021	2.66	1.9955	3.18
<i>Mg-Rich Carbonate Roadcut Veins:</i>						
OM09-35C-MS	0.0001	1.43	0.0003	1.31	3.0990	1.93
OM09-36C-MS	0.0001	2.11	0.0003	4.87	2.9332	5.30
OM09-55C-MS	0.0008	1.86	0.0004	2.55	0.4777	3.15
OM09-57C-MS	0.0053	1.46	0.0005	1.45	0.0978	2.06
OM09-58C-MS	0.0148	1.66	0.0002	2.65	0.0111	3.12
OM09-63C-MS	0.00003	4.47	0.00006	5.21	2.1948	6.86
	0.00004	2.72	0.00008	1.54	1.8916	3.13
	0.00003	3.59	0.00007	3.37	2.0121	4.99
OM10-52C-MG	0.0003	2.08	0.0017	6.37	5.6094	6.70
<i>Serpentines:</i>						
OM10-27C-MG-SERP	0.00031	2.66	0.0003	2.06	1.0655	3.37
OM10-54C-MG-SERP	0.00045	2.11	0.0009	1.53	2.0371	2.61
<i>Sediments:</i>						
OM09-8S-MS	0.1210	2.11	0.0686	2.35	0.5669	3.16
OM09-12S-MS	0.9055	3.91	1.3124	6.66	1.4494	7.72
<i>Carbonate Standard:</i>						
JCp-1	2.7300	2.47	0.0502	3.36	0.0184	4.17
	2.7503	1.51	0.0505	2.29	0.0184	2.74

	<i>2.7616</i>	<i>1.91</i>	<i>0.0510</i>	<i>2.17</i>	<i>0.0185</i>	<i>2.38</i>
	<i>2.8075</i>	<i>2.55</i>	<i>0.0572</i>	<i>2.83</i>	<i>0.0204</i>	<i>3.81</i>
	2.7623	2.11	0.0522	2.66	0.0189	3.28

Table 3

Sample Name:	²³⁴ U/ ²³⁸ U (x 10 ⁻⁶):	2σ (%)	²³⁰ Th/ ²³² Th (x 10 ⁻⁶):	2σ (%)	(²³⁴ U/ ²³⁸ U):	2σ (%)	(²³⁰ Th/ ²³² Th):	2σ (%)	(²³⁸ U/ ²³² Th):	2σ (%)	(²³⁰ Th/ ²³⁸ U):	2σ (%)	Uncorrected ²³⁰ Th Age:
Travertines:													
OM09-76C-MS-A	64.815	0.139	7.503	0.226	1.181	0.139	1.389	0.226	2.013	3.460	0.690	3.470	93,000
OM09-76C-MS-C	64.296	0.205	6.344	0.335	1.172	0.205	1.174	0.335	1.252	2.579	0.938	2.600	163,000
OM09-8COPS-MS	62.190	0.159	7.434	0.136	1.133	0.159	1.376	0.136	1.158	2.347	1.188	2.351	--
OM09-84C-MS	45.670	0.234	32.149	0.462	0.832	0.234	5.950	0.462	11.980	1.989	0.497	2.042	104,000
OM09-106C-MS-A	63.466	0.048	20.330	0.098	1.156	0.048	3.763	0.098	8.606	5.765	0.437	5.766	51,000
OM09-106C-MS-B	63.427	0.076	14.707	0.171	1.156	0.076	2.722	0.171	6.912	2.705	0.394	2.710	
	63.390	0.079	14.693	0.090	1.155	0.079	2.719	0.090	6.956	2.258	0.391	2.260	
	63.409	0.077	14.700	0.131	1.155	0.077	2.721	0.131	6.934	2.481	0.392	2.485	45,000
OM09-107C-MS	68.860	0.064	23.603	0.082	1.255	0.064	4.369	0.082	4.294	1.963	1.017	1.964	164,000
OM09-109C-MS	61.355	0.118	7.694	0.218	1.118	0.118	1.424	0.218	1.944	2.552	0.733	2.561	113,000
OM09-10COPS-MS	61.451	0.081	7.649	0.153	1.120	0.081	1.416	0.153	1.476	2.178	0.959	2.183	--
Mg-Rich Outcrop Veins:													
OM09-47C-MS	67.905	0.295	8.666	0.456	1.237	0.295	1.604	0.456	8.553	7.850	0.188	7.863	18,000
OM10-13C-MG	67.111	0.187	31.111	0.800	1.223	0.187	5.758	0.800	18.024	4.782	0.319	4.848	33,000
OM10-53C-MG	66.747	0.198	8.312	0.629	1.216	0.198	1.538	0.629	16.680	3.687	0.092	3.740	9,000
OM09-91C-MS	66.866	0.129	12.481	0.356	1.218	0.129	2.310	0.356	11.551	15.393	0.200	15.397	19,000
OM10-82C-MG	67.717	0.262	--	--	1.234	0.262	--	--	10.450	4.656	--	--	--
OM10-84C-MG	65.139	0.268	6.723	0.374	1.187	0.268	1.244	0.374	1.520	3.180	0.818	3.202	122,000
Mg-Rich Roadcut Veins:													
OM09-35C-MS	--	--	5.650	1.435	--	--	1.046	1.435	0.979	1.934	1.068	2.408	--
OM09-55C-MS	6.630	0.440	32.800	0.785	1.044	0.440	6.071	0.785	6.351	3.153	0.956	3.250	256,000
OM09-57C-MS	31.344	0.109	164.870	0.151	1.010	0.109	30.515	0.151	31.019	2.055	0.984	2.061	>375,000
OM09-58C-MS	54.751	0.085	--	--	0.998	0.085	--	--	272.445	3.121	--	--	--
OM10-52C-MG	61.609	0.377	4.506	0.428	1.123	0.377	0.834	0.428	0.541	6.701	1.542	6.715	Not Determinable
Sediments:													
OM09-8S-MS	56.637	0.067	28.178	0.075	1.032	0.067	5.215	0.075	5.352	3.162	0.975	3.163	--
OM09-12S-MS	60.501	0.057	8.621	0.063	1.102	0.057	1.596	0.063	2.093	7.724	0.762	7.724	--

<i>Carbonate Standard:</i>													
JCp-1	63.002	0.097	6.965	0.849	1.148	0.097	1.289	0.849	164.831	4.173	0.008	4.259	
	62.943	0.080	7.016	0.430	1.148	0.080	1.299	0.430	165.235	2.740	0.008	2.773	
	63.017	0.075	6.898	0.368	1.148	0.075	1.277	0.368	164.308	2.380	0.008	2.408	
	62.899	0.048	7.107	0.420	1.146	0.048	1.315	0.420	148.854	3.809	0.009	3.832	
	62.965	0.075	6.997	0.517	1.148	0.075	1.295	0.517	160.807	3.275	0.008	3.318	