Ultra-depleted melts in olivine-hosted melt inclusions from the Ontong Java Plateau.

M.G. Jackson¹, R.A. Cabral², E.F. Rose-Koga³, K.T. Koga³, A. Price¹, E.H. Hauri⁴, P. Michael⁵

¹ *University of California Santa Barbara, Department of Earth Science, Santa Barbara, CA, 93109-9630, USA (jackson@geol.ucsb.edu)*

² *Boston University, Department of Earth & Environment, Boston, MA, 02215, USA (racabral@bu.edu)*

³ *Université Blaise Pascal - CNRS - IRD, Laboratoire Magmas et Volcans, OPGC, Clermont-Ferrand, France (e.koga@opgc.univ-bpclermont.fr, k.koga@opgc.univ-bpclermont.fr)*

⁴ *Department of Terrestrial Magnetism, Carnegie Institution of Washington, Washington, DC 20015, USA (ehauri@ciw.edu)*

⁵ *Department of Geosciences, University of Tulsa, Tulsa, OK, 74104, USA (pjm@utulsa.edu)*

Abstract

The Ontong Java Plateau (OJP) represents the largest large igneous province (LIP) preserved in the geologic record. The most voluminous volcanic types of the OJP—the Kroenke and Kwaimbaita groups, which dominate the accessible portions of the plateau—have relatively flat primitive mantle normalized rare earth element (REE) patterns. With the exception of relatively small volumes of late-stage melts—referred to as the Singgalo group—that are characterized by slightly enriched REE patterns relative to a chondritic pattern, the volcanic groups that dominated the eruptive history of the OJP exhibit remarkably homogeneous, flat REE patterns. Here we isolate, for the first time, olivine-hosted melt inclusions from OJP. We show that the melt inclusions have two clear populations defined by distinct trace element

characteristics. The first population has relatively flat trace element patterns that are similar to those observed in whole rock lavas from the most voluminous volcanic groups (Kroenke and Kwaimbaita) recorded in the OJP. In contrast, a second group of melt inclusions, referred to as UDM (ultra-depleted melt) inclusions, exhibit (light-REE) LREE-depleted patterns relative to a chondritic pattern; these trace element patterns are far more depleted than any previously reported lava from OJP. The UDM have unique trace element signatures that preclude an origin by re-melting the depleted mantle source left over after melt extraction during construction of the OJP. We interpret the new UDM compositions to be the result of melting a previously unrecognized (in lavas) ultra-depleted component hosted in the OJP mantle source.

1. Introduction

Large igneous provinces (LIPs) represent the largest outpourings of lava in the geologic record (Coffin and Eldholm, 1994). Ontong Java Plateau (OJP) is the largest LIP preserved on Earth and has an estimated volume of 44.4 million km³ (Coffin and Eldholm, 1994; Coffin and Eldholm, 2000), a crustal thickness of approximately 30 km (Richardson et al., 2000; Miura et al., 2004), and a surface area that is roughly equivalent to Western Europe (Fitton and Godard, 2004). The bulk of the plateau was extruded at ~120 Ma (Mahoney et al., 1993, 2001; Tejada et al., 1996, 2002, 2004, 2013; Parkinson et al., 2001; Chambers et al., 2004) on top of young (15-30 Ma) oceanic crust (Larson,

1997), but the Pacific lithospheric upon which the OJP was erupted may have been as old as 40 Ma at the time of eruption (Ishikawa et al., 2011). A later stage of volumetrically-less extensive volcanism may have occurred at 90 Ma (Mahoney et al., 1993; Tejada et al., 1996, 2002), but the occurrence of the 90 Ma episode is still debated (Chambers et al., 2002). Taylor (2006) argued that the OJP represents a portion of an even larger LIP that was composed of the OJP, Manihiki (8.8 million km³) and Hikurangi Plateaus (2.7 million km³), and following eruption, this “super LIP” rifted apart into these three separate plateaus. The origin of the OJP is the source of significant controversy. The OJP has been suggested to be a product of melting a thermally-buoyant mantle plume (Mahoney and Spencer, 1991; Richards et al., 1991; Tejada et al., 2004; Fitton and Godard, 2004; Ito and Clift, 1998), melting of a largely primitive (but non-chondritic) mantle component (Jackson and Carlson, 2011), entrainment and melting of eclogite-bearing mantle near a spreading center (Korenaga et al., 2005), or melting that resulted from a giant bolide impact (Rogers, 1982; Ingle and Coffin, 2004).

Lavas from the OJP have been sampled at eight deep sea drilling sites (Mahoney et al., 1993) and along a > 3.5 km thick section in the Solomon Islands which was thrust upwards when the OJP “docked” with the Solomon arc at 27 to 23 Ma (Coleman and Kroenke, 1981; Cooper and Taylor, 1985; Neal et al., 1997; Petterson et al., 1997; Tejada et al., 2002). OJP lavas record three distinct geochemical types (e.g., Tejada et al., 1996; 2002; 2004; Neal et al., 1997; Fitton and Godard, 2004). The Kwaimbaita- and Kroenke-type lavas dominate the bulk of the OJP. Kwaimbaita-type lavas are voluminous but fractionated, while Kroenke-type lavas are less evolved (9-10.5 wt.% MgO) and may be parental to the Kwaimbaita lavas, which have identical radiogenic isotopic signatures

(Tejada et al., 2004; Fitton and Godard, 2004). Kroenke-type lavas have been found in just two drill sites, ODP Sites 1185 and 1187. The Kroenke and Kwaimbaita-type lavas have relatively flat primitive mantle normalized rare earth element (REE) patterns (Tejada et al., 2004; Fitton and Godard, 2004), Sr and Nd isotope ratios that are less geochemically depleted than MORB (mid-ocean ridge basalt; Gale et al., 2013), and Pb-isotopic compositions that plot near the geochron (Tejada et al., 2004; Jackson and Carlson, 2011).

The trace element and isotopic characteristics indicate that the Kwaimbaita and Kroenke-type lavas are derived from a mantle source that is less geochemically depleted than the depleted MORB mantle (DMM) (Tejada et al., 2004). This mantle source may actually be a primitive (but non-chondritic) mantle domain (Jackson and Carlson, 2011) if the superchondritic $^{142}\text{Nd}/^{144}\text{Nd}$ in accessible terrestrial mantle rocks is the result of higher Sm/Nd in the bulk silicate Earth relative to chondrites (Boyet and Carlson, 2005; Caro et al., 2008; Caro and Bourdon, 2010; Jackson et al., 2010; Jackson and Jellinek, 2013).

Singgalo-type lavas represent a volumetrically smaller volcanic type at OJP, forming a thin late-stage veneer over the Kroenke and Kwaimbaita-type lavas, and are thought to be the result of melting a slightly more enriched mantle source (lower $^{143}\text{Nd}/^{144}\text{Nd}$, higher $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{206}\text{Pb}/^{204}\text{Pb}$, and a slight relative enrichment in incompatible trace elements) than sampled by Kroenke and Kwaimbaita-type lavas (Tejada et al., 2002, 2004; Fitton and Godard, 2004). While Singgalo lavas sample a mantle source that is slightly more enriched than Kroenke and Kwaimbaita-type lavas, to date there is no geochemical evidence in OJP lavas for the presence of a more

97 geochemically-depleted component than observed in the Kroenke and Kwaimbaita-type
98 lavas.

99 Here we present a major, trace, and volatile element study of olivine-hosted melt
100 inclusions hosted in a single Kroenke-type OJP basaltic sample from a drill core at ODP
101 site 1187. Melt inclusions can trap heterogeneous melts prior to complete melt mixing,
102 and thus provide an opportunity to characterize more of the chemical diversity that exists
103 in melts that is not detectable in whole rock compositions. However, melt inclusion
104 studies of the OJP have been hampered by a general lack of olivine-bearing basalt
105 throughout the plateau. Olivines are extremely rare in OJP lavas, and when present, they
106 are often small and/or altered (see Supplementary Materials). Here we report fresh
107 olivines that host small (from 13 to 94 microns, long axis dimension) melt inclusions
108 (Supplementary Fig. 1 and Supplementary Table 1). To our knowledge, these represent
109 the first olivine-hosted melt inclusions identified in the OJP. The inclusions are of two
110 types: The “normal-type” inclusions exhibit relatively flat primitive mantle normalized
111 REE patterns that are similar to those observed in whole rock analyses of Kroenke-type
112 lavas, while the second type of inclusions, UDM (ultra-depleted melt) inclusions—which
113 exhibit greater incompatible element depletion than observed in the host whole rock
114 composition—are evidence for a previously unrecognized geochemical component in the
115 OJP lavas.

117 **2. Methods**

The inclusions analyzed in this study are all hosted in olivines that were separated from a single whole rock sample (ODP drill core sample 192-1187A-006R-06W [110-129 cm]). We hand crushed 100 g of lava (representing most of the available, olivine-rich material) in a mortar and pestle and obtained just 60 mg of olivine grains. Twenty-eight olivine grains hosted visible inclusions, and only 15 olivine grains (hosting 19 inclusions; Supplementary Table 1) had sufficiently large inclusions for homogenization, exposure and analysis. Images of each melt inclusion analyzed in this study are shown in Supplementary Figure 1.

Unfortunately, all melt inclusions in the studied hand sample were crystalline. Melt inclusions were individually homogenized on a Vernadsky-style heating stage mounted under a microscope at the Laboratoire Magmas et Volcans at the Université Blaise Pascal in Clermont-Ferrand, France. This allowed us to visually confirm homogenization of the inclusion and discard inclusions that burst or leaked and annealed during the heating process. As described in Le Voyer et al. (2008), the homogenization was carried out at 1 atm, the temperatures were monitored with a thermocouple, and the samples were bathed in purified helium gas during homogenization and kept at an oxygen fugacity between 10^{-10} and 10^{-9} atm. The inclusions were raised from room temperature to the liquidus temperature within a span of 20 minutes, and then quenched. Chen et al. (2011) and Gaetani et al. (2012) demonstrated that homogenization, following the Clermont procedure on the heating stage, does not result in measurable hydrogen loss from the inclusion. Following homogenization, the inclusions were mounted on glass slides and individually exposed using silicon carbide paper, followed by polishing using diamond suspension and 0.3 μm alumina paste. Following exposure and polishing, each

olivine-hosted inclusion was pressed into an aluminum mount filled with indium. The mount was cleaned using milliQ water and ethanol prior to volatile, trace and major element analyses.

Trace element analyses were carried out using the Cameca 6F ion microprobe at Arizona State University (ASU) and the analytical protocols follow those detailed in Wanless et al. (2014) and Cabral et al. (2014). A $^{16}\text{O}^-$ primary beam was accelerated to 12.5 kV with a current of 10 nA. The sample was held at 9 kV, and an “energy filtering” window of 75 ± 20 eV was used to reduce molecular ions; this approach has been used successfully in other labs (e.g., Shimizu et al., 1978; Zinner and Crozaz, 1986) and at ASU (Wanless et al., 2014; Cabral et al., 2014). Samples were pre-sputtered for 200 seconds prior to analysis, and each analysis consisted of 20 individual cycles of analyses of the following isotopes with the following dwell times: ^{30}Si (1 s), ^{47}Ti (2 s), ^{88}Sr (2 s), ^{89}Y (2 s), ^{90}Zr (2 s), ^{93}Nb (2 s), ^{138}Ba (2 s), ^{139}La (3 s), ^{140}Ce (3 s), ^{141}Pr (3 s), ^{144}Nd (3 s), ^{147}Sm (3 s), ^{174}Yb (5 s), and ^{180}Hf (5 s). Four basaltic reference glasses (KL2-G, BCR-2G, BHVO-2G, and ALV519-4-1; Gale et al., 2013; Jochum and Nohl, 2008; Melson et al., 2002) were measured to generate working curves, and ALV519-4-1 was also used as a secondary standard to monitor drift. Fifteen analyses of glass sample ALV-519-4-1 (trace element abundances reported in Gale et al. [2013] and Melson et al. [2002]) were carried out during the course of the analytical session, and are reported in Supplementary Table 2, together with reproducibility for trace element analysis: Ti ($\pm 2.5\%$ of the value reported, 2σ standard deviation), Sr ($\pm 5.3\%$), Y ($\pm 2.4\%$), Zr ($\pm 4.0\%$), Nb ($\pm 8.9\%$), Ba ($\pm 8.8\%$), La ($\pm 4.3\%$), Ce ($\pm 4.5\%$), Pr ($\pm 9.6\%$), Nd ($\pm 9.5\%$), Sm ($\pm 18.3\%$), Yb ($\pm 10.7\%$), and Hf ($\pm 16.6\%$).

Volatile (H₂O, CO₂, F, S, Cl) and P analyses were carried out at the Department of Terrestrial Magnetism, Carnegie Institution of Washington, using the 6F Cameca ion microprobe (Supplementary Table 1). In short, a 14nA Cs⁺ primary beam was used to sputter samples, and an electron gun was used to compensate for charge buildup. Samples were pre-sputtered for five minutes to eliminate surface contamination prior to analysis, and the masses of interest were measured five times with the following dwell times: ¹²C (10 s), ¹⁷OH (5 s), ¹⁹F (5 s), ³⁰Si (5 s), ³¹P (5 s), ³²S (5 s), and ³⁵Cl (5 s). A ¹²C ion image of each melt inclusion was examined prior to analysis to avoid any visible surface contamination, and analyses that exhibited drift in ¹²C/³⁰Si over the course of an analysis, indicative of surface contamination, were discarded. ALV519-4-1 was used as a secondary standard to correct for drift. Five analyses of submarine glass sample D52-5 (volatiles are reported in Simons et al., 2002) were carried out during the course of the analytical session, and are reported in Supplementary Table 2. Following correction for drift and offset, we report the following reproducibility for each of the following elements in D52-5: H₂O (±9.9%, 2σ standard deviation), CO₂ (±10.3%), F (±6.8%), S (±3.8%), Cl (±2.8%), and P (±1.7%).

Electron microprobe measurements were carried out using the electron microprobe facility at the University of Toronto on a Cameca SX-50, and the major element data are presented in Supplementary Table 1. Olivine analyses were carried out using a focused beam. For glass analyses, a defocused beam was used and employed a 15 kV accelerating voltage and a 10 nA beam current. We report the following precision for major element analyses of glass by electron microprobe: SiO₂ (±0.65 wt.%, 2σ standard deviation), TiO₂ (±0.16 wt.%), Al₂O₃ (±0.26 wt.%), FeO (±0.32wt.%), MgO (±0.25%),

CaO (± 0.28 wt.%), Na₂O (± 0.12 wt.%), MnO (± 0.08 wt.%), K₂O (± 0.04 wt.%), P₂O₅ (± 0.06 wt.%).

CO₂ analyses by ion microprobe preceded electron microprobe analyses in order to avoid surface contamination from the carbon coat that is required for major element analyses by electron probe. Unfortunately, there was insufficient melt inclusion material remaining for electron probe analyses following ion probe analyses for inclusions 2B and 9C in this study (which lack major element data; Supplementary Table 1).

Whole rock major and trace element analyses of a portion of the same sample from which the olivines (and their inclusions) were separated were performed at the GeoAnalytical Lab at Washington State University (WSU). Major elements were measured by X-ray fluorescence (XRF), and trace elements were measured by inductively-coupled plasma mass spectrometry (ICP-MS). Precision for the analyses of SiO₂, Al₂O₃, TiO₂ and P₂O₅ in basalts is 0.11 – 0.33% (1 σ) of the amount present, and 0.38 – 0.71% (1 σ) of the amount present for FeO, MgO, CaO, Na₂O, MnO and K₂O (Johnson et al., 1999). Trace element analyses by ICP-MS have a precision of 0.77 – 3.2% (1 σ) (Knaack et al., 1994).

3. Results and Observations

3.1 Major element abundances in OJP melt inclusions

Like all whole rock lava and glass compositions obtained from OJP drill sites, the melt inclusions are tholeiitic (Fig. 1). The melt inclusions are hosted in olivines with

forsterite compositions ranging from 86 to 89, except for one Fo83 olivine (Supplementary Table 1). After the major element compositions of the inclusions are corrected to be in equilibrium with the host olivine by equilibrium olivine addition or subtraction (assuming an olivine-melt K_d of 0.3 [Roeder and Emslie, 1970; Ford et al., 1983] and 10% of the total Fe is Fe^{3+}), the inclusions have MgO abundances that are similar to those previously identified in Kroenke-type lavas (Fig. 1 and Supplementary Table 1; unless stated otherwise, melt inclusion major, trace and volatile element abundances presented in the text and figures are adjusted for this olivine fractionation correction). With the exception of inclusion #2—which has the lowest MgO in the inclusion suite— SiO_2 abundances of the inclusions fall in the range previously identified for Kroenke-type lavas (Supplementary Table 1). Two inclusions (#25 and #27) have anomalously high alkali abundances compared to the whole rock and the other inclusions; we do not have an explanation for the alkali element enrichment, but we note that these two inclusions plot well within the tholeiite field defined by OJP whole rocks and glasses in Fig. 1. A group of inclusions have anomalously low $(La/Yb)_N$ (where N represents normalization to the primitive mantle composition of McDonough and Sun, 1995) compared to previously analyzed whole rocks and pillow glasses from the OJP. The inclusions with low $(La/Yb)_N$ values are discussed below, and are referred to as UDM inclusions hereafter. In major element variation diagrams (Fig. 1), the UDM inclusions do not exhibit clear major element compositional differences compared to the inclusions that have “normal” $(La/Yb)_N$ values, except for the possible exception of TiO_2 , which tends to be lower in the UDM inclusions. The slightly lower TiO_2 in the UDM inclusions relative to the inclusions with normal $(La/Yb)_N$ is reflected in higher Na_2O/TiO_2 in the

former group (Fig. 1). In a plot of $\text{CaO}/\text{Al}_2\text{O}_3$ versus $\text{Na}_2\text{O}/\text{TiO}_2$ (Fig. 1), the UDM inclusions are resolved from the normal-type inclusions and from previously examined whole rocks and pillow glasses from the OJP.

Finally, Nb and La concentrations in the melt inclusions are plotted against MgO in Fig. 1. These two highly incompatible trace elements are depleted in the UDM inclusions compared to the normal type inclusions, even though both types of inclusions span a similar range of MgO. The normal type inclusions have Nb and La concentrations that are similar to concentrations observed in whole rock analyses from the OJP. In contrast, the UDM inclusions have Nb and La concentrations that are lower than previously observed in OJP lavas.

3.2. Incompatible trace element patterns in OJP melt inclusions

Primitive mantle (McDonough and Sun, 1995) normalized trace element patterns reveal two groups of melt inclusions. The first group of inclusions has trace element patterns that are most similar to those of whole rock analyses of Kroenke-type OJP lavas (Fig. 2). The observation of these “normal-type” (i.e., Kroenke-type) inclusions is expected, as the olivines hosting the melt inclusions were isolated from a Kroenke-type lava. However, seven of the melt inclusions, referred to as UDMs, exhibit $(\text{La}/\text{Yb})_{\text{N}}$ ratios (0.18-0.32) that are up to four times lower than previously observed in any lava erupted at the OJP. Like the Kroenke-type lavas, Kwaimbaita lavas exhibit trace element patterns that are relatively flat, but shifted vertically relative to the Kroenke-type lavas (Fig. 2); this supports the hypothesis that Kroenke-type lavas are parental to (and therefore have lower incompatible trace element concentrations than) the more evolved Kwaimbaita-

type lavas (Fitton and Godard, 2004). Singgalo-type lavas have trace element patterns that exhibit slight enrichment in the incompatible trace elements relative to Kroenke and Kwaimbaita-type lavas (Figs. 2 and 3).

Two different melt inclusions hosted in the same olivine (inclusions 10A and 10B from olivine 10; Table 1) have the same normal-type trace element patterns. Similarly, two different sets of melt inclusions from two different olivines are UDM inclusions (inclusions 7A and 7B from olivine 7, and inclusions 9B and 9C from olivine 9; Supplementary Table 1). No examples of inclusions where normal-type and UDM inclusions are hosted in the same olivine were found.

There are several trace element ratios in the melt inclusion suite that clearly resolve the UDM inclusions from the normal-type inclusions, and demonstrate that the UDM inclusions are more depleted in incompatible trace elements than any melts previously identified in the OJP. In addition to having lower $(\text{La}/\text{Sm})_{\text{N}}$, the UDM inclusions have lower Nb/Zr and higher Ti/Zr than the normal-type inclusions (Fig. 3). The trace element ratios of the normal-type inclusions overlap with the ratios identified in the freshest Kroenke and Kwaimbaita-type lavas, while Singgalo-type lavas define a level of incompatible trace element enrichment that is greater than observed in the melt inclusion suite and in the Kroenke and Kwaimbaita-type lavas. In summary, no previous melts from the OJP have been observed to exhibit the level of extreme incompatible element depletion observed in the UDM inclusions.

One inclusion, UDM inclusion #25, has a large positive Sr anomaly on a primitive mantle normalized trace element pattern (Fig. 2). This inclusion is also enriched in Ba.

Large positive anomalies of Sr and Ba are common in gabbros, which relates to enrichment of Sr and Ba in plagioclase, an abundant phase in gabbros (e.g., Sobolev et al., 2000; Saal et al., 2007; Danyushevsky et al., 2003). Additionally, there is a general shift towards more plagioclase-like (low) CaO/Al₂O₃ ratios in the OJP inclusions with positive Sr anomalies. The inclusions with the largest positive Sr-anomalies (i.e., the highest Sr/Sr* values, where $Sr/Sr^* = Sr_N / (Sm_N * Nd_N)^{0.5}$), all with values from 1.23 to 1.44, also have the lowest CaO/Al₂O₃ ratios (Fig. 3). The Gabal Gerf plagioclase composition from Sobolev et al. (2000) has low CaO/Al₂O₃ (0.40) and high Sr/Sr* (3.3), and anchors the low CaO/Al₂O₃ and high Sr/Sr* portion of the melt inclusion trend (Fig. 3). We find no relationship between indicators of trace element depletion, like (La/Yb)_N, and trace element proxies for a plagioclase-rich lithology, like Sr/Sr* (Fig. 3). A role for a plagioclase signature in the OJP melt inclusions is discussed below.

3.3. Volatile abundances in OJP melt inclusions.

3.3.1. Chlorine

Cl/K ratios range from 0.19 to 4.3 over the entire melt inclusion suite. Cl/K ratios > 0.07 have been suggested to be the result of assimilation of hydrothermally altered material (Jambon et al., 1995; Kent et al. 1999a, 1999b; Lassiter et al., 2002; Michael and Schilling, 1989; Michael and Cornell, 1998; Stroncik and Haase, 2004; Kendrick et al., 2013). A similar range of elevated Cl/K ratios (0.22 to 2.8) was identified in pillow glasses obtained from a drill core of the OJP (Michael, 1999; Roberge, 2004). Both UDM and normal-type inclusions have similarly high Cl/K.

3.3.2. H₂O

The OJP melt inclusions have H₂O concentrations (0.11 to 0.70 wt.%, see Fig. 4 and Supplementary Table 1) that are similar to the range identified in pillow glasses from the OJP drill sites, where H₂O is 0.13 to 0.49 wt.% (Michael, 1999; Roberge et al., 2004, 2005). When H₂O is normalized to Ce, a non-volatile trace element that is thought to have similar solid-melt partition coefficients to H₂O during mantle melting and fractional crystallization (Michael, 1995; Dixon et al., 2002), the H₂O/Ce ratios extend to remarkably high values. The H₂O/Ce ratios in the normal-type inclusions range from 217 to 709, and the UDM inclusions have H₂O/Ce ratios that are even higher, ranging from 1853 to 2508. The higher H₂O/Ce in the UDM inclusions generally reflects lower Ce, not higher H₂O.

3.3.3. CO₂

The melt inclusions exhibit CO₂ concentrations (18-184 ppm, see Fig. 4) that fall within the range measured in OJP glasses (19-223 ppm; Michael, 1999; Roberge et al., 2004). Owing to the low solubility of CO₂ in silicate melts, most of the CO₂ in a primary melt is degassed prior to extrusion of melt on the seafloor or prior to entrapment of melts in olivines. When CO₂ is normalized to Nb, an incompatible trace element assumed to have mineral-melt partition coefficients similar to CO₂ during mantle melting and fractional crystallization (e.g., Saal et al., 2002), the CO₂/Nb ratios vary from 9 to 69 in the normal-type inclusions, and from 92 to 351 in the UDM inclusions. The CO₂/Nb ratios in the normal-type inclusions are all lower than the value of ~300 suggested for the depleted upper mantle (Koleszar et al., 2009). The higher CO₂/Nb ratios observed in the UDM inclusions extend above values proposed for the depleted upper mantle, but are still within the range suggested for CO₂-rich popping rocks dredged at mid-ocean ridge

settings (Cartigny et al., 2008). Koleszar et al (2009) found an extremely high CO₂/Nb ratio (1879) in a UDM melt inclusion from the Galapagos Islands, and they suggested that such a high ratio is not driven by unusually high CO₂, but by the low abundances of incompatible trace elements.

3.3.4. Fluorine

The F abundances reported in the melt inclusion suite are the first reported in lavas from the OJP. Concentrations of F are lower in the UDM inclusions (56-60 ppm) than in the normal-type inclusions (74-131 ppm) (Fig. 4 and Supplementary Table 1). The average F/Nd ratio for the melt inclusion suite is 21.3 ± 3.4 (1 σ , standard deviation), and this is similar to the average F/Nd ratio of ~21 suggested for global oceanic lavas (Workman et al., 2006).

3.3.5. Sulfur

Previous work showed that S concentrations in OJP pillow glasses are in the range of 692 to 1156 ppm (Roberge et al., 2004). Roberge et al. (2004) observed that ratios of S to other incompatible elements (S/K and S/Ti) slightly decrease with decreased MgO. The concentrations of S in the OJP inclusions range from 327 to 948 ppm, and show a clear, positive relationship with MgO (Fig. 4). We note that sulfides have not been observed in the OJP inclusions.

4. Discussion

OJP lavas are noted for their trace element and isotopic homogeneity. Excluding the small volume Singgalo melts, which are slightly enriched geochemically (relative to the Kroenke and Kwaimbaita-type lavas) and thought to comprise a late-stage of volcanism on the plateau (Tejada et al., 2004; Fitton and Godard, 2004), the bulk of the accessible plateau is composed of Kroenke-type and Kwaimbaita-type lavas that have indistinguishable Sr-Nd-Pb-Hf isotopic compositions and relatively flat primitive mantle normalized REE patterns (Mahoney, 1987; Mahoney and Spencer, 1991; Mahoney et al., 1993; Tejada et al., 1996, 2002, 2004, 2013; Neal, 1997; White et al., 2004; Shafer et al., 2004). These melts are thought to be the result of 20-30% melting of a homogeneous mantle source that is less depleted than the depleted MORB mantle (Mahoney et al., 1993; Neal et al., 1997; Tejada et al., 2002, 2004, 2013; Fitton and Godard, 2004; Herzberg, 2004). There has never been a report of lavas from the OJP with strongly depleted light REE signatures. Thus, the discovery of the UDM inclusions hosted in a Kroenke-type OJP lava are all the more perplexing, and demonstrate that OJP magma compositions are more diverse than previously thought.

Based on the whole rock geochemical analyses of basalts, it has been known for decades that the mantle sources of hotspots and flood basalts host geochemically-depleted components (e.g., Zindler and Hart, 1986; Gee et al., 1998; Chauvel and Hemond, 2000; Kerr et al., 1995; Fitton et al., 1997; Salters et al., 2011; Shorttle et al., 2013). For example, the presence of a depleted component distinct from the normal MORB source has been identified in Hawaii (Frey et al., 2005). Similarly, the presence of a depleted component in the Iceland plume that is distinct from the upper mantle source of normal MORB was advanced by Thirlwall (1995) and Kerr et al. (1995) on the basis

of Pb isotopes, by Fitton et al. (1997, 2003) using Nb-Zr-Y systematics, and by Kempton et al. (2000) and Fitton et al. (2003) using evidence from $^{143}\text{Nd}/^{144}\text{Nd}$ and $^{176}\text{Hf}/^{177}\text{Hf}$. In fact, basalts erupted at most major hotspots exhibit evidence for sampling a mantle source that is geochemically depleted (Hart et al., 1992), and basalts erupted at LIPs globally exhibit evidence for sampling a geochemically depleted mantle (Zindler and Hart, 1986; Jackson and Carlson, 2011).

While geochemically-depleted lavas are not unusual at hotspot and flood basalt localities in the world's ocean basins, UDM inclusions are unusual in that they exhibit greater degrees of incompatible trace element depletion than the whole rock from which they were separated. UDM melt inclusions have been reported in magmatic suites at ocean island hotspots (e.g., Sobolev et al., 2000; Koleszar et al., 2009; Gurenko and Chaussidon, 1995; Slater et al., 2001), mid-ocean ridge localities (e.g., Laubier et al., 2012; Sobolev et al., 2011; Sobolev and Shimizu, 1993; Danyushevsky et al., 2003; Kamenetsky et al., 1998) and komatiites (Shimizu et al., 2009; Kamenetsky et al., 2010).

With the exception of Iceland, the UDM inclusions from the OJP represent, to our knowledge, the first report of UDM inclusions at a LIP locality. Like the OJP melt inclusions here, Fig. 3 shows that Icelandic melt inclusions reveal a range of melt inclusion compositions that exhibit highly depleted trace element compositions. Indeed, the Icelandic melt inclusions trend to much more geochemically-depleted $(\text{La}/\text{Sm})_{\text{N}}$ values (~ 0.05) than identified in OJP inclusions (~ 0.18). Curiously, the Icelandic lavas also host melt inclusions with compositions that are have significantly more enriched $(\text{La}/\text{Sm})_{\text{N}}$ values (~ 6.5) than the OJP (~ 1.18) inclusions. Similarly, the Icelandic inclusions sample more extreme plagioclase signatures (i.e., higher Sr/Sr^*) than

identified in the OJP inclusions. While we do not have an explanation for the greater degree of geochemical variability observed in the Icelandic melt inclusions, we do note that, compared to the number of melt inclusions analyzed from the OJP, many more Iceland melt inclusions have been analyzed, from a larger suite of lavas. It is possible that the relatively small number of OJP melt inclusions may not have captured the full range of melt inclusion compositions present OJP lavas.

The origin of the incompatible element depletion observed in UDM inclusions in oceanic lavas globally is not well understood, but has been the subject of some discussion. Below, we discuss the origin of the UDM inclusions from OJP in the context of existing models for UDM formation. We argue that a distinct, depleted component inherent to the OJP mantle plume generated melts that were trapped as UDM inclusions.

4.1. Origin of the depleted OJP mantle component sampled by UDM inclusions.

4.1.1. UDM inclusions generation by repeated melt extraction from the same portion of upwelling mantle?

Following melt extraction and depletion of the OJP mantle source, subsequent melts extracted from the same portion of (now highly depleted) mantle would be depleted in incompatible elements (e.g., Sobolev and Shimizu, 1993; Thirwall et al., 2004), and we examine whether this might explain the origin of the UDM inclusions at the OJP. Sobolev and Shimizu (1993) showed that such a mechanism might explain the origin of

olivine-hosted UDM inclusions discovered in a MORB lava. They argued that the UDM inclusion represented the last fraction of melt produced by the upwelling mantle column beneath the ridge, and the same portion of the melting column had experienced continuous melt extraction, from greater depths beneath the ridge, to shallower depths where UDM is generated. As a result of melting a mantle source that has already experienced melt extraction and incompatible element depletion, the melt (i.e., the UDM) is depleted in incompatible elements. Sobolev and Shimizu (1993) suggested that MORB lavas are composed of mixtures of polybaric melts with different trace element characteristics, including inclusions with UDM characteristics. The UDM inclusions are generally mixed into bulk MORB magmas and their unique, incompatible element depleted signatures are lost during melt aggregation and homogenization with more geochemically-enriched melts. However, sometimes UDMs are trapped in growing olivines before melt homogenization, thereby preserving depleted instantaneous melts that are products of the final stages of polybaric melting of a highly depleted source.

One possible explanation for the formation of OJP UDM inclusions is that they were produced by a mechanism similar to that proposed by Sobolev and Shimizu (1993), whereby the UDM inclusions were formed by the final (shallower) stages of melting of a homogeneous upwelling OJP plume source that had become depleted in highly incompatible elements during earlier (deeper) stages of melting. In such a model, instantaneous melts of the final stages of melting would have been trapped in growing olivines in magma conduits beneath the OJP before mixing with earlier, deeper melts. The lower abundance of UDM inclusions (only just over one-third of the OJP inclusions are UDM inclusions; Supplementary Table 1) might be explained by such a model, as the

mantle source of the OJP would be expected to be more refractory during the final stages of polybaric melting, thereby producing a lower proportion of depleted melts compared to the less depleted melts extracted at greater depth. However, one potential problem with this model is that it does not explain why the OJP inclusions tend to cluster in two relatively separate groups: In trace element space (Fig. 3, panels A through D), the two inclusion types—normal-type and UDM inclusions—populate two relatively distinct groups, with perhaps one inclusion plotting between the two groups. However, if UDMs sample instantaneous polybaric melts from a homogenous upwelling OJP mantle plume, a continuous spectrum of melt inclusion compositions ranging from normal-type inclusions to UDM inclusions might be expected as the plume undergoes progressive depletion during upwelling. This is not observed.

4.1.2 A melt model to generated normal type inclusions at the OJP.

Before modeling the source of the UDM inclusions, it is first important to constrain the genesis of normal-type inclusions. Such a melt model can be used to place broad constraints on whether a single OJP mantle source can generate melts like the normal-type inclusions and, following depletion by melt extraction, generate melts similar to the UDM inclusions. The relatively flat REE patterns of the Kwaimbaita and Kroenke-type lavas, combined with the observation that the $^{143}\text{Nd}/^{144}\text{Nd}$, $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{176}\text{Hf}/^{177}\text{Hf}$ isotopic composition of the mantle source for these lavas is less depleted than the source for MORB, suggest that the trace element source of OJP Kroenke-type lavas is also less depleted than the MORB source (Tejada et al., 2004; Fitton et al., 2004).

Jackson et al. (2010) and Jackson and Carlson (2011) argued that the mantle sources of many LIPs globally, including the Kroenke lavas from the OJP, are sourced by an early-formed, depleted reservoir (EDR) generated by global differentiation event of the silicate Earth (Boyet and Carlson, 2005)

Adopting the model of Jackson and Carlson (2011), which argues that OJP lavas are melts of the EDR, we employ an aggregated near-fractional melt model of an EDR mantle source to match the trace element composition of Kroenke-type OJP lavas. We employ the alphaMELTS software (Smith and Asimow, 2005) and adopt a mantle source major element composition from Sun and McDonough (1989) and the EDR mantle source from Jackson and Jellinek (2013). The default trace element routine of alphaMELTS was used to calculate trace element fractionation, in which mineral-melt partition coefficients for clinopyroxene, garnet, and feldspar vary as a function of pressure, temperature and composition (Wood and Blundy, 1997; van Westrenen et al, 1999; Blundy and Wood 1994), while mineral-melt partition coefficients for spinel and orthopyroxene are constant values compiled by MacKenzie and O'Nions (1991; 1995). The melting calculation starts at 1600°C and 40 kilobars and proceeds until the degree of melting reaches 25-30%. The trace element composition of Kroenke-type lavas (and therefore, the normal-type inclusions, which have trace element patterns that are similar to Kroenke-type lavas) are best matched when the degree of melting of the EDR mantle is 25-30% (see Fig. 5 for melt model details). This degree of melting is in agreement with previous suggestions that Kwaimbaita and Kroenke-type OJP lavas are the result of 20-30% melting (Mahoney et al., 1993; Neal et al., 1997; Tejada et al., 2002, 2004, 2013; Fitton and Godard, 2004; Herzberg, 2004); for comparative purposes, the EDR mantle

source used here (0.65 ppm Nb and 7.7 ppm Zr; Jackson and Jellinek, 2013) is similar to a previously suggested source for Kroenke-type OJP lavas (~0.6 ppm Nb and 9.5 ppm Zr; Fitton and Goddard, 2004). Notably, the EDR mantle source that generates the Kroenke-type lavas and normal-type inclusions has concentrations of the most incompatible elements (e.g., La) that are similar to or slightly higher than the UDM inclusions (Fig. 2), and therefore cannot source the UDM inclusions. The UDM inclusions require a more incompatible element depleted mantle source.

4.1.3. A distinct mantle source for the UDM melt inclusions?

While melting the EDR mantle source does a reasonable job of generating the trace element patterns observed in the Kroenke-type lavas and the normal-type inclusions, a key trace element signature of the UDM inclusions suggests that they require a distinct mantle source unrelated to the EDR source: Unlike the normal-type OJP inclusions, which are thought to represent entrapped portions of Kroenke-type lavas, the UDM inclusions from the OJP have Zr/Zr^* (where $Zr/Zr^* = Zr_N/(Sm_N * Nd_N)^{0.5}$) values that are generally displaced to lower values than found in the normal-type OJP inclusions (Fig. 6). The low Zr/Zr^* in the UDM inclusions is difficult to generate from the same mantle source that produces the normal-type inclusions, which have generally higher Zr/Zr^* .

While low Zr/Zr^* values in UDM inclusions from other hotspot and mid-ocean ridge localities have been attributed to a gabbro signature (e.g., Sobolev et al., 2000; Saal et al., 2007; Danyushevsky et al., 2003), the low Zr/Zr^* values in the OJP UDM

inclusions cannot be the result of interaction with gabbro: there is no relationship between Zr/Zr^* and Sr anomalies (where the latter is an indicator of a gabbro signature). Thus, the lack of any relationship between Sr/Sr^* and Zr/Zr^* argues against the negative Zr anomalies in the UDM inclusions from the OJP resulting from interaction with gabbro melts. The low Zr/Zr^* signatures in the OJP UDM inclusions are interpreted to be a primary signature of the melts.

We explore whether the lower Zr/Zr^* values in OJP UDM inclusions is a geochemical signature can be generated by melting a mantle source residue left over following melt extraction of Kroenke-type lavas (and normal-type inclusions) from the EDR mantle. Fig. 6 shows the same aggregated fractional melt model of the EDR source that was explored in Fig. 5, and shows that the $(La/Yb)_N$ and Zr/Zr^* compositions of the normal-type inclusions and Kroenke-type basalts are best matched with 25-30% aggregated fractional melting of the EDR source (blue line in Fig. 6), which is consistent with the melt model result shown in Fig. 5. Melts of the EDR source do not generate the low Zr/Zr^* values observed in the UDM inclusions. However, the composition of the mantle residue (grey line in Fig. 6) following melt extraction from the EDR source has diminished $(La/Yb)_N$, which is a feature observed in UDM inclusions. Unfortunately, Zr/Zr^* in the melt model does not match the UDM inclusions: melts extracted from this depleted residue have higher Zr/Zr^* than the residue (red line in Fig. 6), while the UDM inclusions have lower Zr/Zr^* values than the residue. To demonstrate this effect, we employ the alphaMELTS software using similar assumptions employed above. However, instead of extracting melt from the EDR source, melt is re-extracted from the depleted residue following 10% fractional melting of the EDR source (red line in Fig. 6), and the

melt re-extraction starts at 29 kilobars and 1555°C and proceeds to lower pressures and temperatures. The melts have the low $(\text{La/Yb})_N$ values observed in the UDM inclusions, but the Zr/Zr^* of the model melts are higher than the UDM inclusions (Fig. 6). In general, a family of melt models show that re-melting the depleted mantle source left over after melt extraction during construction of the OJP does not generate the combination of low Zr/Zr^* and low $(\text{La/Yb})_N$ values observed in the UDM inclusions.

Multiple melt extraction events from the same portion of upwelling mantle may successfully explain the generation of an ultra-depleted source that gave rise to the UDM inclusion identified in a mid-ocean ridge basalt (Sobolev and Shimizu, 1993). However, given the generally lower Zr/Zr^* observed in the UDMs compared to normal-type inclusions and Kroenke-type basalts, the UDMs from OJP are not explained as melts extracted from the EDR or melt sextracted from the depleted mantle residue left over following extraction of Kroenke-type melts (and the associated normal-type inclusions). Instead, the UDM melts must have been extracted from a mantle source that is unrelated to (and has lower Zr/Zr^* and La/Yb than) the mantle source of the Kroenke-type lavas and normal-type inclusions. Without isotopic information on the UDM inclusions, we cannot speculate on the timing of the melt extraction event that generated the depleted mantle source of the UDMs. However, based on the arguments above, this depletion event is unlikely to have occurred during melt extraction to form the lavas that constitute the OJP.

4.1.4. Distinct mantle sources for UDM and normal-type melt inclusions

Instead of representing melts of a single plume source that has undergone progressive depletion by polybaric melt extraction during plume upwelling, the two groups of melt inclusions may represent melts of two distinct mantle sources within the upwelling plume. Recent work on mantle xenoliths from the island of Malaita has identified several lithologies—depleted harzburgites, fertile lherzolites and pyroxenites—that were suggested to have sourced OJP melts at 120 Ma (Ishikawa et al., 2007, 2011). Ishikawa et al. (2011) argued that the lithological diversity of Malaita xenoliths indicate that the mantle upwelling beneath the OJP was heterogeneous. Ishikawa et al. (2011) suggest that the homogeneity in OJP lavas could result from mixing and homogenization of melts from these diverse lithologies. If their hypothesis is correct, and the OJP magmas sampled a heterogeneous mantle that is represented by the diverse xenoliths from Malaita, then the UDM inclusions may offer a first glimpse at some of the heterogeneity that exists in primary melts prior to melt aggregation. In this model, the UDM inclusions represent melts of a more depleted portion of a geochemically-heterogeneous OJP mantle source than the normal-type inclusions. A model in which there are two distinct mantle sources for the inclusions—normal-type and UDM—might explain why the melt inclusions tend to form two rather distinct groups in Figs 3 and 6.

Isotopic measurements of the OJP melt inclusions could constrain whether UDM inclusions are melts of a mantle source formed by the final stages of polybaric melting of a homogeneous OJP plume, or melts of a more depleted portion of a heterogeneous OJP plume. If UDM inclusions were formed by the final stages of melting of a homogeneous upwelling OJP plume source that had already been depleted in highly incompatible elements during the early stages of melting (Sobolev and Shimizu, 1993), the isotopic

composition of the UDM would be expected to be similar to the normal-type inclusions. However, if the UDM inclusions are melts of a more depleted portion of a geochemically heterogeneous OJP mantle source, then the two inclusion groups would be expected to have distinct isotopic compositions. Unfortunately, the OJP inclusions in this study were too small to permit a systematic isotopic study (e.g., Pb-isotopic analysis by ion probe; Saal et al., 1998; Cabral et al., 2014).

4.2. Incompatible trace element depletion in UDMs is the result from assimilation of plagioclase-rich (gabbroic) assemblages or altered oceanic crust?

Significant discussion surrounds the origin of the gabbro signature that is apparent in a subset of melt inclusions isolated from MORB and hotspot lavas. Sobolev et al. (2000) argued that the gabbro signature in rare Hawaiian inclusions results from melting recycled gabbro incorporated into the Hawaiian mantle. However, Danyushevsky et al., (2003) argued that UDM inclusions at some localities may result from assimilation of plagioclase-rich assemblages, including gabbros, during melt ascent. While we cannot distinguish between these two models for the origin of the gabbro signature in the OJP melt inclusions, we argue that the mechanism responsible for generating the gabbro signature did not generate the UDM signatures.

Gabbro cumulate xenoliths have been found at the OJP (Kinman and Neal, 2006), and gabbro cumulates are predicted to lie in the deep portions of the OJP (Neal et al., 1997). However, trace element evidence suggests that interaction of OJP melts with gabbros did not generate the UDMs. Only one UDM inclusion from the OJP has a large

positive Sr anomaly, which suggests interaction with a plagioclase-assemblage, but the other UDM inclusions do not exhibit clear Sr anomalies within measurement error (Fig. 3). This observation indicates that the processes associated with the generation of the positive Sr anomaly (for example, gabbro assimilation in the Danyushevsky et al. [2003] model) are decoupled from the process that generates the incompatible element depletion noted in the UDM inclusions from the OJP. Danyushevsky et al. (2003) noted that, while some UDM inclusions at other localities might result from plagioclase-bearing gabbro assimilation, not all UDMs acquire their incompatible trace element depleted signature in this way. For example, they indicated that the lack of a positive Sr anomaly in the UDM measured by Sobolev and Shimizu (1993) argued against the UDM signature originating by gabbro assimilation. Laubier et al. (2012) made a similar argument for a suite of MORB melt inclusions, and argued that, while assimilation can generate positive Sr anomalies, it is unlikely to generate the incompatible trace element depletion observed in the same melt inclusions.

Similarly, Cl/K exhibits no relationship with indicators of incompatible element depletion (e.g., La/Yb) (Fig. 4). If elevated Cl/K ratios indicate interaction with hydrothermally-altered oceanic crust, then the lack of relationship between Cl/K and La/Yb argues against shallow-level assimilation of lithospheric material as the mechanism responsible for the trace element patterns in the UDM inclusions.

4.3. Interpreting the water abundances in OJP inclusions

The H₂O contents of the OJP inclusions are not correlated with any incompatible element ratios or concentrations. The H₂O concentrations of the UDM and normal-type inclusions fall in a fairly narrow range (Fig. 4) that is similar to that measured in OJP glasses (Roberge et al., 2004), while the Ce concentrations in the UDM inclusions are lower than the normal type inclusions. This leads to large variations in H₂O/Ce. The best explanation for these observations relates to the high diffusivity of H₂O in olivine. Diffusion of H₂O (but not Ce) from the host liquid through the host olivine and into H₂O-poor UDM inclusions may elevate the H₂O/Ce ratio of the inclusion (Kolezar et al., 2009; Gaetani et al., 2012). If so, the inclusions may have started out with similar H₂O/Ce ratios when they were trapped, but addition of H₂O via diffusion into the olivine raised their H₂O/Ce ratios. If the host olivines experienced relatively long residence times in large magma chambers, it would have been difficult for inclusions to avoid such diffusive equilibration.

The high Cl/K ratios provide evidence for extensive assimilation of hydrothermally altered material in much of the OJP inclusion suite, consistent with previous suggestions of widespread assimilation in the OJP (Neal and Davidson, 1989; Michael, 1999; Roberge, 2004; Kinman and Neal, 2006); both UDM and normal-type inclusions have elevated Cl/K, indicating that assimilation occurred in both suites prior to melt aggregation (Fig. 4). Assimilation of altered, and therefore hydrous, crust can increase the H₂O abundances of affected melts (e.g., Kent et al., 1999a, 1999b). Therefore, we do not argue that the H₂O abundances and the elevated H₂O/Ce ratios in the OJP melt inclusions reflect mantle source ratios, a conclusion made in Roberge et al.'s (2004) study of OJP pillow glasses. However, while the Cl/K is high in the OJP

inclusion suite, there is not a clear correlation between Cl/K and H₂O/Ce. Thus, it is difficult to argue for assimilation as the sole cause for the elevated H₂O/Ce in the inclusions. Therefore, we argue that diffusive incorporation of H₂O may operate alone or in tandem with assimilation to increase the H₂O/Ce of the OJP inclusions, and that the H₂O abundances in the inclusions do not reflect primary magmatic concentrations.

4.4. A widespread, but diluted depleted component in OJP lavas: evidence from UDM inclusions

An important question is whether melts like those sampled in UDM inclusions are relatively common in the OJP but were simply mixed together with Kroenke-type melts and became obscured by magma mixing and dilution. The trace element patterns of the least altered Kroenke-type lavas are generally smooth (Figs. 1, 2 and 3). However, the extremely limited variability may permit a small proportion of melts like the UDM inclusions to comprise (i.e., be “hidden” within) the bulk whole rock chemistry of Kroenke-type basalts without causing their trace element patterns to deviate outside the range of measured (La/Yb)_N ratios in whole rock basalts. If so, then mixing and homogenization in magma chambers could have mixed the depleted component thoroughly into the bulk lava such that the depleted component is only visible in the UDM inclusions that were trapped before complete magma mixing. A mixture of approximately 7% of a UDM melt component (using the average of the UDM inclusions) and 93% of an average Kroenke-type melt component generates a primitive mantle normalized trace element pattern that is similar to the pattern of the most depleted

Kroenke-type lava. This suggests that a small quantity of UDM melts may be “hidden” in the average bulk rock Kroenke-type lavas without forcing the trace element variability to depart from levels observed in whole rock Kroenke lavas.

The crustal portion of the OJP is estimated to be 30 km thick (Miura et al., 2004; Richardson et al., 2000), but only 3.5 km of the plateau’s stratigraphy is exposed in the Solomon islands and has been sampled (e.g., Petterson et al., 1997), while even less has been sampled by drilling. With nearly 90% of the stratigraphy of the OJP left unsampled, the possibility exists that melts like the UDM inclusions comprise a larger fraction than 7% of the bulk rock in samples from deeper in the plateau. However, only the upper stratigraphy of the OJP has been characterized. The bulk of the plateau is still beyond the reach of direct sampling efforts, and the Kroenke lava type (which we find to host UDM inclusions) was only recently discovered (Tejada and Mahoney, 2004; Fitton and Godard, 2004) as a result of a major ocean drilling effort. The deeper portions of the OJP have been suggested to be heterogeneous (Kerr and Mahoney, 2007; Kerr, 2014). If the UDM component reflects a more refractory mantle source (suggested by its trace element depletion), and if the earlier stages of OJP volcanism sampled a hotter portion of the upwelling plume, then earlier stages of volcanism at the OJP might sample this component in purer form. We suggest that, if sampled in purer form in as-yet-undiscovered OJP bulk lavas, this lava type be called Mahoney-type, to complement the Kroenke, Kwaimbaita and Singgalo-type lavas already discovered in the OJP.

5. Conclusions

675 We conducted the first geochemical study of melt inclusions from the OJP. All of
676 the olivine-hosted melt inclusions were isolated from a single hand sample from ODP
677 Site 1187. Measurements of major, trace, and volatile elements on the inclusions reveal
678 that over one third of the inclusions are highly depleted in incompatible elements (with
679 low $(\text{La/Yb})_N$ ratios, referred to as UDM inclusions), while the remaining inclusions have
680 primitive mantle normalized trace element patterns that are similar to whole rock
681 Kroenke-type lavas from the OJP. The inclusions all have elevated Cl/K (indicative of
682 assimilation of altered oceanic crust or brines), and many of the inclusions have elevated
683 Sr/Sr* (indicative of a gabbro component), but neither Cl/K nor Sr/Sr* correlates with
684 indicators of incompatible element depletion, like $(\text{La/Yb})_N$. Thus, we argue that the
685 geochemically depleted signatures of the UDM inclusions are not the result of
686 assimilation.

687 The trace element characteristics of the UDM inclusions suggest that their mantle
688 source is not related to the mantle source of the normal-type inclusions or Kroenke-type
689 lavas. First, the UDM inclusions have unique trace element signatures (e.g., low Zr/Zr*
690 values) that preclude an origin by re-melting the depleted mantle source left over after
691 melt extraction during construction of the OJP. Second, sampling of melts from a
692 progressively depleted OJP mantle component during upwelling and polybaric melt
693 extraction from a mantle plume might be expected to produce a continuous spectrum of
694 inclusion compositions between normal-type and UDM inclusions. However, there is a
695 distinct lack of inclusions that have compositions intermediate between the UDM and
696 normal-type inclusions, which suggests the presence of two distinct mantle sources that
697 give rise to the two groups of melt inclusions observed in this study. We argue that the

UDM inclusions are melts of a previously unrecognized depleted component in the OJP mantle source.

We argue that the two distinct groups of OJP inclusions sample two (perhaps lithologically) distinct mantle sources in the OJP plume. If the lithologically heterogeneous xenoliths previously reported from the Island of Malaita represent geochemical diversity in the OJP mantle source, the heterogeneous trace element compositions in the OJP inclusions represent the first hints of heterogeneity that exists in primary Kroenke and Kwaimbaita-type melts prior to melt aggregation.

Acknowledgements

MGJ acknowledges invaluable ion probe support from Rick Hervig at Arizona State University, and NSF EAR-0948878 that supports the ion probe facility. MGJ also thanks the staff at the IODP Gulf Coast Repository for assistance during a two-day stay at the facility. John MacLennan provided important insights and access to a wealth of Icelandic melt inclusion data. We acknowledge editorial handling efforts of Catherine Chauvel and reviews from Godfrey Fitton, Andrew Kerr, Allison Gale, Tim Elliott, Matthew Thirwall, Clive Neal and an anonymous reviewer. MGJ acknowledges support from NSF proposals EAR-1145202, EAR-1348082, OCE-1153894, and EAR-1347377. PJM acknowledges support from NSF grant OCE-1132038. EFR-K and KTK acknowledge French ANR grant no. 2011 Blanc SIMI 5-6 003 (project DEGAZMAG) that partly supported this work. This is Laboratory of Excellence *ClerVolc* contribution number _____.

Figure captions:

Fig. 1. Major and trace element compositions of the OJP melt inclusions compared to representative whole rock basaltic compositions (Fitton and Godard, 2004) and pillow glass compositions from OJP drill sites (Michael, 1999; Roberge et al., 2004).

In order to avoid the effects of alteration, we follow the alteration filter of Fitton and Godard (2004) and have excluded whole rock basalts with LOI > 0.5 wt.% and K_2O/P_2O_5 > 2. The composition of the Kroenke-type lava hosting the melt inclusions (192-1187A-006R-06W [110-129 cm]) is also shown. The major and trace element concentrations in the melt inclusions have been corrected for olivine fractionation to be in equilibrium with the host olivine. Whole rock and melt inclusion major element data are normalized to 100 wt.% on a volatile-free basis.

Fig. 2. Primitive mantle (McDonough and Sun, 1995) normalized trace element patterns for ultra-depleted and normal-type OJP melt inclusions. UDM inclusions are shown as red lines and normal-type inclusions are shown as blue lines. The two inclusions with the largest Sr/Sr* values, one UDM inclusion (#25) and one normal-type inclusion (#23), are highlighted with diamond symbols. The melt inclusions are compared to whole rock trace element patterns measured on Kroenke-type lavas (top panel, solid grey lines), and Kawaimbaita-type (bottom panel, grey large-dashed lines) and Singgalo-type (bottom panel, solid black lines) lavas. The whole rock analysis for this lava from which inclusions were extracted is shown (top panel, bold black large-dashed line). In the bottom panel, several reference compositions are shown: MORB

(light grey thick solid line; Gale et al., 2013), depleted MORB mantle (DMM, dark bold grey large-dashed line; Workman and Hart, 2005), average Gabal Gerf gabbro composition (Zimmer et al., 1995; small-dashed black line). The EDR composition used in the melt model is shown (grey small-dashed lines). Whole rock lava data come from Fitton and Godard (2004), and we plot only the freshest lavas following the filter outlined in the caption of Fig. 1. The trace element concentrations of the inclusions have been adjusted to reflect major element correction for olivine fractionation/addition to be in equilibrium with the host olivine (see Fig. 1 caption).

Fig. 3. Major and trace element ratios are used to distinguish the UDM inclusions and the normal-type inclusions from the OJP and to evaluate interaction with plagioclase-rich (i.e., plagioclase-rich gabbro) lithologies. The expanded views show olivine-hosted melt inclusions from Iceland, for comparison. Panels A through D use incompatible trace element ratios to show that the UDM and normal-type inclusions form separate groups. In panels E and F, the Gabal Gerf plagioclase composition from Sobolev et al. (2000) plots off the figure, and has low $\text{CaO}/\text{Al}_2\text{O}_3$ (0.40) and high Sr/Sr^* (3.3). Panels G and H show that there is no relationship between a plagioclase-rich gabbro signature (where high values of Sr/Sr^* suggest the strongest gabbro signature) and $(\text{La}/\text{Yb})_N$ signatures of the OJP inclusions. All whole rock data are from Fitton and Godard (2004). The whole-rock data (Kroenke, Kwaimbaita and Singgalo-type lavas) are filtered for alteration as discussed in the caption of Fig. 1. MORB is from Gale et al. (2013). The whole rock composition of the Kroenke-type lava hosting the melt inclusions is also shown. Error bars are 1σ , standard deviation. Symbols are the same as in previous

figures except for the addition of the Iceland melt inclusions (grey circles); Iceland data are from Gurenko and Chaussidon (1995, 1997), Slater et al. (2001) and MacLennan et al. (2003a, 2003b, 2008a, 2008b), Moune et al. (2012), Hartley et al. (2013, 2014) and Neave et al. (2014).

Fig. 4. Plots showing the behavior of CO₂, H₂O, Cl, F and S in OJP melt inclusions.

Symbols as in previous figures. OJP pillow glass compositions are from Michael (1999) and Roberge et al. (2004). For all melt inclusion data, the trace element data are corrected for olivine fractionation/addition so that the major elements are in equilibrium with the host olivine (see Fig. 1 caption); thus, two samples without major element data (2B and 9C) are excluded from the three panels where elemental concentrations are plotted. Error bars are 1 σ , standard deviation.

Fig. 5. A melt model showing that the EDR mantle source (solid grey line, no symbols) from Jackson and Jellinek (2013) generates melts similar to the Kroenke-type OJP lavas (black lines) and normal-type inclusions (blue lines). The UDM inclusions are also plotted (red lines). Two melt models are shown (described in Fig. 6 and Section 4.1.2 of the text): One represents 25% aggregated fractional melting of the EDR mantle source (grey line with circles), and the other represents 30% aggregated fractional melting of the same source (grey line with triangles). The details of the melt model are provided in the text. Kroenke-type lavas from Fitton and Goddard (2004) and the lava hosting the inclusions in this study are shown (black lines); whole rocks are filtered after Fig. 1, and are olivine fractionation corrected to be in equilibrium with Fo₉₂

olivine. Melt inclusions are also corrected for olivine fractionation to be in equilibrium with Fo₉₂ olivine to simulate primary mantle melts.

Fig. 6. A melt model shows that the UDM inclusions cannot be extracted from the same mantle source as the normal-type inclusions or from the depleted residual mantle left over following extraction of normal-type melts. The details of the melt

model are provided in the text. The melt model mantle source (EDR) is shown.

Aggregated near-fractional melt extraction (0.05 to 2% residual porosity in mantle) from this mantle source (blue line with tic marks at 1% aggregated fractional melting intervals, starting at 0% on the far right) best matches the composition of the normal-type inclusions with 25-30% aggregated fractional melting. The composition of the mantle residue following melt extraction is shown as a grey line with tic marks marking 1% fractional melt extraction (with 15% fractional melt extraction on the far left, and 0% fractional melt extraction on the right) from the EDR mantle. Melts extracted from this residue have higher Zr/Zr* than the residue, and therefore have higher Zr/Zr* than the UDM inclusions from the OJP. Following extraction of 10% fractional melt from the EDR to make a depleted residue (grey line, starting at tic labeled F=10%), we extract aggregated fractional melt from this residue (red line with aggregated fractional melting intervals at 1%, starting at 0% on the far right): while the low (La/Yb)_N values of the UDM inclusions are matched by melt extraction from a depleted source (red line), the Zr/Zr* of the model melt is too high. Error bar represents 1σ, standard deviation.

References Cited

Blundy, J., Wood, B., 1994. Prediction of crystal-melt partition-coefficients from elastic-moduli. *Nature* 372, 452-454.

Boyet, M., Carlson, R.W., 2005. ^{142}Nd evidence for early (>4.53 Ga) global differentiation of the silicate Earth. *Science* 309, 576–581.

Cabral, R.A., Jackson, M.G., Koga, K.T., Rose-Koga, E.F., Hauri, E.H., Whitehouse, M.J., Price, A.A., Day, J.M.D., Shimizu, N., Kelley, K.A., 2014. Volatile cycling of H_2O , CO_2 , F, and Cl in the HIMU mantle: A new window provided by melt inclusions from oceanic hotspot lavas at Mangaia, Cook Islands. *Geochemistry, Geophysics, Geosystems*, DOI: 10.1002/2014GC005473

Caro, G., Bourdon, B., Halliday, A.N., Quitte, G., 2008. Super-chondritic Sm/Nd ratios in Mars, the Earth and the Moon. *Nature* 452, 336-339. doi:10.1038/nature06760.

Caro, G., Bourdon, B., 2010. Non-chondritic Sm/Nd ratio in the terrestrial planets: consequences for the geochemical evolution of the mantle–crust system. *Geochim. Cosmochim. Acta* 74, 3333–3349.

Cartigny, P., Pineau, F., Aubaud, C., Javoy, M., 2008. Towards a consistent mantle carbon flux estimate: Insights from volatile systematics ($\text{H}_2\text{O}/\text{Ce}$, δD , CO_2/Nb) in the North Atlantic mantle (14°N and 34°N). *Earth Planet. Sci. Lett.* 265, 672-685. doi:

834 10.1016/j.epsl.2007.11.011.

835

836 Chambers, L.M., Pringle, M.S., Fitton, J.G., 2002. Age and duration of magmatism on
837 the Ontong Java Plateau: ^{40}Ar - ^{39}Ar results from ODP Leg 192. Abstract V72B-1271. Eos,
838 Transactions American Geophysical Union, 93, F47.

839

840 Chauvel, C., Hémond, C., 2000. Melting of a complete section of recycled oceanic crust:
841 Trace element and Pb isotopic evidence from Iceland. *Geochem. Geophys. Geosyst.* 14,
842 1999GC000002.

843

844 Chen, Y., Provost, A., Schiano, P., and Cluzel, N., 2011. The rate of water loss from
845 olivine-hosted melt inclusions. *Contrib. Mineral. Petrol.* 162, 625-636.

846

847 Coffin, M.C., Eldholm, O., 1994. Large igneous provinces: crustal structure, dimensions,
848 and external consequences. *Rev. Geophys.* 32, 1–36.

849

850 Coffin, M.C., Eldholm, O., 2000. Large igneous provinces and plate tectonics.
851 *Geophysical Monograph 121: The history and dynamics of global plate motions*, pp. 309
852 – 326.

853

854 Coleman, P.J., Kroenke, L.W., 1981. Subduction without volcanism in the Solomon
855 Islands arc. *Geo-Mar. Lett.*, 1, 129-134.

856

857 Cooper, P., Taylor, B., 1985. Polarity reversal in the Solomon Islands arc. *Nature* 314,
858 428-430.

859

860 Danyushevsky, L.V., Perfit, M.R., Eggins, S.M., Falloon, T.J., 2003. Crustal origin for
861 coupled ‘ultra-depleted’ and ‘plagioclase’ signatures in MORB olivine-hosted melt
862 inclusions: evidence from the Siqueiros Transform Fault, East Pacific Rise. *Contrib.*
863 *Mineral. Petrol.* 144, 619–637.

864

865 Dixon, J.E., L., Leist, C., Langmuir, and J.-G., Schilling, 2002. Recycled dehydrated
866 lithosphere observed in plume-influenced mid-ocean-ridge basalt, *Nature*, 420, 385-389.

867

868 Fitton, J.G., Saunders, A.D., Norry, M.J., Hardarson, B.S., Taylor, R.N, 1997. Thermal
869 and chemical structure of the Iceland plume. *Earth Planet. Sci. Lett.*, 153, 197-208.

870

871 Fitton, J.G., Saunders, A.D., Kempton, P.D., Hardarson, B.S., 2003. Does depleted
872 mantle form an intrinsic part of the Iceland plume? *Geochem. Geophys. Geosyst.*, 4,
873 2002GC000424.

874

875 Fitton, J.G., Godard, M., 2004. Origin and evolution of magmas on the Ontong Java
876 Plateau. In: Fitton, J.G., Mahoney, J.J., Wallace, P.J., Saunders, A.D. (Eds.), *Origin and*
877 *Evolution of the Ontong Java Plateau*. In: *Geol. Soc. Spec. Publ.*, vol. 229. Geological
878 Society, London, pp. 151–178.

879

880 Ford, C.F., Russell, D.G., Craven, J.A., Fisk, M.R., 1983. Olivine-liquid equilibria:
 881 Temperature, pressure and composition dependence of the crystal/liquid cation partition
 882 coefficients for Mg, Fe²⁺, Ca, and Mn. *J. Petrol.* 24, 256–265.
 883
 884 Frey, F.A., Huang, S., Blichert-Toft, J., Regelous, M., Boyet, M., 2005. Origin of
 885 depleted components in basalt related to the Hawaiian hot spot: Evidence from isotopic
 886 and incompatible element ratios. *Geochem. Geophys. Geosyst.* 6, Q02L07.
 887
 888 Gaetani, G.A., O'Leary, J.A., Shimizu, N., Bucholz, C.E., Newville, M., 2012. Rapid
 889 reequilibration of H₂O and oxygen fugacity in olivine-hosted melt inclusions. *Geology*.
 890 40, 915-918. doi: 10.1130/G32992.1.
 891
 892 Gale, A., Dalton, C.A., Langmuir, C.H., Su, Y., Schilling, J-G., 2013. The mean
 893 composition of ocean ridge basalts. *Geochem. Geophys. Geosyst.* 14, 489–518.
 894 doi:10.1029/2012GC004334.
 895
 896 Gale, A., Laubier, M., Escrig, S., Langmuir, C.H., 2013. Constraints on melting processes
 897 and plume-ridge interaction from comprehensive study of the FAMOUS and North
 898 Famous segments, Mid-Atlantic Ridge. *Earth Planet. Sci. Lett.* 365, 209-220.
 899
 900 Gee, M.A.M, Thirwall, M.F., Taylor, R.N., Lowry, D., Murton, B.J., 1998. Major
 901 controls on Reykjanes Peninsula lava chemistry. *J. Pet.* 39, 819-839.
 902

903 Gurenko, A.A., Chaussidon, M., 1995. Enriched and depleted primitive melts included in
 904 olivine from Icelandic tholeiites – origin by continuous melting of a single mantle
 905 column. *Geochim. Cosmochim. Acta.* 59, 2905–2917.
 906
 907 Gurenko, A. A., & Chaussidon, M., 1997. Boron concentrations and isotopic composition
 908 of the Icelandic mantle: evidence from glass inclusions in olivine. *Chemical Geology*,
 909 135, 21–34.
 910
 911 Hart, S.R., Hauri, E.H., Oschmann, L.A., Whitehead, J.A., 1992. Mantle Plumes and
 912 Entrainment: Isotopic Evidence. *Science* 256, 517-520.
 913
 914 Hartley, M.E., MacLennan, J., Edmonds, M., Thordarson, T., 2014. Reconstructing the
 915 deep CO₂ degassing behaviour of large basaltic fissure eruptions. *Earth Planet. Sci. Lett.*,
 916 393, 120–131.
 917
 918 Hartley, M. E., Thordarson, T., Fitton, J. G., 2013. Oxygen isotopes in melt inclusions
 919 and glasses from the Askja volcanic system, North Iceland. *Geochim. Cosmochim. Acta*,
 920 123, 55–73.
 921
 922 Hauri, E., 2002. SIMS analysis of volatiles in silicate glasses, 2: isotopes and abundances
 923 in Hawaiian melt inclusions. *Chem. Geol.* 183, 115-141.
 924
 925 Herzberg, C., 2004. Partial melting below the Ontong Java Plateau. In: Fitton, J.G.,

926 Mahoney, J.J., Wallace, P.J., Saunders, A.D. (Eds.), Origin and Evolution of the Ontong
 927 Java Plateau. In: Geol. Soc. Spec. Publ., vol. 229. Geological Society, London, pp. 179–
 928 183.
 929
 930 Ingle, S., Coffin, M.F., 2004. Impact origin for the greater Ontong Java Plateau? Earth
 931 Planet. Sci. Lett. 218, 123–134.
 932
 933 Ishikawa, A., Kuritani, T., Makishima, A., Nakamura, E., 2007. Ancient recycled crust
 934 beneath the Ontong Java Plateau: Isotopic evidence from the garnet clinopyroxenite
 935 xenoliths, Malaita, Solomon Islands. Earth Planet. Sci. Lett. 259, 134–148
 936
 937 Ishikawa, A., Pearson, D.G., Dale, C.W., 2011. Ancient Os isotope signatures from the
 938 Ontong Java Plateau lithosphere: Tracing lithospheric accretion history. Earth Planet. Sci.
 939 Lett. 301, 159–170.
 940
 941 Ito, G., Clift, P.D., 1998. Subsidence and growth of Pacific Cretaceous plateaus. Earth
 942 Planet. Sci. Lett., 161, 85–100.
 943
 944 Jackson, M.G., Carlson, R., 2011. An ancient recipe for flood basalt genesis. Nature 476,
 945 316–319.
 946
 947 Jackson, M.G., Carlson, R., Kurz, M.D., Kempton, P.D., Francis, D., Blusztajn, J., 2010.
 948 Evidence for the survival of the oldest terrestrial mantle reservoir. Nature 466, 853–856.
 949

950 Jackson, M.G., Jellinek, A.M., 2013. Major and trace element composition of the high
 951 $^3\text{He}/^4\text{He}$ mantle: Implications for the composition of the bulk silicate Earth. *Geochem.*
 952 *Geophys. Geosyst.* 14. doi: 10.1002/ggge.20188
 953
 954 Jambon, A., Déruelle, B., Dreibus, G., Pineau, F., 1995. Chlorine and bromine abundance
 955 in MORB: the contrasting behaviour of the Mid-Atlantic Ridge and East Pacific Rise and
 956 implications for chlorine geodynamic cycle. *Chem. Geol.* 126, 101-117.
 957
 958 Johnson, D.M., Hooper, P.R., Conrey, R.M., 1999. XRF analysis of rocks and minerals
 959 for major and trace elements in a single low dilution Li-tetraborate fused bead. *Advances*
 960 *in X-Ray Analysis* 41, 843–867.
 961
 962 Jochum, K.P., Nohl, U., 2008. Reference materials in geochemistry and environmental
 963 research and the GeoReM database. *Chem. Geol.* 253, 50-53.
 964
 965 Kamenetsky, V.S., Eggins, S.M., Crawford, A.J., Green, D.H., Gasparon, M., Falloon,
 966 T.J., 1998. Calcic melt inclusions in primitive olivine at 43°N MAR: evidence for melt-
 967 rock reaction/melting involving clinopyroxene-rich lithologies during MORB generation.
 968 *Earth Planet. Sci. Lett.* 160, 115–132.
 969
 970 Kamenetsky, V.S., Gurenko, A.A., Kerr, A.C., 2010. Composition and temperature of
 971 komatiite melts from Gorgona Island, Colombia, constrained from olivine-hosted melt
 972 inclusions. *Geology* 38, 1003-1006.

973
 974 Kempton, P.D., Fitton, J.G., Saunders, A.D., Nowell, G.M., Taylor, R.N., Hardarson,
 975 B.S., Pearson, G., 2000. The Iceland plume in space and time: A Sr-Nd-Hf-Pb study of
 976 the NorthAtlantic rifted margin. *Earth Planet. Sci. Lett.*, 177, 255-271.
 977
 978 Kendrick, M.A., Arculus, R., Burnard, P., Honda, M., 2013. Quantifying brine
 979 assimilation by submarine magmas: Examples from the Galápagos Spreading Centre and
 980 Lau Basin. *Geochim. Cosmoch. Acta* 123, 150-165. doi: 10.1016/j.gca.2013.09.012.
 981
 982 Kent, A.J.R., Norman, M.D., Hutcheon, I.D., Stopler, E.M., 1999. Assimilation of
 983 seawater-derived components in an oceanic volcano: evidence from matrix glasses and
 984 glass inclusions from Loihi seamount, Hawaii. *Chem. Geol.* 156, 299-319.
 985
 986 Kent, A.J., Clague, D.A., Honda, M., Stolper, E.M., Hutcheon, I.D., Norman, M.D.,
 987 1999. Widespread assimilation of a seawater-derived component at Loihi Seamount,
 988 Hawaii. *Geoch. Cosmochim. Acta* 63, 2749-2761.
 989
 990 Kerr, A. C. 2014. [Oceanic plateaus](#). In: Holland, H. D. and Turekian, K. K. eds. *Treatise*
 991 *on Geochemistry: Vol. 4: The crust. 2nd ed.* Amsterdam: Elsevier, pp. 631-667.
 992
 993 Kerr, A.C., Mahoney, J.J., 2007. [Oceanic plateaus: Problematic plumes, potential](#)
 994 [paradigms](#). *Chem. Geol.* 241, 332-353.
 995
 996 Kerr, A.C., Saunders, A.D., Tarney, J., Berry, N.H., Hards, B.H., 1995. [Depleted mantle-](#)

997 [plume geochemical signatures: No paradox for plume theories](#). *Geology* 23, 843-846.
 998
 999 Kinman, W., Neal, C.R., 2006. Magma evolution revealed by anorthite-rich plagioclase
 1000 cumulate xenoliths from the Ontong Java Plateau: Insights into LIP magma dynamics and
 1001 melt evolution. *J. Volc. Geotherm. Res.* 154, 131-157.
 1002
 1003 Knaack, C.M., Cornelius, S.B., Hooper, P.R., 1994. Trace element Analysis of rocks and
 1004 minerals by ICP-MS. Open File Report, GeoAnalytical Lab. Washington State
 1005 University, p. 10.
 1006
 1007 Koleszar, A.M., Saal, A.E., Hauri, E.H., Nagle, A.N., Liang, Y., Kurz, M.D., 2009. The
 1008 volatile contents of the Galapagos plume; evidence for H₂O and F open system
 1009 behavior in melt inclusions. *Earth Planet. Sci. Lett.* 287, 442–452.
 1010
 1011 Korenaga, J., 2005. Why did not the Ontong Java Plateau form subaerially? *Earth Planet.*
 1012 *Sci. Lett.* 234, 385–399.
 1013
 1014 Larson, R.L., 1997. Superplumes and ridge interactions between Ontong Java and
 1015 Manihiki plateaus and the Nova Canton Trough. *Geology* 25, 779– 782.
 1016
 1017 Lassiter, J.C., Hauri, E.H., Nikogosian, I.K., Barsczus, H.G., 2002. Chlorine–potassium
 1018 variations in melt inclusions from Raivavae and Rapa, Austral Islands: constraints on
 1019 chlorine recycling in the mantle and evidence for brine-induced melting of oceanic crust.

1020 Earth Planet. Sci. Lett. 202, 525-540.

1021

1022 Laubier, M., Gale, A., Langmuir, C.H., 2012. Melting and crustal processes at the

1023 FAMOUS segment (mid-Atlantic ridge): New insights from olivine-hosted melt

1024 inclusions from multiple samples, J. Pet., 53, 665-698.

1025

1026 Le Voyer, M., Rose-Koga, E.F., Laubier, M., and Schiano, P., 2008. Petrogenesis of arc

1027 lavas from the Rucu Pichincha and Pan de Azucar volcanoes (Ecuadorian arc): Major,

1028 trace element, and boron isotope evidences from olivine-hosted melt inclusions.

1029 Geochem., Geophys., Geosyst. 9, doi: 10.1029/2008GC002173.

1030

1031 MacLennan, J., 2008a. Concurrent mixing and cooling of melts under Iceland. J. Pet., 49,

1032 1931–1953.

1033

1034 MacLennan, J., 2008b. Lead isotope variability in olivine-hosted melt inclusions from

1035 Iceland. Geochim. Cosmochim. Acta, 72, 4159–4176.

1036

1037 MacLennan J., McKenzie, D., Gronvöld, K., Shimizu, N., Eiler, J. M., & Kitchen, N.,

1038 2003a. Melt mixing and crystallization under Theistareykir, northeast Iceland. Geochem.

1039 Geophys. Geosyst., 4. doi:10.1029/2003GC000558

1040

1041 MacLennan, J., McKenzie, D., Hilton, F., Gronvöld, K., & Shimizu, N., 2003b.

1042 Geochemical variability in a single flow from northern Iceland. J. Geophys. Res.,

1043 108(B1). doi:10.1029/2000JB000142

1044

1045 Mahoney, J.J., 1987. An isotopic survey of Pacific oceanic plateaus: implications for
 1046 their nature and origin. In: Keating, B., Fryer, P., Batiza, R., Boehlert, G. (Eds.),
 1047 Seamounts, Islands, and Atolls. In: Am. Geophys. Monogr., vol. 43. AGU, Washington,
 1048 DC, pp. 207–220.

1049

1050 Mahoney, J.J., Fitton, J.G., Wallace, P.J., and the Leg 192 Scientific Party, 2001.
 1051 Proceedings of the Ocean Drilling Program, Initial Report, 192.

1052

1053 Mahoney, J. J., Storey, M. Duncan, R.A., Spencer, K.J., Pringle, M. Geochemistry and
 1054 age of the Ontong Java Plateau, in In: Pringle, M., Sager, W., Sliter, W., Stein, S. (Eds.),
 1055 *The Mesozoic Pacific: Geology, Tectonics and Volcanism, Geophys. Monogr. Ser.*, vol.
 1056 77, edited by M.S. Pringle et al., pp. 233-262, AGU, Washington, D. C, 1993.

1057

1058 Mahoney, J.J., Spencer, K.J., 1991. Isotopic evidence for the origin of the Manihiki and
 1059 Ontong Java oceanic plateaus. *Earth Planet. Sci. Lett.* 104, 196-210.

1060

1061 McDonough, W.F., Sun, S.S., 1995. The composition of the Earth. *Chem. Geol.* 120,
 1062 223–253. doi:10.1016/0009-2541(94)00140-4.

1063

1064 Melson, W.G., O'Hearn, T., and Jarosewich, E., 2002. A data brief on the Smithsonian
 1065 Abyssal Volcanic Glass Data File. *Geochem., Geophys., Geosyst.* 3, doi:
 1066 10.1029/2001GC000249.

1067

1068 McKenzie, D., O'Nions, R.K., 1991. Partial Melt Distributions from Inversion of Rare-

1069 Earth Element Concentrations. *J. Petrol.*, 32, 1021-1091.

1070

1071 McKenzie, D., O'Nions, R.K., 1995. The source regions of ocean island basalts. *J. Petrol.*,

1072 36, 133-159.

1073

1074 Michael, P.J., 1999. Implications for magmatic processes at Ontong Java Plateau from

1075 volatile and major element contents of Cretaceous basalt glasses. *Geochem. Geophys.*

1076 *Geosyst.* 1, 1999GC000025.

1077

1078 Michael, P., 1995. Regionally distinctive sources of depleted MORB: Evidence from

1079 trace elements and H₂O. *Earth Planet. Sci. Lett.*, 131, 301-320.

1080

1081 Michael, P.J., 1999. Implications for magmatic processes at Ontong Java Plateau from

1082 volatile and major element contents of Cretaceous basalt glasses. *Geochem. Geophys.*

1083 *Geosyst.* 1, 1999GC000025.

1084

1085 Michael, P.J., Cornell, W.C., 1998. Influence of spreading rate and magma supply on

1086 crystallization and assimilation beneath mid-ocean ridges: Evidence from chlorine and

1087 major element chemistry of mid-ocean ridge basalts. *J. Geophys. Res.* 103, 18325-18356.

1088

1089 Michael, P.J., Schilling, J.G., 1989. Chlorine in mid-ocean ridge magmas: evidence for

1090 assimilation of seawater-influenced components. *Geochim. Cosmochim. Acta* 53, 3131-
 1091 3143.
 1092
 1093 Miura, S., Suyehiro, K., Shinohara, M., Takahashi, N., Araki, E., Taira, A., 2004.
 1094 Seismological structure and implications of collision between the Ontong Java Plateau
 1095 and Solomon Island Arc from ocean bottom seismometer-airgun data. *Tectonophysics*
 1096 389, 191– 220.
 1097
 1098
 1099 Moune, S., Sigmarsson, O., Schiano, P., Thordarson, T., & Keiding, J. K., 2012. Melt
 1100 inclusion constraints on the magma source of Eyjafjallajökull 2010 flank eruption. *J.*
 1101 *Geophys. Res.: Solid Earth*, 117(B9). doi:10.1029/2011JB008718
 1102
 1103 Neal, C.R., Davidson, J.P., 1989. An unmetasomatized source for the Malaitan alnöite
 1104 (Solomon Islands): Petrogenesis involving zone refining, megacryst fractionation, and
 1105 oceanic crustal assimilation. *Geochim. Cosmochim. Acta* 53, 1975-1990.
 1106
 1107 Neal, C.R., Mahoney, J.J., Kroenke, L.W., Duncan, R.W., Petterson, M.G. The Ontong
 1108 Java Plateau, in *Large Igneous Provinces: Continental, Oceanic and Planetary Flood*
 1109 *Volcanism, Geophys. Monogr. Ser., vol. 100*, edited by J. J. Mahoney and M. F. Coffin,
 1110 pp. 183-216, AGU, Washington, D. C., 1997.
 1111
 1112

1113 Neave, D.A., MacLennan, J., Edmonds, M., & Thordarson, T., 2014. Melt mixing causes
 1114 negative correlation of trace element enrichment and CO₂ content prior to an Icelandic
 1115 eruption. *Earth Planet. Sci. Lett.*, 400, 272–283. doi:10.1016/j.epsl.2014.05.050
 1116
 1117 Parkinson, I.J., Schaefer, B.F., and ODP Leg 192 Shipboard Scientific Party, 2001. A
 1118 lower mantle origin for the world's biggest LIP? A high-precision Os isotope isochron
 1119 from Ontong Java Plateau basalts drilled on ODP Leg 192: *Eos* (Transactions, American
 1120 Geophysical Union), v. 82, p. F47.
 1121
 1122 Petterson M.G., Neal C.R., Mahoney J.J., Kroenke L.W., Saunders A.D., Babbs T.L.,
 1123 Duncan R.A., Tolia D., McGrail, B., 1997. Structure and deformation of North and
 1124 Central Malaita, Solomon Islands: Tectonic implications for the Ontong Java Plateau -
 1125 Solomon Arc collision, and for the fate of oceanic plateaus. *Tectonophysics* 283, 1-33.
 1126
 1127 Richards, M.A., Jones, D.L., Duncan, R.A., DePaolo, D.J., 1991. A mantle plume
 1128 initiation model for the Wrangellia flood basalt and other oceanic plateaus. *Science* 254,
 1129 263-267.
 1130 Richardson, W.P., Okal, E.A., Van der Lee, S., 2000. Rayleigh-wave tomography of
 1131 the Ontong Java Plateau. *Phys. Earth Planet. Inter.* 118, 29–51.
 1132 Roberge, J., Wallace, P.J., White, R.V., Coffin, M.F., 2005 Anomalous uplift and
 1133 subsidence of the Ontong Java Plateau inferred from CO₂ contents of submarine basaltic
 1134 glasses, *Geology* 33, 501–504. doi: 10.1130/G21142.1
 1135

1136 Roberge, J., White, R.V., Wallace, P.J., 2004. Volatiles in submarine basaltic glasses
 1137 from the Ontong Java Plateau (ODP Leg 192): implications for magmatic processes and
 1138 source region compositions. In: Fitton, J.G., Mahoney, J.J., Wallace, P.J., Saunders, A.D.
 1139 (Eds.), Origin and Evolution of the Ontong Java Plateau. In: Geol. Soc. Spec. Publ., vol.
 1140 229. Geological Society, London, pp. 239-257.
 1141
 1142 Roeder, P.L., Emslie, R.F., 1970. Olivine-liquid equilibrium. Contrib. Mineral. Petrol. 29,
 1143 275–289. doi:10.1007/ BF00371276.
 1144
 1145 Rogers, G.C., 1982. Oceanic plateaus as meteorite impact signatures. Nature 299, 341–
 1146 342.
 1147
 1148 Saal, A.E., Hauri, E.H., Langmuir, C.H., Perfit, M.R., 2002. Vapour undersaturation in
 1149 primitive mid-ocean-ridge basalt and the volatile content of Earth's upper mantle, *Nature*,
 1150 419, 451-455.
 1151
 1152 Saal, A.E., Kurz , M.D., Hart, S.R., Blusztajn, J.S., Blichert-Toft, J., Liang, Y., Geist,
 1153 D.J., 2007. The role of lithospheric gabbros on the composition of Galapagos lavas. Earth
 1154 Planet. Sci. Lett. 257, 391–406.
 1155
 1156 Saal, A.E., Hart, S.R., Shimizu, N., Hauri, E.H., Layne, G.D., 1998. Pb isotopic
 1157 variability in melt inclusions from oceanic island basalts, Polynesia. Science 282, 1481-
 1158 1484.

1159

1160 Salters, V.J.M., Mallick, S., Hart, S.R., Langmuir, C.E., Stracke, A., 2011. Domains of
 1161 depleted mantle: New evidence from hafnium and neodymium isotopes. *Geochem.*
 1162 *Geophys. Geosyst.* 12, [doi:10.1029/2011GC003617](https://doi.org/10.1029/2011GC003617).

1163

1164 Shafer, J.T., Neal, C.R., Castillo, P.R. Compositional variability in lavas from the Ontong
 1165 Java Plateau: results from basalt clasts within the volcanoclastic succession at Ocean
 1166 Drilling Program Site 1184, in: J.G. Fitton, et al., (Eds.), *Origin and Evolution of the*
 1167 *Ontong Java Plateau*, Geological Society, London, Spec. Pub., vol. 229, 2004, pp. 133–
 1168 150.

1169

1170 Shimizu, N., Semet, M.P., Allègre, C.J., 1978. Geochemical applications of quantitative
 1171 ion-microprobe analysis. *Geochim. Cosmochim. Acta* 42, 1324-1334.

1172

1173 Shimizu, K., Shimizu, N., Komiya, T., Suzuki, K., Maruyama, S., Tatsumi, Y., 2009.
 1174 CO₂-rich komatiitic melt inclusions in Cr-spinels within beach sand from Gorgona Island,
 1175 Colombia *Earth Planet. Sci. Lett.* 288, 33-43.

1176

1177 Shorttle, O., MacLennan, J., Piotrowski, A.M., 2012. Geochemical provincialism in the
 1178 Iceland plume. *Geochim. Cosmochim. Acta* 122, 363–397.

1179

1180 Simons, K., Dixon, J., Schilling, J.-G., Kingsley, R., and Poreda, R. 2002. Volatiles in
 1181 basaltic glasses from the Easter-Salas y Gomez Seamount Chain and Easter Microplate:

1182 Implications for geochemical cycling of volatile elements. *Geochem. Geophys. Geosyst.*, 3,
1183 doi: 10.1029/2001GC000173.

1184

1185 Slater, L., McKenzie, D., Gronvold, K., Shimizu, N. 2001. Melt generation and
1186 movement beneath Theistareykir, NE Iceland. *J. Pet.* 42, 321-354.

1187

1188 Smith, P.M., Asimow, P.D. 2005. Adiaabat_1ph: A new public front-end to the MELTS,
1189 pMELTS, and pHMELTS models. *Geochem. Geophys. Geosyst.* 6. Q02004,
1190 doi:10.1029/2004GC000816.

1191

1192 Sobolev, A.V., Hofmann, A.W., Nikogosian, I.K., 2000. Recycled oceanic crust observed
1193 in 'ghost plagioclase' within the source of Mauna Loa lavas. *Nature* 404, 986-990.

1194

1195 Sobolev, A.V., Hofmann, A.W., Jochum, K.P., Kuzmin, D.V., Stoll, B., 2011. A young
1196 source for the Hawaiian plume. *Nature* 476, 434-437.

1197

1198 Sobolev, A.V., Shimizu, N., 1993. An ultra-depleted primary melt included in an olivine
1199 from the Mid-Atlantic Ridge. *Nature* 363, 151-154.

1200

1201 Stroncik, N.A., Haase, K.M., 2004. Chlorine in oceanic intraplate basalts: Constraints on
1202 mantle sources and recycling processes. *Geology* 32, 945. doi: 10.1130/G21027.1.

1203

1204 Sun, S.S., McDonough, W. 1989. Chemical and isotopic systematics of oceanic basalts:

1205 implications for mantle composition and processes. Geological Society, London, Special
 1206 Publications 42, 313-345.
 1207
 1208 Taylor, B., 2006. The single largest oceanic plateau: Ontong Java-Manihiki-Hikurangi.
 1209 Earth Planet. Sci. Lett. 241, 372–380.
 1210
 1211 Tejada, M.L.G., Suzuki, K., Hanyu, T., Mahoney J.J., Ishikawa, A., Tatsumi, Y.,
 1212 Chang, Q., Nakai, S. 2013. Cryptic lower crustal signature in the source of the Ontong
 1213 Java Plateau revealed by Os and Hf isotopes, Earth Planet. Sci. Lett. 377-378, 84-96.
 1214
 1215 Tejada, M.L.G., Mahoney, J.J., Neal, C.R., Duncan, R.A., Petterson, M.G., 2002.
 1216 Basement geochemistry and geochronology of Central Malaita, Solomon Islands, with
 1217 implications for the origin and evolution of the Ontong Java Plateau J. Petrol. 43, 449–
 1218 484.
 1219
 1220 Tejada, M.L.G., J.J. Mahoney, P.R. Castillo, S.P. Ingle, H.C. Sheth, D. Weis. Pin-
 1221 pricking the elephant: evidence on the origin of the Ontong Java Plateau from Pb–Sr–Hf–
 1222 Nd isotopic characteristics of ODP Leg 182 basalts, in: J.G. Fitton, et al., (Eds.), Origin
 1223 and Evolution of the Ontong Java Plateau, Geological Society, London, Spec. Pub., vol.
 1224 229, 2004, pp. 133–150.
 1225
 1226 Tejada, M.L.G., Mahoney, J.J., Duncan, R.A., Hawkins, M.P., 1996. Age and
 1227 geochemistry of basement and alkalic rocks of Malaita and Santa Isabel, Solomon

1228 Islands, southern margin of Ontong Java Plateau. *J. Petrol.* 37, 361–394.

1229

1230 Thirwall, M.F., Gee, M.A.M., Taylor, R.N., Murton, B.J., 2004. Mantle components in
 1231 Iceland and adjacent ridges investigated using double-spike Pb isotope ratios. *Geochim*
 1232 *Cosmoch Acta*, 68, 361-386.

1233

1234 Thirlwall, M.F., 1995. Generation of the Pb isotopic characteristics of the Iceland plume.
 1235 *J. Geol. Soc. Lond.*, 152, 991-996.

1236

1237 van Westrenen, W., Blundy, J., Wood, B., 1999. Crystal-chemical controls on trace
 1238 element partitioning between garnet and anhydrous silicate melt. *Am. Mineral.* 84, 838-
 1239 847.

1240

1241 Wanless, V.D., Shaw, A.M., Behn, M., Plank, T., 2014. Variations in melting dynamics
 1242 and mantle compositions along the Eastern Volcanic Zone of the Gakkel Ridge: Insights
 1243 from olivine-hosted melt inclusions. *Contrib. Min. Pet.* 167, 1-22.

1244

1245

1246 White, R.V., Castillo, P.R., Neal, C.R. Phreatomagmatic eruptions on the Ontong Java
 1247 Plateau: chemical and isotopic relationship to Ontong Java Plateau basalts, in: J.G. Fitton,
 1248 et al., (Eds.), *Origin and Evolution of the Ontong Java Plateau*, Geological Society,
 1249 London, Spec. Pub. 229, 2004, pp. 307-323.

1250

1251 Wood, B. J., Blundy, J. D. 1997. A predictive model for rare earth element partitioning

1252 between clinopyroxene and anhydrous silicate melt. *Contrib. Mineral. Petr.* 129, 166-181.

1253

1254 Workman, R.K., Hauri, E., Hart, S.R., Wang, J., Blusztajn, J., 2006. Volatile and trace

1255 elements in basaltic glasses from Samoa: Implications for water distribution in the

1256 mantle. *Earth Planetary Sci. Lett.* 241, 932-951, doi: 10.1016/j.epsl.2005.10.028.

1257 Zimmer, M., Kröner, A., Jochum, K. P., Reischmann, T., Todt, W., 1995. The Gabal Gerf

1258 complex: A Precambrian N-MORB ophiolite in the Nubian Shield, NE Africa. *Chem.*

1259 *Geol.* 123, 29-51.

1260 Zinner, E., Crozaz, G., 1986. A method for the quantitative measurement of rare earth

1261 elements in the ion microprobe. *Int. J. Mass Spectrom. Ion Processes* 69, 17-38.

1262 Zindler, A., Hart, S.R., 1986. Chemical geodynamics. *Annu. Rev. Earth Planet. Sci.*, 14,

1263 493–571, doi:10.1146/annurev.ea.14.050186.002425.

1264

1265

Figures 1 to 6

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6