Matrix-Isolated Infrared Absorption Spectrum of CH₂BrOO Radical

Xu Zhang* and Stanley P. Sander*

Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, California 91109-8099, United States

Lan Cheng, Venkatesan S. Thimmakondur and John F. Stanton*

Institute for Theoretical Chemistry, Department of Chemistry, University of Texas, Austin, Texas 78712, United States

Abstract:

The bromomethylperoxy radical, CH₂BrOO, has been generated in cryogenic matrices. Six fundamental bands for CH₂BrOO have been observed in an argon matrix at 5 K. The experimental frequencies (cm⁻¹) are: $\nu_4 = 1274.3$, $\nu_5 = 1229.4$, $\nu_6 = 1086.7$, $\nu_7 = 961.8$, $\nu_8 = 879.9$, and $\nu_{10} = 515.4$, two of which are detected for the first time. Ab initio calculations have been performed employing coupled-cluster methods. The experimental frequencies are shown to be in good agreement with the computation as well as the four bands (ν_4 , ν_6 , ν_7 and ν_8) observed by Huang and Lee in the gas phase.

Introduction:

Alkylperoxy radicals, ROO, are important intermediates in the oxidation of alkanes in the atmosphere.¹ Halogenated alkyl peroxy radicals have received attention due to the importance of chlorine (Cl) and bromine (Br) atoms in the degradation of stratospheric ozone.²⁻⁵ For instance, bromomethylperoxy radical (CH₂BrOO) can be generated *via* reaction CH₂Br + O₂ in the atmosphere (especially in the marine boundary layer), while the CH₂Br radicals are formed *via* abstraction of a hydrogen atom from CH₃Br by OH or Cl.^{6,7} The UV absorption spectrum of gaseous CH₂BrOO has been studied, revealing a broad and structureless feature at 240 nm.^{2,4,8} The rate coefficient⁴ of its self-reaction has been determined to be $(1.1 \pm 0.4) \times 10^{-12}$ cm³molecule⁻¹ s⁻¹ at 298 K. Recently, the IR spectrum of CH₂BrOO has been studied by Huang and Lee.⁹ CH₂BrOO radicals were produced by 248 nm photolysis of a flowing mixture of CH₂Br₂ and O₂. A step-scan Fourier-transform spectrometer coupled to a multipass absorption cell was employed to record temporally resolved infrared (IR) absorption spectra of reaction intermediates. In the IR spectrum, four vibrational bands were observed and assigned to *syn*-CH₂BrOO; they are $\nu_4 = 1276.1$, $\nu_6 = 1088.3$, $\nu_7 = 961.0$, $\nu_8 = 884.9$ cm⁻¹. The assignment was made to *syn*-CH₂BrOO as a carrier according to the comparison of observed vibrational wavenumbers, relative IR intensities, and rotational contours with those predicted using the B3LYP/aug-cc-pVTZ method.

In this paper, we have used a different method to generate CH₂BrOO radicals, specifically *via* the reaction CH₂Br + O₂ when the reactants in tandem supersonic nozzles are deposited in a 10 K Ar matrix. Here we report the infrared absorption spectrum of the CH₂BrOO radical in an Ar matrix. Fundamental vibrational frequencies and infrared intensities were calculated using scalar-relativistic coupled-cluster methods. The experimental frequencies and intensities are compared to computational results as well as the results obtained by Huang and Lee in the gas phase.

Experimental Section:

Analogous to earlier studies^{10,25} of matrix-isolated CH₃OO and CH₂IOO radicals, CH₂BrOO can be generated in the Ar matrix *via* reaction (1):

In this work CH₂Br radicals were generated *via* thermal decomposition of CH₂BrI (reaction (2)).

Tandem nozzles were used. The dosing nozzle for CH₂Br production is the hyperthermal nozzle described elsewhere.¹¹ Briefly, this consists of a resistively heated 1 mm diameter SiC tube at the output of a Parker General Valve Series 9 pulsed solenoid valve (100 Hz frequency and about 1 msec pulse width). The hyperthermal nozzle can be heated to *ca.* 1800 K to thermally dissociate a radical precursor; residence time for the radicals in the hyperthermal nozzle is approximately 100 μs. The CH₂BrI precursor (97% purity) was purchased from Aldrich. The dosing nozzle for O₂ was a continuous stainless steel tube at room temperature; a needle valve was used to adjust the dosing rate. The hyperthermal nozzle was mounted to the vacuum shroud of an Advanced Research System cryostat inside a water-cooled mini-chamber. The second valve was attached to the vacuum shroud by attaching the stainless steel tube *via* an O-ring-sealed compression fitting on the side of the shroud. These valves were positioned at a 45° angle with respect to each other and were approximately 2.5 cm away from the cryogenic CsI sample window. Behind the pulsed valve for the hyperthermal nozzle was the gas mixture of CH₂BrI and Ar (CH₂BrI : Ar ≈ 1 : 200) in a 1.2 L stagnation reservoir with 1.1 atm stagnation pressure. The gas mixture for the O₂ nozzle was made at approximately 0.8 atm stagnation pressure in a 4 L flask with the O₂ : Ar ratio at about 1 : 4. The pressure drop in each reservoir was measured using a capacitance manometer to determine the gas flow rate. The dosing rate for each nozzle was about 0.4 mmol per minute. The O₂ (Research grade: 99.999%) and Ar (Ultra High Purity: 99.999%) gases were purchased from Airgas. The CsI sample window was kept at 10 K during dosing (dosing time is usually a couple of hours); once the radicals were trapped in the matrix, the temperature was lowered to about 5 K and the IR spectrum of the sample was measured. Matrix spectra were recorded using a Nicolet 6700 Fourier transform infrared spectrometer equipped with a mercury/cadmium/telluride (MCT-A) detector and a Deuterated Triglycine Sulfate (DTGS) detector. The spectral resolutions of the two detectors are 0.5 and 1 cm⁻¹. The cryostat is equipped with a pair of CsI side windows to accommodate the FTIR beam.

Ab Initio Calculations:

Employing the CFOUR program package,¹² the fundamental frequencies and infrared intensities of CH₂BrOO have been computed at the coupled-cluster singles and doubles (CCSD) augmented¹³ with a perturbative treatment of triple excitations (CCSD(T)) level.¹⁴ The spin-free exact two-component theory in its one-electron variant (SFX2C-1e) has been used^{15,16} to provide a cost-effective treatment of scalar-relativistic effects. Harmonic frequencies and intensities have been calculated using atomic-natural-orbital (ANO) basis sets of triple-zeta quality (ANO1) specifically contracted for the SFX2C-1e scheme.¹⁷⁻¹⁹ The anharmonic corrections to frequencies and intensities have been obtained from second-order perturbation theory²⁰ using linear, quadratic, and cubic force fields as well as the semi-diagonal elements of the quartic force fields computed with the corresponding double-zeta ANO (ANO0) basis sets. Thirty-four core electrons (the 1s electrons of C and O, as well as the 1s, 2s, 2p, 3s, 3p, 3d electrons of Br) have been kept frozen in all the CC calculations presented here. The geometry optimization and the computations of force fields have been greatly facilitated by the availability of the analytic gradients for the SFX2C-1e scheme^{21,22} and for the CCSD(T) method^{23,24} in the CFOUR program package.

Results:

Our computational results have confirmed that the equilibrium structure of CH₂BrOO is the syn-CH₂BrOO conformer and has C₁ symmetry with 12 vibrational modes, all of which are infrared active. The anti-CH₂BrOO conformer (with the BrCOO dihedral angle equal to 180 degree) lies around 0.8 Kcal/mol higher in energy, as shown in Figure 1. Consequently, in an Ar matrix at 5K, only the syn-CH₂BrOO conformer can be observed. Figure 2 shows the equilibrium structure of CH₂BrOO (syn-CH₂BrOO conformer) obtained at the SFX2C-1e-CCSD(T)/ANO1 level. Two of the twelve modes are high frequency C-H vibrations with vibrational frequencies of about 3000 cm⁻¹. The difference between the vibrational frequencies of the two isotopologues CH₂⁷⁹BrOO and CH₂⁸¹BrOO is less than 1 cm⁻¹ for all the modes (for example, the frequency for the Br-C stretching mode in CH₂⁸¹BrOO is 0.8 cm⁻¹ smaller than that in CH₂⁷⁹BrOO) and thus is insignificant for the present study. This is in line with the finding by Huang and Lee. Intuitively this can be attributed to the fact that the substitution of ⁷⁹Br with ⁸¹Br does not significantly change the reduced mass of the Br-C diatomic fragment and thus does not

affect the Br-C stretching frequency much. In the following we focus our discussion on $\text{CH}_2^{79}\text{BrOO}$.

The IR spectra of CH_2BrOO in the $490 - 990 \text{ cm}^{-1}$ region are shown in Figure 3. The black trace (bottom) is the spectrum resulting from precursor CH_2BrI only at room temperature. The blue trace (middle) shows peaks from the pyrolysis product CH_2Br as well as the precursor. Some secondary pyrolysis products are also observed including $\text{CH}_2\text{Br-CH}_2\text{Br}$ (marked as DB), CH_3Br , CH_2CH_2 and CH_2Br_2 . For 1,2-dibromoethane (or DB), there are two conformers, trans (T) and gauche (G). The assignment for the gauche conformer is tentative due to some intensity discrepancies between the literature²⁶ and our experiment. For example, in the literature²⁶ a strong band was observed at 836 cm^{-1} for the gauche conformer, but in our spectrum, the band near 836 cm^{-1} is relatively weak. Possible explanation is that in the literature the spectrum was collected at room temperature liquid phase, while our spectrum is measured in 5 K Ar matrix. The differences in the experimental conditions may cause changes in band intensities. However, more studies are needed to better understand this issue. The pink (top) trace in Fig. 3 is the spectrum of CH_2BrI pyrolysis with O_2 added. As described earlier, CH_2Br and O_2 react in the 10 K Ar matrix to produce CH_2BrOO . The peaks marked by a green bullet (●) are assigned to the title peroxy radical (CH_2BrOO). Again $\text{CH}_2\text{Br-CH}_2\text{Br}$ (marked as DB), CH_3Br , CH_2CH_2 and CH_2Br_2 are observed as secondary pyrolysis products.

The IR spectra of CH_2BrOO in the $990 - 1290 \text{ cm}^{-1}$ region are shown in Figure 4. The black trace (bottom) is the spectrum resulting from precursor CH_2BrI only at room temperature. The blue trace (middle) shows peaks from the pyrolysis product CH_2Br as well as the precursor. Also shown are secondary pyrolysis products including $\text{CH}_2\text{Br-CH}_2\text{Br}$ (marked as DB), CH_3Br , CH_2CH_2 and CH_2Br_2 . The pink (top) trace in Fig. 4 is the spectrum of CH_2BrI pyrolysis with O_2 added. The peaks marked by a green bullet (●) are assigned to the title peroxy radical (CH_2BrOO). Again $\text{CH}_2\text{Br-CH}_2\text{Br}$ (or DB), CH_3Br , CH_2CH_2 and CH_2Br_2 are observed as secondary pyrolysis products.

Table 1 lists the matrix IR bands of CH_2BrOO along with our calculated band positions and intensities. Six fundamental vibrational modes (ν_4 , ν_5 , ν_6 , ν_7 , ν_8 and ν_{10}) are observed. All experimental frequencies are in reasonable agreement with our calculations. For the four bands

(ν_4 , ν_6 , ν_7 and ν_8) previously observed by Huang and Lee, our experimental results compare favorably with the previous report as well as our computational results (all within 5 cm^{-1}). The two new bands that we observed, ν_5 and ν_{10} , are in good agreement with our calculations. Since the band intensity for ν_{10} is small, it is marked as tentative (or T). ν_5 is also marked as tentative due to some interference by a peak from an unknown species. The band intensities are in reasonable agreement among our experimental results, our calculations and the results by Huang and Lee.

To conclude, the matrix-isolated IR absorption spectrum of the bromomethylperoxy radical CH_2BrOO has been detected in an Ar matrix. Six vibrational bands of the CH_2BrOO radical in a 5 K Ar matrix have been observed: $\nu_4 = 1274.3$, $\nu_5 = 1229.4$, $\nu_6 = 1086.7$, $\nu_7 = 961.8$, $\nu_8 = 879.9$, and $\nu_{10} = 515.4$. Two of them, ν_5 and ν_{10} , are observed for the first time.

Acknowledgement

This work was supported by the NASA Tropospheric Chemistry and Upper Atmosphere Research Programs. Additional support for this work to J.F.S. comes from the U.S. Department of Energy (Contract Number DE-FG02-07ER15884) and the Robert A. Welch Foundation of Houston, TX (Grant F-1284). The authors would also like to thank Dr. Kyle Bayes and Prof. Barney Ellison for their helpful discussions. The research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. Copyright 2016 California Institute of Technology. Government sponsorship is acknowledged.

References

- (1) Tyndall, G. S.; Cox, R. A.; Granier, C.; Lesclaux, R.; Moortgat, G. K.; Pilling, M. J.; Ravishankara, A. R.; Wallington, T. J. "Atmospheric chemistry of small organic peroxy radicals", *Journal of Geophysical Research-Atmospheres* **2001**, *106*, 12157 -12182.
- (2) Nielsen, O. J.; Munk, J.; Locke, G.; Wallington, T. J. "ULTRAVIOLET-ABSORPTION SPECTRA AND KINETICS OF THE SELF-REACTION OF CH_2BR AND CH_2BrO_2 RADICALS IN THE GAS-PHASE AT 298-K", *Journal of Physical Chemistry* **1991**, *95*, 8714 - 8719.
- (3) Sehested, J.; Nielsen, O. J.; Wallington, T. J. "ABSOLUTE RATE CONSTANTS FOR THE REACTION OF NO WITH A SERIES OF PEROXY-RADICALS IN THE GAS-PHASE AT 295-K", *Chemical Physics Letters* **1993**, *213*, 457 - 464.
- (4) Villenave, E.; Lesclaux, R. "THE UV ABSORPTION-SPECTRA OF CH_2Br AND CH_2BrO_2 AND THE REACTION-KINETICS OF CH_2BrO_2 WITH ITSELF AND WITH HO_2 AT 298K", *Chemical Physics Letters* **1995**, *236*, 376 - 384.

- (5) Wallington, T. J.; Ball, J. C.; Nielsen, O. J.; Bartkiewicz, E. "SPECTROSCOPIC, KINETIC, AND MECHANISTIC STUDY OF CH_2FO_2 RADICALS IN THE GAS-PHASE AT 298-K", *Journal of Physical Chemistry* **1992**, *96*, 1241 - 1246.
- (6) Mellouki, A.; Talukdar, R. K.; Schmoltner, A. M.; Gierczak, T.; Mills, M. J.; Solomon, S.; Ravishankara, A. R. "ATMOSPHERIC LIFETIMES AND OZONE DEPLETION POTENTIALS OF METHYL-BROMIDE (CH_3Br) AND DIBROMOMETHANE (CH_2Br_2)", *Geophysical Research Letters* **1992**, *19*, 2059 - 2062.
- (7) Orlando, J. J.; Tyndall, G. S.; Wallington, T. J.; Dill, M. "Atmospheric chemistry of CH_2Br_2 : Rate coefficients for its reaction with Cl atoms and OH and the chemistry of the CHBr_2O radical", *International Journal of Chemical Kinetics* **1996**, *28*, 433 - 442.
- (8) Sehested, J.; Bilde, M.; Mogelberg, T.; Wallington, T. J.; Nielsen, O. J. "Kinetics and mechanism of the reaction of F atoms with CH_3Br ", *Journal of Physical Chemistry* **1996**, *100*, 10989 - 10998.
- (9) Huang, Y.-H.; Lee, Y.-P. "Infrared absorption of gaseous CH_2BrOO detected with a step-scan Fourier-transform absorption spectrometer", *Journal of Chemical Physics* **2014**, *141*, 164302-1-8.
- (10) Nandi, S.; Blanksby, S. J.; Zhang, X.; Nimlos, M. R.; Dayton, D. C.; Ellison, G. B. "Polarized infrared absorption spectrum of matrix-isolated methylperoxyl radicals, $\text{CH}_3\text{OO}(\text{X})$ over-tilde $2\text{A}''$ ", *Journal of Physical Chemistry A* **2002**, *106*, 7547 - 7556.
- (11) Zhang, X.; Friderichsen, A. V.; Nandi, S.; Ellison, G. B.; David, D. E.; McKinnon, J. T.; Lindeman, T. G.; Dayton, D. C.; Nimlos, M. R. "Intense, hyperthermal source of organic radicals for matrix-isolation spectroscopy", *Review of Scientific Instruments* **2003**, *74*, 3077 - 3086.
- (12) *CFOUR, Coupled-Cluster techniques for Computational Chemistry, a quantum-chemical program package by Stanton J. F.; Gauss J.; Harding M. E.; Szalay P. G.; with contributions from Auer A. A.; Bartlett R. J.; Benedikt U.; Berger C.; Bernholdt D. E.; Bomble Y. J.; Cheng L.; Christiansen O.; Heckert M.; Heun O.; Huber C.; Jagau T.-C.; Jonsson D.; Jusélius J.; Klein K.; Lauderdale W. J.; Matthews D. A.; Metzroth T.; Mück L. A.; O'Neill D. P.; Price D. R.; Prochnow E.; Puzzarini C.; Ruud K.; Schiffmann F.; Schwalbach W.; Stopkowitz S.; Tajti A.; Vázquez J.; Wang F.; Watts J. D. and the integral packages MOLECULE (Almlöf J.; Taylor P. R.), PROPS (Taylor P. R.), ABACUS (Helgaker T.; Jensen H.J. Aa.; Jørgensen P.; Olsen J.), and ECP routines by Mitin A. V.; van Wüllen C.. For the current version, see <http://www.cfour.de>.*
- (13) Purvis, G. D.; Bartlett, R. J. "A FULL COUPLED-CLUSTER SINGLES AND DOUBLES MODEL - THE INCLUSION OF DISCONNECTED TRIPLES", *Journal of Chemical Physics* **1982**, *76*, 1910 - 1918.
- (14) Raghavachari, K.; Trucks, G. W.; Pople, J. A.; Headgordon, M. "A 5TH-ORDER PERTURBATION COMPARISON OF ELECTRON CORRELATION THEORIES", *Chemical Physics Letters* **1989**, *157*, 479 - 483.
- (15) Dyall, K. G. "Interfacing relativistic and nonrelativistic methods. IV. One- and two-electron scalar approximations", *Journal of Chemical Physics* **2001**, *115*, 9136 - 9143.
- (16) Kutzelnigg, W.; Liu, W. J. "Quasirelativistic theory equivalent to fully relativistic theory", *Journal of Chemical Physics* **2005**, *123*, 241102.
- (17) *The ANO basis sets for the SFX2C-1e scheme used in the present calculations have been obtained by recontracting the primitive functions in Ref. 18 and 19 using SFX2C-1e CCSD atomic density matrices. They are available on www.cfour.de.*
- (18) Faegri, K. "Relativistic Gaussian basis sets for the elements K-Uuo", *Theoretical Chemistry Accounts* **2001**, *105*, 252-258.
- (19) Roos, B. O.; Veryazov, V.; Widmark, P. O. "Relativistic atomic natural orbital type basis sets for the alkaline and alkaline-earth atoms applied to the ground-state potentials for the corresponding dimers", *Theoretical Chemistry Accounts* **2004**, *111*, 345 - 351.

- (20) Mills, I. M. In *In Modern Spectroscopy: Modern Research*; Rao, C. K., Mathews, C.W., Ed.; Academic Press: New York, 1972; Vol. 1, p 115.
- (21) Zou, W.; Filatov, M.; Cremer, D. "Development and application of the analytical energy gradient for the normalized elimination of the small component method", *Journal of Chemical Physics* **2011**, *134*, 244117-1-11.
- (22) Cheng, L.; Gauss, J. "Analytic energy gradients for the spin-free exact two-component theory using an exact block diagonalization for the one-electron Dirac Hamiltonian", *Journal of Chemical Physics* **2011**, *135*, 084114-1-7.
- (23) Gauss, J.; Stanton, J. F.; Bartlett, R. J. "COUPLED-CLUSTER OPEN-SHELL ANALYTIC GRADIENTS - IMPLEMENTATION OF THE DIRECT-PRODUCT DECOMPOSITION APPROACH IN ENERGY GRADIENT CALCULATIONS", *Journal of Chemical Physics* **1991**, *95*, 2623 -2638.
- (24) Watts, J. D.; Gauss, J.; Bartlett, R. J. "OPEN-SHELL ANALYTICAL ENERGY GRADIENTS FOR TRIPLE EXCITATION MANY-BODY, COUPLED-CLUSTER METHODS - MBPT(4), CCSD+T(CCSD), CCSD(T), AND QCISD(T)", *Chemical Physics Letters* **1992**, *200*, 1 - 7.
- (25) Zhang, X.; Sander, S. P.; Cheng, L.; Thimmakondur, V. S.; Stanton, J. F.; "Matrix-Isolated Infrared Absorption Spectrum of CH₂IOO Radical ", *J. Phys. Chem. A.* **2016**, *120*, 260-265.
- (26) NIST Chemistry website. And Shimanouchi, T., "Tables of Molecular Vibrational Frequencies Consolidated", National Bureau of Standards, **1972**, *Volume I*, 1-160.

Figure Captions:

Figure 1. Energy as a function of the torsional angle computed at SFX2C-1e-CCSD(T)/ANO0 level. Other geometrical parameters have been optimized for each torsional angle.

Figure 2. The equilibrium structure of CH₂BrOO (syn-CH₂BrOO conformer) with C₁ symmetry computed at the SFX2C-1e-CCSD(T)/ANO1 level.

Figure 3. IR spectra of CH₂BrOO in the 490 – 990 cm⁻¹ region. The black trace (bottom) is the spectrum of CH₂BrI at room temperature. The blue trace (middle) is the spectrum of CH₂BrI pyrolysis only at around 900 °C with no added O₂. The pink (top) trace is the spectrum of CH₂BrI pyrolysis at around 1000°C with O₂ added. As described in the text, CH₂Br and O₂ react in the 10 K Ar matrix producing CH₂BrOO. The peaks marked by a green bullet (●) are assigned to CH₂BrOO. The peaks marked by **DB(T)** are assigned to BrCH₂CH₂Br (1,2-dibromoethane) with trans conformation. The peaks marked by **DB(G)** are assigned to BrCH₂CH₂Br (1,2-dibromoethane) with gauche conformation. And the peaks marked by **P** are assigned to the precursor CH₂BrI. The peaks marked by (?) are unknown.

Figure 4. IR spectra of CH₂BrOO in the 990 – 1290 cm⁻¹ region. The black trace (bottom) is the spectrum of CH₂BrI at room temperature. The blue trace (middle) is the spectrum of CH₂BrI pyrolysis only at around 900 °C with no added O₂. The pink (top) trace is the spectrum of CH₂BrI pyrolysis at around 1000°C with O₂ added. As described in the text, CH₂Br and O₂ react in the 10 K Ar matrix producing CH₂BrOO. The peaks marked by a green bullet (●) are assigned to CH₂BrOO. The peaks marked by **DB(T)** are assigned to BrCH₂CH₂Br (1,2-dibromoethane) with trans conformation. The peaks marked by **DB(G)** are assigned to BrCH₂CH₂Br (1,2-dibromoethane) with gauche conformation. And the peaks marked by **P** are assigned to the precursor CH₂BrI. The peaks marked by (?) are unknown.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Table 1: CH₂BrOO Fundamental Vibrational Modes from This Work and Earlier Study by Huang and Lee.

		This Work					Huang and Lee [ref. 9] ^a	
Mode	Description	Ar matrix IR/10 K		CCSD(T)			gas-phase/300K	Calc.
		ν/cm^{-1}	A ratio/%	ν/cm^{-1}	A/km mol ⁻¹	A ratio/%	ν/cm^{-1}	ν/cm^{-1}
1	C-H asym stretch	na		3058	0	0	na	3054.1 (3)
2	C-H sym stretch	na		2989	7	22	na	2985.8 (11)
3	C-H scissor/umbrella	na		1424	2	7	na	1413.5 (8)
4	C-H umbrella/scissor	1274.3 ± 1.5	68 ± 13	1278	34	98	1276.1 (109)	1265.8 (84)
5	C-H rocking	1229.4 ± 2.6 (T)	28 ± 10	1229	8	23	1243 (?)	1233.5 (26)
6	O-O stretch	1086.7 ± 1.5	35 ± 7	1086	26	56	1088.3 (51)	1103.3 (42)
7	C-O stretch	961.8 ± 2.2	75 ± 20	957	35	75	961.0 (66)	943.6 (61)
8	C-H, C-Br scissor	879.9 ± 1.7	100	882	38	100	884.9 (100)	858.1 (100)
9	C-Br stretch	na		637	10	109	na	609.9 (150)
10	O-O-C bend/torsion and C-Br stretch	515.4 ± 2.0 (T)	14 ± 3	504	1	28	na	500.3 (26)
11	torsion/Br-C-O bend/C-Br stretch	na		286	1	2	na	279.2 (3)
12	torsion	na		89	1	3	na	82.6 (3)

T: Tentative assignment. A: Integrated IR intensities (in km mol⁻¹); A ratio: Ratio of integrated IR intensities relative to ν_8 mode (in %)

^a: A ratio values (in %) are listed in parentheses. And (?) means tentative assignment because of small intensity.

Black trace (bottom): spectrum of CH_2BrI
at room temperature.

Blue trace (middle): spectrum of CH_2BrI pyrolysis .

Pink trace (top): spectrum of CH_2BrI pyrolysis
with O_2 added.

Green bullet (●): CH_2BrOO ; P: CH_2BrI ;

DB (T): $\text{BrCH}_2\text{CH}_2\text{Br}$ with trans conformation;

DB (G): $\text{BrCH}_2\text{CH}_2\text{Br}$ with gauge conformation;