Cholesterol's Location in Lipid Bilayers

Drew Marquardt^{a,b,*}, Norbert Kučerka^{c,d}, Stephen R. Wassall^e, Thad A. Harroun^f, John Katsaras^{f,g,h,i,*}

^a*University of Graz, Institute of Molecular Biosciences, Biophysics Division, NAWI Graz, Humboldtstr. 50/III, 8010 Graz, Austria*

^b*BioTechMed-Graz, Austria.*

^c*Frank Laboratory of Neutron Physics, Joint Institute for Nuclear Research, 141980 Dubna - Moscow Region, Russia*

^d*Department of Physical Chemistry of Drugs, Faculty of Pharmacy, Comenius University, 832 32 Bratislava, Slovakia*

^e*Department of Physics, Indiana UniversityPurdue University Indianapolis, Indianapolis, Indiana 46202, United States*

^f*Department of Physics, Brock University, St. Catharines, Ontario L2S 3A1, Canada.*

^g*Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831, United States.*

^h*Joint Institute for Neutron Sciences, Oak Ridge, Tennessee 37831, United States.*

ⁱ*Department of Physics, University of Tennessee, Knoxville, Tennessee 37996, United States.*

Abstract

It is well known that cholesterol modifies the physical properties of lipid bilayers. For example, the much studied liquid-ordered L_o phase contains rapidly diffusing lipids with their acyl chains in the all *trans* configuration, similar to gel phase bilayers. Moreover, the L_o phase is commonly associated with cholesterol-enriched lipid rafts, which are thought to serve as platforms for signaling proteins in the plasma membrane. Cholesterol's location in lipid bilayers has been studied extensively, and it has been shown – at least in some bilayers – to align differently from its canonical upright orientation, where its hydroxyl group is in the vicinity of the lipid-water interface. In this article we review recent works describing cholesterol's location in different model membrane systems with emphasis on results obtained from scattering, spectroscopic and molecular dynamics studies.

Keywords: cholesterol, sterol, membrane dynamics, membrane structure, lipid domains

2010 MSC: 00-01, 99-00

☆Fully documented templates are available in the elsarticle package on CTAN.

*Corresponding author

Email addresses: drew.marquardt@uni-graz.at (Drew Marquardt), katsarasj@ornl.gov (John Katsaras)

1. Introduction

Few biomolecules have been scrutinized as much as cholesterol. Numerous works have been written about its life-cycle in the human body, its association with other biomolecules, and its role in human health (Myant, 1981). Moreover, many interesting physical properties have been attributed to it. Although a focused review, as this, cannot do proper justice to cholesterol’s biological importance, we will focus on recent results specifying cholesterol’s location in different model membranes, with special emphasis on data derived from scattering, spectroscopic and molecular dynamics studies.

Cholesterol is found in all animal cell membranes and is needed for their proper function, including membrane permeability and fluidity (Smith, 1991; Parasassi et al., 1995). It is also thought to act as an antioxidant, and has been implicated in cell signaling processes associated with functional domains in the plasma membrane (Petrie et al., 2000; Papanikolaou et al., 2005). As much as 90% of total cholesterol is found in the plasma membrane (PM) (Lange and Ramos, 1983), and accounts for up to 45 mol% of the membrane’s total lipid content (Yeagle, 1993) – it should be pointed out, however, that organelle membranes are almost devoid of it (Mouritsen, 2005). Since its discovery in 1769 by F.P. de la Salle (Stillwell, 2013), cholesterol has become one of the most studied biomolecules (Brown and Goldstein, 1992). However, despite the wealth of research data, many questions remain about cholesterol’s role in membranes.

Cholesterol is made up of a hydrocarbon tail, a fused planar 4-ring assembly – common to steroid hormones (i.e., testosterone and estrogen) – and a hydroxyl headgroup that helps it orient at the membrane-water interface. Although a planar molecule, cholesterol has methyl groups on its two quaternary centres, making up its so-called “rough” or β face, while their absence on the molecule’s other side account for its “smooth” or α face (Figure 1). Although there are different cholesterol isoforms (i.e., 256), only one is naturally occurring (i.e., SRSSSRRR-cholesterol) (Westover et al., 2003; Kristiana et al., 2012).

In water cholesterol aggregates, forming micelles at concentrations between 20 and 40 nM (Haberlan and Reynolds, 1973; Gilbert et al., 1975). However, due to its hydrophobic nature it has a solubility of only 1.8 mg/L Haberlan and Reynolds (1973). Cholesterol excluded from the membrane adopts a crystalline monohydrate form (Brzustowicz et al.,

Figure 1: (a) The structure of cholesterol showing its steroid skeleton of four fused rings, and its fatty acid tail. Molecular dimensions were taken from Marquardt et al. (2015d); Kučerka et al. (2008). (b) Schematic showing differences between the molecule's α - and β -faces. Hydrogen atoms are not shown for better viewing.

2002b). In this review we will report on cholesterol's location and influence on different lipid bilayers as studied by neutron scattering and nuclear magnetic resonance (NMR). Understanding cholesterol's influence on model membranes may eventually help us to resolve some of the open questions associated with this important biomolecule.

2. Cholesterol in Different Bilayer Systems

Cholesterol alters lipid membranes in complex ways, and over the years a number of experimental techniques have been used to construct phase diagrams of cholesterol-containing membranes. In this section we will review some of the experimental techniques used to determine the location of cholesterol in model membranes.

Determining cholesterol's location in model membranes has made use of a number of different sample preparations. For example, commonly used and easy to prepare multilamellar vesicles (MLVs) have the advantage of being easily hydrated under different buffer conditions. Unilamellar vesicles (ULVs) are in many ways similar to MLVs, but with the addition that they have to be extruded or sonicated from a parent MLV solution. An advantage of ULVs is that when extruded, their size is well defined, allowing for studies probing the effects of curvature on membrane organization. Model membranes can also be aligned on

rigid substrates (e.g., using single crystal silicon or glass), forming smectic layers. Breaking the symmetry allows the experiment to probe orientation with respect to the bilayer normal, but also allows for a better signal-to-noise ratio.

2.1. Scattering

Elastic neutron and x-ray scattering have been used to determine the location of cholesterol in model membrane systems (Worcester and Franks, 1976; Franks and Lieb, 1979). A thorough list of literature determining the location cholesterol via scattering can be found in Marquardt and Harroun (2014). Neutron scattering is a particularly powerful method for examining materials inherently rich in hydrogen, due to the neutron’s ability to distinguish between hydrogen (^1H) and deuterium (^2H) atoms – e.g., 63 mol% of cholesterol is hydrogen. Individual or groups of hydrogen atoms within a molecule can selectively be replaced with deuterium, and because of the large difference in neutron scattering power between ^2H and ^1H , this substitution provides the necessary contrast needed to locate groups of interest. For example, the difference in measured scattering between ^2H -cholesterol and ^1H -cholesterol identifies the location and distribution of the ^2H label within the membrane (Harroun et al., 2006a, 2008; Kučerka et al., 2010).

X-ray scattering is in many ways a complementary technique to neutron scattering. Compared to neutrons, X-ray sources offer more intense beams, greater instrumental resolution, and wavelength spreads that are approximately 2 orders of magnitude finer (Marquardt et al., 2015b). These features allow for better data quality through significantly sharper peaks and a better signal-to-noise, especially at higher scattering angles, which translates in to higher resolution scattering density profiles of the membrane. For a detailed review of scattering on membranes see Marquardt et al. (2015b).

2.2. Nuclear Magnetic Resonance

Nuclear magnetic resonance (NMR) spectroscopy is a technique that can be used in a number of different ways, allowing for studies of cholesterol in model membranes under biologically relevant conditions. Like neutron scattering, isotopic enrichment is used to “highlight” a specific functional moiety in a biomolecule. For example, analysis of solid state ^2H NMR spectra of deuterated lipid analogs have yielded the fluctuation and orientation of

acyl chain C-²H bonds with respect to the bilayer normal (Seelig, 1977; Davis, 1983). How cholesterol orients in different membranes has also been mapped with phospholipid analogs having selectively deuterated and perdeuterated acyl chains (Vist and Davis, 1990; Yasuda et al., 2014; Mihailescu et al., 2011), while cholesterol tilt in a membrane has been found to be responsive to lipid environments with deuterated analogs of cholesterol (Brzustowicz et al., 1999; Shaikh et al., 2006)

The order parameter is the quantity most often measured in ²H NMR studies. It is defined according to

$$S_{CD} = \frac{1}{2} \langle 3 \cos^2 \beta - 1 \rangle \quad (1)$$

where β is the angle for a C-²H bond relative to the bilayer normal, which constitutes the axis of motional averaging for the reorientation of lipid molecules, and the angular brackets designate a time average (Seelig, 1977). The value of the order parameter describes the degree of anisotropy of molecular motion at the site of isotopic substitution. Values typically fall in the range $0 \leq S_{CD} \leq \frac{1}{2}$ for methylene segments labeled on a phospholipid chain, the lower limit corresponding to isotropic motion and the upper limit corresponding to fast axial rotation in the all-trans configuration. However, this interpretation must be modified in cases when the most probable orientation for a C-²H bond is not perpendicular to the bilayer normal, due to conformational constraints such as on labeled sites in the vicinity of a double bond in an unsaturated phospholipid chain (Seelig and Waespe-Sarcevic, 1978) or in the rigid steroid moiety of cholesterol Marsan et al. (1999).

2.3. Molecular Dynamics Simulations

The molecular origin of how cholesterol affects lipids has been extensively probed by molecular dynamics (MD) simulations, both from a structural and thermodynamic perspective (Chiu et al., 2002; Hofsäb et al., 2003; Róg and Pasenkiewicz-Gierula, 2003; Aittoniemi et al., 2006; Tieleman and Marrink, 2006; Zhang et al., 2008; Kučerka et al., 2008; Bennett et al., 2009; Khelashvili and Harries, 2013). With MD simulation, the location, orientation and putative energy of cholesterol in different environments can be monitored over timescales as long as microseconds. Unique to MD simulations, molecular interactions are analyzed on an individual molecular basis, rather than by measuring macroscopic sample averages, and

individual events can be directly observed, rather than making inferences of actions that are too fast to measure.

3. Cholesterol's Location in Phospholipid Bilayers

As mentioned, a number of experimental techniques have been used to determine cholesterol's location in lipid bilayers (Marquardt and Harroun, 2014). Early and subsequent studies have determined cholesterol's hydroxyl group locating near the lipid-water interface with its acyl chain buried deep in the bilayer's hydrocarbon region – this is the canonical upright orientation (Franks and Lieb, 1979; Dufourc et al., 1984; Leonard et al., 2001; Kessel et al., 2001; Harroun et al., 2006a, 2008; Shrivastava et al., 2009; Subczynski et al., 2009; Kučerka et al., 2010). This arrangement has been associated with cholesterol's ability to modulate membrane properties by steric interactions with the lipid's acyl chains. In particular, cholesterol affects the lipids transmembrane thickness, lateral area and headgroup hydration monotonically as a function of the biomolecule's concentration – up to 45 mol % (Kučerka et al., 2007; Pan et al., 2008). Acyl chain hydrophobic mismatch is one obvious cause of this effect. However, another reason for bilayer thickening is the well known condensation effect induced by cholesterol (Róg and Pasenkiewicz-Gierula, 2001), whereby it increases the lipid hydrocarbon chain order. In fact, this interaction appears to be more profound when associated with hydrophobic matching, as bilayer thickening has also been reported for long-chain lipids (Kučerka et al., 2008). Interestingly, this implies that cholesterol's ordering effect on lipid hydrocarbon chain organization dominates over the ability of long-chain lipid membranes to reduce their hydrophobic mismatch. In other words, it is more favourable for cholesterol to order acyl chains and thicken membranes, rather than reduce the hydrophobic mismatch.

Compared to ester lipids, Pan et al. have determined a different mode of interaction between cholesterol and ether lipids – while maintaining its upright orientation. In ether bilayers, cholesterol is found to reside higher in the membrane and hydrogen bonds with the phosphate oxygens, behaviors that can be attributed to the absence of the ester oxygens (Pan et al., 2012). Although this change in location is only a few angstroms difference, the end result is that the ether lipid headgroups are less hydrated and the hydrocarbon chain packing

is altered (again, compared to ester lipids), such that it causes cholesterol to tilt (Pan et al., 2012).

3.1. Cholesterol Ordering of Membranes

The ordering effect that cholesterol has on lipid bilayers is best observed by ^2H -NMR experiments, that determine acyl chain order parameters in essentially, a non-invasive manner. A signature profile of order parameter exists along the saturated chains of phospholipids in the lamellar liquid crystalline phase (Seelig, 1977). As exemplified by the smoothed profile generated for the perdeuterated palmitoyl *sn*-1 chain in 1- $[\text{}^2\text{H}_{31}]$ palmitoyl-2-oleoylphosphatidylcholine (POPC-d31), there is a plateau region of slowly varying order ($S_{CD} \sim 0.2$) in the top half of the chain followed by a progressively more rapid drop-off in the bottom half, towards the disordered terminal methyl group ($S_{CD} \sim 0.02$) (Shaikh et al., 2015), see Figure 2.

POPC-d31 – a lipid representative of a naturally occurring membrane phospholipid – exhibits a continuous increase in order that almost doubles in magnitude as a function of cholesterol content over 0-45 mol% (Lafleur et al., 1990). This is because cholesterol’s fused ring structure imposes a restriction on reorientation in the upper motion part of the phospholipid chains, that then propagates into the lower part. There are, however, differences in the magnitude of the response and its dependence upon the concentration of the sterol with different lipids.

The ordering effect exerted on the perdeuterated N-palmitoyl chain in $[\text{}^2\text{H}_{31}]$ -N-palmitoylsphingomyelin (PSM-d₃₁) flattens off in the presence of ≥ 25 mol% cholesterol (Bartels et al., 2008; Bunge et al., 2008) (Figure 2). This behavior was attributed to a more favorable interaction with the saturated sphingolipid than the monounsaturated phospholipid. Greater order is also attained in raft-like mixtures of PSM-d₃₁ with cholesterol, where order parameters $S_{CD} \geq 0.4$ within the plateau region approach the value ($S_{CD}=0.5$) for an all-trans acyl chain undergoing fast axial rotation in an idealized L_o phase. The differential in order between saturated N-stearoylsphingomyelin (SSM) and 1-palmitoyl-2-stearoylphosphatidylcholine (PSPC) bilayers containing 50 mol% cholesterol, however, is small as judged by quadrupolar splittings measured for analogs of these structurally similar lipids selectively deuterated in N-acyl and

Figure 2: Order parameters (S_{CD}) vs carbon number for chain-deuterated PSM-d₃₁ and POPC-d₃₁ without (open circles) and with 33 mol% cholesterol at 60 °C. Note the dramatic increase in order along the fatty acid chain away from the terminal methyl. S_{CD} data were reproduced from Bartels et al. (2008).

sn-2 acyl chains, respectively (Yasuda et al., 2014). A less steep dependence upon temperature for quadrupolar splittings in SSM, compared to PSPC, was taken to signify greater thermal stability for SM-cholesterol bilayers, from which it was inferred that SM would be favored in rafts over saturated PC.

At the opposite extreme are polyunsaturated fatty acid (PUFA) containing phospholipids that have a low affinity for cholesterol due to the highly disordered PUFA chains (Wassall and Stillwell, 2008). Less condensation of 1- $^{[2}\text{H}_{35}$]stearoyl-2-docosahexaenoylphosphatidylcholine (SDPC- d_{35}) than saturated and monounsaturated bilayers is indicated by smoothed order parameter profiles constructed for the perdeuterated stearoyl *sn*-1 chain following the addition of 29 mol% cholesterol (Mihailescu et al., 2011). Reflecting a redistribution of DHA chains towards the middle of the bilayer, as identified by neutron scattering experiments, the effect is slightly more marked in the lower portion of the chain. Support for the notion that cholesterol preferentially interacts with saturated chains was derived from moment analysis of ^2H NMR spectra in a study of another polyunsaturated phospholipid that revealed a similar incremental increase in order for the perdeuterated palmitic acid chain in 1- $^{[2}\text{H}_{31}$]palmitoyl-2-arachidonylphosphatidylcholine (PAPC- d_{31}) and 1- $^{[2}\text{H}_{31}$]palmitoyl-2-linoleoylphosphatidylcholine (PLPC- d_{31}) with four and two double bonds, respectively (Jackman et al., 1999).

3.2. Cholesterol Volumes in Lipid Bilayers

Molecular volumes of cholesterol are required as an input for evaluating the structural details of lipid bilayers – for the detailed evaluation of x-ray or neutron scattering data, for example. The volumes of cholesterol in different lipid bilayers are listed in Table 1. The nomenclature n1:u1-n2:u2 PC is used, where n1 and n2 are the number of carbons in the *sn*-1 and *sn*-2 chains, respectively, and u1 and u2 are the number of unsaturated bonds in the respective chain. Cholesterol volumes increase significantly at concentrations beyond the liquid-disordered (L_d) phase boundary – which varies from system-to-system. The generally accepted concentration where this volume increase takes place is between 20 and 30 mol% cholesterol.

From volume measurements (Greenwood et al., 2006; Gallová et al., 2015) it has been

Table 1: Cholesterol volumes in lipid bilayers

“Solvent”	V_{cho}	
H ₂ O	627 ^{*a}	
H ₂ O	606 ^{**a}	
di16:0ePC (40°C)	676.3 ^b	
di16:0ePC (50°C)	575.5 ^b	
	$\underline{L_d}$	$\underline{L_o}$
di16:0PC	574 ^c , 606 ^a	637.1 ^c , 647.1 ^a
di14:0PC	565.1 ^a	637.5 ^a
sphingomyelin	575.2 ^a	639.5 ^a
16:0-18:1PC	622.6 ^a	
di18:1PC	632.9 ^{a,d}	
di20:1PC	627 ^d	
di22:1PC	621 ^d	
di24:1PC	620 ^d	

*anhydrous cholesterol species; **monohydrate cholesterol species

^aref. Greenwood et al. (2006); ^bref. Pan et al. (2012); ^cref. Miyoshi et al. (2014); ^dref. Gallová et al. (2015)

surmised that cholesterol mixes ideally with monounsaturated phosphocholines in the fluid phase. Average thermal expansivity for diN:1PC with N = 18, 20 and 22 is $(71.5 \pm 1.1) 10^{-5} \text{ K}^{-1}$. The partial volume of cholesterol also decreases slightly with the lengthening of the diN:1PC host molecule (Gallová et al., 2015). Lipids with saturated chains, on the other hand, exhibit two mixing regimes, and cholesterol partial volumes clearly show the molecule’s well known condensation effect (Greenwood et al., 2006).

3.3. Cholesterol-PUFA Interaction

Neutron scattering studies have enabled us to locate cholesterol in PUFA bilayers, namely DAPC (20:4-20:4PC). They revealed cholesterol preferentially sequestering near the bilayer center (Figure 3 C), in contrast to its usual location, where its hydroxyl group locates near the lipid-water interface (i.e., upright orientation, Figure 3 A). Of note were experiments detailing cholesterol’s orientation in mixed lipid bilayer systems (Kučerka et al., 2009). For

Figure 3: Using different experimental techniques (i.e., neutron scattering and NMR) and MD simulations, the location of cholesterol has been determined in different membrane environments. A) The “canonical location”, B) cholesterol interdigitated between leaflets, and C) cholesterol sequestered in the center of polyunsaturated fatty acid (PUFA) bialyers.

example, when “titrating” POPC (16:0-18:1PC) into DAPC bilayers, it was found that it took ~ 50 mol% of POPC to cause cholesterol to revert to its upright orientation, while only 5 mol% of the disaturated DMPC (14:0-14:0 PC) lipid was needed to achieve the same effect. This result is also consistent with previous data showing cholesterol’s reduced solubility and binding affinity in PUFA membranes (Niu and Litman, 2002; Shaikh et al., 2006; Williams et al., 2013). The demonstration of cholesterol’s affinity for lipids with saturated hydrocarbon chains, and its aversion for PUFAs, suggests the structural, and perhaps even functional, significance of lipid diversity exhibited by the lipidome.

Insensitivity to the degree of unsaturation in the *sn*-2 chain is revealed by tilt angles (α_0) calculated from order parameters measured with $[3\alpha\text{-}^2\text{H}_1]\text{cholesterol}$ – an analog labeled at the 3α position – incorporated at an equimolar concentration into bilayers comprised of phospholipids possessing a saturated *sn*-1 chain (Murari et al., 1986; Brzustowicz et al., 1999; Shaikh et al., 2006). The tilt angle α_0 , representing the most probable orientation of the long molecular axis, was extracted by invoking a Gaussian distribution to describe the angular fluctuations (Oldfield et al., 1978). A value of $\alpha_0 = 16\text{--}17^\circ$ was found in membranes

ranging from 1,2-dipalmitoylphosphatidylcholine (DPPC) with saturated palmitic acid at the *sn*-1 and *sn*-2 positions to SDPC with saturated stearic acid, and polyunsaturated DHA at the *sn*-1 and *sn*-2 positions, respectively. Preferential interaction with saturated chains is implied.

The planar facade of the tetracyclic ring is, however, more compatible with close proximity to a saturated chain that adopts a largely linear configuration as opposed to, most acutely, a PUFA chain that rapidly interconverts through all conformational space (Gawrisch and Soubias, 2008). An appreciably larger tilt angle, $\alpha_0 = 24 - 25^\circ$, was determined in 1,2-diarachidonylphosphatidylcholine (DAPC) and 1,2-didocosa-hexaenoylphosphatidylcholine (DDPC) membranes prepared from 1:1 mol mixtures of phospholipid and $[3\alpha\text{-}^2\text{H}_1]\text{cholesterol}$ (Brzustowicz et al., 2002a,b). The aversion cholesterol has for PUFA in these membranes, where intimate contact cannot be avoided, reduces its solubility by more than a factor of 3 relative to most phospholipids, including their counterparts 1-stearoyl-2-arachidonylphosphatidylcholine (SAPC) and SDPC with a saturated *sn*-1 chain. Moreover, a non-standard location is identified by neutron scattering experiments that show both the head group and tail of cholesterol reside near the middle of DAPC bilayers (Harroun et al., 2006b, 2008). The non-standard location of cholesterol has only been observed in phospholipids where the *sn*-1 and *sn*-2 positions are populated with PUFA chains, and not in mixed chain lipids (Harroun et al., 2006b; Mihailescu et al., 2011). It should also be noted here that PUFA chains are extremely prone to the oxidation. The original studies therefore utilized various precautions to reduce lipid oxidation, including conducting the experiments in an oxygen free environment, monitoring bilayer break-down throughout the experiments, and post data collection analysis of the samples conducted to evaluate sample integrity.

Despite its aversion to PUFAs, cholesterol has been proposed as an antioxidant. From a chemical standpoint, cholesterol can serve as a free radical scavenger. Unlike α -tocopherol whose primary function is widely thought to be the protection of PUFA containing phospholipids (Traber and Atkinson, 2007), cholesterol's hydroxyl group is not the sacrificial moiety. Instead the unsaturated bond at the Δ^5 -6 position on the steroid moiety is responsible for antioxidant activity (Smith, 1991; Girotti, 2001). A deeper location of cholesterol within the bilayer means that the reactive center on the sterol is better placed to intercept lipid peroxyl

radicals than the hydroxyl group in alpha-tocopherol, which typically sits near the aqueous interface (DMPC bilayers being the exception) (Marquardt et al., 2013, 2014, 2015c). It should be borne in mind, however, that recent MD simulations reveal alpha-tocopherol can easily tunnel into the bilayer’s interior, as a result of the biomolecule’s methyl groups shielding its hydroxyl group (Leng et al., 2015).

Chemical activity does not necessarily signify *in vivo* function, as illustrated, for example, by DNA (Azzi, 2007). Smith (1991) makes several arguments for cholesterol having a protective *in vivo* role, namely: (i) it exists in environments which are susceptible to oxidative stress; (ii) it can be readily oxidized by reactive oxygen species; (iii) oxidized cholesterol is commonly found *in vivo*; and (iv) oxidized cholesterol is metabolized and excreted from the body. Other studies attribute the apparent protective properties of cholesterol purely to structural interactions and modifications induced to the bilayer (Parasassi et al., 1995; Pandey and Mishra, 1999). Knowing the impact that cholesterol has on membrane organization, the structural interaction hypothesis seems plausible. Although much of the sterols’s role in biological membranes is still not clear, it is generally accepted that when created in the distant past, cholesterol’s biological function was to protect against oxygen – lung surfactant, for example, is rich in cholesterol (Brown and Galea, 2010).

3.4. Bilayer Leaflet Distribution

Although cholesterol’s location(s) in biological membranes is now reasonably well established, its distribution between bilayer leaflets is less so. Compositional asymmetry is central for biological membranes, and it is generally accepted that cholesterol is (not surprisingly) asymmetrically distributed between bilayer leaflets (Marquardt et al., 2015a; Nickels et al., 2015). For example, plasma membrane data suggests a greater concentration of cholesterol in the cytosolic facing monolayer (Wood et al., 2011), and theoretical considerations put forth by Giang and Schick (2014) support these findings. Cholesterol, it seems, prefers the high negative curvature offered by the inner bilayer leaflet, in effect lowering the bilayer’s bending free energy. Its asymmetric distribution has also indirectly been observed in mono-unsaturated PC lipids (Kučerka et al., 2009). However, in those studies they were not able to resolve in which bilayer leaflet cholesterol was most abundant.

Coarse-grain MD simulations have also produced some interesting results. For example, Yesylevskyy and Demchenko (2012) showed that cholesterol distributes asymmetrically in a lipid bilayer after $\sim 10 \mu s$. Of note, is that its equilibrium distribution was dependent on the individual bilayer leaflet chemical composition, highlighting again the importance of lipid diversity in biomembranes.

3.5. Bilayer Leaflet Re-distribution

As mentioned, in erythrocytes and nucleated cells cholesterol is believed to be enriched on the cytosolic leaflet of the plasma membrane (negative curvature), which when one considers the typical membrane content of cholesterol (40 mol%) implies a cholesterol-to-lipid ratio of greater than 1 (van Meer, 2011). This mol% value is near the solubility limit of cholesterol in PC bilayers. Thus a continually balanced redistribution of cholesterol between leaflets is key to maintaining an adequate liquid L_o phase. Asymmetry is stabilized by the relatively quick – compared to lipids with their larger and/or more polar headgroups – flip-flop rate of cholesterol across the lipid bilayer. In many cases, however, protein intervention is needed to shuttle cholesterol across the bilayer (van Meer, 2011).

The flip-flop rates of cholesterol have been determined by a number of different experimental methods and by MD simulations, which have proven very valuable due to the complexity of probe-free studies. MD studies performed by Jo et al. (2010) and Bennett et al. (2009) yielded consistent average single molecule flip-flop ranging from 1.4 ms to 80 ns – depending on temperature and membrane composition. The prevailing trend in those studies is that cholesterol flip-flop increases with membrane disorder, which in turn increases with temperature and unsaturated lipid content, but decreases with lipid chain length. The thinner and more disordered lipid bilayers experience a larger degree of cholesterol interdigitation between the two bilayer leaflets – as their hydroxyl’s hydrogen bonding partners are closer to the bilayer center – and are significantly more permeable to water. These waters cause cholesterol to reorient into the nonpolar bilayer center (Figure 4). Kučerka et al. (2008) speculate that these features create a locally polar environment, promoting cholesterol flip-flop and stabilizing cholesterol’s hydroxyl group within the nonpolar bilayer core.

Figure 4: Sequential snapshots show dynamically changing interactions of cholesterol and lipids during flip-flop. Cholesterol's reorientation is promoted by changing its hydrogen bonding from the lipid's phosphate to its carbonyl, a moiety that is significantly closer to the bilayer center (suggested by the blue line). Cholesterol's hydroxyl is accompanied by water molecules that further facilitate this transition. Figure adapted from Kučerka et al. (2008)

Experimentally determined cholesterol flip-flop rates are often higher than predicted by MD. A 1981 study using cholesterol oxidase to monitor cholesterol flip-flop put an upper limit of 1 minute on flip-flop half-time (Backer and Dawidowicz, 1981). This value is consistent with more recent data from Leventis and Silvius (2001) who determined a 1-2 minute flip-flop half-time using ^3H labeled cholesterol partitioning studies. Other estimates of cholesterol flip-flop are on the order of seconds to nanoseconds (Steck et al., 2002; Bruckner et al., 2009; Hamilton, 2003; Endress et al., 2002). In contrast, a recent neutron scattering study of cholesterol flip-flop has yielded a half-time of 200 min, highlighting the need for further study regarding this subject (Garg et al., 2011).

4. Cholesterol’s Lateral Organization and Phospholipid Phase Behaviour

It is well known that cholesterol has a significant effect on the phase behaviour of phospholipid bilayers (Figure 5). Single component phospholipid bilayers exist, to a first approximation, in two different phases: the low temperature gel phase (L_β) and the high temperature liquid crystalline phase (L_α). The L_β phase is characterized by high segmental fatty acid chain ordering (i.e., lipid hydrocarbon chains are almost fully extended) and the phospholipids are well-ordered in the two-dimensional plane of the bilayer (Smith et al., 1988; Katsaras et al., 1992; Raghunathan and Katsaras, 1995; Heberle and Feigenson, 2011). The transition from L_β to the higher temperature L_α phase is a first order transition (Heberle and Feigenson, 2011), with the first direct observation of phase co-existence in a single component system made by Armstrong et al. (2012). Compared to the L_β phase, the L_α phase exhibits increased hydrocarbon chain disorder (trans/gauche) – causing the chains to sample a larger conformation space – experiences the loss of two-dimensional long range order, and the lipids undergo increased lateral diffusion (Ipsen et al., 1987; Heberle and Feigenson, 2011; Kučerka et al., 2011; Rubenstein et al., 1979). However, when cholesterol is introduced to these single component lipid systems, an entirely new phase is introduced, namely the so-called liquid ordered phase (L_o) – the liquid crystalline L_α phase is now termed the liquid disordered phase (L_d), although not entirely correct. The L_o is unique in that it has characteristics of both the L_β and L_α phases. For example, in the L_o phase the lipids do not form an extended two-dimensional hydrocarbon chain lattice in the plane of the bilayer plane, yet the lipid acyl chains exhibit a high degree of chain segmental ordering and undergo relatively fast lipid lateral diffusion (Heberle and Feigenson, 2011; Veatch and Keller, 2005; Kahya et al., 2003; Ipsen et al., 1987). It is this so-called cholesterol rich L_o phase that is believed to provide the platform for functional domains or lipid rafts.

The existence of lipid rafts in living systems has been contentious, with a plethora of work devoted to their study over the last 30 years (Simons and Toomre, 2000; Hancock, 2006; Lingwood and Simons, 2010; Ipsen et al., 1987). Lipid raft compositions have been developed using ternary (de Almeida et al., 2003; Feigenson and Buboltz, 2001) and quaternary lipid mixtures, and cholesterol (Konyakhina et al., 2013). Studies of these lipid mixtures, and the resultant phase diagrams, have allowed for the identification of different phases (i.e., L_o and

Figure 5: Venn diagram illustrating the properties shared between the liquid ordered (L_o) (contains cholesterol), gel (L_β) and liquid disordered (L_d) phases. Figure reproduced from Marquardt et al. (2015b)

L_d) as a function of cholesterol content. To digress for a moment, Armstrong et al. (2013), however, recently observed the existence of cholesterol induced lipid domains in a nominally single phase lipid bilayer (i.e., DMPC). Importantly, the dynamical properties imparted by cholesterol to this L_o phase were rather unique, in that the observed nanoscale dynamics amplified certain properties of the L_β and L_d phases (Meinhardt et al., 2013; Shlomovitz et al., 2014; Toppozini et al., 2014). Specifically, the nanoscopic cholesterol containing domains appeared softer than L_d bilayers but more ordered than even L_β bilayers (Armstrong et al., 2012). This amplification ability by cholesterol is thought to be an important driver in raft formation.

Heberle et al. (2013) studied the presence and size of nanoscopic domains in 60 nm diameter ULVs made up of a saturated phospholipid, a variable ratio of mono- and di-unsaturated phospholipids, and a constant mol% of cholesterol. As the amount of the di-unsaturated lipid was increased (i.e., mono-unsaturated lipid decreased), domain size and bilayer thickness mismatch between the L_d and L_o both increased. This observation of increasing thickness mismatch with increasing domain diameter is consistent with theories espousing line tension as the driving force behind domain formation (Heberle et al., 2013; Kuzmin et al.,

2005; Frolov et al., 2006). These experiments, carried out at constant temperature, also
lend insight as to how functional domains in biological membranes may be regulated, i.e.,
by altering the membrane’s chemical composition.

5. Concluding Remarks

Cholesterol is an important precursor for the biosynthesis of certain hormones and vitamins, and is an essential biomolecule of animal cell membranes. In this review we have highlighted some of the effects that cholesterol has on model membranes. In turn, the “altered” membranes then influence the energetics of protein conformational transitions due to interactions between the protein’s hydrophobic domains and the bilayers acyl chain region (Andersen and Koeppe, 2007). A better understanding of cholesterol’s significance in membranes will eventually help us elucidate some of the open structure-function questions currently debated in biology.

Acknowledgment

DM thanks Georg Pabst for his support. NK is supported through the VEGA grant 1/1224/12 and collaborative SR-JINR program under theme 04-4-1121-2015/2017, JK is supported through the Scientific User Facilities Division of the DOE Office of Basic Energy Sciences under US DOE Contract No. DE-AC05-00OR22725.

References

- Aittoniemi, J., Róg, T., Niemelä, P., Pasenkiewicz-Gierula, M., Karttunen, M., Vattulainen, I., 2006. Tilt: major factor in sterols’ ordering capability in membranes. *J. Phys. Chem. B.* 110 (51), 25562–25564.
- Andersen, O. S., Koeppe, R. E., 2007. Bilayer thickness and membrane protein function: an energetic perspective. *Annu. Rev. Biophys. Biomol. Struct.* 36, 107–130.
- Armstrong, C. L., Barrett, M. A., Toppozini, L., Kučerka, N., Yamani, Z., Katsaras, J., Fragneto, G., Rheinstaedter, M. C., 2012. Co-existence of gel and fluid lipid domains in single-component phospholipid membranes. *Soft Matter* 8 (17), 4687–4694.

387 Armstrong, C. L., Marquardt, D., Dies, H., Kučerka, N., Yamani, Z., Harroun, T. A.,
 388 Katsaras, J., Shi, A. C., Rheinstaedter, M. C., 2013. The observation of highly ordered
 389 domains in membranes with cholesterol. *PLoS One* 8 (6).

390 Azzi, A., 2007. Molecular mechanism of alpha-tocopherol action. *Free Radicals Biol. Med.*
 391 43 (1), 16–21.

392 Backer, J. M., Dawidowicz, E. A., 1981. Transmembrane movement of cholesterol in small
 393 unilamellar vesicles detected by cholesterol oxidase. *J. Biol. Chem.* 256 (2), 586–588.

394 Bartels, T., Lankalapalli, R. S., Bittman, R., Beyer, K., Brown, M. F., 2008. Raftlike mix-
 395 tures of sphingomyelin and cholesterol investigated by solid-state 2h nmr spectroscopy. *J.*
 396 *Am. Chem. Soc.* 130 (44), 14521–14532.

397 Bennett, W. F. D., MacCallum, J. L., Hinner, M. J., Marrink, S. J., Tieleman, D. P.,
 398 2009. Molecular view of cholesterol flip-flop and chemical potential in different membrane
 399 environments. *J. Am. Chem. Soc.* 131 (35), 12714–12720.

400 Brown, A. J., Galea, A. M., 2010. Cholesterol as an evolutionary response to living with
 401 oxygen. *Evolution (Hoboken, NJ, U. S.)* 64 (7), 2179–2183.

402 Brown, M. S., Goldstein, J. L., 1992. A receptor-mediated pathway for cholesterol homeosta-
 403 sis. In: Lindsten, J. (Ed.), *Nobel lectures: including presentation speeches and laureates’*
 404 *biographies. Nobel lecture. World Scientific, Singapore*, pp. 284–324.

405 Bruckner, R. J., Mansy, S. S., Ricardo, A., Mahadevan, L., Szostak, J. W., 2009. Flip-flop-
 406 induced relaxation of bending energy: Implications for membrane remodeling. *Biophys. J.*
 407 97 (12), 3113–3122.

408 Brzustowicz, Stillwell, W., Wassall, 1999. Molecular organization of cholesterol in polyunsat-
 409 urated phospholipid membranes: a solid state h-2 nmr investigation. *FEBS Lett.* 451 (2),
 410 197–202.

411 Brzustowicz, M. R., Cherezov, V., Caffrey, M., Stillwell, W., Wassall, S. R., 2002a. Molec-
 412 ular organization of cholesterol in polyunsaturated membranes: Microdomain formation.
 413 *Biophys. J.* 82 (1), 285–298.

414 Brzustowicz, M. R., Cherezov, V., Zerouga, M., Caffrey, M., Stillwell, W., Wassall, S. R.,
415 2002b. Controlling membrane cholesterol content. a role for polyunsaturated (docosahe-
416 aenoate) phospholipids. *Biochemistry* 41 (41), 12509–12519.

417 Bunge, A., Müller, P., Stöckl, M., Herrmann, A., Huster, D., 2008. Characterization of the
418 ternary mixture of sphingomyelin, popc, and cholesterol: support for an inhomogeneous
419 lipid distribution at high temperatures. *Biophys. J.* 94 (7), 2680–2690.

420 Chiu, S. W., Jakobsson, E., Mashl, R. J., Scott, H. L., 2002. Cholesterol-induced modifica-
421 tions in lipid bilayers: A simulation study. *Biophys. J.* 83 (4), 1842–1853.

422 Davis, J. H., 1983. The description of membrane lipid conformation, order and dynamics by
423 2h-nmr. *Biochim. Biophys. Acta, Rev. Biomembr.* 737 (1), 117–171.

424 de Almeida, R. F., Fedorov, A., Prieto, M., 2003. Sphin-
425 gomyelin/phosphatidylcholine/cholesterol phase diagram: Boundaries and composition of
426 lipid rafts. *Biophys. J.* 85 (4), 2406–2416.

427 Dufourc, E. J., Parish, E. J., Chitrakorn, S., Smith, I., 1984. Structural and dynamical details
428 of cholesterol lipid interaction as revealed by deuterium nmr. *Biochemistry* 23 (25), 6062–
429 6071.

430 Endress, E., Heller, H., Casalta, H., Brown, M. F., Bayerl, T. M., 2002. Anisotropic motion
431 and molecular dynamics of cholesterol, lanosterol, and ergosterol in lecithin bilayers studied
432 by quasi-elastic neutron scattering †. *Biochemistry* 41 (43), 13078–13086.

433 Feigenson, G. W., Buboltz, J. T., 2001. Ternary phase diagram of dipalmitoyl-pc/dilauroyl-
434 pc/cholesterol: nanoscopic domain formation driven by cholesterol. *Biophys. J.* 80 (6),
435 2775–2788–.

436 Franks, N. P., Lieb, W. R., 1979. The structure of lipid bilayers and the effects of general
437 anaesthetics. *J. Mol. Biol.* 133 (4), 469–500.

438 Frolov, V., Chizmadzhev, Y. A., Cohen, F. S., Zimmerberg, J., 2006. “entropic traps” in
439 the kinetics of phase separation in multicomponent membranes stabilize nanodomains.
440 *Biophys. J.* 91 (1), 189–205.

- 441 Gallová, J., Klacsová, M., Devínsky, F., Balgavý, P., 2015. Partial volumes of cholesterol
442 and monounsaturated diacylphosphatidylcholines in mixed bilayers. *Chem. Phys. Lipids*
443 190, 1–8.
- 444 Garg, S., Porcar, L., Woodka, A. C., Butler, P. D., Perez-Salas, U., 2011. Noninvasive neutron
445 scattering measurements reveal slower cholesterol transport in model lipid membranes.
446 *Biophys. J.* 101 (2), 370–377.
- 447 Gawrisch, K., Soubias, O., 2008. Structure and dynamics of polyunsaturated hydrocarbon
448 chains in lipid bilayers-significance for gpcr function. *Chem. Phys. Lipids* 153 (1), 64–75.
- 449 Giang, H., Schick, M., 2014. How cholesterol could be drawn to the cytoplasmic leaf of the
450 plasma membrane by phosphatidylethanolamine. *Biophys. J.* 107 (10), 2337–2344.
- 451 Gilbert, D. B., Tanford, C., Reynolds, J. A., 1975. Cholesterol in aqueous-solution - hy-
452 drophobicity and self association. *Biochemistry* 14 (2), 444–448.
- 453 Girotti, A. W., 2001. Photosensitized oxidation of membrane lipids: Reaction pathways,
454 cytotoxic effects, and cytoprotective mechanisms. *J. Photochem. Photobiol., B* 63 (1-3),
455 103–113.
- 456 Greenwood, A. I., Tristram-Nagle, S., Nagle, J. F., 2006. Partial molecular volumes of lipids
457 and cholesterol. *Chem. Phys. Lipids* 143 (1-2), 1–10.
- 458 Haberlan, M. E., Reynolds, J. A., 1973. Self-association of cholesterol in aqueous-solution.
459 *Proc. Natl. Acad. Sci. U. S. A.* 70 (8), 2313–2316.
- 460 Hamilton, J. A., 2003. Fast flip-flop of cholesterol and fatty acids in membranes: implications
461 for membrane transport proteins. *Curr. Opin. Lipidol.* 14 (3), 263–271.
- 462 Hancock, J. F., 2006. Lipid rafts: contentious only from simplistic standpoints. *Nat. Rev.*
463 *Mol. Cell Biol.* 7 (6), 456–462.
- 464 Harroun, T., Wignall, G., Katsaras, J., 2006a. Neutron scattering for biology: 1-18. In: Jörg
465 Fitter, Thomas Gutberlet, John Katsaras (Eds.), *Neutron Scattering in Biology Tech-
466 niques and Applications*. Springer, pp. 1–18.

467 Harroun, T. A., Katsaras, J., Wassall, S. R., 2006b. Cholesterol hydrozyl group is found to
 468 reside in the center of a polyunsaturated lipid membrane. *Biochemistry* 45, 1227–1233.

469 Harroun, T. A., Katsaras, J., Wassall, S. R., 2008. Cholesterol is found to reside in the center
 470 of a polyunsaturated lipid membrane. *Biochemistry* 47 (27), 7090–7096.

471 Heberle, F. A., Feigenson, G. W., 2011. Phase separation in lipid membranes. *Cold Spring*
 472 *Harbor Perspect. Biol.* 3 (4).

473 Heberle, F. A., Petruzielo, R. S., Pan, J., Drazba, P., Kučerka, N., Standaert, R. F., Feigen-
 474 son, G. W., Katsaras, J., 2013. Bilayer thickness mismatch controls domain size in model
 475 membranes. *J. Am. Chem. Soc.* 135 (18), 6853–6859.

476 Hofsäb, C., Lindahl, E., Edholm, O., 2003. Molecular dynamics simulations of phospholipid
 477 bilayers with cholesterol. *Biophys. J.* 84 (4), 2192–2206.

478 Ipsen, J. H., Karlstrom, G., Mouritsen, O. G., Wennerstrom, H., Zuckermann, M. J., 1987.
 479 Phase equilibria in phosphatidylcholine-cholesterol system. *Biochim. Biophys. Acta* 905,
 480 162–172.

481 Jackman, C. S., Davis, P. J., Morrow, M. R., Keough, K. M. W., 1999. Effect of cholesterol on
 482 the chain-ordering transition of 1-palmitoyl-2-arachidonoyl phosphatidylcholine 103 (42),
 483 8830–8836.

484 Jo, S., Rui, H., Lim, J. B., Klauda, J. B., Im, W., 2010. Cholesterol flip-flop: insights from
 485 free energy simulation studies. *J. Phys. Chem. B.* 114 (42), 13342–13348.

486 Kahya, N., Scherfeld, D., Bacia, K., Poolman, B., Schwille, P., 2003. Probing lipid mobility of
 487 raft-exhibiting model membranes by fluorescence correlation spectroscopy. *J. Biol. Chem.*
 488 278 (30), 28109–28115.

489 Katsaras, J., Yang, D. S.-C., Epand, R. M., 1992. Fatty acid chain tilt angles and directions
 490 in dipalmitoyl phosphatidylcholine bilayers. *Biophys. J.* 63, 1170.

491 Kessel, A., Ben-Tal, N., May, S., 2001. Interactions of cholesterol with lipid bilayers: The
 492 preferred configuration and fluctuations. *Biophys. J.* 81 (2), 643–658.

493 Khelashvili, G., Harries, D., 2013. How cholesterol tilt modulates the mechanical properties
494 of saturated and unsaturated lipid membranes. *J. Phys. Chem. B*.

495 Konyakhina, T. M., Wu, J., Mastroianni, J. D., Heberle, F. A., Feigenson, G. W., 2013.
496 Phase diagram of a 4-component lipid mixture: Dspc/dopc/popc/chol. *Biochim. Biophys.*
497 *Acta* 1828 (9), 2204–2214.

498 Kristiana, I., Luu, W., Stevenson, J., Cartland, S., Jessup, W., Belani, J. D., Rychnovsky,
499 S. D., Brown, A. J., 2012. Cholesterol through the looking glass: ability of its enantiomer
500 also to elicit homeostatic responses. *J. Biol. Chem.* 287 (40), 33897–33904.

501 Kučerka, N., Marquardt, D., Harroun, T. A., Nieh, M.-P., Wassall, S. R., Jong, D. H. d.,
502 Schäfer, L. V., Marrink, S. J., Katsaras, J., 2010. Cholesterol in bilayers with pufa chains:
503 doping with dmpe or popc results in sterol reorientation and membrane-domain formation.
504 *Biochemistry* 49 (35), 7485–7493.

505 Kučerka, N., Nieh, M.-P., Katsaras, J., 2011. Fluid phase lipid areas and bilayer thicknesses
506 of commonly used phosphatidylcholines as a function of temperature. *Biochim. Biophys.*
507 *Acta* 1808, 2761–2771.

508 Kučerka, N., Nieh, M.-P., Pencer, J., Sachs, J. N., Katsaras, J., 2009. What determines the
509 thickness of a biological membrane. *Gen. Physiol. Biophys.* 28 (2), 117–125.

510 Kučerka, N., Pencer, J., Nieh, M.-P., Katsaras, J., 2007. Influence of cholesterol on the bilayer
511 properties of monounsaturated phosphatidylcholine unilamellar vesicles. *Eur. Phys. J. E:*
512 *Soft Matter Biol. Phys.* 23 (3), 247–254.

513 Kučerka, N., Perlmutter, J. D., Pan, J., Tristram-Nagle, S., Katsaras, J., Sachs, J. N.,
514 2008. The effect of cholesterol on short- and long-chain monounsaturated lipid bilayers as
515 determined by molecular dynamics simulations and x-ray scattering. *Biophys. J.* 95 (6),
516 2792–2805.

517 Kuzmin, P. I., Akimov, S. A., Chizmadzhev, Y. A., Zimmerberg, J., Cohen, F. S., 2005.
518 Line tension and interaction energies of membrane rafts calculated from lipid splay and
519 tilt. *Biophys. J.* 88 (2), 1120–1133.

520 Lafleur, M., Bloom, M., Cullis, P. R., 1990. Modulation of the orientational order profile of
521 the lipid acyl chain in the l alpha phase. *Eur. Biophys. J.* 19 (2), 55–62.

522 Lange, Y., Ramos, B. V., 1983. Analysis of the distribution of cholesterol in the intact cell.
523 *J. Biol. Chem.* 258 (24), 5130–5134.

524 Leng, X., Kinnun, J. J., Marquardt, D., Ghefli, M., Kučerka, N., Katsaras, J., Atkinson, J.,
525 Harroun, T. A., Feller, S. E., Wassall, S. R., 2015. α -tocopherol is well designed to protect
526 polyunsaturated phospholipids: Md simulations 109 (8), 1608–1618.

527 Leonard, A., Escrive, C., Laguerre, M., Pebay-Peyroula, E., Neri, W., Pott, T., Katsaras,
528 J., Dufourc, E. J., 2001. Location of cholesterol in dmpc membranes. a comparative study
529 by neutron diffraction and molecular mechanics simulation. *Langmuir* 17 (6), 2019–2030.

530 Leventis, R., Silvius, J. R., 2001. Use of cyclodextrins to monitor transbilayer movement and
531 differential lipid affinities of cholesterol. *Biophys. J.* 81 (4), 2257–2267.

532 Lingwood, D., Simons, K., 2010. Lipid rafts as a membrane-organizing principle. *Science*
533 327 (5961), 46–50.

534 Marquardt, D., Geier, B., Pabst, G., 2015a. Asymmetric lipid membranes: towards more
535 realistic model systems. *Membranes* 5 (2), 180–196.

536 Marquardt, D., Harroun, T., 2014. Locations of small biomolecules in model membranes. In:
537 Pabst, G., Kučerka, N., Nieh, M.-P., Katsaras, J. (Eds.), *Liposomes, Lipid Bilayers and*
538 *Model Membranes*. CRC Press, pp. 199–216.

539 Marquardt, D., Heberle, F. A., Nickels, J. D., Pabst, G., Katsaras, J., 2015b. On scattered
540 waves and lipid domains: Detecting membrane rafts with x-rays and neutrons. *Soft Matter*
541 11 (47), 9055–9072.

542 Marquardt, D., Kučerka, N., Katsaras, J., Harroun, T. A., 2015c. α -tocopherol’s location in
543 membranes is not affected by their composition. *Langmuir* 31 (15), 4464–4472.

544 Marquardt, D., van Oosten, B., Heberle, F. A., Kučerka, N., Wassall, S., Standaert, R.,
 545 Katsaras, J., Harroun, T. A., 2015d. Hydrocarbon thickness dictates cholesterol's location,
 546 orientation and motion in a phospholipid bilayer 108 (2), 86a.

547 Marquardt, D., Williams, J. A., Kinnun, J. J., Kučerka, N., Atkinson, J., Wassall, S. R., Kat-
 548 saras, J., Harroun, T. A., 2014. Dimyristoyl phosphatidylcholine: a remarkable exception
 549 to α -tocopherol's membrane presence. *J. Am. Chem. Soc.* 136 (1), 203–210.

550 Marquardt, D., Williams, J. A., Kučerka, N., Atkinson, J., Wassall, S. R., Katsaras, J.,
 551 Harroun, T. A., 2013. Tocopherol activity correlates with its location in a membrane: A
 552 new perspective on the antioxidant vitamin e. *J. Am. Chem. Soc.* 135 (20), 7523–7533.

553 Marsan, M. P., Muller, I., Ramos, C., Rodriguez, F., Dufourc, E. J., Czaplicki, J., Milon, A.,
 554 1999. Cholesterol orientation and dynamics in dimyristoylphosphatidylcholine bilayers: A
 555 solid state deuterium nmr analysis. *Biophys. J.* 76 (1, Part 1), 351–359.

556 Meinhardt, S., Vink, R. L. C., Schmid, F., 2013. Monolayer curvature stabilizes nanoscale
 557 raft domains in mixed lipid bilayers. *Proc. Natl. Acad. Sci. U.S.A.* 110 (12), 4476–4481.

558 Mihailescu, M., Soubias, O., Worcester, D., White, S. H., Gawrisch, K., 2011. Structure and
 559 dynamics of cholesterol-containing polyunsaturated lipid membranes studied by neutron
 560 diffraction and nmr. *J. Membr. Biol.* 239 (1-2), 63–71.

561 Miyoshi, T., Lönnfors, M., Peter Slotte, J., Kato, S., 2014. A detailed analysis of partial
 562 molecular volumes in dppc/cholesterol binary bilayers. *Biochim. Biophys. Acta* 1838 (12),
 563 3069–3077.

564 Mouritsen, O. G., 2005. Life - as a matter of fat: The emerging science of lipidomics. Frontiers
 565 collection. Springer, Berlin.

566 Murari, R., Murari, M. P., Baumann, W. J., 1986. Sterol orientations in phosphatidylcholine
 567 liposomes as determined by deuterium nmr. *Biochemistry* 25 (5), 1062–1067.

568 Myant, N. B., 1981. The Biology of Cholesterol and Related Steroids. Elsevier Science,
 569 Burlington.

570 Nickels, J. D., Smith, J. C., Cheng, X., 2015. Lateral organization, bilayer asymmetry, and
571 inter-leaflet coupling of biological membranes. *Chem. Phys. Lipids*.

572 Niu, S. L., Litman, B. J., 2002. Determination of membrane cholesterol partition coeffi-
573 cient using a lipid vesicle-cyclodextrin binary system: Effect of phospholipid acyl chain
574 unsaturation and headgroup composition. *Biophys. J.* 83 (6), 3408–3415.

575 Oldfield, E., Meadows, M., Rice, D., Jacobs, R., 1978. Spectroscopic studies of specifically
576 deuterium labeled membrane systems. nuclear magnetic resonance investigation of the
577 effects of cholesterol in model systems. *Biochemistry* 17 (14), 2727–2740.

578 Pan, J., Cheng, X., Heberle, F. A., Mostofian, B., Kučerka, N., Drazba, P., Katsaras, J.,
579 2012. Interactions between ether phospholipids and cholesterol as determined by scattering
580 and molecular dynamics simulations. *J. Phys. Chem. B.* 116 (51), 14829–14838.

581 Pan, J., Mills, T. T., Tristram-Nagle, S., Nagle, J. F., 2008. Cholesterol perturbs lipid
582 bilayers nonuniversally. *Phys. Rev. Lett.* 100 (19).

583 Pandey, B. N., Mishra, K. P., 1999. Radiation induced oxidative damage modification by
584 cholesterol in liposomal membrane. *Radiat. Phys. Chem.* 54 (5), 481–489.

585 Papanikolaou, A., Papafotika, A., Murphy, C., Papamarcaki, T., Tsolas, O., Drab, M.,
586 Kurzchalia, T. V., Kasper, M., Christoforidis, S., 2005. Cholesterol-dependent lipid as-
587 semblies regulate the activity of the ecto-nucleotidase cd39. *J. Biol. Chem.* 280 (28),
588 26406–26414.

589 Parasassi, T., Giusti, A. M., Raimondi, M., Ravagnan, G., Sapor, O., Gratton, E., 1995.
590 Cholesterol protects the phospholipid bilayer from oxidative damage. *Free Radicals Biol.*
591 *Med.* 19 (4), 511–516.

592 Petrie, R. J., Schnetkamp, P. P. M., Patel, K. D., Awasthi-Kalia, M., Deans, J. P., 2000.
593 Transient translocation of the b cell receptor and src homology 2 domain-containing inos-
594 itol phosphatase to lipid rafts: Evidence toward a role in calcium regulation. *J. Immunol.*
595 165 (3), 1220–1227.

- 596 Raghunathan, V., Katsaras, J., 1995. Structure of the l(c') phase in a hydrated lipid multil-
597 amellar system. *Phys. Rev. Lett.* 74 (22), 4456–4459.
- 598 Róg, T., Pasenkiewicz-Gierula, M., 2001. Cholesterol effects on the phospholipid conden-
599 sation and packing in the bilayer: A molecular simulation study. *FEBS Lett.* 502 (1-2),
600 68–71.
- 601 Róg, T., Pasenkiewicz-Gierula, M., 2003. Effects of epicholesterol on the phosphatidylcholine
602 bilayer: A molecular simulation study. *Biophys. J.* 84 (3), 1818–1826.
- 603 Rubenstein, J. L., Smith, B. A., McConnell, H. M., 1979. Lateral diffusion in binary mixtures
604 of cholesterol and phosphatidylcholines. *Proc. Natl. Acad. Sci. U.S.A.* 76 (1), 15–18.
- 605 Seelig, J., 1977. Deuterium magnetic resonance: theory and application to lipid membranes.
606 *Q. Rev. Biophys.* 10 (3), 353–418.
- 607 Seelig, J., Waespe-Sarcevic, N., 1978. Molecular order in cis and trans unsaturated phospho-
608 lipid bilayers. *Biochemistry* 17 (16), 3310–3315.
- 609 Shaikh, Cherezov, V., Caffrey, M., Soni, S. P., LoCascio, D., Stillwell, W., Wassall, 2006.
610 Molecular organization of cholesterol in unsaturated phosphatidylethanolamines: X-ray
611 diffraction and solid state h-2 nmr reveal differences with phosphatidylcholines. *J. Am.*
612 *Chem. Soc.* 128 (16), 5375–5383.
- 613 Shaikh, S. R., Kinnun, J. J., Leng, X., Williams, J. A., Wassall, S. R., 2015. How polyunsat-
614 urated fatty acids modify molecular organization in membranes: insight from nmr studies
615 of model systems. *Biochim. Biophys. Acta* 1848 (1 Pt B), 211–219.
- 616 Shlomovitz, R., Maibaum, L., Schick, M., 2014. Macroscopic phase separation, modulated
617 phases, and microemulsions: A unified picture of rafts. *Biophysical Journal* 106 (9), 1979–
618 1985.
- 619 Shrivastava, S., Haldar, S., Gimpl, G., Chattopadhyay, A., 2009. Orientation and dynamics
620 of a novel fluorescent cholesterol analogue in membranes of varying phase 113 (13), 4475–
621 4481.

622 Simons, K., Toomre, D., 2000. *Nat. Rev. Mol. Cell Biol.* 1 (1), 31–39.

623 Smith, G. S., Sirota, E. B., Safinya, C. R., Clark, N. A., 1988. Structure of the l_β phase in
624 a hydrated phosphatidylcholine multimembrane. *Phys. Rev. Lett.* 60, 813–816.

625 Smith, L. L., 1991. Another cholesterol hypothesis: Cholesterol as antioxidant. *Free Radicals*
626 *Biol. Med.* 11 (1), 47–61.

627 Steck, T. L., Ye, J., Lange, Y., 2002. Probing red cell membrane cholesterol movement with
628 cyclodextrin. *Biophys. J.* 83 (4), 2118–2125.

629 Stillwell, W., 2013. Chronology of membrane studies. In: Stillwell, W. (Ed.), *An Introduction*
630 *to Biological Membranes*. Elsevier, San Diego, pp. 357–360.

631 Subczynski, W. K., Widomska, J., Feix, J. B., 2009. Physical properties of lipid bilayers from
632 epr spin labeling and their influence on chemical reactions in a membrane environment.
633 *Free Radicals Biol. Med.* 46 (6), 707–718.

634 Tieleman, D. P., Marrink, S.-J., 2006. Lipids out of equilibrium: energetics of desorption
635 and pore mediated flip-flop. *J. Am. Chem. Soc.* 128 (38), 12462–12467.

636 Toppozini, L., Meinhardt, S., Armstrong, C. L., Yamani, Z., Kučerka, N., Schmid, F.,
637 Rheinstädter, M. C., 2014. Structure of cholesterol in lipid rafts. *Physical Review Letters*
638 113 (22).

639 Traber, M. G., Atkinson, J., 2007. Vitamin e, antioxidant and nothing more. *Free Radical*
640 *Biol. Med.* 43 (1), 4–15.

641 van Meer, G., 2011. Dynamic transbilayer lipid asymmetry. *Cold Spring Harbor Perspect.*
642 *Biol.* 327, 3:a004671.

643 Veatch, S. L., Keller, S. L., 2005. Seeing spots: Complex phase behavior in simple mem-
644 branes. *Biochim. Biophys. Acta, Mol. Cell Res.* 1746 (3), 172–185.

645 Vist, M. R., Davis, J. H., 1990. Phase equilibria of choles-
646 terol/dipalmitoylphosphatidylcholine mixtures: Deuterium nuclear magnetic resonance
647 and differential scanning calorimetry. *Biochemistry* 29 (2), 451–464.

648 Wassall, S. R., Stillwell, W., 2008. Docosahexaenoic acid domains: the ultimate non-raft
649 membrane domain. *Chem. Phys. Lipids* 153 (1), 57–63.

650 Westover, E. J., Covey, D. F., Brockman, H. L., Brown, R. E., Pike, L. J., 2003. Cholesterol
651 depletion results in site-specific increases in epidermal growth factor receptor phosphory-
652 lation due to membrane level effects. studies with cholesterol enantiomers. *J. Biol. Chem.*
653 278 (51), 51125–51133.

654 Williams, J., Wassall, C. D., Kemple, M. D., Wassall, S. R., 2013. Dependence of cholesterol-
655 phospholipid affinity on degree of acyl chain unsaturation as determined by epr. *Biophys.*
656 *J.* 104 (2), 588a.

657 Wood, W. G., Igbavboa, U., Müller, W. E., Eckert, G. P., 2011. Cholesterol asymmetry in
658 synaptic plasma membranes. *J. Neurochem.* 116 (5), 684–689.

659 Worcester, D. L., Franks, N. P., 1976. Structural analysis of hydrated egg lecithin and
660 cholesterol bilayers ii. neutron diffraction. *J. Mol. Biol.* 100 (3), 359–378.

661 Yasuda, T., Kinoshita, M., Murata, M., Matsumori, N., 2014. Detailed comparison of deu-
662 terium quadrupole profiles between sphingomyelin and phosphatidylcholine bilayers. *Bio-*
663 *phys. J.* 106 (3), 631–638.

664 Yeagle, P. L., 1993. The biophysics and cell biology of cholesterol: an hypothesis for the
665 essential role of cholesterol in mammalian cells. In: Finegold, L. (Ed.), *Cholesterol in*
666 *membrane models*. CRC Press, Boca Raton, Fla., pp. 1–12.

667 Yesylevskyy, S. O., Demchenko, A. P., 2012. How cholesterol is distributed between mono-
668 layers in asymmetric lipid membranes. *Eur. Biophys. J.* 41 (12), 1043–1054.

669 Zhang, Z., Lu, L., Berkowitz, M. L., 2008. Energetics of cholesterol transfer between lipid
670 bilayers. *J. Phys. Chem. B.* 112 (12), 3807–3811.