On the Controlling Mechanism of the Upper Turnover States in the NTC Regime

Weiqli Ji^a, Peng Zhao^{b,c,*,†}, Tanjin He^d, Xin He^{a,d}, Aamir Farooq^e, Chung K. Law^{a,c,*,‡}

^a Center for Combustion Energy, Tsinghua University, Beijing 100084, China

^b Department of Mechanical Engineering, Oakland University, Rochester, MI, 48309, USA

^c Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, NJ 08544, USA

^d State Key Laboratory of Automotive Safety and Energy, Tsinghua University, Beijing 100084, China

^e Clean Combustion Research Center, Division of Physical Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal 23955-6900, Saudi Arabia

*Corresponding authors

[†]Email: pengzhao@oakland.edu

[‡]Email: cklaw@princeton.edu

Abstract

Using *n*-butane, *n*-heptane and *iso*-octane as representative fuels exhibiting NTC (negative temperature coefficient) behavior, comprehensive computational studies with detailed mechanisms and theoretical analysis were performed to investigate the upper stationary point, denoted as turnover states, on the NTC curve near the higher temperature regime, where the ignition delay τ exhibits a local maximum. It is found that the global behavior of the turnover states exhibits distinctive thermodynamic and kinetic characteristics under different pressures, in that the ignition delay at the turnover states shows an Arrhenius dependence on the temperature T and an approximate inverse quadratic power law dependence on the pressure P . These global behaviors imply that the temperature and pressure of the turnover states are not independent and can be correlated by an Arrhenius dependence, as $\ln P \propto 1/T$. Further theoretical analyses demonstrate that such turnover states result from the competition between the low-temperature chain branching reactions and the decomposition of the intermediate species, and therefore correspond to a critical value, α , of the ratio of OH production from low-temperature chemistry. In addition, the ignition delay at the turnover state can be well correlated by the analytical expression derived by Peters *et al.*, with the further demonstration that the pressure dependence of the turnover ignition delay mainly result from the H_2O_2 decomposition reaction. Comparison of the present results with the literature experimental data of *n*-heptane ignition delay time shows very good agreement.

Key Words: NTC; Turnover; Auto-ignition; Chain branching ratio; Low temperature chemistry

1. Introduction

The phenomenon of negative temperature coefficient (NTC) is an essential feature of the oxidation kinetics of large hydrocarbons, and is closely related to such practically important phenomena and processes as low-temperature ignition [1], cool flames [2, 3], flame stabilization [4], fuel processing [5], engine knock [6] and the development of HCCI (homogenous charge compression ignition) engines [7]. It refers to the effect that, for a given pressure and in an initial temperature regime much lower than the adiabatic flame temperature of the reacting mixture, the ignition delay could increase with increasing initial temperature. The NTC behavior in auto-ignition has been extensively observed and studied in homogenous systems such as the shock tube [8], jet-stirred reactor [9], flow reactor [10] and rapid compression machine (RCM) [11].

It is widely accepted that the NTC behavior is due to the peroxy-chemistry pathways in the low-to-intermediate temperature regimes, as discussed in [12-14]. As an example, Fig. 1 shows a schematic reaction path for the oxidation of large alkanes in such regimes. Specifically, fuel consumption occurs via H-atom abstraction from the fuel molecule to form the alkyl radical, R. Since β scission is slow at low temperatures, O_2 addition to the R radical is dominant, forming the RO_2 radical. Then isomerization of RO_2 forms the hydroperoxy-alkyl radical (QOOH). Further O_2 addition to QOOH forms a transition-ring complex (O_2QOOH) that isomerizes and decomposes to form ketohydroperoxide (KET) and the hydroxyl radical (OH) (red arrows in Fig. 1). The KET then decomposes to carbonyl, aldehyde and generates another OH radical, leading to the key chain branching of low-temperature chemistry. However, with increasing temperature, all the oxygen addition reactions shift towards the decomposition side, and as such the extent of low-T chain

1 branching is reduced. Meanwhile, at higher, intermediate temperatures ($800\text{K} < T < 900\text{K}$, 20
2
3 atm for *n*-heptane), dissociation of the low-temperature intermediates (*e.g.*, RO_2 and QOOH)
4
5 and the R radical through β scission gradually becomes important and forms different
6
7 products, including alkenes, HO_2 and other alkyl radicals (blue arrows in Fig. 1), which
8
9 further compete with the low-temperature chain branching pathway. The backward shift of
10
11 the low-temperature reaction channel, together with the favored β scission at intermediate
12
13 temperatures, results in globally reduced chemical reactivity at moderate temperatures in the
14
15 NTC regime.
16
17
18
19
20
21

22
23 At intermediate-to-high initial temperatures ($900\text{K} < T < 1200\text{K}$, 20 atm for *n*-heptane),
24
25 the major chain branching that drives the system runaway is hydrogen peroxide (H_2O_2)
26
27 decomposition [15] forming two OH radicals, with its rate constant increasing with
28
29 temperature. Consequently, the ignition delay starts to decrease again with increasing
30
31 temperature beyond a threshold value, termed the “turnover temperature”, corresponding to
32
33 the local maximum of the ignition delay time.
34
35
36
37
38

39 In this work, the characteristics of the turnover states including the thermodynamic and
40
41 chemical kinetic behaviors are studied. The motivations are three-fold: First, since the
42
43 turnover state also corresponds to the intermediate-to-high temperature boundary of the NTC
44
45 regime, identification of the mechanism controlling it will lead to new insights on the NTC
46
47 behavior, and can bring about significant convenience in the description of auto-ignition.
48
49 Second, sensitivity of the ignition delay to temperature variation is one of the key factors
50
51 affecting the flame propagation mode in HCCI engines [16] and relevant RCM experiments
52
53 [17, 18], deeper understanding of the underlying mechanism of the temperature-insensitive
54
55
56
57
58
59
60
61
62
63
64
65

1 nature of the ignition delay at the turnover state could be helpful in interpreting the various
2
3 combustion modes under HCCI conditions. Third, correlations of the ignition delay time as a
4
5 function of initial pressure and temperature usually show large deviations near the turnover
6
7 temperature [8, 19]. Thus, developing global scaling for the turnover temperature can help to
8
9 reduce the uncertainties and provide an accurate reference for the selection of experimental
10
11 data at the relevant experimental conditions.
12
13
14
15

16
17 The present study is computational. The fuels investigated are *n*-butane, *n*-heptane and
18
19 *iso*-octane, which represent typical short and long straight-chain alkanes as well as branched
20
21 alkanes. The mechanisms used are the NUIG *n*-butane mechanism [20], the LLNL *n*-heptane
22
23 mechanism version 3.1 [12], and the LLNL *iso*-octane mechanism version 3 [21],
24
25 respectively. The global behaviors of the ignition delay at turnover temperatures under
26
27 different pressures are investigated for all of the three fuels. It is noted that although the
28
29 pressure dependence of the rate constants of the oxygen addition reactions, alkylperoxy
30
31 radical (RO_2) dissociation and isomerization reactions, are not considered, it has been
32
33 recently demonstrated in [22, 23] that the high-pressure limits for these reactions are
34
35 sufficient to describe the low-temperature chemistry at the conditions of this study. The
36
37 mechanism for *n*-heptane updated by Karwat *et al.* [24], based on the rate rules from Villano
38
39 *et al.* [22, 23], is also referred to show that the behavior demonstrated in this study is
40
41 mechanism independent and with general implications on the low-temperature kinetics.
42
43
44
45
46
47
48
49
50
51
52

53 In the following sections, we shall first identify the global system behavior at the
54
55 turnover states under different pressures. We shall then analyze the coupling between
56
57 pressure and the turnover temperature, as well as the power-law dependence on the pressure
58
59
60
61
62
63
64
65

for the ignition delay at the turnover states. Finally, the characteristics of the turnover states are validated against available ignition delay data in the literature, demonstrating good agreement.

2. Global behavior of the turnover states

Figure 2 shows the ignition delay as a function of temperature for a homogeneous, stoichiometric mixture of *n*-heptane and air under 2-60 atm pressure, using detailed mechanism [12]. It is seen that for each pressure, the total ignition delay, τ , first decreases and then increases with increasing initial temperature, T , in the NTC regime, and finally decreases again when the temperature is above the turnover temperature, T_m , which, marked by symbols, is seen to shift to higher values as pressure is increased.

We next note that the ignition delay, τ_m , at the turnover state exhibits unique dependences on the corresponding temperature, T_m , and the system pressure, P . Specifically, its linear variation with T_m in the plot of Fig. 2 suggests an Arrhenius correlation, namely $\ln \tau_m \propto 1/T_m$, while Fig. 3 shows it also varies nearly quadratically with the system pressure, as $\tau_m \propto P^{-2}$. These results then suggest that T_m and P can be related by $\ln P \propto 1/T_m$, as shown in Fig. 4, yielding a *thermodynamic* correlation between the turnover temperature and the system pressure. It is further noted that the data in Figs. 3 and 4 include those for $\phi = 0.5, 1$ and 2, demonstrating that these observations are also insensitive to the equivalence ratio. We shall demonstrate in the following the underlying kinetic mechanism affecting such a behavior.

3. Chemical Kinetic Analysis on the Turnover States

3.1 Low-Temperature Chain Branching Ratio at the Turnover State

Figure 5 shows the evolution of the intermediate species at the turnover states, using stoichiometric *n*-butane/air mixtures under 20 atm for demonstration because less isomers are involved in its oxidation process. It is seen that many radicals of interest, largely remain quasi-steady for an extended period between the starting chain initiation and the final ignition runaway. Take OH as an example, the OH consumption rate is about $10^{-5} \sim 10^{-4}$ mole/cm³s as shown in Fig. 6, while the net OH production rate at the same condition is below 10^{-9} mole/cm³s, four orders of magnitude lower. As such, the OH mole fraction is kept at a low level (less than 10^{-8}) before ignition as shown in Fig. 5 and could be assumed to be in quasi-steady in the plateau period. Therefore, the total production and consumption rates of OH should approximately balance. Figure 6 then plots the rates of the important reactions consuming OH and the fuel molecule, showing that OH is mainly consumed by H-abstraction from the fuel molecule (nearly 90% of OH consumption at the instant of 75% τ_m in Fig. 6a), while fuel consumption is also mainly through the H-abstraction by OH (nearly 90% of fuel consumption at the instant of 75% τ_m by the four reactions in Fig. 6b). Consequently, the OH production rate can be estimated by the consumption rate of the fuel molecule. We shall demonstrate in the following how these observations identify the major kinetic characteristics at the turnover states.

The controlling reactions for the turnover temperature were identified by performing sensitivity analysis, with the sensitivity coefficient calculated by perturbing the A-factor of each elemental reaction rate by 30%, following the definition:

$$S = \frac{d \ln(T)}{d \ln(k)} = \frac{\ln(T_{1.3k}) - \ln(T_k)}{\ln(1.3k) - \ln(k)} \quad (1)$$

A step size of 0.1 K was used to search the turnover temperature, and the perturbation of 30% was selected to allow sufficient change of the turnover temperature for accurate evaluation. We shall only present the sensitivity for *n*-butane as the mechanisms for *n*-heptane and *iso*-octane are based on similar reaction classes and rate rules.

Figure 7 shows the sensitivity for the stoichiometric *n*-butane/air mixture at 2 and 10 atm. The most sensitive reactions are shown to be the chain termination reactions from the β scission of RO₂ and QOOH, as well as the isomerization from RO₂ to QOOH and the subsequent chain branching and propagation reactions. However, it is seen that the turnover temperature is not sensitive to the HO₂ chain termination reaction and the H₂O₂ chain branching reaction. Consequently, the turnover state is mainly controlled by the competition between the β scission of low-temperature intermediates as chain termination reactions producing the stable intermediates and HO₂, and the low-temperature chain branching/propagation reactions producing OH. It should be noted that the turnover temperature is insensitive to the perturbation of the rate of oxygen addition to QOOH reaction as long as its equilibrium constant is not perturbed. Such a behavior is expected as the oxygen addition to QOOH is in the partial equilibrium state, and the low-T chain branching is only affected by the equilibrium constant. Therefore, the result by perturbing the oxygen addition to QOOH using reversible reactions is not presented in Fig. 7, which otherwise might cause misleading and underestimate its role at the turnover states.

We note in passing the worthwhile studies on such competitions in the first-stage ignition controlled by the low-temperature chemistry. Specifically, Peters *et al.* [25] defined a ratio

between the $R + O_2 \rightleftharpoons RO_2$ with subsequent isomerization $RO_2 \rightleftharpoons QOOH$ and the β scission of R to determine the crossover temperature, at which the concentration of KET reaches its peak value and the first-stage ignition occurs. Merchant *et al.* [26] used a similar concept to study the first-stage ignition of propane, and included the HO_2 chemistry, namely $RO_2 \rightarrow$ Alkene + HO_2 , which was not included in [25, 27]. However, these ratios were derived from a reduced mechanism, and only specific isomers of R, RO_2 and QOOH were taken into account. Thus, these definitions are applicable only for limited conditions and for certain mechanisms, in the context of two-stage ignition. However, for the concept of the turnover state of current interest, sensitivity analysis has shown that the turnover temperature is also sensitive to the decomposition of QOOH to produce OH. Consequently, we shall define a ratio of the OH production rate, instead of the KET production rate, to the total production rate of R, as a criterion to evaluate the net contribution from the low-temperature chain branching:

$$\begin{aligned}\alpha &= \frac{\text{OH production rate from low-T chain-branching}}{\text{total production rate of R}} \\ &= \frac{\text{OH production rate from low-T chain-branching}}{\text{total rate of H-abstraction from RH}}\end{aligned}$$

The ratio α can be readily determined from the detailed mechanism including all isomers of R, RO_2 , and QOOH. In particular, we have followed the reaction flux from RH to OH, as shown in Fig. 8, and have considered the effect from β scission at each step, defined by the following ratios:

$$\begin{aligned}\eta_i &= \frac{\text{rate of } RH \rightarrow i\text{-R}}{\text{total rates of } RH \rightarrow R} \approx \frac{\text{rate of } RH+OH \rightarrow i\text{-R}+H_2O}{\text{total rates of } RH+OH \rightarrow R+H_2O} \\ \beta_i &= \frac{\text{net rate of } i\text{-R} \rightarrow i\text{-RO}_2}{\text{total production rate of } i\text{-R}}\end{aligned}$$

$$\gamma_{i,j} = \frac{\text{net rate of } i\text{-RO}_2 \rightarrow i,j\text{-QOOH}}{\text{total production rate of } i\text{-RO}_2}$$

$$\theta_{i,j} = \frac{\text{net rate of } i,j\text{-QOOH} \rightarrow \text{OH}}{\text{total production rate of } i,j\text{-QOOH}}$$

The subscript or prefix i indicates the index of the isomer of R, and those of i, j indicate the index of the isomer of QOOH. Thus, α is determined by

$$\alpha = \sum_i \eta_i \beta_i \sum_j \gamma_{i,j} \theta_{i,j} \quad (2)$$

In the Appendix, we present a systematic determination of α by first deriving it using a simplified skeletal mechanism with major reaction steps involving only one isomer, shown in Fig. 1 and Table 1. We shall then extend the analytical expressions to detailed kinetics involving all of the isomers.

We shall now show that the turnover state can be defined well by a constant value of α under different pressures. Following the definition of Eq. (2), the computed α as a function of temperature under various pressures for the three fuels are shown in Figs. 9 a, b and c, respectively. As expected, α decreases with increasing temperature, and eventually to zero at high enough temperatures. This shows the progressively decreasing significance of the low-temperature chain branching reactions with increasing temperature. Similar to the shift of NTC, the α curves shift to higher temperature with increasing pressure.

Based on α , we define a second ratio, α_{RO_2} , which corresponds to the ideal case where all R radicals participate in the oxygen addition reaction without β scission, *i.e.*, $\beta_i = 1$:

$$\alpha_{\text{RO}_2} = \sum_i \eta_i \sum_j \gamma_{i,j} \theta_{i,j} \quad (3)$$

1 which is also plotted in Fig. 9. Consequently, this definition isolates the effect of the
2
3 competition between β scission and oxygen addition to R. It is seen that α_{RO_2} is equal to α at
4
5 low temperatures, but becomes larger than α when the temperature reaches a critical value at
6
7 intermediate to high temperatures. These results then indicate that β scission of the R radical is
8
9 negligible in the low-to-intermediate temperature regime before the turnover state is reached.
10
11
12
13

14 In Fig. 9, the values of α corresponding to the turnover temperatures are marked by
15
16 symbols in the evolution with increasing temperature. It is then of interest to note that the
17
18 turnover temperatures for different pressures largely correspond to a narrow range of α ,
19
20 which, for a given fuel, slightly depends on the pressure range. Figure 10 further shows that
21
22 the turnover states calculated using a constant α agree well with those obtained with detailed
23
24 mechanism, thereby supporting the suggestion that the turnover states can be correlated with
25
26 a critical state of the low-temperature chain branching ratio, with the associated reactions
27
28 involved in the determination of α . The critical value of α is found to be less than one, and is
29
30 different for each fuel studied here, potentially depending on the fuel structure and the range of
31
32 the turnover temperatures. The pressure effect on α is mainly through the equilibrium shift of
33
34 oxygen addition reactions, as implied from Eq. (A11) in the Appendix.
35
36
37
38
39
40
41
42
43
44

45 It is worth noting that the α ratio developed in the current work is based on the widely used
46
47 rate rules in low temperature chemistry, and it can be readily extended to future mechanisms
48
49 with more accurate rate coefficients. Recently, extensive work has been done on the
50
51 uncertainty quantification in kinetic analysis and combustion modeling [28-30]. Following the
52
53 idea of uncertainty propagation, the value of α in this study is rate dependent and will be
54
55 affected by the uncertainty in the rate coefficients, however, such uncertainty will not affect the
56
57
58
59
60
61
62
63
64
65

scaling characteristics and the controlling mechanism identified for the turnover states. On the other hand, there are some other factors that could affect the quantitative determination of the α value, such as the transient effects of species concentration and other possible reaction channels, however, those effects are secondary with negligible influence compared with the leading order quasi-steady assumptions and dominant reaction channels. It is found that the correlation of the turnover state with a constant α value is valid and general for different fuels and mechanisms, as demonstrated by both detailed calculations and experimental validations in Sec. 4.

3.2 Pressure Scaling of Ignition Delay at Turnover States

With the thermodynamic condition identified at the turnover state, we next investigate the reactions controlling the pressure-scaling factor for the turnover ignition delay, shown in Fig. 3. The sensitivity on the turnover ignition delay, defined as $S = d(\ln \tau) / d(\ln k)$ and shown in Fig. 11 for *n*-butane and *n*-heptane (results for *iso*-octane are similar), demonstrates that the most sensitive reactions are those involving the HO₂ and H₂O₂ chemistry, while the secondary important reactions are those involved in the low-T chemistry that determine α .

Peters *et al.* [25] have derived an analytical solution for the ignition delay time τ in the intermediate temperature range based on the quasi-steady-state approximation for HO₂, as

$$\tau = \zeta \left(\frac{(1-\alpha)^3 k_{11}}{2C_{Fuel} (\sum_i k_{12,i})^2 k_{13}^2} \right)^{1/3} \quad (4)$$

where ζ is a proportional constant, α is defined by Eq. (2), C_{Fuel} is the fuel concentration, and k_{11} , $k_{12,i}$, k_{13} correspond to the rate constants of $2\text{HO}_2 \rightarrow \text{H}_2\text{O}_2 + \text{O}_2$ (R11), $\text{RH} + \text{HO}_2 \rightarrow i\text{-R} + \text{H}_2\text{O}_2$ (R12), and $\text{H}_2\text{O}_2 (+\text{M}) \rightarrow 2\text{OH} (+\text{M})$ (R13), respectively. Figure 12 shows the

turnover ignition delay calculated using both detail mechanism and the correlation of Eq. (4).

It is seen that very good agreement is achieved between the two, and the pressure-scaling factor for all of three fuels are well reproduced.

Equation (4) employs the rates of three types of elemental reactions, namely R11, R12, and R13 to correlate the HO_2 and H_2O_2 chemistry on the ignition delay. The ratio α is also employed in Eq. (4) to account for the contribution from low-temperature chemistry on the ignition delay. Since ξ and α are both constants, the change of τ_m is therefore dominated by the rate change of R11, R12, and R13. Especially, the k_{13} corresponding to the decomposition of H_2O_2 with inherent pressure dependence is described by a Troe formula [28] and varies nearly quadratically with changes of the turnover temperature and pressure in the fall-off regime, shown in Fig. 13. Therefore, given the fact that τ_m and k_{13} vary nearly quadratically in terms of pressure with power index being about -2 and 2 respectively, it could be shown from Eq. (4) that the rate change of H_2O_2 decomposition reaction R13 contributes to about two-thirds of the pressure dependence for τ_m , while the rate changes of R12 by the C_{Fuel} , i.e. $C_{Fuel} \sim P$, and the effect of temperature variation on k_{11} , $k_{12,i}$ contribute to the rest one-third.

4. Experimental validation

We now assess the validity of the above two global behaviors and correlations against the ignition delay data available in literature, with the ignition delay for *n*-heptane/air mixture with $\phi = 1$, covering the turnover temperature under 4 – 55 atm, reported in shock tube [8, 31, 32] and in RCM [11].

Gauthier *et al.* [8] reported a pressure scaling factor of -1.64 for the ignition delay under the pressure range of 13-55 atm by using a quadratic fitting to determine the conditions at the turnover state, while the high temperature ignition data show a scaling factor of -0.55 [33]. However, simulation using the LLNL *n*-heptane mechanism [12] shows that the ignition delay at the turnover state acquired from quadratic fitting covering the data points in regimes of both high temperature and near the turnover states is lower than the actual value. Thus, a cubic fitting covering only the points near the turnover temperature is adopted here, as shown in Fig. 14, because computation shows that the cubic relation also cannot hold far from the turnover state. Higher order fitting is also precluded due to the limited number of data points of each dataset.

The conditions corresponding to the turnover states derived from the fitting curve are plotted in Figs. 15 and 16. Strong Arrhenius dependence is found between pressure and temperature. The predictions using the LLNL *n*-heptane mechanism [12] and the updated *n*-heptane mechanism from Karwat *et al.* [24] are also plotted in Fig. 15 for comparison. Both mechanisms reproduce the Arrhenius behavior between the turnover temperature and pressure, and the “slopes” predicted from both mechanisms are close to those from the experimental data. The prediction using $\alpha = 0.64$ from the LLNL *n*-heptane mechanism [12] is plotted in Fig. 14, and agrees with the experimental data very well. Strong power law dependence is also found between ignition delay and pressure with $\tau_m \propto P^{-1.79}$, and the correlation using Eq. (4) agrees well with the measurements in Fig. 16.

It is nevertheless also noted that the fitted turnover temperature under 12 atm from Shen *et al.* [31] and under 20 atm from Gauthier *et al.* [8] show large deviations from the Arrhenius

correlation with pressure, due to the lack of data near the turnover state. The deviation is smaller for the power law between the ignition delay and pressure, as the ignition delay is not sensitive to temperature near the turnover state.

The validation demonstrates the potential to use these two behaviors and correlations to check the consistence among different ignition delay data. The correlations for the two behaviors can also be used for interpolating measured sample data to other conditions.

5. Conclusions

The current work investigated the thermodynamic and chemical kinetic behavior of the upper turnover states of the NTC regime, namely, the Arrhenius behavior between the turnover temperature and pressure, and the power law between the ignition delay and pressure at the turnover states. The underlying mechanism for these two behaviors was further derived from the detailed mechanism. It is found that the turnover states under different pressures correspond to a near-constant ratio α between the OH production from the low-temperature chemistry and the total fuel consumption, and that the pressure-scaling between the ignition delay at the turnover states can be correlated well using the analytical solution from Peters *et al.* [25]. The analytical expression for the ignition delay only employs the rates of three types of reactions on HO₂ and H₂O₂ chemistry and the ratio α .

The two behaviors and correlations were further validated against the ignition delay data for *n*-heptane/air mixture with $\varphi = 1$, covering the turnover states under 4 – 55 atm; demonstrating quantitative agreement.

The correlations for the two global behaviors at turnover states provide insights into the mechanism controlling the transition from low-temperature to intermediate-to-high

temperature regimes. The behaviors and their controlling mechanism can also assist in the design and validation of measured ignition delay data near turnover states, provide reference points for developing correlation of ignition delay from experimental data, and help to identify the regime where the ignition delay is not sensitive to temperature. Furthermore, these correlations can provide targets for optimizing the reaction rate rules for the NTC chemistry.

Acknowledgements

This work was primarily supported by the National Natural Science Foundation of China 91441202. The work at Princeton University was supported by the Combustion Energy Frontier Research Center, an Energy Frontier Research Center funded by the US Department of Energy, Office of Basic Energy Sciences under Award Number DESC0001198. Peng Zhao was also supported by the startup funding at Oakland University. The guidance on the NUIG *n*-butane mechanism from Dr. Kuiwen Zhang from NUIG is very much appreciated.

References

1. C. K. Law; P. Zhao, NTC-affected ignition in nonpremixed counterflow, *Combust. Flame* 159 (2012) 1044-1054.
2. S. Deng; P. Zhao; D. Zhu; C. K. Law, NTC-affected ignition and low-temperature flames in nonpremixed DME/air counterflow, *Combust. Flame* 161 (2014) 1993-1997.
3. P. Zhao; W. Liang; S. Deng; C.K. Law, Initiation and propagation of laminar premixed cool flames, *Fuel* (2015) <http://dx.doi.org/10.1016/j.fuel.2015.11.025>.
4. S. Deng; P. Zhao; M. E. Mueller; C. K. Law, Autoignition-affected stabilization of laminar nonpremixed DME/air coflow flames, *Combust. Flame* 162 (2015) 3437-3445.
5. A. Naidja; C. R. Krishna; T. Butcher; D. Mahajan, Cool flame partial oxidation and its role in combustion and reforming of fuels for fuel cell systems, *Prog. Energy Combust. Sci.* 29 (2003) 155-191.
6. J. F. Griffiths; J. P. MacNamara; C. G. W. Sheppard; D. A. Turton; B. J. Whitaker, The

relationship of knock during controlled autoignition to temperature inhomogeneities and fuel reactivity, *Fuel* 81 (2002) 2219-2225.

7. M. Yao; Z. Zheng; H. Liu, Progress and recent trends in homogeneous charge compression ignition (HCCI) engines, *Prog. Energy Combust. Sci.* 35 (2009) 398-437.

8. B. M. Gauthier; D. F. Davidson; R. K. Hanson, Shock tube determination of ignition delay times in full-blend and surrogate fuel mixtures, *Combust. Flame* 139 (2004) 300-311.

9. P. Dagaut; M. Reuillon; M. Cathonnet, High pressure oxidation of liquid fuels from low to high temperature. 1. n-Heptane and iso-Octane, *Combust. Sci. Technol.* 95 (1993) 233-260.

10. T. Wada; A. Sudholt; H. Pitsch; N. Peters, Analysis of first stage ignition delay times of dimethyl ether in a laminar flow reactor, *Combustion Theory and Modelling* 17 (2013) 906-936.

11. R. Minetti; M. Carlier; M. Ribaucour; E. Therssen; L. R. Sochet, A rapid compression machine investigation of oxidation and auto-ignition of n-Heptane: Measurements and modeling, *Combust. Flame* 102 (1995) 298-309.

12. M. Mehl; W. J. Pitz; C. K. Westbrook; H. J. Curran, Kinetic modeling of gasoline surrogate components and mixtures under engine conditions, *Proc. Combust. Inst.* 33 (2011) 193-200.

13. J. A. Miller; M. J. Pilling; J. Troe, Unravelling combustion mechanisms through a quantitative understanding of elementary reactions, *Proc. Combust. Inst.* 30 (2005) 43-88.

14. H. J. Curran; P. Gaffuri; W. J. Pitz; C. K. Westbrook, A Comprehensive Modeling Study of n-Heptane Oxidation, *Combust. Flame* 114 (1998) 149-177.

15. C. K. Westbrook, Chemical kinetics of hydrocarbon ignition in practical combustion systems, *Proc. Combust. Inst.* 28 (2000) 1563-1577.

16. J. H. Chen; E. R. Hawkes; R. Sankaran; S. D. Mason; H. G. Im, Direct numerical simulation of ignition front propagation in a constant volume with temperature inhomogeneities: I. Fundamental analysis and diagnostics, *Combust. Flame* 145 (2006) 128-144.

17. H. G. Im; P. Pal; M. S. Wooldridge; A. B. Mansfield, A Regime Diagram for Autoignition of Homogeneous Reactant Mixtures with Turbulent Velocity and Temperature Fluctuations, *Combust. Sci. Technol.* 187 (2015) 1263-1275.

18. K. P. Grogan; S. Scott Goldsborough; M. Ihme, Ignition regimes in rapid compression machines, *Combust. Flame* 162 (2015) 3071-3080.

19. S. S. Goldsborough, A chemical kinetically based ignition delay correlation for iso-octane covering a wide range of conditions including the NTC region, *Combust. Flame* 156 (2009) 1248-1262.

20. D. Healy; N. S. Donato; C. J. Aul; E. L. Petersen; C. M. Zinner; G. Bourque; H. J. Curran, n-Butane: Ignition delay measurements at high pressure and detailed chemical kinetic simulations, *Combust. Flame* 157 (2010) 1526-1539.

21. M. Mehl; H. Curran; W. Pitz; C. Westbrook, Chemical kinetic modeling of component mixtures relevant to gasoline, *European Combustion Meeting* (2009)

22. S. M. Villano; L. K. Huynh; H.-H. Carstensen; A. M. Dean, High-Pressure Rate Rules for Alkyl + O₂ Reactions. 2. The Isomerization, Cyclic Ether Formation, and β -Scission Reactions of Hydroperoxy Alkyl Radicals, *J. Phys. Chem. A* 116 (2012) 5068-5089.

23. S. M. Villano; L. K. Huynh; H.-H. Carstensen; A. M. Dean, High-Pressure Rate Rules for

Alkyl + O₂ Reactions. 1. The Dissociation, Concerted Elimination, and Isomerization Channels of the Alkyl Peroxy Radical, *J. Phys. Chem. A* 115 (2011) 13425-13442.

24. D. M. A. Karwat; S. W. Wagnon; M. S. Wooldridge; C. K. Westbrook, Low-temperature speciation and chemical kinetic studies of n-heptane, *Combust. Flame* 160 (2013) 2693-2706.

25. N. Peters; G. Paczko; R. Seiser; K. Seshadri, Temperature cross-over and non-thermal runaway at two-stage ignition of n-heptane, *Combust. Flame* 128 (2002) 38-59.

26. S. S. Merchant; C. F. Goldsmith; A. G. Vandeputte; M. P. Burke; S. J. Klippenstein; W. H. Green, Understanding low-temperature first-stage ignition delay: Propane, *Combust. Flame* 162 (2015) 3658-3673.

27. J. Beeckmann; L. Cai; A. Berens; N. Peters; H. Pitsch, An analytical approximation for low- and high-temperature autoignition for dimethyl ether–air mixtures, *Proc. Combust. Inst.* 35 (2015) 275-281.

28. L. Cai; H. Pitsch, Optimized chemical mechanism for combustion of gasoline surrogate fuels, *Combust. Flame* 162 (2015) 1623-1637.

29. L. Cai; H. Pitsch, Mechanism optimization based on reaction rate rules, *Combust. Flame* 161 (2014) 405-415.

30. H. Wang; D. A. Sheen, Combustion kinetic model uncertainty quantification, propagation and minimization, *Prog. Energy Combust. Sci.* 47 (2015) 1-31.

31. H.-P. S. Shen; J. Steinberg; J. Vanderover; M. A. Oehlschlaeger, A shock tube study of the ignition of n-heptane, n-decane, n-dodecane, and n-tetradecane at elevated pressures, *Energy Fuels* 23 (2009) 2482-2489.

32. H. K. Ciezki; G. Adomeit, Shock-tube investigation of self-ignition of n-heptane-air mixtures under engine relevant conditions, *Combust. Flame* 93 (1993) 421-433.

33. D. C. Horning; D. F. Davidson; R. K. Hanson, Study of the High-Temperature Autoignition of n-Alkane/O/Ar Mixtures, *Journal of Propulsion and Power* 18 (2002) 363-371.

Appendix: Determination of the Low Temperature Chain Branching Ratio α

Species time histories plotted in Fig. 5 show that it is justified to set R, RO₂, QOOH, O₂QOOH and KET into quasi-steady state, for which the production rates balance the consumption rates, yielding:

$$w_1 = w_2 + w_3 - w_{-3} \quad w_{-3} = w_4 + w_5 - w_{-5} \quad (A1, A2)$$

$$w_5 = w_{-5} + w_6 + w_7 + w_8 - w_{-8} \quad (A3)$$

$$w_8 = w_{-8} + w_9, \quad w_9 = w_{10} \quad (A4, A5)$$

From the definition of θ and applying Eq. (A3-A5), we have

$$\theta = \frac{w_7 + w_9 + w_{10}}{w_5} = \frac{w_7 + 2w_9}{w_{-5} + w_6 + w_7 + w_9} \quad (A6)$$

Introducing Eq. (A4) and the rate expressions into Eq. (A6), and further define

$K_8 = k_8 / (k_{-8} + k_9)$, we obtain

$$\theta = \frac{k_7 + 2k_9 K_8 [O_2]}{k_{-5} + k_6 + k_7 + k_9 K_8 [O_2]} \quad (A7)$$

Recall the definition of γ , and assume $w_5 \gg w_{-5}$, we obtain

$$\gamma = \frac{w_5 - w_{-5}}{w_3 - w_{-3}} \approx \frac{w_5}{w_4 + w_5} = \frac{k_5}{k_4 + k_5} \quad (A8)$$

which is only temperature dependent.

Recall the definition of β :

$$\beta = \frac{w_3 - w_{-3}}{w_1} = \frac{w_3 - w_{-3}}{w_2 + w_3 - w_{-3}} \quad (A9)$$

Substitute Eq. (A2) into Eq. (A9) with rate expressions, and further define

$K_3 = k_3 / (k_{-3} + k_4 + k_5)$, we have β as

$$\beta = \frac{(k_4 + k_5)K_3[\text{O}_2]}{k_2 + (k_4 + k_5)K_3[\text{O}_2]} \quad (\text{A10})$$

As there is only one isomer of R is involved, we have $\eta = 1$. By substituting Eq. (A7), Eq. (A8) and Eq. (A10) into Eq. (2), we obtain

$$\alpha = \frac{(k_4 + k_5)K_3[\text{O}_2]}{k_2 + (k_4 + k_5)K_3[\text{O}_2]} \frac{k_5}{k_4 + k_5} \frac{k_7 + 2k_9K_8[\text{O}_2]}{k_{-5} + k_6 + k_7 + k_9K_8[\text{O}_2]} \quad (\text{A11})$$

Clearly, it depends on both temperature and pressure, with the pressure dependence mainly through the oxygen concentration and therefore oxygen addition reactions.

To determine α using detailed mechanism, the reaction path and rate corresponding to each isomer should be taken into account for different low-temperature intermediates. In the following, we shall present the adaptive definitions of β_i , $\gamma_{i,j}$ and $\theta_{i,j}$ considering isomeric effects.

Merchant *et al.* [26] noticed that the decomposition of O_2QOOH to KET and subsequent reactions may lead to three OH in some cases, *e.g.*, the case where CH_2CHO is in the products of KET decomposition. Besides, detailed mechanism shows that R7 and subsequent reactions may lead to more than one OH in some cases, *e.g.*, the reaction of $\text{C}_7\text{H}_{14}\text{OOH1-3} \rightarrow \text{C}_4\text{H}_7\text{OOH1-4} + \text{NC}_3\text{H}_7$ will lead to two OH radicals in the LLNL *n*-heptane mechanism [12]. For generality, the number of OH produced by R7 and R9 with their subsequent reactions are denoted as n_7 , n_9 , respectively, and $\theta_{i,j}$ is determined by rewriting Eq. (A7) as

$$\theta_{i,j} = \left\{ \frac{n_7 k_7^{i,j} + n_9 k_9^{i,j} K_8^{i,j}[\text{O}_2]}{k_{-5}^{i,j} + k_6^{i,j} + k_7^{i,j} + k_9^{i,j} K_8^{i,j}[\text{O}_2]} \right\}_{i,j-\text{QOOH}} \quad (\text{A12})$$

where the superscript i,j indicates that the rate constant corresponds to the specific isomer of i,j -QOOH.

As $i\text{-RO}_2$ can isomerize to several isomers of QOOH, $\gamma_{i,j}$ can be determined by including

all of the isomerization reactions from $i\text{-RO}_2$ and rewriting Eq. (A8) as

$$\gamma_{i,j} = \left\{ \frac{k_5^{i,j}}{k_4^i + \sum_m k_5^{i,m}} \right\}_{i\text{-RO}_2 \rightarrow i,j\text{-QOOH}} \quad (\text{A13})$$

where $k_5^{i,j}$ corresponds to the rate of isomerization from $i\text{-RO}_2$ to $i,j\text{-QOOH}$.

The β_i can be determined by specifying the rates in Eq. (A10) for each isomer of $i\text{-R}$ as.

$$\beta_i = \left\{ \frac{(k_4^i + \sum_m k_5^{i,m})K_3^i[\text{O}_2]}{k_2^i + (k_4^i + \sum_m k_5^{i,m})K_3^i[\text{O}_2]} \right\}_{i\text{-R}} \quad (\text{A14})$$

Similarly, η_i is determined by

$$\eta_i = \frac{w_1^i}{\sum_m w_1^m} = \frac{k_1^i}{\sum_m k_1^m} \quad (\text{A15})$$

List of Tables

Table 1. Reactions used for determining α .

Label	Reaction	Rate coefficient	Rate
R1	$\text{RH} + \text{OH} \rightarrow \text{R} + \text{H}_2\text{O}$	k_1	w_1
R2	$\text{R} \rightarrow \text{R1} + \text{Alkene}$	k_2	w_2
R3	$\text{R} + \text{O}_2 \rightleftharpoons \text{RO}_2$	k_3, k_{-3}^{a}	w_3, w_{-3}^{a}
R4	$\text{RO}_2 \rightarrow \text{Alkene} + \text{HO}_2$	k_4	w_4
R5	$\text{RO}_2 \rightleftharpoons \text{QOOH}$	k_5, k_{-5}	w_5, w_{-5}
R6	$\text{QOOH} \rightarrow \text{Alkene} + \text{HO}_2^{\text{b}}$	k_6	w_6
R7	$\text{QOOH} \rightarrow \text{Epoxide} + \text{OH}^{\text{c}}$	k_7	w_7
R8	$\text{QOOH} + \text{O}_2 \rightleftharpoons \text{O}_2\text{QOOH}$	k_8, k_{-8}	w_8, w_{-8}
R9	$\text{O}_2\text{QOOH} \rightarrow \text{KET} + \text{OH}$	k_9	w_9
R10	$\text{KET} \rightarrow \text{P} + \text{OH}$	k_{10}	w_{10}

^a k_{-3} is corresponding to the reverse rate of k_3

^b Decomposition reaction of QOOH that does not lead to OH production is also classified in R6

^c Decomposition reaction of QOOH that leads to one OH radical is also classified in R7

Figure Captions

Fig. 1. Schematic mechanism for the low temperature oxidation of a large alkane, RH.

Fig. 2. Total ignition delay as a function of the initial temperature T for stoichiometric n -heptane/air mixture under various pressure. The turnover ignition delays are marked by the triangle symbols, showing an Arrhenius dependence on the turnover temperature for different pressures.

Fig. 3. Computed ignition delays at turnover temperatures for stoichiometric n -butane, n -heptane and iso -octane in air. The turnover ignition delays show a power law dependence on the pressure, to the order of -2, approximately.

Fig. 4. Dependence of pressure and temperature at the turnover states for stoichiometric n -butane, n -heptane and iso -octane in air.

Fig. 5. Computed species and temperature profiles for stoichiometric n -butane/air mixture at the turnover temperature under 20 atm.

Fig. 6. Magnitudes of the rates of important reactions producing or consuming OH (a) and fuel molecule (b) for stoichiometric n -butane/air mixture at the turnover temperature under 20 atm.

Fig. 7. The brute force sensitivity on the turnover temperature for stoichiometric n -butane/air mixture.

Fig. 8. Reaction flux from RH to OH and the definition of η_i , β_i , γ_{ij} , θ_{ij} . Only one of the isomers at every step is presented as an example.

Fig. 9. Computed α and α_{RO2} as a function of temperature under various pressures.

Fig. 10. Turnover temperature calculated using detailed mechanism and obtained at a constant α .

Fig. 11. The brute force sensitivity on the ignition delay at turnover temperature for *n*-butane and *n*-heptane.

Fig. 12. The ignition delays from simulation using detailed mechanism and from the correlation using Eq. (4).

Fig. 13. The rates of $\text{H}_2\text{O}_2 (+\text{M}) \rightarrow 2\text{OH} (+\text{M})$ as a function of pressure at turnover temperature, and those rates at various temperature and pressure.

Fig. 14. Ignition delay data for stoichiometric *n*-heptane/air mixture under various pressure.

All the data are scaled to the listed pressures using $\tau \propto P^{-1}$ to account for deviations in experimental pressure [31]. Each dataset is fitted using a cubic polynomial and covering the points near turnover temperature.

Fig. 15. The pressure and turnover temperature of each dataset.

Fig. 16. The ignition delay from measurements and those from correlation using Eq. (4).

Fig. 1. Schematic mechanism for the low temperature oxidation of a large alkane, RH.

Fig. 2. Total ignition delay as a function of the initial temperature T for stoichiometric n -heptane/air mixture under various pressure. The turnover ignition delays are marked by the triangle symbols, showing an Arrhenius dependence on the turnover temperature for different pressures.

Fig. 3. Computed ignition delays at turnover temperatures for stoichiometric n -butane, n -heptane and iso -octane in air. The turnover ignition delays show a power law dependence on the pressure, to the order of -2, approximately.

Fig. 4. Dependence of pressure and temperature at the turnover states for stoichiometric n -butane, n -heptane and iso -octane in air.

Fig. 5. Computed species and temperature profiles for stoichiometric *n*-butane/air mixture at the turnover temperature under 20 atm.

(a)

(b)

Fig. 6. Magnitudes of the rates of important reactions consuming OH (a) and fuel molecule (b) for stoichiometric *n*-butane/air mixture at the turnover temperature under 20 atm. The presented reactions contribute nearly 90% to the total consumption rate at the instant of 75% τ_m .

Fig. 7. The brute force sensitivity on the turnover temperature for stoichiometric *n*-butane/air mixture.

Fig. 8. Reaction flux from RH to OH and the definition of $\eta_i, \beta_i, \gamma_{ij}, \theta_{ij}$. Only one of the isomers at every step is presented as an example.

(a)

(b)

(c)

Fig. 9. Computed α and α_{RO_2} as a function of temperature under various pressures for different fuels.

Fig. 10. Turnover temperature calculated using detailed mechanism (symbols) and obtained at a constant α (lines).

(a)

(b)

Fig. 11. The brute force sensitivity on the ignition delay at turnover temperature for *n*-butane and *n*-heptane.

Fig. 12. The ignition delays from simulation using detailed mechanism (symbols) and from the correlation using Eq. (4).

Fig. 13. The rates of $\text{H}_2\text{O}_2 (+\text{M}) \rightarrow 2\text{OH} (+\text{M})$ as a function of pressure at turnover temperature for stoichiometric *n*-butane, *n*-heptane and *iso*-octane in air (symbols), and those rates at various temperature and pressure (dash lines). The rate constants are adopted from each detailed mechanism.

Fig. 14. Ignition delay data for stoichiometric *n*-heptane/air mixture under various pressure.

All the data are scaled to the listed pressures using $\tau \propto P^{-1}$ to account for deviations in experimental pressure [31]. Each dataset is fitted using a cubic polynomial and covering the points near turnover temperature.

Fig. 15. The dependence of pressure on turnover temperatures for stoichiometric n -heptane/air mixture obtained from different methods.

Fig. 16. The ignition delay from measurements and those from correlation using Eq. (4).