**Cryopreservation of Lipid Bilayers by LEA Proteins from *Artemia franciscana* and
Trehalose**

Daniel S. Moore and Steven C. Hand*

Department of Biological Sciences
Division of Cellular, Developmental, and Integrative Biology
Louisiana State University
Baton Rouge, Louisiana 70803 USA

*To whom correspondence should be addressed. e-mail: shand@lsu.edu.

Running Head: Protection of lipid bilayers against freezing damage

Key Words: late embryogenesis abundant protein, trehalose, liposome, *Artemia franciscana*,
freezing, intrinsically disordered protein, mitochondria.

Abbreviation: Cho, Cholesterol; CL, cardiolipin; PC, phosphatidylcholine; PE,
phosphatidylethanolamine; PI, phosphatidylinositol; PS, phosphatidylserine; mitochondrial
permeability transition pore (MPTP); large unilamellar vesicles (LUVs); 5(6)-
carboxyfluorescein (CF).

Abstract

The capacity of Late Embryogenesis Abundant (LEA) proteins and trehalose to protect liposomes against freezing-induced damage was examined by measuring the leakage of 5(6)-carboxyfluorescein (CF). Liposomes were prepared to simulate the lipid compositions of the inner leaflet of the plasma membrane, outer mitochondrial membrane (OMM), and inner mitochondrial membrane (IMM). Two recombinant LEA proteins belonging to Group 3 (AfrLEA2 and AfrLEA3m) were expressed and purified from embryos of *Artemia franciscana*. Only OMM-like liposomes were significantly protected by AfrLEA2 and AfrLEA3m against freeze-thaw damage; at the highest protein:lipid mass ratio tested, leakage of CF was 56.3% of control with AfrLEA3m and 29.3% with AfrLEA2. By comparison, trehalose provided protection to all compositional types. The greatest stabilization during freezing occurred when trehalose was present on both sides of the bilayer. When mitochondria isolated from rat liver were freeze-thawed in trehalose solution, the OMM remained intact based on the absence of increased oxygen consumption when cytochrome *c* was added during oxidative phosphorylation (OXPHOS). Respiratory control ratios (OXPHOS/LEAK) were depressed by only 30% after freeze-thawing in trehalose compared to non-frozen controls, which indicated some retention of OXPHOS capacity by the IMM. Trehalose then was loaded into the matrix (0.24 $\mu\text{mol/mg}$ mitochondrial protein) by transient opening of the permeability transition pore, a procedure optimized for retention of OMM integrity. Surprisingly, respiratory control ratios were not improved after freeze-thawing with external plus matrix trehalose, when compared to external trehalose alone. This result could perhaps be explained by insufficient accumulation of matrix trehalose.

Introduction

Anhydrobiotic organisms like embryos of brine shrimp, *Artemia franciscana*, are able to survive the virtual absence of water [15,20,24]. Many organisms that tolerate extreme water stress utilize common protective mechanisms like depressed cellular metabolism and accumulation of specialized proteins and solutes of low molecular weight [14,16,17,20,33,64]. Prominent among the protective molecules found to accumulate in embryos of brine shrimp are late embryogenesis abundant (LEA) proteins [10,34,41,54,57,58] and the sugar trehalose [12,13,27,31,43]. LEA proteins are a family of hydrophilic and often intrinsically disordered proteins expressed in desiccation-tolerant plants and animals [35,55]. Trehalose is a non-reducing disaccharide that accumulates in desiccation tolerant organisms in response to water deficit or is constitutively maintained in developmental stages commonly encountering water stress [13,17,39,49]. In the present study, we evaluate the ability of two LEA proteins from *A. franciscana* to protect liposomes of different lipid compositions against freeze-induced damage, and we compare their effectiveness to the well-established capacity of trehalose for liposome stabilization. Finally, we expand upon existing information regarding the ability of trehalose to preserve the function of isolated mitochondrial during freezing and thawing.

Several LEA proteins form amphipathic alpha-helices capable of interacting with lipid bilayers [30,42,44,45,51,52,53]. However, stabilization of lipid bilayers during water stress has only been experimentally demonstrated for three LEA proteins from plants and two from animals. COR15a and COR15b (LEA23 and LEA24) from *Arabidopsis thaliana* protect chloroplast membranes from freeze-induced damage both *in vitro* and *in vivo* [4,47,50]. PsLEAm from the pea plant (*Pisum sativum*) improves the stability of either dried or frozen liposomes [52]. AfrLEA2 and AfrLEA3m from embryos of *A. franciscana* protect liposomes

87 from desiccation-induced damage [42]. These latter two LEA proteins were used in this
88 current freezing study with liposomes that mimic the lipid content the inner leaflet of the
89 plasma membrane (ILPM), outer mitochondrial membrane (OMM) and inner mitochondrial
90 membrane (IMM).

91 Previous studies have demonstrated the ability of trehalose to protect liposomes and
92 membranes from freezing and desiccation-induced damage [11,17,21,23,38,48,62,63]. Like
93 many sugars, trehalose forms a glass at low water contents and/or temperatures [18].
94 Trehalose is sometimes superior to other sugars at protecting macromolecules due to its
95 relatively high glass transition temperature [22,23]. Sugar glasses provide structural support
96 and immobilize cellular components, which are important for preventing deleterious
97 interactions. In addition to sugar glass formation, there is evidence that trehalose has the
98 ability to bind to dry membranes in the place of water [25,59]. This “water replacement”
99 favors the retention of the liquid crystalline phase of lipid bilayers. For maximal protection of
100 a membrane trehalose must be present on both sides of a lipid bilayer [23,25]. There is some
101 evidence that trehalose protection during drying is dependent on the lipid composition of the
102 target membrane [36,37].

103 For cells to survive water stress, it is critical that cellular components such as the
104 mitochondria be preserved [5,32]. To optimally preserve mitochondria, it would appear
105 important that trehalose contact both surfaces of the relevant bilayers [26,38]. The outer
106 membrane of mitochondria is permeable to solutes with a molecular weight less than 5 kDa
107 due to non-specific protein channels like porin [40,56]. While chemically unmodified
108 trehalose is apparently not permeable across the IMM of mammalian cells [cf. 1], it is possible
109 to load disaccharides into the matrix of isolated mammalian mitochondria by transient opening

110 of the mitochondrial permeability transition pore (MPTP) [2,38]. The MPTP is a mega-pore of
111 the inner membrane that is comprised of dimers of the ATP synthase [7,28]. Based on (1)
112 results below that show protection by trehalose is greatest when present on both sides of IMM-
113 like liposomes during freeze/thawing and (2) previous experiments on drying of isolated
114 mitochondria [38], we viewed it appropriate to test whether loading trehalose into the matrix of
115 mitochondria isolated from rat liver prior to freezing improves retention of respiratory function
116 after thawing.
117

Materials and Methods

Biochemicals

Lipids and cholesterol were purchased from Avanti Polar Lipids (Alabaster, AL). All lipids were naturally derived in order to mimic the fatty acid distribution found in eukaryotic cells. Phosphatidylcholine (PC), phosphatidylethanolamine (PE), and phosphatidylinositol (PI) were derived from bovine liver (Avanti Polar Lipids; product numbers 840055, 840026, 840042). Because these phospholipids were naturally derived, they contain a mixture of fatty acids. However, the predominate phospholipid species in each sample is reported by Avanti to be 1-stearoyl-2-oleoyl-sn-glycero-3-phosphocholine, 1-stearoyl-2-arachidonoyl-sn-glycero-3-phosphoethanolamine, and 1-stearoyl-2-arachidonoyl-sn-glycero-3-phosphoinositol. Phosphatidylserine (PS; predominantly 1-stearoyl-2-oleoyl-sn-glycero-3-phospho-L-serine) was purified from bovine brain tissue (Avanti Polar Lipids; product number 840032) and cardiolipin (CL; predominantly 1',3'-bis[1,2-dilinoleoyl-sn-glycero-3-phospho]-sn-glycerol) from bovine heart (Avanti Polar Lipids; product number 840012). Cholesterol was isolated from ovine wool (Avanti Polar Lipids; product number 700000P). 5(6)-Carboxyfluorescein (CF) (product number 21877) was obtained from Sigma-Aldrich (St. Louis, MO). Trehalose (product number T-104-4) was purchased from Pfanstiehl (Waukegan, IL). Bovine serum albumin (product number A6003, essentially fatty acid free) was purchased from Sigma-Aldrich.

Liposome Preparation

Lipids were combined with mass ratios simulating the inner leaflet of the plasma membrane (ILPM), outer mitochondrial membrane (OMM) and inner mitochondrial membrane (IMM) inner mitochondrial membrane of mammalian cells as previously reported [42]: ILPM

141 (50 % PC, 30 % PE, 10 % PS, 10 % Cho), IMM (50 % PC, 30 % PE, 20 % CL), OMM (55 %
142 PC, 30 % PE, 15 % PI). For comparison to a non-physiological bilayer, liposomes composed
143 entirely of phosphatidylcholine were also prepared. Large unilaminar vesicles (LUVs)
144 extruded and rinsed precisely as described previously [42]. Liposomes were mixed with 250
145 mM trehalose (19:1 sugar:lipid mass ratio) and/or LEA proteins to yield various protein:lipid
146 mass ratios between 0.1 and 0.4. Recombinant AfrLEA3m and AfrLEA2 were purified as
147 described by Boswell et al. [10]. Liposomes were mixed with equal amounts of BSA in order
148 to compare protection by LEA proteins to that of a protein not associated with membrane
149 stabilization. For some experiments (i.e., Fig. 2B), 250 mM trehalose was included during the
150 preparation of LUVs, so that the sugar also was present inside of the liposomes. The final
151 concentration of liposomes was approximately 5 mg total lipid/ml.

152 *Liposome Freezing and Measurements of CF Leakage*

153 For CF leakage assays, 2 μ l droplets of liposome mixtures were placed in the wells of
154 opaque 96-well plates (OptiPlate 96-F, PerkinElmer, Waltham, MA). Plates were covered in
155 plastic wrap and frozen at -80°C for at least 1 week. Fluorescence of CF is strongly quenched
156 at the high concentration of 100 mM contained inside of the liposomes, but increases when CF
157 is released into the medium [60]. CF fluorescence was measured as described previously [42].
158 Due to inherent differences in control leakage across compositional types, experimental
159 treatments were normalized to the controls (no protectant) when comparing stabilization of
160 liposomes with various lipid compositions.

161 *Isolation of Mitochondria*

162 Mitochondrial isolation from rat liver was performed as outlined by Gnaiger et al. [29].
163 Briefly, male Sprague-Dawley rats were euthanized by CO₂ asphyxiation in accordance with

164 protocol 12-061 approved by the LSU Institutional Animal Care and Use and Committee. The
165 liver (approx. 10-15 g) was excised and immediately placed in ice-cold homogenization
166 medium A (HMA), which was composed of 250 mM trehalose, 10 mM Tris, 1 mM EGTA (pH
167 7.4). The large lobes of the liver were finely diced and any connective tissue observed was
168 removed. The minced liver was rinsed several times in HMA and transferred to a 40 ml
169 Dounce homogenizer (Kontes Glass Co., Vineland, NJ). The liver was suspended in 30 ml of
170 homogenization medium B (HMB), which was HMA fortified with 0.2% BSA. The
171 homogenate was centrifuged at 700 x g for 10 min at 4°C. The supernatant was centrifuged at
172 9,000 x g for 10 min at 4°C. The pellet was resuspended in 30 ml of fresh HMB and
173 centrifuged once more at 9,000 x g for 10 min at 4°C. The resulting mitochondrial pellet was
174 resuspended in HMA at 40-60 mg protein/ml as determined by Coomassie Plus (Bradford)
175 protein assay (Thermo Fisher Scientific Inc., Rockford, IL).

176 *Transient Opening of the MPTP*

177 Loading the matrix of isolated mitochondria with trehalose via transient opening of the
178 mitochondrial permeability transition pore (MPTP) was performed as described by Al-Nassar
179 and Crompton [2]. Mitochondria suspended in HMA were diluted to 0.5-1 mg/ml in HMA that
180 contained no EGTA, which resulted in a buffer composed of 250 mM trehalose, 10 mM Tris,
181 and 25 μ M EGTA (pH 7.4). MPTP formation was induced by the addition of glutamate (10
182 mM), malate (2 mM), KH_2PO_4 (1 mM), and Ca^{2+} [6,8,65]. For freezing studies, MPTP
183 opening was induced by the addition of 35 μ M Ca^{2+} . After 1 min the pore was closed by the
184 addition of 2 mM EGTA, 1 mM MgCl_2 , 1 mM NAD^+ , and 2 mg/ml BSA [9,65]. Mitochondria
185 were incubated at room temperature for ~8 min after the addition of EGTA to allow complete
186 closure of the pore [2,65]. Control (i.e. non-permeabilized) mitochondria were incubated

187 under identical conditions with the exception that calcium was not added to the medium. For
188 comparison of trehalose-incubated mitochondria to ones that were not exposed to trehalose,
189 mitochondria were isolated and frozen in a buffer composed of 150 mM KCl, 10 mM Tris, 2
190 mM EGTA, and 2 mg/ml BSA (pH 7.4).

191 *Quantification of Trehalose by HPLC*

192 After opening and reclosing of the MPTP, mitochondria were sedimented by
193 centrifugation (10,000 x g for 10 min at 4°C). To remove extra-mitochondrial trehalose,
194 mitochondrial pellets were resuspended in buffer composed of 250 mM sucrose, 10 mM Tris,
195 and 1 mM EGTA (pH 7.4) and centrifuged once more at 10,000 x g for 10 min at 4°C, after
196 which the supernatant was discarded and the mitochondrial pellet resuspended in water. The
197 suspension was frozen at -80°C and thawed to ensure complete release of trehalose from the
198 matrix. The sample was heated at 95°C for 20 min then centrifuged at 20,000 x g for 30 min at
199 4°C to remove precipitated protein. HPLC analysis was performed as described by Patil et al.
200 [43]. Values were expressed as μ mol trehalose per mg mitochondrial protein.

201 *Freezing of Mitochondria*

202 Mitochondria were frozen as described by Yamaguchi et al. [63]. Mitochondria were
203 suspended at 40 mg mitochondrial protein/ml in HMB. Plastic microcentrifuge tubes (0.6 ml)
204 were placed on a bed of dry ice pellets, and a 5 μ l droplet of mitochondrial suspension was
205 pipetted into each tube. The tubes were transferred in to liquid nitrogen and then placed in a -
206 80°C freezer overnight. Droplets were thawed quickly by warming between fingers.

207 *Measurements of Mitochondrial Respiration*

208 Oxygen consumption of mitochondria was measured at 37°C with an Oxygraph 2K
209 (OROBOROS Instruments, Innsbruck, Austria), and the data were analyzed with DatLab

210 software (OROBOROS Instruments). The rate of oxygen consumption was calculated as the
211 time derivative of oxygen concentration and normalized per mg mitochondrial protein (nmol
212 $\text{O}_2 \text{ s}^{-1} \text{ mg mitochondrial protein}^{-1}$). Complex II-dependent respiration was measured in the
213 presence of succinate (10 mM) and rotenone (0.5 μM ; inhibitor of Complex I). Proton leak
214 respiration (LEAK state, or state 4) was measured in the presence of substrate without ADP.
215 The addition of ADP (5 mM) stimulated maximum oxidative phosphorylation (OXPHOS state,
216 or state 3). The respiratory control ratio (RCR) was calculated by dividing OXPHOS by
217 LEAK respiration. Cytochrome *c* (cyt *c*; 4 mM) was added to the respiration medium during
218 the OXPHOS state in order to assess the integrity of the outer membrane; an increase in
219 oxygen consumption was indicative of compromised membrane integrity. Finally, carbonyl
220 cyanide-4-(trifluoromethoxy) phenylhydrazone (FCCP) was titrated step-wise to estimate the
221 maximum respiratory capacity of the electron transport system in the uncoupled state (ETS
222 state).

223 *Statistics*

224
225 Two-way ANOVA paired with a Tukey post-hoc test were used to compare protection
226 of each type of liposome by AfrLEA3m, AfrLEA2, or BSA across the range of protein:lipid
227 mass ratios tested (Prism 6; GraphPad Software, La Jolla, CA). A Bonferroni multiple
228 comparisons test of sample means was used to identify statistical differences between trehalose
229 and protein combinations compared to each component alone (Prism 6). 1-way ANOVA
230 paired with a Tukey multiple comparisons test was used to compare protection of each type of
231 liposome by trehalose alone. Paired and unpaired t-tests were used to compare trehalose
232 content and mitochondrial respiration rates and RCRs before and after freezing or
233 permeabilization (Prism 6). Significance level was set at $p \leq 0.05$.

Results

Stability of Liposomes Frozen with LEA Proteins

Protection of liposomes by LEA proteins varied depending on the lipid composition of the liposomes (Fig. 1). The only statistically significant (2-way ANOVA, Bonferroni multiple comparisons test, $p > 0.05$, $n = 3-6$) improvement in stability during freezing and thawing occurred with liposomes that simulated the composition of the outer mitochondrial membrane (OMM). When quantified at the highest protein:lipid mass ratio tested, leakage of CF from OMM-like liposomes was $56.3 \pm 2.8\%$ (mean $\pm$ SD, $n = 6$) of control leakage in the presence of AfrLEA3m and $29.3 \pm 2.5\%$ (mean $\pm$ SD, $n = 6$) in the presence of AfrLEA2 (Fig. 1A and B; 2-way ANOVA, Bonferroni multiple comparisons test, $p > 0.05$, $n = 3-6$). After freeze-thawing in the presence of AfrLEA3m, there were no statistical differences compared to controls in CF loss from liposomes that mimicked the inner leaflet of the plasma membrane (ILPM), the inner mitochondrial membrane (IMM), or in those composed of 100% phosphatidylcholine (Fig. 1A, PC; 2-way ANOVA, Bonferroni multiple comparisons test, $p > 0.05$, $n = 3-6$). Likewise, AfrLEA2 did not significantly diminish CF leakage after freeze-thawing from IMM, ILPM, or PC liposomes (Fig. 1B; 2-way ANOVA, Bonferroni multiple comparisons test, $p > 0.05$, $n = 3-6$). The control protein, BSA, did not depress CF leakage from any liposomes tested except at the highest protein:lipid mass ratio tested with OMM-like liposomes, and this depression was very modest (Fig. 1C; 2-way ANOVA, Bonferroni multiple comparisons test, $p > 0.05$, $n = 3-6$). It is noteworthy that the ILPM-like liposomes generally displayed greater variability in leakage during freezing experiments compared to other compositional types for reasons that are unclear.

Stability of Liposomes Frozen with Trehalose

257 Relative to controls (no sugar added), 250 mM external trehalose significantly
258 depressed CF leakage from liposomes for all compositional types tested (Fig. 2A; 2-way
259 ANOVA, Bonferroni multiple comparisons test, $p < 0.05$). Trehalose protected non-biological
260 liposomes composed of 100% PC and liposomes with lipid compositions similar to the OMM
261 and ILPM with equal efficacy. However, CF loss from IMM-like liposomes in the presence of
262 trehalose was significantly greater than for the other liposomes (1-way ANOVA, Tukey
263 multiple comparisons test, $p < 0.05$, $n = 3-6$). Therefore, lipid composition did affect the degree
264 to which trehalose improved liposome stability. The relatively low protection afforded to
265 IMM-like liposomes is consistent with loss of inner membrane function by isolated
266 mitochondria frozen with external trehalose [63]. This result may indicate that the requirement
267 for trehalose to be present on both sides of a membrane to provide maximal protection could be
268 particularly important for preservation of the IMM during freeze/thawing. In order to evaluate
269 maximal protection by trehalose, IMM-like liposomes were prepared with 250 mM trehalose
270 inside and outside. Loss of CF from IMM-like liposomes significantly decreased (1-way
271 ANOVA, Tukey multiple comparisons test, $p < 0.05$, $n = 3$) from $22.9 \pm 1.3\%$ (mean $\pm$ SD, $n = 6$)
272 in the presence of external trehalose alone to $7.1 \pm 0.2\%$ (mean $\pm$ SD, $n = 3$) when liposomes
273 were provisioned with both internal and external trehalose (Fig. 2B). Low leakage was
274 observed in liposomes even without protectant when frozen at -80°C as performed in this study
275 (Fig. 2B), an issue to which we will return in the Discussion section.

276 Stabilization of liposomes by LEA proteins in combination with trehalose was also
277 examined. No significant increase in protection beyond that seen with trehalose alone was
278 detected for any of the liposomes when LEA proteins were added in conjunction with trehalose
279 (data not shown).

280 *Optimization of Transient Opening of the MPTP*

281 Because the presence of trehalose on both sides of the lipid bilayer improved the integrity
282 of IMM model liposomes during freezing, we wished to compare stabilization of isolated
283 mitochondria frozen with and without trehalose in the matrix. Transient opening of the MPTP
284 has been used to load sugars into the matrix of mitochondria isolated from rat liver [2,38].
285 However, the opening and closing of the pore required optimization to preserve the respiratory
286 function of mitochondria. Mitochondria that were permeabilized by the addition of 35 μM Ca^{2+}
287 for 1 min and then resealed accumulated significant trehalose and also retained respiratory
288 function and outer membrane integrity (Fig. 3). By inducing transient opening of the MPTP in
289 this manner, 0.24 ± 0.03 μmol trehalose/mg mitochondrial protein (mean $\pm$ SD, $n = 6$) was
290 incorporated into the matrix (Fig. 3A). The increase in OXPHOS respiration after the addition
291 of exogenous cytochrome *c* was less than 10% and not significantly different from that
292 measured for non-permeabilized mitochondria (Fig. 3B; Student's paired t-test, $p > 0.05$, $n=6$),
293 which indicated that the outer membrane remained substantially intact after transient
294 permeabilization with 35 μM Ca^{2+} . The respiratory control ratio (RCR) was unchanged as well
295 after permeabilization with 35 μM Ca^{2+} (Fig. 3C; Student's paired t-test, $p > 0.05$, $n=6$). In
296 contrast, when 100 μM Ca^{2+} for 1 min was used to induce pore formation, RCR values
297 significantly decreased (Fig. 3C; Student's paired t-test, $p < 0.05$, $n=6$), which was a result of
298 diminished OXPHOS with no change in LEAK (data not shown). Stimulation of OXPHOS by
299 cyt *c* was increased (Fig. 3B; Student's paired t-test, $p < 0.05$, $n=6$). The amount of trehalose
300 loaded into the matrix of mitochondria after permeabilizing with 100 μM Ca^{2+} (0.31 ± 0.03
301 μmol trehalose/mg mitochondrial protein) was significantly greater than achieved with 35 μM
302 Ca^{2+} (Fig. 3A; Student's paired t-test, $p < 0.05$, $n=3-5$). Nevertheless, to test whether matrix

303 trehalose could improve mitochondrial function after freeze-thawing, 35 μM Ca^{2+} was chosen
304 for MPTP opening due the lack of impact on OMM integrity and RCR.

305 *Cryoprotection of Isolated Mitochondria by Trehalose*

306 Respiratory function of mitochondria before and after freezing was measured to assess
307 protection of mitochondria by trehalose (Fig. 4). Prior to freezing, all RCR values were > 4
308 when succinate was used as a substrate. In the absence of trehalose (replaced by 150 mM
309 KCl), the respiratory control ratio was diminished to 0.98 ± 0.36 (mean $\pm$ SD, $n = 4$) after
310 freezing and thawing (Fig. 4A, right panel). This RCR value reflects the lack of respiratory
311 stimulation above LEAK upon ADP addition; there was both an increase in LEAK and a
312 decrease in OXPHOS after freezing in KCl medium compared to non-frozen controls (Fig. 4A,
313 left panel; unpaired t-tests, Holm-Sidak post-hoc, $p < 0.05$, $n=4$). The addition of cyt *c* resulted
314 in an increase in respiration above OXPHOS after freeze-thawing, which indicated
315 compromised integrity of the outer membrane when frozen with KCl (Fig. 4A, left panel;
316 unpaired t-tests, Holm-Sidak post-hoc, $p < 0.05$, $n=4$). In contrast, mitochondria frozen with
317 300 mM external trehalose showed improved RCR values (3.8 ± 0.4 , mean $\pm$ SD, $n=4-5$)
318 compared to those frozen in KCl (Fig. 4A and B, right panels; Student's unpaired t-test, $p \leq$
319 0.05 , $n = 4$). Additionally, the external trehalose eliminated any cyt *c* effect (compared to
320 OXPHOS) post-freezing (Fig. 4B, left panel; unpaired t-tests, Holm-Sidak post-hoc, $p > 0.05$,
321 $n=4-5$); thus, the outer membrane was protected. Providing trehalose to the matrix
322 compartment along with external trehalose did not improve freeze tolerance compared to
323 mitochondria frozen with external trehalose alone (Fig. 4C). Indeed, the RCR value post
324 freezing (2.6 ± 0.2 , mean $\pm$ SD, $n = 4$) was actually less than that measured with external
325 trehalose alone (Student's unpaired t-test, $p \leq 0.05$, $n = 4-5$). It may be that the amount of

326 trehalose loaded into the matrix with 35 μM Ca^{2+} was insufficient to improve the RCR or to
327 compensate for any negative effect that opening/closing the MPTP caused in the RCR that
328 only became evident after freezing (see Discussion).
329

Discussion

In this study we have shown that cryopreservation of liposomes by AfrLEA2 and AfrLEA3m was dependent on the lipid composition of the bilayer. Only liposomes mimicking the outer mitochondrial membrane were significantly stabilized by LEA proteins from *A. franciscana* during freezing and thawing. External trehalose significantly decreased damage to all liposomes tested. However, when trehalose was present on both sides of the lipid bilayer during freezing, as tested with IMM-like liposomes, the stabilization was much greater. Based on this result, we investigated whether isolated mitochondria provisioned with trehalose in the matrix would retain better respiratory function than mitochondria frozen with trehalose present only outside of the matrix. Isolated mitochondria maintained outer membrane integrity and exhibited improved respiratory function when frozen in a medium that contained 300 mM trehalose, compared to mitochondria in a KCl solution without trehalose. However, loading trehalose into the matrix (at least to the modest levels obtained by transient opening of the MPTP) did not improve the RCR after freeze-thawing compared to external trehalose alone.

LEA Proteins and the Stability of Frozen Liposomes

Freezing damage to liposomes with a lipid composition mimicking the OMM was substantially reduced by AfrLEA2 and AfrLEA3m. However, these LEA proteins did not have any effect on freeze-induced damage to IMM-like, ILPM-like, and PC liposomes, which indicated that protection is dependent on the lipid composition of the bilayer. This lack of stabilization of PC liposomes was unexpected because Tolleter et al. [52] report a ~15-25% decrease in CF leakage from PC liposomes frozen with PsLEAm at similar protein:lipid mass ratios. Additionally, COR15a and COR15b offset freeze-induced damage to chloroplast-like liposomes [50]. Liposomes were frozen at -18°C in the study by Tolleter et al. [52] and at -

20°C by Thalhammer et al. [50]. In our study liposomes were frozen at -80°C, which may have been less damaging. Indeed, our leakage values for control liposomes (without protectants) across all compositional types were similar to the smallest amount of leakage reported by Tolleter et al. [52] for liposomes protected with PsLEAm. Thus, in order to detect broader protection of liposomes by AfrLEA2 or AfrLEA3m, perhaps freezing conditions that promote greater damage are needed. It may also be that AfrLEA2 and AfrLEA3m are simply less effective stabilizers of liposomes during freezing than other LEA proteins, or that freezing at -80°C alters the nature of the final secondary structure gained by LEA proteins. Slow versus fast drying modifies the final percentages of α -helix versus β -sheet for plant LEA proteins for example [61].

Protection of Liposomes by Trehalose

In the present study, liposomes frozen with 250 mM trehalose retained significantly more CF than liposomes frozen without trehalose, regardless of the lipid composition of the liposomes. However, IMM-like liposomes were not protected as well compared to all others. Therefore, lipid composition can affect the ability of trehalose to stabilize lipid bilayers. It must be emphasized that most experiments herein were performed with trehalose only on the outside of liposomes. Many studies with trehalose have demonstrated that virtually no damage occurs to liposomes during freezing if at least 0.2 M trehalose is present on both sides of the lipid bilayer [e.g.,(3,19,46)]. However, liposomes with lipid compositions similar to the IMM of mammalian mitochondria were not previously investigated. Consequently, IMM-like liposomes were frozen with internal and external trehalose (Fig. 2B), and CF leakage was significantly reduced, which extends the established paradigm to this compositional type as well.

376 *Trehalose Loading Via Transient Opening of the MPTP*

377 The impact of transient pore opening on the respiratory function of mitochondria has
378 not been the primary focus of studies involving the MPTP. Often respiratory measurements
379 before and after pore opening in isolated mitochondria are not reported. Al-Nasser and
380 Crompton [2] found that mitochondria isolated from rat liver retained RCR values after loading
381 sucrose via transient opening of the MPTP comparable to those of non-permeabilized
382 mitochondria. However, respiration was measured in the presence of exogenous cyt *c*, and
383 therefore any damage to the outer membrane would not have been detected. Liu et al. [38]
384 demonstrated that mitochondria loaded with trehalose in the matrix via transient MPTP
385 formation retained a greater capacity to generate membrane potential after drying compared to
386 mitochondria dried without trehalose or with trehalose present only outside of the matrix.
387 However, RCR values were diminished from >5 before drying to less than 3 after rehydration.
388 Respiratory measurements were not taken after transient MPTP opening prior to drying, so it is
389 possible the OMM was compromised to some degree before desiccation. Nevertheless, matrix
390 trehalose does appear able to stabilize the inner mitochondrial membrane during drying based
391 on the results of Liu et al. [38]. Our results indicate that OXPHOS can be compromised by cyt
392 *c* loss if high Ca^{++} concentrations (100 μM) are used for pore opening.

393 *Stability of Isolated Mitochondria Frozen with Trehalose*

394 Mitochondria isolated from rat liver retain significant respiratory function when frozen
395 in 300 mM trehalose, similar to that reported for mouse mitochondria by Yamaguchi et al.
396 [63]. The fact that mammalian mitochondria exhibit some degree of coupling after freeze-
397 thawing is noteworthy. Mitochondria isolated from mouse liver also retain outer membrane

398 integrity when frozen in 300 mM trehalose medium [63], as shown here for rat liver
399 mitochondria by the cyt *c* test (Fig. 4B).

400 Surprisingly, loading trehalose into the matrix of mitochondria did not improve the
401 respiratory performance of the organelle after freeze-thawing. We speculate that the lower
402 RCR after freezing was largely due to compromised OXPHOS, although the apparent decrease
403 in the mean value did not reach statistical significance (unpaired t-tests, Holm-Sidak post-hoc,
404 $p > 0.05$, $n = 4-5$). In any case, the RCR result is possibly due to an insufficient amount of
405 trehalose loaded into the matrix. Transient opening of the MPTP was optimized to preserve
406 outer membrane integrity and respiratory function. As Figure 3 shows, there is a trade-off
407 between retention of these characteristics and the quantity of trehalose loaded into the matrix.
408 Stabilization of mammalian cells and mitochondria during freezing is dependent on the
409 concentration of trehalose [11,63]. Yamaguchi et al [63] reported that mitochondria isolated
410 from mouse liver retained outer membrane integrity when frozen in 300 mM trehalose but not
411 in 250 mM trehalose. Thus small differences in effective trehalose concentration in the matrix
412 could explain why stabilization is observed in some cases and not others, coupled with
413 differences in the type and severity of water stress imposed.

414 **Acknowledgements.** This work was supported by National Science Foundation grants IOS-
415 0920254 and IOS-1457061/IOS-1456809. We would like to thank Prof. Ramachandra
416 Devireddy, Department of Mechanical and Industrial Engineering at LSU, for his advice with
417 preparation of liposomes.

418

Literature Cited

- [1] Abazari A, Meimetis LG, Budin G, Bale SS, Weissleder R, Toner M (2015) Engineered trehalose permeable to mammalian cells. PLoS ONE 10(6): e0130323. doi:10.1371/journal.pone.0130323
- [2] Al-Nasser, I., and Crompton, M. (1986). The reversible Ca²⁺-induced permeabilization of rat liver mitochondria. Biochem J 239, 19-29.
- [3] Anchordoguy, T.J., Rudolph, A.S., Carpenter, J.F., and Crowe, J.H. (1987). Modes of interaction of cryoprotectants with membrane phospholipids during freezing. Cryobiol 24, 324-331.
- [4] Artus, N.N., Uemura, M., Steponkus, P.L., Gilmour, S.J., Lin, C.T., and Thomashow, M.F. (1996). Constitutive expression of the cold-regulated *Arabidopsis thaliana* COR15a gene affects both chloroplast and protoplast freezing tolerance. Proc Natl Acad Sci U S A 93, 13404-13409.
- [5] Atkin, O.K., and Macherel, D. (2009) The crucial role of plant mitochondria in orchestrating drought tolerance. Ann Bot 103 (4), 581-597.
- [6] Bernardi, P. (1996). The permeability transition pore. Control points of a cyclosporin A-sensitive mitochondrial channel involved in cell death. Biochim Biophys Acta 1275, 5-9.
- [7] Bernardi, P. (2013). The mitochondrial permeability transition pore: a mystery solved? Front Physiol 4, 95.
- [8] Bernardi, P., and Forte, M. (2007). The mitochondrial permeability transition pore. Novartis Found Symp 287, 157-164; discussion 164-159.
- [9] Bernardi, P., Krauskopf, A., Basso, E., Petronilli, V., Blachly-Dyson, E., Di Lisa, F., and Forte, M.A. (2006). The mitochondrial permeability transition from in vitro artifact to disease target. FEBS J 273, 2077-2099.
- [10] Boswell, L.C., Moore, D.S., and Hand, S.C. (2014). Quantification of cellular protein expression and molecular features of group 3 LEA proteins from embryos of *Artemia franciscana*. Cell Stress Chaperon 19, 329-341.
- [11] Buchanan, S.S., Gross, S.A., Acker, J.P., Toner, M., Carpenter, J.F., and Pyatt, D.W. (2004). Cryopreservation of stem cells using trehalose: evaluation of the method using a human hematopoietic cell line. Stem Cells Dev 13, 295-305.
- [12] Clegg, J. S. (1962). Free glycerol in dormant cysts of the brine shrimp *Artemia salina*, and its disappearance during development. Biol Bull, Woods Hole, 123, 295-301.
- [13] Clegg, J.S. (1965). Origin of trehalose and its significance during formation of encysted dormant embryos of *Artemia Salina*. Comp Biochem Physiol 14, 135- 143.

454 [14] Clegg, J.S. (1978). Interrelationships between water and cellular metabolism in
455 Artemia cysts. VIII Sorption isotherms and derived thermodynamic quantities. J Cell
456 Physiol 94, 123-137.

457 [15] Clegg, J.S. (2005). Desiccation tolerance in encysted embryos of the animal
458 extremophile, Artemia. Integr Comp Biol 45, 715-724.

459 [16] Clegg, J.S. (2011). Stress-related proteins compared in diapause and in activated,
460 anoxic encysted embryos of the animal extremophile, *Artemia franciscana*. J Insect
461 Physiol 57, 660-664.

462 [17] Crowe, J.H. (2011). Water, membranes, and life without water. In: Water: The
463 Forgotten Molecule. Le Bihan, D and Hidenao Fukuyama, H (eds.). Pan Stanford
464 Publishing Pte. Ltd., Singapore. pp. 205-232.

465 [18] Crowe, J.H., Carpenter, J.F., and Crowe, L.M. (1998). The role of vitrification in
466 anhydrobiosis. Annu Rev Physiol 60, 73-103.

467 [19] Crowe, J.H., Carpenter, J.F., Crowe, L.M., and Anchordoguy, T.J. (1990). Are freezing
468 and dehydration similar stress vectors? A comparison of modes of interaction of
469 stabilizing solutes with biomolecules. Cryobiol 27, 219-231.

470 [20] Crowe, J.H., Crowe, L.M., Carpenter, J.F., Prestrelski, S.J., Hoekstra, F.A., de Araujo,
471 P.S., and Panek, A.D. (1997). Anhydrobiosis: Cellular adaptation to extreme dehydration.
472 (Oxford: Oxford University Press).

473 [21] Crowe, J.H., Crowe, L.M., and Chapman, D. (1984). Preservation of membranes in
474 anhydrobiotic organisms: the role of trehalose. Science 223, 701-703.

475 [22] Crowe, J.H., Crowe, L.M., Oliver, A.E., Tsvetkova, N., Wolkers, W., and Tablin, F.
476 (2001). The trehalose myth revisited: introduction to a symposium on stabilization of
477 cells in the dry state. Cryobiol 43, 89-105.

478 [23] Crowe, J.H., Crowe, L.M., Wolkers, W.F., Oliver, A.E., Ma, X., Auh, J.H., Tang, M., Zhu, S.,
479 Norris, J., and Tablin, F. (2005). Stabilization of dry Mammalian cells: lessons from nature.
480 Integr Comp Biol 45, 810-820.

481 [24] Crowe, J.H., Hoekstra, F.A., and Crowe, L.M. (1992). Anhydrobiosis. Annu Rev Physiol
482 54, 579-599.

483 [25] Crowe, J.H., Spargo, B.J., and Crowe, L.M. (1987). Preservation of dry liposomes does
484 not require retention of residual water. Proc Natl Acad Sci U S A 84, 1537-1540.

485 [26] Crowe, L.M., and Crowe, J.H. (1992). Stabilization of dry liposomes by carbohydrates.
486 Dev Biol Stand 74, 285-294.

487 [27] Dutrieu, J. (1960). Observations biochimiques et physiologiques sur le development
488 d'*Artemia salina*. Leach Arch Zool Exp Gen 99,1-133.

489 [28] Giorgio, V., von Stockum, S., Antoniel, M., Fabbro, A., Fogolari, F., Forte, M., Glick, G.D.,
490 Petronilli, V., Zoratti, M., Szabo, I., et al. (2013). Dimers of mitochondrial ATP synthase
491 form the permeability transition pore. Proc Natl Acad Sci U S A 110, 5887-5892.

492 [29] Gnaiger, E., Kuznetsov, A.V.S., S., Seiler, R., Brandacher, G., Steurer, W., and
493 Margreiter, R. (2000). Mitochondria in the cold. In Life in the Cold. K.M. Heldmaier G, ed.
494 (Heidelberg, Berlin: Springer), pp. 431-442.

495 [30] Grelet J, Benamar A, Teyssier E, Avelange-Macherel MH, Grunwald D, Macherel D
496 (2005). Identification in pea seed mitochondria of a late embryogenesis abundant protein
497 able to protect enzymes from drying. Plant Physiol 137(1):157-167.

498 [31] Hand, S.C., and Carpenter, J.F. (1986). pH-Induced metabolic transitions in *Artemia*
499 embryos mediated by a novel hysteretic trehalase. Science 232, 1535-1537.

500 [32] Hand, S.C., and Hagedorn, M. (2008). New approaches for cell and animal
501 preservation: lessons from aquatic organisms. In Oceans and Human Health: Risks and
502 Remedies from the Seas. S.S. Walsh PJ, Fleming LE, Solo-Gabriele HM, Gerwick WH, ed.
503 (Academic Press), pp. 613-631.

504 [33] Hand, S.C., and Hardewig, I. (1996). Downregulation of cellular metabolism during
505 environmental stress: Mechanisms and implications. Annu Rev Physiol 58, 539-536.

506 [34] Hand, S.C., Jones, D., Menze, M.A., and Witt, T.L. (2007). Life without water:
507 expression of plant LEA genes by an anhydrobiotic arthropod. J Exp Zool A Ecol Genet
508 Physiol 307, 62-66.

509 [35] Hand, S.C., Menze, M.A., Toner, M., Boswell, L., and Moore, D. (2011). LEA proteins
510 during water stress: not just for plants anymore. Annu Rev Physiol, 73, 115-134.

511 [36] Hinch, D.K., and Crowe, J.H. (1998). Trehalose increases freeze-thaw damage in
512 liposomes containing chloroplast glycolipids. Cryobiol 36, 245-249.

513 [37] Hinch, D.K., Oliver, A.E., and Crowe, J.H. (1998). The effects of chloroplast lipids on
514 the stability of liposomes during freezing and drying. Biochim Biophys Acta 1368, 150-
515 160.

516 [38] Liu, X.H., Aksan, A., Menze, M.A., Hand, S.C., and Toner, M. (2005). Trehalose loading
517 through the mitochondrial permeability transition pore enhances desiccation tolerance
518 in rat liver mitochondria. Biochim Biophys Acta 1717, 21-26.

519 [39] Madin, K.A.C., and Crowe, J.H. (1975). Anhydrobiosis in nematodes- carbohydrate
520 and lipid-metabolism during dehydration. J Exp Zool 193, 335-342.

521 [40] Mannella, C.A. (1992). The 'ins' and 'outs' of mitochondrial membrane channels.
522 Trends Biochem Sci 17, 315-320.

523 [41] Menze, M.A., Boswell, L., Toner, M., and Hand, S.C. (2009). Occurrence of
524 Mitochondria-targeted Late Embryogenesis Abundant (LEA) Gene in Animals Increases
525 Organelle Resistance to Water Stress. J Biol Chem 284, 10714-10719.

526 [42] Moore, D.S., Hansen, R., and Hand, S.C. (2016) Liposomes with diverse compositions
527 are protected during desiccation by LEA proteins from *Artemia franciscana* and trehalose.
528 Biochim Biophys Acta 1858 (1): 104-115.

529 [43] Patil, Y.N., Marden, B., Brand, M.D., and Hand, S.C. (2013). Metabolic downregulation
530 and inhibition of carbohydrate catabolism during diapause in embryos of *Artemia*
531 *franciscana*. Physiol Biochem Zool 86, 106-118.

532 [44] Popova, A.V., Hundertmark, M., Seckler, R., and Hinch, D.K. (2011). Structural
533 transitions in the intrinsically disordered plant dehydration stress protein LEA7 upon
534 drying are modulated by the presence of membranes. Biochim Biophys Acta 1808, 1879-
535 1887.

536 [45] Pouchkina-Stantcheva, N.N., McGee, B.M., Boschetti, C., Toller, D., Chakraborty, S.,
537 Popova, A.V., Meersman, F., Macherel, D., Hinch, D.K., and Tunnacliffe, A. (2007).
538 Functional divergence of former alleles in an ancient asexual invertebrate. Science 318,
539 268-271.

540 [46] Rudolph, A.S., and Crowe, J.H. (1985). Membrane stabilization during freezing: the
541 role of two natural cryoprotectants, trehalose and proline. Cryobiol 22, 367-377.

542 [47] Steponkus, P.L., Uemura, M., Joseph, R.A., Gilmour, S.J., and Thomashow, M.F. (1998).
543 Mode of action of the COR15a gene on the freezing tolerance of *Arabidopsis thaliana*. Proc
544 Natl Acad Sci U S A 95, 14570-14575.

545 [48] Sun, W.Q., Leopold, A.C., Crowe, L.M., and Crowe, J.H. (1996). Stability of dry
546 liposomes in sugar glasses. Biophys J 70, 1769-1776.

547 [49] Tapia, H., and Koshland, D.E. (2014). Trehalose is a versatile and long-lived
548 chaperone for desiccation tolerance. Curr Biol 24, 2758-2766.

549 [50] Thalhammer, A., Bryant, G., Sulpice, R., and Hinch, D.K. (2014). Disordered Cold
550 Regulated15 proteins protect chloroplast membranes during freezing through binding
551 and folding, but do not stabilize chloroplast enzymes in vivo. Plant Physiol 166, 190-201.

552 [51] Thalhammer, A., Hundertmark, M., Popova, A.V., Seckler, R., and Hinch, D.K. (2010).
553 Interaction of two intrinsically disordered plant stress proteins (COR15A and COR15B)
554 with lipid membranes in the dry state. Biochim Biophys Acta 1798, 1812-1820.

555 [52] Tolleter, D., Hinch, D.K., and Macherel, D. (2010). A mitochondrial late
556 embryogenesis abundant protein stabilizes model membranes in the dry state. *Biochim*
557 *Biophys Acta* 1798, 1926-1933.

558 [53] Tolleter, D., Jaquinod, M., Mangavel, C., Passirani, C., Saulnier, P., Manon, S., Teyssier,
559 E., Payet, N., Avelange-Macherel, M.H., Macherel, D. (2007). Structure and function of a
560 mitochondrial late embryogenesis abundant protein are revealed by desiccation. *Plant*
561 *Cell* 19(5):1580-1589.

562 [54] Toxopeus, J., Warner, A.H., and MacRae, T.H. (2014). Group 1 LEA proteins
563 contribute to the desiccation and freeze tolerance of *Artemia franciscana* embryos during
564 diapause. *Cell Stress Chaperones* 19, 939-948.

565 [55] Tunnacliffe, A., and Wise, M.J. (2007). The continuing conundrum of the LEA
566 proteins. *Naturwissenschaften* 94, 791-812.

567 [56] Walther, D.M., and Rapaport, D. (2009). Biogenesis of mitochondrial outer
568 membrane proteins. *Biochim Biophys Acta* 1793, 42-51.

569 [57] Warner, A.H., Chakrabortee, S., Tunnacliffe, A., and Clegg, J.S. (2012). Complexity of
570 the heat-soluble LEA proteome in *Artemia* species. *Comp Biochem Physiol* 7, 260-267.

571 [58] Warner, A.H., Miroshnychenko, O., Kozarova, A., Vacratsis, P.O., MacRae, T.H., Kim, J.,
572 and Clegg, J.S. (2010). Evidence for multiple group 1 late embryogenesis abundant
573 proteins in encysted embryos of *Artemia* and their organelles. *J Biochem* 148, 581-592.

574 [59] Webb, S.J. (1965). *Bound Water in Biological Integrity*. (Springfield: Charles C
575 Thomas Publications).

576 [60] Weinstein, J.N., Blumenthal, R., and Klausner, R.D. (1986). Carboxyfluorescein
577 leakage assay for lipoprotein-liposome interaction. *Methods Enzymol* 128, 657-668.

578 [61] Wolkers, W.F., McCready, S., Brandt, W.F., Lindsey, G.G., and Hoekstra, F.A. (2001)
579 Isolation and characterization of a D-7 LEA protein from pollen that stabilizes glasses in
580 vitro. *Biochim Biophys Acta* 1544, 196-206.

581 [62] Wolkers, W.F., Walker, N.J., Tablin, F., and Crowe, J.H. (2001). Human platelets loaded
582 with trehalose survive freeze-drying. *Cryobiol* 42, 79-87.

583 [63] Yamaguchi, R., Andreyev, A., Murphy, A.N., Perkins, G.A., Ellisman, M.H., and
584 Newmeyer, D.D. (2007). Mitochondria frozen with trehalose retain a number of biological
585 functions and preserve outer membrane integrity. *Cell Death Differ* 14, 616-624.

586 [64] Yancey, P.H., Clark, M.E., Hand, S.C., Bowlus, R.D., and Somero, G.N. (1982). Living
587 with water stress: evolution of osmolyte systems. *Science* 217, 1214-1222.

588 [65] Zoratti, M., and Szabo, I. (1995). The mitochondrial permeability transition. *Biochim*
589 *Biophys Acta* 1241, 139-176.

Figure Legends

Figure 1. Leakage of carboxyfluorescein (CF) from liposomes frozen and thawed in the presence of LEA proteins and BSA. Lipid compositions simulated those of the inner mitochondrial membrane, the outer mitochondrial membrane, the inner leaflet of the plasma membrane, or a non-biological composition of 100 % phosphatidylcholine. Liposomes were frozen at the indicated protein:lipid mass ratios for AfrLEA3m (**A**), AfrLEA2 (**B**), or BSA (**C**). Data are expressed as means $\pm$ SD (n = 3-6 samples). Where error bars are absent, the SD was less than the size of the symbol. Asterisks indicate significantly less CF leakage than that of liposomes frozen without protectants (2-way ANOVA, Bonferroni multiple comparisons test, $p \leq 0.05$, n = 3-6).

Figure 2. (A) Cryoprotection by external trehalose of liposomes with different lipid compositions. Liposome stability was assessed by leakage of carboxyfluorescein (CF) Lipid compositions simulated those of the inner mitochondrial membrane (IMM), the outer mitochondrial membrane (OMM), the inner leaflet of the plasma membrane (ILPM), or a non-biological composition of 100 % phosphatidylcholine (PC). Data are expressed as means $\pm$ SD (n = 3-6 samples). Asterisks indicate significant decrease of CF loss compared to controls (1-way ANOVA, Tukey multiple comparisons test, $p < 0.05$). **(B)** IMM-like liposomes were prepared with and without 250 mM trehalose present on the inside of the liposomes and frozen in a 250 mM trehalose solution. Control liposomes were frozen without any trehalose. Stability of liposomes was determined by CF leakage. Data are expressed as means $\pm$ SD (n = 3-6 samples). An asterisk indicates significant decrease of CF loss compared to controls. Double asterisks indicate significant decrease of CF leakage compared to liposomes frozen

613 with external trehalose only (1-way ANOVA, Tukey multiple comparisons test, $p < 0.05$, $n = 3$ -
614 6).

615 **Figure 3.** Matrix trehalose content (**A**), stimulation of OXPHOS by exogenous cytochrome *c*
616 (**B**), and respiratory control ratios (RCR) (**C**) after transient opening of the MPTP for
617 mitochondria isolated from rat liver. Addition of 35 or 100 μM Ca^{2+} was used to induce MPTP
618 opening for 1 min prior to closing the pore by chelation of Ca^{2+} with 2 mM EGTA. Control
619 mitochondria were incubated under identical conditions as permeabilized mitochondria except
620 that calcium was omitted. As a test for integrity of the OMM, OXPHOS was measured before
621 and after addition of 4 mM cyt *c* and expressed as percent increase after the addition. An
622 asterisk indicates significant difference from control values, and double asterisks indicate
623 significant difference from values obtained with 35 μM Ca^{2+} (Student's unpaired t-test, $p < 0.05$,
624 $n = 3$ -6).

625 **Figure 4.** Respiratory measurements of mitochondria isolated from rat liver before and after
626 freezing. Mitochondria were frozen in media that contained 150 mM KCl (**A**) or 300 mM
627 trehalose (**B** and **C**). Trehalose was loaded into the matrix of mitochondria in (**C**) by transient
628 (1 min) opening of the permeability transition pore as promoted with 35 μM Ca^{2+} (see legend
629 to Fig. 3). Cytochrome *c* (+ cyt *c*) was added during the OXPHOS state and represents a test
630 for integrity of the OMM. Data represent the mean $\pm$ SD of 4-5 samples. Asterisks indicate a
631 significant change from non-frozen values and the double dagger symbol ($\ddagger$) indicates a
632 significant increase in OXPHOS respiration after the addition of cytochrome *c* (Multiple
633 unpaired t-test, Holm-Sidak post hoc, $p \leq 0.05$, $n=4$ -5).

634

Figure 1

Figure 2

Figure 3

Figure 4