Transport and reaction of iron and iron stable isotopes in glacial meltwaters on Svalbard near Kongsfjorden: From rivers to estuary to ocean.

Ruifeng Zhang^{a,*}, Seth G. John^b, Jing Zhang^a, Jingling Ren^c, Ying Wu^a, Zhuoyi Zhu^a, Sumei Liu^c, Xunchi Zhu^a, Chris M. Marsay^b, Fred Wenger^d

a State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062, China

b Department of Earth and Ocean Sciences, University of South Carolina, Columbia, SC 29208, USA

c Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education, Ocean University of China, Qingdao 266100, China

d Norwegian Water Resources and Energy Directorate, Middelthunsgate 29, Postboks 5091 Majorstua, 0301 Oslo, Norway

* Corresponding author.

Address: State Key Laboratory of Estuarine and Coastal Research, East China Normal University, Shanghai 200062, China

Telephone: +86-21-62233890

E-mail address: rfzhang@sklec.ecnu.edu.cn (R. Zhang).

ABSTRACT

1 Glacial meltwater has been suggested as a significant source of potentially
2 bioavailable iron to the oceans. However, the supply of dissolved iron (dFe) in glacial
3 meltwaters is poorly constrained as few sites have been studied, and because the
4 chemical processing of Fe during transport from glaciers to the adjacent coastal ocean
5 is not well understood. In order to better constrain glacial fluxes of dFe to the ocean,
6 iron concentrations, iron stable isotopes ($\delta^{56}\text{Fe}$), and other supporting chemical and
7 physical measurements were made along a ~4 km long glacial meltwater river on
8 Svalbard and in estuarine waters that it flows into. Dissolved iron concentrations in
9 the Bayelva River decreased from a maximum of 734 nM near the glacier to an
10 average value of 116 nM near the mouth of the river. Measurements in the
11 Kongsfjorden estuary suggest that 3 to 10 nM of dFe from the Bayelva River is
12 stabilized in glacial waters by the time it mixes into the ocean. Incubation of Bayelva
13 River waters over two weeks in both the light and dark show similar results, with the
14 majority of dFe being quickly precipitated and 4 to 7 nM Fe stabilized in the
15 dissolved phase. Evidence suggests that Fe is most likely lost from the dissolved
16 phase by aggregation and adsorption of nanoparticulate and colloidal Fe to particles.
17 Dissolved $\delta^{56}\text{Fe}$ was between -0.11‰ and +0.09‰ for all river samples and did not
18 vary systematically with dFe concentrations. We infer that the Fe is lost from the
19 dissolved phase by a process that fractionates Fe isotopes by less than 0.05‰,
20 indicating that the Fe bonding environment does not change during precipitation. This

is consistent with DOC loss that is much faster than predicted photo-oxidation rates, suggesting that DOC is also lost through adsorption and precipitation. Dissolved Fe concentrations in the Bayelva River (15-734 nM), and Fe concentrations which are stabilized in the dissolved phase (4-7 nM) are much lower than some previous estimates of Fe in glacial meltwaters, with roughly 80 % of dFe lost during transit in the Bayelva River and roughly 90% of the remaining dFe lost in the estuary. This may mean that glaciers are a less significant source of dissolved Fe to the global oceans than has been previously hypothesized, that cold base glaciers of the type studied here do not contribute significantly to the dissolved Fe flux, or that the flux of reactive particulate Fe to the oceans is more important than the dissolved flux. In Arctic regions with similar proglacial environments, bedrock composition, weathering intensity, and as precipitation of colloidal and nanoparticulate Fe may all play an important role in regulating the glacial meltwater iron flux to the ocean.

Keywords: Arctic; glacier; meltwater; dissolved iron; iron isotopes

36

37 **1. Introduction**

38 Despite the high abundance of Fe in the earth's crust (~5%), concentrations of Fe
39 in the modern oxidized oceans are very low (typically <1 nM in the open ocean) due
40 to the low solubility of Fe in seawater (Bruland and Lohan, 2003). Low Fe
41 concentrations, combined with physiological requirement for Fe by phytoplankton for
42 processes such as photosynthesis and nitrogen fixation, means that Fe is the limiting
43 nutrient for phytoplankton growth in much of the surface ocean (Moore et al., 2002).
44 The residence time of Fe in seawater is much shorter than the timescale of ocean
45 circulation (e.g. Moore et al 2004; Parekh et al., 2004), meaning that Fe
46 concentrations in the ocean are highly dependent on local sources and sinks of the
47 element. Deposition of dust transported from the continents has often been considered
48 the most important source of Fe to the remote oceans (Jickells et al., 2005), though
49 continental-margin sediments (Elrod et al., 2004; Lam and Bishop, 2008) and
50 hydrothermal vents (Saito et al., 2013; Tagliabue et al., 2010) may also represent
51 important sources of biologically available Fe. Recently, glacial Fe inputs in the form
52 of icebergs and glacial meltwater have been proposed as significant sources of
53 bioavailable iron to the ocean (e.g. Bhatia et al., 2013; Death et al., 2014; Smith et al.,
54 2007). As the rate of glacial melting increases due to climate change, this flux of
55 glacial Fe to the oceans may increase, potentially providing an important climate
56 feedback.

Glacial freshwater input to the oceans, in the form of glacial meltwater runoff and iceberg calving, delivers both dissolved Fe (dFe) and particulate Fe to the high-latitude oceans. Dissolved soluble Fe species are typically considered bioavailable, and very small Fe particles within the colloidal (0.02 – 0.45 μm) or nanoparticulate (0.02 – 1 μm) size range, which can be produced in the glacial environment, are also thought to provide bioavailable Fe because they are more soluble and reactive than larger particles (Raiswell, 2011). Similarly, free drifting icebergs may release iron to support primary production (Smith et al., 2007). These findings have implications for global productivity because Fe is a limiting nutrient throughout the Southern Ocean and large parts of the subarctic Pacific, and there is evidence that Fe may also limit the extent of seasonal nutrient drawdown in the high latitude North Atlantic (Achterberg et al., 2013; Boyd et al. 2007, Nielsdottir et al 2009). Recently, field observations have supported the idea that glacial Fe is bioavailable by showing that it supports a phytoplankton bloom along a glacier meltwater plume in the Amundsen Sea (Gerringa et al., 2012).

Despite the possible importance of glacial Fe, dissolved Fe supply from glacial meltwaters is poorly constrained due to the limited number of measurements and the great variability in dFe concentrations in these few samples. For example, concentrations of up to 9,950 nM were observed at a location in Greenland, leading to an estimated annual flux of biologically available Fe from Greenland meltwater which is comparable to dust-derived soluble iron inputs to the North Atlantic (Bhatia et al.,

2013). Just 100 km to the south in Greenland, however, Statham et al. (2008) observed glacial meltwater with an average dFe concentration of 52 nM. At a location 30 km further south at the Leverett Glacier, soluble iron ($<0.02\ \mu\text{m}$) concentrations in meltwater averaged 7 nM, however, iron concentrations in the colloidal size range ($0.02 - 0.45\ \mu\text{m}$) spanned an order of magnitude, from 232 nM to 4701 nM (Hawkings et al., 2014).

Determining the importance of glacial Fe inputs to the oceans also requires understanding the chemical processing of glacial Fe. Glacial meltwater runoff is characterized by a high physical to chemical erosion ratio (Raiswell et al., 2006), oxygen depletion in the subglacial environment (Hodson et al., 2002), and often by the presence of proglacial basins. Much of the glacial particulate Fe load is in the form of nanoparticulate Fe oxyhydroxides, which have been hypothesized to be directly or indirectly bioavailable, and which may be more easily solubilized in seawater (Raiswell et al., 2006; Raiswell et al., 2009; Raiswell, 2013). Indeed, glacial meltwaters in western Greenland were found having a much higher percentage of dFe in the colloidal fraction ($>98\%$) (Hawkings et al., 2014) than typical non-glacial rivers (e.g. Dai et al., 1995; Escoube et al., 2009; Stolpe et al., 2010). Glacial fed tributaries of the Copper River, Alaska, which were shown to have a higher sediment load than other types of tributaries, also had much higher colloidal Fe concentrations (Schroth et al., 2011).

Iron stable isotope ratios ($\delta^{56}\text{Fe}$) can be a valuable tool for tracking the

biogeochemical cycling of Fe in rivers and the marine environment, though they have so far been minimally applied to the study of glacial systems. The total observed range in riverine $\delta^{56}\text{Fe}$ is between -1.2 and +2.8 ‰, with variations dependent both on river chemistry and filter pore size (e.g. Ilina et al., 2013; Schroth et al., 2011). Continental weathering has been inferred to preferentially release light Fe into the riverine dissolved phase under some conditions (Fantle and DePaolo, 2004). However, the amount of dissolved organic material can also be important in determining $\delta^{56}\text{Fe}$. For example, $\delta^{56}\text{Fe}$ values of -0.2 ± 0.1 ‰ were measured for dFe in main channel sites of the Amazon River (Bergquist and Boyle, 2006), while isotopically heavy dFe (+0.30‰) was measured in waters of an organic-rich tributary. Positive values of dissolved $\delta^{56}\text{Fe}$ have also been reported for the North River, in the United States (Escoube et al., 2009), and some boreal rivers in Russia (Ilina et al., 2013). Recent work on a glacial river showed that the isotopic signature of dFe was found to be similar to crustal values in a glacier fed tributary of the Copper River (Schroth et al., 2011). Better constraints on the $\delta^{56}\text{Fe}$ of dissolved and particulate Fe in glacial meltwaters may make it possible to use Fe isotopes to trace the global distribution of glacial Fe in the oceans, as has recently been done for other Fe sources to the North Atlantic (Conway and John, 2014).

The aim of this study is to investigate the transportation and transformation of dFe between a glacial source and the ocean, via a glacial meltwater river. In addition to measurements of dFe concentration and $\delta^{56}\text{Fe}$, ancillary parameters including

dissolved oxygen (DO), dissolved Mn (dMn), dissolved organic carbon (DOC), total suspended matter (TSM), and pH were measured. These data are used to better understand the chemical cycling of glacial Fe in the Bayelva River, and to refine estimates of glacial Fe delivery to the ocean.

2. Materials and methods

2.1 Study area

Sampling for this study was performed in the vicinity of the Ny-Ålesund International Research Facility next to Kongsfjorden, Svalbard (Fig. 1). Svalbard is located in the southern portion of the Arctic Ocean, at the northern end of the Greenland Sea. The region is an important site of North Atlantic Deep Water (NADW) formation, warm North Atlantic waters travel along the western coast of Svalbard before sinking to form NADW (Jungclaus et al., 2005).

Fig. 1. Map of the study area, with the Bayelva River flowing from supraglacial terrain of the Austre Brøggerbreen glacier for ~4km through proglacial terrain before draining into Kongsfjorden. Red circles represent sampling sites along the Bayelva River. The hatched region represents the sampling area of Bayelva River estuary. Inset: regional map showing the location of the study area in the Svalbard Archipelago (blue square), with local circulation patterns.

About 60% of the land in the Svalbard archipelago is covered by glaciers, with estimated fluxes of meltwater and iceberg calving of $25 \pm 5 \text{ km}^3 \text{ a}^{-1}$ and $4 \pm 1 \text{ km}^3 \text{ a}^{-1}$, respectively (Hagen et al., 2003). The Bayelva River, which this study focuses on, is approximately 4 km long, with a catchment area of 32 km^2 , of which 50% is glaciated by the cold-based valley glaciers Austre and Vestre Brøggerbreen, both of which have thicknesses less than 100 m (Krawczyk et al., 2003). Although meltwater from

cold-based glaciers is not typically thought to have subglacial drainage flowpaths, the water chemistry in Bayelva River at the end of ablation season, when our sampling occurred, was found to be similar to that of subglacial meltwaters (Nowak and Hodson, 2014). The Bayelva River catchment is bounded by steep mountain ridges to the south and the Kongsfjorden fjord to the north. The annual discharge of the river is estimated to be $\sim 30 \cdot 10^6 \text{ m}^3$ (Hagen et al., 2003), with suspended sediment transport of between $5.1 \cdot 10^9$ to $22.8 \cdot 10^9 \text{ g}$ per year (Bogen and Bonsnes, 2003).

Sampling expeditions were carried out in August 2012 and 2013. Sampling locations along the Bayelva River were the HBR (head of Bayelva River) site, near the Austre Brøggerbreen glacier and $\sim 0.5 \text{ km}$ downstream from the visible supraglacial output, the MBR (middle-reach of Bayelva River) station located in the middle reach of Bayelva River, $\sim 1.8 \text{ km}$ from glacier, and the NVE (Norwegian Water Resources and Energy Directorate) station situated 2 km northwest of Ny-Ålesund and 300 m from the outlet of the Bayelva River into Kongsfjorden. All three locations were sampled at least once during both the 2012 and 2013 campaigns, and the NVE site was sampled 5 times during 2012. The Bayelva River estuary was sampled only during 2012.

2.2 Sample collection and handling

All plasticware, including high density polyethylene (HDPE) bottles (Nalgene) and Teflon PFA filtration assemblies (Savillex) were cleaned in a class 1000 clean lab

at East China Normal University by rinsing with ultrapure water (Milli-Q), soaking in 2% Citranox detergent for 24 hours, rinsing five to seven times with ultrapure water, leaching for 7 days in 10% HCl at room temperature or for 2 days in 10% HCl at 60 °C, and rinsing again five times with ultrapure water. In addition to the cleaning steps described above, PFA was boiled in 10% aqua regia for 30 min before Citranox and 10% HCl cleaning steps. HDPE bottles were filled with 0.1% (0.012 M) HCl using acid purified by quadruple distillation in a quartz still (QD HCl), and sealed in double bags until use. Filters (0.4 µm PC membranes, Whatman) were cleaned in 10% QD HCl at 60°C for 24 hours, rinsed ten times with ultrapure water, cleaned at 60 °C for another week in ultrapure water, then rinsed five times with ultrapure water.

Water samples were collected using a custom made “pole sampler” consisting of a PVC holder attached to the end of a fishing rod. This 2 m long rod made it possible to avoid contamination from the side of the ship when sampling in Kongsfjorden, and allowed samples to be taken from the middle of the main stream of the Bayelva River. To collect samples, a 1 L HDPE bottle was attached to the pole sampler, rinsed several times with the natural waters, and submerged until full. After sampling, bottles were sealed and stored in double plastic bags in a dark cool box until filtration in an onshore lab in Ny-Ålesund.

Samples were filtered through a 0.4 µm pore size membrane, meaning our samples included both soluble Fe (<0.02 µm) and colloidal Fe as typically defined operationally in the chemical oceanography literature (0.02-0.45 µm) (Wu et al.,

2001). These samples will also include a portion of the Fe occurring in the size range typically described as nanoparticulate (0.02 – 0.1 μm) (Raiswell and Canfield, 2012). Samples were filtered in a class 100 portable clean flow bench (Air Control). The filtration system was rinsed with copious quantities of 0.1% QD HCl, ultrapure water, and a small amount of sample, before filtering the bulk of the sample. Subsamples were collected in cleaned 30 mL or 500 mL HDPE bottles and acidified to pH 2 with QD HCl before storage.

2.3 Meltwater incubation experiment

Incubation of meltwater was carried out at the Ny-Ålesund research labs in order to determine how dFe changed over time and to investigate the role of light and DOC on dFe. Five liters of meltwater were collected at the NVE station by pole sampling from the main channel of the Bayelva River. These samples were filtered and homogenized, then 1 L was added to each of four 2 L acid-cleaned PTFE-lined plastic bags, sealed with a PTFE plastic rod sealer, and incubated on an exterior balcony. Two incubations were performed in the dark and another two incubations were exposed to natural sunlight, which was visible 24 hours a day. Samples were exposed to ambient temperature fluctuations (0-6 $^{\circ}\text{C}$). From each sample bag, 100 ml subsamples were taken on days 0, 2, 6, 8, and 13, filtered, and stored for analysis of dissolved Fe and DOC.

2.4 Fe concentration analyses

Dissolved iron concentrations were analyzed by ICPMS after single-batch extraction onto a nitrilotriacetate resin (NTA, Qiagen) according to the procedures of Lee et al. (2011). Briefly, samples were spiked with an enriched ^{54}Fe solution, oxidized with H_2O_2 , and extracted onto ~100 clean NTA resin beads. After separating Fe from other ions on the NTA resin, samples were reconstituted in 0.5 M HNO_3 for analysis on a multi-collector ICPMS (Neptune, Thermo) at East China Normal University. The Neptune was operated in high-resolution mode with a minimum resolution of 6000 in order to separate ^{54}Fe and ^{56}Fe beams from $^{40}\text{Ar}^{14}\text{N}$ and $^{40}\text{Ar}^{16}\text{O}$ interferences, respectively. Three replicates from each 30 ml subsample were prepared for analysis, and in rare cases where the three replicates did not agree another 3 replicates were prepared. The typical procedural blank was 0.15 nM, with a detection limit of ~0.12 nM (based on three-times the standard deviation of the procedural blank). Accuracy of the method was tested by analysis of three SAFe open ocean intercalibration samples, and our measurements compared favorably with the consensus values (provided by Ken Bruland, April, 2013). Our measured Fe concentrations for SAFe S, SAFe D1 and SAFe D2 were 0.094 ± 0.019 , 0.68 ± 0.07 and 0.94 ± 0.07 nM compared to consensus values of 0.094 ± 0.008 , 0.70 ± 0.03 and 0.95 ± 0.02 nM, respectively. Measurements of two high concentration certified reference materials (both NRC, Canada) also compared favorably: estuarine water SLEW-3 (10.4 ± 0.2 vs 10.2 ± 1.2 nM) and river water SLRS-5 (1.25 ± 0.10 vs $1.62 \pm$

0.12 μM). We suspect that the relatively lower accuracy of our technique for samples with higher Fe concentrations is due to low spike/sample ratios. Therefore, Fe concentrations for meltwater samples with high concentrations of dFe were obtained from double-spike Fe isotope analyses (see below).

2.5 Fe isotope analysis

Iron stable isotope analyses were carried out using a double-spike technique according to previously published methods (Conway et al., 2013). Briefly, samples were spiked with an iron double spike consisting of roughly equal amounts of ^{57}Fe and ^{58}Fe at a sample:spike ratio of 1:2. Samples were then dried down in clean PFA vials, reacted with 200 μL each of clean concentrated HNO_3 and H_2O_2 (Aristar, VWR) to dissolve any organic material, and dried down again. Samples were then purified by anion exchange chromatography according to methods developed for marine particulate Fe (Revels et al., in press). Samples were first reconstituted in 0.2 mL 10 M HCl and 0.001 M H_2O_2 , then loaded onto 135 μL AG-MP1 resin. After rinsing with 0.5 mL 10 M HCl and 0.001 M H_2O_2 , iron was eluted from the resin with 0.8 mL 1 M HCl. The eluent was then dried down and reconstituted in 2 mL 0.1 M HNO_3 for isotopic analysis on a Neptune multi-collector ICPMS at the University of South Carolina Center for Elemental Mass Spectrometry. In addition to stable isotope ($\delta^{56}\text{Fe}$) data, the double spike method also allowed for the calculation of ultra-precise Fe concentration data for all Bayelva River samples.

251

252 2.6 Analysis of other physical and chemical parameters

253 Temperature, pH, dissolved oxygen (DO), and salinity were measured in situ with
254 a WTW multiline F/sets3 multi-parameter probe. Total suspended material (TSM)
255 was determined by filtering onto pre-weighed Whatman GF/F filters, which were then
256 freeze dried and reweighed. Dissolved organic carbon (DOC) was analyzed by the
257 high-temperature catalytic oxidation (HTCO) method with a Shimadzu TOC analyzer
258 (Model TOC-CPH) according to the methods of Wu et al. (2013). The method used
259 for determination of dissolved manganese (dMn) is modified from the methods of
260 Resing and Mottl (1992), which determined dMn concentrations by leucomalachite
261 green-sodium periodate catalytic kinetic spectrophotometry (Wang et al., 2015).

262

263 3. Results

264

265 3.1 Bayelva River

266 3.1.1 Bayelva River transect

267 During both the 2012 and 2013 field seasons, samples were taken from three
268 stations along the Bayelva River on the same day (Fig. 2). In both years, the highest
269 dFe concentrations were observed in samples taken closest to the glacier output (734
270 nM in 2012 and 401 nM in 2013) with decreasing concentrations downstream. During
271 2012, dFe was approximately twice as high at all three stations compared to 2013,

though the rate of dFe loss was similar both years with a 71% decrease in dFe between HBR and NVE stations in 2012 and an 84% reduction in dFe between the same stations in 2013. Additionally, dFe was measured in two samples taken from tributaries to the Bayelva River in 2012, with 226 nM in a supraglacial tributary and 157 nM in a tributary near NVE station.

A variety of other physical and chemical parameters were also measured along the transect. Little variation of pH was observed, with an average pH of 8.23 ± 0.41 in 2012 and 8.27 ± 0.68 in 2013. Dissolved oxygen (DO) increased slightly with distance from the glacier during both years, from 12.6 mg L^{-1} at HBR to a supersaturated value of 20.1 mg L^{-1} at NVE in 2012, and an increase from 12.5 mg L^{-1} to 15.3 mg L^{-1} at the same stations in 2013. DOC and dMn both decreased away from the glacier in 2012, from 275 nM to 153 nM for dMn and from $167 \text{ }\mu\text{M}$ to $98 \text{ }\mu\text{M}$ for DOC. TSM increased 17% from 389 mg L^{-1} at station HBR to 455 mg L^{-1} at station MBR, then decreased to a value of 115 mg L^{-1} at station NVE. Dissolved $\delta^{56}\text{Fe}$ did not vary in any systematic fashion along the transect, with all values between -0.11‰ and 0.09‰ .

Fig. 2. Dissolved Fe (dFe) concentrations, $\delta^{56}\text{Fe}$ values, and ancillary parameters including pH, dissolved Mn (dMn), dissolved oxygen (DO), dissolved organic carbon (DOC), and total suspended matter (TSM) in the Bayelva River. Open circles represent data from 2012 and open squares represent data from 2013. Dashed grey lines connect data from samples taken during the same day.

3.1.2 Bayelva River temporal variability

Over twelve days in August 2012, the same parameters which had been measured on the transect were monitored at the NVE station (Fig. 2). Over this time, discharge rates varied from $0.8 \text{ m}^3 \text{ s}^{-1}$ to $6.4 \text{ m}^3 \text{ s}^{-1}$ (Fig. 3). Dissolved Fe concentration varied from 15 nM to 213 nM, with an average value of $116 \pm 84 \text{ nM}$ ($n=5$). These dFe values were similar to the range observed at this station during the transect sampling, and all values were lower than dFe measured closer to the glacier, at station HBR. No clear relationship was found between dFe and discharge at the NVE station.

Fig. 3. Dissolved iron concentrations (open circles) and discharge (grey line) monitored at NVE station on the Bayelva River in August 2012. Sampling date for each dFe measurement is given beside dFe data symbols.

3.2 Kongsfjorden estuary

Dissolved Fe concentrations were measured at a variety of salinities in the Bayelva River estuary where fresh glacial waters mix with saline coastal seawater

(Fig. 4). Assuming that the average dFe concentration measured at the NVE station, ~300 m upstream from Kongsfjorden, represents the freshwater end-member (116 nM), there is a significant loss of Fe immediately upon mixing of fresh and saline waters, with dFe of 27.4 nM at a salinity of 1.5, dFe of 11.3 nM at a salinity of 6.9, and an average dFe of 4.7 ± 1.6 nM ($n=7$) at salinities $\sim >10$.

Fig. 4. Dissolved Fe (dFe) concentrations compared to salinity in Bayelva River estuary (triangles) and in the Bayelva River roughly 300 m upstream of the estuary (circles).

3.3 Freshwater incubations

Incubations of Bayelva River waters were performed for two weeks in both natural light conditions and in the dark, during which time dFe and DOC concentrations were monitored (Fig. 5). dFe in filtered waters from the NVE station at the beginning of the experiment (incubation day 0) was 15 nM. This decreased sharply to ~7 nM by day 2 in both dark and light incubations. Subsequently, dFe concentrations in the dark incubations remained relatively constant, decreasing only ~1 nM over the next 12 days of the incubation experiment, with differences between

the two replicate dark incubations within analytical error. In contrast, dFe in the light incubations decreased further, to about 5 nM by the end of the experiment. DOC also decreased dramatically within the first 2 days of the experiment from 75 μ M at day 0 to ~15 μ M at day 2. Between days 2 and 12 there did not appear to be much further loss of DOC, though it is possible that samples were contaminated during filtration, especially considering the low DOC present. Nonetheless, the ~75% loss of DOC at the beginning of the experiment was observed in all four samples and therefore appears robust.

Fig. 5. Dissolved Fe and DOC concentrations in light and dark incubations in duplicates. Open circles are bags incubated in daylight, while solid circles represent bags kept in the dark.

4. Discussion

A primary aim of this study is to investigate the transportation and transformation of dFe between a glacial source and the ocean, via a glacial meltwater river, and to evaluate the importance of glacial meltwaters as a source of Fe to the oceans. To do so requires both a quantitative measure of the amount of dissolved Fe which is

344 stabilized in glacial meltwaters, where it can ultimately be delivered to the oceans,
345 and an understanding of the chemical transformations by which Fe is lost from glacial
346 waters. Below we discuss both our observations about Fe concentrations in the
347 Bayelva River and estuary, and hypothesize about the processes responsible for Fe
348 precipitation or stabilization.

350 4.1 Sources of water and Fe to the Bayelva River

351 Though we regard the Bayelva River as being mostly closed to new inputs along
352 its course to the ocean, there are likely small inputs from groundwater and tributary
353 streams. For example, Mn concentrations were ~8% higher at the MBR station
354 compared to HBR, suggesting either rapid timescale variability in Mn output from the
355 glacier or input from hyporheic groundwater (Green et al., 2005). Inputs from longer
356 residence-time groundwater to the Bayelva are not thought to be significant (Nowak
357 and Hodson, 2013). During the sampling period, we did not observe any large
358 tributaries to the main stream of Bayelva River. dFe concentrations in the two
359 tributaries which we measured (~200 nM) were not low enough to account for the loss
360 of Fe along the path of the Bayelva by dilution alone. Discharge data from many
361 different years has suggested that meltwater from Austre Brøggerbreen contributes
362 >50% of total discharge in Bayelva River (Hodson et al., 1998). We therefore
363 conclude that while lateral inputs of both water and Fe to the Bayelva River may
364 occur, they are not a dominant factor in dFe and cannot explain the dramatic dFe

decrease observed downstream in the Bayelva River.

4.2 Rates of Fe precipitation from Fe in glacial meltwaters

Several lines of evidence suggest that most dissolved Fe is quickly precipitated within hours after emerging from the glacier, but that a small fraction (4-7 nM) is present in a chemical form that is stable over a period of days to weeks. In both 2012 and 2013, Fe concentrations decreased markedly downstream in the Bayelva River (Fig. 2). Estimates of river velocity suggest that water travels the entire ~4 km from the glacier to Kongsfjorden in 1 to 3 hours (Hodson et al., 2005), suggesting that the half-life for dFe in the river is about an hour or less. While most of the dFe was precipitated from river waters during this short transect, significant quantities remained even in the samples taken downstream at the NVE station (15-213 nM). Considering that our first sampling site located at the highest accessible location was still ~0.5 km from the glacier output, our measurements of dFe and our estimates of Fe loss in the proglacial environment represent likely minimum values.

Incubation experiments also suggest that a small portion of glacial meltwater Fe is present in a stable dissolved form, regardless of whether samples are exposed to sunlight or not. When Bayelva River waters were incubated, a rapid loss of Fe was observed over the first two days of the experiment. In the dark incubations, very little additional Fe was lost from the dissolved phase. The rate of Fe loss from the dissolved phase in the light incubations was also quite slow from days 2 to 12, compared to rates of Fe loss in the Bayelva River and during the first two days of incubation. The

slight loss of Fe in the light incubations is consistent with the work of Shiller et al. (2006), who found that photooxidization could destroy Fe binding organic ligands leading to precipitation of Fe(III). We cannot rule out the possibility that the rapid loss of dFe over the first two days was due to Fe adsorption to the PTFE bag lining, though PTFE is not very surface reactive to either inorganic Fe oxyhydroxides or organically bound Fe. Even if a significant amount of Fe was lost by adsorbing to the bag walls, rather than by precipitation and removal during filtration, this may still reflect a large pool of surface-reactive Fe in the Bayelva River that would be lost either by precipitation or adsorption during transit or during flocculation upon mixing with the estuary. The relative surface area in the incubation bags is not much different from the surface area exposed within the stream bed. Thus, we expect that $\sim 4 - 7$ nM of Fe which remained dissolved in the incubation experiments would also remain dissolved under natural conditions.

A similar pattern is observed in the Kongsfjorden estuary, with $\sim 90\%$ of the dissolved Fe present at the NVE station in the Bayelva River lost during the early stages of estuarine mixing (based on an average ~ 120 nM dFe at the NVE station). This amount of loss is similar to other, non-glacial river systems and mixing experiments, where much of the riverine Fe is lost as the ionic strength of the waters increase (Bergquist and Boyle, 2006; Buck et al., 2007; Dai et al., 1995; Krachler et al., 2005; Laglera and van den Berg, 2009). Fe concentrations in the estuary were not significantly different between salinities of 7 to 11 (7.3 ± 3.9 nM, $n=3$) compared to

salinities of 19 to 27 (4.4 ± 1.5 nM, $n=5$). This suggests mixing of brackish water ($S = 7-11$) with seawater ($S=34$), both of which had Fe concentrations of about 5 nM, with little loss or addition of Fe during mixing. During 2012 an endmember seawater sample was collected from the mouth of the Kongsfjorden bay, where it enters the Arctic Ocean, which contained ~ 3 nM dFe at a salinity of 34 (data not shown).

Considering together all incubation samples after 2 days of incubation, and all estuarine samples at salinities $\sim >10$, we suggest that between 4 nM and 7 nM of glacial Fe in the Bayelva River is stabilized in the dissolved phase (5.8 ± 1.4 nM, $n=20$). This represents roughly 1% of the Fe concentrations measured near to the glacial outflow.

4.3 Mechanism of Fe precipitation

4.3.1 Chemical form of Fe in meltwaters

The high dissolved Fe concentrations observed near the glacial outlet may be either truly dissolved chemical species (<0.02 μm), or small particles which fall in the size range typically described as colloidal or nanoparticulate. One possible mechanism by which Fe may dissolve underneath the glacier or within the moraines of the glacial forefield is by the formation of reduced, soluble Fe(II) from sulphide oxidation, weathering, or microbial respiration (Wadham et al., 2013). Because Austre Brøggerbreen is a cold-based glacier there will be a minimum of delayed subglacial meltwater, though even in cold-based glaciers there can be temperate ice

and thin films of water at the ice-rock interface (Raiswell and Canfield, 2012). Observations of water chemistry in Bayelva River have suggested the meltwater does have a subglacial signature during the late ablation season (Nowak and Hodson, 2014). The oxidation half-life of Fe(II) in natural fresh water is typically seconds to minutes (Davison and Seed, 1983), however Fe (II) may be stabilized by complexation with organic ligands (Statham et al., 2012). While we do not observe consistently lower dissolved oxygen concentrations near the glacier, we do observe high dissolved Mn concentrations (153-296 nM) that persist for 4km downstream of the glacier output (Fig. 2). Reduced Mn(II), like Fe(II), is easily water soluble while oxidized Mn(III) is particle reactive, however the oxidation half-life of Mn is significantly longer than for Fe (Morgan, 2005). Higher dissolved Mn concentrations near the glacier may therefore be indicative of reducing conditions in subglacial waters. Dissolved Fe in meltwaters could be comprised either of dissolved Fe(II), or of nanoparticulate Fe(III) minerals formed after oxidation of the river waters. Alternatively, iron in the dissolved fraction may be present as nanoparticulate minerals, such as the schwertmannite, which can form directly under glaciers when oxic waters weather Fe sulfides (Raiswell et al., 2009).

4.3.2 Mechanism of Fe loss

The study of Fe isotopes can help to constrain the mechanism by which Fe is lost from the dissolved phase. The average $\delta^{56}\text{Fe}$ was statistically equivalent at stations

HBR, MBR and NVE, $0.01 \pm 0.03\text{‰}$, $-0.04 \pm 0.10\text{‰}$ and $0.02 \pm 0.02\text{‰}$, respectively, and

there was no apparent relationship between dFe and $\delta^{56}\text{Fe}$ in the Bayelva River.

Assuming that the Bayelva River is a closed system, the relationship between dFe and

$\delta^{56}\text{Fe}$ can be modeled by Rayleigh distillation as:

$$\delta^{56}\text{Fe}_{\text{river}} = \delta^{56}\text{Fe}_{\text{glacier}} + \Delta\delta^{56}\text{Fe}_{\text{precipitate-dissolved}} \cdot \ln(F) \quad (1)$$

where $\delta^{56}\text{Fe}_{\text{river}}$ is the Fe isotope composition measured in the Bayelva River,

$\delta^{56}\text{Fe}_{\text{glacier}}$ is the dissolved $\delta^{56}\text{Fe}$ at the glacier output, $\Delta\delta^{56}\text{Fe}_{\text{precipitate-dissolved}}$ is the

isotope effect of Fe precipitation, and F is the fraction of Fe remaining in the

dissolved phase. Doing so, we find that all of the $\delta^{56}\text{Fe}$ data can be fit with Rayleigh

distillation curves assuming that the process by which Fe precipitates fractionates Fe

isotopes by less than -0.05 to $+0.05\text{‰}$ from a starting dFe of 700 nM and a starting

$\delta^{56}\text{Fe}_{\text{glacier}}$ of -0.02‰ (Fig. 6). Dissolved $\delta^{56}\text{Fe}$ is also similar to the crustal average of

around 0.07‰ (Poitrasson, 2006), suggesting that Fe isotopes were not greatly

fractionated during dissolution under the glacier before entering the Bayelva River.

Fig. 6. $\delta^{56}\text{Fe}$ versus dFe concentration. Open circles represent data from 2012, while open squares are data from 2013. Error bars show the 2σ standard analytical errors. Grey line represents the Rayleigh distillation curves for closed system isotope fractionation from an initial glacial outflow with 700 nM dFe and $\delta^{56}\text{Fe} = -0.02\text{‰}$.

In many natural environments, Fe precipitation occurs by processes that are associated with large Fe isotope fractionations, a pattern that is inconsistent with Bayelva River $\delta^{56}\text{Fe}$ measurements. For example, Fe(II) is roughly 3 ‰ lighter than Fe(III) at isotopic equilibrium (Johnson et al., 2002; Welch et al. 2003), and environments where Fe(II) and Fe(III) are both present are generally characterized by large variations in $\delta^{56}\text{Fe}$ of up to several permil. Examples include reducing sediment porewaters (Severmann et al., 2006; Severmann, et al., 2010), hydrothermal vents (Bennett et al., 2009; Severmann et al., 2004), and submarine groundwaters (Rouxel et al., 2008), all of which exhibit $\delta^{56}\text{Fe}$ at least several permil different from the crustal average. Non-redox chemical changes in Fe speciation are also typically associated with large kinetic isotope effects. For example, a series of inorganic Fe(III) precipitation experiments exhibited isotopic fractionations from -0.22 to -2.12 ‰ with an average $\delta^{56}\text{Fe}$ of -0.95 ‰ (Balci et al., 2006). Kinetic isotope effects associated with the binding and unbinding of Fe to organics have not been well studied, but organic complexation of Fe by the organic siderophore DFB leads to an equilibrium isotope offset of 0.6 ‰ fractionation in $\delta^{56}\text{Fe}$ compared to inorganic Fe, suggesting

that the kinetic isotope effects associated with binding and unbinding may be even larger (Dideriksen et al., 2008). We cannot rule out the possibility that Fe precipitation in the Bayelva River involves a change in Fe speciation associated with a very small kinetic or equilibrium isotope effect. However, we consider it more likely that Fe speciation does not change during precipitation.

Two processes by which Fe could precipitate without any change in Fe bonding environment (speciation) are the adsorption of organically bound Fe to particles, and the aggregation of nanoparticulate/colloidal Fe. Complexation of Fe by organic ligands is common in natural environments, and such complexes are known to keep Fe(III) dissolved in solution (Rue and Bruland, 1995). Both dFe and DOC are present at much higher concentrations close to the glacier, and concentrations of both decrease downstream (Fig. 2, Fig. 7). However, given a DOC photodegradation rate of $\sim 0.25 \mu\text{M h}^{-1}$ (Mostofa et al., 2007; Shiller et al., 2006) and the short water travel time (1-3 h, Hodson et al., 2005), there should be minimal loss of DOC by photodegradation in the Bayelva River. Instead we presume that the DOC is lost predominantly by adsorption onto particulates. This is supported by the finding that particulate organic carbon increased by $\sim 80\%$ during the transit from station HBR to station NVE (Zhu et al., in prep), suggesting that DOC loss and dFe precipitation may occur simultaneously. The relationship between DOC and dFe is not entirely linear however, suggesting either that the organics which bind Fe are precipitated more quickly than bulk DOC, or that dissolved Fe is not bound to organics. Alternatively,

much of the dissolved Fe may be present as nanoparticulate/colloidal minerals. Operationally, colloidal materials are those in the 0.02 to 0.4 μm size range, meaning that they can pass through a filter to be included in the dissolved phase but still have relatively high molecular weights compared to most “truly dissolved” species. Chemically, colloids are large enough that atoms on their interior do not chemically interact with the environment. Thus, aggregation and precipitation of nanoparticulate/colloidal Fe is a pathway by which Fe could leave the dissolved phase without any significant change in Fe bonding environment. Considering the lack of isotopic fractionation during Fe precipitation in the Bayelva River, we propose that precipitation of nanoparticulate/colloidal Fe is the most likely pathway for the loss of dissolved Fe. The suggestion that much of the dFe in Bayelva River meltwaters occurs as nanoparticles/colloids matches similar findings from Greenland (Hawkings et al., 2014) and Alaska (Schroth et al., 2011).

Fig. 7. A. dFe versus DOC in Bayelva River. B. dFe to DOC ratio versus distance from glacier output.

4.4 Glacial Fe contribution to the global marine Fe budget

The dissolved Fe concentrations measured here are on the low end of the global average. The total range for glacial waters reported from previous studies is from a minimum of 20 nM to a maximum of 9,950 nM, with an average concentration of $\sim 1000 \pm 1600$ nM for all published meltwater concentrations ($n=110$) (Bhatia et al., 2013; Green et al., 2005; Hawkings et al., 2014; Statham et al., 2008). The average dFe concentration for all glacial river samples from this study was 233 ± 218 nM ($n=10$), which is at the lower end of the global range. Some of this variability may be driven by differences in sampling protocol. The highest previously reported Fe concentrations were for samples filtered directly in the field (Bhatia et al., 2013), while lower Fe concentrations were observed by Statham et al. (2008) and in this study, where filtration occurred in the lab several hours after sampling. However, no observable precipitate observed during filtration of samples in this study. Also, if such rapid precipitation were occurring it would be consistent with our overall finding that the half-life for dFe in glacial meltwater is quite short.

Crucially for estimating the flux of glacial Fe to seawater, we hypothesize that dissolved Fe in the Bayelva River occurs in two very different forms. Most of the dissolved Fe measured near the glacier outflow is precipitated within hours to days, and does not enter the Kongsfjorden estuary where it might ultimately mix into the North Atlantic. We propose that this Fe is predominantly Fe nanoparticles/colloids. A second species of Fe, present at concentrations of 5 – 10 nM appears to be stable on timescales of days to weeks. We propose that this portion of Fe is organically

547 complexed. Thus, when attempting to calculate glacial Fe fluxes to the ocean we
548 suggest that total dissolved Fe concentrations may be a much less important parameter
549 than the concentration of Fe which is stabilized in the dissolved phase for long enough
550 to be mixed into the coastal ocean.

551 In contrast to previous studies, extrapolating the lower dFe concentrations found
552 in the Bayelva River to global glacial meltwaters would not support glaciers as being
553 an important source of Fe to the North Atlantic. Based on dFe concentrations of 3.7
554 μM in glacial meltwaters measured in Greenland, and assuming a 90% loss of dFe
555 during estuarine removal processes, Bhatia et al. (2013) extrapolate to the 0.5 Tm^3/y
556 Greenland Ice Sheet runoff volume and predict an annual flux of $\sim 0.01 \text{ Tg}$ of
557 dissolved iron from the Greenland ice sheet to the North Atlantic. In contrast,
558 applying the same calculations to our average Bayelva River dFe concentration of 223
559 nM, we would predict fluxes more than an order of magnitude lower ($6 \cdot 10^{-4} \text{ Tg/y}$).
560 Similarly, applying our 223 nM to the global glacial meltwater discharge of 1.4 Tm^3/y
561 (Raiswell et al., 2006), we would calculate a global dFe input (before accounting for
562 estuarine removal processes) of 0.017 Tg/y , which is insignificant compared to other
563 global Fe fluxes, calculated to be of the order of $\sim 1\text{-}100 \text{ Tg/y}$ (Jickells et al., 2005).
564 Furthermore, the decrease in dFe concentration between our mean glacial runoff
565 samples (223 nM) and concentrations measured in mid salinity estuarine waters (a
566 mean of 4.4 nM over a salinity range of 19 to 27) suggests that the assumed 90%
567 attenuation of dFe due to estuarine removal processes may be an underestimate. If we

were to extrapolate from the ‘stabilized dFe’ (Fe which is assumed to remain in the dissolved phase following the increase in salinity as it is delivered to the ocean) concentrations of 4-7 nM from the Bayelva River to Greenland, we would estimate a glacial flux of dFe from the Greenland Ice Sheet of between $1.4 \cdot 10^{-4}$ and $2.8 \cdot 10^{-4}$ Tg/y, which is much smaller than the 0.07 - 0.7 Tg/y soluble Fe input estimated from dust deposition models (Boyd et al., 2010, Fan et al., 2006).

However, there are several reasons why the dFe measurements from the Bayelva River in Svalbard may not be globally representative. First, The Bayelva River originates from a cold-base glacier where the subglacial processes may be very different from larger ice sheets in Greenland and Antarctica (Nowak and Hodson, 2014). Second, there is large variability in glacial bedrock types, catchment size, and meltwater residence time, and these factors can affect Fe concentration and speciation in meltwaters (e.g. Hawkings et al., 2014). Third, we find that processing in the proglacial environment has a large impact on Fe concentrations and speciation, and considering that our first sample was taken ~500 m from the outflow of the glacier there might be much higher dFe in anoxic englacial meltwaters, which in Antarctica and Greenland can be delivered directly into the ocean providing both soluble and reactive nanoparticulate Fe (Raiswell and Canfield, 2012). Finally, even though we observed precipitation of dissolved Fe, the Bayelva River may still be delivering high quantities of reactive particulate Fe to the ocean. Phytoplankton may acquire Fe directly from such particles to support their growth (e.g. Thuróczy et al., 2012).

Alternatively, reactive particulate Fe can be dissolved from within the sediments to contribute Fe to the overlying water column (Wehrmann et al., 2014). While our data add significantly to the global dataset on glacial Fe delivery to the oceans, and provide some of the first measurements of Fe delivery from cold-based glaciers, studies with greater spatial and temporal coverage will be needed to fully constrain the possible importance of glacial Fe to the oceans.

While the concentration measurements made here cannot directly support glaciers as a significant source of Fe to the North Atlantic, it is intriguing to note that the $\delta^{56}\text{Fe}$ data may be consistent with a large glacial input. Measurements of seawater dissolved $\delta^{56}\text{Fe}$ from the North Atlantic show a distinct isotope signature in Upper Labrador Sea Water (ULSW) near 0 to +0.2 ‰ (Conway and John, 2014). This Fe isotope signature was originally proposed to result from the interaction of ULSW with resuspended oxic seafloor sediments. However this $\delta^{56}\text{Fe}$ also matches our measurements of glacial $\delta^{56}\text{Fe}$, suggesting that glacial meltwater input may be contributing to North Atlantic dFe.

5. Conclusions

Our data represent the first study of the transport of dissolved Fe in meltwaters all the way from a glacier through a marine estuary (Fig. 8). In this study, a significant loss of dFe in a glacial meltwater runoff was observed during transport through the proglacial environment. In the Bayelva River, roughly ~80% of Fe was lost between a

station 0.5 km from the glacial outlet and a station 3.2 km further downstream. A further ~90% of the remaining dFe was lost in the Kongsfjorden estuary. Overall, only ~2% of the Fe output from the Austre Brøggerbreen Glacier was stabilized in the estuary where it could eventually mix into the North Atlantic, excluding the possible redissolution of particulate Fe within estuarine sediments. Based on analyses of various physical and chemical parameters, especially DOC and $\delta^{56}\text{Fe}$, we propose that most of the dissolved Fe in the Bayelva River near the glacial output is in a colloidal/nanoparticulate phase which rapidly precipitates from solution, while roughly 4 -7 nM Fe is stabilized by complexation with organic ligands and can be delivered into the Kongsfjorden estuary and eventually the North Atlantic. Given the great variability in glacial Fe fluxes and glacial Fe geochemistry observed to date, our results do not necessarily mean that glaciers are not an important source of Fe to the oceans. However they do suggest that cold-based glaciers contribute relatively low concentrations of dFe, and that bedrock type, weathering intensity below the glacier, and the character of the proglacial environment will all have a significant impact on Fe concentration and speciation in meltwater, complicating efforts to extrapolate the global flux of glacial Fe to the oceans from a limited number of measurements.

Anthropogenic climate change and global warming are predicted to greatly impact the Arctic. Under such conditions, proglacial process will play an important role in controlling the amount of Fe that is delivered to the oceans. Our study implies that accurate estimation of glacial Fe fluxes depends both on sampling a wide variety

of glacial locations, and on understanding more fully the processes which lead to the precipitation and stabilization of Fe in proglacial environments.

Fig. 8. Schematic diagram of dFe transportation from glacier to coastal ocean on Svalbard.

Acknowledgements

This work was supported by: the National Program on Key Basic Research Project of China (No. 2011CB409801), the National Natural Science Foundation of China (No. 41006043, 44106290), the Ocean Public Welfare Scientific Research Project (2012YR11011), the State Key Laboratory of Estuarine and Coastal Research innovation funding (2011KYYW02). John and Marsay were funded by the National Science Foundation (NSF OCE-1334029). We would like to thank the Chinese Antarctic and Arctic Administration of National Oceanic Bureau for the logistic support for field sampling. We thank Zhongqiao Li and others for their efforts on the

648 sample collection. We thank Yun Xue for her effort on maintaining the clean lab and

649 Neptune at East China Normal University.

650

650

651 **References**

652

653 Achterberg, E.P., Moore, C.M., Henson, S.A., Steigenberger, S., Stohl, A., Eckhardt,
654 S., Avendano, L.C., Cassidy, M., Hembury, D., Klar, J.K., Lucas, M.I., Macey,
655 A.I., Marsay, C.M., Ryan-Keogh, T.J., 2013. Natural iron fertilization by the
656 Eyjafjallajökull volcanic eruption. *Geophys. Res. Lett.*, 40, 921–926.

657 Balci, N., Bullen, T.D., Witte-Lien, K., Shanks, W.C., Motelica, M., Mandernack,
658 K.W., 2006. Iron isotope fractionation during microbially stimulated Fe(II)
659 oxidation and Fe(III) precipitation. *Geochim. Cosmochim. Acta* 70, 622–639.

660 Bennett, S.A., Rouxel, O., Schmidt, K., Garbe-Schönberg, D., Statham, P.J., German,
661 C.R., 2009. Iron isotope fractionation in a buoyant hydrothermal plume, 5°S
662 Mid-Atlantic Ridge. *Geochim. Cosmochim. Acta* 73, 5619-5634.

663 Bergquist, B.A., Boyle, E.A., 2006. Iron isotopes in the Amazon River system:
664 Weathering and transport signatures. *Earth Planet. Sci. Lett.* 248, 54-68.

665 Bhatia, M.P., Kujawinski, E.B., Das, S.B., Breier, C.F., Henderson, P.B., Charette,
666 M.A., 2013. Greenland meltwater as a significant and potentially bioavailable
667 source of iron to the ocean. *Nat. Geosci.* 6, 274-278.

668 Bogen, J., Bonsnes, T.E., 2003. Erosion and sediment transport in High Arctic rivers,
669 Svalbard. *Polar Research* 22, 175-189.

670 Boyd, P.W., Jickells, T., Law, C.S., Blain, S., Boyle, E.A., Buesseler, K.O., Coale,

671 K.H., Cullen, J.J., de Baar, H.J.W., Follows, M., Harvey, M., Lancelot, C.,
672 Levasseur, M., Owens, N.P.J., Pollard, R., Rivkin, R.B., Sarmiento, J.,
673 Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., Watson, A.J.,
674 2007. Mesoscale iron enrichment experiments 1993-2005: Synthesis and future
675 directions. *Science* 315, 612-617.

676 Boyd, P.W., Mackie, D.S., Hunter, K.A., 2010. Aerosol iron deposition to the surface
677 ocean_Modes of iron supply and biological responses. *Mar. Chem.* 120, 128-143.

678 Bruland, K.W., Lohan, M.C., 2003. Controls of trace metals in seawater. (Holland,
679 H.D., Turekian, K.H. Eds.) *Treatise on Geochemistry* (Vol. 6, p. 7800). Elsevier.

680 Buck, K.N., Lohan, M.C., Berger, C.J.M., Bruland, K.W., 2007. Dissolved iron
681 speciation in two distinct river plumes and an estuary: Implications for riverine
682 iron supply. *Limnology and Oceanography* 52, 843-855.

683 Conway, T.M., John, S.G., 2014. Quantification of dissolved iron sources to the North
684 Atlantic Ocean. *Nature* 511, 212-215

685 Conway, T.M., Rosenberg, A.D., Adkins, J.F., John, S.G., 2013. A new method for
686 precise determination of iron, zinc and cadmium stable isotope ratios in seawater
687 by double-spike mass spectrometry. *Anal. Chim. Acta* 793, 44-52.

688 Dai M.H., Martin, J.M., 1995. First data on the trace metal level and behavior in two
689 major Arctic river/estuarine systems (Ob & Yenisey) and in the adjacent Kara
690 Sea, *Earth Planet. Sci. Lett.* 131, 127-141.

691 Davison, W., Seed, G., 1983. The kinetics of the oxidation of ferrous iron in synthetic

692 and natural waters. *Geochim. Cosmochim. Acta* 47, 67-79.

693 Death, R., Wadham, J.L., Monteiro, F., Le Brocq, A.M., Tranter, M., Ridgwell, A.,
694 Dutkiewicz, S., Raiswell, R., 2014. Antarctic ice sheet fertilises the Southern
695 Ocean. *Biogeosciences* 11, 2635-2643.

696 Dideriksen, K., Baker, J.A., Stipp, S.L.S., 2008. Equilibrium Fe isotope fractionation
697 between inorganic aqueous Fe(III) and the siderophore complex,
698 Fe(III)-desferrioxamine B. *Earth Planet. Sci. Lett.* 269, 280–290.

699 Elrod, V.A., Berelson, W.M., Coale, K.H., Johnson, K.S., 2004. The flux of iron from
700 continental shelf sediments: A missing source for global budgets. *Geophys. Res.*
701 *Lett.* 31, L12307, doi:10.1029/2004GL020216.

702 Escoube, R., Rouxel, O.J., Sholkovitz, E., Donard, O.F.X., 2009. Iron isotope
703 systematics in estuaries: The case of North River, Massachusetts (USA).
704 *Geochim. Cosmochim. Acta* 73, 4045-4059.

705 Fan, S.M., Moxim, W.J., Levy, H., 2006. Aeolian input of bioavailable iron to the
706 ocean. *Geophys. Res. Lett.* 33, L07602, doi:10.1029/2005GL024852.

707 Fantle, M.S., DePaolo, D.J., 2004. Iron isotopic fractionation during continental
708 weathering. *Earth Planet. Sci. Lett.* 228, 547-562.

709 Gerringa, L.J.A., Alderkamp, A.C., Laan, P., Thuroczy, C.E., De Baar, H.J.W., Mills,
710 M.M., van Dijken, G.L., van Haren, H., Arrigo, K.R., 2012. Iron from melting
711 glaciers fuels the phytoplankton blooms in Amundsen Sea (Southern Ocean): Iron
712 biogeochemistry. *Deep-Sea Res. Part II* 71-76, 16-31.

713 Green, W.J., Stage, B.R., Preston, A., Wagers, S., Shacat, J., Newell, S., 2005.
714 Geochemical processes in the Onyx River, Wright Valley, Antarctica: Major ions,
715 nutrients, trace. *Geochim. Cosmochim. Acta* 69, 839-850.

716 Hagen, J.O., Kohler, J., Melvold, K., Winther, J.G., 2003. Glaciers in Svalbard: mass
717 balance, runoff and freshwater flux. *Polar Res.* 22, 145-159.

718 Hawkings, J.R., Wadham, J.L., Tranter, M., Raiswell, R., Benning, L.G., Statham,
719 P.J., Tedstone, A., Nienow, P., Lee, K., Telling, J., 2014. Ice sheets as a
720 significant source of highly reactive nanoparticulate iron to the oceans. *Nat*
721 *Commun* 5.

722 Hodson, A.J., Gurnell, A.M., Washington, R., Tranter, M., Clark, M.J., Hagen, J.O.,
723 1998. Meteorological and runoff time-series characteristics in a small,
724 high-Arctic glaciated basin, Svalbard. *Hydrological Processes* 12, 509-526.

725 Hodson, A., Tranter, M., Gurnell, A., Clark, M., Hagen, J.O., 2002. The
726 hydrochemistry of Bayelva, a high Arctic proglacial stream in Svalbard. *Journal*
727 *of Hydrology* 257, 91-114.

728 Hodson, A.J., Mumford, P.N., Kohler, J., Wynn, P.M., 2005. The High Arctic glacial
729 ecosystem: new insights from nutrient budgets. *Biogeochemistry* 72, 233-256.

730 Laglera, L.M., van den Berg, C.M.G., 2009. Evidence for geochemical control of iron
731 by humic substances in seawater. *Limnology and Oceanography* 54, 610-619.

732 Ilina, S.M., Poitrasson, F., Lapitskiy, S.A., Alekhin, Y.V., Viers, J., Pokrovsky, O.S.,
733 2013. Extreme iron isotope fractionation between colloids and particles of boreal

734 and temperate organic-rich waters. *Geochim. Cosmochim. Acta* 101, 96-111.

735 Jickells, T.D., An, Z.S., Andersen, K.K., Baker, A.R., Bergametti, G., Brooks, N.,
736 Cao, J.J., Boyd, P.W., Duce, R.A., Hunter, K.A., Kawahata, H., Kubilay, N.,
737 laRoche, J., Liss, P.S., Mahowald, N., Prospero, J.M., Ridgwell, A.J., Tegen, I.,
738 Torres, R., 2005. Global iron connections between desert dust, ocean
739 biogeochemistry, and climate. *Science* 308, 67-71.

740 Johnson, C.M., Skulan, J.L., Beard, B.L., Sun, H., Nealson, K.H., Brateman, P.S.,
741 2002. Isotopic fractionation between Fe(III) and Fe(II) in aqueous
742 solutions. *Earth Planet. Sci. Lett.* 195, 141-153.

743 Jungclaus, J.H., Haak, H., Latif, M., Mikolajewicz, U., 2005. Arctic-North Atlantic
744 interactions and multidecadal variability of the meridional overturning circulation.
745 *J. Clim.* 18, 4013-4031.

746 Krachler, R., Jirsa, F., Ayromlou, S., 2005. Factors influencing the dissolved iron
747 input by river water to the open ocean. *Biogeosciences*. 2, 311-315.

748 Krawczyk, W.E., Lefauconnier, B., Pettersson, L.E., 2003. Chemical denudation rates
749 in the Bayelva Catchment, Svalbard, in the Fall of 2000. *Physics and Chemistry*
750 *of the Earth* 28, 1257-1271.

751 Lam, P.J., and Bishop, J.K., 2008. The continental margin is a key source of iron to
752 the HNLC North Pacific Ocean. *Geophys. Res. Lett.* 35, L07608,
753 doi:10.1029/2008GL033294.

754 Lee, J.M., Boyle, E.A., Echegoyen-Sanz, Y., Fitzsimmons, J.N., Zhang, R.F., Kayser,

755 R.A., 2011. Analysis of trace metals (Cu, Cd, Pb, and Fe) in seawater using
756 single batch nitrilotriacetate resin extraction and isotope dilution inductively
757 coupled plasma massspectrometry. *Anal. Chim. Acta.* 686, 93-103.

758 Moore, J.K., Doney, S.C., Glover, D.M., Fung, I.Y., 2002. Iron cycling and
759 nutrient-limitation patterns in surface waters of the world ocean. *Deep-Sea Res.*
760 *Part II-Top. Stud. Oceanogr.* 49, 463-507.

761 Moore, J.K., Doney, S.C., and Lindsay, K., 2004. Upper ocean ecosystem dynamics
762 and iron cycling in a global three dimensional model. *Global Biogeochem. Cy.* 18,
763 GB4028,doi:10.1029/2004GB002220.

764 Morgan, J.J., 2005. Kinetics of reaction between O₂ and Mn(II) species in aqueous
765 solutions. *Geochim. Cosmochim. Acta* 69, 35-48.

766 Mostofa, K.M.G., Yoshioka, T., Konohira, E., Tanoue, E., 2007. Photodegradation of
767 fluorescent dissolved organic matter in river waters. *Geochemical J.* 41, 323-331.

768 Nielsdottir, M.C., Moore, C.M., Sanders, R., Hinz, D.J., Achterberg, E.P., 2009. Iron
769 limitation of the postbloom phytoplankton communities in the Iceland Basin.
770 *Glob. Biogeochem. Cycles* 23, GB3001, doi:10.1029/2008GB003410.

771 Nowak, A., Hodson, A., 2013. Hydrological response of a High-Arctic catchment to
772 changing climate over the past 35 years: a case study of Bayelva watershed,
773 Svalbard. *Polar Research* 32, 19691, <http://dx.doi.org/10.3402/polar.v32i0.19691>

774 Nowak, A., Hodson, A., 2014. Changes in meltwater chemistry over a 20-year period
775 following a thermal regime switch from polythermal to cold-based glaciation at

776 Austre Brøggerbreen, Svalbard. Polar Research 33, 22779,
 777 <http://dx.doi.org/10.3402/polar.v33.22779>
 778 Parekh, P., Follows, M.J., Boyle, E., 2004. Modeling the global ocean iron cycle.
 779 Glob. Biogeochem. Cycle 18, GB1002, doi:10.1029/2003GB002061.
 780 Poitrasson, F., 2006. On the iron isotope homogeneity level of the continental crust.
 781 Chemical Geology 235, 195-200.
 782 Raiswell, R., 2013. Biogeochemistry: Rusty meltwaters. Nat. Geosci. 6, 251-252.
 783 Raiswell, R., 2011. Iceberg-hosted nanoparticulate Fe in the Southern Ocean:
 784 Mineralogy, origin, dissolution kinetics and source of bioavailable Fe. Deep Sea
 785 Research Part II: Topical Studies in Oceanography 58, 1364-1375.
 786 Raiswell, R., Benning, L.G., Davidson, L., Tranter, M., Tulaczyk, S., 2009.
 787 Schwertmannite in wet, acid, and oxic microenvironments beneath polar and
 788 polythermal glaciers. Geology 37, 431-434.
 789 Raiswell, R., Canfield, D.E., 2012. The iron biogeochemical cycle past and present.
 790 Geochemical Perspectives 1, 1-2.
 791 Raiswell, R., Tranter, M., Benning, L.G., Siegert, M., De'ath, R., Huybrechts, P.,
 792 Payne, T., 2006. Contributions from glacially derived sediment to the global iron
 793 (oxyhydr)oxide cycle: Implications for iron delivery to the oceans. Geochim.
 794 Cosmochim. Acta 70, 2765-2780.
 795 Resing, J.A., Mottl, M.J., 1992. Determination of manganese in seawater using flow
 796 injection analysis with on-line preconcentration and spectrophotometric detection.

797 Anal. Chem. 64, 2682-2687.

798 Revels, B. N., Ohnemus, D. C., Lam, P. J., Conway, T. M. & John, S. G. The isotope
799 signature and distribution of particulate iron in the north Atlantic ocean. Deep
800 Sea Res. Part II Top. Stud. Oceanogr. doi:10.1016/j.dsr2.2014.12.004 (in press).

801 Rue, E.L., Bruland, K.W., 1995. Complexation of iron(III) by natural organic-ligands
802 in the central North Pacific as determined by a new competitive ligand
803 equilibration adsorptive cathodic stripping voltammetric
804 method. Mar. Chem. 50, 117-138.

805 Rouxel, O., Sholkovitz, E., Charette, M., Edwards, K.J., 2008. Iron isotope
806 fractionation in subterranean estuaries. Geochim. Cosmochim. Acta 72,
807 3413-3430.

808 Saito, M.A., Noble, A.E., Tagliabue, A., Geopfert, T.J., Lamborg, C.H., Jenkins, W.J.,
809 2013. Slow-spreading submarine ridges in the South Atlantic as a significant
810 oceanic iron source. Nat. Geosci. 6, 775-779.

811 Schroth, A.W., Crusius, J., Chiver, F., Bostick, B.C., Rouxel, O.J., 2011. Glacial
812 influence on the geochemistry of riverine iron fluxes to the Gulf of Alaska and
813 effects of deglaciation. Geophys. Res. Lett. 38, L16605,
814 doi:10.1029/2011GL048367.

815 Severmann, S., Johnson, C.M., Beard, B.L., German, C.R., Edmonds, H.N., Chiba, H.,
816 Green, D.R.H., 2004. The effect of plume processes on the Fe isotope
817 composition of hydrothermally derived Fe in the deep ocean as inferred from the

818 Rainbow vent site, Mid-Atlantic Ridge, 36°14'N. *Earth Planet. Sci. Lett.* 225, 63
819 –76.

820 Severmann, S., Johnson, C.M., Beard, B.L., McManus, J., 2006. The effect of early
821 diagenesis on the Fe isotope compositions of porewaters and authigenic minerals
822 in continental margin sediments. *Geochim. Cosmochim. Acta* 70, 2006-2022.

823 Severmann, S., McManus, J., Berelson, W.M., Hammond, D.E., 2010. The
824 continental shelf benthic iron flux and its isotope composition. *Geochim.*
825 *Cosmochim. Acta* 74, 3984-4004.

826 Shiller, A.M., Duan, S.W., van Erp, P., Bianchi, T.S., 2006. Photo-oxidation of
827 dissolved organic matter in river water and its effect on trace element speciation.
828 *Limno. Oceanogr.* 51, 1716-1728.

829 Smith, K.L., Robison, B.H., Helly, J.J., Kaufmann, R.S., Ruhl, H.A., Shaw, T.J.,
830 Twining, B.S., Vernet, M., 2007. Free-Drifting Icebergs: Hot Spots of Chemical
831 and Biological Enrichment in the Weddell Sea. *Science* 317, 478-482.

832 Statham, P.J., Jacobson, Y., van den Berg, C.M.G., 2012. The measurement of
833 organically complexed FeII in natural waters using competitive ligand reverse
834 titration. *Anal. Chim. Acta* 743, 111-116.

835 Statham, P.J., Skidmore, M., Tranter, M., 2008. Inputs of glacially derived dissolved
836 and colloidal iron to the coastal ocean and implications for primary productivity.
837 *Glob. Biogeochem. Cycle* 22. GB3013, doi:10.1029/2007GB003106.

838 Stolpe, B., Guo, L., Shiller, A., Hasselov, M., 2010. Size and composition of

839 colloidal organic matter and trace elements in the Mississippi River, Pearl River
840 and the northern Gulf of Mexico as characterized by flow field-flow fractionation.
841 Mar. Chem. 118, 119-129.

842 Tagliabue, A., Bopp, L., Dutay, J.C., Bowie, A.R., Chever, F., Jean-Baptiste, P.,
843 Bucciarelli, E., Lannuzel, D., Remenyi, T., Sarthou, G., Aumont, O., Gehlen, M.,
844 Jeandel, C., 2010. Hydrothermal contribution to the oceanic dissolved iron
845 inventory. Nat. Geosci. 3, 252-256.

846 Thuróczy, C.-E., Alderkamp, A.-C., Laan, P., Gerringa, L.J.A., Mills, M.M., Van
847 Dijken, G.L., De Baar, H.J.W., Arrigo, K.R., 2012. Key role of organic
848 complexation of iron in sustaining phytoplankton blooms in the Pine Island and
849 Amundsen Polynyas (Southern Ocean). Deep Sea Research Part II: Topical
850 Studies in Oceanography 71–76, 49-60.

851 Wang, Z.W., Ren, J.L., Liu, S.M., 2015. Determination method of dissolved
852 manganese in natural water by catalytic kinetic spectrophotometry. Marine
853 Environmental Science. (accepted manuscript, in Chinese)

854 Wadham, J.L., De'ath, R., Monteiro, F.M., Tranter, M., Ridgwell, A., Raiswell, R.,
855 Tulaczyk, S., 2013. The potential role of the Antarctic Ice Sheet in global
856 biogeochemical cycles. Earth and Environmental Science Transactions of the
857 Royal Society of Edinburgh 104, 55-67.

858 Wehrmann, L.M., Formolo, M.J., Owens, J.D., Raiswell, R., Ferdelman, T.G.,
859 Riedinger, N., Lyons, T.W., 2014. Iron and manganese speciation and cycling in

860 glacially influenced high-latitude fjord sediments (West Spitsbergen, Svalbard):
861 Evidence for a benthic recycling-transport mechanism. *Geochim. Cosmochim.*
862 *Acta* 141, 628-655.

863 Welch, S.A., Beard, B.L., Johnson, C.M., Braterman, P.S., 2003. Kinetic and
864 equilibrium Fe isotope fractionation between aqueous Fe(II) and
865 Fe(III). *Geochim. Cosmochim. Acta* 67, 4231-4250.

866 Wu, Y., Bao, H.Y., Unger, D., Herbeck, L.S., Zhu, Z.Y., Zhang, J., Jennerjahn, T.C.,
867 2013. Biogeochemical behavior of organic carbon in a small tropical river and
868 estuary, Hainan, China. *Cont. Shelf Res.* 57, 32-43.

869 Wu, J.F., Boyle, E., Sunda, W., Wen, L.S., 2001. Soluble and colloidal iron in the
870 olgotrophic North Atlantic and North Pacific. *Science* 293, 847-849.

871