Distinctly different parental magmas for calc-alkaline plutons and tholeiitic lavas in the central and eastern Aleutian arc

Yue Cai^{1*}, Matthew E. Rioux², Peter B. Kelemen^{1,3}, Steven L. Goldstein^{1,3}, Louise Bolge¹,
Andrew R. C. Kylander-Clark⁴

¹*Lamont-Doherty Earth Observatory of Columbia University, 61 Rt. 9W, Palisades, NY, 10964,
USA*

²*Earth Research Institute, University of California, Santa Barbara, CA, 93106, USA*

³*Department of Earth and Environmental Sciences, Columbia University, 61 Rt. 9W, Palisades,
NY, 10964, USA*

⁴*Department of Earth Science, University of California, Santa Barbara, CA, 93106*

*email of the corresponding author: cai@ldeo.columbia.edu, merrycai@gmail.com,

ABSTRACT

Cenozoic calc-alkaline plutons that comprise the middle crust of the central and eastern Aleutians have distinct isotopic and elemental compositions compared to Holocene tholeiitic lavas in the same region, including those from the same islands. Therefore the Holocene lavas are not representative of the net magmatic transfer from the mantle into the arc crust. Compared to the lavas, the Eocene to Miocene (9-39 Ma) intermediate to felsic plutonic rocks show higher SiO₂ at a given Fe/Mg ratio, and have higher ϵ_{Nd} - ϵ_{Hf} values and lower Pb-Sr isotope ratios. However, the plutonic rocks strongly resemble calc-alkaline Holocene volcanics with “depleted” isotope ratios in the western Aleutians, whose composition has been attributed to significant contributions from partial melting of subducted basaltic oceanic crust. These data could reflect a

temporal variation of central and eastern Aleutian magma source compositions, from predominantly calc-alkaline compositions with more “depleted” isotope ratios in the Paleogene, to tholeiitic compositions with more “enriched” isotopes more recently. Alternatively, the differences between central Aleutian plutonic and volcanic rocks may reflect different transport and emplacement processes for the magmas that form plutons versus lavas. Calc-alkaline parental magmas, with higher SiO₂ and high viscosity, are likely to form plutons after extensive mid-crustal degassing of initially high water contents. This conclusion has overarching importance because the plutonic rocks are chemically similar to bulk continental crust. Formation of similar plutonic rocks worldwide may play a key role in the genesis and evolution of continental crust.

1. INTRODUCTION

Arc magmatism is the most important process that generates continental crust today and probably throughout Earth history. It is commonly inferred that compositions of primitive arc lavas are representative of the bulk composition of arc crust (e.g., Rudnick 1995; Tatsumi et al. 2008). However, the composition of average arc lavas is basaltic, while the bulk continental crust is andesitic with high molar Mg/(Mg+Fe) ratio or Mg# (e.g., Taylor and McLennan, 1985; Rudnick and Fountain, 1995; Rudnick and Gao, 2003; Kelemen et al., 2003, 2014). Many hypotheses have been proposed to resolve this discrepancy, such as lower crustal foundering (Herzberg et al. 1983; Arndt and Goldstein, 1989; Kay and Kay, 1991); crustal formation from primary mantle-derived andesitic magmas (Kelemen, 1995); mixing of basaltic rock with silicic magma derived by partial melting of mafic, subducting crust (Martin, 1986); relamination of felsic, differentiated subducted crustal material (Hacker et al., 2011) and fractionation of water-

rich magmas at moderate to high pressure conditions (Sisson and Grove, 1993; Grove et al., 2003; Jagoutz et al., 2011). A key aspect of several of these hypotheses is that the compositions of the largely unsampled plutonic arc middle crust may differ from the erupted arc lavas, and instead more closely resemble the continental crust (e.g., Perfit, 1980; Kay et al., 1990).

Low seismic wave-speeds suggest that plutonic arc middle crust may be more silicic than surficial volcanic rocks in some arcs (e.g. Suyehiro et al., 1996; Tatsumi et al., 2008). Similarly, combined geobarometry and geochemistry for the Mesozoic Talkeetna and Kohistan intra-oceanic arc sections show that felsic rocks are abundant in the middle crust, and that the crustal bulk composition may have been andesitic rather than basaltic (Behn and Kelemen, 2006; Hacker et al., 2008; Jagoutz and Schimdt, 2012; Rioux et al., 2010). In the Aleutian arc, seismic data require a mafic bulk composition for the arc lower crust (Holbrook et al. 1999; Flidner & Klemperer, 1999; Shillington et al., 2004); however, the upper and middle crust in the Aleutians contain abundant intermediate composition plutons (e.g., Kay et al., 1982, 1990, Kay and Kay, 1994; Perfit et al., 1980; Yogodzinski et al., 1993).

This paper reports new data on plutonic rocks (and two lavas) from the central and eastern Aleutian arc, and compares them with the large existing dataset on Holocene Aleutian volcanics. We define the central and eastern Aleutians as east of 180°W, which includes the oceanic arc extending from the western end of the Alaska Peninsula to the Delarof Islands; and the western Aleutians as west of 180°W, which includes the active arc extending from the Rat Islands west to the Komandorsky Islands and adjacent submarine volcanoes (Fig. 1A). In order to constrain the composition and timing of plutonism, we report results from whole rock major and trace elements; Sm-Nd, Rb-Sr and Lu-Hf isotopic analyses; and U-Pb zircon geochronology.

2. METHODS

Zircon U-Pb geochronology was conducted by laser ablation on a Nu plasma multi-collector ICP-MS at the University of California at Santa Barbara (UCSB), following the procedures described in Kylander-Clark et al. (2013). Reported $^{206}\text{Pb}/^{238}\text{U}$ dates are Th- and common Pb-corrected, as described in the U-Pb zircon geochronology methods and data interpretations sections of the Supplementary Material (Table 2S, Text S1).

All other geochemical analyses were carried out at Lamont-Doherty Earth Observatory (LDEO). Major element analyses were carried out using an Agilent ICP-OES on rock powders digested with $\text{Li}_2\text{B}_4\text{O}_7$ flux fusion. Trace element analyses were carried out on a VG PlasmaQuad ExCell quadrupole ICP-MS on rock powders digested using both hotplate HF-HNO_3 acid digestion (Ba, U, Th, Rb, Sr, Li, Ce, Pb, Nb, Ta) and Na_2O_2 sintering (La, Nd, Sm, Eu, Pr, Zr, Hf, Tb, Gd, Dy, Yb, Lu, Y) to ensure complete digestion of zircons and other accessory minerals. Sintering was also used for preparation of samples for isotopic analyses. A comparison of the effects of hotplate digestions and sintering demonstrated its importance for elements such as Zr and Hf, and for Hf isotopes in older samples. Details, and the data, are in the Supplementary Materials (Tables 1S-4S, Text S2).

Nd, Pb, Hf, and Sr isotopes were measured on a ThermoScientific Neptune Plus multi-collector MC-ICP-MS in static mode; some Sr isotope ratios were measured on a VG-Sector 54 TIMS in multi-dynamic mode (Table S1). The MC-ICP-MS instrumental drift is monitored by standard-sample bracketing using JNdi, a Johnson-Matthey lab Hf standard with Hf isotope ratios indistinguishable from JMC 475, NBS 987 Sr, and NBS 981 Pb. Typically 30-70 standards and 20-30 samples are analyzed during a 24-hour long run period, and reported 2SD external errors reflect the reproducibility of the standards. Reported values are corrected to those of the international standards. Quality control was further evaluated through measurements of the La

Jolla Nd solution standard, and rock standards BCR-2 and BHVO-2. The analytical metadata (standards data, amounts per sample, signal intensities, values used for international standards, results, and reproducibility) are listed in Table S5. Hf isotope ratios are often listed as ϵHf , even on modern day and recent samples, therefore we do so here, where the deviation is in parts per 10,000 from a “Chondritic Uniform Reservoir” (CHUR) value ($^{176}\text{Hf}/^{177}\text{Hf}_{\text{CHUR}} = 0.282785$, Bouvier et al., 2008), and also for Nd isotope ratios (where $^{143}\text{Nd}/^{144}\text{Nd}_{\text{CHUR}} = 0.512638$, Jacobsen and Wasserburg, 1980).

3. RESULTS

Weighted mean laser ablation-ICP-MS Th-corrected $^{206}\text{Pb}/^{238}\text{U}$ ages of the plutonic samples range from 9.20 ± 0.04 to 39.1 ± 0.2 Ma (Eocene to Miocene), which means they are older than lavas from the same region that have been the focus of most previous studies. The plutonic rocks are predominantly calc-alkaline. They show higher average SiO_2 and higher trace element concentrations than lavas from the same region at a given Fe/Mg ratio (calculated using wt% MgO and FeO, with all Fe as FeO) (Fig. 1A-C). The major and trace element compositions of the studied plutonic rocks resemble lavas from the western Aleutians, as well as estimates for the bulk continental crust (Fig. 1B-C). Except for two samples, the plutonics show negligible Eu anomalies. They have similar or higher Na+K (mol%), and similar or lower CaO and Al_2O_3 (wt%) at a given Mg# compared to the lavas (Fig. 2). These data suggest that the plutonic rocks represent fully crystallized magmas, rather than crystal cumulates resulting from partial crystallization followed by removal of evolved melt. The two samples with positive Eu anomalies (marked with red rims in the figures) show clear signatures of plagioclase accumulation (e.g., high Al). However, we include these samples in the following discussion

because they have similar isotope compositions as other plutonic samples, and because plagioclase accumulation should not change the isotopic composition of the magma.

The Eocene-Miocene central and eastern Aleutian plutons show higher present-day ϵ_{Nd} and ϵ_{Hf} , and lower Pb isotope ratios than the Holocene volcanic rocks from the same area (Figs. 3 and 4). Age-corrected Sr isotope ratios in the plutonic samples are also generally lower than in Holocene lavas from the same islands (Fig. S4), though we place less weight on the Sr data due to the possible effects of alteration. The isotopic compositions of the plutonic rocks fall between the fields defined by central-eastern and western Aleutian lavas (Fig. 4). There is no clear temporal trend in Nd-Hf-Pb isotope compositions of Aleutian plutonic samples (Fig. 3). Without sintering, Hf isotope ratios and concentrations reflected incomplete digestion of zircons (shown in Supplementary Materials). After sintering, Nd-Hf isotope ratios lie close to the mantle crust array (Vervoort et al., 2011) and show negative correlations with Pb isotope ratios (Fig. 4). Age corrections do not change these relationships (Figs. S4-S5 and Table S4).

4. DISCUSSION

4.1 Central-eastern Aleutian plutons and lavas are derived from distinct sources

This study is the first that demonstrates statistically significant and internally consistent differences in isotope and elemental compositions between lavas and the plutons in any arc. It has been shown previously that in the central and eastern Aleutians the erupted lavas are predominantly tholeiitic while the plutons are predominantly calc-alkaline (e.g., Kay, 1978; Kay et al., 1982; Kay and Kay, 1990; Fig. 4B in Kelemen et al. 2003a; Fig. 4 and 5 in Kelemen et al. 2003b, 2014), with the exception of the tholeiitic Finger Bay pluton on Adak Island. Isotopic differences between plutons and lavas have been inferred previously based on a limited number of samples (Kay et al., 1986).

The differences in isotope signatures between the plutonic rocks studied here, and lavas from the same islands in the central and eastern Aleutians, demonstrate that the intrusive and extrusive suites in this area are derived from different sources. These isotopic differences cannot be explained by chemical differentiation or crustal assimilation, although these processes may affect some other compositional characteristics of Aleutian magmas.

Elevated parental magma water contents and oxygen fugacity can drive the differentiating magma towards the calc-alkaline trend by suppressing plagioclase fractionation and bringing oxide and amphibole fractionation closer to the liquidus (e.g., Sisson and Grove, 1993). This process is often called upon to explain co-existing calc-alkaline and tholeiitic arc magmas (e.g., Osborn, 1959; Gill 1981, Sections 11.3 and 12.1; Grove et al., 2003, 2005; Zimmer et al., 2010; Jagoutz et al., 2011). However, it cannot explain the isotopic differences between the plutons and the lavas.

High ϵ_{Nd} and ϵ_{Hf} , and lower Pb isotope ratios might be generated via assimilation of arc crust or pre-existing oceanic crust (e.g. Kay and Kay 1994). However, AFC (Assimilation during Fractional Crystallization) modeling shows that to generate the isotopic compositions of the plutons via crustal assimilation requires large extents of fractionation (up to 90%, Fig. 5) and the resulting magma would have high Nd and Hf contents and strong negative Eu anomalies. These signatures are not observed in the plutonic rocks, which plot far above the AFC trend. Thus, the isotopic differences between the plutons and the lavas in the Aleutians were not generated through crustal assimilation.

One may also argue that the plutons could represent deeper crustal magma chambers that have experienced more mafic magmatic recharge (e.g. Lee et al., 2014), a process that could buffer the isotopic compositions of the magma towards “isotopically depleted” compositions

(referring to Nd-Hf-Pb-Sr isotopic values that reflect long term depletions in incompatible elements, thus low Nd/Sm, Hf/Lu, U/Pb and Th/Pb). However, mafic magma recharge cannot explain the major element characteristics of the plutons. For example, compared to the volcanics, the plutons show elevated SiO₂ contents for a given Fe/Mg ratio (Fig. 1B) and lower MgO contents for a given Mg# (Fig S6A). Additionally, mixing between more primitive, isotopically depleted magmas and more evolved, isotopically enriched magmas should generate negative correlations between Nd isotope ratios and SiO₂ and positive correlations between Nd isotopes and MgO, which is not the case for the plutons (Fig S6 B-C). Instead, the isotopic differences between central-eastern Aleutian plutons and lavas persist across a large range of major element compositions.

4.2 Coeval formation of calc-alkaline and tholeiitic parental magmas? Highly likely

Instead, we propose two alternative hypotheses based on the data presented. (1) The sources of central and eastern Aleutian magmas changed over time, with the early source characterized by “depleted” Nd, Hf and Pb isotope ratios, represented by the plutons, which are Miocene and older; changing to a present day source characterized by relatively more “enriched” Nd, Hf and Pb isotope ratios, represented by Holocene central and eastern Aleutian lavas. In this view, the isotopic shift was accompanied by a change in major element magma composition, from more calc-alkaline to more tholeiitic, perhaps due to decreasing SiO₂, *f*O₂, and/or H₂O contents in central-eastern Aleutian parental magmas over time. (2) The differences between plutonic and volcanic compositions reflect coeval input of distinct parental magma types from the mantle into the Aleutian crust. In this case, the dominance of calc-alkaline compositions among the plutons could reflect higher SiO₂, *f*O₂, and/or H₂O contents in their parental magmas,

182 compared to the basaltic, more reduced magmas with lower H₂O contents that are parental to
183 Aleutian tholeiitic lavas (Kay et al., 1990; Kelemen et al., 2003a,b, 2014).

184 The scarcity of isotope data on pre-Holocene volcanics in the central and eastern Aleutian
185 islands that host our plutonic samples means that we cannot definitively distinguish between
186 these two hypotheses. However, the isotopic and geochronological data from the plutons do not
187 show systematic temporal variations from 39Ma to 9Ma (Fig. 3C-D). If central and eastern
188 Aleutian lavas and plutons have shared a common source composition through time, then the
189 data require an abrupt change in the magma source between ~9 Ma and the Holocene, from
190 predominantly calc-alkaline to predominantly tholeiitic. We are not aware of any large changes
191 in the regional tectonics that would cause such a shift. Moreover, coeval calc-alkaline plutons
192 and tholeiitic lavas are observed at Adak, where calc-alkaline plutons intrude ca. 38 Ma tholeiitic
193 lavas, suggesting that the observed compositional differences between the plutons and the lavas
194 have persisted through time (Jicha et al., 2006; Kay et al., 1983).

195 Therefore, we prefer the hypothesis that the distinct isotopic and geochemical signatures
196 of the central and eastern Aleutian plutons and lavas reflect distinct, coeval magma sources. The
197 observed differences may be generated by the addition of different types and amounts of
198 subducted material to calc-alkaline and tholeiitic magmas, mirroring the differences between
199 western vs. central and eastern Aleutian lavas.

200 **4.3 Central-eastern Aleutian plutons resemble western Aleutian calc-alkaline lavas**

201 The oceanic part of the Aleutian arc has a well-defined, along-strike variation in
202 Holocene lava compositions, ranging from dominantly calc-alkaline compositions with
203 “depleted” Sr-Nd-Pb-Hf isotopic compositions in the western Aleutians, to tholeiitic
204 compositions with more “enriched” isotopic compositions in the east (e.g., Kay and Kay, 1994;

Kelemen et al., 2003; Singer et al., 2007). End-member western Aleutian lava compositions are primitive andesites and dacites showing light rare earth element (LREE) enrichment, heavy REE (HREE) depletion, and high Mg# (>0.6), Ni (>150 ppm) and Sr/Nd (>100). These can be understood as the result of reaction of mantle peridotite and hydrous, low degree partial melts of subducted material that formed in eclogite facies (e.g., Kay, 1978; Yogodzinski et al., 1995, 2001, 2015; Yogodzinski and Kelemen 1998, 2007; Kelemen and Yogodzinski, 2007). A quantitative model of the reaction process is given in Kelemen et al. (2003a). Primitive calc-alkaline high Mg# andesites and dacites with strong adakitic signatures are found erupted through thin oceanic lithosphere in the western Aleutians, rendering it unlikely that deep crustal assimilation is responsible for their characteristic compositional signatures (Yogodzinski et al., 2015).

Western Aleutian lavas have the highest Nd and lowest Pb isotope ratios of any arc magmas worldwide (Kelemen et al., 2003a,b, 2014; Singer et al., 2007; Yogodzinski et al., 2010, 2015). Both western Aleutian lavas and the plutonic rocks from the central and eastern Aleutians show high ϵ_{Nd} and ϵ_{Hf} values (Fig. 6) coupled with higher Th/Nd and Hf/Lu ratios than the “depleted MORB mantle” (DMM) source for mid-ocean ridge basalts (Salters and Stracke, 2004). These compositional characteristics, that is high Nd-Hf isotope ratios and incompatible element enrichment, are consistent with addition of low degree partial melts of subducted material in eclogite facies, where garnet and omphacite preferentially retain Lu and Nd in the solid residue compared to Hf and Th, respectively (e.g., Cai et al., 2014; Kay, 1978; Klemme et al., 2002; Yogodzinski et al., 2015).

The eclogite melt component derived from the subducted material could comprise ~20% of the parental magma for western Aleutian primitive andesites and dacites (Fig. 6, and Kelemen

et al., 2003a; Yogodzinski et al., 2015). The SiO₂-rich nature of this component poses a bit of a conundrum. The high Mg# and Ni contents of the primitive andesites and dacites suggest extensive interaction with mantle olivine, which should limit SiO₂ contents of the resulting magmas to <60 wt% depending on pressure, alkali content, and H₂O content (Carroll et al., 1989; Draper et al., 1999). Furthermore, reaction with the hot mantle wedge might be expected to raise MgO contents, whereas primitive Aleutian dacites have <5 wt% MgO despite their high Mg#'s. One possible explanation is that reacting eclogitic melts rise in cold diapirs that pond just beneath the crust and evolve by combined reaction and crystal fractionation in the shallow mantle (e.g., Gerya et al., 2003; Kelemen et al., 2003b; Ringwood and Green, 1966). Instead or in addition, hydrous, high SiO₂ eclogite melts may be sufficiently abundant to convert some regions of the sub-arc mantle from peridotite to olivine-free pyroxenite by overwhelming the olivine-pyroxene SiO₂ buffer (Macpherson et al., 2006; Yogodzinski et al., 1994). Independent of the cause, the link between a substantial eclogite melt component and primitive andesites and dacites seems clear, not only in the western Aleutians but worldwide (e.g. Cai et al., 2014; Defant and Drummond, 1990; Kay, 1978; Shimoda et al., 1998; Stern and Kilian, 1996; Straub et al., 2011).

The calc-alkaline nature of western Aleutian lavas suggests high magmatic water contents. Zimmer et al. (2010) and Ruscitto et al. (2011) demonstrated a correlation between high H₂O content and strongly calc-alkaline magma series in the Aleutians and the Cascades. It is possible that calc-alkaline magma series also have higher fO_2 in their parental liquids, compared to tholeiitic magma series. Thus, we infer that parental magmas for typical western Aleutian primitive andesites and dacites likely have higher SiO₂, H₂O and fO_2 compared to parental magmas for typical central and eastern Aleutian tholeiitic basalts.

In comparison, central and eastern Aleutian volcanics have lower ϵ_{Nd} and ϵ_{Hf} values, and higher Sr and Pb isotope ratios, all of which are consistent with incorporation of a subducted sediment component similar to near-trench sediments sampled at DSDP Site 183 (Fig. 6 and Peucker-Ehrenbrink et al., 1994). The sediment component in the central and eastern Aleutian lavas is thought to contribute just a few percent of the mass of the parental magmas, with the balance coming from fluxed melting of the mantle wedge (Class et al., 2000; Kay et al., 1978; Miller et al., 1994; Singer et al., 2007). The isotopic signals associated with this sediment component are much weaker in western Aleutian lavas and systematically decrease with distance west of Adak (Kelemen et al., 2003a; Yogodzinski et al., 2015).

4.4 Calc-alkaline magmas may preferentially form plutons

If the distinct chemical and isotopic compositions of the plutonic and volcanic rocks in the central and eastern Aleutians are related to the addition of different types and amounts of subducted material, the intrusive versus extrusive nature of the lavas are likely also related to different transport mechanisms. Based on the major element and isotopic characteristics, the parental magmas are similar in the central and eastern Aleutian plutonic rocks and western Aleutian lavas. However, plutonics and lavas in central and eastern Aleutians have distinctly different parental magmas. Primitive ($\text{Mg\#} > 0.6$) andesites and dacites in the western Aleutians have low MgO, indicative of low melt-mantle equilibration temperatures, whereas primitive basalts in the central and eastern Aleutians have higher MgO, indicative of high melt-mantle equilibration temperatures. Magmas with high SiO_2 and H_2O become viscous, and preferentially stall during mid-crustal degassing (e.g., Cann, 1970). It is reasonable to infer that in the central and eastern Aleutians, such magmas predominantly stall to form the calc-alkaline plutons, rather than erupting as previously proposed (e.g., Kay et al., 1990; Kay and Kay, 1994; Kelemen et al.,

2003a). In the thinner, faulted crust of the western Aleutians, such magmas erupt to form many small cones on the seafloor. In contrast, the parental magmas to central and eastern Aleutian tholeiitic lavas likely have higher temperatures and lower SiO₂ and H₂O contents. Such lower viscosity magmas may readily erupt.

If it turns out that parental magmas for both calc-alkaline and tholeiitic magma series are simultaneously present in the Holocene central and eastern Aleutian arc, then hot, parental tholeiites from the core of the mantle wedge may form conduits that “pierce through” shallow mantle pyroxenites and ponded parental calc-alkaline magmas during relatively rapid ascent to the surface. Such a process in arcs worldwide would result in the bulk arc crust having higher SiO₂ than erupted arc volcanics. This has important implications for our understanding of continent formation, since arc lavas are routinely used to estimate the average composition of magmas passing from the mantle into arc crust (e.g., Plank and Langmuir, 1993; Plank et al. 2002), and the bulk composition of arc crust (e.g., Tatsumi 2000; Tatsumi et al. 2008).

5. CONCLUSIONS

In summary, Eocene to Miocene plutonic rocks in the central and eastern Aleutians show stronger calc-alkaline signatures than Holocene volcanics from the same region, and are similar to Holocene volcanics from the western Aleutians. The central and eastern Aleutian plutonics have higher ϵ_{Nd} and ϵ_{Hf} , and lower Pb and Sr isotope ratios than Holocene central and eastern Aleutian lavas. The isotopic signatures of the plutonic samples cannot be generated by crustal assimilation or magma fractionation in the Aleutians. Instead, they demonstrate distinct sources for the Holocene lavas and Cenozoic plutons in the central and eastern Aleutians. The parental magmas for the calc-alkaline plutons and lavas likely have higher SiO₂, H₂O and $f\text{O}_2$ than the parental magmas for tholeiitic lavas. This indicates that the major element differences between

calc-alkaline and tholeiitic magma series in the central and eastern Aleutians arise, in part, from parental magmas with distinct initial compositions. The difference between central and eastern Aleutian lavas and plutonic rocks may reflect a temporal variation in the nature of parental magmas, or the effect of magma transport processes alone. More detailed studies of Aleutian plutons and pre-Holocene central and eastern Aleutian lavas are needed to resolve this uncertainty. However, the available evidence favors the hypothesis that the calc-alkaline magmas preferentially form plutons as they undergo a rapid viscosity increase during mid-crustal degassing. In contrast, dryer and hotter tholeiitic parental magmas with lower initial SiO₂ and H₂O contents preferentially erupt. Regardless of the cause, a first-order implication is that erupted magmas alone are not an accurate guide to the composition of the bulk arc crust. Furthermore, Aleutian plutonic rocks – and similar plutonic rocks worldwide – are compositionally similar to continental crust. Such plutons may be important building blocks for the continental crust. Understanding the processes of their formation is a key to understanding the genesis and evolution of the continents.

6. ACKNOWLEDGEMENTS

We would like to thank Jean Hanley for careful preparation of the samples, Gwendolyn Hicks for enthusiastic assistance during her Federal work-study internship at Lamont, and Karen Benedetto for help with all the logistics. This study was funded by NSF OCE-1144759, NSF OCE-1144648 and EAR-1457293.

7. REFERENCES CITED

318 Behn, M. D., and Kelemen, P. B., 2006, Stability of arc lower crust: Insights from the Talkeetna
319 arc section, south central Alaska, and the seismic structure of modern arcs: *Journal of*
320 *Geophysical Research-Solid Earth*, v. 111, no. B11.

321 Cann, J. R., 1970, Upward movement of granitic magma, *Geological Magazine*, 335-340.

322 Carroll, M. R., and Wyllie, P. J., 1989, Experimental phase relations in the system tonalite-
323 peridotite-H₂O at 15 kb: implications for assimilation and differentiation processes near the
324 Crust-Mantle Boundary: *Journal of Petrology*, v. 30, p. 1351-1382.

325 Class, C., Miller, D. M., Goldstein, S. L., and Langmuir, C. H., 2000, Distinguishing melt and
326 fluid subduction components in Umnak Volcanics, Aleutian Arc: *Geochemistry Geophysics*
327 *Geosystems*, v. 1.

328 Defant, M. J., and Drummond, M. S., 1990, Derivation of some modern arc magmas by melting
329 of young subducted lithosphere: *Nature*, v. 347, no. 6294, p. 662-665.

330 DePaolo, D.J., 1981. Trace element and isotopic effects of combined wallrock assimilation and
331 fractional crystallization. *Earth and Planetary Science Letters* 53, 189-202.

332 Draper, D. S., and Green, T. H., 1999, P-T phase relations of silicic, alkaline, aluminous liquids:
333 new results and applications to mantle melting and metasomatism: *Earth and Planetary*
334 *Science Letters*, v. 170, no. 3, p. 255-268.

335 Fliedner, M.M., Klemperer, S.L., 1999. Structure of an island-arc: Wide-angle seismic studies in
336 the eastern Aleutian Islands, Alaska. *Journal of Geophysical Research-Solid Earth* 104,
337 10667-10694.

338 Gale, A., Dalton, C.A., Langmuir, C.H., Su, Y., Schilling, J.-G., 2013. The mean composition of
339 ocean ridge basalts. *Geochemistry, Geophysics, Geosystems* 14, 489-518.

340 Gerya, T. V., and Yuen, D. A., 2003, Rayleigh-Taylor instabilities from hydration and melting
341 propel 'cold plumes' at subduction zones: *Earth and Planetary Science Letters*, v. 212, no. 1-
342 2, p. 47-62.

343 Gill, J. B. (1981) *Orogenic Andesites and Plate Tectonics*, Springer Verlag, Berlin-Heidelberg,
344 392 pp.

345 Grove, T.L., Elkins-Tanton, L.T., Parman, S.W., Chatterjee, N., Muntener, O., Gaetani, G.A.,
346 2003. Fractional crystallization and mantle-melting controls on calc-alkaline differentiation
347 trends. *Contributions to Mineralogy and Petrology* 145, 515-533.

348 Grove, T.L., Baker, M.B., Price, R.C., Parman, S.W., Elkins-Tanton, L.T., Chatterjee, N.,
349 Muntener, O., 2005. Magnesian andesite and dacite lavas from Mt. Shasta, northern
350 California: products of fractional crystallization of H₂O-rich mantle melts. *Contributions to*
351 *Mineralogy and Petrology* 148, 542-565.

352 Hacker, B. R., Mehl, L., Kelemen, P. B., Rioux, M., Behn, M. D., and Luffi, P., 2008,
353 Reconstruction of the Talkeetna intraoceanic arc of Alaska through thermobarometry:
354 *Journal of Geophysical Research*, v. 113, no. B3.

355 Herzberg, C.T., Fyfe, W.S., Carr, M.J., 1983. Density constraints on the formation of the
356 continental Moho and crust. *Contributions to Mineralogy and Petrology* 84, 1-5.

357 Holbrook, W.S., Lizarralde, D., McGeary, S., Bangs, N., Diebold, J., 1999. Structure and
358 composition of the Aleutian island arc and implications for continental crustal growth.
359 *Geology* 27, 31-34.

360 Jagoutz, O., Muntener, O., Schmidt, M.W., Burg, J.P., 2011. The roles of flux- and
361 decompression melting and their respective fractionation lines for continental crust
362 formation: Evidence from the Kohistan arc. *Earth and Planetary Science Letters* 303, 25-36.

363 Jagoutz, O., and Schmidt, M. W., 2012, The formation and bulk composition of modern juvenile
364 continental crust: The Kohistan arc: *Chemical Geology*, v. 298, p. 79-96.

365 Janiszewski, H. A., Abers, G. A., Shillington, D. J., and Calkins, J. A., 2013, Crustal structure
366 along the Aleutian island arc: New insights from receiver functions constrained by active-
367 source data: *Geochemistry, Geophysics, Geosystems*, v. 14, no. 8, p. 2977-2992.

368 Jicha, B. R., Singer, B. S., Brophy, J. G., Fournelle, J. H., Johnson, C. M., Beard, B. L., Lapen,
369 T. J., and Mahlen, N. J., 2004, Variable impact of the subducted slab on Aleutian island arc
370 magma sources: Evidence from Sr, Nd, Pb, and Hf isotopes and trace element abundances:
371 *Journal of Petrology*, v. 45, no. 9, p. 1845-1875.

372 Jicha, B. R., Scholl, D. W., Singer, B. S., Yogodzinski, G. M., and Kay, S. M., 2006, Revised
373 age of Aleutian Island Arc formation implies high rate of magma production: *Geology*, v.
374 34, no. 8, p. 661-664.

375 Kay, R.W., 1978. Aleutian magnesian andesites: Melts from subducted Pacific ocean crust.
376 *Journal of Volcanology and Geothermal Research* 4, 117-132. Kay, R. W., Sun, S.-S., and
377 Lee-Hu, C. N., 1978, Pb and Sr isotopes in volcanic rocks from the Aleutian islands and
378 Pribilof Islands, Alaska,: *Geochim. Cosmochim. Acta*, v. 42, p. 263-272.

379 Kay, S. M., Kay, R. W., and Citron, G. P., 1982, Tectonic controls on tholeiitic and calc-alkaline
380 magmatism in the Aleutian Arc: *Journal of Geophysical Research*, v. 87, no. B5, p. 4051-
381 4071.

382 Kay, S. M., Kay, R. W., Brueckner, H. K., and Rubenstone, J. L., 1983, Tholeiitic Aleutian arc
383 plutonism - the Finger Bay pluton, Adak, Alaska: *Contributions to Mineralogy and*
384 *Petrology*, v. 82, no. 1, p. 99-116.

385 Kay, R. W., Rubenstone, J. L., and Kay, S. M., 1986, Aleutian terranes from Nd isotopes:
386 Nature, v. 322, no. 6080, p. 605-609.

387 Kay, S. M., Kay, R. W., Citron, G. P., and Perfit, M. R., 1990, Calc-alkaline plutonism in the
388 intra-oceanic Aleutian arc, Alaska: Geological Society of America Special Papers, Kay,
389 S.M., and Rapela, C. W., eds., v. 241, p. 233-256.

390 Kay, S. M., and Kay, R. W., 1994, Aleutian magmas in space and time, in Plafker, G. a. B., H.
391 C., ed., The Geology of Alaska, Volume G-1, Geological Society of America, p. 687-722.

392 Kelemen, P.B., Yogodzinski, G.M., 2007. High-magnesian andesite from Mount Shasta: A
393 product of magma mixing and contamination, not a primitive melt: COMMENT AND
394 REPLY. *Geology* 35, 149. Klemme, S., Blundy, J. D., and Wood, B. J., 2002, Experimental
395 constraints on major and trace element partitioning during partial melting of eclogite:
396 *Geochimica Et Cosmochimica Acta*, v. 66, no. 17, p. 3109-3123.

397 Kelemen, P. B., Yogodzinski, G. M., and Scholl, D. W., 2003a, Along-strike variation in lavas of
398 the Aleutian island arc: Implications for the genesis of high Mg# andesite and the
399 continental crust, in Eiler, J. M., ed., *Inside the Subduction Factory*, AGU Monograph
400 Volume 138, p. 224-276.

401 Kelemen, P. B., Hanghøj, K., and Greene, A. R., 2003b, One View of the Geochemistry of
402 Subduction-Related Magmatic Arcs, with an Emphasis on Primitive Andesite and Lower
403 Crust, in Holland, H. D., and Turekian, K. K., eds., *Treatise on Geochemistry*,: Oxford,
404 Elsevier-Pergamon, v. 3, p. 593-659.

405 Kelemen, P. B., Hanghøj, K., and Greene, A. R., 2014, One view of the geochemistry of
406 subduction-related magmatic arcs, with an emphasis on primitive andesite and lower crust,

407 in Holland, H. D., and Turekian, K. K., eds., *Treatise on Geochemistry* (Second Edition) :
408 Oxford, Elsevier, v. 4, p. 749-806.

409 Macpherson, C. G., Dreher, S. T., and Thirlwall, M. F., 2006, Adakites without slab melting:
410 High pressure differentiation of island arc magma, Mindanao, the Philippines: *Earth and*
411 *Planetary Science Letters*, v. 243, no. 3-4, p. 581-593.

412 Lee, C.-T.A., Lee, T.C., Wu, C.-T., 2014. Modeling the compositional evolution of recharging,
413 evacuating, and fractionating (REFC) magma chambers: Implications for differentiation of
414 arc magmas. *Geochimica et Cosmochimica Acta* 143, 8-22.

415 Miller, D. M., Goldstein, S. L., and Langmuir, C. H., 1994, Cerium lead and lead-isotope ratios
416 in arc magmas and the enrichment of lead in the continents: *Nature*, v. 368, no. 6471, p. 514-
417 520.

418 Osborn, E. F., 1959. Role of oxygen pressure in the crystallization and differentiation of basaltic
419 magma. *American Journal of Science*, 257(9), 609-647

420 Perfit, M. R., Brueckner, H., Lawrence, J. R., and Kay, R. W., 1980, Trace element and isotopic
421 variations in a zoned pluton and associated volcanic rocks, Unalaska Island, Alaska: A
422 model for fractionation in the Aleutian calcalkaline suite: *Contributions to Mineralogy and*
423 *Petrology*, v. 73, no. 1, p. 69-87.

424 Plank, T., and C H . Langmuir, 1993, Tracing trace elements from sediment input to volcanic
425 output at subduction zones, *Nature*, 362, 739-743.

426 Plank, T., V. Balzer and M. Carr, 2002, Nicaraguan volcanoes record paleoceanographic changes
427 accompanying closure of the Panama gateway, *Geology* 30, 1087–1090

428 Plank, T., 2014. 4.17 - The Chemical Composition of Subducting Sediments, in: Holland, H.D.,
 429 Turekian, K.K. (Eds.), *Treatise on Geochemistry* (Second Edition). Elsevier, Oxford, pp.
 430 607-629.

431 Ringwood, A. E., and Green, D. H., 1966, An experimental investigation of the Gabbro-Eclogite
 432 transformation and some geophysical implications: *Tectonophysics*, v. 3, no. 5, p. 383-427.

433 Rioux, M., Mattinson, J., Hacker, B., Kelemen, P., Blusztajn, J., Hanghoj, K., and Gehrels, G.,
 434 2010, Intermediate to felsic middle crust in the accreted Talkeetna arc, the Alaska Peninsula
 435 and Kodiak Island, Alaska: An analogue for low-velocity middle crust in modern arcs:
 436 *Tectonics*, v. 29.

437 Rudnick, R. L. and Gao, S. 2003, Composition of the Continental Crust, in in Holland, H. D.,
 438 and Turekian, K. K., eds., *Treatise on Geochemistry*: Oxford, Elsevier-Pergamon, v. 3, p.
 439 1-64.

440 Ruscitto, D.M., Wallace, P.J., Kent, A.J.R., 2010. Revisiting the compositions and volatile
 441 contents of olivine-hosted melt inclusions from the Mount Shasta region: implications for
 442 the formation of high-Mg andesites. *Contributions to Mineralogy and Petrology* 162,
 443 109-132.

444 Salters, V.J.M., Stracke, A., 2004. Composition of the depleted mantle. *Geochemistry*
 445 *Geophysics Geosystems* 5.

446 Sarbas, B., 2008. The GEOROC database as part of a growing geoinformatics network, in:
 447 Brady, S.R., Sinha, A.K. and Gundersen, L.C., Ed., *Geoinformatics 2008 - Data to*
 448 *Knowledge*, Proceedings: U.S. Geological Survey Scientific Investigations Report 2008-
 449 5172. USGS, pp. 42-43.

450 Shillington, D. J., Van Avendonk, H. J. A., Holbrook, W. S., Kelemen, P. B., and Hornbach, M.
451 J., 2004, Composition and structure of the central Aleutian island arc from arc-parallel wide-
452 angle seismic data: *Geochemistry Geophysics Geosystems*, v. 5.

453 Shimoda, G., Tatsumi, Y., Nohda, S., Ishizaka, K., and Jahn, B. M., 1998, Setouchi high-Mg
454 andesites revisited: geochemical evidence for melting of subducting sediments: *Earth and*
455 *Planetary Science Letters*, v. 160, no. 3–4, p. 479-492.

456 Singer, B. S., Jicha, B. R., Leeman, W. P., Rogers, N. W., Thirlwall, M. F., Ryan, J., and
457 Nicolaysen, K. E., 2007, Along-strike trace element and isotopic variation in Aleutian Island
458 arc basalt: Subduction melts sediments and dehydrates serpentine: *Journal of Geophysical*
459 *Research-Solid Earth*, v. 112, no. B6.

460 Sisson, T.W., Grove, T.L., 1993. Experimental investigations of the role of H₂O in calc-alkaline
461 differentiation and subduction zone magmatism. *Contributions to Mineralogy and Petrology*
462 113, 143-166.

463 Skora, S., Blundy, J., 2010. High-pressure Hydrous Phase Relations of Radiolarian Clay and
464 Implications for the Involvement of Subducted Sediment in Arc Magmatism. *Journal of*
465 *Petrology* 51, 2211-2243.

466 Stern, C. R., and Kilian, R., 1996, Role of the subducted slab, mantle wedge and continental
467 crust in the generation of adakites from the Andean Austral volcanic zone: *Contributions to*
468 *Mineralogy and Petrology*, v. 123, no. 3, p. 263-281.

469 Straub, S.M., Gomez-Tuena, A., Stuart, F.M., Zellmer, G.F., Espinasa-Perena, R., Cai, Y.,
470 Iizuka, Y., 2011. Formation of hybrid arc andesites beneath thick continental crust. *Earth*
471 *and Planetary Science Letters* 303, 337-347.

472 Suyehiro, K., Takahashi, N., Ariie, Y., Yokoi, Y., Hino, R., Shinohara, M., Kanazawa, T.,
473 Hirata, N., Tokuyama, H., and Taira, A., 1996, Continental crust, crustal underplating, and
474 low-Q upper mantle beneath an oceanic island arc: *Science*, v. 272, no. 5260, p. 390-392.

475 Tanaka, T., Togashi, S., Kamioka, H., Amakawa, H., Kagami, H., Hamamoto, T., Yuhara, M.,
476 Orihashi, Y., Yoneda, S., Shimizu, H., Kunimaru, T., Takahashi, K., Yanagi, T., Nakano, T.,
477 Fujimaki, H., Shinjo, R., Asahara, Y., Tanimizu, M., Dragusanu, C., 2000, JNdi-1: a
478 neodymium isotopic reference in consistency with LaJolla neodymium. *Chemical Geology*
479 168, 279-281.

480 Tatsumi, Y., 2000, Continental crust formation by delamination in subduction zones and
481 complementary accumulation of the enriched mantle I component in the mantle,
482 *Geochemistry, Geophysics, Geosystems (G-cubed)*, 1, 1-17.

483 Tatsumi, Y., Shukuno, H., Tani, K., Takahashi, N., Kodaira, S., and Kogiso, T., 2008, Structure
484 and growth of the Izu-Bonin-Mariana arc crust: 2. Role of crust-mantle transformation and
485 the transparent Moho in arc crust evolution: *Journal of Geophysical Research-Solid Earth*, v.
486 113, no. B2.

487 Taylor, S. R., and McLennan, S. M., 1995, The geochemical evolution of the continental-crust:
488 *Reviews of Geophysics*, v. 33, no. 2, p. 241-265.

489 Todt, W., Cliff, R.a., Hanser, A., Hofmann, A.W., 1996. Evaluation of a ^{202}Pb - ^{205}Pb double
490 spike for high-precision lead isotope analysis., in: Basu, A., Hart, S. (Eds.), *Geophysical*
491 *Monograph 95, Earth Processes: Reading the Isotopic Code*. AGU, Washington, D.C., pp.
492 429-437.

493 Vervoort, J.D., Plank, T., Prytulak, J., 2011. The Hf-Nd isotopic composition of marine
494 sediments. *Geochimica Et Cosmochimica Acta* 75, 5903-5926.

495 Yogodzinski, G. M., Rubenstone, J. L., Kay, S. M., and Kay, R. W., 1993, Magmatic and
 496 tectonic development of the western Aleutians: An oceanic arc in a strike-slip setting:
 497 Journal of Geophysical Research, v. 98, no. B7, p. 11807.

498 Yogodzinski, G. M., Kay, R. W., Volynets, O. N., Koloskov, A. V., and Kay, S. M., 1995,
 499 Magnesian Andesite in the Western Aleutian Komandorsky Region - Implications for Slab
 500 Melting and Processes in the Mantle Wedge: Geological Society of America Bulletin, v.
 501 107, no. 5, p. 505-519.

502 Yogodzinski, G. M., and Kelemen, P. B., 1998, Slab melting in the Aleutians: implications of an
 503 ion probe study of clinopyroxene in primitive adakite and basalt: Earth and Planetary
 504 Science Letters, v. 158, no. 1-2, p. 53-65.

505 Yogodzinski, G. M., Lees, J. M., Churikova, T. G., Dorendorf, F., Woerner, G., and Volynets, O.
 506 N., 2001, Geochemical evidence for the melting of subducting oceanic lithosphere at plate
 507 edges: Nature, v. 409, no. 6819, p. 500-504.

508 Yogodzinski, G.M., Kelemen, P.B., 2007. Trace elements in clinopyroxenes from Aleutian
 509 xenoliths: Implications for primitive subduction magmatism in an island arc. Earth and
 510 Planetary Science Letters 256, 617-632.

511 Yogodzinski, G. M., Vervoort, J. D., Brown, S. T., and Gersen, M., 2010, Subduction controls
 512 of Hf and Nd isotopes in lavas of the Aleutian island arc: Earth and Planetary Science
 513 Letters, v. 300, no. 3-4, p. 226-238.

514 Yogodzinski, G.M., Brown, S.T., Kelemen, P.B., Vervoort, J.D., Portnyagin, M., Sims, K.W.W.,
 515 Hoernle, K., Jicha, B.R., Werner, R., 2015. The Role of Subducted Basalt in the Source of
 516 Island Arc Magmas: Evidence from Seafloor Lavas of the Western Aleutians. Journal of
 517 Petrology.

Zimmer, M. M., Plank, T., Hauri, E. H., Yogodzinski, G. M., Stelling, P., Larsen, J., Singer, B., Jicha, B., Mandeville, C., and Nye, C. J., 2010, The role of water in generating the Calc-alkaline trend: new volatile data for Aleutian magmas and a new Tholeiitic Index: Journal of Petrology, v. 51, no. 12, p. 2411-2444.

8. FIGURE CAPTIONS

Figure 1. A) Schematic map of the Aleutian island arc, sampled areas are highlighted in black. B) Wt% SiO₂ versus Fe/Mg ratio of studied Aleutian magmas and literature data downloaded from the GEOROC database (<http://georoc.mpch-mainz.gwdg.de>; Sarbas, 2008). By convention, the Fe/Mg ratio is calculated using wt% MgO and FeO, with all Fe as FeO. The bulk continental crust composition is from Rudnick and Gao (2003). A complete list of references for the literature data is provided in the Supplementary Materials. Samples with Eu/Eu* > 1 are outlined in red (the calculation is given in the Fig. 2). C) Average trace element compositions of Aleutian plutons and lavas are fractionation corrected to Mg# = 0.5 following Behn and Kelemen (2006) and normalized to the composition of bulk continental crust (Rudnick and Gao, 2003). Extended trace element patterns are shown in Fig. S2.

Figure 2. Selected major and trace element compositions versus molar Mg# of Aleutian plutons and lavas, showing that the Eocene-Miocene plutonic rocks resemble western Aleutian lavas and match better with the continental crust than Holocene central and eastern Aleutian lavas. Molar Mg# = molar MgO/(molar MgO + molar FeO), with all Fe as FeO. Of the plutonics, only two of our studied samples and a few of the literature samples (outlined in red) show chemical signatures that indicate incorporation of cumulate plagioclase, such as elevated Al₂O₃ contents

and “positive” Eu anomalies ($\text{Eu}/\text{Eu}^* > 1$, where $\text{Eu}^* = \text{Eu}(\text{N}) / [\text{SQRT}(\text{Sm}(\text{N}) * \text{Gd}(\text{N}))]$ and elemental concentrations are normalized to chondrite compositions from McDonough and Sun, 1990). Overall, at a given Mg# the plutonic rocks show similar to higher mol% Na+K, and similar to lower wt% CaO than central and eastern Aleutian volcanics.

Figure 3. Present-day Nd and Pb isotope ratios of Aleutian magmas vs. longitude (A and B) and vs ages (C and D). Circles are central and eastern Aleutian volcanics (the two larger circles are from this study): Yellow= Rat and Delarof Islands, Green=Adak and Kanaga, Blue=Atka, Purple=Umnak, White=Unalaska. Error bars are smaller than the symbols. The other symbols are the same as in Fig. 1. ADK 53 is a primitive Miocene andesite from Mt Moffet on Adak Island (Kay, 1978; Kay et al. 1998). In C) and D), the Holocene volcanics are separated by location only without age differences.

Figure 4. A) and B) show present-day Nd, Pb, Hf isotopic compositions of Aleutian magmas. Central and eastern (C-E) Aleutian plutons fall between the fields defined by C-E Aleutian lavas and western Aleutian lavas. The ϵNd - ϵHf mantle-crust array is from Vervoort et al. (2011). The error bars are smaller than the symbols. The symbols are listed in Fig. 1.

Figure 5. AFC model showing the effect of assimilating arc crust or pre-existing oceanic crust, following DePaolo (1981), assuming $D_{\text{Nd}} = 0.06$, $D_{\text{Hf}} = 0.1$ (calculated assuming 50% plag, 25% cpx, 10% olivine and 10% opx, using Kd’s from Salters and Stracke, 2004). Assimilated crust compositions are modeled as $[\text{Hf}] = 2.5$, $\epsilon\text{Hf} = 15$; $[\text{Nd}] = 10.6$, $\epsilon\text{Nd} = 10$. The blue lines show the evolution trend of the magma as the fraction (F) of the remaining melt decreases (F values

are shown next to the tick marks along the AFC trend). Only $r = Ma/Mc = 0.8$ is shown, where Ma = mass assimilation rate and Mc = fractionation rate, following DePaolo (1981). Smaller r values would generate lower ϵ_{Nd} and ϵ_{Hf} values at a given F value.

Figure 6. Model results of different slab contributions for (a) ϵ_{Nd} vs Th/Nd and (b) ϵ_{Hf} vs Lu/Hf . Green lines show the effect of adding sediment melt to the subarc mantle wedge; red lines show addition of partial melt of the subducted oceanic crust ($F = 0.5\%$) and subducted sediments ($F = 5\%$) in a 4:1 ratio. Partial melt of subducted oceanic crust is modelled as a 0.5% equilibrium melt of the average MORB (EPR north of the equator segment average, Gale et al., 2013). Sediment melt is modeled as a 5% equilibrium partial melt of DSDP 183 sediments (Plank et al., 2014). For slab melts, we assumed $D_{Th}=0.2$, $D_{Nd}=1.4$, $D_{Hf}=0.5$, $D_{Lu}=50$ (Skora and Blundy, 2010). The mantle wedge is modeled as a spinel lherzolite that would generate typical EPR MORB via 10% equilibrium melting using D -values from Salters and Stracke (2004). After mixing the slab melt with the mantle wedge, the metasomatized mantle undergoes 15% equilibrium partial melting with spinel lherzolite D -values to generate the mixing lines shown in the figure (Salters and Stracke, 2004).

Figure 1. A) Schematic map of the Aleutian island arc, sampled areas are highlighted in black. B) $\text{Wt\% SiO}_2$ versus Fe/Mg ratio of studied Aleutian magmas and literature data downloaded from the GEOROC database (<http://georoc.mpch-mainz.gwdg.de>; Sarbas, 2008). By convention, the Fe/Mg ratio is calculated using $\text{wt\% MgO}$ and FeO , with all Fe as FeO . The bulk continental crust composition is from Rudnick and Gao (2003). A complete list of references for the literature data is provided in the Supplementary Materials. Samples with $\text{Eu}/\text{Eu}^* > 1$ are outlined in red (the calculation is given in the Fig. 2). C) Average trace element compositions of Aleutian plutons and lavas are fractionation corrected to $\text{Mg\#} = 0.5$ following Behn and Kelemen (2006) and normalized to the composition of bulk continental crust (Rudnick and Gao, 2003). Extended trace element patterns are shown in Fig. S2.

Figure 2. Selected major and trace element compositions versus molar Mg# of Aleutian plutons and lavas, showing that the Eocene-Miocene plutonic rocks resemble western Aleutian lavas and match better with the continental crust than Holocene central and eastern Aleutian lavas. Molar Mg# = molar MgO/(molar MgO + molar FeO), with all Fe as FeO. Of the plutonics, only two of our studied samples and a few of the literature samples (outlined in red) show chemical signatures that indicate incorporation of cumulate plagioclase, such as elevated Al₂O₃ contents and “positive” Eu anomalies ($Eu/Eu^* > 1$, where $Eu^* = Eu(N) / [SQRT(Sm(N) \cdot Gd(N))]$ and elemental concentrations are normalized to chondrite compositions from McDonough and Sun, 1990). Overall, at a given Mg# the plutonic rocks show similar to higher mol% Na+K, and similar to lower wt% CaO than central and eastern Aleutian volcanics.

Figure 3. Present-day Nd and Pb isotope ratios of Aleutian magmas vs. longitude (A and B) and vs ages (C and D). Circles are central and eastern Aleutian volcanics (the two larger circles are from this study): Yellow= Rat and Delarof Islands, Green=Adak and Kanaga, Blue=Atka, Purple=Umnak, White=Unalaska. Error bars are smaller than the symbols. The other symbols are the same as in Fig. 1. ADK 53 is a primitive Miocene andesite from Mt Moffet on Adak Island (Kay, 1978; Kay et al. 1998). In C) and D), the Holocene volcanics are separated by location only without age differences.

Figure 4. A) and B) show present-day Nd, Pb, Hf isotopic compositions of Aleutian magmas. Central and eastern (C-E) Aleutian plutons fall between the fields defined by C-E Aleutian lavas and western Aleutian lavas. The $\epsilon_{\text{Nd}}-\epsilon_{\text{Hf}}$ mantle-crust array is from Vervoort et al. (2011). The error bars are smaller than the symbols. The symbols are listed in Fig. 1.

Figure 5. AFC model showing the effect of assimilating arc crust or pre-existing oceanic crust, following DePaolo (1981), assuming $D_{Nd} = 0.06$, $D_{Hf} = 0.1$ (calculated assuming 50% plag, 25% cpx, 10% olivine and 10% opx, using K_d 's from Salters and Stracke, 2004). Assimilated crust compositions are modeled as $[Hf] = 2.5$, $\epsilon_{Hf} = 15$; $[Nd] = 10.6$, $\epsilon_{Nd} = 10$. The blue lines show the evolution trend of the magma as the fraction (F) of the remaining melt decreases (F values are shown next to the tick marks along the AFC trend). Only $r = Ma/Mc = 0.8$ is shown, where Ma = mass assimilation rate and Mc = fractionation rate, following DePaolo (1981). Smaller r values would generate lower ϵ_{Nd} and ϵ_{Hf} values at a given F value.

Figure 6. Model results of different slab contributions for (a) ϵ_{Nd} vs Th/Nd and (b) ϵ_{Hf} vs Lu/Hf . Green lines show the effect of adding sediment melt to the subarc mantle wedge; red lines show addition of partial melt of the subducted oceanic crust ($F = 0.5\%$) and subducted sediments ($F = 5\%$) in a 4:1 ratio. Partial melt of subducted oceanic crust is modelled as a 0.5% equilibrium melt of the average MORB (EPR north of the equator segment average, Gale et al., 2013). Sediment melt is modeled as a 5% equilibrium partial melt of DSDP 183 sediments (Plank et al., 2014). For slab melts, we assumed $D_{Th}=0.2$, $D_{Nd}=1.4$, $D_{Hf}=0.5$, $D_{Lu}=50$ (Skora and Blundy, 2010). The mantle wedge is modeled as a spinel lherzolite that would generate typical EPR MORB via 10% equilibrium melting using D-values from Salters and Stracke (2004). After mixing the slab melt with the mantle wedge, the metasomatized mantle undergoes 15% equilibrium partial melting with spinel lherzolite D-values to generate the mixing lines shown in the figure (Salters and Stracke, 2004).