Bahamian speleothem reveals temperature decrease associated with Heinrich stadials

Monica M. Arienzo^{1,2*}, Peter K. Swart¹, Ali Pourmand³, Kenny Broad¹, Amy C. Clement¹, Lisa N. Murphy¹, Hubert B. Vonhof⁴, Brian Kakuk⁵

¹Rosenstiel School of Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, FL, USA

²Now at Division of Hydrologic Sciences, Desert Research Institute, Reno, NV, USA

³Neptune Isotope Lab, Rosenstiel School of Marine and Atmospheric Science, University of Miami, 4600 Rickenbacker Causeway, Miami, FL, USA

⁴Faculty of Earth and Life Sciences, Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands

⁵Bahamas Underground, Marsh Harbor, Bahamas

*Corresponding author marienzo@dri.edu

Keywords: Speleothem, Heinrich event, stable isotope, fluid inclusion, Bahamas

Abstract:

Temperature reconstructions across Heinrich stadials 1-3 are presented from an absolute-dated speleothem from Abaco Island in the Bahamas to understand the nature of climate change across these intervals in the subtropical Atlantic. The stalagmite carbonate record, dated by the U-Th geochronometry technique, includes higher $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values within Heinrich stadials 1, 2, and 3 followed by rapid declines at the end of the stadials. To aid in the interpretation of these results, the $\delta^{18}\text{O}$ of fluid inclusions associated with the Heinrich stadials were also analyzed. These measurements, which allowed for the relative influence of temperature and $\delta^{18}\text{O}$ of precipitation to be distinguished, demonstrate minimal changes in the $\delta^{18}\text{O}$ of fluid inclusions, suggesting that changes in the $\delta^{18}\text{O}$ values of the speleothem carbonate associated with Heinrich stadials 1 – 3 are principally driven by an average $\sim 4^\circ\text{C}$ temperature decrease, rather than a change in the $\delta^{18}\text{O}$ of the rainfall (hence rainfall amount). These findings support previous work in the North Atlantic and are consistent with the climate response to a weakening of the Atlantic meridional overturning circulation.

1. Introduction

Ice core and deep sea sediment records of the last 65,000 years show 18 periods of millennial scale climate events known as the Dansgaard/Oeschger (D/O) cycles and 6 Heinrich stadials (Dansgaard et al., 1984). Dansgaard/Oeschger events in Greenland are millennial scale alternations between warm (interstadial) and cold (stadial) periods (Bond et al., 1997). Heinrich events are characterized in the North Atlantic by cold periods, and are recognized in the

sedimentary record as eroded terrigenous materials (Ice Rafted Debris, IRD) deposited in the North Atlantic by ice bergs upon melting (Bond et al., 1997; Heinrich, 1988). Heinrich stadials are associated with a slowdown of the Atlantic meridional overturning circulation (AMOC) and an inter-hemispheric climate response (Wolff et al., 2010; McManus et al., 2004). Observations and models (Zhang and Delworth, 2005) support reduction in sea surface temperatures (SSTs) associated with Heinrich stadials, thought to be caused by reduced northward heat transport, driven by the slowdown of the AMOC (Clement and Peterson, 2008) or the increase in sea/land ice (Chiang and Bitz, 2005). Reduced Northern Hemisphere SSTs led to the southward shift in the intertropical convergence zone (ITCZ) and drier conditions in the tropical Northern Hemisphere (Chiang and Bitz, 2005; Stager et al., 2011). Recent studies have also demonstrated the impact of varying height of the Laurentide ice sheet as a climate forcing during Heinrich stadial events (Roberts et al., 2014). During interstadial periods (*i.e.* D/O interstadial events), the inverse occurs with a poleward shift of the Northern Hemisphere summer ITCZ and the jet streams (Asmerom et al., 2010). However, the exact mechanisms driving these events are still not well understood (Clement and Peterson, 2008).

Various types of paleoclimate data support the global response to North Atlantic Heinrich stadials, as well as the abrupt nature of the events. Paleoclimate records suggest that the global signature of Heinrich stadials includes: a drier Europe (Genty et al., 2003), weaker Asian monsoon (Wang et al., 2001), wetter southwestern North America (Asmerom et al., 2010; McGee et al., 2012), drier northern South America (Peterson et al., 2000), wetter southern South America (Kanner et al., 2012), an overall drier tropical Asia and Africa (Stager et al., 2011), and a gradually warming Antarctica (Wolff et al., 2010) (for a review of paleoclimate data across Heinrich stadials see Clement and Peterson (2008)). While a comprehensive picture of climate

across North Atlantic Heinrich stadials is emerging from records in both hemispheres, very few studies have been conducted in the subtropical western Atlantic (Grimm et al., 2006; Lachniet et al., 2013; Sachs and Lehman, 1999), which may be an important area for detecting the global propagation of these events and constraining climate models. In this study, geochemical data obtained from a speleothem spanning Heinrich stadials 1-3 are presented from a cave in Abaco Island, Bahamas. Each Heinrich stadial event exhibited unique characteristics, supporting other studies showing Heinrich stadial 1 to be the strongest event (Stager et al., 2011), with almost complete shutdown in AMOC (McManus et al., 2004).

1.1 Speleothems as paleoclimate archives

Speleothems have proven to be valuable archives for paleoclimate reconstructions, particularly for the study of climate variability on millennial timescales (Asmerom et al., 2010; Wang et al., 2001). Stable isotope ratios of oxygen and carbon ($\delta^{18}\text{O}$ and $\delta^{13}\text{C}$) of the carbonate are the most common geochemical proxies analyzed within speleothems. There are several climate factors that can lead to changes in the $\delta^{18}\text{O}$ values of the carbonate record. In the case of tropical speleothems, the $\delta^{18}\text{O}$ of the carbonate is typically interpreted to be driven primarily by the $\delta^{18}\text{O}$ of the rainfall and/or the temperature of the cave (Kanner et al., 2012; van Breukelen et al., 2008), and studies have shown that cave drip water has $\delta^{18}\text{O}$ values similar to local precipitation (Tremaine et al., 2011; van Breukelen et al., 2008). In the Bahamas and throughout the Caribbean and south Florida, there is an inverse relationship between the amount of rainfall and the $\delta^{18}\text{O}$ of the rainwater (Baldini et al., 2007; van Breukelen et al., 2008), and therefore the amount of rainfall (the amount effect) is considered to exert the main control on the $\delta^{18}\text{O}$ composition of the rainfall precipitation (Dansgaard, 1964). However, distinguishing the

competing influences of temperature and $\delta^{18}\text{O}$ of the rainwater is inherently complex when interpreting the carbonate $\delta^{18}\text{O}$ results. One approach to address the confounding influences of temperature and water $\delta^{18}\text{O}$ is through the $\delta^{18}\text{O}$ analysis of fluid inclusions. These inclusions are microscopic water filled cavities located within the speleothem calcite mineral structure which contain drip water trapped at the time the stalagmite formed. As suggested above, the drip water is thought to directly represent precipitation, and therefore the isotopic analysis of the trapped water provides a direct measure of the temporal changes in the $\delta^{18}\text{O}$ of rainfall (hence rainfall amount). Determining the $\delta^{18}\text{O}$ value of both the trapped fluid and that of the accompanying mineral allows the temperature at the time of speleothem formation to be calculated (van Breukelen et al., 2008).

In contrast, the $\delta^{13}\text{C}$ of the carbonate is a function of the type and amount of vegetation above the cave, root respiration, organic material decomposition, the amount of water/rock interactions and prior calcite precipitation (Fairchild et al., 2006). Variation in the biogenic CO_2 component of the carbon is in turn influenced by precipitation amount and temperature (Genty et al., 2003). Generally the partial pressure of CO_2 ($p\text{CO}_2$) is expected to show an inverse correlation with $\delta^{13}\text{C}$ values in the soil horizon. Changes in the amount of rainfall or temperature could also alter the ratio of C_3 and C_4 plants, causing further isotopic changes, particularly over longer timescales (Fairchild et al., 2006). Cave ventilation may also exert control over the $\delta^{13}\text{C}$ of the CO_2 in the cave and therefore ultimately the $\delta^{13}\text{C}$ in the cave fluids (Tremaine et al., 2011). Through the combination of both $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ of the carbonate as well as $\delta^{18}\text{O}$ of the fluid inclusions, variations in the temperature and amount of rainfall can be estimated over the sample time period.

1.2 Previous work on Bahamian speleothems

Several studies have been conducted on the formation, growth, age, and geochemistry of Bahamian speleothems, with most of the work carried out on samples from caves currently submerged in seawater (Richards et al., 1994 and references therein). These speleothems formed during previous sea level low-stands and stopped forming when rising sea level flooded the caves, making the speleothems potential archives of paleo-sea level. However, it has also been shown in some instances that the speleothems ceased forming prior to seawater flooding, presumably as a result of drier conditions (Richards et al., 1994). Richards et al. (1994) also found no geochemical evidence of alteration in samples from submerged caves. Stalagmites from the Bahamas have been additionally utilized as recorders of past atmospheric $\Delta^{14}\text{C}$ concentration for radiocarbon calibrations (Hoffmann et al., 2010, Beck et al., 2001).

The aim of this work is to determine the paleoclimatic changes associated with Heinrich stadials over the last ~32,000 years by applying multiple geochemical tools. This study represents the first high resolution paleoclimate reconstruction utilizing a Bahamian speleothem and the first multi-collector ICP-MS dated speleothem from this region. Additionally, the combination of both stable isotope analyses of the carbonate and fluid inclusions is unique and the resolution of the fluid inclusion analyses for a stalagmite record from this time period is unprecedented. Finally, considering that relatively few studies have been conducted in the subtropical western Atlantic (Escobar et al., 2012; Grimm et al., 2006; Hagen and Keigwin, 2002; Hodell et al., 2012; Keigwin and Jones, 1994; Sachs and Lehman, 1999), and even fewer studies are from terrestrial records, this study offers the opportunity to better understand abrupt climate change at this location during the Pleistocene and its relationship to the climate of the North Atlantic.

2. Sample Locality and Methods

2.1 Regional setting

The modern climate of the Bahamas is primarily controlled by the easterly trade winds and winds from the west are limited to frontal passages (Baldini et al., 2007). The annual variation in air temperature ranges from 22 to 28 °C (Baldini et al., 2007). There are distinct wet and dry seasons with the wetter period between April to December driven by the Bermuda high.

The stalagmite (sample AB-DC-09) was collected from a currently submerged cave located in the middle of southern Abaco Island, Bahamas (N26°14, W77°10) in July of 2007 from a depth of 16.5 m below current sea level (Figure 1). The cave is accessed through a collapsed sinkhole and consists of three laterally extensive levels at ~22, 33.5 and 45 m below sea level. The overlying bedrock of the cave is composed of Pleistocene limestone aeolianites and marine limestones (Walker et al., 2008). After collection, the speleothem was sectioned along the main growth axis (Figure 2a) and sampled in the central region for U-Th geochronometry, stable C and O isotopes ($\delta^{13}\text{C}_c$ and $\delta^{18}\text{O}_c$), and $\delta^{18}\text{O}_w$ of fluid inclusions. Sample AB-DC-09 has a total length of 23 cm and is comprised of dense milky white calcite with no evidence of post-depositional diagenesis (Figure 2a).

2.2 Geochemical methods

U-Th geochronometry: Twenty-three U-Th dates (Table 1) were measured at the Neptune Isotope Laboratory of the University of Miami - RSMAS utilizing a ThermoFisher-Neptune Plus multi-collector ICP-MS. Details of the U-Th geochronometry technique and propagation of random and systematic uncertainties are discussed elsewhere (Pourmand et al., 2014). Briefly, 100-200 mg of the sample was hand-drilled. Samples were then completely digested in 5 mL of

146 6 mol L⁻¹ HNO₃ followed by the addition of ~0.5 g of a previously calibrated ²²⁹Th-²³³U-²³⁶U
147 spike mixture. The concentrations of U and Th in the spike solution were determined by isotope
148 dilution mass spectrometry against a mixture of U and Th mono-elemental standard solutions
149 and gravimetrically prepared U standard from ~1g of U metal (full details on spike calibration
150 are provided in Pourmand et al. (2014)). To separate U and Th from matrix elements (mainly Ca,
151 Sr and Mg), the sample and spike mixture were loaded on to pre-packed 2-mL cartridges of
152 U/TEVA resin (particle size 50–100 µm) from Eichrom Inc. After removal of the matrix
153 elements, Th and U were quantitatively separated from each other in 10 mL of 3 and 0.1 mol L⁻¹
154 HCl, respectively.

155 Higher abundance U (²³³U, ²³⁵U, ²³⁶U and ²³⁸U) and Th (²²⁹Th, ²³²Th) isotopes were
156 measured in Faraday collectors while ²³⁴U and ²³⁰Th isotopes were measured in the secondary
157 electron multiplier (SEM) after the ion beams passed through a Retardation Potential Quadrupole
158 (RPQ) filter to reduce the abundance sensitivity effect. Each U and Th measurement was
159 bracketed by two measurements of inlet system memory from previous runs and measurements
160 of CRM-112A and IRMM-035 standard solutions spiked with IRMM-3636a ²³³U-²³⁶U double
161 spike. Data acquisition for U and Th isotopes and the half-masses were exported to Excel
162 spreadsheet templates and processed using an open-source algorithm in Mathematica as
163 presented in Pourmand et al. (2014) for calculating the U and Th concentrations, age and
164 uncertainties (reported here as 95% confidence interval). A significant advantage of this
165 technique is consideration for U and Th isotope covariance and propagation of uncertainties on
166 decay constants and other sources of random and systematic uncertainties on corrected ages,
167 which are especially crucial for comparison with age models that are not based on U-Th
168 geochronometry. The half-life values used are those presented in Edwards et al. (2003) and

Fietzke et al. (2005). The concentration of U and Th in procedural blanks, which were processed every five samples, were below 1 and 10 pg, respectively, and therefore no blank corrections were necessary. All ages are reported in years before present (yr BP).

Isotopic sampling on the carbonate: Sampling of the speleothem carbonate for $\delta^{13}\text{C}_\text{c}$ and $\delta^{18}\text{O}_\text{c}$ (calcite) was carried out using a New-Wave computerized micromill at both high and low resolution. Low resolution sampling consisted of one individual sample every 1000 μm (Figure 2a & b). The low resolution analysis revealed three areas of rapid and significant isotopic shifts. These areas were subsequently analyzed at a higher resolution (20 μm sampling interval) for a total of ~550 high resolution samples (Figure 2b). Isotopic measurements were made using a Kiel III interfaced with a Thermo-Finnigan Delta Plus Mass Spectrometer at the University of Miami. All data have been corrected for isobaric inferences at mass 45 and 46 and $\delta^{13}\text{C}_\text{c}$ and $\delta^{18}\text{O}_\text{c}$ values are reported relative to Vienna Pee Dee Belemnite (VPDB). The error for these analyses is <0.1‰ as determined by repeated measurement of an internal standard.

Fluid inclusions: Thirty-one oxygen and hydrogen isotopic analyses of the fluids trapped within inclusions ($\delta^{18}\text{O}_\text{w}$ and $\delta^2\text{H}_\text{w}$) were carried out at the University of Miami using a cavity ring-down spectroscopy instrument (L2130-i Picarro) as outlined in Arienzo et al. (2013). This system allowed for the determination of $\delta^{18}\text{O}$ of the water ($\delta^{18}\text{O}_\text{w}$) and $\delta^2\text{H}_\text{w}$ of the fluid inclusions from approximately 0.2 g samples, which on average yielded 0.4 μL of water from AB-DC-09 speleothem. Sampling for fluid inclusion analysis was conducted along the center of the growth axis, utilizing a band saw to extract 0.6 – 1.0 g cubes of calcite which were subsequently divided into ~0.2 g cubes for replicate analysis. The isotopic compositions of the fluid inclusions are reported relative to the Vienna Standard Mean Ocean Water (VSMOW).

Each sample was analyzed 1-3 times with an average error of 0.5 ‰ for $\delta^{18}\text{O}_w$ and 2.0 ‰ for $\delta^2\text{H}_w$ values (Arienzo et al., 2013). As a result of the larger sample size, average sampling was conducted every 0.75 cm. The $\delta^{18}\text{O}_c$ of the calcite hosting the inclusion was measured on the crushed residue at the University of Miami using a Thermo-Finnigan Delta Plus mass spectrometer (see above).

3. Results

3.1 Age model

Unsupported ^{230}Th : An important variable in U-Th geochronometry is the contribution of excess (unsupported) ^{230}Th to the U series equilibrium clock (Edwards et al., 2003). Isochron techniques and direct measurements from modern cave deposits provide estimates of the initial ^{230}Th . Previous studies of Bahamian speleothems forming over the last glacial maximum (LGM) utilizing isochrons suggested relatively high initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratios, ranging from 7.8 ± 4.0 (Hoffmann et al., 2010) to 18.7 ± 2.9 (Beck et al., 2001). These samples were collected from Grand Bahama Island, located to the north of Abaco Island and were forming over the LGM. Rather than conducting isochrons, which require identifying cogenetic samples of homogeneous parent and daughter composition within the same depositional horizon (Edwards et al., 2003), we have analyzed modern cave calcites from an actively forming cave in Eleuthera, Bahamas in order to determine a representative ^{230}Th detrital activity ratio. The results measured in the depositional slides yielded an average initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratio of 2.2 ± 0.6 (supplementary material). These findings from speleothems suggest that, for the Bahamas, there is a wide range of measured initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratios from 2.2 to 18.7. The published data with higher initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratios were from samples that were forming over the

LGM, suggesting the cave had a higher Th input during this period than the modern samples from Eleuthera. Other studies have characterized the U and Th concentrations in carbonates and waters of the Bahamas, and these studies demonstrate wide variability in concentrations and several potential sources of detrital Th, including aeolian inputs (*e.g.*, Robinson et al., 2004).

Considering that stalagmites from Abaco Island have yet to be studied, comparative data for this island does not exist. Therefore, from the results outlined above, the activity ratio of 3.7 ± 0.6 for initial $^{230}\text{Th}/^{232}\text{Th}$ is considered a reasonable estimate as this value falls in the range of values for the Bahamas, and additionally results in good agreement with previously published records. Nevertheless, we further assess the contribution of the initial, unsupported ^{230}Th by a sensitivity test whereby the ages were propagated at various initial ($^{230}\text{Th}/^{232}\text{Th}$) ratios (Supplementary Table 2). This sensitivity study demonstrates that the initial Th value does not significantly change the timing of Heinrich stadials 1 or 2, however, a lower activity ratio (*e.g.*, 0.6) resulted in the timing of Heinrich stadial 3 to be an unreasonable age with respect to published ages for the event. Varying the initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratio impacts the age calculated on the oldest U-Th ages, and these ages do consist of the largest errors. Given the good agreement with other published records, the activity ratio of 3.7 ± 0.6 for initial $^{230}\text{Th}/^{232}\text{Th}$ is considered a reasonable estimate.

It must be noted that we make the assumption that the initial Th activity ratio did not change throughout the deposition of the stalagmite. When propagating the U-Th ages using an activity ratio of 3.7 ± 0.6 for initial $^{230}\text{Th}/^{232}\text{Th}$, three ages showed reversals (Table 1). Two of the ages located ~5 cm from the top were younger than the previous ages, resulting in a reversal (Table 1). This reversal may be driven by a reduction in the initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratio

component during this time period and therefore we do not include these two dates in the age model. One U-Th age 11 cm from the top was older and resulted in an age reversal, possibly suggesting an increase in the initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratio component at this time. Considering the good agreement between surrounding ages, we discount this age as well and do not include it in the final age model.

Age model: Utilizing twenty U-Th ages the final age model was calculated using the Constructing Proxy-Records from Age models (COPRA) program from Breitenbach et al. (2012) (Figure 3). The age model was calculated in MATLAB[®] using the Piecewise Cubic Hermite Interpolating Polynomial (PCHIP) and 2000 Monte Carlo simulations to assign a precise timescale (Figure 3). When plotting the age model with other paleoproxies from the region, we find good agreement in the timing of isotopic shifts between records (discussed in detail in section 4.4), which further validates the decision to utilize a value of 3.7 ± 0.6 for initial $^{230}\text{Th}/^{232}\text{Th}$ activity ratio.

The age model results indicate that the stalagmite grew between 13.8 to 32 kyr before present (Figures 3 & 4). There is no evidence for any growth hiatuses throughout the sample from either the age measurements or petrographic thin section analysis, however, there is evidence for periods of decreased growth rates from the U-Th geochronometry. The lowest growth rate occurred from 22 to 26 kyr BP (~ 14.5 to 15.5 cm from top) and two additional periods of decreased growth rate from 14.9 to 17.3 kyr BP (~ 4.2 to 6 cm from the top) and 27.8 to 30 kyr BP (~ 21 to 23 cm from the top). There is evidence of three small layers of darker laminations at ~5, 14.5 and 22 cm from the top (Figure 2a & b) which correlate with the three short periods of reduced calcite precipitation. These layers may represent brief reductions in

growth or hiatuses, however do not significantly impact the interpretation of the results (Supplementary Figures 3 & 4).

3.2 Tests for kinetic fractionation

Several studies have shown that the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ can be affected by kinetically driven fractionations during the deposition of the carbonate which can overwhelm environmental and climate signals (Hendy, 1971; Lachniet, 2009). An approach often used to examine the role of kinetic effects is known as the “Hendy test” (Hendy, 1971) in which the lateral variation in a chemical parameter within the speleothem is measured. Assuming that sampling can be carried out along a growth band, the following criteria must be met for the Hendy test: i) an absence of lateral increases in $\delta^{18}\text{O}_\text{c}$ and $\delta^{13}\text{C}_\text{c}$ values, and ii) a low correlation between $\delta^{18}\text{O}_\text{c}$ and $\delta^{13}\text{C}_\text{c}$ values. If these conditions are fulfilled, then it is assumed that minimal kinetic fractionation has taken place. However, studies have demonstrated that this test may not be sufficient to rule out kinetic fractionation (Mühlinghaus et al., 2009). Other authors have suggested a more robust method is to conduct replication tests, in which multiple stalagmites from the same cave are measured and if the $\delta^{13}\text{C}_\text{c}$ and $\delta^{18}\text{O}_\text{c}$ trends are replicated, this supports a lack of kinetic fractionation (Dorale and Liu, 2009). For this study, we have conducted the Hendy test and present two locations across which lateral variation of $\delta^{18}\text{O}_\text{c}$ and $\delta^{13}\text{C}_\text{c}$ values were measured. Sampling for kinetic fractionation was conducted at one millimeter increments from the center of the growth axis progressing outward along a growth band. As shown in Figure 2 c and d, tests for kinetic fractionation demonstrate low correlation and no significant increase for both $\delta^{13}\text{C}_\text{c}$ and $\delta^{18}\text{O}_\text{c}$ across a lamina. This demonstrates that significant fractionation for $\delta^{13}\text{C}_\text{c}$ and $\delta^{18}\text{O}_\text{c}$ has not occurred for this speleothem, additionally supported by visual and petrographic observation of the sampled calcite, which is free of evidence of dissolution. In addition, in a cave

from Eleuthera in the Bahamas where there is currently active stalagmite formation, the measured $\delta^{18}\text{O}$ of the calcite, the temperature of the cave, and drip water $\delta^{18}\text{O}$ all support the idea that the calcite is precipitated in equilibrium (Supplementary Text).

For fluid inclusions, isotopic fractionation after deposition can occur through fractionation between the water and the surrounding calcite, thereby altering the isotopic composition of the trapped water. However, such fractionation in the humid subtropics should be minimal (van Breukelen et al., 2008). The fluid inclusions are representative of paleoprecipitation, therefore results should fall on the global meteoric water line (GMWL), which defines the linear $\delta^{18}\text{O}_w$ versus $\delta^2\text{H}_w$ relationship for all meteoric waters (Craig, 1961). Fluid inclusion results plot near the GMWL, suggesting that the $\delta^{18}\text{O}_w$ and $\delta^2\text{H}_w$ values have not been affected by fractionation and the data are therefore representative of the original meteoric water composition (Figure 5).

Precipitation rate can influence kinetic fractionation if the rate of precipitation is faster than the rate of $\text{HCO}_3^-(\text{aq})$ re-equilibration, which may impact the fluid inclusion results. However, given the lack of evidence for kinetic fractionation from the carbonate isotopes, it is not thought that kinetic fractionation is a significant driver of the chemistry. In addition, Baldini et al. (2007) demonstrated that the $\delta^{18}\text{O}_w$ of rainwater from San Salvador, Bahamas varied between -4.4‰ to -1.2‰ VSMOW, within the range of values measured from the fluid inclusions in this study (Table 2).

3.3 Carbon and oxygen isotopes of carbonate

The speleothem AB-DC-09 was sampled at both high (one sample every ~2 years) and low resolution (one sample every ~50 years). To aid with the analysis of the data, the $\delta^{13}\text{C}$ and

$\delta^{18}\text{O}$ carbonate isotope results are plotted with a 50-year interpolation (Figure 4a & b). A comparison of the $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ of the speleothem calcite ($\delta^{18}\text{O}_c$ and $\delta^{13}\text{C}_c$) with the derived age model demonstrates significant variations in the isotopic value of the calcite (Figure 4a & b). The variation in the $\delta^{13}\text{C}_c$ results are 6.2 ‰ for the sample, with the most positive $\delta^{13}\text{C}_c$ value of +3.2 ‰ VPDB occurring at 31.7 ± 1 kyr BP. Above average $\delta^{13}\text{C}_c$ values are also found at 15 ± 0.2 kyr and 25 ± 0.2 kyr BP (Figure 4a). Similar results are found in the $\delta^{18}\text{O}_c$ record with the most positive $\delta^{18}\text{O}_c$ value of -0.2 ‰ VPDB at 15 ± 0.2 kyr and above average $\delta^{18}\text{O}_c$ values at 24.1 ± 0.2 kyr and 31.5 ± 1 kyr BP (Figure 4b). Overall, a total of 3.5 ‰ variation is observed in the $\delta^{18}\text{O}_c$ results. A similar trend for each positive C and O isotope event is observed, with a gradual increase leading to a maximum, followed by a more rapid decline. For the 15 kyr event, the gradual increase begins at 17.4 kyr BP and reaches a maximum at ~15 kyr BP, followed by a rapid decrease (~200 years) in $\delta^{18}\text{O}_c$ and $\delta^{13}\text{C}_c$ values. The trend is also observed across the 24.5 kyr event with an increase beginning at ~26 kyr BP to a maximum, followed by a decrease. For the oldest event, maximum $\delta^{18}\text{O}_c$ and $\delta^{13}\text{C}_c$ values occurred at ~31.5 kyr and declined over ~200 years after the maximum. In summary, the carbonate results support three unique shifts, of which the least significant in both $\delta^{13}\text{C}_c$ and $\delta^{18}\text{O}_c$ is the 24.5 kyr BP event (Figure 4a & b).

3.4 Fluid inclusions

A comparison of the $\delta^{18}\text{O}_w$ measured on the fluid inclusions with the derived age model from U-Th geochronometry demonstrates minimal variations in the isotopic value of the fluids (Figure 4c). The $\delta^2\text{H}_w$ and $\delta^{18}\text{O}_w$ measured on the fluid inclusions support variations of 11.7 ‰ and 1.6 ‰ respectively, over the studied interval with average values of -8.6 ‰ $\delta^2\text{H}_w$ and -2.5 ‰ $\delta^{18}\text{O}_w$ VSMOW (Table 2). The variability in the $\delta^{18}\text{O}_w$ value is relatively small compared to the changes in the $\delta^{18}\text{O}_c$ value (~3.5 ‰, Figure 4c). The data support a minimum $\delta^{18}\text{O}_w$ value at 25.3

kyr BP and a maximum $\delta^{18}\text{O}_w$ value at 30.7 kyr BP. Fluid inclusion samples formed during Heinrich stadial 1 include two samples at 15.5 and 16.6 kyr BP, Heinrich stadial 2 includes one sample at 25.3 kyr BP and Heinrich stadial three includes one sample at 31.7 kyr BP (Figure 3).

4. Discussion

4.1 Climate interpretation of the oxygen isotope record: Aridity vs temperature

The $\delta^{18}\text{O}_w$ values of the fluid inclusions shows an overall 1.6 ‰ decline from 30.7 to 25.3 kyr BP. Decreases in $\delta^{18}\text{O}_w$ are generally attributed to the ‘amount’ effect (Dansgaard, 1964) and therefore this change suggests an increase in the amount of precipitation over this period, an interpretation supported by a progressively increasing growth rate. The $\delta^{18}\text{O}_c$ record shows a maxima at $\sim 15 \pm 0.2$, 24.1 ± 0.2 , and 31.5 ± 1 kyr BP and the timing of these events is associated with Heinrich stadials 1, 2, and 3 (Figure 6). In contrast to a significant increase of $\delta^{18}\text{O}_c$ values within each of the three Heinrich stadials, the $\delta^{18}\text{O}_w$ data obtained from the fluid inclusions reveals only minimal changes. For example, the average change in the $\delta^{18}\text{O}_w$ value across Heinrich stadial 1, from 17.1 to 14.8 kyr BP, is 0.2 ‰ which is less than the analytical uncertainty. The average change from 26 to 23 kyr BP is 0.4 ‰ for $\delta^{18}\text{O}_w$ value (Figure 4 c). The minimal change in the $\delta^{18}\text{O}_w$ data (Fig. 4 c, g) suggests that the amount of rainfall did not significantly change during Heinrich stadials. The increase in $\delta^{18}\text{O}_c$ values across the Heinrich stadials, which is much greater than that found in the $\delta^{18}\text{O}_w$ results, most likely arises from lower temperatures, rather than a significant change in the $\delta^{18}\text{O}_w$ value or precipitation amount. The $\delta^{18}\text{O}_w$ values, when combined with $\delta^{18}\text{O}_c$ values, allows for the calculation of the temperature at the time of mineral formation. Through the application of the cave-specific water-calcite oxygen

isotope fractionation relationship from Tremaine et al. (2011), temperature has been calculated using the measured $\delta^{18}\text{O}_w$ and $\delta^{18}\text{O}_c$ values. The equation is as follows:

$$1000\ln\alpha = 16.1(10^3 T^{-1}) - 24.6 \quad (1)$$

With α representing the fractionation factor between two substances (in this case the measured water and calcite oxygen isotopes) and T in Kelvin (Tremaine et al., 2011). Equation 1 was developed from a cave monitoring study in Florida, and was found to be in good agreement with caves throughout the world (Tremaine et al., 2011).

While caves generally maintain a constant temperature throughout the year (representative of mean annual temperature), over longer time periods, speleothems can record decadal to centennial scale changes in temperature. Utilizing equation 1, the average temperature (not including the samples forming over Heinrich stadials, as defined in section 3.4) was 17.7 ± 2.7 °C. This average temperature is cooler than the average modern annual temperature for the Bahamas (average annual temperature = 22 to 28 °C (Baldini et al., 2007)) and typical temperatures within modern dry caves in the area (average annual cave temperature = 23.2°C (Arienzo et al., 2013)). Heinrich stadials 1, 2, and 3 are represented by cooler temperatures, with the average temperature across Heinrich stadials 1, 2 and 3 of 13.4 ± 2.7 °C, representing an average temperature decrease of 4.3 ± 2.7 °C and Heinrich stadial 1 being the coolest event (Figure 4d).

4.2 Climate interpretation of the carbon isotope record

Both the $\delta^{13}\text{C}_c$ and $\delta^{18}\text{O}_c$ results consists of increased values around 15 ± 0.2 , 24.5 ± 0.2 , and 31.6 ± 1 kyr BP suggesting that similar environmental factors influenced the $\delta^{13}\text{C}_c$ and $\delta^{18}\text{O}_c$

composition of the stalagmite during Heinrich stadials. As the $\delta^{18}\text{O}_c$ value across Heinrich stadial events is primarily driven by changes in the cave temperature, we suggest that the increases in the $\delta^{13}\text{C}_c$ values are indirectly driven by cooling during these Heinrich stadials. Assuming changes in cave ventilation did not occur during sample deposition, the $\delta^{13}\text{C}_c$ ratio of the speleothem is most likely driven by changes in the soil zone or the vegetation above the cave. The increased $\delta^{13}\text{C}_c$ values may be a result of a decline in soil microbial activity, which tends to increase the $\delta^{13}\text{C}_{\text{DIC}}$ ratio (Dissolved Inorganic Carbon, DIC) of the drip water during periods of cooler temperatures. Support for this notion comes from similar results found in a cave in France, where lower $\delta^{13}\text{C}_c$ values during warm periods were driven by drip water having a greater biogenic CO_2 component (Genty et al., 2003). Nevertheless, it is not possible to rule out the potential influence of changes in the amount of precipitation on the $\delta^{13}\text{C}_c$ record. An increase in the $\delta^{13}\text{C}_c$ ratio of the carbonate can be driven by aridity, which leads to a reduction in soil microbial activity, or a greater proportion of C_4 versus C_3 plants. While this is a valid interpretation of the carbon isotope results, the oxygen isotope results from fluid inclusion analysis do not suggest a reduction in the amount of rainfall as a driver of the $\delta^{18}\text{O}_c$ record.

4.3 Climate variability in the context of the Atlantic Basin

Proxy records from throughout the Atlantic not only support significant climate shifts associated with Heinrich stadials, but they are also in good agreement with the timing of Heinrich stadials 1-3 (Figure 6). These reconstructions support the global scale response to Heinrich stadials. The difference in climate expression between Northern Hemisphere records (*e.g.*, Hulu Cave, Cariaco Basin, and the Bahamas) compared to Southern Hemisphere records (*e.g.*, Pacupahuain Cave) demonstrates the anti-phased response across Heinrich stadials (Broecker, 1998; Kanner et al., 2012; Peterson et al., 2000; Wang et al., 2001; Wang et al., 2004)

(Figure 6). This anti-phased relationship between the hemispheres is in part due to a shift in climate across Heinrich stadials and is well documented in the ice core records with the Greenland ice core records supporting cooling and the ice cores from Antarctica supporting warming during Heinrich stadials (Broecker, 1998; Wolff et al., 2010).

The Bahamian speleothem record, along with other proxies from the subtropical/tropical western Atlantic, support overall drying and/or similar cooling associated with Heinrich stadials (Escobar et al., 2012; Lachniet et al., 2013; Peterson et al., 2000) (Figure 6). This coherence in climate expression among the North Atlantic tropical records across Heinrich stadials can be explained by a regional shift in climate. An alkenone-derived SST record from the Bermuda Rise across Heinrich stadials 4 and 5 shows a ~3-5 °C decrease in SSTs (Sachs and Lehman, 1999) suggesting SSTs also may have declined during earlier Heinrich stadials. Further south, a lake record from Guatemala shows an increased aridity and a 6-10 °C cooling associated with Heinrich stadial 1 (Escobar et al., 2012; Hodell et al., 2012). The temperatures observed during Heinrich stadial 1 are lower than the modern mean annual temperature for Guatemala (mean annual temperature = 26 °C) (Hodell et al., 2012) and the increased aridity and cooling in Guatemala is thought to be driven by a southerly shifted ITCZ and decreased tropical Atlantic SSTs (Escobar et al., 2012; Hodell et al., 2012). The amount of cooling observed in the Guatemala and the Bermuda records (between 3-10 °C) across Heinrich stadials is similar to the amount of cooling observed in the Bahamas record presented here (~4 °C). Modeling results also support decreased temperatures for the tropical Northern Hemisphere coincident with Heinrich stadials, however, the modeled temperature changes are much less (Murphy et al., 2014; Zhang and Delworth, 2005).

As suggested above, while the geochemical data in the speleothem record from the Bahamas supports a decrease in temperature, the carbon isotope results do not rule out the possibility of enhanced aridity across these events, which potentially may drive the variability in the carbon isotopic record. A speleothem record from southwestern Mexico demonstrates a reduction in the North American Monsoon associated with Heinrich stadial 1 driven by the southerly shifted ITCZ and reduced AMOC (Lachniet et al., 2013). A southerly shift in the ITCZ would be accompanied by an expanded Bermuda high, which could have led to increased aridity in the Bahamas across Heinrich stadials (Hodell et al., 2000).

In contrast to the data presented here, a record based on pollen from Lake Tulane, Florida in the southeastern U.S. suggests warmer and wetter Heinrich stadials (Grimm et al., 2006). These authors attributed the warming and increased rainfall to a slowdown of the AMOC leading to a reduction in the transport of heat from the Gulf of Mexico and thereby forcing the observed warmer and wetter climate in Florida (Grimm et al., 2006). This climate response may not be observed in the Bahamas as a result of the location of Abaco Island, on the eastern edge of the Bahamas platform, and therefore this study site may not have been influenced by changes in the Gulf of Mexico. Furthermore, the postulated warming of the Gulf of Mexico from the Florida record is not consistent with the simulated climate response to an AMOC shut down rather, most climate models suggest a cooling of the entire North Atlantic, including the Gulf of Mexico (Clement and Peterson (2008) and references therein).

In many paleoclimate proxy reconstructions, Heinrich stadial 1 is a larger magnitude event compared to stadials 2 and 3 (Stager et al., 2011). In the record presented here, Heinrich stadial 1 is characterized by the greatest change in $\delta^{18}\text{O}_c$ and temperature (observed $\sim 5.9^\circ\text{C}$ temperature decrease) (Figure 6). Heinrich stadial 2 exhibits minimal geochemical changes

which may be due to a minimal change in the AMOC during this event (Lynch-Stieglitz et al., 2014; McManus et al., 2004). While Heinrich stadial 1 has been shown to be the strongest event in many paleoclimate reconstructions, it is also a period punctuated with climate changes. For example, in Guatemala a lake record indicates a cold and dry period beginning around 18 kyr BP, interrupted by a wetter and warmer period ~17 kyr BP, and followed by a return to drier and cooler temperatures at approximately 16.1 kyr BP (Escobar et al., 2012; Hodell et al., 2012). The authors suggest the two cold and dry periods observed in Guatemala are driven by two separate North Atlantic IRD events (Escobar et al., 2012). Similar to the observations in Guatemala, a multi-proxy study of a core from the Cariaco Basin supports the drier climate associated with Heinrich stadial 1 is interrupted by a shift to wetter climate conditions (Escobar et al., 2012; Yurco, 2010). In this speleothem record from the Bahamas, geochemical results suggest a temperature increase and/or precipitation change from ~17.5 to ~17.1 kyr BP followed by a progressively cooler Heinrich stadial 1 period (Table 2, Figure 4 g & h). These results suggest that Heinrich stadial 1 in the tropical North Atlantic may exhibit centennial scale climate oscillations associated with high latitude IRD events.

Additional $\delta^{18}\text{O}_c$ and $\delta^{13}\text{C}_c$ variations, not associated with Heinrich stadials, are observed at 19.1 ± 0.2 and 22.6 ± 0.2 kyr before present. The magnitude of the isotopic shifts at 19 and 22.6 kyr BP are less than the isotopic shifts observed during Heinrich stadial events (at 19 kyr BP, a 1.7‰ shift in the $\delta^{13}\text{C}_c$ value and at 22.6 kyr BP a 2 ‰ shift for $\delta^{13}\text{C}_c$). Both of these time periods are also associated with a minor increase in temperature and $\delta^{18}\text{O}_w$ value from the fluid inclusion results. Similarly, Bond et al. (1997) observed increased IRD deposits at similar time periods potentially suggesting these isotopic variations are associated with high latitude events.

The resumption of AMOC at the end of Heinrich stadial 1 is believed to have resulted in an increase in SSTs in the North Atlantic (McManus et al., 2004), leading to the northward shift of the ITCZ and the observed shift to wetter and warmer conditions in the northern tropics and the start of the Bølling-Allerød period (Peterson et al., 2000). This is also supported by the increase in temperatures after Heinrich stadial 1 from the fluid inclusion results (Figure 4d).

While other climate records for the same time period from nearby localities such as the Cariaco Basin (Peterson et al., 2000) and stalagmites from the Peruvian Andes (Kanner et al., 2012) have recorded millennial scale variations (*e.g.*, D/O events) there is no evidence of D/O variability in the isotopic record of sample AB-DC-09. A similar observation was made from speleothems collected from Borneo in which the geochemistry of the speleothems recorded Heinrich stadials, but not the D/O events (Carolin et al., 2013). The lack of evidence for D/O events suggests that these are different to Heinrich stadials either in the forcing mechanism, the global propagation, the geographic expression, or how these events are recorded in the speleothems, and also supports the significant impact of Heinrich stadials on tropical climate (Carolin et al., 2013; Wang et al., 2004).

5. Conclusion

The stalagmite record reported in this paper reveals three distinctive climate changes during the last glacial period, each coincident with a Heinrich stadial. The $\delta^{18}\text{O}_c$ and $\delta^{13}\text{C}_c$ data support significant shifts within the Heinrich stadials, while minimal changes occur in the $\delta^{18}\text{O}$ of the water contained in fluid inclusions. The $\delta^{18}\text{O}$ of the carbonate and the fluid inclusion results demonstrate that Heinrich stadials 1- 3 are characterized by a temperature decrease, rather than an increase in aridity at this time. The observed temperature decreases are consistent with

paleoclimate records from the Western Atlantic and we propose that the average temperature decline of ~4 °C during Heinrich stadials is due to the meridional shift in the Northern Hemisphere climate that led to cooler subtropics. Finally, the results support the teleconnection between the subtropical Atlantic and cooling in the North Atlantic and the sensitivity of the Bahamas to changes in AMOC.

Acknowledgement

This research was supported with funds from NSF grant AGS-1103489 (PKS, AC, KB, and AP) and a Geological Society of America Graduate Student Research Grant to MMA. The authors would like to thank the Stable Isotope Laboratory members at the University of Miami for insightful discussions and assistance with sample analysis and G. MacKenzie for her editorial advice. The authors would also like to thank the reviewers for their comments, which greatly enhanced the paper.

References

- Arienzo, M.M., Swart, P.K., Vonhof, H.B., 2013. Measurement of $\delta^{18}\text{O}$ and $\delta^2\text{H}$ values of fluid inclusion water in speleothems using cavity ring-down spectroscopy compared with isotope ratio mass spectrometry. *Rapid Communications in Mass Spectrometry* 27, 2616-2624.
- Asmerom, Y., Polyak, V.J., Burns, S.J., 2010. Variable winter moisture in the southwestern United States linked to rapid glacial climate shifts. *Nature Geoscience* 3, 114-117.
- Baldini, L.M., Walker, S.E., Railsback, L.B., Baldini, J.U.L., Crowe, D.E., 2007. Isotopic ecology of the modern land snail *Cerion*, San Salvador, Bahamas: Preliminary advances toward establishing a low-latitude island paleoenvironmental proxy. *Palaios* 22, 174-187.
- Beck, J.W., Richards, D.A., Edwards, R.L., Silverman, B.W., Smart, P.L., Donahue, D.J., Herrera-Osterheld, S., Burr, G.S., Calsoyas, L., Jull, A.J., Biddulph, D., 2001. Extremely large variations of atmospheric ^{14}C concentration during the last glacial period. *Science* 292, 2453-2458.
- Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen, H., Hajdas, I., Bonani, G., 1997. A pervasive millennial-scale cycle in North Atlantic Holocene and glacial climates. *Science* 278, 1257-1266.
- Breitenbach, S., Rehfeld, K., Goswami, B., Baldini, J., Ridley, H., Kennett, D., Prufer, K., Aquino, V., Asmerom, Y., Polyak, V., Cheng, H., Kurths, J., Marwan, N., 2012. COConstructing Proxy Records from Age models (COPRA). *Climate of the Past* 8, 1765-1779.

511 Broecker, W.S., 1998. Paleoocean circulation during the last deglaciation: A bipolar seesaw?
512 *Paleoceanography* 13, 119-121.

513 Carolin, S.A., Cobb, K.M., Adkins, J.F., Clark, B., Conroy, J.L., Lejau, S., Malang, J., Tuen,
514 A.A., 2013. Varied response of western Pacific hydrology to climate forcings over the last
515 glacial period. *Science* 340, 1564-1566.

516 Chiang, J.C.H., Bitz, C.M., 2005. Influence of high latitude ice cover on the marine Intertropical
517 Convergence Zone. *Climate Dynamics* 25, 477-496.

518 Clement, A.C., Peterson, L.C., 2008. Mechanisms of abrupt climate change of the last glacial
519 period. *Reviews of Geophysics* 46.

520 Craig, H., 1961. Isotopic variations in meteoric waters. *Science* 133, 1702-1703.

521 Dansgaard, W., 1964. Stable isotopes in precipitation. *Tellus* 16, 436-468.

522 Dansgaard, W., Johnsen, S.J., Clausen, H.B., Dahl-Jensen, D., Gundestrup, N., Hammer, C.U.,
523 Oeschger, H., 1984. North Atlantic climatic oscillations revealed by deep Greenland ice cores,
524 *Climate Processes and Climate Sensitivity*. AGU, p. 10.

525 Dorale, J.A., Liu, Z.H., 2009. Limitations of Hendy Test criteria in judging the paleoclimatic
526 suitability of speleothems and the need for replication. *J Cave Karst Stud* 71, 73-80.

527 Edwards, R.L., Gallup, C.D., Cheng, H., 2003. U-series dating of marine and lacustrine
528 carbonates, in: Bourdon, B. (Ed.), *Reviews in Mineralogy and Geochemistry: Uranium Series*
529 *Geochemistry*. Mineralogical Society of America, Washington, DC.

530 Escobar, J., Hodell, D.A., Brenner, M., Curtis, J.H., Gilli, A., Mueller, A.D., Anselmetti, F.S.,
531 Ariztegui, D., Grzesik, D.A., Perez, L., Schwab, A., Guilderson, T.P., 2012. A ~43-ka record of
532 paleoenvironmental change in the Central American lowlands inferred from stable isotopes of
533 lacustrine ostracods. *Quaternary Science Reviews* 37, 92-104.

534 Fairchild, I.J., Smith, C.L., Baker, A., Fuller, L., Spötl, C., Mathey, D., McDermott, F., E.I.M.F.,
535 2006. Modification and preservation of environmental signals in speleothems. *Earth-Science*
536 *Reviews* 75, 105-153.

537 Fietzke, J., Liebetrau, V., Eisenhauer, A., Dullo, C., 2005. Determination of uranium isotope
538 ratios by multi-static MIC-ICP-MS: method and implementation for precise U- and Th-series
539 isotope measurements. *Journal of Analytical Atomic Spectrometry* 20, 395-401.

540 Genty, D., Blamart, D., Ouahdi, R., Gilmour, M., Baker, A., Jouzel, J., Van-Exter, S., 2003.
541 Precise dating of Dansgaard-Oeschger climate oscillations in western Europe from stalagmite
542 data. *Nature* 421, 833-837.

543 Grimm, E.C., Watts, W.A., Jacobson Jr, G.L., Hansen, B.C.S., Almquist, H.R., Dieffenbacher-
544 Krall, A.C., 2006. Evidence for warm wet Heinrich events in Florida. *Quaternary Science*
545 *Reviews* 25, 2197-2211.

546 Hagen, S., Keigwin, L.D., 2002. Sea-surface temperature variability and deep water
547 reorganisation in the subtropical North Atlantic during Isotope Stage 2-4. *Marine Geology* 189,
548 145-162.

549 Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the Northeast Atlantic-
550 Ocean during the past 130,000 years. *Quaternary Research* 29, 142-152.

551 Hendy, C.H., 1971. The isotopic geochemistry of speleothems—I. The calculation of the effects
552 of different modes of formation on the isotopic composition of speleothems and their
553 applicability as palaeoclimatic indicators. *Geochimica et Cosmochimica Acta* 35, 801-824.

554 Hodell, D.A., Brenner, M.K., Curtis, J.H., 2000. Climate change in the northern American
555 tropics and subtropics since the last ice age: Implications for environment and culture, in: Lentz,
556 D.L. (Ed.), *An Imperfect Balance : Landscape Transformations in the Pre-Columbian Americas*.
557 Columbia University Press, New York, p. 547 p.

558 Hodell, D.A., Turchyn, A.V., Wiseman, C.J., Escobar, J., Curtis, J.H., Brenner, M., Gilli, A.,
559 Mueller, A.D., Anselmetti, F., Ariztegui, D., Brown, E.T., 2012. Late Glacial temperature and
560 precipitation changes in the lowland Neotropics by tandem measurement of $\delta^{18}\text{O}$ in biogenic
561 carbonate and gypsum hydration water. *Geochimica et Cosmochimica Acta* 77, 352-368.

562 Hoffmann, D.L., Beck, J.W., Richards, D.A., Smart, P.L., Singarayer, J.S., Ketchmark, T.,
563 Hawkesworth, C.J., 2010. Towards radiocarbon calibration beyond 28ka using speleothems from
564 the Bahamas. *Earth and Planetary Science Letters* 289, 1-10.

565 Kanner, L.C., Burns, S.J., Cheng, H., Edwards, R.L., 2012. High-latitude forcing of the South
566 American summer monsoon during the Last Glacial. *Science* 335, 570-573.

567 Keigwin, L.D., Jones, G.A., 1994. Western North-Atlantic evidence for millennial-scale changes
568 in ocean circulation and climate. *Journal of Geophysical Research-Oceans* 99, 12397-12410.

569 Lachniet, M.S. 2009. Climatic and environmental controls on speleothem oxygen-isotope values.
570 *Quaternary Science Reviews* 28, 412-432.

571 Lachniet, M.S., Asmerom, Y., Bernal, J.P., Polyak, V.J., Vazquez-Selem, L., 2013. Orbital
572 pacing and ocean circulation-induced collapses of the Mesoamerican monsoon over the past
573 22,000 y. *Proceedings of the National Academy of Sciences of the United States of America* 110,
574 9255-9260.

575 Lynch-Stieglitz, J., Schmidt, M.W., Henry, L.G., Curry, W.B., Skinner, L.C., Mulitza, S., Zhang,
576 R., Chang, P., 2014. Muted change in Atlantic overturning circulation over some glacial-aged
577 Heinrich events. *Nature Geoscience* 7, 144–150.

578 McGee, D., Quade, J., Edwards, R.L., Broecker, W.S., Cheng, H., Reiners, P.W., Evenson, N.,
579 2012. Lacustrine cave carbonates: Novel archives of paleohydrologic change in the Bonneville
580 Basin (Utah, USA). *Earth and Planetary Science Letters* 351, 182-194.

581 McManus, J.F., Francois, R., Gherardi, J.M., Keigwin, L.D., Brown-Leger, S., 2004. Collapse
582 and rapid resumption of Atlantic meridional circulation linked to deglacial climate changes.
583 *Nature* 428, 834-837.

584 Mühlinghaus, C., Scholz, D., Mangini, A., 2009. Modelling fractionation of stable isotopes in
585 stalagmites. *Geochimica et Cosmochimica Acta* 73, 7275-7289.

586 Murphy, L.N., Clement, A.C., Albani, S., Mahowald, N.M., Swart, P.K., Arienzo, M.M., 2014.
587 Simulated changes in atmospheric dust in response to a Heinrich stadial. *Paleoceanography* 29,
588 30-43.

589 Peterson, L.C., Haug, G.H., Hughen, K.A., Rohl, U., 2000. Rapid changes in the hydrologic
590 cycle of the tropical Atlantic during the last glacial. *Science* 290, 1947-1951.

591 Pourmand, A., Tissot, F.L.H., Arienzo, M.M., Sharifi, A., 2014. Introducing a comprehensive
592 data reduction and uncertainty propagation algorithm for U-Th geochronometry with extraction
593 chromatography and isotope dilution MC-ICP-MS. *Geostandards and Geoanalytical Research*
594 38, 129-148.

595 Richards, D.A., Smart, P.L., Edwards, R.L., 1994. Maximum sea levels for the last glacial period
596 from U-series ages of submerged speleothems. *Nature* 367, 357-360.

597 Roberts, W., Valdes, P., Payne, A., 2014. Topography's crucial role in Heinrich Events.
598 *Proceedings of the National Academy of Sciences of the United States of America* 111, 16688-
599 16693.

600 Robinson, L.F., Belshaw, N.S., Henderson, G.M., 2004. U and Th concentrations and isotope
601 ratios in modern carbonates and waters from the Bahamas. *Geochimica et Cosmochimica Acta*
602 68, 1777-1789.

603 Sachs, J.P., Lehman, S.J., 1999. Subtropical North Atlantic temperatures 60,000 to 30,000 years
604 ago. *Science* 286, 756-759.

605 Stager, J.C., Ryves, D.B., Chase, B.M., Pausata, F.S., 2011. Catastrophic drought in the Afro-
606 Asian monsoon region during Heinrich event 1. *Science* 331, 1299-1302.

607 Tremaine, D.M., Froelich, P.N., Wang, Y., 2011. Speleothem calcite farmed in situ: Modern
608 calibration of $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ paleoclimate proxies in a continuously-monitored natural cave
609 system. *Geochimica et Cosmochimica Acta* 75, 4929-4950.

610 van Breukelen, M.R., Vonhof, H.B., Hellstrom, J.C., Wester, W.C.G., Kroon, D., 2008. Fossil
611 dripwater in stalagmites reveals Holocene temperature and rainfall variation in Amazonia. *Earth*
612 *and Planetary Science Letters* 275, 54-60.

- Walker, L.N., Mylroie, J.E., Walker, A.D., Mylroie, J.R., 2008. The caves of Abaco Island, Bahamas: Keys to geologic timelines. *Journal of Cave and Karst Studies* 70, 108-119.
- Wang, Y.J., Cheng, H., Edwards, R.L., An, Z.S., Wu, J.Y., Shen, C.C., Dorale, J.A., 2001. A high-resolution absolute-dated late Pleistocene monsoon record from Hulu Cave, China. *Science* 294, 2345-2348.
- Wang, X., Auler, A.S., Edwards, R.L., Cheng, H., Cristalli, P.S., Smart, P.L., Richards, D.A., Shen, C. C., 2004. Wet periods in northeastern Brazil over the past 210 kyr linked to distant climate anomalies. *Nature* 432, 740-743.
- Wolff, E.W., Chappellaz, J., Blunier, T., Rasmussen, S.O., Svensson, A., 2010. Millennial-scale variability during the last glacial: The ice core record. *Quaternary Science Reviews* 29, 2828-2838.
- Yurco, L.N., 2010. A multi-proxy investigation of the late glacial "Mystery Interval" (17.5-14.5 ka) in the Cariaco Basin, Venezuela, Marine Geology and Geophysics. University of Miami. MS Thesis.
- Zhang, R., Delworth, T.L., 2005. Simulated tropical response to a substantial weakening of the Atlantic thermohaline circulation. *Journal of Climate* 18, 1853-1860.

Figure captions

Figure 1: Sample AB-DC-09 was collected from Abaco Island in the Bahamas (N26°14, W77°10). Cave location is indicated by the star. Contours show 120 m bathymetry.

Figure 2: a) Photo of sample AB-DC-09 showing the sampling locations: dashed line represents low resolution analysis, black bars represent high resolution analysis, and horizontal dotted lines represent the location of Hendy tests. b) Photo of thin section after sampling for geochemical analysis of the upper most section of the speleothem. This demonstrates areas of both high and low resolution $\delta^{18}\text{O}_c$ & $\delta^{13}\text{C}_c$ analyses. Along the center axis, millimeter scale laminations of the speleothem calcite are found. Microscopic analysis displays no evidence of post depositional alteration or evidence of a hiatus in the center. However, along the outer most portion of the stalagmite, blocky calcite is observed with minimal layering, suggesting this area may be affected by post-depositional diagenesis. c) Hendy test results from 3 cm from the top showing variation in carbon and oxygen isotopes (black circles and open squares respectively) from the center of the growth axis outward along a growth band, showing little variation in both carbon and oxygen isotopes. Also shown is low correlation between carbon and oxygen isotopes. d) Hendy test results from 23 cm from the top showing carbon and oxygen isotopes (black circles and open squares respectively) from the center of the growth axis outward, with little variation and low correlation across a growth band.

647 Figure 3: Age model (black line) based on twenty U-Th ages. The U-Th ages (black dots) are
648 reported with 95% confidence interval which includes random and systematic sources of
649 uncertainty, and the uncertainties on decay constants (see Pourmand et al. (2014) for details).
650 The age model was developed utilizing the COPRA program from Breitenbach et al. (2012).
651 2000 Monte Carlo simulations were conducted, with the black line representing the mean value
652 and the dotted line representing the 95% confidence interval (Breitenbach et al., 2012).

653 Figure 4: a and b) Carbon isotope values of the carbonate (green) time series from AB-DC-09
654 plotted with $\delta^{18}\text{O}_c$ (blue) from the same sample, with 50 year interpolation in black. c) Dark blue
655 squares are fluid inclusion $\delta^{18}\text{O}_w$ water results. d) Average temperature from fluid inclusion
656 results is shown in the solid black line, calculated using the equation from Tremaine et al.
657 (2011). e-g) Same as a-d from 18,500 to 14,500 years BP. Yellow bars are Heinrich stadials (HS)
658 1, 2 and 3.

659 Figure 5: The $\delta^{18}\text{O}_w$ and $\delta^2\text{H}_w$ values from fluid inclusion water isotopes plotted with the GMWL
660 in black and $\pm 5\text{‰}$ $\delta^2\text{H}$ of the GMWL in grey. Samples shown in green represent fluid inclusion
661 samples analyzed across Heinrich stadials. Samples all fall near the GMWL suggesting that the
662 water isotopes have not been significantly fractionated due to evaporation.

663 Figure 6: Compilation of paleoclimate studies from throughout the Atlantic. a) $\delta^{18}\text{O}$ time series
664 from NGRIP ice core (black) data from Wolff et al. (2010) plotted with a 5 point running
665 average. b) In orange, $\delta^{18}\text{O}$ time series from Hulu Cave, China data from PD and MSD samples
666 from Wang et al. (2001) note the axis is reversed. c) Oxygen isotope time series from this study
667 (sample AB-DC-09) plotted with a 50 year interpolation. Note the axis is reversed. d) Cariaco
668 Basin reflectance (red) at 550 nm, axis reversed from Peterson et al. (2000) with a 5 point
669 smoothing. e) Green is $\delta^{18}\text{O}$ record from Pacupahuain Cave, Peru from Kanner et al. (2012).

Tables

670 Table 1: Stalagmite AB-DC-09 ^{230}Th dating results for samples dated in the Neptune Isotope Laboratory at the University of Miami
671 (see Pourmand et al. (2014) for detailed methodology), ages have been corrected for a $^{230}\text{Th}/^{232}\text{Th}$ activity ratio of 3.7 ± 0.6 to account
672 for detrital ^{230}Th . All uncertainties are calculated as 95% confidence interval (95% CI). Ratios in parentheses are activities.

Distance from Top μm	^{238}U (ppb)	(95% CI) $\pm$	($^{230}\text{Th}/^{238}\text{U}$)	(95% CI) $\pm$	($^{230}\text{Th}/^{232}\text{Th}$)	(95% CI) $\pm$	Uncorr. Age (yr)	(95% CI) $\pm$	($^{234}\text{U}/^{238}\text{U}$) initial (corrected)	(95% CI) $\pm$	Corr. Age (yr)	(95% CI) $\pm$
17000	201.838	0.124	0.1332	0.0007	57.6	0.94	15394	77	1.0131	0.00259	14468	166
35000	179.526	0.097	0.1360	0.0006	54.5	1.56	15882	71	1.0047	0.00267	14872	178
42500	209.472	0.103	0.1355	0.0009	66.8	2.30	15717	110	1.0112	0.00249	14902	174
60000	150.938	0.086	0.1552	0.0010	47.9	1.12	18572	123	0.9921	0.00278	17244	246
72500	188.150	0.113	0.1545	0.0014	102.5	1.90	18054	176	1.0145	0.00251	17453	198
89000	204.483	0.108	0.1607	0.0009	43.5	0.27	18956	109	1.0087	0.00265	17465	260
95000	203.633	0.110	0.1569	0.0007	81.8	2.81	18527	78	1.0053	0.00250	17753	146
103000	191.743	0.103	0.1615	0.0009	95.4	4.01	19043	111	1.0093	0.00248	18363	156
115000	245.881	0.129	0.1683	0.0008	144.4	8.89	19852	96	1.0129	0.00256	19385	126
124000	214.688	0.106	0.1732	0.0008	80.7	0.52	20586	96	1.0085	0.00274	19722	169
125000	212.781	0.109	0.1739	0.0009	86.9	3.56	20597	105	1.0117	0.00255	19795	168
133000	180.454	0.083	0.1781	0.0010	77.4	0.96	21486	120	0.9972	0.00245	20551	191
137000	183.408	0.109	0.1876	0.0022	128.6	1.69	22309	284	1.0159	0.00253	21727	299
145000	164.362	0.082	0.1923	0.0010	48.3	0.36	23646	124	0.9875	0.00258	22007	294
149000	179.101	0.104	0.2067	0.0009	47.0	0.85	25148	100	1.0051	0.00249	23366	299
156000	192.366	0.113	0.2210	0.0011	110.9	0.83	26649	126	1.0215	0.00258	25859	177
178000	202.090	0.127	0.2235	0.0011	139.6	0.93	27030	127	1.0201	0.00252	26394	162
199000	204.470	0.150	0.2427	0.0010	38.2	0.17	29794	122	1.0170	0.00271	27239	433
210000	229.838	0.113	0.2435	0.0010	44.5	0.86	30055	124	1.0125	0.00271	27849	378
222000	202.531	0.126	0.2879	0.0013	19.0	0.11	36204	172	1.0209	0.00261	30025	1036
Not used in age model:												
50000	128.045	0.084	0.1324	0.0009	53.9	0.92	15640	112	0.9918	0.00248	14633	194
55000	129.585	0.074	0.1281	0.0006	69.1	2.56	15145	62	0.9888	0.00251	14384	138
110000	211.462	0.127	0.1779	0.0009	79.5	0.55	20926	110	1.0206	0.00263	20038	177

673 Table 2: Fluid inclusion oxygen and hydrogen isotope values with an average error of 0.5
674 and 2.0 ‰ respectively. Temperature was calculated from Tremaine et al. (2011), the
675 average error of 2.7 °C was calculated by accounting for the errors on the fluid inclusion
676 analyses.

cm from top	Age	Average $\delta^{18}\text{O}_w$	Average $\delta^2\text{H}_w$	n	$\delta^{18}\text{O}_c$	Temperature ¹
	yr BP	‰ (VSMOW)	‰ (VSMOW)		‰ (VPDB)	(°C)
		± 0.5 ‰	± 2.0 ‰			± 2.7 °C
0.9	14125	-2.7	-9.8	2	-2.3	17.3
3.0	14770	-2.3	-2.4	2	-1.6	15.8
3.6	14877	-2.2	-4.8	2	-1.6	16.1
4.8	15459	-2.4	-8.7	3	-1.0	12.2
5.5	16648	-2.3	-7.8	1	-0.7	11.3
6.0	17125	-2.2	-11.1	3	-2.2	19.7
6.6	17290	-2.5	-12.0	1	-2.1	17.5
7.2	17384	-2.8	-5.8	2	-2.0	15.3
8.0	17458	-3.1	-13.3	2	-2.0	13.9
8.5	17517	-2.5	-7.1	2	-2.2	18.1
10.1	18223	-2.7	-5.7	2	-2.4	17.7
10.6	18634	-2.8	-11.4	2	-2.3	16.8
11.1	19111	-2.4	-5.7	3	-2.3	18.6
11.6	19416	-3.1	-9.4	2	-2.1	14.8
12.7	20060	-3.2	-11.8	1	-2.5	15.9
13.6	21470	-2.6	-10.2	2	-2.5	18.8
14.5	22044	-2.4	-8.7	2	-1.9	17.0
14.9	23191	-3.1	-14.2	1	-2.1	14.4
15.4	25259	-3.3	-12.1	1	-2.0	12.8
16.0	26011	-2.9	-7.9	3	-1.6	12.6
17.7	26369	-2.7	-8.7	2	-2.1	16.6
18.3	26550	-2.3	-8.2	1	-2.0	17.9
19.1	26846	-2.0	-7.5	2	-2.4	21.7
19.4	26972	-2.6	-6.7	2	-2.5	18.7
19.9	27176	-1.9	-9.0	2	-2.2	20.7
20.5	27483	-1.9	-7.8	2	-2.5	22.4
21.0	27877	-2.2	-10.3	3	-2.2	19.5
21.5	28549	-2.2	-7.8	2	-2.0	18.1
22.0	29639	-2.1	-2.4	2	-2.2	19.8
22.4	30672	-1.8	-10.2	2	-2.1	21.5
22.8	31733	-1.8	-8.3	2	-1.4	17.4

¹ Temperature calculated from Tremaine et al. (2011)

678 Figure 1: Sample AB-DC-09 was collected from Abaco Island in the Bahamas (N26°14,
679 W77°10). Cave location is indicated by the star. Contours show 120 m bathymetry.

Figure 2: a) Photo of sample AB-DC-09 showing the sampling locations: dashed line represents low resolution analysis, black bars represent high resolution analysis, and horizontal dotted lines represent the location of Hendy tests. b) Photo of thin section after sampling for geochemical analysis of the upper most section of the speleothem. This demonstrates areas of both high and low resolution $\delta^{18}\text{O}_c$ & $\delta^{13}\text{C}_c$ analyses. Along the center axis, millimeter scale laminations of the speleothem calcite are found. Microscopic analysis displays no evidence of post depositional alteration or evidence of a hiatus in the center. However, along the outer most portion of the stalagmite, blocky calcite is observed with minimal layering, suggesting this area may be affected by post-depositional diagenesis. c) Hendy test results from 3 cm from the top showing variation in carbon and oxygen isotopes (black circles and open squares respectively) from the center of the growth axis outward along a growth band, showing little variation in both carbon and oxygen isotopes. Also shown is low correlation between carbon and oxygen isotopes. d) Hendy test results from 23 cm from the top showing carbon and oxygen isotopes (black circles and open squares respectively) from the center of the growth axis outward, with little variation and low correlation across a growth band.

Figure 3: Age model (black line) based on twenty U-Th ages. The U-Th ages (black dots) are reported with 95% confidence interval which includes random and systematic sources of uncertainty, and the uncertainties on decay constants (see Pourmand et al. (2014) for details). The age model was developed utilizing the COPRA program from Breitenbach et al. (2012). 2000 Monte Carlo simulations were conducted, with the black line representing the mean value and the dotted line representing the 95% confidence interval (Breitenbach et al., 2012).

Figure 4: a and b) Carbon isotope values of the carbonate (green) time series from AB-DC-09 plotted with $\delta^{18}\text{O}_c$ (blue) from the same sample, with 50 year interpolation in black. c) Dark blue squares are fluid inclusion $\delta^{18}\text{O}_w$ water results. d) Average temperature from fluid inclusion results is shown in the solid black line, calculated using the equation from Tremaine et al. (2011). e-g) Same as a-d from 18,500 to 14,500 years BP. Yellow bars are Heinrich stadials (HS) 1, 2 and 3.

738 Figure 5: The $\delta^{18}\text{O}_w$ and $\delta^2\text{H}_w$ values from fluid inclusion water isotopes plotted with the
 739 GMWL in black and $\pm 5\text{‰}$ $\delta^2\text{H}$ of the GMWL in grey. Samples shown in green represent
 740 fluid inclusion samples analyzed across Heinrich stadials. Samples all fall near the
 741 GMWL suggesting that the water isotopes have not been significantly fractionated due to
 742 evaporation.

759 Figure 6: Compilation of paleoclimate studies from throughout the Atlantic. a) $\delta^{18}\text{O}$ time
760 series from NGRIP ice core (black) data from Wolff et al. (2010) plotted with a 5 point
761 running average. b) In orange, $\delta^{18}\text{O}$ time series from Hulu Cave, China data from PD and
762 MSD samples from Wang et al. (2001) note the axis is reversed. c) Oxygen isotope time
763 series from this study (sample AB-DC-09) plotted with a 50 year interpolation. Note the
764 axis is reversed. d) Cariaco Basin reflectance (red) at 550 nm, axis reversed from
765 Peterson et al. (2000) with a 5 point smoothing. e) Green is $\delta^{18}\text{O}$ record from
766 Pacupahuain Cave, Peru from Kanner et al. (2012).

767
768
769
770
771
772
773
774