Knickpoint Evolution on the Yarlung River: Evidence for late Cenozoic Uplift of the Southeastern Tibetan Plateau Margin

Jennifer L Schmidt^{1*}, Peter K Zeitler¹, Frank J Pazzaglia¹, Marissa M Tremblay^{2,3}, David L Shuster^{2,3},
Matthew Fox^{2,3}

¹Department of Earth and Environmental Sciences, Lehigh University, 1 W. Packer Ave., Bethlehem, PA
18015, USA

²Department of Earth and Planetary Science, University of California, Berkeley, 307 McCone Hall #4767,
Berkeley, CA 94720-4767, USA

³Berkeley Geochronology Center, 2455 Ridge Road, Berkeley, CA 94709, USA

*Corresponding author: jl711@lehigh.edu

Abstract

A salient geomorphic feature of the Yarlung River is its abundance of large knickpoints, which in many cases coincide with north-south trending rifts. Across one of these rifts, near the town of Jiacha, the Yarlung falls nearly 500 m from an elevation of ~3500 m over 80 river kilometers, making this the second largest knickpoint on the river. We propose that the Jiacha knickpoint represents a wave of incision migrating upstream through the drainage network in response to a downstream base level fall, not a disturbance in the channel to due rift tectonics.

Longitudinal profile slope-area and chi (χ) analysis of Yarlung River tributaries and those of several major rivers in southeastern Tibet indicate several knickpoints are present at ~3500 m elevation, all resulting from a single regional-scale base level fall. Retreat rates calculated from celerity modeling indicate that the Jiacha knickpoint was located at the upstream edge of the Namche Barwa massif at ~10 Ma, a history consistent with apatite $^4\text{He}/^3\text{He}$ thermochronometry data and thermokinematic modeling from that region. These data suggest the Yarlung River has flowed in its present course through this area since at least 10 Ma and imply that at least 500 m of incision occurred within this canyon over this time period. The spatial scale of these observations suggests that these knickpoints resulted from surface uplift of southeastern Tibet of 500 to 2500 m just prior to ~10 Ma. Additionally, our mapped knickpoint locations indicate that reorganization of the drainage network just east of the Namche Barwa massif occurred prior to this time.

Keywords: Yarlung River; landscape evolution; knickpoint celerity modeling; $^4\text{He}/^3\text{He}$ apatite thermochronology; thermokinematic modeling.

1. Introduction

The Yarlung River flows strike parallel to the Himalayan orogen for ~1,700 km in southern Tibet before it abruptly drops over an ~2 km knickpoint and changes trajectory by ~180° in a gorge only 200 km long in the eastern Himalayan syntaxis (Fig. 1). The river drains >200,000 km² of the Tibetan

plateau and delivers ~2000 m³/s of discharge through the syntaxis (Montgomery et al., 2004), making the Yarlung River one of the major drainage catchments in Tibet and the largest in the southern plateau through which eroded sediments are evacuated. The topographic evolution of the southern Tibetan landscape is thus largely controlled by the base level established on the Yarlung drainage.

The longitudinal profile of the Yarlung River contains several large knickpoints, river reaches with convex elevation profiles and anomalously steep slopes (Fig. 2A). In the eastern portion of the catchment, near the town of Jiacha, the Yarlung channel navigates a narrow bedrock gorge, falling ~500 m in elevation over 80 river km from an elevation of 3500 m to form the second largest knickpoint on the river. In a detachment-limited stream power model (Howard and Kerby, 1983; Howard et al., 1994; Whipple and Tucker, 1999), in which elevation is increased due to rock uplift rate (U) and lowered through erosion ($E = KA^m S^n$: where A is upstream drainage area; S is local along channel slope; K is bedrock erodibility), such deviations from a graded, concave-up profile in a long-lived river imply either spatial variations in rock uplift rate or transience in the system associated with major changes in base level and upstream migration of erosional waves. As with several other knickpoints on the Yarlung River, the Jiacha knickpoint locally coincides with a late Cenozoic N–S trending rift that accommodates E–W extension in southern Tibet (Kapp et al., 2008; Yin et al., 1999; Fig. 1). An obvious inference is that dynamic support and/or active rift-flank uplift is locally steepening the Yarlung channel, forming the knickpoint (Zhang, 1998). However, the lack of major offset on this rift (Yin et al. 2010) precludes pinning of the Jiacha knickpoint to this location for an extended period of time. An alternative hypothesis that we test here is that the Jiacha knickpoint formed as a result of a regional base level fall downstream of its present position, propagating through the drainage network and triggering a wave of incision that is presently carving through the eastern Jiacha rift flank. Knickpoints originating in this manner travel upstream with a celerity of KA^m (assuming $n = 1$; Crosby and Whipple, 2006; Rosenbloom and Anderson, 1994; Weissel and Seidl, 1998; Whipple and Tucker 1999; Whipple et al., 2000) and should be evident in rivers throughout the drainage network. Furthermore, solutions to the stream power model indicate that transient features with a constant travel time (τ) should be present at the same elevation (Weissel and Seidl, 1998;

Pritchard, 2009; Royden and Perron, 2013; Goren et al, 2014). As there is no signature of K within the landscape, it is possible to calculate a K -independent travel time with units of length (Perron and Royden, 2013), termed χ , where $\chi(x) = KA_0^m \tau(x)$. By this method, knickpoints are identified at a specific elevation and corresponding χ value. These represent idealized scenarios and in reality, U and K can be spatially and temporally variable due to complex tectonics or variations in lithology or climate, respectfully. Nevertheless, this simplified stream power model provides a framework to assess the origin of this knickpoint, its relationship to knickpoints of similar elevation in southeastern Tibet, and implications for late Cenozoic drainage reorganization and landscape evolution.

We test these ideas through a combination of longitudinal profile and knickpoint celerity modeling, $^4\text{He}/^3\text{He}$ apatite thermochronometry, and thermokinematic modeling to constrain the Cenozoic incision history of the Yarlung downstream of the Jiacha knickpoint. Our results provide new perspective on the timing of river capture east of the eastern Himalayan syntaxis. We highlight the integration of knickpoint mapping with $^4\text{He}/^3\text{He}$ apatite thermochronometry and thermokinematic modeling as a novel approach to interpreting paleo-drainage patterns. Finally, we speculate on the origin of the Jiacha knickpoint by examining river profiles throughout eastern Tibet.

2. Identifying Transient Features in the Yarlung Drainage Network

We analyzed longitudinal profiles of the Yarlung and its tributaries upstream of the Namche Barwa massif to document evidence for migration of the Jiacha knickpoint in the reach immediately downstream, where knickpoint preservation is most likely. We used a 30 m ASTER digital elevation model (DEM) of southern Tibet as our base data set downloaded from the USGS Global Data Explorer (gdex.cr.usgs.gov/gdex). Using ESRI ArcGIS software, we extracted longitudinal elevation and drainage area profiles of the Yarlung and several of its tributaries, smoothing profiles using a loess filter in order to decrease noise and highlight the general form of the profile.

As erosion rate in the stream power model is equal to $KA^m S^n$, assuming the river is in steady state, erosion rate is equal to uplift rate ($U = KA^m S^n$) and we can therefore gain insight into relative erosion rates

by analyzing the relationship between S and A . This is achieved by rearranging this expression and regressing linear segments of the streams in $\log S - \log A$ space to provide a channel steepness index ($k_s = [U/K]^{1/n}$) and concavity ($\theta = m/n$). This slope-area analysis objectively highlights knickpoints in the profile (Hack, 1957; Kirby and Whipple, 2001; Snyder et al., 2000). We identified knickpoints where there is a change in a linear array of values to negative concavity in $\log S - \log A$ space, i.e. where a particular reach has a convex profile (Fig. 2B).

37 notable knickpoints exist within the Yarlung and 18 of its tributaries in the reach between the Namche Barwa massif and Jiacha knickpoint (Fig. 3). One of these knickpoints coincides with a mapped lithologic boundary. Of the others, 13 occur at elevations of $3500 \pm \sim 150$ m, the approximate elevation of the Jiacha knickpoint. The lithology of the tributaries north of the Yarlung is more variable than that of the southern tributaries and those lacking knickpoints at ~ 3500 m are focused in the north. This additional lithologic variation likely stalled or enhanced headward migration of these northern knickpoints, affecting their elevation. Cross-valley profiles from above and below the tributary knickpoints do not significantly differ, indicating the common elevation of these features does not reflect a regional erosive process change from fluvially dominated to glacially dominated at ~ 3500 m elevation. The morphology of this landscape at or below this elevation is therefore not controlled by glacial erosion.

A strong correlation exists between drainage area upstream of these 13 knickpoints and their distance from tributary confluence with the Yarlung (Fig. 4A). Since the celerity of knickpoint retreat in the stream power model is proportional to the drainage area raised to the power of m , this drainage area dependence on the horizontal distance a knickpoint will travel predicts this relationship when knickpoints are the result of a common base level fall (Berlin and Anderson, 2007; Goren et al., 2014; Harkins et al., 2007; Niemann et al., 2001; Van der Beek and Bishop, 2003; Whipple and Tucker, 1999; Zaprowski et al., 2001). We can further investigate this relationship by looking for commonalities across entire tributary channels and not solely the knickpoints. To remove the dependence of drainage area and negate the need to calculate slopes, as those derived from DEMs are prone to noise, we calculate χ -elevation profiles for the Yarlung River and its tributaries (Perron and Royden, 2013). If the tributaries share a common base

level history they should overlap in χ -elevation space. Divergence of the profiles may be due to spatial variation in channel incision processes or non-uniform precipitation expressed in K (Perron and Royden 2013). For the Yarlung and its tributaries, steps corresponding to knickpoints at ~3500 m elevation exhibit significant overlap, suggesting that these knickpoints are the result of a single, downstream base level fall. The dissimilarity in the overall morphology of the χ -elevation profiles between the northern and southern tributaries may result from a difference in hillslope and fluvial erosion processes arising from aspect and/or rock type.

The profile in Figure 4B shown in green contains a knickpoint at ~3500 m, however has significantly different χ values for a given elevation, likely due to recent capture by that tributary as suggested by its anomalous morphology observable in the DEM (see Fig. 3 for tributary location). This indicates that a substantial change in the drainage area occurred after the knickpoint migrated into the area, which increased χ values along this tributary. It is important to note that the absolute χ values are insignificant here given the uncertainty in the magnitude of the total elevation change across the Jiacha knickpoint (see section 6.3). Placing time constraints on the age of these knickpoint and commonalities requires external data and is discussed in section 3.

The detection of downstream knickpoints within streams east of the Namche Barwa massif, provides possible tests for mechanisms for this base level fall. Longitudinal profiles from six tributaries in the Siang River, along with the Yigong and Parlun rivers and 11 of their tributaries extracted from a 90 m SRTM DEM (downloaded from www.viewfinderpanoramas.org) highlight fourteen knickpoints occurring at 3500 ± 200 m (Fig. 3). Three of these knickpoints are located in the Namche Barwa massif within currently glaciated valleys. Longitudinal profiles from these tributaries thus do not reflect the valley bottom profile, but rather a combination of the glacier surface at high elevations and the valley bottom at lower elevations. They are therefore not related to the Jiacha knickpoint. Major rivers analyzed draining the eastern plateau margin and the Himalaya including the Subansiri, Dibang, Lhuit, Irrawaddy, Salween, Mekong, Yangtze, and Yalong (Fig. S1) also have knickpoints at ~3500 m, with the exception of the Lhuit

(Fig. 3). We discuss the implications of common elevation knickpoints in drainage networks spanning such a large area in Sections 6.2 and 6.3.

3. $^4\text{He}/^3\text{He}$ Apatite Thermochronometry

The development of high-relief topography such as that observed within the Yarlung River valley will perturb shallow crustal isotherms and thus shape the thermal history of the valley (Mancktelow and Grasemann, 1997; Braun, 2002). If a wave of sufficiently deep incision migrated headward along the Yarlung River, bedrock downstream of Jiacha should record exhumation-related cooling associated with passage of the knickpoint. Apatite $^4\text{He}/^3\text{He}$ thermochronometry constrains cooling histories through the uppermost few km of the crust. Schildgen et al. (2010) and Shuster et al. (2011) have demonstrated the potential of this method in identifying incision propagating through high-relief valleys. Here, we apply the method to samples downstream of the Jiacha knickpoint.

Retention of radiogenic ^4He , produced by alpha decay of U, Th, and Sm begins below $\sim 85^\circ\text{C}$ and is complete by $\sim 30^\circ\text{C}$ in apatite, depending on the size, cooling path, and the U, Th, and Sm concentrations of a given apatite crystal (Flowers et al., 2009; Shuster et al., 2006). A bulk (U-Th)/He apatite age is calculated from the total abundances of ^{238}U , ^{232}Th , ^{147}Sm and radiogenic ^4He in an apatite, and provides a non-unique constraint on the timing and rate of cooling between these temperatures. In $^4\text{He}/^3\text{He}$ thermochronometry, stepwise degassing analysis of a sample containing a uniform distribution of proton-induced ^3He resolves the spatial distribution of ^4He within a crystal, which more restrictively constrains an individual sample's thermal history (Shuster et al. 2004); the ^4He distribution and the (U-Th)/He age can thus be used to exclude thermal histories that are permitted on the basis of the (U-Th)/He age alone.

Our topographic analysis predicts that background exhumation within the valley will be punctuated by a period of rapid incision associated with knickpoint migration. If the knickpoint propagated upstream, this transient pulse of incision will have migrated with time. Bedrock apatite (U-Th)/He ages should therefore correlate with downstream distance from the Jiacha knickpoint and $^4\text{He}/^3\text{He}$ modeled thermal histories will provide constraints on the cooling rate. We applied these methods to two sample

sets collected within the Yarlung River valley to (1) ascertain the relationship between bedrock age and distance along the river, and (2) use modeled thermal histories to determine if samples experienced a pulse of rapid cooling.

We collected two groups of bedrock samples along the Yarlung River valley bottom (Table S1). The first group was collected near the base of the Jiacha knickpoint, while the second group spans a 100 km transect beginning ~200 km downstream from the lip of the Jiacha knickpoint (Fig. 5). Sample collection at river level maximizes the potential for recording knickpoint migration-induced incision, as these samples would experience the greatest amount of exhumation from this process. For analysis, we selected individual euhedral apatite grains free of visible inclusions and with a spherical equivalent radius of at least 50 microns from each sample. Following methods described by Shuster et al. (2004), samples were subjected to proton irradiation to produce a uniform distribution of ^3He . The $^4\text{He}/^3\text{He}$ ratio was measured by mass spectrometry during sequential, stepwise degassing of individual crystals (Table S2). The evolution of this ratio with cumulative ^3He release reflects the spatial distribution of ^4He within each grain. In these cases, we calculate the (U-Th)/He age from the sum of the ^4He released in all steps, and the U, Th, and Sm molar abundances in the same grain, measured by isotope dilution. The finite difference model described by Schildgen et al. (2010) was used to model cooling histories that predict the (U-Th)/He age within one standard error. Histories that satisfy this criterion are scored based on their misfit with the observed $^4\text{He}/^3\text{He}$ ratio evolution of these samples (Schildgen et al., 2010). Modeled histories with mean squared error (normalized to mean ratio uncertainty) of <1.5 represent good fits to our observed ratios (see Schildgen et al., 2010).

Figure 5 shows observed apatite $^4\text{He}/^3\text{He}$ data, modeled cooling histories, and sample locations with respect to the Jiacha knickpoint (see Fig. S3 for overlaid thermal histories). Within the valley transect, (U-Th)/He ages are 8.8 ± 0.88 Ma at the farthest downstream location and young to 4.4 ± 0.44 Ma ~100 km upstream (Fig. 5 & 6), demonstrating that cooling of these samples occurred earlier downstream than upstream. These data are consistent with a wave of cooling associated with incision propagating upstream as predicted by our knickpoint model. Apatite (U-Th)/He ages from the base of the knickpoint cluster

around 5 Ma. Because of the proximity of these samples to the present-day Jiacha knickpoint, they would have been far above the He partial retention zone (PRZ) when it migrated through this location. We therefore interpret these apatite ages as having been set during pre-knickpoint background erosion.

Thermal histories consistent with the observed $^4\text{He}/^3\text{He}$ data for upstream samples ML01, ML06, and ML18 (Fig. 5, Table S2) do not record a major change in cooling rate but do permit rapid cooling, suggesting they were below or entering the PRZ when knickpoint migration occurred. Because these histories do not record the onset of rapid cooling, we can constrain the timing of initiation of rapid incision at their location only to at least 4 Ma. The samples farthest downstream from the Jiacha knickpoint (ML19, and ML21) display a range of permissible model histories that is broad enough that we cannot resolve if these samples experienced fast or slow cooling. However, rapid cooling related to knickpoint migration by at least 8 Ma is permissible. Collectively, our modeled cooling histories from the valley transect samples constrain a time window (4 – 8 Ma) over which the samples cooled that young with upstream distance, which is consistent with our model of knickpoint migration. Predicted knickpoint migration rates from these data are 20-25 km/Ma.

In general, $^4\text{He}/^3\text{He}$ data from samples from the base of the knickpoint have uncertainties on the individual heating steps that are too large to provide quantitative constraints. Two samples from this area (LB12-ZD08 and LB12-ZD07) have lower uncertainties; nonetheless we are unable to distinguish between fast and slow cooling histories using these data (Fig.S4). We therefore present mean ages for these knickpoint samples in Figure 5. Additionally, we conducted conventional single crystal apatite (U-Th)/He analyses, calculating the (U-Th)/He age from the total release of ^4He in a single step (Table S3). U, Th, and Sm molar abundances were measured from the same grain by isotope dilution. These ages agree with those determined by $^4\text{He}/^3\text{He}$ thermochronometry for each sample, supporting the trend of decreasing age with distance upstream.

4. Knickpoint Celerity Modeling

We applied the stream power based knickpoint migration model for the Jiacha knickpoint following the general methodology of Berlin and Anderson (2007) to predict the knickpoint celerity and therefore mean retreat rate of valley incision directly downstream of Jiacha.

We selected three, adjacent north-flowing tributaries directly downstream of the Jiacha that have prominent knickpoints at ~3500 m (Fig. 3). Knickpoint celerity modeling is best constrained by using a large number of tributaries that experience instantaneous and coeval base level fall. The Yarlung drainage network geometry makes it difficult satisfy this requirement. Furthermore, differences in rock type among tributaries influence K and differences in aspect influence watershed hydrology (m), as observed in the χ -elevation plot (Fig. 4). For these reasons, solutions to the celerity model that combined north-flowing and south-flowing tributaries resulted in very high residual values for the best-fitting K and m . Although not ideal, the three tributaries used to constrain the celerity model have little to no variation in rock type, are of similar size and hypsometry, and are all confluent with the Yarlung at nearly the same location.

We assumed that these three tributaries experienced a base level fall resulting from Jiacha knickpoint migration at the exact same time, which is reasonable given the proximity of the tributary mouths where they join the Yarlung and the rapid migration rate expected. We modeled m and K values assuming the three tributary knickpoints initiated at 4.4 Ma (t) based on the tributary locations and our interpretation of sample ML01, the closest sample affected by knickpoint-related cooling. The distance from a tributary mouth to the top of the 3500m elevation knickpoint (x) and the corresponding upstream drainage area for a tributary channel reach (A) are known.

Using the celerity equation, we calculate a travel time through each cell of the tributary and cumulatively step upstream through the profile until the total distance is equal to the knickpoint travel distance, x . The difference between the calculated travel times through that distance and the initiation time (4.4 Ma) is the resulting misfit. The two-parameter exhaustive search approach applied searches parameter space linearly for values of m between 0.1 and 1.0 and logarithmically for K between 10^{-4} and 10^{-12} , to identify the best combination of m and K that minimizes the misfit among the three tributaries

(Berlin and Anderson, 2007; Crosby and Whipple, 2006). The residual-minimized m and K for these tributaries are 0.42 and $9.33 \times 10^{-7} \text{ m}^{-0.16} \text{ yr}^{-1}$ respectively.

We applied these best-fit values of m and K using the celerity equation in a forward sense to the Yarlung main channel to predict the location of the Jiacha knickpoint as a function of time. The Jiacha knickpoint projects back in time to the base of the three modeled tributaries (Fig. 3) at 4.4 Ma which is consistent with the notion that the tributary knickpoints were formed by migration of the Jiacha knickpoint. Following the same approach, the Jiacha knickpoint projects just upstream of the Namche Barwa massif at ~10 Ma (Fig. 3) and migrated at an average rate of ~40 km/Ma.

5. Thermokinematic Modeling

Our bulk apatite ages, $^4\text{He}/^3\text{He}$ -based thermal histories, and celerity modeling suggest that a wave of incision migrated through the Yarlung River valley from the Namche Barwa massif starting at ~10 Ma. We use a forward 3D thermokinematic model to evaluate geologically plausible incision scenarios consistent with our observed apatite (U-Th)/He data (using a modified version of Pecube; Braun, 2003). For a prescribed model of topographic evolution, the model solves the heat transport equation through time incorporating variable kinematic parameters to predict time-temperature history of each node in a 3-dimensional grid.

Models of canyon incision require integrating both vertical background surface erosion and the development of topography through time. As an initial condition of the model, we create a simplified topography to simulate a flat surface cut by a 1.5 km deep valley with an 80 km long knickpoint (Fig. S5). With successive time steps, the knickpoint advances horizontally upstream to cut the valley to a total of 2 km deep. The background erosion rate, and its variance through time, is a prescribed parameter in our model, as are knickpoint migration rate and geothermal gradient. We explored the effects of broadly varying each parameter on the time-temperature history of the downstream canyon. Prescribed knickpoint retreat rates dictate the model run times and we varied these between 15-35 km/Ma. We assigned typical geothermal gradients of either 25°C/km or 30°C/km. High-elevation (U-Th)/He apatite ages from near the

Yarlung-Nyang confluence indicate background exhumation rates are on the order of 0.25 km/Ma (Zeitler et al. 2014). However, we explored an order of magnitude variation in background erosion rates possible in this region by assigning it to be either 0.01 km/Ma or 0.25-0.2 km/Ma.

For each model run, we extracted time-temperature paths for nodes downstream of the ending knickpoint location that correspond to our sample location distance from the Jiacha knickpoint. We then use HeFTy (Ketchum, 2005) to predict apatite (U-Th)/He ages for the t-T paths of each of the five sample nodes, assuming ^4He diffusion kinetics prescribed by the radiation damage accumulation and annealing model (RDAAM; Flowers et al., 2009). We then calculated the sum of squared residuals between the predicted and observed (U-Th)/He ages for each model run, and use its minimum to identify the best fit of the explored Pecube models (Table S4).

We find that a knickpoint retreat rate of 30 km/Ma succeeds in predicting the youngest (U-Th)/He ages and an increasing trend with downstream distance (Fig. 6), assuming a geothermal gradient of 25°C/km and a continuous background exhumation rate of 0.2 km/Ma. Models with much faster rates of retreat predict apatite (U-Th)/He ages far less than those observed and those with prescribed background erosion rates an order of magnitude slower predict (U-Th)/He ages that are far older.

6. Discussion

Our results indicate that the geomorphic evolution of the Yarlung River and therefore erosion in southern Tibet between the Namche Barwa Massif and the Jiacha knickpoint has been influenced by a downstream base level fall in the Late Miocene. Furthermore, our data provide insights into the evolution of the drainage network surrounding the eastern Himalayan Syntaxis. Finally, we examine major river profiles in eastern Tibet to explore proposed river capture models for this region and speculate on the origin of the Jiacha knickpoint.

6.1 Timing of Knickpoint Retreat in the Upper Yarlung Valley

In our knickpoint celerity model, the time at which the three modeled tributaries experienced a base level fall is assumed to be roughly 4.4 Ma (t). This is based on the closest apatite $^4\text{He}/^3\text{He}$ sample affected by knickpoint migration (sample ML01). In this model, the Jiacha knickpoint projects to the upstream edge of the Namche Barwa massif at ~ 10 Ma. Solutions to the knickpoint celerity model are linear, meaning that a doubling of the assumed initiation time for the tributary knickpoints to 8.8 Ma relocates the Jiacha knickpoint to this point at ~ 20 Ma. We find that scenarios where t is far less than 4.4 Ma predict past locations that are inconsistent with our $^4\text{He}/^3\text{He}$ apatite data. Apatite (U-Th)/He ages of 7-12 Ma downstream of our sample locations (Zeitler et al., 2014) are also consistent with propagation of the Jiacha knickpoint over ~ 10 Ma.

Knickpoint retreat rates derived from our interpretation of $^4\text{He}/^3\text{He}$ apatite spectra and (U-Th)/He ages are 20-25 km/Ma. Our celerity modeling results suggest rates of ~ 40 km/Ma, whereas the best-fitting Pecube result was modeled with a retreat rate of 30 km/Ma. Although there is a degree of dependence of the celerity modeling on the apatite data, given the uncertainty in these analyses, the rates calculated among the different methods do not vastly differ and highlight the consistency of our model predictions. The discrepancy between retreat rates predicted from the thermokinematic and celerity modeling are likely due to the simplified nature of this version of Pecube, which does not include a full landscape evolution model.

The bestfitting of the explored Pecube model predicts ages for the three samples farthest downstream that are slightly too young when compared to our observed data. This may result from these samples having been collected from valley-wall locations that do not accurately represent paleo-river level. Wang et al. (2014) demonstrated that the bedrock river valley is buried beneath ~ 500 m of sediment in this portion of the Yarlung. Samples collected from 500 m above river level would produce ages that were slightly too old for our given model assumptions.

Collectively, the results presented here demonstrate that the knickpoints at ~ 3500 m are present throughout the present-day Yarlung River drainage network and that a wave of incision migrated westward upstream from the western border of the Namche Barwa massif. This indicates that the base

level fall that formed the Jiacha knickpoint occurred downstream along the present course of the Yarlung River and that the Yarlung has flowed eastward along the Indus-Yarlung suture from Jiacha through the Eastern Syntaxis since at least ~10 Ma. This interpretation and the findings of Lang and Huntington (2014) contrast with the idea that the Yarlung River flowed westward through the Himalaya-traversing Subansiri River for any significant amount of time (Cina et al., 2009; Zhang et al., 2012). Notwithstanding, short-duration glacial or landslide damming events on the Yarlung River (see Korup and Montgomery, 2008; Montgomery et al., 2004) may have temporarily impeded eastward flow and retreat of the knickpoint and forced intermittent trans-Himalayan routing.

Apatite (U-Th)/He and fission track ages from farther downstream near the Nyang confluence decrease to 4-5 Ma and within the Yarlung gorge are <1 Ma (Seward and Burg, 2008; Zeitler et al., 2014). Young apatites outside of the massif are likely affected by lateral heat flow, which advects isotherms towards the surface (Zeitler et al., 2014); consequently any evidence of knickpoint migration directly adjacent to the massif has been removed. Recent discussion regarding tectonic-erosional feedbacks in this region suggests that incision of the Yarlung promotes rapid rock uplift in the massif that began at 8-10 Ma (Zeitler et al., 2014). As localized rapid exhumation progressed, the position of the large ~2 km knickpoint within the Namche Barwa massif was fixed, holding upstream base level to ~3000 m, which is consistent with a slowing of upstream erosion rates after 5 Ma (Finnegan et al., 2008; Zeitler et al., 2014). Our calculated retreat rates suggest the Jiacha knickpoint migrated out of the massif just prior to or coincident with the initiation of rapid rock uplift. It is interesting to note this nearly synchronous series of events and we speculate on a relationship between knickpoint migration and initiation of rapid rock uplift in the syntaxis in Section 6.3. Our knickpoint migration model is consistent with recent measurement of an ~500 m thick sediment package within the Yarlung River downstream of the Jiacha knickpoint (Wang et al. 2014) in that our apatite data and knickpoint celerity modeling require the river to occupy a deeply incised canyon that formed over the last ~10 Ma. Formation of the ~2 km knickpoint within the Yarlung gorge and backfilling of the Yarlung canyon however may have initiated anytime between 2.5 Ma (Wang et al. 2014) and ~10 Ma.

6.2 Implications for Drainage Evolution

Barbed tributaries and wind gaps along the Parlung River indicate that it once flowed to the southeast through the Lhuit and/or Irrawaddy Rivers, contrasting its present-day northwest flow direction (Fig. 3; Burchfiel et al., 2000; Clark et al., 2004). Two end-member models for the reversal of the Parlung suggest very different paleo-drainage patterns within the region and thus different sediment evacuation routes. Some authors have postulated the existence of an antecedent Yarlung-Siang-Brahmaputra River, flowing in its present configuration since at least the Early Miocene (Fig. 7a), with capture of a separate Yigong-Parlung-Lhuit River occurring by northeastward migration of the deforming Namche Barwa antiform (Hallet and Molnar, 2001; Lang and Huntington, 2014; Seward and Burg, 2008). Others suggest a paleo-Yarlung-Parlung-Lhuit connection with headward cutting of the Siang-Brahmaputra capturing the Parlung (Fig. 7b; Brookfield, 1998; Clark et al., 2004; Liang et al., 2008; Robinson et al., 2014).

Our results provide new perspective on the timing of these drainage integration models just east of the Namche Barwa area. Long profiles and slope-area analysis of the Siang, Parlung, and Yigong rivers and their tributaries highlight the presence of ~3500 m elevation knickpoints (Fig. 3) indicating that these rivers have experienced the same base level fall as the Yarlung River. Based on these observations, we constrain knickpoint migration pathways and the timing of previously proposed scenarios for river reorganization consistent with the locations of knickpoints at ~3500 m in the drainage network.

If integration of the Yarlung, Parlung, and Siang occurred after the Jiacha knickpoint reached the upper Yarlung valley, both the Siang-Brahmaputra and the Parlung-Lhuit would have experienced this base level fall independently as we find evidence for knickpoints at ~3500 m elevation in both drainages. In a model in which the Yarlung was connected to the Parlung-Lhuit prior to this integration, the Jiacha knickpoint would have traveled up the Lhuit, through the Parlung, and into the upper Yarlung valley. However, we find evidence of an ~3500 m elevation knickpoint in the Parlung longitudinal profile presently located upstream of tributaries also containing knickpoints at ~3500 m elevation, indicating the Parlung knickpoint propagated from the northwest. If it had migrated from the southeast, we would not

expect to find ~3500 m elevation knickpoints on tributaries that meet the Parlung northwest of the present day Parlung knickpoint (Fig. 3). This suggests that the Parlung has flowed in its present northwesterly course since at least ~10 Ma, which is inconsistent with a post-10 Ma integration scenario. In an antecedent Yarlung-Siang-Brahmaputra model, the Jiacha knickpoint would have originated on the Siang or Brahmaputra portion of the drainage. The ~3500 m elevation knickpoints on the tributaries of the Parlung and Yigong would therefore be the result of the same base level fall but would have originated on the Lhuit rather than the Siang or Brahmaputra. An early connection between the Yarlung and Siang-Brahmaputra is consistent with Gangdese detritus in Miocene foreland units (Cina et al., 2009; Lang and Huntington, 2014). However, the location of the Parlung knickpoint is again inconsistent with migration from the Lhuit.

We note that the Lhuit does not contain a major knickpoint at ~3500 m and acknowledge that after Jiacha knickpoint migration, localized uplift of the Parlung-Lhuit wind gap could have obscured evidence of it. This would suggest that the Parlung knickpoint did not result from the base level fall discussed in this paper, but rather represents a later knickpoint that was subsequently uplifted to ~3500 m. However, we are presently unaware of any evidence for such localized uplift in this region.

If integration of the Yarlung, Parlung, and Siang rivers occurred prior to ~10 Ma, the Jiacha knickpoint would have originated from within the Siang or Brahmaputra portion of the drainage (Fig. 7a, b). This is consistent with knickpoints at ~3500 m elevation present within tributaries of the Siang, close to its confluence with the Yarlung. Gangdese Batholith-derived detritus present within Early through Late Miocene units of the Himalayan foreland basin is also consistent with pre-Late Miocene connection of the Yarlung and Siang-Brahmaputra (Chirouze et al., 2013; Cina et al., 2009; Lang and Huntington, 2014). Moreover, evidence of fish species diversification at 19-24 Ma (Britz, 2009; Ruber et al., 2004) and an increase in sediment flux into the Bengal Basin in the Eocene to Early Miocene (Alam et al., 2003) indicate an early capture event as suggested by Robinson et al. (2014).

Because knickpoints at ~3500 m elevation exist on both the Parlung and Yigong rivers and evidence of uplift in the Parlung-Lhuit wind gap is lacking, these knickpoints likely migrated up the Parlung and Yigong

via the Siang. This requires that the Yarlung, Yigong, Parlung, and Siang were integrated at the time of headward migration through the Namche Barwa region. In this case, either pre-integration drainage configuration previously proposed is consistent with the mapped knickpoint locations, as long as integration occurred prior to ~10 Ma (Fig. 7).

6.3 Origin of 3500 m elevation Knickpoint

From our knickpoint mapping, we infer that a major base level fall occurred downstream of the Yarlung-Siang confluence that caused knickpoint formation and generated a wave of incision that propagated upstream as previously suggested by Zeitler et al. (2014). We find evidence of this base level fall in knickpoints in tributaries on the Siang downstream until the maximum elevations of its tributaries do not reach 3500 m. Below this point, the tributaries may have experienced a base level fall, but the stream has fully adjusted to the new base level, precluding preservation and observation of a knickpoint.

A base level fall could be the result of a drainage capture event, deformation or uplift, or an abrupt change in precipitation. However, knickpoints at ~3500 m elevation are evident on other major rivers on the eastern margin of the Tibetan plateau including the Irrawaddy, Salween, Mekong, Yangtze, and Yalong (Fig. S1). Harkins et al. (2007) also describe a knickpoint at ~3500 m elevation on the Yellow river and several of its tributaries. Given the elevation change at the Jiacha knickpoint is 500 m, and that we find knickpoints of its approximate elevation throughout the eastern Tibetan plateau, we speculate that these knickpoints resulted from uplift of at least 500 m of the entire eastern margin just prior to ~10 Ma. Clark et al. (2005) and Duvall et al. (2012) describe a similar wave of incision at 9-13 Ma constrained by apatite (U-Th)/He thermochronology in the three-rivers region and elsewhere on the eastern margin of Tibet, which they relate to eastward expansion of the plateau. Additional knickpoint celerity modeling and thermochronologic constraints on the rivers of eastern Tibet are required to address the scale and potential synchronicity of this base level change.

Localized rapid rock uplift within the Namche Barwa massif pinned an ~2 km knickpoint to this location (Zeitler et al. 2001, Koons et al. 2013). As our celerity modeling suggests, the Jiacha knickpoint

migrated out of this region at ~10Ma, concurrent with or just prior to development of rapid rock uplift (Zeitler et al. 2014). The Jiacha and Namche Barwa knickpoints may have originated as a single feature resulting from a base level fall and were segmented with pinning of the Namche Barwa knickpoint. A 500 m elevation change (i.e. the Jiacha knickpoint) continued to migrate upstream to its present position. If resulting from the uplift of the southeastern margin of Tibet just prior to ~10 Ma, the full magnitude of this base level fall may have been the combined elevation change of these two features. Total uplift of the margin could therefore be upwards of 2.5 km.

7. Conclusions

We present thermochronology, knickpoint celerity modeling, and thermokinematic modeling from southeastern Tibet that demonstrates the mechanism and timing of incision of the Yarlung River. A series of ~3500 m elevation knickpoints exist throughout the region and slope-area and χ analysis of several longitudinal profiles from the drainage basin confirms their genetic relationship, suggesting a wave of knickpoint migration incised the Yarlung canyon. $^4\text{He}/^3\text{He}$ apatite thermochronology data from transect along the Yarlung are consistent with a wave of incision propagating upstream towards the Jiacha knickpoint. Celerity modeling projects the Jiacha knickpoint to just upstream of the Namche Barwa massif at ~10 Ma. Thermokinematic modeling of a simplified Jiacha knickpoint incision scenario predicts apatite (U-Th)/He ages similar to our observed data.

The spatial arrangement of multiple knickpoints at ~3500 m elevation in this region indicates that previously-recognized reorganization of the major rivers in the drainage just east of the Namche Barwa massif occurred prior to ~10 Ma and that the Yarlung flowed predominantly eastward since at least this time. Our results suggest a major base level fall resulting from uplift of 500-2500 m of the southeastern Tibetan margin occurred just prior to ~10 Ma.

Acknowledgements

This work was supported by an NSF Continental Dynamics program grant (EAR-1111853) to P.K.Z. and D.L.S. and a Geological Society of America Graduate Student Research Grant to J.L.S. We thank FanyiMeng, Jin Yue, and Yue Chen for collecting and sharing samples and Zhidan Zhao, Mo Xuanxue, and Di-Cheng Zhu for help with field logistics.

References

- Alam, M., Alam, M. M., Curray, J. R., Chowdhury, M. L. R., Gani, M. R., 2003. An overview of the sedimentary geology of the Bengal Basin in relation to the regional tectonic framework and basin-fill history. *Sedimentary Geology*. 155, 179-208.
- Berlin, M. M. and Anderson, R. S., 2007. Modeling of knickpoint retreat on the Roan Plateau, western Colorado. *Journal of Geophysical Research*. 112, F03S06.
- Braun, J., 2003. Pecube; a new finite-element code to solve the 3D heat transport equation including the effects of a time-varying, finite amplitude surface topography. *Computers & Geosciences*. 29, 787-794.
- Braun, J., 2002. Quantifying the effect of Recent relief changes on age-elevation relationships. *Earth and Planetary Science Letters*. 200, 331-343.
- Britz, R., 2009. *Danionellapriapus*, a new species of miniature cyprinid fish from West Bengal, India (Teleostei: Cypriniformes: Cyprinidae). *Zootaxa*. 53-60.
- Brookfield, M., 2008. Evolution of the great river systems of southern Asia during the Cenozoic India-Asia collision: Rivers draining north from the Pamir syntaxis. *Geomorphology*. 100, 296-311.

- 458 Burchfiel, B. C., Clark, M., Wang, E., Chen, Z., Liu, Y., Pan, G., 2000. Tectonic framework of the
459 Namche Barwa region, Eastern Himalayan Syntaxis, SE Tibet. Abstracts with Programs -
460 Geological Society of America. 32, 33.
- 461 Chirouze, F., Huyghe, P., van der Beek, P., Chauvel, C., Chakraborty, T., Dupont-Nivet, G., Bernet, M.,
462 2013. Tectonics, exhumation, and drainage evolution of the eastern Himalaya since 13 Ma from
463 detrital geochemistry and thermochronology, Kameng River section, Arunachal
464 Pradesh. Geological Society of America Bulletin. 125, 523-538.
- 465 Cina, S., Yin, A., Grove, M., Dubey, C., Shukla, D., Lovera, O., Kelty, T., Gehrels, G., Foster, D., 2009.
466 Gangdese arc detritus within the eastern Himalayan Neogene foreland basin: Implications for the
467 Neogene evolution of the Yalu-Brahmaputra River system. Earth and Planetary Science Letters.
468 285, 150-162.
- 469 Clark, M. K., House, M. A., Royden, L. H., Whipple, K. X., Burchfiel, B. C., Zhang, X., Tang, W., 2005.
470 Late Cenozoic uplift of southeastern Tibet. Geology (Boulder). 33, 525-528.
- 471 Clark, M. K., Schoenbohm, L. M., Royden, L. H., Whipple, K. X., Burchfiel, B. C., Zhang, X., Tang, W.,
472 Wang, E., Chen, L., 2004. Surface uplift, tectonics, and erosion of eastern Tibet from large-scale
473 drainage patterns. Tectonics. 23, 20.
- 474 Crosby, B. T. and Whipple, K. X., 2006. Knickpoint initiation and distribution within fluvial networks;
475 236 waterfalls in the Waipaoa River, North Island, New Zealand. Geomorphology. 82, 16-38.
- 476 Duvall, A. R., Clark, M. K., Avdeev, B., Farley, K. A., Chen, Z., 2012. Widespread late Cenozoic
477 increase in erosion rates across the interior of eastern Tibet constrained by detrital low
478 temperature thermochronometry. Tectonics. 31, Citation TC3014.

- 479 Finnegan, N. J., Hallet, B., Montgomery, D. R., Zeitler, P. K., Stone, J. O., Anders, A. M., Liu Yuping,
480 2008. Coupling of rock uplift and river incision in the Namche Barwa-Gyala Peri Massif, Tibet.
481 Geological Society of America Bulletin.120, 142-155.
- 482 Flowers, R. M., Ketcham, R. A., Shuster, D. L., Farley, K. A., 2009. Apatite (U-Th)/He
483 thermochronometry using a radiation damage accumulation and annealing
484 model.GeochimicaEtCosmochimicaActa. 73, 2347-2365.
- 485 Goren, L., Fox, M., Willett, S. D., 2014.Tectonics from fluvial topography using formal linear inversion;
486 theory and applications to the Inyo Mountains, California. Journal of Geophysical Research:
487 Earth Surface. 119, 1651-1681.
- 488 Hack, J. T., 1957. Studies of longitudinal stream profiles in Virginia and Maryland. U.S.Geological
489 Survey Professional Paper. 45-97.
- 490 Hallet, B. and Molnar, P., 2001. Distorted drainage basins as markers of crustal strain east of the
491 Himalaya. Journal of Geophysical Research.106, 13-13,709.
- 492 Harkins, N., Kirby, E., Heimsath, A., Robinson, R., Reiser, U., 2007. Transient fluvial incision in the
493 headwaters of the Yellow River, northeastern Tibet, China.Journal of Geophysical Research.112,
494 F03S04.
- 495 Howard, A. D., 1994. A detachment-limited model of drainage basin evolution.Water Resources
496 Research.30, 2261-2285.
- 497 Howard, A. D. and Kerby, G., 1983. Channel changes in badlands. Geological Society of America
498 Bulletin.94, 739-752.

- 499 Kapp, P., Taylor, M., Stockli, D., Ding, L., 2008. Development of active low-angle normal fault systems
500 during orogenic collapse; insight from Tibet. *Geology (Boulder)*. 36, 7-10.
- 501 Ketcham, R. A., 2005. Forward and inverse modeling of low-temperature thermochronometry
502 data. *Reviews in Mineralogy and Geochemistry*. 58, 275-314.
- 503 Kirby, E. and Whipple, K., 2001. Quantifying differential rock-uplift rates via stream profile analysis.
504 *Geology (Boulder)*. 29, 415-418.
- 505 Koons, P.O., Zeitler, P.K., and Hallet, B., 2013, Tectonic Aneurysms and Mountain Building, *in*
506 Schroder, J. and Owen, L.A. eds., *Treatise on Geomorphology*, Academic Press, San Diego, p.
507 318–349.
- 508 Korup, O. and Montgomery, D. R., 2008. Tibetan plateau river incision inhibited by glacial stabilization
509 of the Tsangpo gorge. *Nature*. 455, 786-789.
- 510 Lang, K. A. and Huntington, K. W., 2014. Antecedence of the Yarlung-Siang-Brahmaputra River, eastern
511 Himalaya. *Earth and Planetary Science Letters*. 397, 145-158.
- 512 Liang, Y., Chung, S., Liu, D., Xu, Y., Wu, F., Yang, J., Wang, Y., Lo, C., 2008. Detrital zircon evidence
513 from Burma for reorganization of the eastern Himalayan river system. *American Journal of*
514 *Science*. 308, 618-638.
- 515 Mancktelow, N. S. and Grasemann, B., 1997. Time-dependent effects of heat advection and topography
516 on cooling histories during erosion. *Tectonophysics*. 270, 167-195.
- 517 Montgomery, D. R., Hallet, B., Liu, Y., Finnegan, N., Anders, A., Gillespie, A., Greenberg, H. M., 2004.
518 Evidence for Holocene megafloods down the Tsangpo River gorge, southeastern Tibet.
519 *Quaternary Research*. 62, 201-207.

- 520 Niemann, J. D., Gasparini, N. M., Tucker, G. E., Bras, R. L., 2001. A quantitative evaluation of Playfair's
521 law and its use in testing long-term stream erosion models. *Earth Surface Processes and*
522 *Landforms*. 26, 1317-1332.
- 523 Perron, J. T. and Royden, L., 2013. An integral approach to bedrock river profile analysis. *Earth Surface*
524 *Processes and Landforms*. 38, 570-576.
- 525 Pritchard, D., Roberts, G. G., White, N. J., Richardson, C. N., 2009. Uplift histories from river profiles.
526 *Geophysical Research Letters*. 36.
- 527 Robinson, R. A. J., Brezina, C. A., Parrish, R. R., Horstwood, M. S. A., Oo, N. W., Bird, M. I., Thein, M.,
528 Walters, A. S., Oliver, G. J. H., Zaw, K., 2014. Large rivers and orogens; the evolution of the
529 Yarlung Tsangpo-Irrawaddy system and the eastern Himalayan syntaxis. *Gondwana Research*. 26,
530 112-121.
- 531 Rosenbloom, N. A. and Anderson, R. S., 1994. Hillslope and channel evolution in a marine terraced
532 landscape, Santa Cruz, California. *Journal of Geophysical Research*. 99, 13-14,029.
- 533 Royden, L. and Taylor Perron, J., 2013. Solutions of the stream power equation and application to the
534 evolution of river longitudinal profiles. *Journal of Geophysical Research: Earth Surface*. 118,
535 497-518.
- 536 Rüber, L., Britz, R., Kullander, S. O., Zardoya, R., 2004. Evolutionary and biogeographic patterns of the
537 Badidae (Teleostei: Perciformes) inferred from mitochondrial and nuclear DNA sequence data.
538 *Molecular Phylogenetics and Evolution*. 32, 1010-1022.
- 539 Schildgen, T. F., Balco, G., Shuster, D. L., 2010. Canyon incision and knickpoint propagation recorded
540 by apatite (super 4) He/ (super 3) He thermochronometry. *Earth and Planetary Science Letters*.
541 293, 377-387.

- 542 Seward, D. and Burg, J. P., 2008. Growth of the Namche Barwa Syntaxis and associated evolution of the
543 Tsangpo Gorge; constraints from structural and thermochronological data. *Tectonophysics*. 451,
544 282-289.
- 545 Shuster, D. L., Cuffey, K. M., Sanders, J. W., Balco, G., 2011. Thermochronometry reveals headward
546 propagation of erosion in an alpine landscape. *Science*. 332, 84-88.
- 547 Shuster, D. L., Flowers, R. M., Farley, K. A., 2006. The influence of natural radiation damage on helium
548 diffusion kinetics in apatite. *Earth and Planetary Science Letters*. 249, 148-161.
- 549 Shuster, D. L., Farley, K. A., Sistierson, J. M., Burnett, D. S., 2004. Quantifying the diffusion kinetics and
550 spatial distributions of radiogenic (super 4) He in minerals containing proton-induced (super 3)
551 He. *Earth and Planetary Science Letters*. 217, 19-32.
- 552 Snyder, N. P., Whipple, K. X., Tucker, G. E., Merritts, D. J., 2000. Landscape response to tectonic
553 forcing; digital elevation model analysis of stream profiles in the Mendocino triple junction
554 region, Northern California. *Geological Society of America Bulletin*. 112, 1250-1263.
- 555 van der Beek, P. and Bishop, P., 2003. Cenozoic river profile development in the upper Lachlan
556 catchment (SE Australia) as a test of quantitative fluvial incision models. *Journal of Geophysical*
557 *Research*. 108, .
- 558 Wang, P., Scherler, D., Liu-Zeng, J., Mey, J., Avouac, J., Zhang, Y., Shi, D., 2014. Tectonic control of
559 Yarlung Tsangpo Gorge revealed by a buried canyon in southern Tibet. *Science*. 346, 978-981.
- 560 Weissel, J. K. and Seidl, M. A., 1998. Inland propagation of erosional escarpments and river profile
561 evolution across the Southeast Australian passive continental margin. *Geophysical Monograph*.
562 107, 189-206.

- 563 Whipple, K. X., Hancock, G. S., Anderson, R. S., 2000. River incision into bedrock; mechanics and
564 relative efficacy of plucking, abrasion, and cavitation. *Geological Society of America*
565 *Bulletin*. 112, 490-503.
- 566 Whipple, K. X. and Tucker, G. E., 1999. Dynamics of the stream-power river incision model;
567 implications for height limits of mountain ranges, landscape response timescales, and research
568 needs. *Journal of Geophysical Research*. 104, 17-17,674.
- 569 Yin, A., Kapp, P. A., Murphy, M. A., Manning, C. E., Harrison, T. M., Grove, M., Ding, L., Deng, X.,
570 Wu, C., 1999. Significant late Neogene east-west extension in northern Tibet. *Geology*
571 *(Boulder)*. 27, 787-790.
- 572 Yin, A., Dubey, C.S., Kelty, T.K., Webb, A.A.G., Harrison, T.M., Chou, C.Y. and Clrier, J.,
573 2010. Geologic correlation of the Himalayan orogen and Indian craton: Part 2. Structural geology,
574 geochronology, and tectonic evolution of the Eastern Himalaya. *Geological Society of America*
575 *Bulletin*. 122, 3/4, 360–395.
- 576 Zaprowski, B. J., Evenson, E. B., Pazzaglia, F. J., Epstein, J. B., 2001. Knickzone propagation in the
577 Black Hills and northern High Plains; a different perspective on the late Cenozoic exhumation of
578 the Laramide Rocky Mountains. *Geology (Boulder)*. 29, 547-550.
- 579 Zeitler, P. K., Meltzer, A. S., Koons, P. O., Craw, D., Hallet, B., Chamberlain, C. P., Kidd, W. S. F., Park,
580 S. K., Seeber, L., Bishop, M., Shroder, J., 2001. Erosion, Himalayan geodynamics, and the
581 geomorphology of metamorphism. *GSA Today*. 11, 4-9.
- 582 Zeitler, P. K., Meltzer, A. S., Brown, L., Kidd, W. S. F., Lim, C., Enkelmann, E., 2014. Tectonics and
583 topographic evolution of Namche Barwa and the easternmost Lhasa Block, Tibet. *Special Paper -*
584 *Geological Society of America*. 507, 23-58.

Zhang, D. D., 1998. Geomorphological problems of the middle reaches of the Tsangpo River, Tibet. *Earth Surface Processes and Landforms*. 23, 889-903.

Zhang, J., Yin, A., Liu, W. C., Wu, F. Y., Lin, D., Grove, M., 2012. Coupled U-Pb dating and Hf isotopic analysis of detrital zircon of modern river sand from the Yalu River (Yarlung Tsangpo) drainage system in southern Tibet; constraints on the transport processes and evolution of Himalayan rivers. *Geological Society of America Bulletin*. 124, 1449-1473.

Figure Captions

Figure 1. Overview map of the Yarlung River drainage basin (location highlighted in inset). The drainage area includes a significant portion of southern Tibet. Major rifts (from Kapp and Guynn 2004) locally coincide with knickpoints along the river. The Nari Yun Chu Rift coincides with the Yarlung River at the Jiacha knickpoint (highlighted in red). Dashed ellipse illustrates approximate area of rapid rock uplift in the Namche Barwa massif (adapted from Stewart et al. 2008).

Figure 2. A) Elevation, drainage area, and SL index plotted against river distance for the Yarlung River. B) Slope-area plot for Yarlung River. Filled circles denote knickpoint reaches. Both A and B highlight the magnitude of the Jiacha (in red) and other knickpoints on the Yarlung River.

Figure 3. Analyzed tributaries (black, red, and green) and major rivers (blue) of southeastern Tibet (location highlighted in Figure 1). Circles indicate locations of identified knickpoints at ~3500 m elevation. Black circles indicate reaches with convex profiles that are not associated with fluvial knickpoints (see Section 2). Tributaries used in knickpoint celerity modeling are highlighted in red. Green

tributary has undergone a recent capture event as discussed in Section 2 and Figure 4b. Green star indicates the projected Jiacha knickpoint location at ~10 Ma, based on our knickpoint celerity model.

Figure 4. A) Knickpoint distance plotted against tributary drainage area for Yarlung tributaries containing knickpoints at ~3500 m elevation. Each knickpoint is plotted as filled circle. Note: Nyang river knickpoint located out of plot range. B) χ -elevation plot for tributaries on the north (top) and south (bottom) side of the Yarlung containing knickpoints at ~3500 m (red circles). We assume a concavity of 0.5. South tributary shaded in green (see Figure 3 for location) displays clear offset in χ , resulting from recent river capture as discussed in Section 2.

Figure 5. Sample ages and $^4\text{He}/^3\text{He}$ results. A) Map showing sample locations (yellow circles) and corresponding (U-Th)/He apatite age. The Jiacha knickpoint is highlighted in red. Samples located at the base of the Jiacha knickpoint are labeled with mean age of all aliquots (shown with 1 standard deviation uncertainty), whereas downstream transect samples are calculated from individual aliquot in shown in B. B) Modeled $^4\text{He}/^3\text{He}$ time-temperature (right panel) and ratio evolution spectra (left panel) results. Ratios of $^4\text{He}/^3\text{He}$ for each step relative to the bulk $^4\text{He}/^3\text{He}$ of a sample are plotted against the cumulative ^3He release fraction ($\Sigma^3\text{He}$; black boxes). Gray paths represent modeled cooling histories that do not predict the sample (U-Th)/He age to within the observed variance between analyzed crystals. Green paths are those that predict the (U-Th)/He age and have good fits to the observed spectra for that sample. Yellow and red paths also predict the age but have progressively worse fits. Black paths are the best fitting of the explored Pecube models.

Figure 6. Apatite (U-Th)/He data from this study plotted with approximate distance from the lip of the Jiacha knickpoint. Open circles are (U-Th)/He age predicted from HeFTy using thermal histories of corresponding sample nodes extracted from the best fitting of the explored Pecube models (see Section 5). Black circles are our observed $^4\text{He}/^3\text{He}$ sample ages from aliquots in Figure 5 with estimated 10%

uncertainty in age. Black triangles are our observed average (U-Th)/He ages with standard deviation for samples from the base of the knickpoint. Gray circles are apatite (U-Th)/He data from Zeitler et al. (2014) and were not included in Pecube model. Approximate sediment depth in Yarlung canyon estimated from Wang et al. (2014).

Figure 7. Schematic temporal progression of Yarlung-Parlung-Siang integration scenarios consistent with present day knickpoint locations. Unlabeled tributaries are those analyzed here that contain knickpoints at ~3500 m. Black circles are locations of ~3500 m elevation knickpoints on major rivers of the region. Black arrows depict flow directions. Panels A, C, and D demonstrate an antecedent Yarlung-Siang drainage captures the Parlung prior to migration of the Jiacha knickpoint from the Siang. Panels B, C, and D show a similar scenario with a Yarlung-Parlung drainage prior to its capture by a headward-cutting Siang. Regardless of pre-integration configuration, integration by ~10 Ma is required to account for Parlung and tributary knickpoints.

Figure

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7