A process-based model for non-equilibrium clumped isotope effects in carbonates

J.M. Watkins^{*1} and J.D. Hunt²

1. watkins4@uoregon.edu / Dept. of Geological Sciences, University of Oregon

2. Ramboll Environ

August 26, 2015

Abstract

The equilibrium clumped isotope composition of carbonate minerals is independent of the composition of the aqueous solution. However, many carbonate minerals grow at rates that place them in a non-equilibrium regime with respect to carbon and oxygen isotopes with unknown consequences for clumped isotopes. We develop a process-based model that allows one to calculate the oxygen, carbon, and clumped isotope composition of calcite as a function of temperature, crystal growth rate, and solution pH. In the model, carbon and oxygen isotope fractionation occurs through the mass-dependent attachment/detachment kinetics of the isotopologues of HCO_3^- and CO_3^{2-} to and from the calcite surface, which in turn, influence the clumped isotope composition of calcite. At experimental and biogenic growth rates, the mineral is expected to inherit a clumped isotopic composition that is similar to that of the DIC pool, which helps to explain (1) why different organisms share the same clumped isotope versus temperature calibration curves, (2) why many inorganic calibration curves are slightly but significantly different from one another, and (3) why foraminifera, coccoliths, and deep sea corals can have near-equilibrium clumped isotope compositions but far-from-equilibrium carbon and oxygen isotope compositions. Some aspects of the model can be generalized to other mineral systems and should serve as a useful reference in future efforts to quantify kinetic clumped isotope effects.

Keywords: clumped isotopes, oxygen isotopes, carbon isotopes, equilibrium, kinetic, calcite,

1 Introduction

Clumped isotope geochemistry, or the study of bond ordering in molecules, is at the frontier of efforts to use temperature proxies for reconstructing ocean chemistry (Came et al., 2007), estimating the uplift rate of mountains (Ghosh et al., 2006b), studying diagenetic processes (Dennis and Schrag, 2010), sourcing methane (Stolper et al., 2014, 2015; Wang et al., 2015), identifying signatures of biology in the atmosphere (Yeung et al., 2015) and even investigating the thermal physiology of dinosaurs (Eagle et al., 2011). The most widely-used clumped isotope thermometer is based on the homogeneous reaction (Ghosh et al., 2006a):

which describes the extent to which the rare isotopes ${}^{13}\text{C}$ and ${}^{18}\text{O}$ are bound together as ‘clumps’ among the carbonate molecules. The equilibrium constant for this reaction, which can be applied to either dissolved carbonate molecules or carbonate molecules in minerals such as calcite, has a well-defined temperature-dependence based on equilibrium thermodynamics (Wang et al., 2004; Schauble et al., 2006) and is measured by expressing the proportion of mass 63 isotopologues relative to what one would expect for a stochastic distribution of isotopes among the carbonate isotopologues (Ghosh et al., 2006a).

Because the clumped isotope thermometer is based on equilibrium thermodynamics, it potentially suffers from the same issue as oxygen isotope thermometry: many carbonates that form in nature grow at rates that place them in a non-equilibrium regime (Dietzel et al., 2009). There are clear instances, for example in speleothems and surface corals, where clumped isotope compositions yield significant over- or under-estimates of temperature, representing clear departures from equilibrium (Affek et al., 2008; Daëron et al., 2011; Saenger et al., 2012; Affek, 2012; Eiler et al., 2014). There are other cases, namely in deep-sea corals, where clumped isotope compositions are apparently equilibrated even when there are significant departures from carbon and oxygen isotope equilibrium (Thiagarajan et al., 2011; Eiler et al., 2014). Several ideas have been put forth to explain these observations, but because of the complexity of clumped isotope systematics relative to oxygen or

carbon isotope systematics, the models have remained largely conceptual (Tang et al., 2014; Tripathi et al., 2015) or based on analogies to gas phase processes such as Knudsen diffusion and mixing of indestructible molecules (Thiagarajan et al., 2011; Henkes et al., 2013).

In this paper we employ a process-based crystal growth model to investigate the complex interplay between carbon, oxygen, and clumped isotope systematics during carbonate growth under non-equilibrium conditions. The non-equilibrium isotope effects attending carbonate precipitation are quantified by considering the mass-dependent transport of isotopologues of CO_3^{2-} and HCO_3^- to and from the mineral surface. In this framework, the carbon isotope composition is calculated from the ratio of $^{13}\text{C}^{16}\text{O}^{16}\text{O}$ (mass 61) to $^{12}\text{C}^{16}\text{O}^{16}\text{O}$ (mass 60) ions incorporated into the mineral, and the oxygen isotope composition is calculated from the ratio of $^{12}\text{C}^{18}\text{O}^{16}\text{O}$ (mass 62) to $^{12}\text{C}^{16}\text{O}^{16}\text{O}$ (mass 60) incorporated into the mineral. Since each dissolved inorganic carbon (DIC = $\text{CO}_2(\text{aq}) + \text{HCO}_3^- + \text{CO}_3^{2-}$) species has a different equilibrium carbon, oxygen, and clumped isotopic composition (Zeebe and Wolf-Gladrow, 2001; Beck et al., 2005; Hill et al., 2014), the net carbon and oxygen isotopic composition of calcite depends on the relative proportions of HCO_3^- and CO_3^{2-} participating in calcite growth, the isotopic compositions of dissolved HCO_3^- and CO_3^{2-} , and any isotopic fractionations (equilibrium or kinetic) between calcite and each species that arise during ion transport onto, or away from, the mineral surface. The processes could include aqueous diffusion, isotope exchange reactions in aqueous solution, desolvation, surface diffusion and isotope exchange reactions at or near the aqueous-mineral interface. For clumped isotopes, the $^{13}\text{C}^{18}\text{O}^{16}\text{O}$ isotopologue (mass 63) is postulated to follow a mass fractionation law that is mathematically related to the behavior of the mass 62 and mass 61 isotopologues.

2 Notation

2.1 Notation for carbon and oxygen isotopes

The difference in isotope composition between any two chemical species or phases can be expressed in terms of an isotopic fractionation factor (α). The fractionation factors between calcite and DIC (for carbon isotopes) and calcite and water (for oxygen isotopes) are given by:

$$\alpha_{\text{xtl-DIC}}^{\text{C}} = \frac{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{xtl}}}{\left(\frac{^{13}\text{C}}{^{12}\text{C}}\right)_{\text{DIC}}}, \quad (2)$$

75 and

$$\alpha_{\text{xtl-w}}^{\text{O}} = \frac{\left(\frac{^{18}\text{O}}{^{16}\text{O}}\right)_{\text{xtl}}}{\left(\frac{^{18}\text{O}}{^{16}\text{O}}\right)_{\text{w}}}, \quad (3)$$

76 where the subscript ‘xtl’ refers to calcite and ‘w’ refers to water. The *equilibrium* isotope compo-
 77 sitions for species dissolved in aqueous solution are plotted as curves in Fig. 1. The fractionation
 78 factors used to construct these curves are provided in Table 1 along with the appropriate references.
 79 Note that the oxygen isotope composition of DIC varies with pH because it represents a weighted
 80 average of the isotopic composition of individual species. The carbon isotope composition of DIC
 81 is also a weighted average of DIC, but is constant with pH because DIC is the only carbon in the
 82 system (there is no carbon in H₂O). Hence, the carbon isotope composition of each dissolved species
 83 must vary as a consequence of mass balance (cf. Zeebe and Wolf-Gladrow, 2001).

84 All model calculations in this paper will be done under the assumption that the system is wa-
 85 ter buffered such that the equilibrium ¹⁸O/¹⁶O values of the DIC species are controlled solely by
 86 temperature and the ¹⁸O/¹⁶O of the water. We also assume an infinite reservoir such that the DIC
 87 concentration and ¹³C/¹²C of DIC are constant in all calculations; i.e., there is no distillation of
 88 carbon isotopes or change in pH as the carbonate precipitates.

89 2.2 Notation for clumped isotopes

90 The clumped isotope composition of calcite, denoted Δ_{xtl}^{63} , is given by

$$\Delta_{\text{xtl}}^{63} = 1000 \left(\frac{r_{\text{xtl}}^{63}}{r_{\text{xtl}}^{63*}} - 1 \right), \quad (4)$$

91 where

$$r_{\text{xtl}}^{63} = \frac{^{13}\text{C}^{18}\text{O}^{16}\text{O}^{16}\text{O}}{^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}, \quad (5)$$

92 and the asterisk in Eq. 4 denotes the stochastic ratio, which can be calculated from knowledge
 93 of the oxygen and carbon isotope compositions (Eiler and Schauble, 2004). Note that clumped
 94 isotope compositions are commonly expressed as Δ^{47} because the measurement involves ¹³C-¹⁸O
 95 bond ordering in CO₂(g) released upon acid digestion of carbonates. Conversion of Δ^{63} to Δ^{47} is
 96 accomplished via an acid digestion fractionation factor (AFF):

$$\Delta^{47} = \Delta^{63} + \text{AFF}, \quad (6)$$

where the AFF depends on the temperature of acid digestion (Defliese et al., 2015), the sample size (Wacker et al., 2013), and perhaps also the clumped isotope composition of the carbonate mineral itself (Guo et al., 2009). These details are important to consider when comparing model results to measurements on natural samples.

2.3 Isotope versus isotopologue ratios

If we consider that only the singly-substituted ^{18}O isotopologue contributes to the oxygen isotope composition of calcite, we have the following relationships between isotope ratios (R) and isotopologue ratios (r):

$$R_{\text{xtl}}^{\text{O}} = \left(\frac{[^{18}\text{O}]}{[^{16}\text{O}]} \right)_{\text{xtl}} \approx \frac{1}{3} r_{\text{xtl}}^{62} = \frac{1}{3} \frac{\text{Ca}^{12}\text{C}^{18}\text{O}^{16}\text{O}^{16}\text{O}}{\text{Ca}^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}, \quad (7)$$

and

$$R_{\text{xtl}}^{\text{C}} = \left(\frac{[^{13}\text{C}]}{[^{12}\text{C}]} \right)_{\text{xtl}} \approx r_{\text{xtl}}^{61} = \frac{\text{Ca}^{13}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}{\text{Ca}^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}. \quad (8)$$

The factor of 1/3 for oxygen isotopes comes from the fact that there are three oxygen atoms in the CO_3^{2-} molecule, and the approximate signs come from the recognition that the singly-substituted isotopologues are not the only, but by far the most abundant, ^{18}O - or ^{13}C -bearing carbonate molecules (Usdowski and Hoefs, 1993; see Appendix in Watkins et al., 2014 for more details).

For the dissolved species in solution, we use the shorthand notation $\text{B}_1 = \text{CO}_3^{2-}$ and $\text{B}_2 = \text{HCO}_3^-$ and write

$$R_{\text{B}_1}^{\text{O}} = \left(\frac{[^{18}\text{O}]}{[^{16}\text{O}]} \right)_{\text{B}_1} \approx \frac{1}{3} r_{\text{B}_1}^{62} = \frac{1}{3} \frac{^{12}\text{C}^{18}\text{O}^{16}\text{O}^{16}\text{O}}{^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}, \quad (9)$$

$$R_{\text{B}_1}^{\text{C}} = \left(\frac{[^{13}\text{C}]}{[^{12}\text{C}]} \right)_{\text{B}_1} \approx r_{\text{B}_1}^{61} = \frac{^{13}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}{^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}, \quad (10)$$

$$R_{\text{B}_2}^{\text{O}} = \left(\frac{[^{18}\text{O}]}{[^{16}\text{O}]} \right)_{\text{B}_2} \approx \frac{1}{3} r_{\text{B}_2}^{62} = \frac{1}{3} \frac{\text{H}^{12}\text{C}^{18}\text{O}^{16}\text{O}^{16}\text{O}}{\text{H}^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}, \quad (11)$$

and

$$R_{\text{B}_2}^{\text{C}} = \left(\frac{[^{13}\text{C}]}{[^{12}\text{C}]} \right)_{\text{B}_2} \approx r_{\text{B}_2}^{61} = \frac{\text{H}^{13}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}{\text{H}^{12}\text{C}^{16}\text{O}^{16}\text{O}^{16}\text{O}}. \quad (12)$$

Note that here, and throughout the paper, we ignore the contribution of hydrogen to the mass of HCO_3^- isotopologues such that mass 63 refers to the doubly-substituted isotopologue of HCO_3^- as

117 well as CO_3^{2-} .

118 3 Ion-by-ion model for carbon and oxygen isotopes

119 Kinetic isotope effects in carbonates can arise from incomplete carbon and oxygen isotope exchange
 120 among DIC species or during transfer of HCO_3^- and CO_3^{2-} ions to the mineral surface and into
 121 the mineral lattice. In the simplest scenario, HCO_3^- and CO_3^{2-} exchange kinetics in the aqueous
 122 solution are fast enough such that the isotopic compositions of HCO_3^- and CO_3^{2-} are the known equi-
 123 librium values (Beck et al., 2005; Zeebe, 2007; Wang et al., 2013). The picture is considerably more
 124 complicated when the DIC species are not isotopically equilibrated because, at any given instant,
 125 the isotopic compositions of HCO_3^- and CO_3^{2-} are poorly constrained due to limited knowledge of
 126 the isotope exchange kinetics and pathways associated with (de-)hydration and (de-)hydroxylation
 127 of dissolved CO_2 (Guo, 2009). It is therefore constructive to focus on the case where the isotopic
 128 composition of HCO_3^- and CO_3^{2-} are known and constant (i.e., equilibrated) and then discuss the
 129 consequences of an unequilibrated DIC pool.

130 Our starting point is a model for a cubic crystal (an approximation to the calcite rhombohedron)
 131 where the attachment and detachment fluxes (moles $\text{m}^{-2} \text{s}^{-1}$) of calcium and (bi-)carbonate ions
 132 to and from the mineral surface are given by (Wolthers et al., 2012):

$$J_{\text{attach}}^{\text{ion}} = \frac{\rho_c (\text{AR}^{\text{ion}}) a^2 d}{y_0} \quad (13)$$

133 and

$$J_{\text{detach}}^{\text{ion}} = \frac{\rho_c (\text{DR}^{\text{ion}}) a^2 d}{y_0}, \quad (14)$$

134 where ρ_c is the steady state kink site density (dimensionless), AR and DR are attachment rates and
 135 detachment rates (s^{-1}) of each ion, $a = 3.199 \times 10^{-10}$ is the closest spacing between adjacent Ca^{2+}
 136 and CO_3^{2-} sites on the calcite surface, $d = 27100$ moles m^{-3} is the molar density of calcite and y_0 (m)
 137 is the step spacing. The attachment and detachment rates are calculated from the model of Wolthers
 138 et al. (2012) and depend on factors such as temperature, pH, salinity, and the calcium:carbonate
 139 ratio of the solution (see next section). The net flux of an individual ion is given by the difference

$$J_{\text{net}}^{\text{ion}} = J_{\text{attach}}^{\text{ion}} - J_{\text{detach}}^{\text{ion}}, \quad (15)$$

140 and the net growth or dissolution rate of the crystal (R_c) is given by the sum

$$R_c = J_{\text{net}}^{\text{Ca}^{2+}} = J_{\text{net}}^{\text{B}_1} + J_{\text{net}}^{\text{B}_2}. \quad (16)$$

141 3.1 Expressions for the carbon and oxygen isotope composition of non- 142 equilibrium calcite

143 In the ion-by-ion growth model for oxygen isotopes derived in our previous work (Watkins et al.,
144 2014), the difference in oxygen isotope composition between calcite and water is related to the
145 rates of isotopologue attachment to (AR^i ; i is isotopologue mass), and detachment from (DR^i),
146 the mineral surface, as well as the relative contribution of each DIC species to the overall kinetic
147 fractionation:

$$\alpha_{\text{xtl-w}}^{\text{O}} = \frac{\alpha_{\text{xtl-w}}^{\text{eq,O}} \text{AR}^{60}}{\alpha_{\text{xtl-w}}^{\text{eq,O}} (\text{AR}^{60} - \text{DR}^{60}) \eta + \text{DR}^{60}}, \quad (17a)$$

148 where

$$\eta = \frac{1 + \phi}{\alpha_{\text{xtl-B}_1}^{\text{f,O}} \alpha_{\text{B}_1-\text{w}}^{\text{eq,O}} + \phi \alpha_{\text{xtl-B}_2}^{\text{f,O}} \alpha_{\text{B}_2-\text{w}}^{\text{eq,O}}} \quad (17b)$$

149 and ϕ is the ratio of bicarbonate to carbonate dissolved in solution. The α 's are the calcite-water
150 fractionation factors in the slow growth limit ($\alpha_{\text{xtl-w}}^{\text{eq,O}}$) and fast growth limit ($\alpha_{\text{xtl-w}}^{\text{f,O}}$), as discussed
151 further below. These expressions are identical to Eq. 18a and 18b in Watkins et al., 2014. The
152 attachment and detachment rates (s^{-1}) of the major isotopologues are given by (Wolthers et al.,
153 2012):

$$\text{AR}^{60} = k_{\text{B}_1^{60}} [\text{B}_1^{60}] P_{\text{A}} + k_{\text{B}_2^{60}} [\text{B}_2^{60}] P_{\text{A}}, \quad (18)$$

154 and

$$\text{DR}^{60} = \nu_{\text{B}_1}^{60} P_{\text{B}_1}^{60} + \nu_{\text{B}_2}^{60} P_{\text{B}_2}^{60}, \quad (19)$$

155 where the k 's ($M^{-1}s^{-1}$) and ν 's (s^{-1}) are mass-dependent attachment and detachment frequencies
156 and the P 's are probabilities of a kink site being occupied by a bicarbonate, carbonate, or calcium
157 ion on the crystal surface. All of the values in Eqs. 18 and 19, which apply only to the mass 60
158 isotopologue, are treated as known quantities taken from Wolthers et al. (2012). Note that AR^i

includes the attachment rates of both B_1 and B_2 and DR^i includes the detachment rates of both B_1 and B_2 .

Because clumped isotope compositions are defined in terms of isotopologues, it is useful to cast Eqs. 17 in terms of isotopologue ratios instead of isotope ratios. It is also useful to have a consistent expression for the oxygen and carbon isotope fractionation factors; rather than cast oxygen isotope compositions relative to water and carbon isotope compositions relative to DIC, we hereafter write the non-equilibrium calcite isotope compositions relative to the carbonate ion for both carbon and oxygen isotopes. This makes for easier comparison between the expressions for carbon and oxygen isotopes and leads to cleaner expressions when we adapt the model for clumped isotopes. Expanding the α 's and combining Eqs. 17a and 17b leads to:

$$\frac{r_{\text{xtl}}^{62}}{3R_w^O} = \frac{\frac{r_{\text{xtl}}^{\text{eq},62}}{3R_w^O} AR^{60}}{\frac{r_{\text{xtl}}^{\text{eq},62}}{3R_w^O} (AR^{60} - DR^{60}) (1 + \phi) + DR^{60}} + \frac{\alpha_{\text{xtl}-B_1}^{\text{f},62} \frac{r_{B_1}^{\text{eq},62}}{3R_w^O} + \phi \alpha_{\text{xtl}-B_2}^{\text{f},62} \frac{r_{B_2}^{\text{eq},62}}{3R_w^O}}{AR^i}, \quad (20)$$

where the factors of $3R_w^O$ cancel out. Note that $\alpha_{\text{xtl}-B_1}^{\text{f},O} = \alpha_{\text{xtl}-B_1}^{\text{f},62}$ because the factor of 3 that relates oxygen isotope ratios to isotopologue ratios in the crystal also shows up in the expressions for DIC species, and cancels out. The molar ratio of any isotopologue with mass i relative to the mass 60 isotopologue can then be given by:

$$r_{\text{xtl}}^i = \frac{r_{\text{xtl}}^{\text{eq},i} AR^{60}}{r_{\text{xtl}}^{\text{eq},i} (AR^{60} - DR^{60}) \frac{AR^{60}}{AR^i} + DR^{60}}, \quad (21a)$$

where (see Appendix A.1 for derivation)

$$\frac{AR^{60}}{AR^i} = \frac{(1 + \phi)}{\alpha_{\text{xtl}-B_1}^{\text{f},i} r_{B_1}^{\text{eq},i} + \phi \alpha_{\text{xtl}-B_2}^{\text{f},i} r_{B_2}^{\text{eq},i}}, \quad (21b)$$

which states that the non-equilibrium isotopic composition of calcite is a function of the specified equilibrium composition and the relative rates of ion attachment versus detachment of each isotopologue during crystal growth. The new expression for the oxygen isotope composition of calcite is:

$$\alpha_{\text{xtl-B}_1}^{\text{O}} = \frac{\text{AR}^{60} \alpha_{\text{xtl-B}_1}^{\text{eq,O}}}{\frac{\alpha_{\text{xtl-B}_1}^{\text{eq,O}} (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl-B}_1}^{\text{f,O}} + \phi \alpha_{\text{xtl-B}_2}^{\text{f,O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,O}}} + \text{DR}^{60}} \quad (22)$$

and the analogous expression for the carbon isotope composition of calcite is:

$$\alpha_{\text{xtl-B}_1}^{\text{C}} = \frac{\text{AR}^{60} \alpha_{\text{xtl-B}_1}^{\text{eq,C}}}{\frac{\alpha_{\text{xtl-B}_1}^{\text{eq,C}} (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl-B}_1}^{\text{f,C}} + \phi \alpha_{\text{xtl-B}_2}^{\text{f,C}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,C}}} + \text{DR}^{60}}, \quad (23)$$

where all terms with a superscript ‘eq’ are treated as knowns (Table 1). The AR^{60} and DR^{60} terms can be calculated using parameter values (attachment/detachment frequencies, kink formation energy, edge work, and speciation on the crystal surface) from the ion-by-ion calcite growth model of Wolthers et al. (2012) (see Tables 2 and 3 in Watkins et al., 2014, for a summary). This leaves the kinetic fractionation factors as the only ‘unknown’ quantities needed to solve for the isotopic composition of non-equilibrium calcite. The two functions above are plotted versus the model-derived crystal growth rate (cf. Watkins et al., 2014) in Figure 2 for arbitrary choices of the kinetic fractionation factors. The equations recover the equilibrium carbon and oxygen isotopic compositions in the limit of extremely slow growth. At fast growth rates, the model-derived isotopic compositions are dictated by the specified kinetic fractionation factors. More specifically, $\alpha_{\text{xtl-B}_1}^{\text{f}}$ represents a kinetic limit to the degree of isotopic fractionation between calcite and CO_3^{2-} while $\alpha_{\text{xtl-B}_2}^{\text{f}}$ represents a kinetic limit to the degree of isotopic fractionation between calcite and HCO_3^- . This is most clearly seen at pH values where only one of the two dissolved species is participating in calcite growth. At pH values between 7 and 12, both HCO_3^- and CO_3^{2-} are involved and the isotopic composition of calcite in the kinetic limit reflects the relative contributions of HCO_3^- and CO_3^{2-} (and their respective kinetic fractionation factors) to the isotopic composition of the mineral.

4 Ion-by-ion model for clumped isotopes

As is the case for carbon and oxygen isotopes, each DIC species in aqueous solution has a unique equilibrium clumped isotope composition (Hill et al., 2014). A potentially useful framework for constructing a process-based model for what governs bond ordering in carbonates is to consider the abundance of mass 63 isotopologues in HCO_3^- and CO_3^{2-} and the extent to which each species

200 contributes mass 63 molecules to the crystal. This is the approach that is followed below. Before we
 201 begin, it is important to emphasize that although the isotopic ion-by-ion growth model is constructed
 202 on the basis of molecular building blocks of calcite, it does not require that HCO_3^- and CO_3^{2-}
 203 be viewed as indestructible molecules throughout the steps involved in ion incorporation into the
 204 growing mineral (Watkins et al., 2014).

205 4.1 Calculating the Δ_{xtl}^{63} of non-equilibrium calcite

206 4.1.1 Expression for r_{xtl}^{63}

207 For the doubly-substituted isotopologue, we use Eq. 21 where $i = 63$:

$$r_{\text{xtl}}^{63} = \frac{\text{AR}^{60} r_{\text{xtl}}^{\text{eq},63}}{r_{\text{xtl}}^{\text{eq},63} (\text{AR}^{60} - \text{DR}^{60}) \frac{\text{AR}^{60}}{\text{AR}^{63}} + \text{DR}^{60}}, \quad (24)$$

208 which states that the non-equilibrium abundance of mass 63 isotopologues in calcite is a function of
 209 the specified equilibrium composition and the relative rates of ion attachment versus detachment.
 210 As is the case for oxygen and carbon isotopes, the attachment-detachment kinetics for the heavier
 211 isotopologues differ slightly from those for the mass 60 isotopologue. The ratio of isotopologue-
 212 specific attachment rates is folded into the $\text{AR}^{60}/\text{AR}^{63}$ term:

$$\frac{\text{AR}^{60}}{\text{AR}^{63}} = \frac{1 + \phi}{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},63} r_{\text{B}_1}^{\text{eq},63} + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},63} r_{\text{B}_2}^{\text{eq},63}}, \quad (25)$$

213 where

$$r_{\text{B}_1}^{\text{eq},63} = r_{\text{B}_1*}^{\text{eq},63} \left(\frac{\Delta_{\text{B}_1}^{\text{eq},63}}{1000} + 1 \right) \quad (26)$$

214 and

$$r_{\text{B}_2}^{\text{eq},63} = r_{\text{B}_2*}^{\text{eq},63} \left(\frac{\Delta_{\text{B}_2}^{\text{eq},63}}{1000} + 1 \right). \quad (27)$$

215 By substitution we now have

$$r_{\text{xtl}}^{63} = \frac{\text{AR}^{60} r_{\text{xtl}*}^{\text{eq},63} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right)}{r_{\text{xtl}*}^{\text{eq},63} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right) (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi) + \text{DR}^{60} \left(\alpha_{\text{xtl}-\text{B}_1}^{\text{f},63} r_{\text{B}_1*}^{\text{eq},63} \left(\frac{\Delta_{\text{B}_1}^{\text{eq},63}}{1000} + 1 \right) + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},63} r_{\text{B}_2*}^{\text{eq},63} \left(\frac{\Delta_{\text{B}_2}^{\text{eq},63}}{1000} + 1 \right) \right)}. \quad (28)$$

216 Values for $\Delta_{\text{B}_1}^{\text{eq},63}$, $\Delta_{\text{B}_2}^{\text{eq},63}$, $\Delta_{\text{xtl}}^{\text{eq},63}$ can be taken from Hill et al. (2014), so all that is needed to obtain
 217 a model-derived Δ_{xtl}^{63} are expressions for the stochastic ratios for B_1 , B_2 , equilibrium calcite and
 218 non-equilibrium calcite.

219 4.1.2 Expressions for the equilibrium stochastic ratios

220 Computation of the stochastic ratio of mass 63 to mass 60 isotopologues in each of the dissolved
 221 species and/or calcite requires knowledge of the $^{18}\text{O}/^{16}\text{O}$ and $^{13}\text{C}/^{12}\text{C}$ ratios. Without ^{17}O , there
 222 are three indistinguishable $^{13}\text{C}^{16}\text{O}^{16}\text{O}^{18}\text{O}$ molecules of mass 63 (the ^{18}O can be in any of the three
 223 sites), making the conversion straightforward. The stochastic ratio is

$$r_*^{63} = \frac{3 \left[^{13}\text{C} \right] \left[^{16}\text{O} \right] \left[^{16}\text{O} \right] \left[^{18}\text{O} \right]}{\left[^{12}\text{C} \right] \left[^{16}\text{O} \right] \left[^{16}\text{O} \right] \left[^{16}\text{O} \right]} = 3R^{\text{C}} R^{\text{O}}, \quad (29)$$

224 which can be calculated for a given T , pH, and growth rate from the output of the ion-by-ion growth
 225 model for carbon and oxygen isotopes. The stochastic ratio for each of the equilibrated DIC species
 226 and equilibrium calcite can be written in terms of their oxygen and carbon isotopic compositions,
 227 which in turn, can be written in terms of the oxygen and carbon isotope compositions of CO_3^{2-} and
 228 conventional fractionation factors:

$$r_{\text{B}_1*}^{\text{eq},63} = 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \quad (30)$$

229

$$r_{\text{B}_2*}^{\text{eq},63} = 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq},\text{O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq},\text{C}} \quad (31)$$

230

$$r_{\text{xtl}*}^{\text{eq},63} = 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq},\text{O}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq},\text{C}} \quad (32)$$

231 The equilibrium α 's in the above equations are known from experiments (Beck et al., 2005; Zeebe,
 232 2007; Wang et al., 2013) and natural cave calcites (Coplen, 2007; Kluge et al., 2014).

233 4.1.3 Non-stochastic ratio for non-equilibrium calcite

234 We can now substitute the equations from the previous section into Eq. 28 to obtain:

$$r_{\text{xtl}}^{63} = \frac{\text{AR}^{60} 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,O}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right)}{3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,O}} \alpha_{\text{c}-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right) (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)} + \text{DR}^{60} \\ \frac{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},63} 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \left(\frac{\Delta_{\text{B}_1}^{\text{eq},63}}{1000} + 1 \right) + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},63} 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{B}_2}^{\text{eq},63}}{1000} + 1 \right)}{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},63} 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \left(\frac{\Delta_{\text{B}_1}^{\text{eq},63}}{1000} + 1 \right) + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},63} 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{B}_2}^{\text{eq},63}}{1000} + 1 \right)} + \text{DR}^{60} \quad (33)$$

235 where $3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}}$ cancels out of the denominator, leaving:

$$r_{\text{xtl}}^{63} = \frac{\text{AR}^{60} 3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,O}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right)}{\alpha_{\text{xtl}-\text{B}_1}^{\text{eq,O}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right) (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)} + \text{DR}^{60} \\ \frac{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},63} \left(\frac{\Delta_{\text{B}_1}^{\text{eq},63}}{1000} + 1 \right) + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},63} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{B}_2}^{\text{eq},63}}{1000} + 1 \right)}{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},63} \left(\frac{\Delta_{\text{B}_1}^{\text{eq},63}}{1000} + 1 \right) + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},63} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{B}_2}^{\text{eq},63}}{1000} + 1 \right)} + \text{DR}^{60} \quad (34)$$

236 4.1.4 Stochastic ratio for non-equilibrium calcite

237 The stochastic ratio for non-equilibrium calcite can be obtained by using Eq. 29 and the definitions
238 of conventional fractionation factors to obtain:

$$r_{\text{xtl}*}^{63} = 3R_{\text{B}_1}^{\text{C}} R_{\text{B}_1}^{\text{O}} \alpha_{\text{xtl}-\text{B}_1}^{\text{C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{O}}. \quad (35)$$

239 Further substitution of Eq. 22 and 23 into Eq. 35 leads to:

$$r_{\text{xtl}*}^{63} = \frac{3R_{\text{B}_1}^{\text{O}} R_{\text{B}_1}^{\text{C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,C}} \alpha_{\text{xtl}-\text{B}_1}^{\text{eq,O}} (\text{AR}^{60})^2}{\left(\frac{\alpha_{\text{xtl}-\text{B}_1}^{\text{eq,C}} (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},\text{C}} + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},\text{C}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,C}}} + \text{DR}^{60} \right) \left(\frac{\alpha_{\text{xtl}-\text{B}_1}^{\text{eq,O}} (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl}-\text{B}_1}^{\text{f},\text{O}} + \phi \alpha_{\text{xtl}-\text{B}_2}^{\text{f},\text{O}} \alpha_{\text{B}_2-\text{B}_1}^{\text{eq,O}}} + \text{DR}^{60} \right)}. \quad (36)$$

240 This general expression links directly, for the first time, the stochastic ratio in calcite to physical
241 quantities that describe the growth of calcite under non-equilibrium conditions. Validation of this
242 expression is provided in Appendix A.2.

243 4.1.5 Generate final Δ_{xtl}^{63}

244 We can now combine the non-stochastic ratio (Eq. 34) and the stochastic ratio (Eq. 36) to get the
245 final Δ_{xtl}^{63} :

$$\Delta_{\text{xtl}}^{63} = 1000 \left(\frac{r_{\text{xtl}}^{63}}{r_{\text{xtl}*}^{63}} - 1 \right), \quad (37)$$

246 where

$$\frac{r_{\text{xtl}}^{63}}{r_{\text{xtl}*}^{63}} = \frac{\left(\frac{\alpha_{\text{xtl}-B_1}^{\text{eq,C}} (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl}-B_1}^{\text{f,C}} + \phi \alpha_{\text{xtl}-B_2}^{\text{f,C}} \alpha_{B_2-B_1}^{\text{eq,C}}} + \text{DR}^{60} \right) \left(\frac{\alpha_{\text{xtl}-B_1}^{\text{eq,O}} (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl}-B_1}^{\text{f,O}} + \phi \alpha_{\text{xtl}-B_2}^{\text{f,O}} \alpha_{B_2-B_1}^{\text{eq,O}}} + \text{DR}^{60} \right) \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right)}{\text{AR}^{60} \left(\text{DR}^{60} + \frac{\alpha_{\text{xtl}-B_1}^{\text{eq,O}} \alpha_{\text{xtl}-B_1}^{\text{eq,C}} \left(\frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1 \right) (\text{AR}^{60} - \text{DR}^{60}) (1 + \phi)}{\alpha_{\text{xtl}-B_1}^{\text{f},63} \left(\frac{\Delta_{B_1}^{\text{eq},63}}{1000} + 1 \right) + \phi \alpha_{\text{xtl}-B_2}^{\text{f},63} \alpha_{B_2-B_1}^{\text{eq,O}} \alpha_{B_2-B_1}^{\text{eq,C}} \left(\frac{\Delta_{B_2}^{\text{eq},63}}{1000} + 1 \right)} \right)}. \quad (38)$$

247 Most of the symbols in Eq. 38 can be treated as known quantities; if the equilibrium and kinetic
248 fractionation factors for carbon and oxygen isotopes are known or specified, then the only two free
249 parameters are the kinetic fractionation factors for the mass 63 isotopologue. As we show in the
250 next section, however, the kinetic parameters for the mass 63 isotopologue are in fact constrained
251 by the kinetic fractionation factors for the mass 62 and mass 61 isotopologues.

252 It should be noted that the equilibrium condition is recovered when the attachment rate is equal
253 to the detachment rate. Mathematically, $\text{AR} \rightarrow \text{DR}$, so $\text{AR} - \text{DR} \rightarrow 0$ and $\text{AR}/\text{DR} \rightarrow 1$, leading
254 to:

$$\frac{r_{\text{xtl}}^{63}}{r_{\text{xtl}*}^{63}} = \frac{\Delta_{\text{xtl}}^{\text{eq},63}}{1000} + 1. \quad (39)$$

255 In the kinetic limit, $\text{DR} \rightarrow 0$ and AR cancels out, leading to:

$$\frac{r_{\text{xtl}}^{63}}{r_{\text{xtl}*}^{63}} = \frac{(1 + \phi) \left(\alpha_{B_2-B_1}^{\text{eq,C}} \alpha_{B_2-B_1}^{\text{eq,O}} \alpha_{\text{xtl}-B_2}^{\text{f},63} \phi \left(\frac{\Delta_{B_2}^{63}}{1000} + 1 \right) + \alpha_{\text{xtl}-B_1}^{\text{f},63} \left(\frac{\Delta_{B_1}^{63}}{1000} + 1 \right) \right)}{\left(\alpha_{B_2-B_1}^{\text{eq,C}} \alpha_{\text{xtl}-B_2}^{\text{f,C}} \phi + \alpha_{\text{xtl}-B_1}^{\text{f,C}} \right) \left(\alpha_{B_2-B_1}^{\text{eq,O}} \alpha_{\text{xtl}-B_2}^{\text{f,O}} \phi + \alpha_{\text{xtl}-B_1}^{\text{f,O}} \right)}. \quad (40)$$

256 Although this expression does not lend itself to a simple interpretation, its behavior is remarkably
257 straightforward to interpret once the kinetic fractionation factors for the mass 63 isotopologue are
258 properly constrained.

5 What are the kinetic fractionation factors for clumped isotopes?

The fractionation factors for the mass 63 isotopologue are not independent of the fractionation factors for the mass 62 and 61 isotopologues. To see why this must be the case, we consider here the fractionation factors in the equilibrium limit and then apply the insights gained to the treatment of kinetic fractionation factors.

5.1 Relationship between doubly- and singly-substituted fractionation factors in the equilibrium limit

The equilibrium constant for the clumped isotope exchange reaction (Eq. 1) is

$$K_{\text{eq}}^{63} = \frac{[^{13}\text{C}^{18}\text{O}^{16}\text{O}_2^{2-}][^{12}\text{C}^{16}\text{O}_3^{2-}]}{[^{13}\text{C}^{16}\text{O}_3^{2-}][^{12}\text{C}^{18}\text{O}^{16}\text{O}_2^{2-}]} = \frac{[^{63}\text{CO}_3^{2-}][^{60}\text{CO}_3^{2-}]}{[^{61}\text{CO}_3^{2-}][^{62}\text{CO}_3^{2-}]} \quad (41)$$

Dividing all terms on the right-hand side of Eq. 41 by $^{60}\text{CO}_3^{2-}$ we can write the above expression in terms of ‘conventional’ isotopologue ratios:

$$K_{\text{eq,xtl}}^{63} = \frac{\left(\frac{[^{63}\text{CO}_3^{2-}]}{[^{60}\text{CO}_3^{2-}]}\right)_{\text{xtl}}}{\left(\frac{[^{61}\text{CO}_3^{2-}]}{[^{60}\text{CO}_3^{2-}]}\right)_{\text{xtl}} \left(\frac{[^{62}\text{CO}_3^{2-}]}{[^{60}\text{CO}_3^{2-}]}\right)_{\text{xtl}}}, \quad (42)$$

where the denominator now corresponds to the equilibrium carbon isotope composition multiplied by the equilibrium oxygen isotope composition of the crystal. Using the definition of equilibrium fractionation factors between calcite and dissolved CO_3^{2-} , Eq. 42 can be recast as

$$K_{\text{eq,xtl}}^{63} = \frac{\alpha_{\text{xtl-B}_1}^{\text{eq},63} \left(\frac{[^{63}\text{CO}_3^{2-}]}{[^{60}\text{CO}_3^{2-}]}\right)_{\text{B}_1}}{\alpha_{\text{xtl-B}_1}^{\text{eq},61} \left(\frac{[^{61}\text{CO}_3^{2-}]}{[^{60}\text{CO}_3^{2-}]}\right)_{\text{B}_1} \alpha_{\text{xtl-B}_1}^{\text{eq},62} \left(\frac{[^{62}\text{CO}_3^{2-}]}{[^{60}\text{CO}_3^{2-}]}\right)_{\text{B}_1}}, \quad (43)$$

which is the same as

$$K_{\text{eq,xtl}}^{63} = K_{\text{eq,B}_1}^{63} \frac{\alpha_{\text{xtl-B}_1}^{\text{eq},63}}{\alpha_{\text{xtl-B}_1}^{\text{eq},61} \alpha_{\text{xtl-B}_1}^{\text{eq},62}}. \quad (44)$$

274 Rearranging, we arrive at an expression for $\alpha_{\text{xtl-B}_1}^{\text{eq},63}$ as a function of known quantities:

$$\alpha_{\text{xtl-B}_1}^{\text{eq},63} = \frac{K_{\text{eq,xtl}}^{63}}{K_{\text{eq,B}_1}^{63}} \alpha_{\text{xtl-B}_1}^{\text{eq},61} \alpha_{\text{xtl-B}_1}^{\text{eq},62} \approx \alpha_{\text{xtl-B}_1}^{\text{eq},61} \alpha_{\text{xtl-B}_1}^{\text{eq},62}. \quad (45)$$

275 From the definitions of $K_{\text{eq,xtl}}^{63}$ (Eq. 42) and $\Delta_{\text{eq,xtl}}^{63}$ (Eq. 39) we can write

$$K_{\text{eq,xtl}}^{63} = \left(\frac{\Delta_{\text{eq,xtl}}^{63}}{1000} + 1 \right), \quad (46)$$

276 which allows us to recast Eq. 45 in terms of Δ^{63} values:

$$\alpha_{\text{xtl-B}_1}^{\text{eq},63} = \underbrace{\left(\frac{\Delta_{\text{eq,xtl}}^{63} + 1000}{\Delta_{\text{eq,B}_1}^{63} + 1000} \right)}_{\approx 1+10^{-5}} \alpha_{\text{xtl-B}_1}^{\text{eq},61} \alpha_{\text{xtl-B}_1}^{\text{eq},62}. \quad (47)$$

277 A similar expression exists for the equilibrium calcite-B₂ fractionation factor:

$$\alpha_{\text{xtl-B}_2}^{\text{eq},63} = \underbrace{\left(\frac{\Delta_{\text{eq,xtl}}^{63} + 1000}{\Delta_{\text{eq,B}_2}^{63} + 1000} \right)}_{\approx 1+10^{-5}} \alpha_{\text{xtl-B}_2}^{\text{eq},61} \alpha_{\text{xtl-B}_2}^{\text{eq},62}. \quad (48)$$

278 Eqs. 47 and 48 show how the fractionation factor for the mass 63 isotopologue is very nearly equal
 279 to the product of the fractionation factors for the mass 61 and mass 62 isotopologues. Physically,
 280 this reflects the fact that the ¹⁸O in the mass 63 isotopologue should fractionate from ¹⁶O in nearly
 281 the exact same manner as the ¹⁸O in the mass 62 isotopologue. That is, during ion transport
 282 to the mineral surface, the oxygen isotopes of the doubly-substituted mass 63 isotopologue will
 283 be fractionated by an amount given approximately by $\alpha_{\text{xtl-B}_1}^{\text{eq},62}$ while the carbon isotopes will be
 284 fractionated by an amount given approximately by $\alpha_{\text{xtl-B}_1}^{\text{eq},61}$. The relationship is approximate because
 285 secondary isotope effects (e.g., ¹³C affects the bonding behavior of ¹⁸O and vice versa) may create
 286 deviations from the approximation. The prefactors in Eqs. 47 and 48, which are nearly equal to
 287 unity, are important for ensuring that the equilibrium clumped isotope composition is recovered in
 288 the equilibrium (slow growth) limit.

5.2 Relationship between doubly- and singly-substituted fractionation factors in the kinetic limit

Using the expressions for the equilibrium fractionation factors as a guide, we propose the following form for the kinetic fractionation factors:

$$\alpha_{\text{xtl-B}_1}^{\text{f},63} = \alpha_{\text{xtl-B}_1}^{\text{f},61} \alpha_{\text{xtl-B}_1}^{\text{f},62} + \epsilon_{\text{B}_1} \quad (49)$$

and

$$\alpha_{\text{xtl-B}_2}^{\text{f},63} = \alpha_{\text{xtl-B}_2}^{\text{f},61} \alpha_{\text{xtl-B}_2}^{\text{f},62} + \epsilon_{\text{B}_2}, \quad (50)$$

where the ϵ_{B_1} and ϵ_{B_2} are expected to be small, on the order of 10^{-5} , and could plausibly be positive, negative, or zero. We treat $\epsilon_{\text{B}_1} = \epsilon_{\text{B}_2} = 0$ as the reference case before demonstrating that even small values of ϵ_{B_1} and ϵ_{B_2} have a large influence on the modeled kinetic clumped isotope effects.

6 Results and discussion

6.1 What is the true equilibrium clumped isotope composition?

The first step in performing calculations based on Eq. 38 is to specify the equilibrium carbon and oxygen isotope composition of calcite relative to dissolved CO_3^{2-} as well as the equilibrium clumped isotope composition of calcite. A conclusion drawn from recent laboratory experiments is that most biogenic and laboratory-grown carbonates form in a regime where their stable isotope compositions are not representative of true equilibrium (Dietzel et al., 2009; Watkins et al., 2013). In natural settings, it is possible for inorganic calcite to grow slowly enough for equilibrium to be achieved, and one such example appears to be inorganic calcite from Devils Hole cave, Nevada, which formed under conditions of very low supersaturation at 33.7°C (Coplen, 2007). The clumped isotope composition of Devils Hole calcite determined by Kluge et al. (2014) is in agreement with the most recent, experimentally-produced inorganic calibration curve of Zaarur et al. (2013), suggesting that (1) the slowly-grown calcites from Devils Hole are the best natural laboratory for anchoring the true equilibrium composition at their formation temperature of 33.7°C , and (2) it is possible to obtain near-equilibrium clumped isotope compositions in the laboratory.

However, there are now several conflicting clumped isotope calibration curves that have been

313 proposed, some of which are shown in Fig. 4. The Zaarur et al. (2013) calibration is similar to the
 314 Ghosh et al. (2006a) calibration whereas the Hill et al. (2014) calibration is similar to the Dennis and
 315 Schrag (2010) and Kluge et al. (2015) calibrations (see Fig. 4 and Hill et al. (2014) for comparisons).
 316 The curves for these different calibrations intersect at about 34°C, such that the Devils Hole cave
 317 cannot be used to argue in favor of one calibration over the others. Therefore, until other natural
 318 calcites can be identified as having formed under conditions that approach true thermodynamic
 319 equilibrium, assumptions must be made regarding the temperature dependence of the equilibrium
 320 $\Delta_{\text{calcite}}^{47}$. In the absence of available experimental data, calculated temperature dependencies of DIC
 321 species and calcite can be used (Hill et al., 2014). The following equations are fits to the calculated
 322 data of Hill et al. (2014) between 0 and 100 °C for calcite, CO_3^{2-} , and HCO_3^- :

$$\Delta_{\text{eq,xtl}}^{63} = 37.55 \frac{1000}{T_K^2} - 0.0296, \quad (51a)$$

$$\Delta_{\text{eq,B}_1}^{63} = 36.36 \frac{1000}{T_K^2} - 0.0397, \quad (51b)$$

324 and

$$\Delta_{\text{eq,B}_2}^{63} = 38.35 \frac{1000}{T_K^2} - 0.0291. \quad (51c)$$

325 We use these equations for self-consistency while recognizing that the Hill et al. (2014) results may
 326 in fact underestimate the difference between the clumped isotope composition of dissolved HCO_3^-
 327 and CO_3^{2-} . The Hill et al. (2014) study predicts $\Delta_{\text{HCO}_3^-}^{63} - \Delta_{\text{CO}_3^{2-}}^{63} \approx 0.033\text{‰}$, which is considerably
 328 less than the value of $0.063 \pm 0.006\text{‰}$ deduced from recent witherite experiments (Tripathi et al.,
 329 2015).

330 6.2 Kinetic fractionation factors

331 Once the equilibrium fractionation factors are specified, one must then input the kinetic fractionation
 332 factors for carbon and oxygen isotopes. For oxygen isotopes, the kinetic fractionation factors have
 333 been inferred from the pH-dependence of $\alpha_{\text{xtl-w}}^{\text{O}}$ measured from calcites precipitated in the presence
 334 of an isotopically equilibrated DIC pool (Watkins et al., 2014). For carbon isotopes, the existing
 335 data are restricted to $\text{pH} < 7.8$ (Romanek et al., 1992) from which the kinetic fractionation factors
 336 are inferred to be close to unity (Fig. 3). Although it is not immediately obvious, the choice
 337 of kinetic fractionation factors for oxygen and carbon isotopes is actually not important when it

comes to modeling the clumped isotope composition of non-equilibrium calcite because the kinetic fractionation factor for the mass 63 isotopologue is written as the product of the kinetic fractionation factors for carbon and oxygen isotopes (Eqs. 49 and 50). An outcome of this relationship is that the results for clumped isotopes have an extremely weak dependence on the input values for the carbon and oxygen kinetic fractionation factors because of how clumped isotope compositions depend upon the carbon and oxygen isotope composition of calcite via the stochastic distribution. For any reasonable range of carbon and oxygen kinetic fractionation factors, the results for clumped isotopes are essentially independent of those factors. As shown in the model outputs presented below, the important parameters dictating the rate- and pH-dependence of kinetic Δ^{63} effects in non-equilibrium calcite are the small ϵ values in Eqs. 49 and 50.

6.3 Model behavior

Although Eq. 38 is a complicated algebraic expression, the behavior that emerges is relatively straightforward. The modeled rate-dependence of Δ_{xtl}^{63} is presented in Fig. 5a for the reference case where ϵ_{B_1} and ϵ_{B_2} are equal to zero. The curves have a similar shape to those for oxygen isotopes and carbon isotopes; in the limit of extremely slow growth, the clumped isotope composition is equal to the specified equilibrium composition, and with increasing growth rate, there is an asymptotic approach to a kinetic limit on Δ_{xtl}^{63} . The pH-dependence for both the slow- and fast-growth limits is shown in Fig. 5b. When the ϵ_{B_1} and ϵ_{B_2} parameters are equal to zero, calcite acquires the clumped isotope composition of DIC in the limit of fast growth. This is analogous to the case for oxygen (or carbon) isotopes where the forward fractionation factors are equal to one.

There is no reason to believe *a priori* that the ϵ_{B_1} and ϵ_{B_2} parameters should be exactly equal to zero based on Eqs. 47 - 50. Figure 6 shows the pH-dependence at two different (relatively fast) growth rates for additional scenarios where ϵ_{B_1} and $\epsilon_{\text{B}_2} \neq 0$. Also shown are the corresponding curves for carbon and oxygen isotopes for arbitrary choices of the kinetic fractionation factors. A positive value of ϵ_{B_1} or ϵ_{B_2} describes an increase in bond ordering at the mineral surface relative to the dissolved species (curves labeled “1”). Conversely, a negative ϵ_{B_1} or ϵ_{B_2} corresponds to a decrease in bond ordering relative to the dissolved species (curves labeled “3”). It is also possible for ϵ_{B_1} and ϵ_{B_2} to have opposite signs but these scenarios are not shown to retain clarity in the figure. As illustrated in these figures, the same general principles that apply to oxygen and carbon isotope discrimination also apply to clumped isotopes: the isotopic composition of non equilibrium calcite

depends on (1) the clumped isotope compositions of HCO_3^- and CO_3^{2-} dissolved in solution, (2) the relative proportions of HCO_3^- and CO_3^{2-} participating in calcite growth, and (3) any increase or decrease in bond ordering that arises during all of the steps involved in ion transport into the mineral lattice, as quantified by the signs and magnitudes of ϵ_{B_1} and ϵ_{B_2} .

6.4 Implications

6.4.1 Inorganic calibrations of the clumped isotope thermometer

It is possible that many of the calcites grown experimentally have isotopic compositions that are offset from the true equilibrium value due to some combination of isotopic disequilibrium among DIC species, isotopic distillation of the DIC pool, pH effects, and growth rate effects. Two recent studies have made progress toward assessing the influence of one or more of these factors. Tang et al. (2014) grew inorganic calcite in an environment where pH and growth rate were controlled. They observed a relatively wide range in Δ_{xtl}^{47} at a given temperature of about 0.006‰, comparable to other experimental studies that lacked such controls (e.g., Ghosh et al., 2006a; Dennis and Schrag, 2010; Zaarur et al., 2013; Defliese et al., 2015). This observation, in combination with the fact that the time required to achieve clumped isotope equilibrium among DIC species is about 10 hours at 25°C (cf. Affek, 2013; Clog et al., 2015; Tang et al., 2014), suggests that an out-of-equilibrium DIC pool may be contributing to the scatter in the data from many previous studies in ways that are not well understood. According to our model, the data from Tang et al. (2014) would be unable to resolve the expected pH- or rate-dependence on Δ_{xtl}^{47} , consistent with their findings. More recently, Tripathi et al. (2015) grew calcite in the presence of the enzyme carbonic anhydrase, which promotes isotopic equilibration among DIC species, but since their experiments controlled neither pH nor growth rate, these results are also not directly comparable to model outputs. We anticipate that ongoing experiments designed to measure the pH- and rate-dependence of Δ^{47} of calcite grown from an isotopically equilibrated DIC pool will be useful for constraining the kinetic fractionation factors and interpreting data from previous inorganic calcite precipitation experiments. Meanwhile, the calculations presented thus far offer a semi-quantitative explanation for why the various calibrations of the clumped isotope thermometer are broadly similar despite the lack of experimental control of pH, growth rate, and the isotopic composition of the DIC pool; surface reaction controlled kinetic effects restrict Δ_{xtl}^{63} to a range of values between $\Delta_{\text{eq,HCO}_3^-}^{63}$ and $\Delta_{\text{eq,CO}_3^-}^{63}$ (assuming ϵ_{B_1} and $\epsilon_{\text{B}_2} = 0$)

that differ from the $\Delta_{\text{eq,xtl}}^{63}$ by only a few 0.001‰. Larger differences, where observed, would require that ϵ_{B_1} and $\epsilon_{\text{B}_2} \neq 0$, which could be a manifestation of isotopic disequilibrium among DIC species, as discussed further below.

6.4.2 Kinetic effects in forams, coccoliths, and deep-sea corals

An observation that has warranted considerable discussion is that planktic foraminifera, coccoliths, benthic foraminifera, and deep-sea corals exhibit large ‘vital effects’ with respect to $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ but little to no vital effect with respect to clumped isotopes (Tripathi et al., 2010; Thiagarajan et al., 2011). A number of hypotheses have been considered to explain this observation, including isotopic mixing of samples with different bulk isotopic compositions, incomplete exchange of carbon and oxygen isotopes among DIC species, diffusive transport of DIC species, and direct inheritance of isotopes from bulk DIC (Tripathi et al., 2010; Thiagarajan et al., 2011). The latter hypothesis has perhaps received the most support since it can explain (a) why so many different types of carbonates fall on the same Δ^{63} versus T calibration line (Tripathi et al., 2010; Affek, 2012; Henkes et al., 2013; Hill et al., 2014), (b) why there is no evidence for species-specific offsets in Δ_{xtl}^{47} , and (c) why $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ are more sensitive to pH than Δ_{xtl}^{47} . While this hypothesis may be approximately correct, there are clear instances where carbon and oxygen isotopes in carbonates are offset from the isotopic composition of DIC (Spero et al., 1997; Watkins et al., 2014), which precludes the conceptual model wherein carbonates quantitatively precipitate DIC from a finite reservoir near the site of calcification (Tripathi et al., 2010). Our calculations show that quantitative precipitation is not required for a carbonate mineral to have the same clumped isotope composition as that of DIC. Instead, we propose that carbonate minerals may acquire a clumped isotope composition similar to that of DIC as a consequence of mass balance and the fact that clumped isotope compositions are calculated relative to a stochastic distribution based on the actual $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values.

6.4.3 Kinetic effects in surface corals

The above conclusion does not directly address why some mollusks and brachiopods, surface corals, and speleothems are displaced from the main calibration line shared by other types of carbonates (Daëron et al., 2011; Saenger et al., 2012; Henkes et al., 2013; Eiler et al., 2014). Focusing on surface corals, for which pH and growth rate are known (Saenger et al., 2012), Fig. 7 shows the difference between the measured clumped isotope composition ($\Delta_{\text{meas}}^{47}$) and the expected clumped

isotope composition ($\Delta_{\text{Hill}}^{47}$). Here, the calculated Δ_{xtl}^{63} were converted to Δ_{xtl}^{47} values using an AFF of 0.28 (acid digestion at 25°C; Tripathi et al., 2015) for the comparison to experimental data. The observation that has been made previously is that surface corals are shifted systematically to higher Δ^{47} values, and therefore lower apparent temperatures, than would be expected from several calibrations of the Δ^{47} -temperature relationship (Saenger et al., 2012; Eiler et al., 2014). The current explanation for this shift is that hydration of excess CO_2 near the calcification site leads to a higher-than-equilibrium Δ^{47} of HCO_3^- which is then inherited by the organism. This conceptual model is quantified by the best-fit ϵ values being greater than 0, which means that the HCO_3^- molecules (and also perhaps the CO_3^{2-}) adsorbed on the mineral surface are less isotopically scrambled than their counterparts in the bulk solution. The ion-by-ion model cannot distinguish whether a positive ϵ is due to processes near the mineral surface such as desolvation, or whether it reflects an elevated Δ^{47} of HCO_3^- due to kinetics of (de-)hydroxylation and (de-)hydration reactions in the bulk solution. Like the kinetic fractionation factors for carbon and oxygen isotopes, the ϵ parameters are macroscopic in the sense that their values describe the net result of all processes contributing to mass fractionation during isotope exchange and mass transport to the growing mineral, including bond reordering at the mineral-solution interface (Tripathi et al., 2015).

7 Summary

We modified an existing ion-by-ion calcite growth model, which describes steady-state calcite growth through the attachment-detachment of Ca^{2+} , HCO_3^- and CO_3^{2-} ions to the crystal surface, to include the isotopologues of HCO_3^- and CO_3^{2-} . The reference scenario for the model is calcite precipitation from an equilibrated DIC pool, wherein the isotopic compositions of dissolved HCO_3^- and CO_3^{2-} are treated as known quantities. Carbon and oxygen isotopic fractionation between calcite and aqueous solution arises through the isotopologue-specific attachment and detachment rates. The clumped isotope composition of calcite is calculated from the carbon and oxygen isotope composition and the mass fractionation behavior of the doubly-substituted isotopologue. Although the model is constructed on the basis of molecular building blocks of calcite, this does not imply that the isotopologues of HCO_3^- and CO_3^{2-} are indestructible molecules that maintain their isotopic identity throughout the steps (e.g., aqueous diffusion, ion desolvation, surface diffusion and isotope exchange/reordering reactions at the aqueous-mineral interface) involved in ion incorporation into

the growing mineral. The equations represent a self-consistent framework for calculating the carbon, oxygen, and clumped isotope composition of calcite as a function of temperature, crystal growth rate, and solution pH. The derivations and several example outputs presented herein lead to the following key points:

1. The equilibrium fractionation factor for the mass 63 isotopologue can be written as a function of the equilibrium fractionation factors for the mass 62 and mass 61 isotopologues.
2. We propose that the kinetic fractionation factor for the mass 63 isotopologue is equal to the product of the kinetic fractionation factors for the mass 62 and mass 61 isotopologues plus or minus some small deviation, denoted with the symbol ϵ , that is on the order of 10^{-5} .
3. The ϵ parameters describe the extent to which the clumped isotope composition of HCO_3^- and CO_3^{2-} become scrambled or unscrambled during the many steps involved in ion transport to and from the mineral surface. Like the kinetic fractionation factors for oxygen and carbon isotopes, the ϵ parameters are macro-scale parameters that describe the net result of all processes contributing to isotopic fractionation or re-ordering.
4. If the ϵ parameters are equal to zero, the clumped isotope composition of a rapidly-grown carbonate mineral is nearly equal to that of DIC. This may or may not be a common situation. Nevertheless, at the normal pH of seawater, the clumped isotope composition of DIC and equilibrium calcite differ by only 0.01‰. Further, calcites grown at typical growth rates will have a clumped isotope composition between that of rapidly-grown carbonates and equilibrium calcite. This result offers a quantitative rationalization for how foraminifera, coccoliths, and deep sea corals can have near-equilibrium clumped isotope compositions but far-from-equilibrium carbon and oxygen isotope compositions (Tripathi et al., 2010; Thiagarajan et al., 2011). It may also help explain why so many different organisms (including those that precipitate aragonite instead of calcite) fall on the same clumped isotope-temperature curve(s).
5. In cases where the ϵ parameters are non-zero, the clumped isotope composition of a rapidly-grown carbonate mineral may deviate significantly from that of DIC. Such deviations may be the simple result of an unequilibrated DIC pool, but it is also conceivable that the ϵ parameters could be non-zero due to processes involved in the transport of HCO_3^- and CO_3^{2-} to and from

- 483 the mineral surface; i.e., surface reaction-controlled kinetic effects such as those described by
484 the Tripathi et al. (2015) conceptual model.
- 485 6. It is possible to match the observed rate-dependence observed in surface corals with positive ϵ
486 parameters. This result does not conflict with previous proposals that kinetic clumped isotope
487 effects in surface corals may be due to rapid hydration of excess CO_2 (Saenger et al., 2012;
488 Eiler et al., 2014).
- 489 7. For the model curves presented herein, we have assumed that the kinetic fractionation factors
490 for carbon, oxygen, and clumped isotopes do not depend temperature or other factors such as
491 pH or solution composition. This crude assumption may need to be relaxed if warranted by
492 new experimental data.
- 493 8. An particularly useful set of experiments for comparison to previous inorganic calcite calibra-
494 tions would involve monitoring the pH and growth rate, using a water buffered system with
495 respect to oxygen isotopes, ensuring there is no isotopic distillation of the DIC pool, and en-
496 suring that equilibrium among DIC species is at least approximately maintained during calcite
497 growth.
- 498 9. While the model is built upon an existing framework that is specific to the mineral calcite,
499 the steps involved in adapting the model for clumped isotopes should be useful to future
500 developments of quantitative descriptions of non-equilibrium clumped isotope effects in other
501 systems.

502 **Acknowledgments:** Mathieu Daeron’s online wiki and responses to inquiries on clumped isotopes
503 were very helpful. Section 5.1 benefited from a conversation with D. Stolper at the 2014 AGU
504 conference. We gratefully acknowledge the constructive reviews by Cedric John and Daniel Stolper.
505 This research was supported by the University of Oregon through startup funds to JMW.

Carbon isotopes

Compounds	Equation	α (25°C)	References
CO ₂ (g) - HCO ₃ ⁻	-9.483/ T_K +1.0239	0.9921	Mook et al. (1974) as in Mook (1986)
CO ₂ (aq) - HCO ₃ ⁻	-9.866/ T_K +1.0241	0.9910	Vogel et al. (1970) as in Mook (1986)
CO ₃ ²⁻ - HCO ₃ ⁻	-0.867/ T_K +1.0025	0.9996	Turner (1982) as in Mook (1986)
CO ₂ (g) - Calcite	$\Delta = -2.988 \cdot 10^6/T_K^2$ $+7.666 \cdot 10^3/T_K - 2.461$	0.9897	Bottinga (1968)

Oxygen isotopes

Compounds	Equation	α (25°C)	References
CO ₂ (g) - H ₂ O	17.611/ T_K + 0.9821	1.0412	Zeebe (2007)
CO ₂ (aq) - H ₂ O	17.54/ T_K + 0.9827	1.0415	Wang et al. (2013)
HCO ₃ ⁻ - H ₂ O	17.76/ T_K + 0.9725	1.0321	Wang et al. (2013)
CO ₃ ²⁻ - H ₂ O	21.72/ T_K + 0.9539	1.0268	Wang et al. (2013)
Calcite - H ₂ O	$\Delta = 17747/T_K - 29.777$	1.0302	Coplen (2007), Watkins et al. (2013)

Table 1: Compilation of equilibrium fractionation factors (α_{i-j}^{eq} unless otherwise noted) for carbon and oxygen isotopes in aqueous solution. These equations are presented graphically in Fig. 1 (note: $\Delta = 1000 \ln \alpha$).

Table 2: Descriptions of symbols used to calculate the isotopic composition of non-equilibrium calcite.

Symbol	Description
AR^{60}	Attachment rate of mass 60 isotopologue
DR^{60}	Detachment rate of mass 60 isotopologue
ϕ	HCO_3^- / CO_3^{2-} in solution
R_{xtl}^C	Molar ratio of $^{13}C/^{12}C$ in the crystal
R_{xtl}^O	Molar ratio of $^{18}O/^{16}O$ in the crystal
r_{xtl}^{61}	Molar ratio of mass 61 to mass 60 isotopologues in the crystal
$r_{xtl}^{eq,61}$	Equilibrium molar ratio of mass 61 to mass 60 isotopologues in the crystal
r_{xtl}^{62}	Molar ratio of mass 62 to mass 60 isotopologues in the crystal
r_{xtl}^{63}	Molar ratio of mass 63 to mass 60 isotopologues in the crystal
r_{xtl*}^{63}	Stochastic molar ratio of mass 63 to mass 60 isotopologues in the crystal
$\alpha_{xtl-B_1}^{eq,C}$	Equilibrium carbon isotope fractionation factor between calcite and dissolved CO_3^{2-}
$\alpha_{xtl-B_2}^{eq,C}$	Equilibrium carbon isotope fractionation factor between calcite and dissolved HCO_3^-
$\alpha_{xtl-B_1}^{eq,O}$	Equilibrium oxygen isotope fractionation factor between calcite and dissolved CO_3^{2-}
$\alpha_{xtl-B_2}^{eq,O}$	Equilibrium oxygen isotope fractionation factor between calcite and dissolved HCO_3^-
$\alpha_{xtl-B_1}^{eq,63}$	Equilibrium clumped isotope fractionation factor between calcite and dissolved CO_3^{2-}
$\alpha_{xtl-B_2}^{eq,63}$	Equilibrium clumped isotope fractionation factor between calcite and dissolved HCO_3^-
$\alpha_{B_2-B_1}^{eq,C}$	Equilibrium carbon isotope fractionation factor between dissolved HCO_3^- and CO_3^{2-}
$\alpha_{B_2-B_1}^{eq,O}$	Equilibrium oxygen isotope fractionation factor between dissolved HCO_3^- and CO_3^{2-}
$\alpha_{xtl-B_1}^{f,C}$	Kinetic carbon isotope fractionation factor between calcite and dissolved CO_3^{2-}
$\alpha_{xtl-B_2}^{f,C}$	Kinetic carbon isotope fractionation factor between calcite and dissolved HCO_3^-
$\alpha_{xtl-B_1}^{f,O}$	Kinetic oxygen isotope fractionation factor between calcite and dissolved CO_3^{2-}
$\alpha_{xtl-B_2}^{f,O}$	Kinetic oxygen isotope fractionation factor between calcite and dissolved HCO_3^-
$\alpha_{xtl-B_1}^{f,63}$	Kinetic clumped isotope fractionation factor between calcite and dissolved CO_3^{2-}
$\alpha_{xtl-B_2}^{f,63}$	Kinetic clumped isotope fractionation factor between calcite and dissolved HCO_3^-
ϵ_{B_1}	Describes the extent of isotopic scrambling (negative values) or unscrambling (positive values) of CO_3^{2-} during transport from the solution to the crystal
ϵ_{B_2}	Describes the extent of isotopic scrambling (negative values) or unscrambling (positive values) of HCO_3^- during transport from the solution to the crystal

Figure 1: Equilibrium stable isotope composition of calcite and dissolved inorganic carbon (DIC) species in aqueous solution. *Top*: Distribution of DIC species in aqueous solution as a function of pH. *Middle*: Oxygen isotope compositions expressed relative to water. Since water is by far the predominant oxygen-bearing species in aqueous solution, mass balance dictates that the oxygen isotope composition of DIC vary with pH. *Bottom*: Carbon isotope compositions expressed relative to DIC. Since all of the carbon in aqueous solution is stored in DIC, mass balance dictates that the carbon isotope composition of DIC be constant with pH. Also shown are proposed curves for the equilibrium stable isotope composition of calcite. See Table 2 for the equilibrium fractionation factors used to construct these curves.

Figure 2: Modeled stable isotope compositions as a function of growth rate and pH at 25°C. *Top*: Oxygen isotope systematics. *Bottom*: Carbon isotope systematics. All curves show compositions expressed relative to the stable isotope composition of dissolved CO_3^{2-} following Eqs. 22 and 23. At very low supersaturations (slow growth), the equilibrium composition of calcite is determined by the specified equilibrium fractionation factors. At high growth rates (kinetic limit), the composition of non-equilibrium calcite is determined by the kinetic fractionation factors and the relative proportions of HCO_3^- and CO_3^{2-} participating in calcite growth. For example, at pH = 7, HCO_3^- is the predominant species and the composition of calcite in the fast growth limit is offset from HCO_3^- by $1000 \ln \alpha_{\text{xtl-B}_2}^f$. Similarly, at pH = 12, CO_3^{2-} is the predominant species and the composition of calcite in the fast growth limit is offset from CO_3^{2-} by $1000 \ln \alpha_{\text{xtl-B}_1}^f$.

Figure 3: Experimental constraints on kinetic oxygen and carbon isotope fractionation factors. For both isotopic systems, the equilibrium isotopic composition is inferred from slowly-grown calcites at Devils Hole cave, Nevada (Coplen, 2007). (A) Results from experiments at $T = 25^\circ\text{C}$ and $\text{pH} = 8.3$ where inorganic calcite was precipitated in the presence of carbonic anhydrase (Watkins et al., 2014). (B) Results from experiments at $T = 25^\circ\text{C}$ and $\text{pH} = 6.6$ to 7.3 where inorganic calcite was precipitated in the presence of a DIC pool with known $\delta^{13}\text{C}$ (Romanek et al., 1992). The kinetic fractionation factors used to construct the model curves are $\alpha_{\text{xtl-B}_1}^{\text{f,O}} = 0.9980$, $\alpha_{\text{xtl-B}_2}^{\text{f,O}} = 0.9964$, $\alpha_{\text{xtl-B}_1}^{\text{f,C}} = 1.0000$, and $\alpha_{\text{xtl-B}_2}^{\text{f,C}} = 1.0000$.

Figure 4: Four different calibrations of the clumped isotope thermometer with 95% confidence intervals, presented in the absolute reference frame (Dennis et al., 2011) and normalized to a 25 °C acid digestion temperature. The agreement between slowly-grown Devils Hole calcite with the calibration curves has been used to argue that Devils Hole represents the best natural laboratory for measuring the true equilibrium composition (Kluge et al., 2014). Because the curves intersect at the approximate temperature of the Devils Hole samples, however, Devils Hole calcite cannot be used to argue in favor one calibration over the others.

Figure 5: Modeled rate- and pH-dependence of the clumped isotope composition of non-equilibrium calcite. *Top*: The rate-dependence behaves in much the same way as for carbon and oxygen isotopes (see Fig. 2) insofar as the clumped isotope composition in the fast-growth limit depends on the weighted sum of the clumped isotope compositions of DIC species participating in calcite growth. *Bottom*: The pH-dependence for the two limiting cases of slow growth and fast growth. If the ϵ_{B_1} and $\epsilon_{B_2} = 0$, fast-grown calcite acquires the clumped isotope composition of DIC, regardless of the kinetic fractionation factors for oxygen and carbon isotopes (see text). Although this figure only depicts processes at 25 °C, the second y-axis shows the apparent equivalent temperatures for several chosen Δ_i^{63} values using the Hill et al. (2014) equilibrium calcite calibration. Note that since calcite dissolves at low pH, the model only applies at pH above about 7.

Figure 6: Modeled pH-dependence of the non-equilibrium isotopic composition of calcite. *Left:* Composition of non-equilibrium calcite in the limit of extremely rapid crystallization. *Right:* Composition of non-equilibrium calcite at $10^{-6.4} \text{ mol m}^{-2} \text{ s}^{-1}$, which is representative of experimentally precipitated calcites (Dietzel et al., 2009; Tang et al., 2014). For carbon and oxygen isotopes, we use the following kinetic fractionation factors for illustrative purposes: $\alpha_{\text{xtl-B}_1}^{\text{f},\text{O}} = 0.9980$, $\alpha_{\text{xtl-B}_2}^{\text{f},\text{O}} = 0.9964$, $\alpha_{\text{xtl-B}_1}^{\text{f},\text{C}} = 0.9995$, $\alpha_{\text{xtl-B}_2}^{\text{f},\text{C}} = 0.9990$. For clumped isotopes, the carbon and oxygen fractionation factors are arbitrary for reasons discussed in the main text.

Figure 7: Measured versus modeled oxygen and clumped isotope compositions of coral calcites from Saenger et al. (2012). The surface corals are displaced to higher Δ_{xtl}^{47} values (or lower apparent temperatures) than expected based on numerous calibrations of the clumped isotope thermometer (cf. Eiler et al., 2014). These deviations can be fit with positive ϵ_{B_1} and ϵ_{B_2} values, which represent a net increase in bond ordering within dissolved HCO_3^- and CO_3^{2-} as a result of all transport processes, including possible disequilibrium among DIC species, that ultimately lead to ion incorporation into the mineral.

A Appendix

A.1 Derivation of the $\text{AR}^{60}/\text{AR}^i$ expression

The attachment-detachment kinetics for the higher mass isotopologues differ slightly from those for the mass 60 isotopologue. The ratio of isotopologue-specific attachment frequencies can be used to define two kinetic (forward) fractionation factors, one for B_1 and one for B_2 as (see Eq. 16a and 16b in Watkins et al., 2014):

$$\alpha_{\text{xtl-B}_1}^{\text{f},i} = \frac{k_{\text{B}_1}^i}{k_{\text{B}_1}^{60}}, \quad (\text{A.1})$$

and

$$\alpha_{\text{xtl-B}_2}^{\text{f},i} = \frac{k_{\text{B}_2}^i}{k_{\text{B}_2}^{60}}. \quad (\text{A.2})$$

We can then start with the expression for the attachment rates of each isotopologue:

$$\frac{\text{AR}^{60}}{\text{AR}^i} = \frac{k_{\text{B}_1}^{60}[\text{B}_1^{60}]P_{\text{A}} + k_{\text{B}_2}^{60}[\text{B}_2^{60}]P_{\text{A}}}{k_{\text{B}_1}^i[\text{B}_1^i]P_{\text{A}} + k_{\text{B}_2}^i[\text{B}_2^i]P_{\text{A}}}. \quad (\text{A.3})$$

Expanding the denominator we can write

$$\frac{\text{AR}^{60}}{\text{AR}^i} = \frac{k_{\text{B}_1}^{60}[\text{B}_1^{60}]P_{\text{A}} + k_{\text{B}_2}^{60}[\text{B}_2^{60}]P_{\text{A}}}{\alpha_{\text{xtl-B}_1}^{\text{f},i} k_{\text{B}_1}^{60}[\text{B}_1^{60}]r_{\text{B}_1}^{\text{eq},i}P_{\text{A}} + \alpha_{\text{xtl-B}_2}^{\text{f},i} k_{\text{B}_2}^{60}[\text{B}_2^{60}]r_{\text{B}_2}^{\text{eq},i}P_{\text{A}}}. \quad (\text{A.4})$$

From the Wolthers (2012) model, we use the following relationships:

$$k_{\text{B}_1}^{60} = k_{\text{B}_2}^{60} \quad (\text{A.5})$$

and

$$[\text{B}_2^{60}] = \phi [\text{B}_1^{60}], \quad (\text{A.6})$$

which leads to

$$\frac{\text{AR}^{60}}{\text{AR}^i} = \frac{(1 + \phi)}{\alpha_{\text{xtl-B}_1}^{\text{f},i} r_{\text{B}_1}^{\text{eq},i} + \phi \alpha_{\text{xtl-B}_2}^{\text{f},i} r_{\text{B}_2}^{\text{eq},i}}. \quad (\text{A.7})$$

Table A.1: Values used for the example calculation of the stochastic ratio from the ion-by-ion model using Eq. 36. The AR^{60} , DR^{60} , and ϕ parameters correspond to $T = 25^\circ\text{C}$, $\text{pH} = 8.3$ and $\Omega = 7.4$. For these calculations, we use $r_w = 0.0020672$ and $r_{\text{DIC}} = 0.0112372$, which represent the oxygen and carbon isotopic composition of Pee Dee Belemnite.

Ion-by-ion model		Canonical approach		
Parameter	Value	Isotope/isotopologue	Mass	Mole fraction
$r_{B_1}^{61}$	0.011216144632026	C_{12}	12	0.988901710623719
$r_{B_1}^{62}$	0.002122496022646	C_{13}	13	0.011098289376281
AR^{60}	14.031689198469467	C_{16}	16	0.997878246772305
DR^{60}	1.904996407891161	C_{18}	18	0.002121753227695
$\alpha_{\text{xtl}-B_1}^{\text{eq},C}$	1.004469000100333	$C_{2666} = C_{12}C_{16}C_{16}C_{16}$	60	0.982620440610989
$\alpha_{\text{xtl}-B_1}^{\text{eq},O}$	1.003354613133830	$C_{2668} = C_{12}C_{16}C_{16}C_{18} \cdot 3$	62	0.006267933282070
$\alpha_{\text{xtl}-B_1}^{\text{f},C}$	1.0000	$C_{2688} = C_{12}C_{16}C_{18}C_{18} \cdot 3$	64	$1.332728488192658 \times 10^{-5}$
$\alpha_{\text{xtl}-B_2}^{\text{f},C}$	0.9980	$C_{2888} = C_{12}C_{18}C_{18}C_{18}$	66	$9.445778181893845 \times 10^{-9}$
$\alpha_{\text{xtl}-B_1}^{\text{f},O}$	0.9980	$C_{3666} = C_{13}C_{16}C_{16}C_{16}$	61	0.011027795664416
$\alpha_{\text{xtl}-B_2}^{\text{f},O}$	0.9964	$C_{3668} = C_{13}C_{16}C_{16}C_{18} \cdot 3$	63	$7.034403582107305 \times 10^{-5}$
$\alpha_{B_2-B_1}^{\text{eq},C}$	1.001995685541594	$C_{3688} = C_{13}C_{16}C_{18}C_{18} \cdot 3$	65	$1.495700357586258 \times 10^{-7}$
$\alpha_{B_2-B_1}^{\text{eq},O}$	1.005179546014960	$C_{3888} = C_{13}C_{18}C_{18}C_{18}$	67	$1.060084925737459 \times 10^{-10}$
ϕ	$1.070823878831513 \times 10^2$	-	-	-
$r_{\text{xtl}*}^{63}$	$7.158820732177469 \times 10^{-5}$	$r_{\text{xtl}*}^{63}$		$7.158820732177469 \times 10^{-5}$

A.2 Example calculation of the stochastic ratio

A.2.1 Process-based stochastic ratio

For simplicity we continue to assume that there is no ^{17}O . From Eq. 36 we can compute the stochastic ratio using parameter values provided in Table A.1 for $T = 25^\circ\text{C}$, $\text{pH} = 8.3$ and $\Omega = 7.4$. This yields $r_{\text{xtl}*}^{63} = 7.158820732177469 \times 10^{-5}$ (first two columns of Table A.1).

A.2.2 Non-process-based stochastic ratio

With knowledge of the carbon and oxygen isotope composition of the crystal one can calculate the molar abundances (denoted with the symbol C) of isotopes and isotopologues (last three columns of Table A.1). For the parameters listed ($T = 25^\circ\text{C}$, $\text{pH} = 8.3$ and $\Omega = 7.4$), the isotopic composition of non-equilibrium calcite is $^{13}\text{C}/^{12}\text{C} = 0.011222843743774$ and $^{18}\text{O}/^{16}\text{O} = 0.006378793909653$. Using this as an input and following the canonical approach for calculating the stochastic distribution, we get $r_{\text{xtl}*}^{63} = 7.158820732177469 \times 10^{-5}$, in perfect agreement with the value calculation from Eq. 36. This example calculation is intended to instill confidence in the expressions derived herein for the clumped isotope composition of calcite.

References

- Affek, H., 2013. Clumped isotopic equilibrium and the rate of isotope exchange between CO_2 and water. *American Journal of Science* 313, 309–325.
- Affek, H. P., 2012. Clumped isotope paleothermometry: Principles, applications, and challenges. *Reconstructing Earth's Deep-Time Climate - The State of the Art in*.
- Affek, H. P., Bar-Matthews, M., Ayalon, A., Matthews, A., Eiler, J. M., 2008. Glacial/interglacial temperature variations in Soreq cave speleothems as recorded by 'clumped isotope' thermometry. *Geochimica et Cosmochimica Acta* 72 (22), 5351–5360.
- Beck, W., Grossman, E., Morse, J., 2005. Experimental studies of oxygen isotope fractionation in the carbonic acid system at 15, 25, and 40°C. *Geochimica et Cosmochimica Acta* 69 (14), 3493–3503.
- Bottinga, Y., 1968. Calculation of fractionation factors for carbon and oxygen isotopic exchange in the system calcite-carbon dioxide-water. *The Journal of Physical Chemistry* 72 (3), 800–808.
- Came, R. E., Eiler, J. M., Veizer, J., Azmy, K., Brand, U., Weidman, C. R., 2007. Coupling of surface temperatures and atmospheric CO_2 concentrations during the Palaeozoic era. *Nature* 449 (7159), 198–201.
- Clog, M., Stolper, D., Eiler, J. M., 2015. Kinetics of $\text{CO}_2(\text{g})\text{-H}_2\text{O}$ isotopic exchange, including mass 47 isotopologues. *Chemical Geology* 395, 1–10.
- Coplen, T., 2007. Calibration of the calcite–water oxygen-isotope geothermometer at Devils Hole, Nevada, a natural laboratory. *Geochimica et Cosmochimica Acta* 71 (16), 3948–3957.
- Daëron, M., Guo, W., Eiler, J., Genty, D., Blamart, D., Boch, R., Drysdale, R., Maire, R., Wainer, K., Zanchetta, G., 2011. $^{13}\text{C}\text{-}^{18}\text{O}$ clumping in speleothems: Observations from natural caves and precipitation experiments. *Geochimica et Cosmochimica Acta* 75 (12), 3303–3317.
- Defliese, W. F., Hren, M. T., Lohmann, K. C., 2015. Compositional and temperature effects of phosphoric acid fractionation on δ_{47} analysis and implications for discrepant calibrations. *Chemical Geology* 396, 51–60.
- Dennis, K., Schrag, D., 2010. Clumped isotope thermometry of carbonatites as an indicator of diagenetic alteration. *Geochimica et Cosmochimica Acta* 74 (14), 4110–4122.

559 Dennis, K. J., Affek, H. P., Passey, B. H., Schrag, D. P., Eiler, J. M., 2011. Defining an absolute
560 reference frame for ‘clumped’ isotope studies of CO₂. *Geochimica et Cosmochimica Acta* 75 (22),
561 7117–7131.

562 Dietzel, M., Tang, J., Leis, A., Köhler, S., 2009. Oxygen isotopic fractionation during inorganic
563 calcite precipitation: Effects of temperature, precipitation rate and pH. *Chemical Geology* 268 (1),
564 107–115.

565 Eagle, R. A., Tütken, T., Martin, T. S., Tripathi, A. K., Fricke, H. C., Connely, M., Cifelli, R. L.,
566 Eiler, J. M., 2011. Dinosaur body temperatures determined from isotopic (¹³C-¹⁸O) ordering in
567 fossil biominerals. *Science* 333 (6041), 443–445.

568 Eiler, J. M., Bergquist, B., Bourg, I., Cartigny, P., Farquhar, J., Gagnon, A., Guo, W., Halevy, I.,
569 Hofmann, A., Larson, T. E., et al., 2014. Frontiers of stable isotope geoscience. *Chemical Geology*
570 372, 119–143.

571 Eiler, J. M., Schauble, E., 2004. ¹⁸O¹³C¹⁶O in Earth’s atmosphere. *Geochimica et Cosmochimica*
572 *Acta* 68 (23), 4767–4777.

573 Ghosh, P., Adkins, J., Affek, H., Balta, B., Guo, W., Schauble, E., Schrag, D., Eiler, J., 2006a. ¹³C-
574 ¹⁸O bonds in carbonate minerals: A new kind of paleothermometer. *Geochimica et Cosmochimica*
575 *Acta* 70 (6), 1439–1456.

576 Ghosh, P., Garzione, C. N., Eiler, J. M., 2006b. Rapid uplift of the Altiplano revealed through
577 ¹³C-¹⁸O bonds in paleosol carbonates. *Science* 311 (5760), 511–515.

578 Guo, W., 2009. Carbonate clumped isotope thermometry: application to carbonaceous chondrites
579 and effects of kinetic isotope fractionation. Ph.D. thesis, California Institute of Technology.

580 Guo, W., Mosenfelder, J. L., Goddard, W. A., Eiler, J. M., 2009. Isotopic fractionations associated
581 with phosphoric acid digestion of carbonate minerals: insights from first-principles theoretical
582 modeling and clumped isotope measurements. *Geochimica et Cosmochimica Acta* 73 (24), 7203–
583 7225.

584 Henkes, G. A., Passey, B. H., Wanamaker Jr, A. D., Grossman, E. L., Ambrose Jr, W. G., Carroll,
585 M. L., 2013. Carbonate clumped isotope compositions of modern marine mollusk and brachiopod
586 shells. *Geochimica et Cosmochimica Acta* 106, 307–325.

587 Hill, P. S., Tripathi, A. K., Schauble, E. A., 2014. Theoretical constraints on the effects of pH,
588 salinity, and temperature on clumped isotope signatures of dissolved inorganic carbon species and
589 precipitating carbonate minerals. *Geochimica et Cosmochimica Acta* 125, 610–652.

590 Kluge, T., Affek, H. P., Dublyansky, Y., Spötl, C., 2014. Devils Hole paleotemperatures and im-
591 plications for oxygen isotope equilibrium fractionation. *Earth and Planetary Science Letters* 400,
592 251–260.

593 Kluge, T., John, C. M., Jourdan, A.-L., Davis, S., Crawshaw, J., 2015. Laboratory calibration of the
594 calcium carbonate clumped isotope thermometer in the 25-250°C temperature range. *Geochimica*
595 *et Cosmochimica Acta* 157, 213–227.

596 Mook, W., 1986. ^{13}C in atmospheric CO_2 . *Netherlands Journal of Sea Research* 20 (2-3).

597 Mook, W., Bommerson, J., Staverman, W., 1974. Carbon isotope fractionation between dissolved
598 bicarbonate and gaseous carbon dioxide. *Earth and Planetary Science Letters* 22.

599 Romanek, C. S., Grossman, E. L., Morse, J. W., 1992. Carbon isotopic fractionation in synthetic
600 aragonite and calcite: Effects of temperature and precipitation rate. *Geochimica et Cosmochimica*
601 *Acta* 56 (1), 419–430.

602 Saenger, C., Affek, H. P., Felis, T., Thiagarajan, N., Lough, J. M., Holcomb, M., 2012. Carbonate
603 clumped isotope variability in shallow water corals: Temperature dependence and growth-related
604 vital effects. *Geochimica et Cosmochimica Acta* 99, 224–242.

605 Schauble, E. A., Ghosh, P., Eiler, J. M., 2006. Preferential formation of ^{13}C - ^{18}O bonds in carbonate
606 minerals, estimated using first-principles lattice dynamics. *Geochimica et Cosmochimica Acta*
607 70 (10), 2510–2529.

608 Spero, H. J., Bijma, J., Lea, D. W., Bemis, B. E., 1997. Effect of seawater carbonate concentration
609 on foraminiferal carbon and oxygen isotopes. *Nature* 390 (6659), 497–500.

610 Stolper, D., Lawson, M., Davis, C., Ferreira, A., Neto, E. S., Ellis, G., Lewan, M., Martini, A.,
611 Tang, Y., Schoell, M., et al., 2014. Formation temperatures of thermogenic and biogenic methane.
612 *Science* 344 (6191), 1500–1503.

613 Stolper, D., Martini, A., Clog, M., Douglas, P., Shusta, S., Valentine, D., Sessions, A., Eiler, J., 2015.
614 Distinguishing and understanding thermogenic and biogenic sources of methane using multiply
615 substituted isotopologues. *Geochimica et Cosmochimica Acta* 161, 219–247.

616 Tang, J., Dietzel, M., Fernandez, A., Tripathi, A., Rosenheim, B., 2014. Evaluation of kinetic ef-
617 fects on clumped isotope fractionation (Δ_{47}) during inorganic calcite precipitation. *Geochimica*
618 *et Cosmochimica Acta*.

619 Thiagarajan, N., Adkins, J., Eiler, J., 2011. Carbonate clumped isotope thermometry of deep-sea
620 corals and implications for vital effects. *Geochimica et Cosmochimica Acta* 75 (16), 4416–4425.

621 Tripathi, A. K., Eagle, R. A., Thiagarajan, N., Gagnon, A. C., Bauch, H., Halloran, P. R., Eiler,
622 J. M., 2010. ^{13}C - ^{18}O isotope signatures and ‘clumped isotope’ thermometry in foraminifera and
623 coccoliths. *Geochimica et Cosmochimica Acta* 74 (20), 5697–5717.

624 Tripathi, A. K., Hill, P. S., Eagle, R. A., Mosenfelder, J. L., Tang, J., Schauble, E. A., Eiler, J. M.,
625 Zeebe, R. E., Uchikawa, J., Coplen, T. B., et al., 2015. Beyond temperature: Clumped isotope sig-
626 natures in dissolved inorganic carbon species and the influence of solution chemistry on carbonate
627 mineral composition. *Geochimica et Cosmochimica Acta* 166, 344–371.

628 Turner, J. V., 1982. Kinetic fractionation of carbon-13 during calcium carbonate precipitation.
629 *Geochimica et Cosmochimica Acta* 46 (7), 1183–1191.

630 Usdowski, E., Hoefs, J., 1993. Oxygen isotope exchange between carbonic acid, bicarbonate, car-
631 bonate, and water: a re-examination of the data of Mccrea (1950) and an expression for the
632 overall partitioning of oxygen isotopes between the carbonate species and water. *Geochimica et*
633 *Cosmochimica Acta* 57 (15), 3815–3818.

634 Vogel, J., Grootes, P., Mook, W., 1970. Isotopic fractionation between gaseous and dissolved carbon
635 dioxide. *Zeitschrift für Physik* 230.

636 Wacker, U., Fiebig, J., Schoene, B. R., 2013. Clumped isotope analysis of carbonates: comparison
637 of two different acid digestion techniques. *Rapid Communications in Mass Spectrometry* 27 (14),
638 1631–1642.

- 639 Wang, D. T., Gruen, D. S., Lollar, B. S., Hinrichs, K.-U., Stewart, L. C., Holden, J. F., Hristov,
640 A. N., Pohlman, J. W., Morrill, P. L., Könneke, M., et al., 2015. Nonequilibrium clumped isotope
641 signals in microbial methane. *Science* 348 (6233), 428–431.
- 642 Wang, Z., Gaetani, G., Liu, C., Cohen, A., 2013. Oxygen isotope fractionation between aragonite
643 and seawater: Developing a novel kinetic oxygen isotope fractionation model. *Geochimica et*
644 *Cosmochimica Acta* 117, 232–251.
- 645 Wang, Z., Schauble, E. A., Eiler, J. M., 2004. Equilibrium thermodynamics of multiply substituted
646 isotopologues of molecular gases. *Geochimica et Cosmochimica Acta* 68 (23), 4779–4797.
- 647 Watkins, J. M., Hunt, J. D., Ryerson, F. J., DePaolo, D. J., 2014. The influence of temperature,
648 pH, and growth rate on the $\delta^{18}\text{O}$ composition of inorganically precipitated calcite. *Earth and*
649 *Planetary Science Letters* 404, 332–343.
- 650 Watkins, J. M., Nielsen, L. C., Ryerson, F. J., DePaolo, D. J., 2013. The influence of kinetics on
651 the oxygen isotope composition of calcium carbonate. *Earth and Planetary Science Letters* 375,
652 349–360.
- 653 Wolthers, M., Nehrke, G., Gustafsson, J., Van Cappellen, P., 2012. Calcite growth kinetics: Modeling
654 the effect of solution stoichiometry. *Geochimica et Cosmochimica Acta* 77, 121–134.
- 655 Yeung, L. Y., Ash, J. L., Young, E. D., 2015. Biological signatures in clumped isotopes of O_2 . *Science*
656 348 (6233), 431–434.
- 657 Zaarur, S., Affek, H. P., Brandon, M. T., 2013. A revised calibration of the clumped isotope ther-
658 mometer. *Earth and Planetary Science Letters* 382, 47–57.
- 659 Zeebe, R., 2007. An expression for the overall oxygen isotope fractionation between the sum of
660 dissolved inorganic carbon and water. *Geochem. Geophys. Geosyst.* 8 (9), Q09002.
- 661 Zeebe, R., Wolf-Gladrow, D., 2001. CO_2 in seawater: Equilibrium, kinetics, isotopes. Vol. 65. Elsevier
662 Science Limited.