Comparing the impacts of Miocene-Pliocene changes in inter-ocean gateways on climate: Central American Seaway, Bering Strait, and Indonesia.

Chris M. Brierley^a and Alexey V. Fedorov^b

^aDepartment of Geography, University College London, London, WC1E 6BT, United Kingdom,

^bDepartment of Geology and Geophysics, Yale University, New Haven, Connecticut, 06511, United States

Corresponding Author: Chris Brierley c.brierley@ucl.ac.uk

Abstract

Changes in inter-ocean gateways caused by tectonic processes have been long considered an important factor in climate evolution on geological timescales. Three major gateway changes that occurred during the Late Miocene and Pliocene epochs are the closing of the Central American seaway (CAS) by the uplift of the Isthmus of Panama, the opening of the Bering Strait, and the closing of a deep channel between New Guinea and the Equator. This study compares the global climatic effects of these changes within

the same climate model framework. We find that the closure of the CAS and the opening of the Bering Strait induce the strongest effects on the Atlantic meridional overturning circulation (AMOC). However, these effects potentially compensate, as the closure of the CAS and the opening of the Bering Strait cause similar AMOC changes of around 2 Sv (strengthening and weakening respectively). Previous simulations with an open CAS consistently simulated colder oceanic conditions in the Northern hemisphere - contrasting with the evidence for warmer sea surface temperatures 10-3 million years ago. Here we argue that this cooling is overestimated because (a) the models typically simulated too strong an AMOC change not yet in equilibrium, (b) used a channel too deep and (c) lacked the compensating effect of the closed Bering Strait - a factor frequently ignored despite its potential influence on northern high latitudes and ice-sheet growth. Further, we discuss how these gateway changes affect various climatic variables from surface temperature and precipitation to ENSO characteristics.

Keywords: gateways, Bering, Panama, onset glaciation, palaeoclimate

1. Introduction

The ultimate driving forces behind the global climate cooling from the late Miocene through the mid-Pliocene and culminating in the onset of modern glacial cycles remain enigmatic. The general consensus is that atmospheric CO₂ concentration was a major factor (DeConto et al., 2008; Lunt et al., 2008). Yet uncertainties in its values (Fedorov et al. 2013) and examples of divergent trends in CO₂ and temperature (LaRiviere et al. 2012) necessitate considering additional factors, such as the effects of tectonic changes on the climate evolution. The climate system is especially sensitive to tectonic changes of its inter-ocean flows (gateways). Numerical modelling of the role of gateways has been performed for over two decades (Hirst and Godfrey, 1993; Maier Reimer et al., 1990).

The closure of the Central American Seaway (CAS) that linked the tropical Atlantic to the Pacific has been the predominant focus of research on Plio-Pleistocene gateway changes, as it was suggested that the closure might have acted as a trigger for the onset of Northern Hemisphere glaciation (Haug and Tiedemann, 1998). The opening of the Bering Strait created a high latitude connection between the Pacific and the Arctic. This has previously thought to have occurred prior to 4.8 million years ago (Ma) (Marincovich and Gladenkov, 1999), yet recent boundary conditions provided for the Pliocene Model Intercomparison Project still have it closed at ~3 Ma (Haywood et al., 2014). This work builds on that of Fedorov *et al.* (2013), where several proposed explanations for the Early Pliocene's weakened temperature gradient in the tropical Pacific were compared. These included two proposed gateway changes: the closing of the Central American Seaway and alterations of the Indonesian passages, to which we add the opening of the Bering Strait.

Previous studies that have looked at the impact of multiple ocean gateways have included the Southern Ocean (e.g. Mikolajewicz et al., 1993), which was closed long before the Pliocene. As far as we know, no one has previously compared the impacts of opening the Bering Strait, closing the Central American Seaway and altering the Indonesian passages within the same model framework. As such, we will first briefly review each separately below. The model setup will then be presented, along with our altered boundary conditions. The climate impacts will be investigated; first locally; then globally and finally we will explore the changes in the El Niño Southern Oscillation. The conclusions of the intercomparison will then be summarised and its implications for Plio-Pleistocene climate evolution discussed.

1.1 Central American Seaway

The established view of the closure of the Central American Seaway and creation of the Isthmus of Panama is of a slow process taking many millions of years. The first step was the creation of a volcanic arc around 17 Ma leading to the creation of an archipelago by 12 Ma (Coates et al., 1992). The critical condition from an oceanographic perspective is the extent of constriction of deep and shallow water flow. The deep water connection was already cut by the Pliocene. The upper-ocean flow through the CAS curtailed between 4.7 and 4.2 Ma; as evidenced by the developing contrast in ocean surface $\delta^{18}\text{O}$ values between the Caribbean Sea and the Pacific (Haug et al. 2001). Finally, the shallow link is commonly thought to have been severed around 3.5Ma (Coates et al., 1992). The similarity of this date to that of the onset of Northern Hemisphere Glaciation has led to much discussion of the closing of the seaway as preconditioning the glaciation (Haug and Tiedemann, 1998). However, debate continues on the timing of closure (Molnar, 2008, provides a comprehensive review of the problems of determining this timing). Recent suggestions of a much earlier closure in the middle Miocene (e.g. Montes et al., 2015) have further complicated matters.

93

94 Numerical modeling experiments looking at the role of the Isthmus of Panama on
95 the global circulations have been performed by many authors over the past two decades
96 (many compiled by Zhang et al., 2012). From the outset, it was recognized that the
97 Atlantic meridional overturning circulation (AMOC) is weaker with an open seaway
98 (Maier Reimer et al., 1990). How much weaker depends on both the details of the
99 seaway changes and the climate model used (Zhang et al., 2012) as these factors affect
100 the salinity contrast between the North Pacific and Atlantic, which in turn influences the
101 strength of the AMOC. This salinity contrast is maintained largely by atmospheric
102 freshwater transport from the Caribbean into the Eastern Pacific. However, the Central
103 American Seaway provides an oceanic counterbalance to this freshwater flux,
104 weakening the salinity difference and hence the AMOC. Mestas Nuñez and Molnar
105 (2014) note this salinity contrast can also be influenced by other climate changes, such
106 as long-term trends in Pacific sea surface temperatures (SSTs; Fedorov et al. 2013), so
107 salinity changes may not relate solely to tectonic movement.

108

109 Haug and Tiedemann (1998) hypothesize that a strong AMOC (caused by a recent
110 closure of the CAS) was a primer for the onset of Northern Hemisphere glaciation. The
111 consequences of a CAS closure at ~3.5 Ma would be a coeval warming of the North
112 Atlantic. The warmer Atlantic SSTs would have led to increases in precipitation and
113 presumably ice accumulation (Haug and Tiedemann, 1998). However, this idea
114 contradicts more recent paleoclimate reconstructions (Lawrence et al., 2010) that
115 suggest gradual cooling in northern high latitudes over the same time period. It is
116 indicative that this cooling had a similar magnitude in northern and southern high
117 latitudes (Fedorov et al. 2013), whereas an AMOC change would typically produce a
118 seesaw SST anomaly about the equator. Furthermore, climate model experiments with
119 interactive continental-ice sheets suggest the increased precipitation does not lead to

greater ice cover over Greenland (Lunt et al., 2008), implying the role of CAS closure for Northern Hemisphere glaciation may be overstated.

Indications of an earlier closure of the CAS (~4.4 Ma) have led to suggestions that it caused a shoaling of the tropical thermocline (Steph et al., 2010), which is tentatively supported by model simulations (Zhang et al., 2012). However, for realistic depths of the CAS (a few hundred meters or less) this effect is moderate, shows both thermocline deepening and shoaling along the equator, and has only weak manifestation in SST (Zhang et al. 2012, Fedorov et al. 2013, and Section 3 of the present study).

1.2 Bering Strait

The Pacific ocean was connected to the Arctic through the Bering Strait in the late Miocene or early Pliocene (Marincovich and Gladenkov, 1999). The Bering Strait is shallow (about 50 m) yet plays an interesting role in the Arctic Ocean circulation as a conduit for fresh water (Woodgate, 2005).

The timing of the opening of the Bering Strait is not well known. Marincovich and Gladenkov (1999) find that the Bering Strait was permanently closed prior to 4.8 Ma from biogeographic evidence. Ocean-only experiments also suggest the closure of the CAS reverses the flow in the Bering Strait (Maier Reimer et al., 1990). The Pliocene Model Intercomparison Project (PlioMIP) provides global land-sea mask reconstructions for the period 3.2-3.0 Ma. The Bering Strait was considered open in the first set of PlioMIP experiments (Dowsett et al., 2012), but is closed in the recent reconstruction (Haywood et al., 2014). As the Bering Strait is/was so shallow, the timing and nature of its opening is also contingent on global sea level changes (Hu et al., 2010). The uncertainty in Pliocene sea levels have a significant impact in this instance, with the observational error of ± 10 m probably being overly optimistic (Dutton et al., 2015).

146

147 Numerical and theoretical studies have shown a role for the Bering Strait in
148 controlling the strength and stability of the AMOC (Shaffer, 1994; Wadley and Bigg,
149 2002; De Boer and Nof, 2004). This occurs as the Bering Strait helps determine the
150 salinity of the Arctic and hence the North Atlantic, by regulating the flow of relatively
151 fresh water from the North Pacific. Simulations by Hu et al. (2010) looked at the impact
152 of rapid changes in the Bering Strait controlled by sea level changes during a glacial
153 cycle. They posited a feedback involving the Laurentide ice sheet, the AMOC and the
154 Bering Strait – suggesting a role in glacial climate variability. Here we are thinking more
155 about its impacts in the Late-Miocene/Pliocene and so do not use a glacial baseline for
156 our experiments.

157 1.3 Indonesian Throughflow

158 Since detaching from Antarctica in the Early Eocene, the continent of Australasia has
159 been moving slowly northwards towards the Equator (Hall, 2002). The most northerly
160 tip of Australasia, namely the Bird's Head of Papua New Guinea, is now within 1° of the
161 Equator. It is so close to the island of Halamahera to the north of the Equator that there
162 are no channels through which deep water may pass between them. It is assumed that
163 this deep channel closed during the Pliocene (Hall, 2002); however more precise, direct
164 dates are not available.

165

166 One effect of this deep channel closing may have been to shift the source of the
167 Indonesian Throughflow from Southern to Northern Pacific subtropical waters (Rodgers
168 et al., 2000). Potential traces of such behavior has been observed in the paleoclimate
169 record (Karas et al., 2009), who find a cooling and freshening of the subsurface waters
170 entering the Indian ocean from the Pacific after 3.5 Ma. It has been suggested that such a
171 change could have had global climate consequences, including the aridification of East

Africa during the Pliocene (Cane and Molnar, 2001), but subsequent model simulations did not support that idea (Jochum et al., 2009; Krebs et al., 2011).

Krebs et al. (2011) found that Indonesian changes could explain some of the large ecological changes observed in Australia during the Plio-Pleistocene. Jochum et al. (2009) found little global climate impact with both coupled and ocean-only mode of CCSM3 (an earlier version of the model used here). They saw a small warming of the Central Equatorial Pacific and alterations in the statistical properties of ENSO. Modeling results of Fedorov et al. (2013) also did not support Indonesian constriction as an explanation for the changes in patterns of tropical Pacific SSTs they had diagnosed. Here, we expand upon that analysis.

Changes in the pathways of the Indonesian Throughflow are not the only method by which changes in topography in this region may affect global climate. Recent work on dynamic topography (Rowley et al., 2013) allows for the exposure of shallow shelves in the region due to mantle convection (Haywood et al., 2014). Exposure of the Sunda and Sahul shelves at the Last Glacial Maximum are thought to have affected the atmospheric Walker circulation (DiNezio et al., 2011). Likewise, Molnar and Cronin (2015) suggest a gradual increase in the exposed landmass of the Maritime Continent since 5 Ma played a role both in the CO₂ drop seen during the Plio-Pleistocene and the evolution of the Walker circulation. Brierley and Fedorov (2011) show changes in tidal mixing in the Banda Sea (arising from changes in bathymetric roughness) could be sufficient to drive changes in throughflow properties, yet hard to constrain from the geologic record.

2. Method

2.1 Community Earth System Model

All the simulations presented in this comparison study use the Community Earth System Model (CESM; Gent et al., 2011). This is the most recent generation of the coupled general circulation model developed by the National Center for Atmospheric Research (NCAR). This model involves fully dynamical atmosphere and ocean components with representations of the land surface and sea ice. These simulations are performed with the (relatively) low resolution version developed for paleoclimate studies (Shields et al., 2012). The atmosphere and land models have a horizontal resolution of T31 ($3.75^\circ \times 3.75^\circ$) with 26 atmospheric vertical levels. The land model involves biochemistry and semi-dynamical vegetation that grows and dies-back with the seasons, however the land-cover proportions are prescribed throughout the simulations. The ocean has a rotated grid with a pole under Greenland. It has a nominal resolution of 3° and 60 vertical levels, along with suitable parameter settings (Shields et al., 2012). More precisely, we use CESM version 1.0.2 at T31_gx3 resolution with component setting B1850_CN.

The treatment of the sub-grid scale mixing has become significantly more sophisticated than previous model generations, which relied predominantly on the Gent-McWilliams parameterization (Gent et al., 2011). Of relevance to this study are parameterizations of Nordic overflows (Danabasoglu et al., 2010), sub-mesoscale mixed layer eddies (Fox-Kemper et al, 2010) and abyssal tidal mixing (Jayne, 2009). The overflow parameterization improves the representation of dense water crossing a sill and entraining water as it sinks. It would only be relevant for the sill in the newly-created Central American Seaway, yet the density gradient here is not sufficient to necessitate its implementation.

The simulation used as a control is a 500 year extension from the preindustrial run described in Shields et al. (2012). The simulation with an altered New Guinea,

“Indonesia”, is also 500 years long starting from the same initial conditions. Assessment of the degree of equilibration was made through inspection of the global average top of the atmosphere heat flux imbalance, and surface and deep ocean temperature trends (Supplementary Figure 1). The simulations involving the closing of the Bering Strait, “Bering”, and the opening of the Central American Seaway, “Panama”, showed trends in the deep ocean and so were integrated for 1500 and 2400 years respectively. Even after such long times neither simulation is fully equilibrated throughout the ocean, yet the trends in the AMOC are hard to distinguish from internal variability at this point (sect. 4.5). The results shown here are the difference in the final 200 years of each simulation compared to the Control.

2.2 Application of boundary condition changes

All of the simulations described here are sensitivity studies: meaning that the gateway change is the only imposed difference from the control simulation. To alter the Central American Seaway and the Indonesian Archipelago, we have converted land grid points into ocean. This requires some assumptions to be made about the ocean bathymetry (Figure 1). The new ocean grid points in Indonesia were set at the average depth of the neighboring locations, leading to a depth of 500m. For the open Central American Seaway, a value of 150m was chosen for the sensitivity test (Fig. 1). This sill depth is shallower than some of the simulations in the multi-model comparison of Zhang et al. (2012), but still three times the depth of the present Bering Strait. The suggestion by De Schepper et al. (2013) that the CAS was temporally closed by the sea level fall during MIS M2 glaciation at ~3.3 Ma implies a sill depth of at most 65 m.

Previous work has suggested that changes in tidal mixing may have been important in Indonesia during the Pliocene (Brierley and Fedorov, 2011). Therefore, rather than turning off the abyssal tidal mixing (as proposed by Paleoclimate Working Group, 2015, for palaeoclimate simulations), the prescribed energy flux field was interpolated over

new ocean locations, although this is probably second-order to the bathymetric changes. Appropriate river routings and other ancillary files were created using the altered bathymetry (Paleoclimate Working Group, 2015). The potential land dataset used to create CESM boundary conditions at finer resolutions has sufficient island data that realistic preindustrial values were prescribed for the new land points after the closing of the Bering Strait (Fig. 1). This prevented arbitrary choices of the new land cover, and results in roughly 40% deciduous broadleaf boreal shrub, 30% bare ground and 30% Arctic C3 grass.

2.3 Statistical Significance

The importance of any changes observed in this article is assessed by comparing to the model's internal variability. Reliably estimating the model's multi-centennial internal variability is problematic and therefore we adopt a conservative approach and use the control run to estimate the internal variability on a shorter timescale. The 500 year long preindustrial run has been subdivided into twenty different segments, each 25 years long. Differences are considered statistically significant if they fall outside the two-tailed 95% confidence range assuming the segments are independent samples of a normally distributed noise caused by internal variability. In the following figures, stippling indicates significant anomalies (in spatial maps) whilst diagonal hatching indicates non-significant anomalies (in the plots of overturning streamfunction).

3. Local impacts

Unsurprisingly, altering each gateway leads to changes in the local oceanographic conditions that are significantly different from the model's internal variability. We present the transports of mass, heat and salt (Table 1) along with velocities of the upper 150 m of the ocean near the gateways (Figures 2 & 3).

3.1 The Central American Seaway

Opening the Central American Seaway leads to a net flow from the Pacific to Atlantic with a volume of 3.7 Sv (Table 1). This flow is at the lower end of the range in Zhang et al. (2012), but is to be expected as its cross-sectional area is substantially less than those with deeper sill-depths. The flow is achieved by a northward current flowing up the Colombia's Pacific coast partly balanced by a westward flow through what is presently Costa Rica (Fig. 2e). In fact, there is some recirculation occurring, as water flowing along the Colombian coast is entrained into the larger Atlantic western boundary current within the Caribbean. In the Pacific, there is an extension of the southward current flowing along the Mexican coast. When this newly strengthened current reaches the mouth of the Central American Seaway, it in turn leads to greater recirculation.

3.2 The Bering Strait

The Bering Strait is presently open, but only reaches depths of up to 40m in the model (Fig. 1) and so does not support as large a flow as the other gateways. Closing the Bering Strait deprives the Arctic Ocean of this inflow, which is a source of freshwater because of the low salinity of the northern Pacific (Woodgate, 2005). The salt transport is provided in Table 1, yet the virtual freshwater transport is perhaps more enlightening. The average density of water flowing through the Bering Strait is 1026.5 kg/m³ meaning that the salt transport is equivalent to a mass transport of 0.869 Sv of 34.8 psu sea water. The Bering Strait in CESM therefore transports approximately 0.037 Sv of freshwater, about half the observational estimate (Woodgate, 2005). Surprisingly the strong changes in upper-ocean flow (Fig. 2c) do not extend past the Aleutian Islands into the North Pacific. There are significant changes in the Arctic Ocean that are more widely felt, primarily downstream of the Strait in the Beaufort Gyre.

Recent palaeoceanographic work has found evidence of a progressive Pliocene cooling south of the Strait (Horikawa et al., 2015). This cooling has been interpreted as a consequence of a flow reversal in the Strait caused by the closing of the Central American Seaway (Maier Reimer et al., 1990). There is a Northward flow of both heat and salt in all the simulations (Tab. 1) suggesting an alternate paleoclimatic interpretation is required – perhaps revolving around changes to the Bering Strait itself.

3.3 Indonesian Throughflow

Observational estimates of the Indonesian Throughflow are 15 Sv emerging into the Indian Ocean, with 13 Sv entering the region directly from the Pacific Ocean (Gordon et al., 2010). The model underestimates the net outflow slightly (13.8 Sv, Tab. 1), but considering its coarser resolution in relation to other recent studies (Jochum et al., 2009; Krebs et al., 2011), it is reasonable.

In the simulation without the northern extension of New Guinea, there is a significant reduction in throughflow (Tab. 1). There is a strong increase in flow entering the Makassar Strait/Molucca Sea (Fig 3b; the model does not resolve Sulawesi so they form a single entity). However, this is negated by an even greater increase in the flow around the northern edge of New Guinea into the Pacific. There is a weakened outflow resulting from this, although the flow through the Tasman Sea has been somewhat strengthened.

Interestingly, changes in all three gateways investigated have significant impacts on the Indonesian Throughflow (Tab. 1). The closing of Bering Strait increases the mass transport by just over 1.5 Sv. The opening of the Central American Seaway causes the throughflow to reduce by 35%. Both of these changes are consistent in sign and relative magnitude with changes to the AMOC (section 4.5) driving a global ocean conveyor.

4. Global climate impacts

The majority of the discussion of gateways in the paleoceanographic literature centres on the role they may play in long-term climate evolution. This assumes that the gateways have remote impacts, leading to either global changes (Haug and Tiedemann, 1998) or at least changes elsewhere (Cane and Molnar, 2001). In this section, we compare several global diagnostics between the simulations. We present these diagnostics in the conventional sense for sensitivity studies (i.e. *perturbed – control*). This is opposite to the chronological sense of the changes, which would be *control – perturbed* (i.e. *after – before*).

4.1 Poleward heat transports

The opening of the Central American Seaway causes the largest alterations in the poleward heat budgets (Fig. 4). It reduces northward ocean heat transports at most latitudes and increases the atmospheric transports at all latitudes. The two changes do not compensate completely leaving a net alteration in the Tropics (Fig. 4b). The anomalous northward atmosphere heat transports is associated with a southward movement of the inter-tropical convergence zone (ITCZ) and reduced temperature gradient between the Northern and Southern hemispheres. This southward ITCZ shift leads to a reduced dominance of the Southern Hadley cell (not shown). The relative strengthening of the Northern Hadley Cell allows it to transport more heat northwards. The reduced ocean heat transport is associated with both AMOC changes (sect 4.2) and changes in the subtropical cells.

Shifting Indonesia causes a marginally significant increase in the atmospheric heat transport near the Equator. The closing of the Bering Strait causes no significant changes in the combined heat transport (Fig. 4b). There is an increase in the ocean heat transport with a maximum at 35 °N, likely associated with changes in the AMOC.

351

352 4.2 Meridional overturning circulation

353 The preindustrial control simulation shows a peak in the zonal mean meridional
354 overturning streamfunction in the Atlantic occurring at a depth of 1 km around 35 °N
355 (Figure 5b). The Antarctic bottom water cell is weak - a feature of all resolutions of
356 CESM (Danabasoglu et al., 2012). A pair of shallow subtropical overturning cells exists in
357 the upper 250 m, with upwelling occurring on the equator.

358

359 The temporal evolution of the maximum of the meridional overturning
360 streamfunction in the North Atlantic shows little evidence of secular trends beyond 100
361 years in either the control or Indonesian simulations (Fig. 5a). There are AMOC changes
362 in both the Bering Strait and Central American Seaway simulations emerging over the
363 first 500 years. These simulations were integrated for a further 1000 and 1900 years
364 respectively, allowing any recovery to occur. The recovery is most notable in the CAS
365 simulation (Fig. 5a) and is predominantly complete after 1500 years (though there is
366 still drift in the deep ocean, Supplementary Fig 1). The AMOC continues to exhibit a
367 strong centennial to multi-centennial internal variability (to be discussed elsewhere).
368 Only changes in the mean state are discussed here.

369

370 A closed Bering Strait causes an increase in the deep overturning cell throughout the
371 Atlantic (Fig. 5d). This reaches a maximum of 2.5 Sv and is statistically distinguishable
372 down to 4 km. An open Central American Seaway has the reverse response (Fig. 5e) and
373 leads to significant weakening and shoaling of the AMOC (as observed by previous
374 authors e.g. Maier Reimer et al., 1990; Zhang et al., 2012). The weakening seen here is
375 among the lowest of the results compiled by Zhang et al. (2012), but it also has one of
376 the shallowest sill depths of their simulations. Interestingly with this shallower sill

depth (more appropriate for the Pliocene) the impact of the CAS on the AMOC has a very similar magnitude as that of the Bering Strait. This was not foreseen in the first 500 years of simulation. The recovery means that if the AMOC response to the opening of the CAS was taken after, say, only 500 years (as in Lunt et al., 2007) it would be over-estimated by factor of two (Fig. 5a). Alterations in the Indonesia bathymetry have no significant impacts on the AMOC (Fig. 5c). In general, the changes in AMOC are accompanied by changes in the North-South surface salinity gradient in the Atlantic of roughly 1 psu (see sect 4.5).

4.3 Surface Temperature Changes

All three simulations show statistically significant changes in surface air temperature at different regions (Fig. 6). These surface air temperature changes are very similar to the underlying SST changes, so the SSTs are not shown here.

In the Indonesia simulation, there are no significant temperature changes directly overlying the boundary alterations (Fig. 6a). There is a cooling over the Yellow Sea and a warming of the Eastern Equatorial Pacific. It is not clear why the Arctic warms just north of the Bering Strait, especially given the lack of significant changes in the flow through the Bering Strait itself (Tab. 1). We suspect this may be part of internal variability in the model or atmospheric teleconnections.

A closed Bering Strait leads to a local cooling, yet the largest region of significant changes is a warming in the North Atlantic (Fig 6b). The impacts of closing the Bering Strait on Pacific surface temperatures appear marginal. The warming in the Southern Ocean is counter to a bipolar seesaw response to a strengthened AMOC, and is instead collocated with weakened surface wind stresses (not shown).

403

404 The opening of the Central American Seaway demonstrably had a global impact (Fig.
405 6c). It alters the inter-hemispheric temperature gradient, by cooling the Northern
406 Hemisphere and warming the Southern. Interestingly the changes in the higher latitudes
407 of the North Atlantic (the sinking region of the AMOC) are not statistically significant
408 according to the metric used here (sect. 2.3). The lack of statistical significance may be
409 influenced by the sea ice edge of the pre-industrial control simulation, which has a
410 southward bias into the region (Shields et al., 2012). Intriguingly, a strong cooling
411 occurs in the Subarctic North Pacific. While it is insufficient to activate a Pacific MOC, it
412 may indicate a tendency towards a modified version of the ocean global conveyor that is
413 known to be impacted by tropical gateways (von der Heydt and Dijkstra, 2008).

414 4.4 Precipitation

415 The changes in annual average precipitation caused by the gateway changes are
416 generally less than those for temperature (Fig. 6). In general, changes in either the
417 Bering Strait or Indonesia do not show systematic changes in precipitation of local or
418 global extent. Specifically, the simulation with alterations to Indonesia (Fig. 6d) does not
419 show either the rainfall changes over Africa hypothesized by Cane and Molnar (2001) or
420 over Australia as modeled by Krebs et al. (2011). The Indonesian simulation does show
421 increased precipitation over the Eastern Equatorial Pacific, probably associated with its
422 altered ENSO properties (see section 4.7). The warming in the North Atlantic caused by
423 the closing of the Bering Strait is also associated with increased precipitation there (Fig.
424 6e).

425

426 Opening the Central American Seaway causes significant regional precipitation
427 changes in the Eastern Tropical Pacific and Tropical Atlantic (fig. 6f). These show the
428 southward shift in the ITCZ associated with the reduction in the inter-hemispheric

temperature gradient discussed above. The reduction in rainfall on either side of the Isthmus of Panama shows a further alteration of the inter-ocean freshwater flux. Not only is there a freshwater flux transport through the opened seaway from the Pacific to the Atlantic (Tab. 1), there is also a weakened freshwater transport going the other direction in the atmosphere.

4.5 Surface salinity

The changes in salt transport through the ocean gateways (Tab. 1) and their impact on the freshwater budget of the ocean would be expected to alter the ocean's surface salinity structure (Fig. 7). The impact of altering the Indonesian gateway leads to changes in the tropical Pacific salinities (Fig. 7a). These changes are not collocated with the temperature (Fig. 6a) and precipitation changes (Fig. 6d), but are presumably related. Preventing the flow of relatively fresher water in the Arctic Ocean by closing the Bering Strait, leads to the Arctic becoming significantly saltier (Fig 7b). This signal is able to propagate into the North Atlantic, causing the Labrador Sea to become 0.5 psu saltier. The rest of the ocean becomes fresher as a consequence, leading to an increased salinity gradient between the North and South Atlantic. The initially counter-intuitive freshening north of the Bering Strait is caused by weaker circulation in region (Fig. 2; Wadley and Bigg, 2002).

Globally the impact of opening the Central American Seaway makes the ocean surface significantly saltier. Intriguingly the salinity reduction in the North Atlantic is relatively weak (Fig. 7c), but joined with the saltier South Atlantic leads to a robust decrease in the salinity gradient consistent with the weakened AMOC (Fig. 5e). The increased surface salinity in the Arctic may arise from increased sea ice formation due to the colder temperatures.

4.6 Thermocline changes in the tropics

Steph et al. (2010) suggest that the closure of the Central American Seaway resulted in the end of the weak SSTs characteristic of the equatorial Pacific in the Pliocene (Fedorov et al., 2013). Zhang et al. (2012) found support for this hypothesis from a model compilation. The thermocline changes in the simulation here are qualitatively similar, but with a reduced magnitude as befits the shallower sill depth (Supplementary Figure 2). Minimal changes to the tropical thermocline are caused by the closure of the Bering Strait (despite its AMOC intensification). However, in all of these experiments surface temperature manifestation of these changes in the equatorial band is weak, on the order of 0.25 °C (Fig. 6).

4.7 El Niño Southern Oscillation

It has previously been observed that changing ocean gateways can significantly alter the modes of interannual variability of a climate model, even with weak changes in the mean climate (von der Heydt et al., 2011; Jochum et al., 2009). We focus our attention on the El Niño Southern Oscillation (ENSO) as it is the dominant mode of climate variability globally. Jochum et al. (2009) found that altering the Indonesia bathymetry led to a more irregular and weaker ENSO. Analysis of 200 years of SST anomaly in the Niño 3.4 region (5°S–5°N, 190–240°E) does not confirm the weakening of ENSO (Table 2), which is instead stronger in this study.

The low-resolution version of CESM used here has a power spectrum of ENSO that is not unreasonable (Shields et al., 2012). There is too little power at periods above 4 years (Figure 8) compared to observations (Smith et al., 2008), with a slight concentration at 2.5 years. The Indonesian alterations tend to smear this concentration out to longer periods (Tab. 2) in agreement with Jochum et al. (2009).

An open Central American Seaway leads to marginally stronger ENSO, but with dominant period stretching towards ~ 3 years instead of ~ 2.5 years (Table 2). It is likely that this is caused by changes in the seasonal cycle in the Eastern Equatorial Pacific as hinted at by the alterations in the annual mean flow patterns (Fig. 2). There may also be a contribution from a shoaling of the thermocline (Supplementary Figure 2), but that is unlikely to be the dominant cause (Steph et al., 2010; Zhang et al., 2012). Previous work with a simple model suggested an open Central American Seaway reduces ENSO amplitudes and shortens its period (Heydt et al., 2011): the opposite to the effects observed here. Interestingly the existence of the Central American Seaway removes the skew in ENSO – making La Niña deviations as strong as El Niño ones (Table 2). The correlation patterns associated with El Niño appear substantively similar in all four integrations (not shown). The opening of the Central American Seaway leads to only a slightly stronger link to ENSO in the Caribbean, despite the new ocean connection.

5. Discussion

We have compared the impacts of three gateway changes that potentially occurred during the late Miocene-Pliocene within the same model framework. In a global sense, the closing of the Central American Seaway through the creation of the Isthmus of Panama has the largest effects. The opening of the Bering Strait, thereby connecting the North Pacific to the Arctic, also had global consequences.

Our simulation with a closed Central American Seaway show changes in SST and SSS over the North Atlantic, but statistically significant only in some regions. There is a ~ 1 °C cooling over the North Atlantic, while surface salinity in the North Atlantic increases in the subtropics, but decreases slightly in higher latitudes. The model does show a reduction in the Atlantic Meridional Overturning Circulation of ~ 2 Sv, but a recovery with a timescale longer than one thousand years means this reduction is much

507 smaller than seen after the first 500 years (~ 5 Sv). The impact of the closed Bering Strait
508 on the AMOC is opposite and approximately equal to that of an open CAS, yet causes
509 significant changes to the surface climate of the North Atlantic (with surface
510 temperature changes in the North Atlantic exceeding 1°C). The global climate impacts of
511 altering the land/ocean configuration of Indonesia appear insignificant. The weakening
512 of the AMOC in the opened CAS experiment and the strengthening of the AMOC in the
513 closed Bering Strait experiments are paralleled by the increase and decrease of the
514 ocean vertical stratification, respectively (Supplementary Fig. 1).

515
516 The changes around Indonesia represent only a gateway constriction, not a complete
517 opening or closing like in the other two simulations. This may explain their small
518 impacts, but the imposed alterations are representative of changes seen over the past
519 several million years. Nevertheless, one could ask whether sufficient land mass has been
520 removed from the model to truly test the hypothesis of Cane and Molnar (2001) that
521 Indonesia change were responsible for African aridification. The western boundary
522 current in the South Pacific that runs along the coast of New Guinea has not been given
523 an opportunity to flow unimpeded into the Banda Sea (as the resolution is too coarse).
524 The two prior studies with coupled climate models are equivocal: Krebs et al. (2011)
525 show an increased throughflow, whilst Jochum et al. (2009) show little change.
526 Nonetheless neither previous studies, nor this one, show substantial remote impacts to
527 the mean climate to support the African aridification hypothesis.

528
529 As the exact bathymetry of the Indonesian Archipelago several million years ago is
530 not known, other alterations may have driven climate changes. An increased tidal
531 mixing is one such possible alteration (Brierley and Fedorov, 2011), although even that
532 does not have strong non-local consequences to the mean climate. A gradual increase in

the areal extent of Indonesia can be potentially important for climate evolution (Molnar and Cronin, 2015), but is not tested here.

Previously the El Niño-Southern Oscillation (ENSO) has been suggested to alter in response to both tropical gateway changes: an open Central American Seaway to make ENSO weaker but more frequent (Heydt et al., 2011) and Indonesian alterations to make ENSO again weaker but less frequent (Jochum et al., 2009). Our simulations show some ENSO changes from the tropical gateway changes, although not those anticipated previously. We find both tropical gateways generally decrease the frequency of ENSO and increase its amplitude. However, the Bering and Indonesia simulations also develop a secondary, biennial peak in the ENSO spectrum. The most notable response to an open CAS appears to be an absence in skew of ENSO. We would be cautious in interpreting these ENSO changes, since their statistical significance is hard to estimate and some may be model dependent.

This study has treated each gateway change in isolation. It is possible that there could be non-linear interactions between gateway changes and broader changes in climate. Examples of these non-linearities have been documented by several studies in other contexts. For example, von der Heydt and Dijkstra (2008; 2006) found the combined impacts of the Central American Seaway and the Drake Passage cause different circulation regimes depending on their configuration. Hu et al. (2010) have found that the closing of the Bering Strait during glacial intervals can lead to greater impacts in the North Atlantic than during an interglacials. In this study, an attempt was made to combine the multiple gateway changes, but this led to unrealistic drifts in the salt budget culminating in numerical errors. However, given the potentially compensating character of the Bering Strait opening and the CAS closure, and a weak effect of Indonesia drift on climate, we anticipate that such non-linear interaction are

not a major factor in Miocene-Pliocene climate evolution. Uncertainties in the depth of gateways appear to be much more important.

The three gateways investigated here had not previously been investigated in a single model. The present model, a CESM version with a relatively low resolution, was chosen for computational efficiency, but potentially these results could be model dependent. However, the responses are qualitatively similar to prior simulations (Zhang et al., 2012; Kerbs et al., 2011; Wadley and Bigg, 2002). Another issue is systematic cold biases in the present and other climate models. However, the relatively small climatic impacts of the imposed gateway alteration (e.g. $\sim 10\%$ change in the AMOC, SST changes on the order of 1°C or less in our experiments) suggest a linear regime, which should not be strongly affected by these biases.

A number of processes that are missing or under-resolved in climate models can be also relevant in a Pliocene and/or Gateway context, including tidal mixing (Brierley and Fedorov, 2011), tropical cyclone feedbacks (Fedorov et al., 2010), changes in tropical convection (Arnold et al., 2015), cloud properties (Burls and Fedorov, 2014), atmospheric chemistry (Unger and Yue, 2014), stratospheric connections (Joshi and Brierley, 2013) and ocean eddies (Viebahn et al., 2015).

6. Conclusions

It is widely thought that changes in the inter-ocean gateways played a role in the development of Northern Hemisphere glaciation during the Plio-Pleistocene (e.g. Haug and Tiedemann, 1998). The onset of glaciation was probably a threshold response to a gradual reduction in atmospheric greenhouse gases (DeConto et al., 2008; Lunt et al., 2008), whose precise timing was controlled by variations in Earth's orbital parameters favorable for ice accumulation. Inter-ocean gateway changes may have altered the Earth

system's response to (orbital) forcing and so helped set the level of that threshold. Specifically, the closing of the Central American Seaway (CAS) has been suggested as preconditioning glaciation through its impact on the North Atlantic (Haug and Tiedemann, 1998). Subsequently, the climate role of the CAS closure has received substantial discussion (e.g. Steph et al., 2010; Zhang et al., 2012; Horikawa et al., 2015), but was questioned by some studies (Molnar, 2008).

Here, for the first time, we have compared the climate impacts of three inter-ocean gateway changes that potentially occurred in the Late Miocene or early Pliocene within a single coupled model. Whilst the impacts of the Bering Strait are not as globally pervasive as those of the Central American Seaway, they have a stronger signature in the high northern latitudes pertinent for glacial inception (Fig. 6). It is uncertain when exactly the Bering Strait opened, although a Pliocene date seems probable. A recent global topography for 3.2 Ma (Haywood et al., 2014) reconstructs a Bering Strait that has not yet opened, although the Central American Seaway has already closed. De Schepper et al. (2013) suggest a shallow CAS was critical in aborting an early attempt at initiating glacial cycles at 3.3 Ma. Yet perhaps the closed Bering Strait is a more likely culprit: it has a greater impact in the North Atlantic and is now reconstructed to have changed after 3.3 Ma.

We hope that this work will provoke further consideration of changes in the Bering Strait - especially given its nearly equal and opposite impact on the deep ocean circulation as that of the CAS closure. The simulations presented here are individual sensitivity studies. Further work is required to test whether the relative impacts of the gateways remain the same with other climate models and with more representative boundary conditions. The role of non-linear interactions between inter-ocean gateways (e.g. von der Heydt and Dijkstra, 2008) and systematic biases in model simulations (e.g.

Burls and Fedorov, 2014) also need investigation. Yet, if indeed the two effects (the CAS closure and the Bering strait opening) were nearly compensating as suggested by these results, one must consider other mechanisms that led to global cooling since the late Miocene. A likely mismatch in the timing of the opening of the Bering Strait and the closure of the CAS is another factor to consider.

Acknowledgements

Financial support was provided by the grants to A.V.F. from the US National Science Foundation (AGS-1405272) and the David and Lucile Packard Foundation. This work was supported in part by the UCL Legion High Performance Computing Facility (Legion@UCL) and associated support services, and the Yale University Faculty of Arts and Sciences High Performance Computing facility.

References

- Arnold, N., Branson, M., Kuang, Z., Randall, D.A., Tziperman, E., 2015. MJO intensification with warming in the Super-Parameterized CESM. *J. Climate*, 28(7), 2706–2724.
- Brierley, C., Fedorov, A.V., 2011. Tidal mixing around Indonesia and the Maritime continent: Implications for paleoclimate simulations. *Geophys. Res. Lett.* 38, L24703. doi:10.1029/2011GL050027
- Burls, N.J., Fedorov, A.V., 2014. Simulating Pliocene warmth and a permanent El Niño-like state: the role of cloud albedo, *Paleoceanography*, 29(10), 893-910, doi:10.1002/2014PA002644.
- Cane, M.A., Molnar, P.H., 2001. Closing of the Indonesian seaway as a precursor to East African aridification around 3-4 million years ago. *Nature* 411(24), 265–271. doi:10.1038/35075500
- Coates, A.G., Jackson, J.B.C., Collins, L.S., Cronin, T.M., Dowsett, H.J., Bybell, L.M., Jung, P.,

637 Obando, J.A., 1992. Closure of the Isthmus of Panama: The near-shore marine record
 638 of Costa Rica and western Panama. Geological Society of America Bulletin 104, 814–
 639 828. doi:10.1130/0016-7606(1992)104<0814:COTIOP>2.3.CO;2
 640 Danabasoglu, G., Bates, S.C., Briegleb, B.P., Jayne, S.R., Jochum, M., Large, W.G., Peacock, S.,
 641 Yeager, S.G., 2012. The CCSM4 Ocean Component. J. Climate 25, 1361–1389.
 642 doi:10.1175/JCLI-D-11-00091.1
 643 Danabasoglu, G., Large, W.G., Briegleb, B.P., 2010. Climate impacts of parameterized
 644 Nordic Sea overflows. J. Geophys. Res. 115, C11005. doi:10.1029/2010JC006243
 645 De Boer, A.M., Nof, D., 2004. The Bering Strait's grip on the northern hemisphere climate.
 646 *Deep Sea Research Part I: Oceanographic Research Papers*, 51(10), 1347–1366.
 647 <http://doi.org/10.1016/j.dsr.2004.05.003>
 648 De Schepper, S., Groeneveld, J., Naafs, B.D.A., Van Renterghem, C., Hennissen, J., Head,
 649 M.J., Louwye, S., Fabian, K., 2013. Northern Hemisphere Glaciation during the
 650 Globally Warm Early Late Pliocene. PLoS ONE 8, e81508.
 651 doi:10.1371/journal.pone.0081508
 652 DeConto, R.M., Pollard, D., Wilson, P.A., Pälike, H., Lear, C.H., Pagani, M., 2008. Thresholds
 653 for Cenozoic bipolar glaciation. Nature 455, 652–656. doi:10.1038/nature07337
 654 DiNezio, P.N., Clement, A.C., Vecchi, G.A., Soden, B.J., Broccoli, A.J., Otto-Bliesner, B.L.,
 655 Braconnot, P., 2011. The response of the Walker circulation to Last Glacial
 656 Maximum forcing: Implications for detection in proxies. Paleoceanogr. 26, n/a–n/a.
 657 doi:10.1029/2010PA002083
 658 Dowsett, H.J., Robinson, M., Haywood, A.M., Salzmann, U., Hill, D.J., Sohl, L.E., Chandler,
 659 M., Williams, M., Stoll, D.K., 2012. The PRISM3D paleoenvironmental reconstruction.
 660 Stratigraphy 7, 123–139.
 661 Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P.,
 662 Rahmstorf, S., Raymo, M.E., 2015. Sea-level rise due to polar ice-sheet mass loss
 663 during past warm periods. Science 349, aaa4019.

664 Fedorov, A.V., Brierley, C.M., Emanuel, K., 2010. Tropical cyclones and permanent El
665 Nino in the early Pliocene epoch. *Nature*, 463 (7284), 1066-1070.
666 doi:10.1038/nature08831

667 Fedorov, A.V., Brierley, C.M., Lawrence, K.T., Liu, Z., Dekens, P.S., Ravelo, A.C., 2013.
668 Patterns and mechanisms of early Pliocene warmth. *Nature* 496, 43–49.
669 doi:10.1038/nature12003

670 Fox-Kemper, B., Danabasoglu, G., Ferrari, R., Griffies, S., Hallberg, R., Holland, M., Maltrud,
671 M., Peacock, S. and Samuels, B., 2011. Parameterization of mixed layer eddies. III:
672 Implementation and impact in global ocean climate simulations. *Ocean Modelling*,
673 39(1-2), 61-78.

674 Gent, P.R., Danabasoglu, G., Donner, L.J., Holland, M.M., Hunke, E.C., Jayne, S.R., Lawrence,
675 D.M., Neale, R.B., Rasch, P.J., Vertenstein, M., Worley, P.H., Yang, Z.-L., Zhang, M.,
676 2011. The Community Climate System Model Version 4. *J. Climate* 24, 4973–4991.
677 doi:10.1175/2011JCLI4083.1

678 Gordon, A.L., Sprintall, J., Van Aken, H.M., Susanto, D., 2010. The Indonesian throughflow
679 during 2004–2006 as observed by the INSTANT program. *Dynamics of Atmospheres*
680 *and Oceans* 50. 115–128.

681 Hall, R., 2002. Cenozoic geological and plate tectonic evolution of SE Asia and the SW
682 Pacific: computer-based reconstructions, model and animations. *Journal of Asian*
683 *Earth Sciences* 20, 353–431.

684 Haug, G.H., Tiedemann, R., 1998. Effect of the formation of the Isthmus of Panama on
685 Atlantic Ocean thermohaline circulation. *Nature* 393, 673–676.

686 Haug, G.H., Tiedemann, R., Zahn, R., Ravelo, A.C., 2001. Role of Panama uplift on oceanic
687 freshwater balance. *Geology*, 29(3), 207–210.

688 Haywood, A.M., Dowsett, H.J., Dolan, A.M., Rowley, D.B., Abe-Ouchi, A., Chandler, M., Lunt,
689 D.J., Salzmann, U., 2014. The Pliocene Model Intercomparison Project (PlioMIP)
690 Phase 2.

691 http://geology.er.usgs.gov/egpsc/prism/data/PlioMIP2_Science_Document_v1.pdf
692 (last accessed: 3rd August 015)
693 Hirst, A.C., Godfrey, J.S., 1993. The Role of Indonesian Throughflow in a Global Ocean
694 GCM. *J. Phys. Oceanogr.* 23, 1057–1086. doi:10.1175/1520-
695 0485(1993)023<1057:TROI>2.0.CO;2
696 Horikawa, K., Martin, E.E., Basak, C., Onodera, J., Seki, O., Sakamoto, T., Ikehara, M., Sakai,
697 S., Kawamura, K., 2015. Pliocene cooling enhanced by flow of low-salinity Bering Sea
698 water to the Arctic Ocean. *Nat Comms* 6, 7587. doi:10.1038/ncomms8587
699 Hu, A., Meehl, G.A., Otto-Bliesner, B.L., Waelbroeck, C., Han, W., Loutre, M.-F., Lambeck, K.,
700 Mitrovica, J.X., Rosenbloom, N., 2010. Influence of Bering Strait flow and North
701 Atlantic circulation on glacial sea-level changes. *Nature Geosci.* 3, 118–121.
702 doi:10.1038/ngeo729
703 Jayne, S.R., 2009. The Impact of Abyssal Mixing Parameterizations in an Ocean General
704 Circulation Model. *J. Phys. Oceanogr.* 39, 1756–1774. doi:10.1175/2009JPO4085.1
705 Jochum, M., Fox-Kemper, B., Molnar, P.H., Shields, C., 2009. Differences in the Indonesian
706 seaway in a coupled climate model and their relevance to Pliocene climate and El
707 Niño. *Paleoceanogr.* 24, n/a–n/a. doi:10.1029/2008PA001678
708 Joshi, M., Brierley, C., 2013. Stratospheric modulation of the Boreal response to Pliocene
709 tropical Pacific sea surface temperatures. *Earth Planet. Sci. Lett.*, 365, 1–6.
710 Karas, C., Nürnberg, D., Gupta, A.K., Tiedemann, R., Bickert, T., 2009. Mid-Pliocene
711 climate change amplified by a switch in Indonesian subsurface throughflow. *Nature*
712 *Geosci.* 2, 434–438. doi:10.1038/ngeo520
713 Krebs, U., Park, W., Schneider, B., 2011. Pliocene aridification of Australia caused by
714 tectonically induced weakening of the Indonesian throughflow. *Palaeogeography,*
715 *Palaeoclimatology, Palaeoecology* 309, 111–117. doi:10.1016/j.palaeo.2011.06.002
716 LaRiviere, J.P., Ravelo, A.C., Crimmins, A., Dekens, P.S., Ford, H.L., Lyle, M., Wara, M.W.,
717 2013. Late Miocene decoupling of oceanic warmth and atmospheric carbon dioxide

718 forcing. *Nature*, 486(7401), 97–100. <http://doi.org/10.1038/nature11200>

719 Lawrence, K.T., Sossdian, S.M., White, H.E., Rosenthal, Y., 2010. North Atlantic climate
720 evolution through the Plio-Pleistocene climate transitions. *Earth and Planetary
721 Science Letters* 300, 329–342. doi:10.1016/j.epsl.2010.10.013

722 Lunt, D.J., Foster, G.L., Haywood, A.M., 2008. Late Pliocene Greenland glaciation
723 controlled by a decline in atmospheric CO₂ levels. *Nature* 454, 1102–1105.
724 doi:10.1038/nature07223

725 Lunt, D.J., Valdes, P.J., Haywood, A.M., Rutt, I.C., 2007. Closure of the Panama Seaway
726 during the Pliocene: implications for climate and Northern Hemisphere glaciation.
727 *Climate Dynam.* 30, 1–18. doi:10.1007/s00382-007-0265-6

728 Maier Reimer, E., Mikolajewicz, U., Crowley, T.J., 1990. Ocean general circulation model
729 sensitivity experiment with an open central American isthmus. *Paleoceanogr.* 5,
730 349–366.

731 Marincovich, L., Gladenkov, A.Y., 1999. Evidence for an early opening of the Bering Strait.
732 *Nature* 397, 149–151.

733 Mestas Nuñez, A.M., Molnar, P.H., 2014. A mechanism for freshening the Caribbean Sea
734 in pre-Ice Age time. *Paleoceanogr.* 29, 508–517. doi:10.1002/2013PA002515

735 Mikolajewicz, U., Maier Reimer, E., Crowley, T.J., Kim, K.-Y., 1993. Effect of Drake and
736 Panamanian Gateways on the circulation of an ocean model. *Paleoceanogr.* 8, 409–
737 426. doi:10.1029/93PA00893

738 Molnar, P.H., 2008. Closing of the Central American Seaway and the Ice Age: A critical
739 review. *Paleoceanogr.* 23, PA2201. doi:10.1029/2007PA001574

740 Molnar, P.H., Cronin, T.W., 2015. Growth of the Maritime Continent and its possible
741 contribution to recurring Ice Ages. *Paleoceanogr.* 30, 196–225.
742 doi:10.1002/2014PA002752

743 Montes, C., Cardona, A., Jaramillo, C., Pardo, A., Silva, J.C., Valencia, V., Ayala, C., Pérez-
744 Angel, L.C., Rodriguez-Parra, L.A., Ramirez, V., Niño, H., 2015. Middle Miocene

745 closure of the Central American Seaway. *Science* 348, 226–229.
 746 doi:10.1126/science.aaa2815
 747 Paleoclimate Working Group, 2015. CESM1 Paleo Users Guide.
 748 <https://www2.cesm.ucar.edu/working-groups/pwg/documentation/cesm1-paleo->
 749 [ug](https://www2.cesm.ucar.edu/working-groups/pwg/documentation/cesm1-paleo-) (last accessed 3rd August 2015).
 750 Phillips, A.S., Deser, C., Fasullo, J., 2014. Evaluating Modes of Variability in Climate
 751 Models. *Eos* 95, 453–455. doi:10.1002/2014EO490002
 752 Rodgers, K.B., Latif, M., Legutke, S., 2000. Sensitivity of equatorial Pacific and Indian
 753 Ocean watermasses to the position of the Indonesian Throughflow. *Geophys. Res.*
 754 *Lett.* 27, 2941–2944. doi:10.1029/1999GL002372
 755 Rowley, D.B., Forte, A.M., Moucha, R., Mitrovica, J.X., Simmons, N.A., Grand, S.P., 2013.
 756 Dynamic Topography Change of the Eastern United States Since 3 Million Years Ago.
 757 *Science* 340, 1560–1563. doi:10.1126/science.1229180
 758 Shaffer, G., 1994. Role of the Bering Strait in controlling North Atlantic. *Nature* 367, 354-
 759 357.
 760 Shields, C.A., Bailey, D.A., Danabasoglu, G., Jochum, M., Kiehl, J.T., Levis, S., Park, S., 2012.
 761 The Low-Resolution CCSM4. *J. Climate* 25, 3993–4014. doi:10.1175/JCLI-D-11-
 762 00260.1
 763 Smith, T.M., Reynolds, R.W., Peterson, T.C., Lawrimore, J., 2008. Improvements to NOAA's
 764 Historical Merged Land–Ocean Surface Temperature Analysis (1880–2006). *J.*
 765 *Climate* 21, 2283–2296. doi:10.1175/2007JCLI2100.1
 766 Steph, S., Tiedemann, R., Prange, M., Groeneveld, J., Schulz, M., Timmermann, A.,
 767 Nürnberg, D., Rühlemann, C., Saukel, C., Haug, G.H., 2010. Early Pliocene increase in
 768 thermohaline overturning: A precondition for the development of the modern
 769 equatorial Pacific cold tongue. *Paleoceanogr.* 25, PA2202.
 770 doi:10.1029/2008PA001645
 771 Unger, N., Yue, X., 2014. Strong chemistry-climate feedbacks in the Pliocene, *Geophys.*

772 Res. Lett., 41, 527–533, doi:10.1002/2013GL058773.
 773 von der Heydt, A.S., Dijkstra, H.A., 2008. The effect of gateways on ocean circulation
 774 patterns in the Cenozoic. *Global Planet. Change* 62, 132–146.
 775 doi:10.1016/j.gloplacha.2007.11.006
 776 von der Heydt, A.S., Nnafie, A., Dijkstra, H.A., 2011. Cold tongue/Warm pool and ENSO
 777 dynamics in the Pliocene. *Clim. Past* 7, 903–915. doi:10.5194/cp-7-903-2011
 778 Viebahn, J.P., von der Heydt, A.S., Le Bars, D., Dijkstra, H.A., 2015. Effects of Drake
 779 Passage on a strongly eddying global ocean. Submitted.
 780 <http://arxiv.org/abs/1510.04141>
 781 Wadley, M.R., Bigg, G.R., 2002. Impact of flow through the Canadian Archipelago and
 782 Bering Strait on the North Atlantic and Arctic circulation: An ocean modelling study.
 783 *Quarterly Journal of the Royal Meteorological Society* 128, 2187–2203.
 784 doi:10.1256/qj.00.35
 785 Woodgate, R.A., 2005. Revising the Bering Strait freshwater flux into the Arctic Ocean.
 786 *Geophys. Res. Lett.* 32, L02602. doi:10.1029/2004GL021747
 787 Zhang, X., Prange, M., Steph, S., Butzin, M., Krebs, U., Lunt, D.J., Nisancioglu, K.H., Park, W.,
 788 Schmittner, A., Schneider, B., Schulz, M., 2012. Changes in equatorial Pacific
 789 thermocline depth in response to Panamanian seaway closure: Insights from a
 790 multi-model study. *Earth and Planetary Science Letters* 317–318, 76–84.
 791 doi:10.1016/j.epsl.2011.11.028

792

Gateway	Simulation	Mass (Sv)	Heat (TW)	Salt (10^6 kg/s)
Bering	Control	0.906 ± 0.017	-1.367 ± 0.356	29.4 ± 0.6
	Indonesia	<i>0.922</i>	<i>-0.935</i>	<i>29.7</i>
	Bering	0 [†]	0 [†]	0 [†]
	Panama	0.588	<i>-1.903</i>	18.7
Indonesia	Control	13.8 ± 0.2	891 ± 14	486 ± 7
	Indonesia	12.0	749	424
	Bering	15.5	959	544
	Panama	9.23	698	325
Panama	Control	0 [†]	0 [†]	0 [†]
	Indonesia	0 [†]	0 [†]	0 [†]
	Bering	0 [†]	0 [†]	0 [†]
	Panama	3.69	116*	151*

794

795 **Table 1. Transport through the gateways.** The total mass, heat and salt
796 transported through each gateway in each simulation (positive is defined as flow out of
797 the Pacific). The error on the preindustrial control simulation represents the standard
798 deviation of 25-year segments during the integration (sect. 2.3). Values in italics do not
799 have statistically significant differences from the control simulation. [†]There must be no
800 transport through closed gateways. *Unfortunately, the diagnostics for the eastward
801 component of heat and salt fluxes caused by isopycnal diffusion nor the sub-mesoscale
802 eddy parameterization were stored during the simulation, so their contributions are
803 assumed to be zero.

804

805

	Niño 3.4 SST Anomaly		
Simulation	Amplitude (°C)	Skew (°C ²)	Dominant Period (years)
Control	0.75	0.45	2.4
Indonesia	0.84	0.40	4.0
Bering	0.74	0.47	3.4
Panama	0.81	0.24	3.1
Observations	0.85	0.27	3.3

806

807 **Table 2. ENSO metrics in the simulations.** The standard deviation and skew of the
808 time series of Niño 3.4 sea surface temperature anomalies. The dominant period is
809 computed as the frequency with the most spectral power between 1.2 and 8 years
810 (computed before smoothing as in Fig 8).

811

Figure Captions

Figure 1. The model bathymetry on the temperature (left) and velocity (right) grids surrounding the three gateways under investigation: Indonesia (top), the Bering Strait (middle) and the Central American Seaway (bottom). Green indicates land points. The red crosshatched area shows the land points that have been removed or created in the each sensitivity study.

Figure 2. The upper ocean (top 150m) average velocities around the Bering Strait and Central American Seaway. The difference between the control (top) and perturbed (middle) simulations is shown at the bottom. The arrows represent the ocean velocity, whilst the color indicates its magnitude. The solid black line delineates the land points (Fig. 1) and the crosshatched area also incorporates the grid points whose velocity is zero (through the non-slip boundary conditions). Only statistically significant velocity changes are shown in the lower panels. *Note color scales differ between the left and right panels.*

Figure 3. The upper ocean (top 150m) average velocities around Indonesia. The flow regime in the control simulations is shown at the top. The flows in the various sensitivity simulations (left) and its difference from the control (right) are shown below. The arrows represent the upper ocean velocity, whilst the color indicates its magnitude. The solid black line delineates the land points (Fig. 1) and the crosshatched area also incorporates the grid points whose velocity is zero (through the non-slip boundary conditions). Only statistically significant velocity changes are shown in the right panels. *Note the magnitude of the change in velocity arrows differs (although the color scales do not).*

838

839 **Figure 4.** The poleward heat transports in the control simulation (a) are shown
840 along with the changes in each sensitivity experiment of total (b), atmospheric (c) and
841 oceanic (d) heat transports. The blue envelope indicates the 5-95% range of internal
842 variability estimated from 25 year segments of the control simulation. The atmospheric
843 term is calculated as a residual.

844

845 **Figure 5.** The Atlantic Meridional Overturning Circulation (AMOC). (a) Maximum of
846 the Atlantic meridional overturning streamfunction in the control (blue), Indonesia
847 (orange), Panama (purple) and Bering (green) simulations. Decadal smoothing has been
848 applied. The mean in the control simulation (dashed) and the 5-95% range (dotted) are
849 shown as horizontal lines. (b) Atlantic meridional overturning streamfunction in the
850 final 200 years of the control simulation. (c-e) Changes in the streamfunction observed
851 in the final 200 years of the sensitivity experiments. Those changes covered with
852 diagonal lines are not statistically significant with respect to the 25-year internal
853 variability in the control simulation.

854

855 **Figure 6.** Global impacts of the gateway changes on annual average surface air
856 temperature (left in °C) and annual average precipitation rates (right as %). The
857 stippling indicates a 200 year average change that is statistically different from the 25-
858 year internal variability in the control simulation. *Note: chronologically the patterns*
859 *would be reversed, so an opening of the Bering Strait would lead to a cooling of the North*
860 *Atlantic.*

861

862 **Figure 7.** The impact of the gateway changes on the sea surface salinity (psu). The
863 stippling indicates a 200 year average change that is statistically different from the 25-
864 year internal variability in the control simulation (sect. 2.3).

865

866 **Figure 8.** El Niño Southern Oscillation (ENSO). (a-d) Progression of the Niño 3.4 sea
867 surface temperature anomalies in the four simulations. (e) Power spectra for the Niño
868 3.4 sea surface temperature anomalies from the ERSST observations (Smith et al., 2008)
869 and in each simulation (following Phillips et al., 2014, but with a five-point smoothing).
870 The area under each line integrates across all periods to give the total variance. *Note:*
871 *whilst the time series are shown for only the final fifty years of each simulation (for*
872 *legibility), the spectra and the statistics quoted in Table 2 are taken over the whole two*
873 *hundred years of the intercomparison.*

874

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8