Continental Flood Basalt Weathering as a Trigger for Neoproterozoic Snowball Earth

Grant M. Cox^{1,2,*}, Galen P. Halverson^{1,2}, Ross K. Stevenson³, Michelle Vokaty¹, André Poirier³,
Marcus Kunzmann¹, Zheng-Xiang Li², Steven W. Denyszyn⁴, Justin V. Strauss⁵, Francis A.
Macdonald⁶

* Corresponding author: grant.cox@curtin.edu.au

1 - Department of Earth and Planetary Sciences, McGill University, Montréal, Québec, Canada.

2 - ARC Centre of Excellence for Core to Crust Fluid Systems and The Institute for Geoscience Research,
Department of Applied Geology, Curtin University, WA, Australia.

3-Département des sciences de la Terre et de l'atmosphère/Geotop, Université du Québec à Montréal, Montréal,
Québec, Canada.

4-School of Earth and Environment/Centre for Exploration Targeting, The University of Western Australia, Crawley,
WA, Australia.

5. Department of Earth Sciences, Dartmouth College, HB6105 Fairchild Hall, Hanover, NH, 03755.

6-Department of Earth and Planetary Sciences, Harvard University, Cambridge MA, USA.

Word count: Main Text excluding references and figure captions = 5631

ABSTRACT

Atmospheric CO₂ levels and global climate are regulated on geological timescales by the silicate weathering feedback. However, this thermostat has failed multiple times in Earth's history, most spectacularly during the Cryogenian (c. 720–635 Ma) Snowball Earth episodes. The unique middle Neoproterozoic paleogeography of a rifting, low-latitude, supercontinent likely favored a globally cool climate due to the influence of the silicate weathering feedback and planetary albedo. Under these primed conditions, the emplacement and weathering of extensive continental flood basalt provinces may have provided the final trigger for runaway global glaciation. Weathering of continental flood basalts may have also contributed to the characteristically high carbon isotope ratios ($\delta^{13}\text{C}$) of Neoproterozoic seawater due to their elevated P contents. In order to test these hypotheses, we have compiled new and previously published Neoproterozoic Nd isotope data from mudstones in northern Rodinia (North America, Australia, Svalbard, and South China) and Sr isotope data from carbonate rocks. The Nd isotope data are used to model the mafic detrital input into sedimentary basins in northern Rodinia. The results reveal a dominant contribution from continental flood basalt weathering during the ca. 130 m.y. preceding the onset of Cryogenian glaciation, followed by a precipitous decline afterwards. These data are mirrored by the Sr isotope record, which reflects the importance of chemical weathering of continental flood basalts on solute fluxes to the early–middle Neoproterozoic ocean, including a pulse of unradiogenic Sr input into the oceans just prior to the onset of Cryogenian glaciation. Hence, our new data support the hypotheses that elevated rates of flood basalt weathering contributed to both the high average $\delta^{13}\text{C}$ of seawater in the Neoproterozoic and to the initiation of the first (Sturtian) snowball Earth.

1. INTRODUCTION

The Neoproterozoic Era (1000–541 Ma) began with the amalgamation of the supercontinent Rodinia and ended with the assembly of Gondwana. Intervening global glaciations, massive perturbations to the carbon cycle, eukaryotic diversification, and presumed oxygenation of the atmosphere and ocean are broadly coincident with the fragmentation of Rodinia (Hoffman et al., 1998; Condon et al., 2005; Swanson-Hysell et al., 2010, Li et al., 2013). Mechanistic interconnections among these events are commonly invoked, but remain conjectural. Middle Neoproterozoic paleogeography (Fig. 1) likely favored both a cool global climate and elevated burial rates of organic carbon (Swanson-Hysell et al., 2010), as evinced in unusually high ($\sim 5\text{‰}$) average carbon isotope compositions of marine carbonates ($\delta^{13}\text{C}_{\text{carb}}$) at this time (Fig. 2). Horton (2015) recently proposed that the high $\delta^{13}\text{C}_{\text{carb}}$ values were driven by the weathering of extensive large igneous provinces (LIPs) during the Neoproterozoic, which would have greatly increased riverine phosphorus fluxes to the global ocean.

The onset of the Sturtian Snowball Earth ca. 717 Ma (Macdonald et al., 2010) brought an end to over one billion years of non-glacial climate and a brief resumption in the deposition of iron formation. Like the subsequent Marinoan glaciation, which ended at 635 Ma (Condon et al., 2005), the Sturtian glaciation was preceded by a deep negative $\delta^{13}\text{C}_{\text{carb}}$ anomaly—the Islay excursion (Fig. 2; Strauss et al., 2014). Several triggers have been proposed for the initiation of snowball glaciations that are linked to these pre-glacial $\delta^{13}\text{C}_{\text{carb}}$ excursions, including a transient increase then abrupt decline in methane fluxes (Schrag et al., 2002) and enhanced organic export production and anaerobic remineralization (Tzipermann et al., 2011). However, recent radiometric age constraints indicate that the Islay excursion preceded Sturtian glaciation by >10 Ma, implicitly decoupling it from the onset of global glaciation (Rooney et al., 2014; Strauss et

al., 2014). A third and still viable hypothesis is that the onset of the Sturtian snowball glaciation was driven by accelerated uptake of CO₂ resulting from the emplacement and weathering of Neoproterozoic LIPs as the rifting supercontinent drifted into the wet tropics—the Fire and Ice hypothesis (Goddéris et al., 2003; Donnadieu et al., 2004).

Because LIP's have geochemical signatures that are distinct from average continental crust, massive weathering of their surficial carapaces, continental flood basalts, should leave a distinct imprint in the sedimentary record. For example, siliciclastic sediments deposited in basins proximal to continental flood basalts (CFB's) should record primitive Nd isotope signatures and other geochemical characteristics typical of a dominantly mafic provenance. Chemical weathering of CFBs should also influence ocean chemistry (Mills et al., 2014), for example through the input of unradiogenic Sr and Os (Taylor and Lasaga, 1999; Ravizza and Peucker-Ehrenbrink, 2003). Consequently, the hypothesis that a flare-up in LIP emplacement in the Neoproterozoic contributed to fertilization of the oceans (Horton, 2015) and triggering of Cryogenian glaciation (Goddéris et al., 2003; Rooney et al., 2014) is testable in the sedimentary record.

To test the predictions made by this hypothesis, we have compiled new and previously published Neoproterozoic Nd isotope data on mudstones from basins proximal to a cluster of early-middle Neoproterozoic LIPs in northwestern North America, Australia, South China, and Svalbard and Sr isotope ratios (⁸⁷Sr/⁸⁶Sr) from well preserved marine carbonates globally. These datasets are coupled to a new $\delta^{13}\text{C}_{\text{carb}}$ compilation for the Neoproterozoic. Together, the data reveal a dominant contribution of CFB weathering to both clastic and solute fluxes in the ca. 130 m.y. preceding Cryogenian glaciation, followed by a dramatic decrease afterwards. These results attest to a direct role of CFB weathering on modulating the extraordinary paleoenvironmental

perturbations of the Neoproterozoic Era and reveal the missing link between supercontinental break-up, high $\delta^{13}\text{C}_{\text{carb}}$, and snowball glaciation.

2. LARGE IGNEOUS PROVINCES IN THE NEOPROTEROZOIC

Large igneous provinces result from the extrusion of $>10^6 \text{ km}^3$ of mafic volcanic rocks, typically in less than a few million years. Although their geodynamic origin remains controversial (Sobolev et al. 2007; Wang et al., 2015), their association with supercontinental break-up is well established (Courtilot et al., 1999). They are also closely coupled to many Phanerozoic environmental catastrophes including mass extinctions and climatic perturbations (Wignall, 2001; Sobolev et al., 2011; Schaller et al., 2011; Blackburn et al., 2013). LIP emplacement, in particular CFB volcanism, perturbs global climate through the massive release of magmatic gases, most notably SO_2 , which can drive global cooling on a timescale of years, and CO_2 , which triggers global warming over 10^1 – 10^5 years (Sobolev et al., 2011; Schaller et al., 2011; Blackburn et al., 2013). CFBs also lower pCO_2 on longer timescales (10^5 – 10^7 years) through enhanced silicate weathering owing to the ~ 5 – 10 times greater weatherability of basalt compared to felsic continental crust (White and Brantley, 1995; Dessert et al., 2001). For example, long-term decline in pCO_2 likely followed the emplacement of both the ca. 66 Ma Deccan Traps (Dessert et al., 2001, Kent and Muttoni, 2013, Lefebvre et al., 2013) and the ca. 201 Ma Central Atlantic Magmatic Province (CAMP) (Schaller et al., 2012).

Multiple large Neoproterozoic CFBs were emplaced on northern Rodinia in the early–middle Neoproterozoic (ca. 850–720 Ma; Figs. 1, 2). The ca. 830 Ma Guibei and Willouran LIP in South China and Australia is estimated to have covered a combined $\sim 10^6 \text{ km}^2$, comparable in size to the end-Permian Siberian Traps (Ernst et al., 2008). Northwestern Laurentia (North

America) hosts the ca. 780 Ma Gunbarrel LIP, which along with the coeval Kanding LIP in South China, covered an estimated original area of $1.6 \times 10^6 \text{ km}^2$ (Ernst et al., 2008). The ca. 755 Ma Mundine Well LIP in western Australia (Wingate and Giddings, 2000) is inferred to have been relatively small ($0.2 \times 10^6 \text{ km}^2$), but roughly coeval LIPs are widespread elsewhere in Rodinia, including the Shaba Gabbro in South China (Ernst et al., 2008). Emplacement of the ca. 725–712 Ma Franklin LIP centered on northern Laurentia temporally overlapped the onset of Sturtian glaciation (Fig. S3) but apparently did not result in sufficient outgassing of CO_2 to prevent an albedo runaway (Nabelek et al., 2014). The Franklin LIP has been estimated at $2.2 \times 10^6 \text{ km}^2$ in original area (Ernst et al., 2008), but was likely much larger given evidence for coeval mafic magmatism in northwestern Canada and Arctic Alaska (Macdonald et al., 2010; Cox et al., 2015), as well as slightly older but possibly cogenetic mafic intrusions in southwestern Siberia (Ariskin et al., 2013; Ernst et al., 2013) (Fig. 1b).

By the time the Franklin LIP was emplaced in northern Canada, northern Rodinia and its abundant CFB's were positioned in the wet tropics (Fig. 1). Combined with the incipient rifting of Rodinia at this time and the resulting exposure of more of the northern Rodinia land area to seaways, this paleogeography would have been uniquely suited to drive rapid basalt weathering. However, even though full break-up of Rodinia did not occur until the Ediacaran, LIP magmatism on the continents appears to have declined precipitously following the Franklin event (Fig. 2). Several relatively small LIP's were emplaced in the middle Ediacaran as the Iapetus Ocean opened (Ernst et al., 2008; McCausland et al., 2011), but combined with the drift of this region across the sub-tropics and towards higher latitudes (Li et al., 2013), their impact on global silicate weathering would have been minimal compared to that of the northern Rodinia LIP's leading up to the Cryogenian glaciation.

3. METHOD SUMMARY

Summaries of the methods used for this study are outlined below. Full details of these methods are provided in the supplementary information.

3.1. Carbon Isotopes

Carbon isotope analyses were performed following conventional procedures (e.g. Halverson et al., 2010). In short, sample powders were drilled from slabs of carbonate hand samples, avoiding fractures, heterogeneities, and weathered surfaces. Aliquots of approximately 50–100 μg of the homogenized, powdered samples were analyzed via dual-inlet mass spectrometry on a Nu Instruments Perspective™ at McGill University. All data are expressed in conventional per mil (‰) notation and calibrated to the Vienna Pee Dee Belemnite (VPDB) reference standard. This new compilation can be found in Table S1 of the accompanying supplementary material.

3.2. Sm-Nd Isotopes

Nd and Sm isotope ratios of mudstone, iron formation, and basalt samples were analyzed via thermal ionization mass spectrometry (TIMS) on a Thermo Triton™ mass spectrometer following standard multi-acid dissolution and anion exchange chromatography protocols at Geotop/Université du Québec à Montréal. Nd isotope ratios ($^{143}\text{Nd}/^{144}\text{Nd}$) are presented as $\epsilon_{\text{Nd}(t)}$, where the $^{144}\text{Nd}/^{143}\text{Nd}$ ratio is normalized to CHUR as a function of age (t) and expressed in parts per ten thousand (Table S2).

3.3. Sr Isotopes

Carbonates were pre-screened based on their Mg/Ca, Mn/Sr, Sr, and where possible, Rb concentrations or Rb/Sr ratios. Although generally samples with low Mg/Ca (<0.01), Mn/Sr

(<0.1), and Rb/Sr (<0.1) and high Sr concentrations (>500 ppm) are optimal, the criteria are variable for each succession, with the degree of alteration a function of the initial mineralogy and post-burial history of each sample (Edwards et al., 2015). Prescreened samples were leached with ammonium acetate and then rinsed three times in MQH₂O. The calcite fraction of the samples was then dissolved in 0.5M acetic acid. Following anion exchange chromatography, the isolated Sr fractions were measured via thermal ionization mass spectrometry (TIMS) on a Thermo Triton™ at Geotop/Université du Québec à Montréal. This data can be found in Table S3.

4. RESULTS

4.1 C isotope compilation

The compilation of $\delta^{13}\text{C}_{\text{carb}}$ values shown in Figure 2 is modified from previous compilations (Halverson et al., 2005, 2010), with additional data from Maloof et al. (2010) and Swanson-Hysell et al. (2010) and new data from Yukon (Macdonald et al., 2012 and this contribution), the southern Urals (Kuznetsov et al., 2006) and Siberia (Knoll et al., 1995 and this contribution), which extend the record back to the Mesoproterozoic–Neoproterozoic boundary. The age constraints for these successions, in particular in the Urals and Siberia, are very poor, but the relative ages are robust. This compilation suggests that moderate fluctuations in $\delta^{13}\text{C}$ (up to 6‰) were a feature of the earliest Neoproterozoic, but extreme ^{13}C -enrichment ($>5\text{‰}$) did not begin until ca. 850 Ma. Another key feature of the revised $\delta^{13}\text{C}_{\text{carb}}$ record is the clear offset (i.e. >10 m.y.) between the minima of the pre-Sturtian Islay anomaly and the onset of the Sturtian glaciation (Strauss et al., 2014). Indeed, the Sturtian glaciation now appears to be presaged instead by a sharp rise in $\delta^{13}\text{C}$ (from $< -5\text{‰}$ to $>5\text{‰}$).

The carbon isotope compilation is used to calibrate both the Nd and Sr isotope records, such that even if ages for individual samples are not always accurate or precise, the relative ages

of the C, Nd, and Sr isotope data sets are harmonized. Age calibrations are revised from age models in Halverson et al. (2010) and Swanson-Hysell et al. (2010), with modifications based on new ages from northwest Canada (Macdonald et al., 2010; Strauss et al., 2014; Rooney et al., 2014, 2015). For each individual succession, ages between calibration points were determined, where possible, by basic thermal subsidence modeling (e.g. Halverson et al., 2002; Le Guerroué et al., 2006). Where this technique was not possible, or over short stratigraphic intervals, age models were constructed by linear interpolation between correlated ages based on the assumption of constant sediment accumulation rates.

4.2 Nd isotope compilation

Samples selected for Nd isotope analyses include mudstones, the matrices of glacial diamictite and iron formation. Our new data from North America, Australia, and Svalbard were combined with published data from North America, Australia, and South China (Table S2) to generate the four regional compilations shown in Figure 3.

The Nd compilation from NW Laurentia displays a strikingly similar pattern to South China and Australia (Barovich and Foden, 2000; Wang et al., 2011), which together define a broad high in ϵ_{Nd} leading into the Cryogenian period, followed by a marked decline from the late Cryogenian to the early Palaeozoic (Fig. 3a–c). These data are consistent with Laurentia's position close to Australia and South China in northern Rodinia (Fig. 1) and highlight an extensive mafic sediment source for 130 million years preceding Sturtian glaciation. The new NW Laurentia data reveal additional detail. Specifically, ϵ_{Nd} is high (>-3) in the early Neoproterozoic, implying a separate source of juvenile Nd, most likely the remains of the $>2.7 \times 10^6 \text{ km}^3$ Mackenzie LIP (Ernst et al., 2008) in NW Laurentia. A short-lived decline in ϵ_{Nd} to -10 at ca. 850 Ma likely reflects an episode of continental extension at this time that initiated

early Neoproterozoic basins in NW Laurentia (Macdonald et al., 2012) and exposed fresh, evolved, continental crust. ϵ_{Nd} values then rise, with a culmination during the Sturtian glaciation, following Franklin magmatism. ϵ_{Nd} values decrease steeply following the Sturtian glaciation, reaching basement values (<-15) by the latter Ediacaran as the western margin of Laurentia fully opened (Colpron et al., 2003).

Data from mudstones in Svalbard, which lay off Laurentia (Fig. 1), also show a rise in ϵ_{Nd} values preceding Sturtian glaciation and an abrupt decline afterwards. These results imply a prominent juvenile source area leading into Cryogenian glaciation, consistent with an extensive Franklin LIP and our paleogeographic reconstruction (Fig. 1). The Australian Nd compilation is distinct from the Laurentian data insofar as values are close to that of Australian continental basement (~ -10) in the early Neoproterozoic but rise to near zero in the matrix of Sturtian glacial deposits. This trend presumably reflects the low abundance of basalt on Australia prior to emplacement of the Willouran and possibly Guibei LIP and the dominance of basaltic provenance by the onset of the Sturtian glaciation.

We also analyzed syn-glacial iron formation (IF) in Australia and Canada, which in both cases comprise a combination of hematitic jaspilite and mudstone (Klein and Beukes, 1993; Cox et al., 2016). Samples from the Rapitan IF in NW Laurentia, which occurs within basalt clast-rich glacial deposits, show a range of relatively high ϵ_{Nd} (-4 to +2), with the most juvenile values overlapping the temporal arrays of the Gunbarrel and Franklin LIPs (Fig. 3). The Holowilena IF in South Australia has higher detrital content and displays a larger range in ϵ_{Nd} (-10 to -3), but still overlaps the Mundine Well LIP array (Fig. 3b). Together, these data reflect the preponderance of juvenile Nd in both the waters and sediments of the basins in which they formed.

4.3 Modeling the mafic contribution to detrital sediments (f_{maf})

To estimate the magnitude of the basaltic component of mudstone samples (f_{maf} = fraction derived from CFB), a mixing lattice was constructed utilizing the measured Nd concentrations and initial Nd isotope ratios ($^{143}\text{Nd}/^{144}\text{Nd}_{(i)}$) of local LIP's and cratonic basement (Fig. 4). For these calculations, we combined data from northwestern Canada and Svalbard to construct a single northern Laurentia dataset. Neoproterozoic LIP's show a relatively restricted range in ϵ_{Nd} and [Nd], especially when compared with values for regional cratonic basement. Therefore we have fixed ϵ_{Nd} and [Nd] values for the basaltic end member for each region (i.e. northern Laurentia, Australia, South China) in the calculations. In contrast, the cratonic basement province (CBP) end member is allowed to vary in [Nd]. The ϵ_{Nd} value used for the CBP end member is estimated from the compilation of ϵ_{Nd} data available for cratonic basement of each region (Fig. 4a, b, c).

The resulting contoured mixing lattices (Fig. 4d, e, f) can be used to calculate the contribution of CFB to the mudstone samples based on their ϵ_{Nd} and Nd concentrations, and hence to calculate an f_{maf} value (Figs. 4g–i, 5) for each sample. From these data, we constructed f_{maf} curves for each region by applying a 10-point locally weighted scatterplot smoothing process (LOESS). The f_{maf} profiles elucidate the prominent shift from mafic to felsic provenance spanning the Cryogenian period (Fig. 5). During the Ediacaran, f_{maf} values from North America and Australia decline towards a dominantly CBP provenance, likely enhanced by the final stages of Rodinia break-up, which was not accompanied by massive CFB volcanism (Fig. 2b). The f_{maf} data also reveal important distinctions between regions. For example, f_{maf} rises abruptly from lower values in the early Neoproterozoic in both South China and Australia, reflecting the importance of ca. 830 Ma Willouran and Guibei LIP in covering old continental crust with young

basalts. In South China, the CFB contribution is >80% prior to the onset of the Cryogenian and declines to the lowest values (~0.2) by the close of the Cryogenian.

4.4 Sr isotope compilation

The compilation of $^{87}\text{Sr}/^{86}\text{Sr}$ data shown in Figure 6 is revised from previous compilations (Halverson et al., 2007, 2010) and includes data from Kaufman et al. (1993), Burns et al. (1994), Misi and Veizer (1998), Calver (2000), Yoshioka et al. (2003) and Pokrovskii et al. (2006), with additional published data from Bartley et al., (2001), Kuznetsov et al. (2006), Melezhik et al. (2009), Rooney et al. (2014), Bold et al. (in press) and our own previously unpublished data (Table S2). For each individual succession, ages were determined, where possible, by calibration with the $\delta^{13}\text{C}_{\text{carb}}$ compilation, such that even where absolute ages are inaccurate, their relative ages should be broadly correct.

The new $^{87}\text{Sr}/^{86}\text{Sr}$ compilation (Fig. 6) reveals a pronounced trough in $^{87}\text{Sr}/^{86}\text{Sr}$ (down to 0.7053) in the early Neoproterozoic that coincides with the ca. 950–850 Ma interval between Rodinia assembly and the onset of break-up and is reminiscent of the late Paleozoic low in $^{87}\text{Sr}/^{86}\text{Sr}$ that followed the assembly of Pangaea (Burke et al., 1982). $^{87}\text{Sr}/^{86}\text{Sr}$ then rises abruptly to values >0.7060 around 850–840 Ma, prior to a decline back to ~0.7055 by 830 Ma, synchronous with the emplacement of the Willouran and Guibei LIP's. A second steep rise in $^{87}\text{Sr}/^{86}\text{Sr}$ to near 0.7070 by 800 Ma is interrupted by a shift to a roughly flat trend and a ca. 780 Ma downturn that coincides broadly with the Gunbarrel and Kanding LIP's. Following a brief recovery to values near 0.7070, $^{87}\text{Sr}/^{86}\text{Sr}$ then drops abruptly to as low as 0.7064 just prior to the onset of Cryogenian glaciation.

This pre-Cryogenian downturn is most clearly seen in data from the Coates Lake Group in the Mackenzie Mountains of NW Canada (Rooney et al., 2014), which fills small basins that

were at least intermittently restricted. A similar decline is also evident in combined data from Svalbard and East Greenland. Hence, it appears that this sharp decline in $^{87}\text{Sr}/^{86}\text{Sr}$ is a true seawater signal that heralds impending snowball glaciation.

During the Cryogenian glacial interlude, $^{87}\text{Sr}/^{86}\text{Sr}$ rises abruptly to new steady state values of ~ 0.7073 , (Fig. 6), attesting to the role of Sturtian glaciation in removing unradiogenic basalt and exposing fresh, old, continental crust (Rooney et al., 2014). Strontium isotope ratios again shift abruptly upward (to > 0.7080) in the early Ediacaran, reflecting further removal of CFB cover from the continents and the transient effect of the silicate weathering drawdown of the extraordinarily large, post-Marinoan atmospheric CO_2 reservoir (Le Hir et al., 2009).

5. DISCUSSION

5.1. Silicate Weathering of CFB's in the Neoproterozoic

The Nd isotope compilation for mudstones and coupled f_{maf} calculations reveal an exceptionally high detrital contribution from mafic lithologies to sedimentary basins in northern Rodinia during the early–middle Neoproterozoic, followed by a sharp decline spanning the Cryogenian glaciations (Figs. 6, 7). These results are consistent with the geological record, which shows a flare-up in LIP's between ca. 850–720 Ma, with many of these clustered around northern Rodinia, followed by a lull in LIP emplacement during the latter Cryogenian and early Ediacaran periods.

Given the global paleogeography leading to Cryogenian glaciation, with the extensive CFB provinces located in the tropics (Fig. 1), it is reasonable to assume that these basalts were not only extensively physically weathered, but also intensely chemically weathered. Indeed, physical and chemical weathering are closely coupled (Gaillardet et al., 1999; Maher and Chamberlain, 2014), and several direct lines of evidence attest to the role of chemical weathering of these

CFB's and their influence on riverine solute fluxes to the early Neoproterozoic oceans. First, ϵ_{Nd} data on marine carbonates from the paleo-Pacific Ocean (Keto and Jacobsen, 1988) show a virtually identical trend to our mudstone data (Fig. 7), with high pre-Cryogenian values and sharply declining Ediacaran values, implicating a prominent early Neoproterozoic source of dissolved juvenile Nd to seawater. Second, a compilation of chemical index of alteration (CIA) data from the studied basins in northern Rodinia shows moderately to strongly chemically weathered signatures in early Neoproterozoic samples, clearly offset from CFB and average upper continental crust (Fig. S2). Finally, the seawater strontium isotope record, which is sensitive to the abundance of CFB's due to their unradiogenic initial Sr-isotopic compositions (Taylor and Lasaga, 1999), mirrors the Nd isotope trends, providing independent evidence of the Cryogenian removal of the flood basalt carapace from northern Rodinia. Consequently the combined dataset of Nd and Sr isotopes supports enhanced silicate weathering of CFB's during the middle Neoproterozoic, and in particular leading into the Cryogenian period.

5.2 Lithological controls on phosphorus supply and the role of CFB versus arc weathering

If it is assumed that phosphorus is the limiting nutrient on geological timescales (Tyrrell, 1999), the variations in phosphorus content within continental lithologies must exert significant control on nutrient availability over geological time scales. To demonstrate the importance of surficial lithology on globally integrated P fluxes from the continents, we have compiled ~14,000 analyses of Archean cratonic basement rocks, ~13,000 CFB samples, and ~68,000 arc volcanics (Fig. 8). Similar to the results of Horton (2015), these datasets reveal that the peak in P_2O_5 concentration occurs between 50–53% SiO_2 , which corresponds to an overall mafic (i.e. primitive) composition. Interestingly, the broad shape of this P_2O_5 vs. SiO_2 distribution closely matches the theoretical fractional crystallization path for a basaltic melt. This suggests that the P-

enrichment observed in these datasets is largely the result of the high solubility of P in basaltic
magmas (Toplis et al. 1994) and that basaltic compositions are reached before apatite saturation.
Adjusting this model to include the effect of assimilation and fractional crystallization (AFC)
results in higher P_2O_5 for any given SiO_2 composition. Importantly, similar peak P_2O_5
concentrations also occur in basalts in arc-related volcanics (Fig. 8b), and this enrichment, as
compared to oceanic equivalents (i.e. MORB; 8c) is also likely the result of AFC processes.
However, whereas arc-basalts are P-enriched, most arc-related volcanics are andesitic in
composition, which contain significantly less P_2O_5 (Fig. 8b). Therefore, although arc weathering
presumably plays an important role in the long-term continental P flux to the oceans, it does not
have the same potential to dramatically alter P fluxes as do more rapidly emplaced and removed
CFBs.

Additional evidence suggests that the variations in f_{maf} and $^{87}Sr/^{86}Sr$ spanning the
Cryogenian glaciations are most likely due to weathering of CFB and not other mafic lithologies.
First, with the possible exception of South China (Zhao et al., 2007), volcanic arcs were absent in
northern Rodinia during the middle Neoproterozoic (Li et al., 2013). And although early
Neoproterozoic basins in northern Laurentia (Macdonald et al., 2012, Turner and Bekker, 2015)
and Australia were the result of continental rifting prior to the onset of Rodinia break-up, these
rifting events were largely non-magmatic. Nor did fluctuations in arc length likely contribute
significantly to a global increase in mafic weathering in the middle Neoproterozoic. Both
theoretical considerations (Silver and Behn, 2008) and global U-Pb zircon age compilations
(Cawood et al., 2013; McKenzie et al., 2014; Roberts and Spencer, 2015) (Fig. 7) suggest that
total continental arc length was near a long-term minimum leading into Cryogenian glaciation
and did not peak until the late Ediacaran—coincident with the Neoproterozoic maximum in

⁸⁷Sr/⁸⁶Sr. Consequently, although we cannot rule out a contribution of juvenile Nd from isolated rift- or arc-related volcanics, we interpret the f_{maf} values (Fig. 6) to reflect principally the contribution of CFB weathering relative to the underlying continental basement.

5.3 The Biogeochemical Impact of CFB Weathering

Previous estimates by Horton (2015) and a compilation of chemical index of alteration (CIA) data from the studied basins (Fig. S1) indicate that approximately half of the initial mass of CFB source rocks was dissolved via chemical weathering during the Neoproterozoic. Given that CFB's have on average ~3 times as much P as typical continental crust, with much of this P residing in the fine-grained matrix (i.e., glass), and that Neoproterozoic LIP's, in particular the ca. 780 Ma Gunbarrel LIP (median P₂O₅ ≈ 0.5), are exceptionally rich in P (Horton, 2015), it follows that the early–middle Neoproterozoic would have experienced a sustained, anomalously high continental flux of phosphorus to the global oceans. This P flux was likely coupled to iron via adsorption onto ferric oxyhydroxides formed during oxidative weathering (Laakso and Schrag, 2014) of the CFB's and subsequent liberation in estuaries by desorption upon mixing of river and seawater (Froelich, 1988) and in porewaters of anoxic continental margin sediments by a combination of dissimilatory iron reduction, which likely played an important role in Precambrian iron cycling (Severmann et al., 2008, Li et al., 2015), and sulfide-catalyzed reduction of iron oxides (Poulton et al., 2015).

Direct evidence for the elevated terrestrial P flux in the Neoproterozoic is seen in the marked phosphorus enrichment observed in Cryogenian iron formation (Planavsky et al., 2010), which is thought to be the result of leaching of continental margin sediments dominated by basaltic detritus (Cox et al. 2016). Indirect evidence of the elevated P flux is found in the exceptionally high $\delta^{13}\text{C}_{\text{carb}}$ values that prevailed through much of the early–middle

Neoproterozoic. Assuming that P limited net primary productivity, then it follows that increased P would have increased organic carbon burial. Because the increased P flux would typically be accompanied by increased alkalinity flux (hence higher inorganic carbon burial) (Holland, 1984; Holland and Bekker, 2012), this factor alone should not strongly influence $\delta^{13}\text{C}_{\text{carb}}$. However, our calculations show that P/(CO₂D) ratios – a measure of the P released compared to CO₂ consumed during dissolution, and ultimately alkalinity generated by this process, is ~3 times greater for average CFB than for either average continental basement or mid-ocean ridge basalt (see supplementary information). Therefore, higher organic carbon burial would also have translated into higher fractional organic carbon burial (f_{org} = organic carbon burial/total carbon burial) and higher $\delta^{13}\text{C}_{\text{carb}}$ values, consistent with typically ¹³C-enriched carbonates during the early–middle Neoproterozoic (up to +8‰; Fig. 7). Conversely, the decrease in f_{maf} from ~0.6 to ~0.3 across the Cryogenian implies a six-fold decrease in total P flux from northern Rodinia, and this decline aligns with the observation that average $\delta^{13}\text{C}_{\text{carb}}$ values are much lower in the Ediacaran than in the early Neoproterozoic (Fig. 2).

Importantly, although it is likely that fertilization via weathering of CFB's helps to explain early-middle Neoproterozoic carbon isotope record (Horton, 2015), this mechanism alone cannot explain the very high average $\delta^{13}\text{C}_{\text{carb}}$ that prevailed for most of this time. Based on a simple model of the exogenic carbon cycle in which f_{org} is controlled by the P flux (see supplementary information), we find that the increased P flux from CFB weathering accounts for ~2‰ of the $\delta^{13}\text{C}_{\text{carb}}$ values above the typical Proterozoic background of 0‰. Consequently, other factors are required to explain the sustained high $\delta^{13}\text{C}_{\text{carb}}$. One explanation for further enhancement in organic carbon burial is the redox-dependent P-feedback (Van Cappellen and Ingall, 1994), by which organic-bound P would have been efficiently recycled during anaerobic

remineralization of primary organic matter under the dominantly anoxic deep waters (Sperling et al., 2015) of the early Neoproterozoic oceans. Another factor contributing to the high $\delta^{13}\text{C}_{\text{carb}}$ values may have been authigenic carbonate production and burial (Schrag et al., 2013), which would have been enhanced on continental shelves experiencing high primary production (Higgins et al., 2009; Thompson et al., 2015).

5.4 A CFB-Weathering Trigger for Snowball Glaciation?

At the onset of the Cryogenian period, the Franklin CFB sat entirely within the tropics (Denyszyn et al., 2009) (Fig. 1) where it would have had maximal impact on atmospheric CO_2 due to the sensitivity of hydrolysis reactions to temperature and rainfall (Dessert et al., 2003; Allègre et al., 2010; Kent and Muttoni, 2013). Seawater proxies from NW Laurentia are consistent with intense weathering of the Franklin LIP leading into the Sturtian Snowball glaciation: $^{87}\text{Sr}/^{86}\text{Sr}$ declined sharply from 0.7071 to 0.7064 (Fig. 2e) and $^{187}\text{Os}/^{188}\text{Os}$ fell synchronously with a rapid return to highly positive $\delta^{13}\text{C}_{\text{carb}}$ values following the ca. 735–730 Ma Islay excursion (Rooney et al., 2014). This drop in $^{87}\text{Sr}/^{86}\text{Sr}$ dwarfs similar declines following emplacement of large Phanerozoic LIP's (Fig. S6), including CAMP, for which an abrupt decline in atmospheric CO_2 has been reported (Schaller et al., 2012). Hence, chemical weathering in the pre-Cryogenian (i.e., Tonian) interval was exceptionally dominated by the hydrolysis of CFB's in the tropics.

In addition to the increase basalt weathering at this time, the accompanying elevated P flux fuelled greater primary productivity, raising f_{org} (as seen in the sharp rise in $\delta^{13}\text{C}_{\text{carb}}$ following the Islay negative anomaly), expanding anoxia on proximal continental margins, and increasing biological drawdown of atmospheric CO_2 . This eutrophication would have both increased P recycling—providing a strong positive feedback to primary productivity—and

transiently drawn down CO₂ due to increased alkalinity generated by bacterial iron and sulfate reduction (Tziperman et al., 2011). Under prevailing conditions of already low atmospheric CO₂ levels due to enhanced silicate weathering, this alkalinity pump to the deep ocean may have provided the final nudge past the critical pCO₂ threshold (Hoffman et al., 1998; Goddris et al., 2003; Donnadieu et al., 2002) for triggering runaway glaciation.

6. CONCLUSIONS

Coupled Nd and Sr isotope data sets for the Neoproterozoic reveal that weathering of unusually extensive continental flood basalts between ca. 850 and 717 Ma dominated the global silicate weathering feedback and continental chemical fluxes to the oceans. This effect was enhanced in the middle Neoproterozoic as the Rodinia supercontinent drifted into the tropics and began to rift apart, exposing more of these CFB's to warm, moist conditions that increased the rates of hydrolysis reactions. These CFB's were largely (but not exclusively) clustered in northern Laurentia and neighboring cratons (Fig. 1). Because CFB's are strongly enriched in phosphorus relative to typical continental crust, the high weathering rate of the CFB's would have fertilized the global oceans (Horton, 2015). High riverine P fluxes, combined with extensive anoxia on continental margins (Sperling et al., 2015), would have promoted efficient P recycling, sustaining high rates of organic carbon burial, as indicated by the carbon isotope record at this time. This additional control of CFB weathering on atmospheric CO₂ levels adds credibility to the Fire and Ice hypothesis (Goddris et al., 2003; Donnadieu et al., 2004), in which it is argued that weathering of CFB lowered pCO₂ sufficiently to trigger snowball glaciation. Our new data suggest specifically that it was the emplacement rapid weathering of the Franklin LIP that pushed the Earth system across this critical threshold and ushered in the extraordinary climatic and geochemical perturbations that characterize the Neoproterozoic Era.

If intensified chemical weathering of Neoproterozoic CFB's and coupled fertilization of the oceans (Horton, 2015) triggered glaciation through rapid consumption of CO₂, Snowball glaciation itself was a powerful negative feedback on CFB weathering. It was previously argued that Cryogenian glaciation would have radically altered the global continental surface by removing a veneer of infertile regolith that developed on continental interiors and leaving behind a fresh, weatherable surface (Swanson-Hysell et al., 2010). Perhaps more importantly, Cryogenian glaciations stripped the continents of their blanket of basalt, removing the source of nutrients that fuelled long-lived positive carbon isotope perturbation and preventing subsequent snowball glaciations.

ACKNOWLEDGMENTS

This research was partially supported by Natural Sciences and Engineering Research Council (NSERC) Discovery grants to GPH and RSS, an NSERC Vanier fellowship to GMC, the Yukon Geological Survey, Polar Continental Shelf Program, and an NSF graduate research fellowship to JVS. GPH acknowledges donors of the American Chemical Society Petroleum Research Fund for partial support of this research. FAM acknowledges NSF Sedimentary Geology and Paleontology and the NASA Astrobiology Institute for support. This manuscript benefited from reviews by Andrey Bekker and 2 anonymous reviewers. Lee Kump and John Hayes provided valuable feedback on earlier drafts of this manuscript.

REFERENCES

Allègre, C.J., Louvat, P., Gaillardet, J., Meynadier, L., Rad, S., Capmas, F., 2010. The fundamental role of island arc weathering in the oceanic Sr isotope budget. *Earth Planet. Sci. Lett.* 292, 51–46.

455 Ariskin, A.A., Kostitsyn, Y.A., Konnikov, E.G., Danyushevsky, L.V., Meffre, S., Nikolaev, G.S.,
 456 McNeill, A., Kislov, E.V., Orsoev, D.A., 2013. Geochronology of the Dovyren Intrusive Complex,
 457 Northwestern Baikal Area, Russia, in the Neoproterozoic. *Geochem. Int.* 51, 859–875.

458 Barovich, K.M., Foden, J., 2000. A Neoproterozoic flood basalt province in southern- central Australia:
 459 geochemical and Nd isotope evidence from basin fill. *Precambrian Res.* 100, 213–234.

460 Blackburn, T.J. et al., 2013. Zircon U-Pb geochronology links the End-Triassic mass extinction with the
 461 Central Atlantic Magmatic Province. *Science* 340, 941–945.

462 Bold, U., Smith, E.F., Rooney, A.D., Bowring, S.A., Buchwaldt, R., Dúdas, F.Ö., Ramezani, J., Crowley,
 463 J.L., Schrag, D.P., Macdonald, F.M. in press. Neoproterozoic stratigraphy of the Zavkhan Terrane of
 464 Mongolia: The backbone for Cryogenian and early Ediacaran chemostratigraphic records, *Am. J. Sci.*

465 Burke, W. H., Denison, R. E., Hetherington, E. A., Koepnick, R. B., Nelson, H. F., Otto, J. B., 1982.
 466 Variation of seawater $^{87}\text{Sr}/^{86}\text{Sr}$ through Phanerozoic time. *Geology* 10, 516–519.

467 Burns, S.J., Haudenschild, U., Matter, A., 1994. The strontium isotopic composition of carbonates from
 468 the Late Precambrian (~560–540 Ma) Huqf Group of Oman. *Chem. Geol.* 111, 269–282.

469 Calver, C., 2000. Isotope stratigraphy of the Ediacaran (Neoproterozoic III) of the Adelaide rift complex,
 470 Australia, and the overprint of water column stratification, *Precambrian Res.* 100 121–150.

471 Canfield, D. E., 1998. A new model for Proterozoic ocean chemistry. *Nature* 396, 450–453.

472 Canfield, D. E., Poulton, S. W., Knoll, A. H., Narbonne, G. M., Ross, G., Goldberg, T., Strauss, H., 2008.
 473 Ferruginous conditions dominated later Neoproterozoic deep-water chemistry. *Science* 321, 949–952.

474 Cawood, P.A., Hawkesworth, C.J., Dhuime, B., 2013. The continental record and the generation of
 475 continental crust. *Geol. Soc. Am. Bull.* 125, 14–32.

476 Colpron, M., Logan, J. M., Mortensen, J. K., 2003. U–Pb zircon age constraint for late Neoproterozoic
 477 rifting and initiation of the lower Paleozoic passive margin of western Laurentia. *Can. J. Earth Sci.* 39,

478 133–143.

479 Condon, D., Zhu, M., Bowring, S.A., Jin, Y., Yang, A., 2005. From the Marinoan glaciation to the oldest
480 bilaterians: U-Pb ages from the Doushantuo Formation, China. *Science* 308, 95–98.

481 Courtillot, V., Jaupart, C., Manighetti, I., Tapponier, P., and Besse, J., 1999. On causal links between
482 flood basalts and continental breakup. *Earth Planet Sci. Lett.* 166, 177–195.

483 Cox, G. M., Halverson, G. P., Poirier, A., Le Heron, D., Strauss, J. V., and Stevenson, R., 2016, A model
484 for Cryogenian iron formation: *Earth Plan. Sci. Lett.* 433, 280–292.

485 Cox, G. M., Strauss, J. V., Halverson, G. P., Schmitz, M. D., McClelland, W. C., Stevenson, R. S.,
486 Macdonald, F. A., 2015b. Kikiktat volcanics of Arctic Alaska—Melting of harzburgitic mantle associated
487 with the Franklin large igneous province. *Lithosphere* 7, 275–295.

488 Denyszyn, S.W., Halls, H.C., Davis, D.W., Evans, D.A.D., 2009a. Paleomagnetism and U-Pb
489 geochronology of Franklin dykes in High Arctic Canada and Greenland: a revised age and paleomagnetic
490 pole constraining block rotations in the Nares Strait region. *Can. J. Earth Sci.* 46, 689–705.

491 Denyszyn, S.W., Davis, D.W., Hall, H.C., 2009b. Paleomagnetism and U-Pb geochronology of the
492 Clarence Head dykes, Arctic Canada: orthogonal emplacement of mafic dykes in a large igneous province.
493 *Can. J. Earth Sci.* 46, 155–167.

494 Dessert, C., Dupré, B., Gaillardet, J., François, L.M., Allègre, C.J., 2003. Basalt weathering laws and the
495 impact of basalt weathering on the global carbon cycle. *Chem. Geol.* 202:257–273.

496 Dessert, C et al., 2001. Erosion of Deccan Traps determined by river geochemistry: impact on the global
497 climate and the $^{87}\text{Sr}/^{86}\text{Sr}$ ratio of seawater. *Earth Planet. Sci. Lett.* 188, 459–474.

498 Donnadieu, Y., Goddérès, Y., Ramstein, G., Nédélec, A., Meert, J., 2004. A snowball Earth climate
499 triggered by continental break-up through changes in runoff. *Nature* 428, 303–306.

500 Edwards, C.T., Saltzman, M.R., Leslie, S.A., Bergström, S.M., Sedlacek, A.R.C., Howard, A., Bauer,
 501 J.A., Sweet, W.C., Young, S.A., 2015. Strontium isotope ($^{87}\text{Sr}/^{86}\text{Sr}$) stratigraphy of Ordovician bulk
 502 carbonate: implications for preservation of primary values. *Geol. Soc. Am. Bull.* 127, 1275–1289.

503 Ernst, R.E., Bleeker, W., Söderland, U., Kerr, A.C., 2013. Large Igneous Provinces and supercontinents:
 504 Towards the complete tectonic revolution. *Lithos* 174:1–14.

505 Ernst, R.E., Wingate, M.T.D., Buchan, K.L., Li, Z.-X. 2008. Global record of 1600– 700 Ma Large
 506 Igneous Provinces (LIPs): Implications for the reconstruction of the proposed Nuna (Columbia) and
 507 Rodinia supercontinents. *Precambrian Res.* 160, 159– 178.

508 Fike, D.A., Grotzinger, J.P., Pratt, L.M., Summons, R.E., 2006. Oxidation of the Ediacaran Ocean. *Nature*
 509 44, 744–747.

510 Froelich, P.N., 1988. Kinetic control of dissolved phosphate in natural rivers and estuaries: a primer on the
 511 phosphate buffer mechanism. *Limnol. Oceanogr.* 33, 649–668.

512 Gaillardet, J., Dupré, Louvat, P., Allègre, C., 1999. Global silicate weathering and CO_2 consumption rates
 513 deduced from the chemistry of large rivers. *Chem. Geol.* 159, 3–30.

514 Ghiorso, M. S., and Sack, R. O., 1995. Chemical mass transfer in magmatic processes IV. A revised and
 515 internally consistent thermodynamic model for the interpolation and extrapolation of liquid-solid
 516 equilibria in magmatic systems at elevated temperatures and pressures: *Contr. Min. Petr.* 119, 197–212.

517 Goddéis, Y., Donnadieu, Y., Nédélec, A., Dupré, B., Dessert, C., Grard, A., Ramstein, G., François, L.,
 518 2003. The Sturtian 'snowball' glaciation: fire and ice. *Earth Plan. Sci. Lett.* 211, 1–12.

519 Halverson, G.P., Dudás, F.Ö., Maloof, A.C., Bowring, S.A. 2007. Evolution of the $^{87}\text{Sr}/^{86}\text{Sr}$ composition
 520 of Neoproterozoic seawater. *Palaeogeogr. Paleoclimatol. Palaeoecol.* 256, 103–129.

521 Halverson, G.P. Hoffman, P.F., Schrag, D.P., Kaufman, A.J., 2002. A major perturbation of the carbon
 522 cycle before the Ghaub glaciation (Neoproterozoic) in Namibia: Prelude to snowball Earth? *Geochem.*
 523 *Geophys. Geosys.* 3, 10.1029/2001GC000244.

524 Halverson, G.P., Hoffman, P.F., Schrag, D.P., Maloof, A.C., Rice, A.H., 2005. Toward a Neoproterozoic
 525 composite carbon-isotope record. *Geol. Soc. Am. Bull.* 117, 1181–1207.

526 Halverson, G.P., Wade, B.P., Hurtgen, M.T., Barovich, K., 2010. Neoproterozoic chemostratigraphy.
 527 *Precambrian Res.* 182, 337–350.

528 Heaman, L.H., LeCheminant, A.N., Rainbird, R.H. 1992. Nature and timing of Franklin igneous events,
 529 Canada: implications for a Late Proterozoic mantle plume and the break-up of Laurentia. *Earth Planet.*
 530 *Sci. Lett.* 109, 117–131.

531 Higgins, J.A., Schrag, D.P., Fischer, W.W., 2009. Oxygenation of the oceans and sediments:
 532 Consequences for the seafloor carbonate factory. *Earth Plan. Sci. Lett.* 284, 25–33.

533 Hoffman, P.F., Kaufman, A.J., Halverson, G.P., Schrag, D.P., 1998. A Neoproterozoic Snowball Earth.
 534 *Science* 281, 1342–1346.

535 Holland, H., 1984. *The Chemical Evolution of the Oceans and Atmosphere*. Princeton University Press.

536 Horton, F., 2015. Did phosphorus derived from the weathering of large igneous provinces fertilize the
 537 Neoproterozoic oceans? *Geochem. Geophys. Geosys.* 16:10.1002/2015GC005792.

538 Jenner, F. E., and O'Neill, H. S. C., 2012. Analysis of 60 elements in 616 ocean floor basaltic glasses:
 539 *Geochemistry, Geophysics, Geosystems*, 13, Q02005.

540 Kaufman, A.J., Jacobsen, S.B., Knoll, A.H. 1993. The Vendian record of Sr and C isotopic variations in
 541 seawater: implications for tectonics and paleoclimate. *Earth Plan. Sci. Lett.* 120, 409–430.

542 Kent, D.V., Muttoni, G., 2013. Modulation of Late Cretaceous and Cenozoic climate by variable
 543 drawdown of atmospheric pCO₂ provinces on continents drifting through the equatorial humid belt. *Clim.*
 544 *Past* 9, 525–546.

545 Keto, L.S., Jacobsen, S.B., 1988. Nd isotopic variation of Phanerozoic paleoceans. *Earth Plan. Sci. Lett.*
 546 90, 395–410.

547 Klein, C., Beukes, N.J., 1993. Sedimentology and geochemistry of the glaciogenic Late Proterozoic
 548 Rapitan iron-formation in Canada. *Econ. Geol.* 88, 542–565.

549 Knoll, A.H., Kaufman, A.J., Semikhatov, M.A., 1995. The carbon-isotopic composition of Proterozoic
 550 carbonates; Riphean successions from northwestern Siberia (Anabar Massif, Turukhansk Uplift). *Am. J.*
 551 *Sci.* 295, 823–850.

552 Kuznetsov, A.B., Semikhatov, M.A., Maslov, A.V., Gorokhov, I.M., Prasolov, E.M., Krupenin, M.T.,
 553 Kislova, I.V., 2006. New data on Sr-and C-isotopic chemostratigraphy of the Upper Riphean type section
 554 (Southern Urals). *Strat. Geol. Correlation* 14, 602–628.

555 Laakso, T.A., Schrag, D.P., 2014. Regulations of atmospheric oxygen during the Proterozoic. *Earth*
 556 *Planet. Sci. Lett.* 388, 81–91.

557 Le Guerroue, E., Allen, P.A., Cozzi, A., 2006. Chemostratigraphic and sedimentological framework of the
 558 largest negative carbon isotopic excursion in Earth history: The Neoproterozoic Shuram Formation, Nafun
 559 Group, Oman. *Precambrian Res.* 146, 68–92.

560 Le Hir, G., Donnadieu, Y., Goddris, Y., Pierrehumbert, R. T., Halverson, G. P., Macouin, M., Ndlec,
 561 A., Ramstein, G., 2009. The snowball Earth aftermath: Exploring the limits of continental weathering
 562 processes. *Earth Planet. Sci. Lett.* 277, 453–463.

563 Lefebvre, V., Donnadieu, Y., Goddris, Y., Fluteau, F., and Hubert-Thou, L., 2013. Was the Antarctic
 564 glaciation delayed by a high degassing rate during the early Cenozoic? *Earth Planet. Sci. Lett.* 371–372,
 565 203–211.

566 Li, W., Beard, B.L., Johnson, C., 2015. Biologically recycled continental iron is a major component in
 567 banded iron formations. *Proc. Nat. Acad. Sci. (USA)* 112, 8193–8198.

568 Li, Z.-X., Evans, D.A.D., Halverson, G.P., 2013. Neoproterozoic glaciations in a revised global
 569 paleogeography. *Sed. Geol.* 294, 219–232.

570 McCausland, P.J.A., Hankard, F., Van der Voo, R., Hall, C.M., 2011. Ediacaran paleogeography of

571 Laurentia: Paleomagnetism and geochronology of the 583 Ma Bae de Moutons syenite, Quebec.
 572 Precambrian Res. 187, 58–78.

573 McKenzie, N.R., Hughes, N.C., Gill, B.C., Myrow, P.M., 2014. Plate tectonic influences on
 574 Neoproterozoic–early Paleozoic climate and animal evolution. *Geology* 42, 127–130.

575 Macdonald, F. A., Schmitz, M. D., Crowley, J. L., Roots, C. F., Jones, D. S., Maloof, A. C., Strauss, J. V.,
 576 Cohen, P. A., Johnston, D. T., Schrag, D. P., 2010. Calibrating the Cryogenian. *Science* 327, 1241–1243.

577 Macdonald, F. M., Halverson, G. P., Strauss, J. V., Smith, E. F., Cox, C. M., Sperling, E., Roots, C. F.,
 578 Schrag, D. P., 2012. Early Neoproterozoic basin formation in the Yukon: Implications for the make-up
 579 and break-up of Rodinia. *Geosci. Can.* 39, 77–97.

580 Maher, K., Chamberlain, C.P., 2014. Hydrologic regulation of chemical weathering and the geologic
 581 carbon cycle. *Science* 343, 1502–1504.

582 Melezhik, V.A., Pokrovsky, B.G., Fallick, A.E., Kuznetsov, A.B., Bujakaite, M.I., 2009. Constraints on
 583 the $^{87}\text{Sr}/^{86}\text{Sr}$ of Late Ediacaran seawater: insights from high-Sr limestones. *J. Geol. Soc. Lon.* 166, 183–
 584 191.

585 Misi, A., Veizer, J. 1998. Neoproterozoic carbonate sequences of the Una Group, Irece Basin, Brazil:
 586 chemostratigraphy, age and correlations. *Precambrian Res.* 89, 87–100.

587 Mills, B., Daines, S.J., Lenton, T.M., 2014. Changing tectonic controls on the long- term climate cycle
 588 from Mesozoic to present. *Geochem. Geophys. Geosys.* 15, 4866– 4884.

589 Nabelek, P.I., Bédard, J.H., Rainbird, R.H., 2014. Numerical constraints on degassing of metamorphic
 590 CO_2 during the Neoproterozoic large igneous event, Arctic Canada. *Geol. Soc. Am. Bull.* 126:759–772.

591 Pehrsson, S.J., Buchan, K.L., 1999. Borden dykes of Baffin Island, Northwest Territories: a Franklin U-Pb
 592 baddeleyite age and a paleomagnetic reinterpretation. *Can. J. Earth Sci.* 36, 65–73.

593 Planavsky, N. J., Rouxel, O. J., Bekker, A., Lalonde, S. V., Konhauser, K. O., Reinhard, C. T., Lyons, T.

594 W., 2010. The evolution of the marine phosphate reservoir. *Nature* 467, 1088–1090.

595 Pokrovskii, B.G., Melezhik, V.A., Bujakaite, M.I. 2006. Carbon, oxygen, strontium, and sulfur isotopic
596 compositions in late Precambrian rocks of the Patom Complex, central Siberia: Communication 1.
597 Results, isotope stratigraphy, and dating problems. *Lith. Min. Res.* 41, 450–474.

598 Poulton, S.W., Henkel, S., März, C., Urquhart, H., Flögel, S., Kasten, S., Sinninghe Damsté, J.S., Wagner,
599 T., 2015. A continental-weathering control on orbitally driven redox-nutrient cycling during Cretaceous
600 Ocean Anoxic Event 2. *Geology* 43, 963–966.

601 Ravizza, G., Peucker-Ehrenbrink, B., 2003. Chemostratigraphic evidence of Deccan volcanism from the
602 marine osmium isotope record. *Science* 302, 1392–1395.

603 Roberts, N.M.W, Spencer, C.J., 2015. The zircon archive of continent formation through time. *Geol. Soc.*
604 *London Spec. Publ.* 389, 197–225.

605 Rooney, A.D., Macdonald, F.A., Strauss, J.V., Dudás, F.Ö., Hallmann, C., Selby, D., 2014. Re-Os
606 geochronology and coupled Os-Sr isotope constraints on the Sturtian snowball Earth. *Proc. Nat. Acad. Sci.*
607 (USA) 111, 51– 56.

608 Rooney, A.D., Strauss, J.V., Brandon, A.D., Macdonald, F.A., 2015. A Cryogenian chronology: Two
609 long-lasting synchronous Neoproterozoic glaciations. *Geology* 43, 459–462.

610 Schaller, M.F., Wright, J.D., Kent, D.V. 2011. Atmospheric PCO₂ perturbations associated with the
611 Central Atlantic Magmatic Province. *Science* 331, 1404–1408.

612 Schaller, M.F., Wright, J.D., Kent, D.V., Olsen, P.E., 2012. Rapid emplacement of the Central Atlantic
613 Magmatic Province as a net sink for CO₂. *Earth Plan. Sci. Lett.* 323-324, 27–39.

614 Schrag, D.P., Berner, R.A., Hoffman, P.F., Halverson, G.P., 2002. On the initiation of snowball Earth.
615 *Geochem. Geophys. Geosys.*, 3:10.1029/2001GC000219.

616 Schrag, D.P., Higgins, J.A., Macdonald, F.A., Johnston, D.T., 2013. Authigenic carbonate and the history

617 of the global carbon cycle. *Science*, 339, 540–543.

618 Severmann, S., Lyons, T.W., Anbar, A., McManus, J., and Gordon, G., 2008. Modern iron isotope
619 perspective on the benthic iron shuttle and the redox evolution of ancient oceans: *Geology* 36, 487–490.

620 Silver, P.G., Behn, M.G., 2008. Intermittent plate tectonics? *Science* 319, 18–85.

621 Smith, P.M., and Asimow, P.D., 2005. Adibat_1ph: A new public front-end to the MELTS, pMELTS,
622 and pHMELTS models: *Geochemistry, Geophysics, Geosystems*, 6, 1–8.

623 Sobolev, S.V., Sobolev, A.V., Kuzmin, D.V., Krivolutskaya, N.A., Petrunin, A.G., Arndt, N.T., Radko,
624 V.A., Vasiliev, Y.R., 2011. Linking mantle plumes, large igneous provinces and environmental
625 catastrophes. *Nature* 477, 312–316.

626 Sobolev, A.V., Hofmann, A.W., Kuzmin, D.V., Yaxley, G.M., Arndt, N.T., Chung, S.-L., Danyushevsky,
627 L.V., Elliott, T., Frey, F.A., Garcia, M.O., Gurenko, A.A., Kamenetsky, V.S., Kerr, A.C., Krivolutskaya,
628 N.A., Matvienkov, V.V., Nikogosian, I.K., Rocholl, A., Sigurdsson, I.A., Sushchevskaya, N.M., Teklay,
629 M., 2007. The amount of recycled crust in sources of mantle-derived melts. *Science* 316, 412–417

630 Sperling, E. A., Halverson, G. P., Knoll, A. H., Macdonald, F. A., Johnston, D. T., 2013. A basin redox
631 transect at the dawn of animal life. *Earth and Planetary Science Letters* 371–372, 143–155.

632 Strauss, J.V., Rooney, A.D., Macdonald, F.A., Brandon, A.D, Knoll, A.H., 2014. 740 Ma vase-shaped
633 microfossils from Yukon, Canada: Implications for Neoproterozoic geochronology and biostratigraphy.
634 *Geology* 42, 659–662.

635 Swanson-Hysell, N. L., Rose, C. V., Calmet, C. C., Halverson, G. P., Hurtgen, M. T., Maloof, A. C.,
636 2010. Cryogenian glaciation and the onset of carbon-isotope decoupling. *Science* 328, 608–611.

637 Tanaka, T., Togashi, S., Kamioka, H., Amakawa, H. Kagami, H., Hamamoto, T., Yuhara, M., Orihashi,
638 Y., Yoneda, S., Shimizu, H., Kunimaru, T., Takahashi, K., Yanagi, T., Nakano, T., Fujimaki, H., Shinjo,
639 R., Asahara, Y., Tanimizu, M., Dragusanu, C. 2000. JNdi-1: a neodymium isotopic reference in
640 consistency with LaJolla neodymium. *Chem. Geol.* 168, 279–281.

641 Taylor, A.S., Lasaga, A.C., 1999. The role of basalt weathering in the Sr isotope budget of the oceans.
642 Chem. Geol. 161, 199–214.

643 Taylor, S. R., and McLennan, S.M., 1985. The continental crust: its composition and evolution, Oxford,
644 Blackwell, 312 p.

645 Taylor, S. R., and McLennan, S.M., 1995. The geochemical evolution of the continental crust: Rev.
646 Geophys. 33, 241–265.

647 Tziperman, E., Halevy, I., Johnston, D.T., Knoll, A.H., Schrag, D.P., 2011. Biologically induced initiation
648 of Neoproterozoic snowball-Earth events. Proc. Nat. Acad. Sci. (USA) 108, 15091–15096.

649 Toplis, M. J., Libourel, G., and Carroll, M. R., 1994, The role of phosphorus in crystallisation processes of
650 basalt: An experimental study: Geochim. Cosmochim. Acta 58, 797–810.

651 Turner, E. C., and Bekker, A., 2015, Thick sulfate evaporite accumulations marking a mid-Neoproterozoic
652 oxygenation event (Ten Stone Formation, Northwest Territories, Canada): Geol. Soc. Am. Bull. 128, 203–
653 222.

654 Tyrrell, T., 1999. The relative influences of nitrogen and phosphorus on oceanic primary production:
655 Nature 400, 525–531.

656 Van Cappellen, P., Ingall, E.D., 1994. Benthic phosphorus regeneration, net primary production, and
657 ocean anoxia: a model of the coupled marine biogeochemical cycles of carbon and phosphorus.
658 Paleoceanography 9, 677–692.

659 Vervoort, J.D., Patchett, J.R., Blichert-Toft, J., Albarède, F., 1999. Relationships between Lu-Hf and Sm-
660 Nd isotopic systems in the global sedimentary system. Earth Plan. Sci. Lett. 168, 79–99.

661 Wang, X.-C., Li, Z.-X., Li, X.-H., Li, W.-L., Zhang, Q.-R., 2011.. Geochemical and Hf-Nd isotope data of
662 Nanhua rift sedimentary and volcanoclastic rocks indicate a Neoproterozoic continental flood basalt
663 provenance. Lithos 127, 427–440.

664 White, A.F., Brantley, S.L., 1995. Chemical weathering rates of silicate minerals: an overview. *Rev. Min.*
665 *Geochem.* 31, 1–22.

666 Wignall P.B., 2001. Large igneous provinces and mass extinctions. *Earth-Sci. Rev.* 53, 1–33.

667 Wingate, M.T.D., Giddings, J.W., 2000. Age and palaeomagnetism of the Mundine Well dyke swarm,
668 Western Australia: Implications for an Australia-Laurentia connection at 755 Ma. *Precambrian Res.* 100,
669 335–357.

670 Yoshioka, H., Asahara, Y., Tojo, B., Kawakami, S. 2003. Systematic variations in C, O, and Sr isotopes
671 and elemental concentrations in Neoproterozoic carbonates in Namibia: implications for glacial to
672 interglacial transition. *Precambrian Res.* 124, 69–85.

673 Zhao, J.-H., Zhou, M.-F., 2007. Geochemistry of Neoproterozoic mafic intrusions in the Panzihua district
674 (Sichuan Province, SW China): implications for subduction- related metasomatism in the upper mantle.
675 *Precambrian Res.* 152, 27–47.

Figure 1
a

720 Ma

Figure 2

Figure 4

Figure 5

Figure 6

Figure 8