Global seismic data reveal little water in the mantle transition zone

C. Houser^{a,*}

^a *Earth-Life Science Institute, Tokyo Institute of Technology, 2-12-1-IE-1 Ookayama, Meguro-ku, Tokyo, 152-8550, Japan.*

Abstract

Knowledge of the Earth's present water content is necessary to constrain the amount of water and other volatiles the Earth acquired during its formation and the amount that is cycled back into the interior from the surface. This study compares 410 and 660 km discontinuity depth with shear wave tomography within the mantle transition zone to identify regions with seismic signals consistent with water. The depth of the 410 and 660 km discontinuities is determined from a large updated dataset of SS-S410S and SS-S660S differential travel times, known as SS precursors. The discontinuity depths measured from binning and stacking the SS precursor data are then compared to the shear velocity model HMSL-S06 in the transition zone. Mapping all the possible combinations, very few locations match the predictions from mineral physics for the effects of water on discontinuity depth and shear velocity. The predictions, although not yet measured at actual transition zone temperatures and pressures, are a shallow 410 km discontinuity, a deep 660 km discontinuity, and a slow shear velocity. Only 8% of the bins with high-quality data are consistent with these predictions, and the calculated average water content within these bins is around 0.6 wt%. A few isolated locations have patterns of velocity/topography that are consistent with water, while there are large regional-scale patterns consistent with cold/hot temperature anomalies. Combining this global analysis of long period seismic data and the current mineral physics predictions for water in transition zone minerals, I find that the mantle transition zone is generally dry, containing less than one Earth ocean of water. Although subduction zones could be locally hydrated, the combined discontinuity and velocity data show no evidence that wadsleyite or ringwoodite have been globally hydrated by subduction or initial Earth conditions.

Keywords: mantle transition zone, mantle discontinuities, mantle shear velocity, SS precursors,

*Corresponding author

Email address: chouser@elsi.jp (C. Houser)

1. Introduction

Understanding the Earth's water budget through time requires constraining the amount of water that is currently present in the Earth's interior. The amount of water in the interior could be decreasing over time by degassing at divergent plate boundaries, increasing by transport within subducting oceanic plates, or in balance between these processes. Water may be able to reach the mantle transition zone within metastable mineral phases in cold subducting oceanic lithosphere (i.e. slabs) (Angel et al., 2001). Mineral physics studies have shown that nominally anhydrous wadsleyite and ringwoodite could contain close to 3% water (Inoue et al., 1995; Kohlstedt et al., 1996), which is a storage capacity of up to 10 oceans of water. Since planetary formation models are sensitive to the Earth's initial water content, it is necessary to put bounds on the amount of water in this potential reservoir. The mantle transition zone is here defined by global seismic reflectors consistent with phase transitions in the olivine component of the mantle at approximately 410 km (olivine to wadsleyite) and 660 km depth (ringwoodite to bridgmanite and ferropericlase).

Several studies have measured the effects of water on the properties of transition zone minerals and will be summarized briefly here. While wadsleyite and ringwoodite can hold up 3 wt.% water at ambient conditions (Smyth et al., 2004; Ohtani et al., 2000), the capacity for water storage decreases with increasing temperature such that 1-2 wt.% is often predicted at transition zone temperatures and pressures (Bolfan-Casanova et al., 2006; Mao et al., 2008), but could be as low as 0.2 wt.% (Dixon et al., 2002; Huang et al., 2005; Hirschmann, 2006; Inoue et al., 2010; Mao et al., 2012). Inoue et al. (2004) found that the seismologically observed bulk sound speed contrast across 410 km discontinuity is consistent with a bulk composition that is 60% olivine and 1.5 wt.% water. They also measured a lower density and bulk sound velocity for hydrous compared to anhydrous wadsleyite. Ab initio calculations indicate that both shear and compressional velocity will decrease by 1.5% for each 1.0 wt.% water added to hydrous wadsleyite (up to the maximum of 3.3 wt.%) (Tsuchiya and Tsuchiya, 2009). Smyth and Frost (2002) observed that water will stabilize wadsleyite relative to olivine, and therefore decrease the depth of the transition such that the 410 km discontinuity would be shallower (lower pressure) in the presence of water. By the same reasoning, since water is more compatible in ringwoodite than bridgmanite and ferropericlase, the 660 km discontinuity is deeper (higher pressure) for a hydrous peridotite composition (Litasov et al., 2005). In addition to these

30 experimental results, Pearson et al. (2014) measured 1.4 wt.% water in a ringwoodite inclusion in
31 a natural diamond.

32 Thio et al. (2016) point out that there have been no experiments that measure properties of
33 transition zone materials at both the high temperatures and pressures for water-bearing phases.
34 From a thermodynamic point of view, since water is more easily incorporated into wadsleyite than
35 olivine at the top of the transition zone and ringwoodite than bridgmanite and ferropericlase at
36 the base of the transition zone, water will expand the stability fields of wadsleyite and ringwoodite
37 and increase the thickness of the transition zone. This scenario could change if other minerals that
38 can incorporate water become stable at the high temperatures and pressures of the transition zone.
39 However, the consensus from mineral physics (Jacobsen and Smyth, 2006; Bolfan-Casanova et al.,
40 2006) is that the seismic signature of 1-2 wt.% water present in the mantle transition zone is slow
41 shear velocity, shallow 410 km discontinuity, and deep 660 km discontinuity, Figure 1.

42 Mineral physics predictions can be tested by comparing global maps of shear wave velocity
43 and discontinuity depth (Helffrich, 2002). Despite the fact that global tomographic models have
44 decreased resolution in the mantle transition zone, most tomographic models are in generally good
45 agreement about the large scale features such as the emergence of subducted slabs (Ritsema et al.,
46 2004). Global maps of discontinuity topography are constructed by measuring the time difference
47 between the surface-reflected SS phase and the S410S and S660S reflections (see Figure 2 and further
48 discussion in the methods section). Although they use different methods, there are several reliable
49 global discontinuity studies from SS precursors which are also in relatively good agreement about
50 the large scale features (Deuss, 2009). Thus, general conclusions can be made by combining any
51 tomography and discontinuity model to examine which regions may be compatible with the presence
52 of water. Here I compare an updated map of discontinuity topography to the shear velocity model
53 HMSL-S06 (Houser et al., 2008b).

54 2. Method

55 I use the same data and methodology as (Houser et al., 2008a) to obtain the SS-S410S and
56 SS-S660S differential times from 2006 to 2010. Since the S410S and S660S arrivals are too faint to
57 be seen above the noise on most individual seismograms, I stack seismograms with high quality SS
58 signals whose bounce points lie within 5° radius bins (Figure 2). The traces are first deconvolved
59 by the main SS pulse and then aligned and stacked according to their arrival times measured using

60 cluster analysis (Houser et al., 2008b). The SS-S410S and SS-S660S differential times are converted
61 to depth using 20 second PREM (Dziewonski and Anderson, 1981). For a full description of the
62 stacking procedure, see Houser et al. (2008a).

Table 1: SS Precursor Arrival Time Measurements

Author	Start	End	Number of Data
Flanagan and Shearer 1998	1976	1995	13,467
Houser et al. 2008a	1995	2005	26,348
Update	2006	2010	31,577
Total for this study	1976	2010	71,392

63 This update increases the amount of measured SS arrival times by over 40% (Table 1) and im-
64 proves the SS precursor stack quality in the northern Pacific, southern Pacific, eastern Asia, Indian
65 ocean, northern South America, and north Atlantic. Figure 3 shows the increase in the number
66 of traces per stack in each bin and representative stacks from the many different regions before
67 and after the update. While the improvement in coverage is dominant in the Pacific and eastern
68 Asia, there is improvement in a variety of tectonic settings. The resulting maps of topography on
69 the 410 and 660 km discontinuity and transition zone thickness are shown in Figure 4. I used the
70 total combined dataset from Flanagan and Shearer (1998), Houser et al. (2008a), and the updated
71 dataset of 2006-2010 for the SS precursor stacking. While the update improves coverage in several
72 regions, there are still many areas in the southern hemisphere that have too few data to obtain a
73 quality stack. In Figures 4, 6 and 7, I do not plot areas where the depth error exceeds 20 km.

74 I compare the updated discontinuity topography to a shear velocity model with the same in-
75 version method as HMSL-S06 but parameterized with 6° equal-area blocks (Figure 5). I use the
76 530-660 km depth layer within the transition zone parameterized at 6° (not 4° as in the original
77 HMSL-S06 (Houser et al., 2008b)) to be consistent with the broader sensitivity of the SS precursors.
78 The upper mantle is constrained by surface wave phase velocity maps of Love Bassin et al. (2000)
79 and Rayleigh (Laske 2004, personal communication) waves at frequencies of 4-15 mHz, where the
80 lowest frequencies are sensitive to transition zone structure. Although HMSL-S06 does not use
81 body waves that turn within the transition zone because the arrival times are complicated by the
82 triplications from the 410 and 660 km discontinuities, the body waves turning in the uppermost

lower mantle are partially sensitive to transition zone structure. I do not include the compressional wave model HMSL-P06 in this analysis since the upper mantle of that model is simply a scaled version of the shear wave model and thus does not add any additional constraint.

To compare the discontinuity topography and shear velocity, I use the analysis first described in Houser and Williams (2010). The combination of 410 and 660 km discontinuity depth and fast/slow seismic velocity anomaly results in 8 possible cases relating these three observations. The left side of Figure 6 shows the western (top) and eastern (bottom) hemispheres, where each circle is the bin location for the highest quality stacked precursor data. For instance, a stacked trace that has a shallower-than-average 410 km discontinuity and a deeper-than-average 660 km discontinuity will plot as dark blue if the seismic velocity is fast, or purple if the seismic velocity is slow. In general, cool colors are used for fast seismic velocities and warm colors are used for slow seismic velocities. The seismic velocity used is the 600 km depth slice of the 6° shear velocity model HMSL-S06 shown in Figure 5. The right side of Figure 6 highlights the 105 of 1301 bin locations (8%) that are consistent with water (slow shear velocity, shallow 410 km discontinuity, deep 660 km discontinuity).

In order to examine any regional velocity-topography relationships, I test the observed patterns against random arrays of points on a globe. For instance, there are a few locations in the right panel of Figure 6 where bins consistent with water appear to cluster, but it is not possible to say a region is consistent with water unless the clustering exceeds the clustering that could occur by random chance. I assign each of the bin locations a random value between 1-8 for the 8 possible relations of velocity and topography shown in Figure 6, then count how many bins with the same value were found within a radius of 20° . This process is performed 1000 times to compute the mean and standard deviation for the number of bins with the same value that would be expected if the values were random. For most locations, the mean value is 4-7 and σ is around 1.5. The maps in Figure 7 show the bin locations for each of the 8 cases separately where $+Vs$ = fast, $-Vs$ = slow, $-410/660$ = shallow, $+410/660$ = deep. The locations for which there are more nearby bins ($mean + 2\sigma$) with the same value compared to random are colored according to the scheme in Figure 6. The bins that are within 2σ are plotted as empty circles. A distance of 20° was chosen since it is small enough to represent regional patterns, but large enough to get good statistics in the poorly sampled regions.

The signals in seismic data are dominated by the effects of temperature in many regions, but there are very few regions hydrated enough to produce a dominant seismic signal. Cases 1-6 show

114 that the velocity-topography relationship can be statistically significant across large regions of the
 115 globe. Case 1 (fast shear velocity, shallow 410 km discontinuity, deep 660 km discontinuity) is
 116 found in subduction areas and is consistent with the expected signals from a cold thermal anomaly.
 117 Likewise, Case 6 with a low shear velocity, deep 410 km discontinuity, and shallow 660 km disconti-
 118 nuity (the signals expected for a vertically continuous warm thermal anomaly (Deuss et al., 2013))
 119 is statistically significant across the mid Pacific. Since the 660 km discontinuity might actually be
 120 depressed in hot regions due to the majorite phase transition (Hirose, 2002; Houser and Williams,
 121 2010), Case 4 with low shear velocity, deep 410 km discontinuity, and deep 660 km discontinuity
 122 may also indicate a temperature anomaly. Indeed, this pattern is observed to be statistically sig-
 123 nificant across the mid Pacific. Case 8 of Figure 7 shows that only a few spots around Antarctica
 124 and under Brazil have a signal for water that is greater than random. Although the eye detects
 125 a possible clustering in the Indian Ocean and New Guinea, this clustering does not appear to be
 126 statistically significant.

127 To calculate the upper-limit of water concentration in the bins with the water signal, I apply
 128 Equation 1 from Suetsugu et al. (2006), without the temperature dependent terms, to the seismic
 129 observations of transition zone topography and velocity.

$$\begin{aligned}\delta d &= \frac{\partial d}{\partial W} \delta W \\ \delta V_s &= \frac{\partial(\delta V_s)}{\partial W} \delta W\end{aligned}\tag{1}$$

130 where δd is the departure from the average depth (km) of the 410 or 660 km discontinuities
 131 shown in Figure 4, $\delta V_s\%$ is the observed shear velocity anomaly shown in Figure 5, and W is the
 132 water content (wt.%). For comparison with other studies, I choose the same values as Mao et al.
 133 (2012):

$$134 \frac{\partial(\delta V/V)}{\partial W} = -4.5 \text{ vel\%/wt.\% (Mao et al., 2012)}$$

$$135 \frac{\partial d_{410}}{\partial W} = -15 \text{ km/wt.\% (Smyth and Frost, 2002)}$$

$$136 \frac{\partial d_{660}}{\partial W} = 6 \text{ km/wt.\% (Higo et al., 2001)}$$

137 Using these values and the measured 410 km and 660 km discontinuity topography and shear
 138 velocity, I calculate the upper limit on the water content since I ignore the temperature-dependent
 139 terms. The distribution of the average water content measured in each of the 105 bins consistent
 140 with water is shown in the left side of Figure 8. The overall average value is 0.6 wt% water in

141 these portions of the mantle transition zone. This is slightly larger than the 0.1% reported by
 142 Mao et al. (2012), but expected since I do not factor in the effects of temperature. Suetsugu et al.
 143 (2006) obtain larger values for water content in subduction zones because they use a smaller value
 144 of $\frac{\partial d}{\partial W}$ (2.7 km/wt.%) and do not use the 410 km discontinuity topography in their calculations.
 145 The right hand side of Figure 8 overlays the water content values separately for each type of data;
 146 410 km discontinuity topography, 660 km discontinuity topography, and shear velocity anomaly.
 147 The very low (0-0.2 wt.%) values predominantly come from the velocity measurements, while the
 148 very high values (1.5-3.0 wt.%) come from the 660 km discontinuity topography measurements.
 149 Mao et al. (2012) report that the sensitivity of shear velocity to temperature increases at transition
 150 zone temperatures compared to room temperature. Since the shear velocity variations are about
 151 0.5% on average, the magnitude of the water anomaly is around 0.1 wt.%. Likewise, although the
 152 410 and 660 km discontinuities have about the same topography range, the 660 km discontinuity
 153 measurements lead to higher estimates of water content since the sensitivity is reported to be half
 154 that of the 410 km discontinuity.

155 **3. Discussion**

156 The global long period seismic data observations reported here are not compatible with the
 157 mineral physics predictions regarding the effects of water on transition zone minerals . Since most
 158 mantle shear velocity models are in general agreement, and most maps of discontinuity topography
 159 are in general agreement, the results shown here are not strongly dependent on the models. The
 160 seismic data presented here are sensitive to lateral variations in rock properties and do not directly
 161 measure the background level of hydration. If the transition zone is completely saturated or com-
 162 pletely devoid of water, then there would be no seismic signals from lateral changes in water content.
 163 Interpreting the seismic signals in terms of changes in water content applies in the intermediate
 164 case where there is some amount of water in the transition zone even if the background level is not
 165 known.

166 The sharpness of the 410 km discontinuity has been considered an indicator of water content
 167 since the phase transition interval will extend from 4 km in the dry case to up to 40 km with
 168 complete saturation (Wood, 1995; Helffrich and Wood, 1996; Smyth and Frost, 2002). Most short
 169 period studies find the 410 km discontinuity has a sharpness less than 10 km (Benz and Vidale,
 170 1993; Neele, 1996; Xu et al., 2003) which can be interpreted as evidence against hydrous wadsleyite.

171 However, the relative proportions of the initial and final phases within the phase loop is often not
172 linear, so a sharp seismic discontinuity can arise even from a broad phase transition (Stixrude,
173 1997). Hydrogen also diffuses quickly which can sharpen the boundary in areas where vertical flow
174 is slower than the diffusion rate (Smyth and Frost, 2002).

175 Recent seismic studies have had mixed results identifying water in the transition zone. Cam-
176 marano and Romanowicz (2007) find that their seismic observations are best explained by an upper
177 mantle that is faster and a transition zone that is slower than predicted for pyrolite. While the
178 reduced seismic velocities within the transition zone could be due to water, water does not explain
179 their upper mantle measurements, and they favor a gradual increase in the proportion of garnet
180 with depth. Meier et al. (2009) jointly invert for surface wave velocity and topography and see
181 different structure in the 410 km discontinuity topography than reported here. Since their 410
182 km discontinuity is elevated under the Pacific, they propose that the uplift and the slow velocity
183 there could be due to water. Their 410 km discontinuity topography is highly correlated with the
184 660 km discontinuity topography and an ocean-continent function, indicating that their data may
185 have difficulty resolving structure of the discontinuities. Their results do not seem compatible with
186 precursor data since the highest quality SS precursors are under the Pacific, with over 1000 traces
187 contributing to the stacks in this region, and a depressed 410 km discontinuity under the Pacific
188 is also observed in other precursor studies (Gu et al., 2003; Lawrence and Shearer, 2008; Deuss,
189 2009). For an overview of earlier seismic studies regarding water in the transition zone, see (Deuss
190 et al., 2013).

191 Globally, the 410 and 660 km discontinuities are not correlated with each other, rather than
192 anti-correlated as expected for both temperature or water (Houser et al., 2008a). If water is not
193 distributed evenly with depth in the transition zone, then perhaps it is collecting either in the
194 ringwoodite or the wadsleyite due to material flowing across the transition zone. In which case,
195 the 410 km discontinuity and the shear velocity should be consistent with water in upwelling
196 regions and likewise for the 660 km discontinuity and shear velocity in downwelling regions. The
197 strongest signals in the transition zone topography and velocities are a thick transition zone and
198 fast velocities in the circum-Pacific and a thin transition zone and slow velocities under the Pacific.
199 The correlation of the 410 km discontinuity topography and shear velocity should be stronger
200 than the 660 km discontinuity and shear velocity since water is more compatible in wadsleyite than
201 ringwoodite. However, the depressed 410 km discontinuity and slow shear velocity where the mantle

202 is flowing upwards under the Pacific plate is not consistent with water, but with temperature. The
 203 corresponding argument that the 660 km discontinuity and shear velocity should be correlated in
 204 downwellings is difficult to assess because of the strong temperature signal in a down going slab.
 205 While studies such as Litasov et al. (2005) say that 660 depth is more sensitive to water than
 206 temperature, such that a thick transition zone is wet, the fast velocities of the slab due to the
 207 cold temperatures could hide the signal from water. Thus, more regional studies are necessary
 208 to evaluate distribution and concentration of water in areas of active subduction (Suetsugu et al.,
 209 2006; Zheng et al., 2007).

210 While the global seismic observations presented here find that the mantle transition zone is
 211 generally dry, the recent observation of hydrous ringwoodite in diamonds from a kimberlite pipe
 212 in Brazil (Pearson et al., 2014) has been interpreted as evidence for close to 1 wt.% water in
 213 ringwoodite. Another interpretation is that water is not compatible at transition zone temperatures
 214 and pressures for long time scales and will result in melting instabilities that lead to the formation of
 215 such kimberlite pipes. Schmerr and Garnero (2007) provide a detailed examination of the transition
 216 zone beneath South America using SS precursors, but do not observe the same topography pattern
 217 under Brazil reported here. They find a slightly depressed 410 km discontinuity and a slightly
 218 elevated 660 km discontinuity. Their observations in the region of the slab suggest local dehydration
 219 of the slab and storage of water 10s of kilometers below the uplifted 410 km discontinuity, but no
 220 regional hydration of the entire transition zone. In addition, Lessing et al. (2014)’s observations
 221 of PP and SS precursors beneath India and western China, an area with 100s of millions of years
 222 of subduction of the oceanic Tethys and Indian plates (Stampfli and Borel, 2002), show no signal
 223 consistent with hydration of the transition zone.

224 The assertion here that the mantle transition zone is generally dry is compatible with recent
 225 results that water could be stable in some mineral structures allowing water to pass into the lower
 226 mantle. Although the hydrous phase H could be stable in the lower mantle at low temperatures
 227 (Tsuchiya, 2013), the stability field is expanded even to the core-mantle boundary for Al-rich phase
 228 H (Nishi et al., 2014). Similarly, Al-rich phase D has been shown to have a stability field that
 229 extends into the lower mantle (Pamato et al., 2014). Thus, these recent findings motivate questions
 230 regarding the properties and stability of hydrous phases and melt at lower mantle conditions.
 231 However, the stability of water in the upper mantle and transition zone is still not resolved (Green
 232 et al., 2010).

233 Smyth et al. (2004) propose that the magnitude of seismic velocity variations due to changes in
 234 water content in ringwoodite could be larger than those from temperature variations. However,
 235 the discontinuity structure is most consistent with water (elevated 410 and depressed 660 km
 236 discontinuity) in the circum-Pacific subduction zones, but the seismic velocities are usually fast,
 237 which is expected for a cold subducting slab. Likewise, a large region of slow seismic velocities exists
 238 under the Pacific as predicted for water, but there is a strong, high quality signal indicating that
 239 the 410 km discontinuity topography is depressed. Since the best quality discontinuity topography
 240 measurements are within the Pacific, the absence of bins consistent with water there is a strong
 241 argument against widespread or global storage of water in the transition zone. While it is possible
 242 that hydration produces seismic signals that are weaker or different than the current mineral physics
 243 data suggest, there is currently almost no evidence in the global seismic data that water from
 244 subduction or the deep mantle from the early Earth is slowly hydrating the mantle transition zone.

245 **4. Conclusion**

246 Examining the long period seismic observations of discontinuity topography from SS precursors
 247 along with shear velocity from a global shear velocity model, there are only a few, scattered locations
 248 consistent with the signal of water as predicted by mineral physics. The amount of water indicated
 249 is around 0.6 wt.% considering both the 410 and 660 km discontinuity topography and shear velocity
 250 in these locations. These results indicate that the mantle transition zone is generally dry, containing
 251 around one ocean of water.

252 **5. Acknowledgements**

253 This work was funded by the National Science Foundation EAR1114445 and the Earth-Life
 254 Science Institute at Tokyo Institute of Technology. The facilities of the Incorporated Research
 255 Institutions for Seismology (IRIS) Data Services, specifically the IRIS Data Management Center
 256 and the EarthScope USArray facility, were used for access to waveforms, related metadata, and/or
 257 derived products used in this study. IRIS Data Services are funded through the Seismological Facil-
 258 ities for the Advancement of Geoscience and EarthScope (SAGE) Proposal of the National Science
 259 Foundation under Cooperative Agreement EAR-1261681. Global Seismographic Network (GSN)
 260 is a cooperative scientific facility operated jointly by the Incorporated Research Institutions for

261 Seismology (IRIS), the United States Geological Survey (USGS), and the National Science Founda-
262 tion (NSF), under Cooperative Agreement EAR-1261681. The author would like to thank Nicholas
263 Guttenberg and members of the Deep Thoughts discussion group for exchanges that contributed
264 to this work.

- Angel, R., Frost, D., Ross, N., Hemley, R., 2001. Stabilities and equations of state of dense hydrous magnesium silicates. *Phys. Earth Planet. Int.* 127, 181–196.
- Bassin, C., Laske, G., Masters, G., 2000. The current limits of resolution for surface wave tomography in North America. *EOS* 81, 897.
- Benz, H., Vidale, J., 1993. The sharpness of upper mantle discontinuities determined by high-frequency P’P’ precursors. *Nature* 365, 147–150.
- Bolfan-Casanova, N., McCammon, C., Mackwell, S., 2006. Water in the transition zone and lower mantle minerals. In: Jacobsen, S., van der Lee, S. (Eds.), *Earth’s Deep Water Cycle*. Vol. 168. *Geophysical Monograph Series*, pp. 57–68.
- Cammarano, F., Romanowicz, B., 2007. Insights into the nature of the transition zone from physically constrained inversion of long-period seismic data. *Proc. Nat. Acad. Sci.* 104, 9139–9144.
- Deuss, A., 2009. Global observations of mantle discontinuities using *SS* and *PP* precursors. *Surv. Geophys.* 30, 301–326.
- Deuss, A., Andrews, J., Day, E., 2013. Seismic observations of mantle discontinuities and their mineralogical and dynamical interpretation. In: Karato, S. (Ed.), *Physics and Chemistry of the Deep Earth*. John Wiley and Sons, Ltd., pp. 297–323.
- Dixon, J., Leist, L., Langmuir, C., Schilling, J., 2002. Recycled dehydrated lithosphere observed in plume-influenced mid-ocean-ridge basalt. *Nature* 420, 385–389.
- Dziewonski, A., Anderson, D., 1981. Preliminary reference Earth model. *Phys. Earth Planet. Int.* 25, 297–356.
- Flanagan, M., Shearer, P., 1998. Global mapping of topography on transition zone discontinuities by stacking of *SS* precursors. *J. Geophys. Res.* 103, 2673–2692.
- Green, H., Chen, W.-P., Brudzinski, M., 2010. Seismic evidence of negligible water carried below 400-km depth in subducting lithosphere. *Nature* 467, 828–831.
- Gu, Y., Dziewonski, A., Ekstrom, G., 2003. Simultaneous inversion for mantle shear velocity and topography of transition zone discontinuities. *Geophys. J. Int.* 154, 559–583.

- Helfrich, G., 2002. Thermal variations in the mantle inferred from 660 km discontinuity topography and tomographic wave speed variations. *Geophys. J. Int.* 151, 935–943.
- Helfrich, G., Wood, B., 1996. 410 km discontinuity sharpness and the form of the olivine $\alpha - \beta$ phase diagram: resolution of apparent seismic contradictions. *Geophys. J. Int.* 126, F7–F12.
- Higo, Y., Inoue, T., Yurimoto, H., 2001. Effect of water on spinel-postspinel transformation in Mg_2SiO_4 . *Geophys. Res. Lett.* 28, 3505–3508.
- Hirose, K., 2002. Phase transitions in pyrolitic mantle around 670-km depth: Implications for upwelling of plumes from the lower mantle. *J. Geophys. Res.* 107, doi:10.1029/2001JB000597.
- Hirschmann, M., 2006. Water, melting, and the deep Earth H_2O cycle. *Annu. Rev. Earth Planet. Sci.* 34, 629–653.
- Houser, C., Masters, G., Flanagan, M., Shearer, P., 2008a. Determination and analysis of long-wavelength transition zone structure using *SS* precursors. *Geophys. J. Int.* 174, 178–194.
- Houser, C., Masters, G., Shearer, P., Laske, G., 2008b. Shear and compressional velocity models of the mantle from cluster analysis of long-period waveforms. *Geophys. J. Int.* 174, 195–212.
- Houser, C., Williams, Q., 2010. Reconciling Pacific 410 and 660 km discontinuity topography, transition zone shear velocity patterns, and mantle phase transitions. *Earth Planet. Sci. Lett.* 296, 255–266.
- Huang, X., Xu, Y., Karato, S., 2005. Water content in the transition zone from electrical conductivity of wadsleyite and ringwoodite. *Nature* 434, 746–749.
- Inoue, T., Tanimoto, Y., Irifune, T., Suzuki, T., Fukui, H., Ohtaka, O., 2004. Thermal expansion of wadsleyite, ringwoodite, hydrous wadsleyite and hydrous ringwoodite. *Phys. Earth Planet. Int.* 143–144, 279–290.
- Inoue, T., Wada, T., Sasaki, R., Yurimoto, H., 2010. Water partitioning in the Earth’s mantle. *Phys. Earth Planet. Int.* 183, 245–251.
- Inoue, T., Yurimoto, H., Kudoh, Y., 1995. Hydrous modified spinel, $Mg_{1.75}SiH_{0.5}O_4$, a new water reservoir in the mantle transition zone. *J. Geophys. Res.* 22, 117–120.

- Jacobsen, S., Smyth, J., 2006. Effect of water on the sound velocities of ringwoodite in the transition zone. In: Jacobsen, S., van der Lee, S. (Eds.), *Earth's Deep Water Cycle*. American Geophysical Union, pp. 131–145.
- Kohlstedt, D., Keppler, H., Rubie, D., 1996. Solubility of water in the α , β , and γ phases of $(\text{Mg,Fe})_2\text{SiO}_4$. *Contributions to Mineral Petrology* 123, 345–357.
- Lawrence, J., Shearer, P., 2008. Imaging mantle transition zone thickness with *SdS-SS* finite-frequency sensitivity kernels. *Geophys. J. Int.* 174, 143–158.
- Lessing, S., Thomas, C., Rost, S., Cobden, L., Dobson, D., 2014. Mantle transition zone structure beneath India and Western China from migration of PP and SS precursors. *Geophys. J. Int.* 197, 396–413.
- Litasov, K., Ohtani, E., Sano, A., Suzuki, A., Funakoshi, K., 2005. In situ X-ray diffraction study of post-spinel transformation in a peridotite mantle: Implication for the 660-km discontinuity. *Earth Planet. Sci. Lett.* 238, 311–328.
- Mao, Z., Jacobsen, S., Jian, F., Smyth, J., Holl, C., Duffy, T., 2008. Elasticity of hydrous wadsleyite to 12 GPa: Implications for Earth's transition zone. *Geophys. Res. Lett.* 35, L21305.
- Mao, Z., Lin, J., Jacobsen, S., Duffy, T., Chang, Y., Smyth, J., Frost, D., Hauri, E., Prakapenka, V., 2012. Sound velocities of hydrous ringwoodite to 16 GPa and 673 K. *Earth Planet. Sci. Lett.* 331–332, 112–119.
- Meier, U., Trampert, J., Curtis, A., 2009. Global variations of temperature and water content in the mantle transition zone from higher mode surface waves. *Earth Planet. Sci. Lett.* 282, 91–101.
- Neele, F., 1996. Sharp 400-km discontinuity from short-period P reflections. *Geophys. Res. Lett.* 23, 419–422.
- Nishi, M., Irifune, T., Tsuchiya, J., Tange, Y., Nishihara, Y., Fujino, K., Higo, Y., 2014. Stability of hydrous silicate at high pressures and water transport to the deep lower mantle. *Nature Geoscience* 7, 224–227.
- Ohtani, E., Mizobata, H., Yurimoto, H., 2000. Stability of dense hydrous magnesium silicate phases in the systems $\text{Mg}_2\text{SiO}_4\text{-H}_2\text{O}$ and $\text{MgSiO}_3\text{-H}_2\text{O}$ at pressures up to 27 GPa. *Physics and Chemistry of Minerals* 27, 533–544.

- Pamato, M., Myhill, R., Ballaran, T., Frost, D., Heidelbach, F., Miyajima, N., 2014. Lower-mantle water reservoir implied by the extreme stability of a hydrous aluminosilicate. *Nature Geoscience* 8, 75–79.
- Pearson, D., Brenker, F., Nestola, F., McNeill, J., Nasdala, L., Hutchison, M., Matveev, S., Mather, K., Siversmit, G., Schmitz, S., Vekemans, B., Vincze, L., 2014. Hydrous mantle transition zone indicated by ringwoodite included within diamond. *Nature* 507, 221–224.
- Ritsema, J., van Heijst, H., Woodhouse, J., 2004. Global transition zone tomography. *J. Geophys. Res.* 109, 10.1029/2003JB002610.
- Schmerr, N., Garnero, E., 2007. Upper mantle discontinuity topography from thermal and chemical heterogeneity. *Science* 318, 623–626.
- Smyth, J., Frost, D., 2002. The effect of water on the 410-km discontinuity: An experimental study. *Geophys. Res. Lett.* 29, doi:10.1029/2001GL014418.
- Smyth, J., Holl, C., Frost, D., Jacobsen, S., 2004. High pressure crystal chemistry of hydrous ringwoodite and water in the Earth’s interior. *Phys. Earth Planet. Int.* 143-144, 271–278.
- Stampfli, G., Borel, G., 2002. A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons. *Earth Planet. Sci. Lett.* 196, 17–33.
- Stixrude, L., 1997. Structure and sharpness of phase transitions and mantle discontinuities. *J. Geophys. Res.* 102, 14835–14852.
- Suetsugu, D., Inoue, T., Yamada, A., Zhao, D., Obayashi, M., 2006. Towards mapping the three-dimensional distribution of water in the transition zone from P-velocity tomography and 660-km discontinuity depths. In: Jacobsen, S., van der Lee, S. (Eds.), *Earth’s Deep Water Cycle*. Vol. 168. *Geophysical Monograph Series*, pp. 237–249.
- Thio, V., Cobden, L., Trampert, J., 2016. Seismic signature of a hydrous mantle transition zone. *Physics Earth. Planet. Int.* 250, 46–63.
- Tsuchiya, J., 2013. First principles prediction of a new high-pressure phase of dense hydrous magnesium silicates in the lower mantle. *Geophys. Res. Lett.* 40, 4570–4573.

- Tsuchiya, J., Tsuchiya, T., 2009. First principles investigation of the structural and elastic properties of hydrous wadsleyite under pressure. *J. Geophys. Res.* 114, doi:10.1029/2008JB005841.
- Wood, B., 1995. The effect of H_2O on the 410-kilometer seismic discontinuity. *Science* 268, 74–76.
- Xu, F., Vidale, J., Earle, P., 2003. Survey of precursors to P'P': Fine structure of mantle discontinuities. *J. Geophys. Res.* 108, doi:10.1029/2001JB000817.
- Zheng, Y., Lay, T., Flanagan, M., Williams, Q., 2007. Pervasive seismic wave reflectivity and metasomatism of the Tonga mantle wedge. *Science* 316, 855–859.

Figure 1: Cartoon representing the predicted behavior (see Jacobsen and Smyth (2006)) of transition zone phases in the presence of a downwelling wet slab or dry mantle upwelling colored red/blue because water reduces shear velocity (ignoring temperature). The schematic phase diagram shows the Clapeyron slopes of dry (black solid) and wet (grey dashed) olivine component of the mantle. Ol = olivine, Gt = garnet, Cpx = clinopyroxene, Wads = wadsleyite, Ri = ringwoodite, CaPv = calcium perovskite, Br = bridgmanite, and Fp = ferropericlase

Figure 2: Ray geometry of *SS* precursors, and an actual precursor stack used in this study from the NW Pacific including 1133 seismograms. There is a time gap between the main *SS* arrival and the precursors. The vertical lines at the pulse peak are the SdS arrival times and the small grey lines above the pulse represent the error in the pulse peak from the bootstrap random resampling.

Improved Stack Quality

Figure 3: Map of the increase in the number of traces per stack for this study using data through 2010 versus the previous Houser et al. (2008a) study which has data through 2005. Dark pink regions have significant improvement in the quality of the SS precursor stacks due to more traces contributing to the stack. The color scale saturates at 50 traces/stack, but stacks in the central Pacific have increases of over 500 traces/stack. Improvement occurs in the northern Pacific, southern Pacific, eastern Asia, Indian ocean, northern South America, and north Atlantic. Light blue regions have almost the same stack quality as the previous study.

Figure 4: Topography of the 410 and 660 km discontinuities and the resulting transition zone thickness where the topography is defined as the elevation above and the below the average of 411 km, 650 km, and 239 km respectively. Only bins for which the number of traces/stack is greater than 15 and the depth error is less than 20 km are shown which reduces the bin number from 1464 to 1301.

Shear Velocity at the Base of the Transition Zone

Figure 5: Shear velocity of the mantle transition zone from 530-660 km depth. The model is the same as HMSL-S06 (Houser et al., 2008b), but the tomographic inversion is performed with 6° instead of 4° equal-area blocks at the equator. This velocity model is compared to the topography plotted in Figure 4 to create the comparison map in Figure 6.

Figure 6: Maps of the western and eastern hemispheres where each circle represents the relative discontinuity topography from the global average in each 5° bin. The color-coding scheme distinguishes bins by the sign seismic velocity anomaly and the elevation of both discontinuities. The purple bins which are consistent with the predictions for water are shown magnified in the right column, but represent only 8% of the 1301 bins used for this analysis.

Figure 7: Maps highlighting where each of the 8 observed discontinuity topography - shear velocity relationship cases have more neighboring bins with the same value than expected from a random process. Bins for which there are more neighboring bins than random are filled using the same color-coding scheme as in Figure 6. Empty circles are bin locations for which the neighboring bins (within 20°) with the same value occur less than $2\sigma + \text{mean}$ times. $+Vs$ = fast, $-Vs$ = slow, $-410/660$ = shallow, $+410/660$ = deep. Case 8 (bottom right, purple) is the velocity/topography relationship that is expected for water in the transition zone. Cases 1 (cold) and 6 (warm) represent the expected relationship for velocity/topography with temperature. This analysis demonstrates that the temperature signal has much stronger regional trends than the water signal.

Figure 8: Left: The distribution of average water concentration values $(410_{value} + 660_{value} + V_{svalue})/3$ from each bin with a seismically consistent water signal. The overall average of these values is 0.6 wt.% water. Right: water concentration values from each type of data; 410 km discontinuity topography, 660 km discontinuity topography, and shear velocity anomaly.