Volatiles beneath mid-ocean ridges: deep melting, channelised transport, focusing, and metasomatism

Tobias Keller^{a,b,*}, Richard F. Katz^a, Marc M. Hirschmann^c

^a*University of Oxford, Oxford, UK.*

^b*Stanford University, Stanford CA, USA.*

^c*University of Minnesota, Minneapolis MN, USA.*

Abstract

Deep-Earth volatile cycles couple the mantle with near-surface reservoirs. Volatiles are emitted by volcanism and, in particular, from mid-ocean ridges, which are the most prolific source of basaltic volcanism. Estimates of volatile extraction from the asthenosphere beneath ridges typically rely on measurements of undegassed lavas combined with simple petrogenetic models of the mean degree of melting. Estimated volatile fluxes have large uncertainties; this is partly due to a poor understanding of how volatiles are transported by magma in the asthenosphere. Here, we assess the fate of mantle volatiles through numerical simulations of melting and melt transport at mid-ocean ridges. Our simulations are based on two-phase, magma/mantle dynamics theory coupled to an idealised thermodynamic model of mantle melting in the presence of water and carbon dioxide. We combine simulation results with catalogued observations of all ridge segments to estimate a range of likely volatile output from the global mid-ocean ridge system. We thus predict global MOR crust production of 66–73 Gt/yr (22–24 km³/yr) and global volatile output of 52–110 Mt/yr, corresponding to mantle volatile contents of 100–200 ppm. We find that volatile extraction is limited: up to half of deep, volatile-rich melt is not focused to the axis but is rather deposited along the LAB. As these distal melts crystallise and fractionate, they metasomatise the base of the lithosphere, creating rheological heterogeneity that could contribute to the seismic signature of the LAB.

Keywords: Mid-ocean ridge degassing, mantle melting, reactive channels, melt focusing, deep volatile cycles, magma/mantle dynamics

*Corresponding author

Email address: tobias.keller@stanford.edu (Tobias Keller)

1. Introduction

1.1. Deep volatile cycles

Deep-Earth volatile cycles are a result of fluxes between near-surface, shallow-mantle and deep-mantle reservoirs. These fluxes are associated with transport processes in the solid Earth that may be modified by the volatiles themselves (e.g., [Hirth and Kohlstedt, 1996](#); [Dasgupta and Hirschmann, 2010](#)). Volatile cycles impact aspects of planetary dynamics including mantle convection and partial melting. Partial melting is the key process by which volatiles are released from the mantle to near-surface reservoirs; it influences deep and shallow phenomena including the formation of the oceanic ([Hirth and Kohlstedt, 1996](#); [Asimow et al., 2004](#)) and continental crust ([Müntener et al., 2001](#); [Ulmer, 2001](#)), the long-term tectonic regime of a planet ([Mian and Tozer, 1990](#); [Regenauer-Lieb, 2001](#)) and, potentially, variations of the surface climate ([Huybers and Langmuir, 2009](#); [Crowley et al., 2015](#); [Burley and Katz, 2015](#); [Tolstoy, 2015](#); [Huybers and Langmuir, 2017](#)). The thermodynamic effect of mantle volatiles on partial melting and, in particular, that of water and carbon dioxide, is now well documented (e.g., [Hirschmann et al., 1999](#); [Asimow and Langmuir, 2003](#); [Dasgupta et al., 2013](#)). A key problem, however, remains almost unexplored: the dynamics of melt transport in the presence of volatiles. A theoretical framework to investigate these dynamics has recently been developed by [Keller and Katz \(2016\)](#).

Volatile-enriched partial melting and melt transport is relevant to all environments in which partial melting occurs; perhaps most of all to subduction zones, where melt production is primarily driven by the input of volatiles. In this manuscript, however, we focus on divergent plate margins, which are a simpler geodynamic context. The mid-ocean ridge (MOR) system is the primary avenue for volatile extraction from the mantle ([Resing et al., 2004](#); [Dasgupta and Hirschmann, 2010](#); [Kelemen and Manning, 2015](#)). A simple estimate of the volatile output from ridges is the product of the formation rate of oceanic crust ($\sim 24 \text{ km}^3/\text{yr}$ or 72 Gt/yr , [Crisp \(1984\)](#)) and the mean volatile concentration in undegassed primary mid-ocean-ridge basalt (MORB) (e.g., [Michael and Graham, 2015](#); [Rosenthal et al., 2015](#)). [Resing et al. \(2004\)](#) constrain the MOR CO_2 emission rate to a range of 22–88 $\text{Mt CO}_2/\text{yr}$, [Dasgupta and Hirschmann \(2010\)](#) produce estimates of 46–324, [Cartigny et al. \(2008\)](#) of 44–220, and [Kelemen and Manning \(2015\)](#) of 29–154 $\text{Mt CO}_2/\text{yr}$. Some of these include contributions from oceanic intraplate volcanism.

Such estimates are the basis for currently accepted volatile output from ridges, but bypass important considerations. For example, relating fluxes to the concentration of volatiles in the

sub-ridge mantle requires constraints on the locus of partial melting in the mantle as well as on the efficiency of volatile transport with ascending melts. Also, simple budget calculations cannot constrain the fraction of volatile-rich melts that are emplaced into the oceanic lithosphere rather than focused to the ridge axis. Any such emplaced melts may influence the properties of the oceanic lithosphere and contribute to volatile release during subduction. Moreover, mean flux estimates do not allow exploration of dynamic controls such as viscosity and permeability (Braun et al., 2000; Kono et al., 2014), channelised flow (Kelemen et al., 1995), and spatial variations in mantle potential temperature and composition (Dalton et al., 2014).

1.2. Volatiles in mid-ocean ridge magmatism

Small concentrations of volatiles greatly expand the volume of mantle that produces partial melt beneath ridges by causing melting to initiate at greater depths. For typical sub-ridge potential temperatures, volatile-free melting may occur above 60–75 km, but as little as 100 ppm H₂O depresses the onset of melting to 100–120 km depth (Hirth and Kohlstedt, 1996; Hirschmann et al., 1999). Comparable amounts of CO₂ lead to an onset of melting between 150 and 300 km depth, depending on the redox state of the mantle (Dasgupta and Hirschmann, 2010). Moreover, due to the widening of the upwelling region with depth (Lachenbruch, 1976; Spiegelman and McKenzie, 1987), deepening the melting regime produces small fractions of melt at considerable lateral distance away from the axis. These distal, low-degree melts are rich in incompatible trace elements as well as in volatiles. The proportion of distal melts that are focused to the ridge axis rather than frozen into growing oceanic lithosphere or erupted at off-axis seamounts is uncertain (Galer and O’Nions, 1986; Plank and Langmuir, 1992). Furthermore, erupted MORB has long been considered a mixture of polybaric melts, the composition of which is sensitive to the shape of the melting regime (O’Hara, 1985; Plank and Langmuir, 1992; Asimow and Langmuir, 2003). Lastly, the consequences of rheological weakening by volatiles are not yet well understood (Hirth and Kohlstedt, 1996; Braun et al., 2000). Resolving the dynamic aspects of MORB petrogenesis therefore requires an improved understanding of the reactive segregation of deep, volatile-rich melts.

Such deep melting produces melt fractions much less than 1% (e.g., Plank and Langmuir, 1992; Hirth and Kohlstedt, 1996; Asimow et al., 2004; Dasgupta and Hirschmann, 2006). Evidence from short-lived radiogenic products of uranium decay (²³⁰Th, ²³¹Pa, and ²²⁶Ra) indicates that very small melt fractions segregate from their deep sources and ascend rapidly to the surface

(Spiegelman and Elliott, 1993; McKenzie, 2000; Elliott and Spiegelman, 2003). This may be facilitated by high-permeability melt channels (Aharonov et al., 1995; Spiegelman et al., 2001). Keller and Katz (2016) showed that such channels may form when deep, volatile-rich melt fluxes the primary basaltic melting regime from below. Quantifying the consequences of incipient melting, channelised melt transport, and magmatic focusing on MOR volatile output is the goal of this study.

1.3. Method summary

We use two-dimensional numerical simulations of two-phase, multi-component, reactive flow in the asthenosphere to model magma genesis and transport beneath a MOR. The model is calibrated to reproduce accepted features of melting in an upwelling mantle containing H₂O and CO₂ in low concentrations. Crustal thickness is used as an observational constraint on MOR models (White et al., 2001); we generate predictions for comparison with data by computing the rate at which magma is delivered to the ridge axis. To scale our model results from individual instances to the ensemble of ridge segments that comprise the global MOR system, we use a catalogue of kinematic, chemical, and thermal parameters compiled by Gale et al. (2014) and Dalton et al. (2014). Empirical scaling laws fitted to the output of dynamic simulations are combined with this catalogued data to create estimates of crust production and volatile extraction for the global MOR system.

Section 2 introduces the computational model used to generate simulations and discusses the leading-order features and parametric controls of their output. Section 3 discusses melt focusing, crust production and volatile extraction predicted by simulations; section 4 develops estimates of global MOR volatile output; metasomatism of the oceanic lithosphere is considered in section 5. Section 6 summarises our findings and offers some conclusions.

2. Mid-ocean ridge simulations

2.1. Theory, parameters, and numerical solutions

Simulations are based on numerical solutions to a system of partial differential equations comprising statements of mass, momentum, and energy conservation for two phases (liquid and solid), and four thermochemical components. Solutions are computed on a two-dimensional domain representing half of a symmetrical spreading center. The domain extends to a depth of 200 km and

a width of 300 km from the axis. Model derivation and numerical implementation are given in Keller and Katz (2016); the theory is built upon that of McKenzie (1984) and Rudge et al. (2011).

Magma and rock compositions are modelled in a four-component compositional space of DUN + MORB + hMORB + cMORB. These chemical components represent the residue of mantle melting (dunite), the product of volatile-free decompression melting (basalt), and the products of hydrated and carbonated melting at depth (hydrated, carbonated basalt), respectively. The latter two components contain volatiles at fixed concentrations of 5 wt% H₂O in hMORB, and 20 wt% CO₂ in cMORB. Linear kinetic reaction rates are calculated with the R_DMC method (Keller and Katz, 2016, 2015), using a constant rate factor of $\mathcal{R} = 3 \text{ kg/m}^3/\text{yr}$ (corresponding to a reaction time of $\sim 1 \text{ kyr}$). Density is taken as constant and uniform for both phases, except in body-force terms where a density difference between melt and solid of $\Delta\rho = 500 \text{ kg/m}^3$ drives segregation. See Appendix A for a summary of the R_DMC model for thermodynamic equilibrium and melt-rock reactions.

The viscosity of mantle rock is reduced by water and melt (Hirth and Kohlstedt, 1996; Mei et al., 2002), and volatile-bearing silicate melt is weakened by both H₂O and CO₂. Permeability depends on grain size and melt fraction according to the Kozeny-Carman relation. For details of constitutive laws see Appendix B.

Plate spreading at a half-rate u_0 is imposed along the top boundary of the model domain; the bottom and off-axis side boundaries are open to inflow and outflow of the mantle in passive response to plate spreading. Symmetry conditions are applied along the vertical boundary beneath the ridge axis. Melt extraction at the axis occurs through an imposed patch of $4 \times 8 \text{ km}$ size where melting and freezing reactions are disallowed. For numerical convenience, a maximum solidus depth of 160 km is imposed to keep the lower boundary of the domain melt-free at all times.

Calculations are initialised with the temperature solution to the plate-cooling half-space problem. Some pre-depletion of the mantle composition is imposed in the melting region. The initial temperature is then limited point-wise by the solidus temperature such that the melt fraction is initially zero. These arrangements minimise the time required for model evolution from the initial condition to an approximately steady state (the “spin-up” time) (Katz, 2008, 2010).

Simulations at reference resolution of 1 km per grid cell have a numerical problem size of 744k degrees of freedom and are computed on 64 cores of the BRUTUS cluster at ETH Zurich, Switzerland. Numerical solutions are obtained using a parallel Newton-Krylov method leveraging

123 the PETSc toolkit (Balay et al., 2011, 2010; Katz et al., 2007).

124 2.2. Reference model

125 The reference model simulates an intermediate-spreading ridge with $u_0 = 3$ cm/yr, a mantle
126 potential temperature $T_m = 1350$ °C, and a mantle bulk composition $\bar{c}^i$ (in wt%) of 74.75 DUN,
127 25 MORB, 0.2 hMORB, and 0.05 cMORB. The concentrations of the latter two correspond to
128 volatile contents of 100 wt ppm each of H₂O and CO₂. A white-noise perturbation field is low-pass
129 filtered to a minimum wavelength of ~ 5 km and used to add compositional heterogeneity in mantle
130 fertility (MORB content) at a relative amplitude of $\pm 10\%$. The identical perturbation is used to
131 introduce mantle heterogeneity in all simulations reported here.

132 Figure 1 shows a snapshot of the reference model after 10 Myr. At this stage, plate spreading
133 and mantle upwelling have swept the full width and depth of the domain and thus dynamics have
134 evolved well past the effects of initial conditions. The general features of the magmatic system
135 are representative of all simulations presented here. A key feature is the coalescing network of
136 magmatic channels focusing melt towards the ridge axis. These channels are most notable across
137 a depth range of 50–80 km below the ridge axis and are characterised by increased melt fraction
138 (panel (a)), magnitude of Darcy flux $|\mathbf{q}| = \phi|\mathbf{v}_\ell - \mathbf{v}_s|$ (panel (b)), and volatile concentrations
139 (panels (c) & (d)).

140 The network is formed of reactive-dissolution channels as described in Keller and Katz (2016).
141 Channelisation is driven by the corrosivity of deep, low-degree, volatile-rich melts. These melts
142 flux the base of the volatile-free melting regime where they create a Reactive Infiltration Instability
143 (Aharonov et al., 1995; Szymczak and Ladd, 2014). Enhanced flux of deep melt leads to dissolution
144 of the MORB component, increased porosity and permeability, and further enhancement of melt
145 flux. Reactive channels thus form approximately parallel to the melt flow direction.

146 Figure 2 shows a comparison of the Darcy flux in simulations with volatile contents of 0, 50,
147 100 (ref. case), and 200 ppm. The distributed flow regime in the volatile-free case of panel (a)
148 indicates that the perturbation of the compositional field does not itself cause channelisation.
149 However, with increasing volatile content (panels (b)–(d)) the degree of flow localisation increases.
150 This trend reinforces the argument that the presence of volatile elements in the mantle source
151 causes channelisation. In our calculations, as little as 50 ppm each of H₂O and CO₂ is sufficient
152 to initiate channelling. This value is at the low end of estimates for the MORB-source mantle.

Figure 1: Output from reference simulation at 10 Myr. Panels show (a) melt fraction, with melt (red) and solid (blue) streamlines; (b) Darcy flux magnitude, with 600, 900 and 1200 °C isotherms; (c) bulk water concentration, with calibrated solidus depths for volatile-free, hydrated and carbonated mantle; (d) bulk carbon dioxide concentration, with contour of primary upwelling and melt producing domain (white) and melt streamline delimiting the melt focusing domain (orange), with equilibrium focusing distance x_e (dotted) for comparison.

2.3. Volatile extraction by melt focusing

Volatile-rich melt is produced over a broad region beneath the volatile-free melting regime. Extraction of these deep melts depends on the efficiency of melt focusing towards the ridge axis. Interpretation of observed volatile contents of MORBs therefore requires a model, whether implicit or explicit, of melt extraction from the melting regime. The canonical model of MORB petrogenesis analyses melting and melt focusing in terms of the residual mantle column (RMC) — the lateral solid outflow from the melting regime, after steady-state melt extraction (e.g., Klein and Langmuir, 1987; McKenzie and Bickle, 1988; Plank and Langmuir, 1992). Figure 3 juxtaposes this model (left-hand side) against a conceptual model derived from simulations presented here (right-hand side).

Figure 2: Darcy flux magnitude $|q|$ at $t = 10$ Myr for simulations with (a) no volatiles, (b) 50 ppm, (c) 100 ppm (reference), and (d) 200 ppm each H_2O and CO_2 in the mantle source. Streamlines for melt (red) and solid (blue) flow, outline of primary upwelling domain (black), and melt focusing domain (orange), with x_e (dotted) for comparison.

Plank and Langmuir (1992) and Asimow and Langmuir (2003) calculate volume and composition of extracted melt using the RMC (Fig. 3, left). All melt produced within a chosen focusing distance from the ridge is assumed to arrive at the axis, a concept referred to by the authors as “melt pooling.” Volume and composition of extracted melt are calculated as a weighted sum of melts produced at various pressures and degrees of melting (Langmuir et al., 1992). Although most of the calculations in Plank and Langmuir (1992) assume perfect focusing of distal melts, they recognised that “it is especially difficult to envision efficient extraction and focusing over hundreds of kilometers;” they discuss these difficulties in some detail. Asimow and Langmuir (2003) do not consider the limits of focusing; their calculations pool melts from the entire melting regime by integrating over the full depth of the RMC.

In the present calculations, melt focusing is not an assumption but rather a consequence of model dynamics. Careful analysis of results suggests that focusing arises by two mechanisms,

Figure 3: Summary of melt focusing beneath a MOR. Shaded areas show volatile-bearing (blue) and volatile-free (red) domains decompression melting; reactive channelling (purple); area of crystallisation and metasomatism (green). Dotted lines mark depths of first melting for volatile-bearing, z_{sol}^{vol} , volatile-free mantle, z_{sol}^{dry} ; lateral melt focusing distance x_f ; active focusing radius r_f . Melt (red) and solid (grey) flow paths are strongly simplified. Left hand side presents simplified standard model, compared to model presented in this study on the right hand side. The key difference is the fate of deep, volatile-rich melt.

represented schematically on the right-hand side of Fig 3 (c.f. Suppl. Figs S1–S3). Melt ascends by vertical porous flow driven by melt buoyancy. As it approaches the thermal boundary layer along the LAB, crystallisation and increasing rock viscosity prohibit further vertical ascent. Instead, melt is diverted towards the axis, continuing as gravity-driven flow in a decompaction channel along the LAB (Sparks and Parmentier, 1991; Montési et al., 2011). We refer to this first mechanism as passive focusing. The passive focusing distance x_f arises from balancing rates of melt supply from beneath and loss to freezing into the base of the spreading plate; it is modulated by the slope of the LAB (e.g., Katz, 2008; Hebert and Montési, 2010).

Under the second mechanism, plate spreading induces a dynamic pressure gradient that focuses melt transport toward the axis (Spiegelman and McKenzie, 1987); we refer to this as active focusing. It extends radially from the ridge to a radius r_f . Inside this radius, melt streamlines curve away from the thermal boundary on the ridge flanks, pointing down the dynamic pressure gradient

187 towards the axis. The focusing radius scales with the compaction length,

$$\delta_c = \sqrt{\zeta K / \mu}, \quad (1)$$

188 where K is the permeability, and η and ζ the shear and compaction viscosities, assuming $\zeta \gg \eta$.
189 In two-phase dynamics, pressure perturbations decay away from an applied forcing over a length
190 scale of approximately the compaction length (McKenzie, 1984). Depending on assumptions about
191 permeability, compaction viscosity and melt viscosity, and for melt fractions of 0.1–1 wt%, δ_c in
192 the asthenosphere may be of order 1–100 km (see Appendix B).

193 Conductive heat flux to the surface causes cooling along the flanks of the melting regime.
194 Some melt pathways will traverse these regions, where cooler temperatures cause deep fractional
195 crystallisation. As a consequence, volatiles (and other incompatibles) become more enriched in
196 these melts. Hence the volatile content of focused melts depends to some degree on their focusing
197 pathways, which in turn are controlled by the two focusing mechanisms above. Previous models
198 did not include these effects or their consequences for volatile extraction.

199 For parameter ranges considered here, the active focusing radius is typically smaller than the
200 passive focusing distance. Thus x_f marks the outer boundary of the melt focusing domain. The
201 melt streamline in Fig. 1(d) approximately marks the path of the most distal melt still focused
202 to the axis. We use the distance at which it intersects the base of the melting regime as a proxy
203 for x_f . Katz (2008) proposed an independent measure of x_f based on integrated mass fluxes; it is
204 termed the equilibrium focusing distance x_e (orange dotted, Fig. 1(d)). Both measures typically
205 produce consistent estimates. x_f correlates well with the lateral distance where the $u_0/3$ isopleth of
206 solid upwelling rate intersects the volatile-free solidus depth (white line in Fig. 1(d)). At reference
207 parameters, we find $x_f \approx 65$ km.

208 Model results suggest that the primary effects of volatiles — deeper onset of melting, a broader
209 melting regime and reactive channelisation — do not have any significant effect on melt focusing.
210 Thus, only volatile-rich melt produced in the central parts of the deep melting regime reaches
211 the MOR axis. This result differs sharply from assumptions of Asimow and Langmuir (2003) and
212 similar models. Furthermore, although volatiles cause channelisation of melt flow, this does not
213 affect melt focusing: channels emerge in alignment with the prior direction of melt transport. The
214 main effect of channelised melt transport is instead to increase the spatial and temporal variability
215 of melt flux and composition.

216 Melt focusing that is independent of deep, volatile-rich melting limits the efficiency of volatile
 217 extraction relative to previous estimates (e.g., [Asimow and Langmuir, 2003](#)). Basalt extracted at
 218 the axis is therefore dominated by low-pressure, high-degree melt. Most of the low-degree melt
 219 produced in the distal wings of the volatile-rich melting regime, conversely, migrates towards the
 220 LAB. There it is collected into lenticular bodies with melt fraction of 5–50 wt% at 20–60 km
 221 depth (Fig. 1(a)). These liquids metasomatise the base of the lithosphere. As they crystallise and
 222 fractionate, their volatile content increases while their freezing point decreases. We discuss these
 223 effects in section 5, below.

224 3. Quantifying volatile extraction

225 3.1. Melt production and focusing

226 To quantify production and focusing of melt in MOR simulation results we calculate the fol-
 227 lowing rates of mass transfer per unit length of MOR axis: q_{base}^i , the flow rate of component mass
 228 into the base of the melting regime ($z_{\text{sol}} = 160$ km); q_{melt}^i , the rate of component mass transfer by
 229 melting over the full melting regime; q_{focus}^i , the rate of component mass delivery to the ridge axis.
 230 Their calculation is discussed in Appendix C.

231 The ratios R_{melt}^i and R_{focus}^i are then introduced to quantify the mean degree of melting beneath
 232 the ridge and the efficiency of melt focusing to the axis. The melt production ratio R_{melt}^i is the
 233 time-averaged melt production rate divided by the time-averaged mantle inflow rate. The melt
 234 focusing ratio R_{focus}^i is the time-averaged melt focusing rate divided by the time-averaged melt
 235 production rate. These ratios are calculated for each component and for the bulk mantle as

$$R_{\text{melt}}^i = \frac{\langle q_{\text{melt}}^i \rangle}{\langle q_{\text{base}}^i \rangle}, \quad (2a)$$

$$R_{\text{focus}}^i = \frac{\langle q_{\text{focus}}^i \rangle}{\langle q_{\text{melt}}^i \rangle}, \quad (2b)$$

236 where $\langle \cdot \rangle$ denotes a time average.

237 Time-averaging is used to analyse the mean, long-term behaviour of the ridge apart from inter-
 238 nal fluctuations. The time interval over which averages are taken corresponds to plate spreading
 239 distances between $x_0 = 180$ km and the full width of the model domain $x_1 = W = 300$ km. By
 240 that interval, plate spreading and solid upwelling have swept once through the melt regime, and
 241 visual inspection suggests that transients associated with model spin-up have disappeared (see

Suppl. Figs S4–S6). Below, we discuss model output in terms of time-averaged rates $\langle q \rangle$ (and derived quantities) and their standard deviations σ across this post-spin-up time interval.

Figure 4: Magmatic ratios from (2) as a function of parameter values. Time-averaged melt production ratios for MORB (circles), hMORB (triangles), and bulk mantle (squares). Results are plotted as a function of: (a) half-spreading rate; (b) compaction length; (c) potential temperature; (d) fertility; (e) volatile content. Time-averaged melt focusing ratios for MORB (circles), and hMORB (triangles), and bulk mantle (squares). Bars indicate $\pm 2\sigma$. Dotted lines are regressions of the simulation output. Open symbols not included in regressions.

Figure 4 shows time-averaged melt production and focusing ratios plotted against model parameters including spreading rate, compaction length, mantle temperature, fertility, and volatile content. Parameter variations are applied one parameter at a time with other parameters held at reference values. Variations span a range of 1–7 cm/yr in half-spreading rate, 8–128 km in compaction length (see Appendix B), 1300–1400 °C in mantle temperature, 15–35 wt% in mantle fertility (MORB content), and 0–200 ppm in volatile content (H₂O and CO₂ added as hMORB and cMORB).

The bulk melt production ratio is $\sim 10\%$. This is comparable to the mean degree of melting discussed by Asimow and Langmuir (2003). R_{melt}^i correlates positively with mantle temperature and fertility but is largely independent of spreading rate (at least for slow to fast spreading ridges), compaction length and volatile content. We do not observe a lowering of R_{melt}^i with the addition of volatiles, as described for the mean degree of melting by Asimow and Langmuir (2003). This difference comes from normalising R_{melt}^i by mantle inflow at a fixed depth (160 km) rather than the volatile-dependent depth of first melting used by Asimow and Langmuir (2003).

The melt production ratios for the hMORB and MORB component illustrate the difference

between volatile-rich and volatile-free melt production. While the former shows degrees of melting of 0.85–0.95, the latter only reaches 0.25–0.4. The dissolution of the incompatible, water-rich component from the partially molten asthenosphere is therefore nearly complete in our models. Results for the carbonated component are similar to those for hMORB, and are therefore not shown in Fig. 4. The melts extracted at the ridge axis show no fractionation of water and carbon from their 1:1 source ratio. This is simply due to the high mean degree of melting that dominates the composition of extracted melts.

Plank and Langmuir (1992) considered focusing of distal, volatile-rich melts to be physically implausible. Here, we assess the efficiency of melt focusing and volatile extraction arising from dynamic simulations. Fig. 4(f)–(j) shows melt focusing ratios for the MORB and hMORB components. 80–100% of MORB melt is delivered to the axis, whereas only 40–70% of hydrated melt is focused. The remaining ~60% of dissolved water is transported towards the LAB and thus remains in the mantle. Volatile extraction efficiency is largely independent of spreading rate in panel (f) and of mantle fertility in panel (i). However, it decreases with increasing volatile content in panel (j). The addition of volatiles enhances deep melting in the distal parts of the melting regime, yet the focusing distance is unaffected. Thus, an increasing proportion of dissolved volatiles is not focused to the axis.

The striking anti-correlation between focusing of MORB and hMORB in Fig. 4(g)–(h) shows the control of melt extraction pathways on extracted melt compositions. With increasing compaction length, active focusing becomes more important relative to passive focusing. Melt streamlines curve away from the vertical toward the ridge axis at greater depth (c.f. Suppl. Fig. S2). Melt is transported away from the thermal boundary layer and hence interaction with the cooler lithosphere is minimised. Active focusing thus leads to less volatile enrichment by fractional crystallisation as compared to passive focusing. Less enrichment is also expected for other incompatible elements in the melt. Some of the compositional diversity of extracted melt may thus be attributed to different pathways of melt extraction, rather than to variations in source composition or degree of melting.

3.2. Crustal thickness

Predicted crustal thickness is an important model metric that can be compared with observations (see Appendix C). Figure 5(a)–(e) shows the time averaged crustal thickness $\langle H_c \rangle$ plotted against the same parameters as in Fig. 4 (full time-series of H_c in Suppl. Fig. S4). Crustal thickness as a function of spreading rate shows the characteristic saturation towards fast spreading ridges.

For comparison, MOR crustal thickness data obtained from seismics and rare-Earth-element in-
versions (White et al., 2001) are plotted alongside simulation results. Motivated by the variation
with spreading rate, we fit the simulation output and observational data with a log-cubic functions.
The results are shown as dotted and dashed lines, respectively, in panel (a). Qualitatively, the
simulation results follow the same trend as the observational data. However, our reference model
result of ~ 12 km crustal thickness is considerably higher than the observed ~ 7 km. We will return
to this point below.

Figure 5: Time-averaged crustal thickness $\langle H_c \rangle$ and volatile extraction rate $\langle q_{\text{focus}}^{\text{vol}} \rangle$ as function of parameter values. Results for $\langle H_c \rangle$ (circles) are plotted as a function of: (a) half-spreading rate; (b) compaction length; (c) potential temperature; (d) fertility; (e) volatile content. Bars indicate $\pm 2\sigma$. Dots in (a) are data from White et al. (2001). (f)–(j) show $\langle q_{\text{focus}}^{\text{vol}} \rangle$ for H_2O (squares), and CO_2 (triangles) as function of same parameter variations. Dotted lines are regressions of the simulation output; dashed lines are adjusted regressions to fit crustal thickness data; open symbols not included in regressions.

Crustal thickness is shown to correlate with compaction length in Figure 5(b). This is a consequence of more efficient melt focussing with increasing compaction length. A log-linear regression of the simulation output shows that a doubling of compaction length leads to a ~ 0.7 km increase in crustal thickness. Crustal thickness also increases significantly with higher potential temperature in panel (c), and mantle fertility in panel (d). A linear regression of the results with temperature and a quadratic regression with fertility show that 1 km of additional crust are produced by a ~ 16 $^\circ\text{C}$ increase in T_m or by a ~ 1.5 wt% increase in $\bar{c}^{\text{MORB}}$, holding everything else constant.

Whereas these trends follow expectations from previous analyses, the absolute melt production in our models is above observational constraints. We use the regression from panel (d) to calculate

the fertility reduction required to shift the fertility in the reference model down such that the misfit with observational constraints is reduced. The required shift in $\bar{c}^{\text{MORB}}$ is -6 wt%. To check this shift, we include an additional simulation with the MORB component concentration reduced to 19 wt%. The result is shown as open circles in panels (a)–(e); as expected, the adjusted crustal thickness is ~ 7 km. Adjusted regression curves for other parameters are shown as dashed lines.

The sensitivity of crustal thickness to mantle volatile content is relatively small: 1 km of additional crust is produced by an increase of ~ 140 wt ppm each of H_2O and CO_2 (more than doubling the reference value). In contrast, the temporal variability of crustal thickness is highly sensitive to volatile content. It follows that variability of magma supply at the ridge axis is not necessarily diagnostic of source heterogeneity in mantle fertility or volatiles — all simulations are seeded with the identical mantle heterogeneity. Rather, the temporal variability seen here is a result of channelised melt transport and its associated intermittency in space and time.

3.3. Volatile extraction

Rates of H_2O and CO_2 extraction by focused magma at the axis are given by $q_{\text{focus}}^{\text{H}_2\text{O}}$, and $q_{\text{focus}}^{\text{CO}_2}$ (see Appendix C). For H_2O , the extraction rate is interpreted as water supplied to the oceanic crust. Most of this water will remain contained within the crust but most of the CO_2 focused to the ridge is degassed from the shallow magmatic system into the ocean–atmosphere reservoir.

Fig. 5(f)–(j) shows time-averaged volatile extraction rates plotted against mantle parameters (full time-series of extraction rates in Suppl. Figs S5–S6). Under most conditions, H_2O and CO_2 extraction rates are nearly identical. As discussed above, volatile dilution by shallow, high-degree melting precludes their fractionation. Furthermore we find that the instances where the behaviour of CO_2 and H_2O differ are best explained by errors associated with insufficient numerical resolution. Simulations with $\delta_c < 32$ km have significant mass conservation errors for the incompatible components. Hence cMORB is more strongly affected while hMORB remains mostly well resolved. MORB and dunite components are not affected (see Suppl. Figs S5(f) & S6(f) for resolution test results). We therefore proceed with discussion of the hMORB component and assert that these results are a meaningful proxy for CO_2 extraction.

Volatile extraction rates increase linearly with half-spreading rate in panel (f). At reference parameters the extraction rate is ~ 1200 kg/m/yr. In the model with shifted reference fertility it is reduced to ~ 950 kg/m/yr (open square). A linear regression through the origin and this shifted reference rate (dashed line) has an increase in volatile extraction rate of ~ 320 kg/m/yr for every

1 cm/yr increase in u_0 . This sensitivity is comparable to idealised models of MOR carbon degassing by Burley and Katz (2015). The correlation with spreading rate is the result of increased melt production, while volatile concentration in the magma remains relatively stable (c.f. time-averaged melt compositions in Suppl. Fig. S7).

The decrease of extraction rates with increasing compaction length in panel (g) is, again, associated with the shift toward more active focussing. As melt is diverted away from the sub-lithospheric crystallisation front, deep fractional crystallisation and the consequent enrichment of incompatibles is reduced. Results at small δ_c suggest insufficient numerical resolution. A log-linear regression of the well-resolved results at higher δ_c has a drop in volatile extraction rate of ~ 110 kg/m/yr for a doubling of δ_c . An increase in mantle temperature in panel (h) and fertility in panel (i) both lead to a moderate increase in volatile extraction, as the deepening and widening of the volatile-free melting regime allows some more distal volatile-rich melt to be focused. Linear regressions of these results show that a rate increase of 100 kg/m/yr is produced by either a temperature increase of ~ 14 °C or an additional ~ 3.4 wt% mantle fertility.

Volatile extraction at the ridge axis is proportional to the mantle volatile content in panel (j). Regression through the origin and the data point from the adjusted fertility reference simulation shows that an additional 10 wt ppm volatiles in the mantle gives an increase of ~ 95 kg/m/yr in ridge extraction rate. The signature of increased channelised transport at higher volatile contents is expressed by higher variability of extraction rates. The relative amplitude of these variations is up to 50%. Again, this variability is not caused by source heterogeneity or variable degrees of melting. Rather it is the signature of reactive melt transport in the asthenosphere.

4. Simulation-based estimates of global MOR volatile output

4.1. Distribution along the global MOR system

The simulations presented here can provide a novel perspective on the globally integrated MOR output. Using the regressions of simulation output in Fig. 5 we interpolate crustal thickness and volatile extraction rate to conditions found along the global MOR system. However, in reading the results, one should bear in mind the limitations of this method: each regression curve was obtained for an isolated parameter variation in a complex, non-linear system; the natural system may be characterised by significant covariation between parameters. Despite this limitation, we expect to gain insights into possible global patterns as well as globally integrated magnitudes of

367 MOR output.

368 Conditions along the global MOR system are available from a catalogue of ridge segments
369 composed by Gale et al. (2014), which gives the location, length, and spreading rate of each
370 segment. Dalton et al. (2014) obtained estimates of potential temperature beneath segments on
371 the basis of inversion of observed ridge depth, S-wave speed, and MORB geochemistry. We use
372 those estimates to assign mantle temperature for each ridge segment in the Gale et al. (2014)
373 catalogue. We shift the temperatures by -25 °C to align their average temperature to our reference
374 temperature of 1350 °C. For ridge segments not included in Dalton et al. (2014), we interpolate
375 between the nearest available constraint and 1350 °C. Figure 6(a)–(b) shows half- spreading rate
376 and mantle potential temperature plotted on the mid-point coordinates of each ridge segment in
377 the combined catalogue.

378 Global average values for mantle fertility, volatile content and compaction length are approx-
379 imately known. Regions of increased volatile content are typically linked to nearby hotspots
380 (e.g., Azores (Asimow et al., 2004)). Yet, emerging information about these variations along the
381 ridge system is still sparse (Le Voyer et al., 2017). Therefore, we will investigate the range of
382 $100 \leq \bar{c}^{\text{H}_2\text{O}} \leq 200$ ppm, along with reference values for $\delta_c = 32$ km, $\bar{c}^{\text{MORB}} = 19$ wt% (the adjusted
383 value). We can now use the adjusted regressions in Fig. 5 to compute crustal thickness and volatile
384 extraction rate for each segment in the catalogue.

385 The predicted global distribution for $\bar{c}^{\text{H}_2\text{O}} = 100$ ppm, is shown in Fig. 6(c)–(d). The pattern
386 of crustal thickness in panel (c) mostly follows the pattern of mantle temperature from Dalton
387 et al. (2014). The thickest crust is found in regions influenced by hot-spots, most pronounced
388 beneath Iceland. There crustal thickness exceeds 12 km. The global average crustal thickness is
389 6.67 km. The lowest values are found at the eastern end of the Southwest Indian Ridge, at < 2 km.
390 The distribution of volatile extraction rate, in contrast, shows a strong dependence on spreading
391 rate, with only a weak influence of mantle temperature. The highest volatile extraction rates of
392 > 2.5 t/m/yr occur along the East Pacific Rise. The global average rate is 0.87 t/m/yr. For
393 cool, slow-spreading ridge sections such as parts of the Mid-Atlantic and Southwest Indian Ridges,
394 values are < 0.25 t/m/yr.

395 4.2. Range of global MOR output

396 Globally integrated MOR output can be obtained from the distributions above. For crust pro-
397 duction, we multiply crustal thickness by $2u_0\rho_c$ and integrate over the length of ridge segments.

Figure 6: MOR magmatic flux estimates for global ridge system with $\bar{c}^{\text{H}_2\text{O}} = 100$ ppm: (a) Half-spreading rates taken from [Gale et al. \(2014\)](#); (b) Mantle potential temperature extrapolated from [Dalton et al. \(2014\)](#); (c) Crustal thickness, and (d) volatile extraction rate calculated from regressions of simulation outputs. Each data point represents one ridge segment in [Gale et al. \(2014\)](#) catalogue.

398 The resulting projection of global crust production is 66.1 Gt/yr— equivalent to 22 km³/yr. Sim-
 399 ilarly, integrating the volatile extraction rate over the length of ridge segments after [Burley and](#)
 400 [Katz \(2015\)](#) predicts the global MOR volatile output to be 52.7 Mt/yr of H₂O and CO₂ each
 401 for reference volatile content. Repeating the global calculation above for a mean mantle volatile
 402 content of 200 ppm gives a global volatile output of 110.2 Mt/yr. The corresponding global crust
 403 production is 73.1 Gt/yr (24 km³/yr). The average volatile concentration in (undegassed, un-
 404 differentiated) focused melt is calculated by dividing the global volatile extraction by the crust
 405 production rate. The calculated values are 0.08 and 0.15 wt% for 100 and 200 ppm H₂O and CO₂
 406 in the mantle. According to our calculations, measured H₂O (0.13–0.77 wt%) and reconstructed

Figure 7: Globally integrated MOR magmatic flux estimates. Parameter distributions (line) and example histogram (bars) for segment-wise sampling of (a) compaction length, (b) mantle fertility, and (c), volatile content. Crustal thickness (d)–(f), and volatile extraction rate (g)–(h) calculated from fitting functions for example parameter distribution. [White et al. \(2001\)](#) data points for comparison (black); Lines show adjusted fitting functions for crustal thickness with best fit parameters (solid), and ± 2 km variation (dotted), and for extraction rate with best fit parameters (solid) and ± 0.5 t/m yr variation (dotted). Normal distributions fitted to population of globally integrated output from 1383 sampling instances (j)–(k). Colours denote individually sampled problem parameters.

CO₂ (0.066–5.75 wt%) concentrations in MORB by [Cartigny et al. \(2008\)](#) would map back to global mean concentrations of >170 ppm H₂O and >80 ppm CO₂; the upper limits would map to values well above 1000 ppm.

In addition to these leading-order global figures, we estimate the uncertainty of global MOR output arising from uncertainty in geographic, segment-wise variations of parameters that are not listed in the above catalogue. To this end we assume that mantle fertility, volatile content and compaction length are not constant in the upper mantle, but are distributed over a plausible range. Black curves in Figure 7 (a)–(c) show distributions of compaction length, fertility, and volatile content that are proposed to represent variation in the asthenosphere beneath the ridge system. Each distribution is centred around the reference values above. We assume compaction length has a log-normal distribution across a range of ~ 10 – 100 km ($\mu_n \pm 2\sigma_n$); fertility is normally distributed with a range of ~ 16 – 22 wt%; volatile content is normally distributed with a range of ~ 20 – 180 wt ppm (truncated at zero).

A population of instances of global MOR output is produced by repeating the following three steps. First, one of the three parameters is randomly sampled according to its distribution to obtain a value for each ridge segment. A histogram of a sample is shown for each parameter in Fig. 7(a)–(c). Second, segment-wise crustal thickness and volatile extraction rate are computed from regressions as above. Calculated values for sampled parameter sets in panels (a)–(c) are plotted against spreading rate in panels (d)–(i). Third, instances of globally integrated MOR output are calculated by integrating over the length of ridge segments. These three steps are repeated separately for each of these three parameters until each parameter-specific population of instances is fitted by a normal distribution to a relative tolerance of 10^{-3} .

The populations for crust production and volatile output obtained from $\gtrsim 10^3$ instances of statistically sampled parameter sets are shown in Fig. 7(j)–(k). The ranges of outcomes are shown as $\mu_n \pm 2\sigma_n$. Global crustal production is most sensitive to mantle fertility variations with a range of 65–68 Gt/yr. Global volatile extraction rate is most sensitive to mantle volatile content with a range of 52–56 Mt/yr. Variations in compaction length and mantle fertility have second-order effects on the range of volatile output.

These projected values of global volatile export from the mantle fall within the range of previous estimates, particularly in comparison to previous estimates of CO₂ emissions from MOR degassing ([Resing et al., 2004](#); [Cartigny et al., 2008](#); [Dasgupta and Hirschmann, 2010](#); [Kelemen](#)

and Manning, 2015). However, limitations of the present method should moderate our confidence in the produced estimates of MOR volatile output. The validity of the physical model and its numerical implementation is limited. For example, a more realistic rheology depending on stress and dynamically evolving grain size could have important effects on melt segregation (Turner et al., 2015); the model of mantle melting is based on simplified thermodynamics that are, at best, a crude approximation of full mantle petrogenesis; and the simulations skirt the limits of spatial resolution, introducing potentially disruptive mass conservations errors for incompatible species. Furthermore, using single-parameter regressions of simulation output neglects possible non-linear effects of parameter co-variations; polynomial degrees for regressions were chosen for convenience, not consistency with physical processes.

Nevertheless, the present method produces estimates of MOR volatile output that are consistent with dynamic simulations taking into account conservation of mass, momentum and energy. The models are calibrated to reproduce accepted features of MORB petrogenesis to leading order. Model parameters are tuned with the most recent observational constraints on the global MOR system, where available. The present estimates of global MOR volatile output are consistent with available constraints on extraction processes. Our method estimates MOR output for a range of global mean mantle properties and it quantifies the uncertainty associated with plausible regional distributions around reference conditions.

5. Deep volatiles and the LAB

From the above it is evident that not all melt produced beneath a MOR is focused to the ridge axis. Instead, a significant fraction of the deep, volatile-rich melt is transported towards the oceanic LAB. As seen in Fig. 1, this transport creates remnants of reactive channels bent away from the axis through plate motion. Over time such channels become stagnant sub-horizontal melt bodies stacked along the LAB. This feature of our results could help explain the sharp negative shear-wave velocity contrasts identified along the oceanic LAB in recent studies (Kawakatsu et al., 2009; Schmerr, 2012; Stern et al., 2015). We find that such melt lenses, which contain melt fractions of 5–25% (some up to 50%), are typically located along the 1200 °C isotherm. Some of this stalling melt could be further extracted by dike propagation (e.g. Havlin et al., 2013), an effect not currently included in our models. With ongoing crystallisation, volatiles are enriched in the fractionated liquid; the oceanic LAB is metasomatised by these volatile-rich liquids. In

contrast to simpler models (e.g. Plank and Langmuir, 1992; Hirth and Kohlstedt, 1996; Asimow and Langmuir, 2003), the volatile content with depth beneath the ocean floor (i.e., the RMC) does not reflect either a fractional or batch melting residue. Instead, the depth profile of volatiles is a result of complex transport processes. In particular, channelised melt transport leads to a heterogeneous distribution of volatiles along the LAB as it moves away from the ridge axis.

Figure 8: Averaged depth profiles of volatile and melt content, and rheology along far flank of ridge. Depth profiles obtained by horizontal averaging over interval from 200–300 km distance from axis, and triangular moving average along profile. Water and carbon dioxide bulk concentrations (a) & (b); melt fraction (c); rock viscosity (d); melt viscosity (e). Given for simulations with 0 (black), 50 (blue), 100 (red), 200 (light blue) wt ppm initial volatile content. Where no melt present, melt viscosity is found from fractional melt composition.

Figure 8 shows depth profiles produced by horizontally averaging simulation results in the interval from 180 to 300 km from the axis (6–10 Myr plate age). These profiles are then filtered with a vertical moving-average to emphasise structures larger than $\gtrsim 5$ km. Figure 8 shows results from simulations with various volatile contents. Moving upwards from 160 to 60 km depth, both H_2O and CO_2 follow a trend of depletion. Above is found an enriched region from 60 to 20 km depth. The enrichment is layered and reaches concentrations up to 1000 ppm; it is stronger for CO_2 than for H_2O , which indicates fractional crystallisation of accumulated low-degree melts. This layered band of metasomatised mantle coincides with the top boundary of the partially molten domain. Below 60 km depth, melt is stable at low fractions of 0.01–0.1%. Along the thermal LAB at ~ 60 –50 km depth, melt is accumulated into decompaction layers with horizontal averages upwards of 1% melt. Some remains of crystallising melt are found to a depth of 30 km. It is

484 conceivable that such melt layers are a source of off-axis seamount volcanism.

485 Volatile enrichment along the LAB has consequences for rock viscosity, as shown in Fig. 8(d).
486 Viscosity is reduced by water; a gradual increase in viscosity above the base of the melting regime
487 reflects volatile depletion by melt extraction. Along the band of metasomatism, however, rehy-
488 dration of the solid creates layers with up to an order of magnitude reduction of viscosity. These
489 represent a structural heterogeneity at the base of the lithosphere. Such rheological structure
490 could help explain the sharp decrease in shear- wave speed observed along the oceanic LAB (see
491 [Olugboji et al., 2016](#), and references therein). Our results support the current hypothesis that
492 these signals reflect the presence of partial melt and/or hydrated material along the LAB. Fur-
493 thermore, a metasomatised oceanic lithosphere has recently been evoked as the potential source of
494 enriched alkaline ocean island basalts ([Workman et al., 2004](#); [Pilet et al., 2011](#)) and highly alkaline
495 petit-spot volcanism ([Hirano, 2011](#)).

496 Lastly, the depth profiles of magma viscosity in Fig. 8(e) show a clear signal of lubrication by
497 volatile enrichment. The signal is seen in deep primitive as well as in shallow fractionated melt.
498 This viscosity variation promotes melt segregation from the mantle residue even at small melt
499 fractions < 0.1 wt%. An appropriate liquid viscosity model with volatile content is thus crucial
500 for understanding the transport of volatiles in the mantle beneath mid-ocean ridges.

501 6. Summary and conclusions

502 We present dynamic models of thermochemically coupled magma/mantle dynamics with volatiles
503 beneath a mid-ocean ridge. The presence of volatiles in low concentrations in the mantle source
504 causes deep, low-degree, volatile-rich melting. Due to the corrosive effect of volatile-rich melt
505 on silicates with decreasing pressure, the flux of deep melt into the volatile-free melting regime
506 triggers reactive channelisation. We find that melt focussing to the axis is not extended by chan-
507 nelled flow. Rather, channelling introduces spatial and temporal heterogeneity in flux and melt
508 composition at the ridge axis.

509 We investigate volatile extraction from the asthenosphere across a range of parameters including
510 spreading rate, compaction length, mantle temperature, fertility and volatile content. Volatile
511 extraction depends on the rate of melt delivery to the axis, as well as the concentration of volatiles
512 therein. The efficiency of melt focusing increases with compaction length, mantle temperature and
513 fertility. The concentration of volatiles in that melt depends on the melt extraction pathways and,

514 in particular, whether they approach the thermal boundary layer along the LAB, where fractional
515 crystallisation leads to volatile enrichment.

516 We calculate crustal thickness and volatile extraction rates for individual ridge segments in a
517 global catalogue using regressions of simulation outputs and crustal thickness data. The resulting
518 distribution of projected crustal thickness is mainly controlled by mantle temperature, whereas the
519 distribution of volatile extraction rate dominantly reflects spreading rate. Global integration gives
520 values of 66–73 Gt/yr ($22\text{--}24\text{ km}^3/\text{yr}$) for global crust production and a global volatile output of
521 53–110 Mt/yr, corresponding to mantle volatile contents of 100–200 ppm. The uncertainty of these
522 estimates relating to the uncertain distribution of mantle conditions along the global ridge system
523 is assessed using statistical sampling from plausible distributions of compaction length, mantle
524 fertility and volatile content. The resulting range of uncertainty for parameters distributed around
525 reference mantle conditions is 65–68 Gt/yr for crust production and 50–55 Mt/yr for volatile
526 output.

527 Finally, and in contrast to [Asimow and Langmuir \(2003\)](#), our models suggest that up to half of
528 dissolved volatiles that cross into the melting regime are not focused to the axis, but are instead
529 extracted towards the flanks of the ridge, where they metasomatise the mantle along the LAB.
530 The resulting rheological heterogeneity caused by water-weakening of mantle rock could explain
531 seismic signals interpreted as evidence of the oceanic LAB depth. Our results show bands of
532 metasomatised rock at a depth range of 20–60 km at spreading ages of 6–10 Myr, consistent with
533 LAB-related signals in seismic studies.

534 *Acknowledgements*

535 The research leading to these results has received funding from the European Research Council
536 under the European Union’s Seventh Framework Programme (FP7/2007–2013)/ERC grant agree-
537 ment number 279925. Hirschmann acknowledges support from NSF grant EAR1426772. The
538 authors thank the Isaac Newton Institute for Mathematical Sciences for its hospitality during the
539 programme Melt in the Mantle which was supported by EPSRC Grant Number EP/K032208/1.
540 Katz is grateful for the support of the Leverhulme Trust. The authors further thank the Geophysi-
541 cal Fluid Dynamics group for access to the BRUTUS cluster at ETH Zurich, Switzerland. Thanks
542 to R. White, C. Dalton, and A. Gale for making their data available.

References

- Aharonov, E., Whitehead, J.A., Kelemen, P.B., Spiegelman, M., 1995. Channeling instability of upwelling melt in the mantle. *J Geophys Res* 100, 20433–20450. doi:[10.1029/95JB01307](https://doi.org/10.1029/95JB01307).
- Asimow, P.D., Dixon, J.E., Langmuir, C.H., 2004. A hydrous melting and fractionation model for midocean ridge basalts: Application to the MidAtlantic Ridge near the Azores. *Geochem Geophys Geosyst* 5. doi:[10.1029/2003GC000568](https://doi.org/10.1029/2003GC000568).
- Asimow, P.D., Langmuir, C.H., 2003. The importance of water to oceanic mantle melting regimes. *Nature* 421, 815–820. doi:[10.1038/nature01429](https://doi.org/10.1038/nature01429).
- Balay, S., Brown, J., Buschelman, K., Eijkhout, V., Gropp, W.D., Kaushik, D., Knepley, M.G., McInnes, L.C., Smith, B.F., Zhang, H., 2010. PETSc Users Manual. Technical Report ANL-95/11 - Revision 3.4. Argonne National Laboratory.
- Balay, S., Brown, J., Buschelman, K., Gropp, W.D., Kaushik, D., Knepley, M.G., McInnes, L.C., Smith, B.F., Zhang, H., 2011. PETSc Web page. URL: <http://www.mcs.anl.gov/petsc>.
- Braun, M., Hirth, G., Parmentier, E.M., 2000. The effects of deep damp melting on mantle flow and melt generation beneath mid-ocean ridges. *Earth Planet Sci Lett* 176, 339–356. doi:[10.1016/S0012-821X\(00\)00015-7](https://doi.org/10.1016/S0012-821X(00)00015-7).
- Burley, J.M.A., Katz, R.F., 2015. Variations in mid-ocean ridge CO₂ emissions driven by glacial cycles. *Earth Planet Sci Lett* 426, 246–258. doi:[10.1016/j.epsl.2015.06.031](https://doi.org/10.1016/j.epsl.2015.06.031).
- Cartigny, P., Pineau, F., Aubaud, C., Javoy, M., 2008. Towards a consistent mantle carbon flux estimate: Insights from volatile systematics (H₂O/Ce, δ D, CO₂/Nb) in the North Atlantic mantle (14 N and 34 N). *Earth Planet Sci Lett* 265, 672–685. doi:[10.1016/j.epsl.2007.11.011](https://doi.org/10.1016/j.epsl.2007.11.011).
- Crisp, J.A., 1984. Rates of magma emplacement and volcanic output. *J Volcan Geotherm Res* 20, 177–211. doi:[10.1016/0377-0273\(84\)90039-8](https://doi.org/10.1016/0377-0273(84)90039-8).
- Crowley, J.W., Katz, R.F., Huybers, P.J., Langmuir, C.H., Park, S.H., 2015. Glacial cycles drive variations in the production of oceanic crust. *Science* 347, 1237–1240. doi:[10.1126/science.1261508](https://doi.org/10.1126/science.1261508).

569 Dalton, C.A., Langmuir, C.H., Gale, A., 2014. Geophysical and geochemical evidence for deep
 570 temperature variations beneath mid-ocean ridges. *Science* 344, 80–83. doi:[10.1126/science.](https://doi.org/10.1126/science.1249466)
 571 [1249466](https://doi.org/10.1126/science.1249466).

572 Dasgupta, R., Hirschmann, M.M., 2006. Melting in the Earth’s deep upper mantle caused by
 573 carbon dioxide. *Nature* 440, 659–662. doi:[10.1038/nature04612](https://doi.org/10.1038/nature04612).

574 Dasgupta, R., Hirschmann, M.M., 2010. The deep carbon cycle and melting in Earth’s interior.
 575 *Earth Planet Sci Lett* 298, 1–13. doi:[10.1016/j.epsl.2010.06.039](https://doi.org/10.1016/j.epsl.2010.06.039).

576 Dasgupta, R., Mallik, A., Tsuno, K., Withers, A.C., Hirth, G., Hirschmann, M.M., 2013. Carbon-
 577 dioxide-rich silicate melt in the Earth’s upper mantle. *Nature* 493, 211–215. doi:[10.1038/](https://doi.org/10.1038/nature11731)
 578 [nature11731](https://doi.org/10.1038/nature11731).

579 Elliott, T., Spiegelman, M., 2003. Melt migration in oceanic crustal production: a U-series per-
 580 spective, in: R.L. Rudnick, H.H., Turekian, K. (Eds.), *The Crust*. Elsevier-Pergamon. volume 3
 581 of *Treatise on Geochemistry*, pp. 465–510.

582 Gale, A., Langmuir, C.H., Dalton, C.A., 2014. The global systematics of ocean ridge basalts and
 583 their origin. *J Petrol* 55, 1051–1082. doi:[10.1093/petrology/egu017](https://doi.org/10.1093/petrology/egu017).

584 Galer, S.J.G., O’Nions, R.K., 1986. Magmagenesis and the mapping of chemical and isotopic
 585 variations in the mantle. *Chem Geol* 56, 45–61. doi:[10.1016/0009-2541\(86\)90109-9](https://doi.org/10.1016/0009-2541(86)90109-9).

586 Havlin, C., Parmentier, E.M., Hirth, G., 2013. Dike propagation driven by melt accumulation at
 587 the lithosphere–asthenosphere boundary. *Earth Planet Sci Lett* 376, 20–28. doi:[10.1016/j.](https://doi.org/10.1016/j.epsl.2013.06.010)
 588 [epsl.2013.06.010](https://doi.org/10.1016/j.epsl.2013.06.010).

589 Hebert, L.B., Montési, L.G.J., 2010. Generation of permeability barriers during melt extraction
 590 at mid-ocean ridges. *Geochem Geophys Geosyst* 11, Q12008. doi:[10.1029/2010GC003270](https://doi.org/10.1029/2010GC003270).

591 Hirano, N., 2011. Petit-spot volcanism: A new type of volcanic zone discovered near a trench.
 592 *Geochem J* 45, 157–167. doi:[10.2343/geochemj.1.0111](https://doi.org/10.2343/geochemj.1.0111).

593 Hirschmann, M.M., Asimow, P.D., Ghiorso, M.S., Stolper, E.M., 1999. Calculation of peridotite
 594 partial melting from thermodynamic models of minerals and melts. III. Controls on isobaric

595 melt production and the effect of water on melt production. *J Petrol* 40, 831–851. doi:[10.1093/](https://doi.org/10.1093/etroj/40.5.831)
596 [etroj/40.5.831](https://doi.org/10.1093/etroj/40.5.831).

597 Hirth, G., Kohlstedt, D., 2003. Rheology of the upper mantle and the mantle wedge: A view from
598 the experimentalists, in: Eiler, J. (Ed.), *Inside the Subduction Factory*. Am Geophys Union.
599 volume 138 of *Geophysical Monograph*, pp. 83–105.

600 Hirth, G., Kohlstedt, D.L., 1996. Water in the oceanic upper mantle: implications for rheology,
601 melt extraction and the evolution of the lithosphere. *Earth Planet Sci Lett* 144, 93–108. doi:[10.](https://doi.org/10.1016/0012-821X(96)00154-9)
602 [1016/0012-821X\(96\)00154-9](https://doi.org/10.1016/0012-821X(96)00154-9).

603 Huybers, P., Langmuir, C., 2009. Feedback between deglaciation, volcanism, and atmospheric
604 CO₂. *Earth Planet Sci Lett* 286, 479–491. doi:[10.1016/j.epsl.2009.07.014](https://doi.org/10.1016/j.epsl.2009.07.014).

605 Huybers, P., Langmuir, C.H., 2017. Delayed CO₂ emissions from mid-ocean ridge volcanism as
606 a possible cause of late-Pleistocene glacial cycles. *Earth Planet Sci Lett* 457, 238–249. doi:[10.](https://doi.org/10.1016/j.epsl.2016.09.021)
607 [1016/j.epsl.2016.09.021](https://doi.org/10.1016/j.epsl.2016.09.021).

608 Katz, R., 2008. Magma dynamics with the enthalpy method: Benchmark solutions and magmatic
609 focusing at mid-ocean ridges. *J Petrol* 49, 2099–2121. doi:[10.1093/etrology/egn058](https://doi.org/10.1093/etrology/egn058).

610 Katz, R., 2010. Porosity-driven convection and asymmetry beneath mid-ocean ridges. *Geochem*
611 *Geophys Geosyst* 10. doi:[10.1029/2010GC003282](https://doi.org/10.1029/2010GC003282).

612 Katz, R., Knepley, M., Smith, B., Spiegelman, M., Coon, E., 2007. Numerical simulation of
613 geodynamic processes with the Portable Extensible Toolkit for Scientific Computation. *Phys*
614 *Earth Planet Inter* 163, 52–68. doi:[10.1016/j.pepi.2007.04.016](https://doi.org/10.1016/j.pepi.2007.04.016).

615 Kawakatsu, H., Kumar, P., Takei, Y., Shinohara, M., Kanazawa, T., Araki, E., Suyehiro, K., 2009.
616 Seismic Evidence for Sharp Lithosphere-Asthenosphere Boundaries of Oceanic Plates. *Science*
617 324, 499–502. doi:[10.1126/science.1169499](https://doi.org/10.1126/science.1169499).

618 Kelemen, P.B., Manning, C.E., 2015. Reevaluating carbon fluxes in subduction zones, what goes
619 down, mostly comes up. *Proc Natl Acad Sci* doi:[10.1073/pnas.1507889112](https://doi.org/10.1073/pnas.1507889112).

620 Kelemen, P.B., Shimizu, N., Salters, V.J.M., 1995. Extraction of mid-ocean-ridge basalt from the
621 upwelling mantle by focused flow of melt in dunite channels. *Nature* 375, 747–753. doi:[10.1038/](https://doi.org/10.1038/375747a0)
622 [375747a0](https://doi.org/10.1038/375747a0).

623 Keller, T., Katz, R.F., 2015. R_DMC Online repository. URL: [https://bitbucket.org/](https://bitbucket.org/tokeller/r_dmc)
624 [tokeller/r_dmc](https://bitbucket.org/tokeller/r_dmc).

625 Keller, T., Katz, R.F., 2016. The role of volatiles in reactive melt transport in the asthenosphere.
626 J Petrol 57, 1073–1108. doi:[10.1093/petrology/egw030](https://doi.org/10.1093/petrology/egw030).

627 Klein, E.M., Langmuir, C.H., 1987. Global correlations of ocean ridge basalt chemistry with axial
628 depth and crustal thickness. J Geophys Res 92, 8089–8115. doi:[10.1029/JB092iB08p08089](https://doi.org/10.1029/JB092iB08p08089).

629 Kono, Y., Kenney-Benson, C., Hummer, D., Ohfuji, H., Park, C., Shen, G., Wang, Y., Kavner, A.,
630 Manning, C.E., 2014. Ultralow viscosity of carbonate melts at high pressures. Nature Commun
631 5, 5091. doi:[10.1038/ncomms6091](https://doi.org/10.1038/ncomms6091).

632 Lachenbruch, A., 1976. Dynamics of a passive spreading center. J Geophys Res 81, 1883–1902.
633 doi:[10.1029/JB081i011p01883](https://doi.org/10.1029/JB081i011p01883).

634 Langmuir, C., Klein, E., Plank, T., 1992. Petrological systematics of mid-oceanic ridge basalts:
635 constraints on melt generation beneath ocean ridges, in: Phipps Morgan, J., Blackman, D.,
636 Sinton, J. (Eds.), Mantle flow and melt generation at mid-ocean ridges. Am Geophys Union.
637 volume 71 of *Geophysical Monograph*, pp. 183–280.

638 Le Voyer, M., Kelley, K.A., Cottrell, E., Hauri, E.H., 2017. Heterogeneity in mantle carbon content
639 from CO₂-undersaturated basalts. Nature Commun 8, 14062. doi:[10.1038/ncomms14062](https://doi.org/10.1038/ncomms14062).

640 McKenzie, D., 1984. The generation and compaction of partially molten rock. J Petrol 25, 713–765.
641 doi:[10.1093/petrology/25.3.713](https://doi.org/10.1093/petrology/25.3.713).

642 McKenzie, D., 2000. Constraints on melt generation and transport from U-series activity ratios.
643 Chem Geol 162, 81–94. doi:[10.1016/S0009-2541\(99\)00126-6](https://doi.org/10.1016/S0009-2541(99)00126-6).

644 McKenzie, D., Bickle, M.J., 1988. The volume and composition of melt generated by extension of
645 the lithosphere. J Petrol 29, 625–679. doi:[10.1093/petrology/29.3.625](https://doi.org/10.1093/petrology/29.3.625).

646 Mei, S., Bai, W., Hiraga, T., Kohlstedt, D.L., 2002. Influence of melt on the creep behavior
647 of olivine–basalt aggregates under hydrous conditions. Earth Planet Sci Lett 201, 491–507.
648 doi:[10.1016/S0012-821X\(02\)00745-8](https://doi.org/10.1016/S0012-821X(02)00745-8).

649 Mian, Z.U., Tozer, D.C., 1990. No water, no plate tectonics: convective heat transfer and the
650 planetary surfaces of Venus and Earth. *Terra Nova* 2, 455–459. doi:[10.1111/j.1365-3121.](https://doi.org/10.1111/j.1365-3121.1990.tb00102.x)
651 [1990.tb00102.x](https://doi.org/10.1111/j.1365-3121.1990.tb00102.x).

652 Michael, P.J., Graham, D.W., 2015. The behavior and concentration of CO₂ in the suboceanic
653 mantle: Inferences from undegassed ocean ridge and ocean island basalts. *Lithosphere* 236-237,
654 338–351. doi:[10.1016/j.lithos.2015.08.020](https://doi.org/10.1016/j.lithos.2015.08.020).

655 Montési, L.G.J., Behn, M.D., Hebert, L.B., Lin, J., Barry, J.L., 2011. Controls on melt migration
656 and extraction at the ultraslow Southwest Indian Ridge 10°–16°E. *J Geophys Res–Solid Earth*
657 116. doi:[10.1029/2011JB008259](https://doi.org/10.1029/2011JB008259).

658 Müntener, O., Kelemen, P.B., Grove, T.L., 2001. The role of H₂O during crystallization of
659 primitive arc magmas under uppermost mantle conditions and genesis of igneous pyroxenites:
660 an experimental study. *Contrib mineral Petrol* 141, 643–658. doi:[10.1007/s004100100266](https://doi.org/10.1007/s004100100266).

661 O’Hara, M.J., 1985. Importance of the ‘shape’ of the melting regime during partial melting of the
662 mantle. *Nature* 314, 58–62. doi:[10.1038/314058a0](https://doi.org/10.1038/314058a0).

663 Olugboji, T.M., Park, J., Karato, S.I., Shinohara, M., 2016. Nature of the seismic lithosphere-
664 asthenosphere boundary within normal oceanic mantle from high-resolution receiver functions.
665 *Geochem Geophys Geosyst* doi:[10.1002/2015GC006214](https://doi.org/10.1002/2015GC006214).

666 Pilet, S., Baker, M.B., Müntener, O., Stolper, E.M., 2011. Monte Carlo Simulations of Metasomatic
667 Enrichment in the Lithosphere and Implications for the Source of Alkaline Basalts. *J Petrol* 52,
668 1415–1442. doi:[10.1093/petrology/egr007](https://doi.org/10.1093/petrology/egr007).

669 Plank, T., Langmuir, C.H., 1992. Effects of the melting regime on the composition of the oceanic
670 crust. *J Geophys Res* 97, 19749–19770. doi:[10.1029/92JB01769](https://doi.org/10.1029/92JB01769).

671 Regenauer-Lieb, K., 2001. The Initiation of Subduction: Criticality by Addition of Water? *Science*
672 294, 578–580. doi:[10.1126/science.1063891](https://doi.org/10.1126/science.1063891).

673 Resing, J.A., Lupton, J.E., Feely, R.A., Lilley, M.D., 2004. CO₂ and ³He in hydrothermal plumes:
674 implications for mid-ocean ridge CO₂ flux. *Earth Planet Sci Lett* 226, 449–464. doi:[10.1016/](https://doi.org/10.1016/j.epsl.2004.07.028)
675 [j.epsl.2004.07.028](https://doi.org/10.1016/j.epsl.2004.07.028).

676 Rosenthal, A., Hauri, E.H., Hirschmann, M.M., 2015. Experimental determination of C, F, and H
677 partitioning between mantle minerals and carbonated basalt, CO₂/Ba and CO₂/Nb systematics
678 of partial melting, and the CO₂ contents of basaltic source regions . *Earth Planet Sci Lett* 412,
679 77–87. doi:[10.1016/j.epsl.2014.11.044](https://doi.org/10.1016/j.epsl.2014.11.044).

680 Rudge, J.F., Bercovici, D., Spiegelman, M., 2011. Disequilibrium melting of a two phase multi-
681 component mantle. *Geophys J Int* 184, 699–718. doi:[10.1111/j.1365-246X.2010.04870.x](https://doi.org/10.1111/j.1365-246X.2010.04870.x).

682 Schmerr, N., 2012. The Gutenberg Discontinuity: Melt at the Lithosphere-Asthenosphere Bound-
683 ary. *Science* 335, 1480–1483. doi:[10.1126/science.1215433](https://doi.org/10.1126/science.1215433).

684 Sparks, D., Parmentier, E., 1991. Melt extraction from the mantle beneath spreading centers.
685 *Earth Planet Sci Lett* 105, 368–377. doi:[10.1016/0012-821X\(91\)90178-K](https://doi.org/10.1016/0012-821X(91)90178-K).

686 Spiegelman, M., Elliott, T., 1993. Consequences of melt transport for uranium series disequilibrium
687 in young lavas. *Earth Planet Sci Lett* 118, 1–20. doi:[10.1016/0012-821X\(93\)90155-3](https://doi.org/10.1016/0012-821X(93)90155-3).

688 Spiegelman, M., Kelemen, P., Aharonov, E., 2001. Causes and consequences of flow organization
689 during melt transport: the reaction infiltration instability in compactible media. *J Geophys*
690 *Res-Solid Earth* 106, 2061–2077. doi:[10.1029/2000JB900240](https://doi.org/10.1029/2000JB900240).

691 Spiegelman, M., McKenzie, D., 1987. Simple 2-D models for melt extraction at mid-ocean ridges
692 and island arcs. *Earth Planet Sci Lett* 83, 137–152. doi:[10.1016/0012-821X\(87\)90057-4](https://doi.org/10.1016/0012-821X(87)90057-4).

693 Stern, T.A., Henrys, S.A., Okaya, D., Louie, J.N., Savage, M.K., Lamb, S., Sato, H., Sutherland,
694 R., Iwasaki, T., 2015. A seismic reflection image for the base of a tectonic plate. *Nature* 518,
695 85–88. doi:[10.1038/nature14146](https://doi.org/10.1038/nature14146).

696 Szymczak, P., Ladd, A.J.C., 2014. Reactive-infiltration instabilities in rocks. Part 2. Dissolution
697 of a porous matrix. *J Fluid Mech* 738, 591–630. doi:[10.1017/jfm.2013.586](https://doi.org/10.1017/jfm.2013.586).

698 Tolstoy, M., 2015. Mid-ocean ridge eruptions as a climate valve. *Geophys Res Lett* 42, 1346–1351.
699 doi:[10.1002/2014GL063015](https://doi.org/10.1002/2014GL063015).

700 Turner, A.J., Katz, R.F., Behn, M.D., 2015. Grain-size dynamics beneath mid-ocean ridges:
701 Implications for permeability and melt extraction. *Geochem Geophys Geosyst* 16, 925–946.
702 doi:[10.1002/2014GC005692](https://doi.org/10.1002/2014GC005692).

703 Ulmer, P., 2001. Partial melting in the mantle wedge — the role of H₂O in the genesis of
 704 mantle-derived ‘arc-related’ magmas. *Phys Earth Planet Inter* 127, 215–232. doi:[10.1016/
 705 S0031-9201\(01\)00229-1](https://doi.org/10.1016/S0031-9201(01)00229-1).

706 White, R.S., Minshull, T.A., Bickle, M.J., Robinson, C.J., 2001. Melt generation at very slow-
 707 spreading oceanic ridges: Constraints from geochemical and geophysical data. *J Petrol* 42,
 708 1171–1196. doi:[10.1093/petrology/42.6.1171](https://doi.org/10.1093/petrology/42.6.1171).

709 Workman, R.K., Hart, S.R., Jackson, M., Regelous, M., Farley, K.A., Blusztajn, J., Kurz, M.,
 710 Staudigel, H., 2004. Recycled metasomatized lithosphere as the origin of the Enriched Mantle
 711 II (EM2) end-member: Evidence from the Samoan Volcanic Chain. *Geochem Geophys Geosyst*
 712 5, Q04008. doi:[10.1029/2003GC000623](https://doi.org/10.1029/2003GC000623).