A Nonmagnetic Differentiated Early Planetary Body

Benjamin P. Weiss^{1*}, Huapei Wang¹, Thomas G. Sharp², Jérôme Gattacceca³, David L. Shuster^{4,5}, Brynna Downey¹, Jinping Hu², Roger R. Fu¹, Aaron T. Kuan⁶, Clément Suavet¹, Anthony J. Irving⁷, Jun Wang⁸, Jiajun Wang⁸

¹Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

²School of Earth and Space Exploration, Arizona State University, Tempe, AZ, USA

³Centre National de la Recherche Scientifique, Aix-Marseille Université, Institut de Recherche pour le Développement, Collège de France, CEREGE, Aix-en-Provence, France

⁴Department of Earth and Planetary Science, University of California, Berkeley, CA 94720, USA

⁵Berkeley Geochronology Center, 2455 Ridge Road, Berkeley, CA 94709, USA

⁶Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

⁷Department of Earth and Space Sciences, University of Washington, Seattle, WA 98195

⁸Photon Sciences Directorate, Brookhaven National Laboratory, Upton, NY, USA,

*To whom correspondence should be addressed: bpweiss@mit.edu

Corresponding author: Benjamin P. Weiss, 77 Massachusetts Avenue 54-622, Cambridge, MA 02139, USA, tel.: 617-324-0224, fax: 617-258-7401, email: bpweiss@mit.edu

Keywords: planetesimals; paleomagnetism; meteorites; dynamo; differentiation

Paleomagnetic studies of meteorites have shown that the solar nebula was likely magnetized, while many early planetary bodies formed advecting metallic cores that generated dynamo magnetic fields. The surface fields on these bodies were recorded by a diversity of chondrites and achondrites, ranging in intensity from several μT to several hundred μT . In fact, an achondrite parent body without evidence for paleomagnetic fields has yet to be confidently identified, hinting that early solar system field generation and the dynamo process in particular may have been common. Here we present paleomagnetic measurements of the ungrouped achondrite NWA 7325 indicating that it last cooled in a near-zero field ($<1.7 \mu\text{T}$), estimated to have occurred at 4563.09 ± 0.26 million years ago (Ma) from Al-Mg chronometry. Because NWA 7325 is highly depleted in siderophile elements, its parent body nevertheless underwent large-scale metal-silicate differentiation and likely formed a metallic core. This makes NWA 7325 the first recognized example of an essentially unmagnetized igneous rock from a differentiated early solar system body. These results indicate that all magnetic fields, including those from any core dynamo on the NWA 7325 parent body, the solar nebula, young Sun, and solar wind, were $<1.7 \mu\text{T}$ at the location of NWA 7325 at 4563 Ma. This supports a recent conclusion that the solar nebula had dissipated by ~ 4 million years after solar system formation. NWA 7325 serves as an experimental control that gives greater confidence in the positive identification of remanent magnetization in other achondrites.

1. Introduction

The existence of dozens of achondrite groups and ungrouped achondrites indicates that numerous planetesimals experienced large-scale melting and differentiation in the early solar system (Scheinberg et al., 2015b). The energy for melting was likely supplied by the decay of the short-lived radionuclide ^{26}Al , which should have substantially heated bodies that accreted a large fraction of their masses within 2 million years (My) after the formation of calcium aluminum-rich inclusions (CAIs) (Weisberg et al., 2006). Hf-W chronometry of iron meteorites confirms that many bodies experienced large-scale metal-silicate fractionation during this period (Kruijer et al., 2014). Thermal and compositional core convection (Nimmo, 2009), possibly combined with mechanical stirring from impacts, likely powered core advection for ten to

perhaps a few hundred My (Bryson et al., 2015; Elkins-Tanton et al., 2011; Formisano et al., 2016; Scheinberg et al., 2015a; Sterenborg and Crowley, 2013).

Scaling relationships derived from magnetohydrodynamic simulations of convecting metallic cores indicate that many of these bodies should have been capable of generating dynamos (Weiss et al., 2010). Indeed, essentially every achondrite group that has yet been studied with modern paleomagnetic methods has been found to contain ancient natural remanent magnetization (NRM): evidence for past core dynamos has been identified in plutonic angrites (Wang et al., 2017), eucrites (Fu et al., 2012), and main group pallasites (Bryson et al., 2015; Tarduno et al., 2012). Remanent magnetization attributed to possible core dynamos has even been found in CV (Carpözen et al., 2011; Gattacceca et al., 2016) and CM chondrites (Cournède et al., 2015). Although a study of five eucrites including ALHA81001 and PCA 82502 concluded that they recorded low-field ($<10 \mu\text{T}$) conditions (Cisowski, 1991), this study lacked precise radiometric ages and thermochronometry, used only relatively low-field alternating field (AF) demagnetization, and was unable to prevent thermochemical alteration of the samples during thermal demagnetization; recent studies of ALHA81001 and PCA 82502 identified stable, unidirectional NRM of extraterrestrial origin acquired in fields of ~ 12 and $\sim 70 \mu\text{T}$, respectively (Fischer et al., 2013; Fu et al., 2012). All told, the available paleomagnetic data indicate that planetesimal dynamos were likely common, if not ubiquitous, in the early solar system.

Nevertheless, it is expected that not all differentiated bodies that underwent large-scale metal-silicate fractionation should have been capable of generating dynamos. There are numerous reasons a dynamo might be inhibited on a differentiated body. First, it is possible that a body underwent large-scale metal-silicate fractionation but formed localized, dispersed metal-rich regions rather than a single metallic core. The low gravity and possibility of incomplete melting on planetesimals means that even if temperatures exceeded that of the Fe-S solidus, metal percolation to the center of planetesimals was not guaranteed (Cerantola et al., 2015). Possible examples of such bodies include the winonaite-IAB iron meteorite parent body and the IIE iron meteorite parent body (Ruzicka, 2014). Second, even if a metallic core formed, it may have not advected. Thermal convection would be inhibited by sufficiently slow cooling rates, as might be expected if the overlying mantle was solid and sufficiently thick; such an outcome may be consistent with the proposal that planetesimals never formed silicate magma oceans but only melted episodically and in small quantities (Wilson and Keil, 2012). Compositional convection

is a potent power source but is only expected for metallic cores containing a light alloying element (Nimmo, 2009). Furthermore, although both outward core crystallization and inward core crystallization in the iron snow (Rückriemen et al., 2015) regime provides a buoyancy flux for driving convection, inward dendritic core crystallization does not (Scheinberg et al., 2015a). Third, even if the metallic core advected, it still might not generate a dynamo. Fluid velocities may be too slow, as expected for core radii smaller than ~80-100 km (Elkins-Tanton et al., 2011; Nimmo, 2009; Weiss et al., 2010) or perhaps even ranging up to radii of 1000 km (Sternborg and Crowley, 2013), or the velocity field may have an unsuitable geometry (e.g., not lead to a field-amplification instability) (Stevenson, 1983). One or more of these reasons can account for why, among all rocky and icy solar system bodies, only the Earth, Mercury, and Ganymede are known to have active dynamos today (Stevenson, 2010). Finally, even if a body generated a dynamo, meteorites from this body may not retain records of this field if the meteorites were last cooled or aqueously altered when the dynamo was not active. In particular, it has recently been argued that thermal blanketing of planetesimal cores by ^{26}Al -enriched mantles should delay thermal convection dynamos until at least several million years after differentiation (Sternborg and Crowley, 2013).

We see that there is every expectation that differentiated planetesimals should have formed in the early solar system that either never generated magnetic fields or at least produced fields not recorded by achondrites from these bodies. Searching for such examples would not only establish the frequency and diversity of planetesimal dynamo activity, but also test the methodologies that have been developed for identifying dynamos from the meteorite record. In short, demonstrating our ability to identify unmagnetized meteorites would build greater confidence in our identification of magnetized meteorites.

Because arbitrarily weak paleofields will produce essentially no detectable NRM, it is not possible to demonstrate that an apparently unmagnetized rock formed in truly zero-field conditions but only to place an upper limit on the paleofield. Even so, meteorites with near-zero paleointensities are difficult to identify because they are unfamiliar: essentially all Earth rocks that have been studied with paleomagnetic methods formed in the geomagnetic field. Furthermore, numerous processes can impart secondary magnetization after the meteorites arrive at Earth and enter the geomagnetic field: surficial heating during atmospheric entry, weathering and associated crystallization of ferromagnetic grains, viscous remanence acquisition, and

application of hand magnets by meteorite collectors (Weiss et al., 2010). Another major problem is the acquisition of spurious remanence during the laboratory demagnetization process. The ferromagnetism of the vast majority of achondrites is dominated by the iron-nickel minerals kamacite and martensite, which usually form low-coercivity, multidomain grains that readily acquire spurious remanence during alternating field (AF) demagnetization (Weiss et al., 2010). This spurious remanence could potentially be mistaken for NRM or, at the very least, limits our ability to put stringent upper limits on the paleointensities for these samples. Furthermore, thermal demagnetization of these samples typically leads to thermochemical alteration of these iron-nickel minerals (Suavet et al., 2014), which can render Thellier-Thellier paleointensities inaccurate. Finally, because individual ferromagnetic grains have spontaneous magnetization even when the bulk rock is not magnetized, paleomagnetic analysis of small samples (as is common in extraterrestrial paleomagnetic studies) is fundamentally limited by the statistics of small numbers of grains (Berndt et al., 2016). As a result of these limitations, only recently have we been able to confidently identify largely unmagnetized rocks from the Moon that place meaningful upper limits ($<4 \mu\text{T}$) on the paleofield after 3.5 Ga (Tikoo et al., 2014).

Identification of null magnetic field conditions on an early planetesimal requires an ancient, well-preserved sample with high-fidelity magnetic recording properties. Here we present paleomagnetic, petrographic, and $^{40}\text{Ar}/^{39}\text{Ar}$ analyses of the ungrouped basaltic achondrite Northwest Africa (NWA) 7325. Our measurements show that this meteorite was last cooled from above the peak magnetic disordering temperature ($\sim 780^\circ\text{C}$) in a field of no greater than $\sim 1.7 \mu\text{T}$.

2. Petrology, thermal history, and age of NWA 7325

NWA 7325 is a dark green achondrite partially covered with a chartreuse-colored fusion crust found as 37 separate pieces in the Western Sahara in 2012 (Barrat et al., 2015; Irving et al., 2013). It is an unbrecciated, fine- to medium-grained (typical grain size 0.5-1 mm) plutonic rock (probably a cumulate) composed of 56 vol.% calcic plagioclase, 27 vol.% diopside, 16 vol.% forsterite and accessory troilite containing Cr-rich lamellae, chromite, and FeNi metal.

NWA 7325 is thought to have been produced as partial melt in the rocky exterior of a highly differentiated parent body (Barrat et al., 2015; Weber et al., 2016). Its abundances of highly siderophile elements are depleted by 3-4 orders of magnitude relative to CI chondrites,

suggesting that its parent body underwent large-scale metal-silicate fraction (Archer et al., 2015). NWA 7325's distinctive elemental composition and unique combination of O and ^{54}Cr isotopes (Barrat et al., 2015; Irving et al., 2013) indicate that it originate from a parent body not sampled by other achondrites. It has even been suggested that this parent body could be Mercury (Irving et al., 2013; Koefoed et al., 2016).

With a pyroxene Pb-Pb age of 4563.4 ± 2.6 Ma and an Al-Mg model age of 4563.09 ± 0.26 Ma (Koefoed et al., 2016), NWA 7325 has one of the oldest crystallization ages amongst known igneous rocks. Although cracks in the meteorite contain secondary carbonate from terrestrial weathering and very rare secondary oxides (Barrat et al., 2015; Weber et al., 2016), NWA 7325 is overall very fresh and nearly all metal grains are unaltered. Nevertheless, multiple lines of evidence suggest that following its formation, NWA 7325 was substantially reheated and partially melted, followed by rapid cooling. Olivines in the meteorite exhibit undulatory extinction and fracturing while plagioclase is not maskelytinized, suggesting the meteorite experienced shock pressures of 5-10 GPa (shock stage 2) (Weber et al., 2016). Despite these relatively low shock pressures, there is abundant evidence for a brief high-temperature event that led to localized melting. A second generation of fine (<20 μm) plagioclase laths crystallized in cracks and at grain boundaries with mafic minerals, in some cases in contact with SiO_2 -normative late-stage melts (Weber et al., 2016). Furthermore, plagioclase contains finely dispersed tiny (<5 μm) metal and iron sulfide grains, suggesting it was melted and then cooled rapidly (Weber et al., 2016) (Fig. 1A, D-E). As described in Section 3.1, our observations of partial desulfurization of FeS provide additional evidence for this remelting event. ^{26}Al (Barrat et al., 2015) and ^{129}Xe (Weber et al., 2016) isotopic data are consistent with this melting event occurring just several tens of My after the formation of CAIs. Such an early age for this thermal event is also consistent with bulk rock and plagioclase $^{40}\text{Ar}/^{39}\text{Ar}$ ages of 4516 ± 160 Ma and 4481 ± 150 Ma (Weber et al., 2016).

In an attempt to further constrain the age and thermal history of NWA 7325, we conducted $^{40}\text{Ar}/^{39}\text{Ar}$ stepwise degassing analysis of a neutron-irradiated bulk chip from the meteorite (Fig. S1). As described in the supplementary text, this experiment was not successful due to a lack of statistical concordance of the results and so we do not discuss these data further.

3. Ferromagnetic grains

3.1. Mineralogy

To determine the mineralogy and constrain the grain size and setting of the ferromagnetic minerals in NWA 7325, we conducted backscattered electron microscopy (BSEM), electron dispersive spectroscopy (EDS), and transmission electron microscopy (TEM) analyses in the MIT Experimental Petrography Lab, Harvard Center for Nanoscale Science, and Le-Roy Eyring Center for Solid State Science at Arizona State University. These collectively data confirm the presence of numerous fine (mostly $<1\ \mu\text{m}$ size, with many grains $<0.1\ \mu\text{m}$) metal and Cr-rich iron sulfide grains showing few secondary alteration textures (Figs. 1 and S2 and Table S2). Wavelength dispersive spectroscopy (WDS) showed that nearly all metal grains are FeNi metal (87-92% Fe alloyed with 5-10% Ni), although we observed one large Cr-rich FeNi grain (Table S2 and Fig. 1B). Multiple WDS analyses taken from within metal grains did not detect significant variations in Ni abundances, while our high-contrast BSEM and bright-field TEM images show no evidence of exsolution textures in the FeNi grains down to the resolution of the images ($\sim 0.1\ \mu\text{m}$ and $<10\ \text{nm}$, respectively). Therefore, the metal grains' Ni-abundance indicates that they should be dominantly in the form of the body-centered cubic (bcc) phase martensite (α_2 -FeNi). Electron diffraction patterns from the FeNi grains confirm a bcc structure (Fig. 1F-G). The presence of lamellar twins and evidence for lattice strain in some of these metal particles (Fig. 1H) are consistent the presence of lath martensite, the expected martensite morphology for the low-Ni, low-C composition of the metal in NWA 7325 that crystallized from the parent γ -FeNi (fcc) phase (Krauss, 2005). The host for the metal particles is polycrystalline plagioclase, consisting of nanocrystals with an apparent lattice-preferred orientation (Fig. 1F). The metal particles are closely associated with FeS grains and voids in the plagioclase (Fig. 1 E-F). This association suggests that the metal formed by the partial desulfidization of the FeS, producing metal and sulfur vapor, providing more evidence for a second melting event after formation of the rock (see Section 2.1).

Our rock magnetic experiments provide further evidence that martensite is the dominant ferromagnetic mineral and additionally constrain its domain size and coercivity. Martensite with the composition found in NWA 7325 should have martensite-start and martensite-finish temperatures ranging between 680 - 510°C and 630 - 430°C , respectively, and austenite-start and austenite-finish temperatures of 760 - 680°C and 780 - 700°C , respectively (Swartzendruber et al., 1991). These temperatures are consistent with our thermomagnetic measurements (Fig. 2),

which show hysteretic behavior indicative of sluggish transformations from α_2 -FeNi to γ -FeNi occurring at 760-800°C during warming and γ -FeNi to α_2 -FeNi at ~550-560°C during cooling. Our measurements of saturation magnetization, M_s , of subsamples SR1 and JG2 of NWA 7325 indicate a metal abundance of 243 and 358 ppm by mass [assuming a M_s for pure Fe of 224 Am² kg⁻¹ (Dunlop and Özdemir, 1997)].

Low-temperature cycling of room-temperature saturation remanence, M_{rs} , using a Quantum Designs Magnetic Properties Measurement System at the Institute for Rock Magnetism identified a weak magnetic transition at a temperature of ~118 K, consistent with the Verwey transition of magnetite [although no Verwey transition was detectable in a field cooled warming of low temperature M_{rs} experiment] (Fig. S3). Given the small observed saturation remanence decrease (average of 2.85×10^{-5} Am² kg⁻¹ across the Verwey transition temperature for two repeat field-cooled warming experiments), magnetite's M_s of 92 Am² kg⁻¹, assuming the magnetite grains are pseudo single domain (PSD) in size and so have $M_{rs}/M_s \sim 0.1$, and assuming demagnetization of magnetite's M_{rs} by 50% across the Verwey transition (Özdemir et al., 2002), this indicates that NWA 7325 contains just ~6 ppm magnetite by mass, consistent with petrographic evidence for very minor terrestrial weathering (see above).

3.2. Thermochemical stability of magnetization carriers

The valence states of Cr, V, and Ti in NWA 7325 indicate the meteorite formed under highly reducing conditions with an estimated oxygen fugacity somewhere between 2 and 7 log units below the iron-wüstite (IW) buffer (Sutton et al., 2014). As a result, we conducted our thermal demagnetization and paleointensity analyses (Sections 4 and 5) under controlled oxygen fugacity conditions to mitigate thermochemical alteration of the ferromagnetic grains. However, because of the large range in oxygen fugacity estimates for NWA 7325, we first heated subsamples of the meteorite under a range of reducing conditions and temperatures to establish the optimal oxygen fugacity for mitigating alteration of the rock magnetic properties. These experiments were conducted in a thermal demagnetization system that employs regulated CO₂-H₂ gas mixing (Suavet et al., 2014).

Prior to each heating step, we conducted repeated rock magnetic experiment to quantify the effect of subsequent heating on the magnetization carrying-capacity of the rock. Beginning with fresh unheated samples, we measured anhysteretic remanent magnetization (ARM)

acquisition in a peak alternating field (AF) of 200 mT and bias fields ranging from 0.05 to 1.5 mT, AF demagnetized this ARM to 400 mT, conducted stepwise IRM acquisition up to 1 T, and finally AF demagnetized this IRM to 400 mT. We then gave the samples another IRM in a 1 T field and then heated them to a particular temperature in zero field; the goal of the latter measurements was to determine when alteration might set in during thermal demagnetization of NRM. We then AF demagnetized this thermally demagnetized IRM and again measured stepwise IRM acquisition up to 1 T; the goal of the latter measurements was to assess the maximum possible stability expected for IRM and, by implication, NRM. This sequence of measurements was then repeated for increasingly higher heating temperatures ranging up to 600°C and in oxygen fugacity conditions ranging from IW to well below IW-4 [the most reducing experiments employed essentially pure (99.995%) lab-grade H₂ gas]. The results show that under all oxygen fugacity conditions studied, thermochemical alteration of the magnetic carriers, manifested as strong increases in magnetization intensity, occurred after temperatures reached 400°C, at which point ARM and SIRM reached values ~3 times that of the original unheated sample (although one subsample heated under IW-3 conditions, 7B3B, remained relatively unaltered up to at least 500°C) (Fig. S4). The condition that resulted in the least alteration at 300°C was IW-3 (for which ARM and IRM increased by several tens of %), although IW-2 and IW conditions yielded very similar results. Even so, after heating in IW-3 conditions to 500°C and 800°C, ARM and IRM increased by factors of ~2-6 and ~7-10, respectively. Our inability to mitigate alteration may reflect the fact NWA 7325 cooled under temporally varying oxygen fugacity conditions; this possibility is consistent with the 5 orders of magnitude range in oxygen fugacity estimates for the meteorite (Sutton et al., 2014). Another likely problem is that we did not control for sulfur fugacity despite the presence of sulfides in the meteorite.

3.3. Grain size and habit

As discussed in Section 3.1, our BSEM and TEM images observed FeNi grains with sizes mainly ranging from 0.1 to 1 µm. This range indicates a dominantly PSD size range (Einsle et al., 2016). To obtain additional information on the domain size and habit of the FeNi grains, we conducted X-ray tomography imaging and hysteresis measurements. Determining domain size and habit is important for establishing whether the grains can retain high-fidelity records of

ancient magnetic fields and also for calibrating our non-thermal paleointensity analyses (Section 5.2).

To image the three dimensional sizes and shapes of the larger FeNi grains, we conducted synchrotron-based full-field transmission hard X-ray microscopy (TXM) using a recently developed transmission hard X-ray microscope at beamline X8C at the National Synchrotron Light Source (NSLS), Brookhaven National Laboratory (BNL) (Wang et al., 2012). TXM has the advantage over TEM of enabling three dimensional analyses over much larger fields of view. We mounted a mm-sized grain from of NWA 7325 on a tungsten carbide pin, which was then placed on a kinematic stage. To target the FeNi minerals, we used an X-ray energy of 7.2 keV, just above the Fe absorption K-edge (7.112 keV) (Baker et al., 2012). We acquired images each with a field-of-view of $40\text{ }\mu\text{m} \times 40\text{ }\mu\text{m}$ and an effective pixel size of 40 nm after using 2×2 pixel binning. We then stitched together the TXM images to construct a $240 \times 240\text{ }\mu\text{m}$ image (Fig. 3A). We then rotated the sample 90° and conducted the same X-ray imaging for a side view (Fig. 3B). For a $40\text{ }\mu\text{m}$ cylindrical volume near the tip of the sample, we also conducted tomographic scans by taking 721 single X-ray images every 0.25° from 0° to 180° , which were then used to reconstruct a three-dimensional rendering of the cylinder (Movie S1). These X-ray data confirm the existence of abundant $<1\text{ }\mu\text{m}$ sized iron mineral grains with similar sizes as those visible in the SEM images (Fig. 3C and D).

We measured hysteresis loops on fresh subsamples (SR1, JG2, and SR3) and samples previously heated to 400°C in IW conditions and 800°C in IW-3 conditions (samples 7B3 and SR1a, respectively) with vibrating sample magnetometers at the Institute of Rock Magnetism (maximum applied field 1.5 T) and CEREGE (maximum applied field 1T) and with a Princeton Measurements Corporation Micromag 2900 alternating gradient force magnetometer (AGFM) in the Rutgers University Paleomagnetism Laboratory. The measured M_{rs}/M_s values and ratios of coercivity (B_c) to coercivity of remanence (B_{cr}) for all subsamples indicate a mean grain PSD grain size with a possible small contribution of superparamagnetic (SP) grains (Dunlop, 2002; Einsle et al., 2016) (Table S3 and Fig. 4). The presence of SP grains and/or multidomain (MD) grains is also indicated by the observation of remanence decay experienced during warming of a low-temperature saturation remanence (Fig. S3B).

We also measured first-order reversal curves (FORC) (Harrison and Feinberg, 2008) for one fresh and two heated subsamples with Rutgers AGFM. The FORC data were processed

using the software package FORCinel v. 2.01 (Harrison and Feinberg, 2008) with a smoothing factor of 6 (for fresh subsample SR3) and 3 (for heated subsamples 7B3C and SR1a) (Fig. 5). The magnetization of the fresh subsample SR3 is relatively weak, with a barely distinguishable narrow central ridge with coercivity B_c ranging up to 80 mT, consistent with PSD FeNi grains and a weak low-coercivity ($B_c < 10$ mT) signal with a wide range in H_u , indicative of MD ferromagnetic grains (Fig. 5A). The FORC diagram of heated subsample 7B3C (Fig. 5B) shows a clear narrow high coercivity central ridge, accompanied by an even stronger low coercivity signal, while subsample SR1a (Fig. 5C) shows an even stronger and wider low coercivity signal and virtually no high coercivity central ridge. The FORC diagrams of the fresh and heated-to-400°C subsamples indicate a mixture of PSD grains, while the heated-to-800°C subsample is dominated by MD grains, indicating that the 800°C heating treatment severely altered the mineralogy of the magnetic grains within NWA 7325 even in the controlled $fO_2 = IW-3$ environment.

These results are consistent with the conclusions in Section 3.1 and 3.2. All told, a diversity of imaging and rock magnetic techniques establishes that NWA 7325 contains dominantly single-phase FeNi martensite grains with sizes of 0.1-1 μm . These grains should be dominantly in the PSD vortex state, which has recently been shown to behave like stable uniaxial single domain particles with high unblocking and blocking temperatures and high resistance to remagnetization by hand magnets and viscous relaxation (Almeida et al., 2016; Einsle et al., 2016). In particular, any remanence in NWA 7325 that formed in the early solar system would likely have been stable against viscous decay over the history of the solar system: Néel theory shows that single domain grains with 1-hour blocking temperatures $>\sim 200^\circ\text{C}$ would be stable over 4.5 Gy at 0°C (Garrick-Bethell and Weiss 2010); a similar conclusion likely hold for the fine PSD grains in NWA 7325 [e.g., Winklhofer et al. (1997)]. This means that NWA 7325 has unusually high magnetic recording fidelity compared to most FeNi-bearing achondrites (Weiss et al., 2010).

4. Demagnetization of NRM

4.1. Methods

We analyzed the NRM of subsamples from 14 parent stones acquired by S. Ralew with masses ranging from 0.3-2.1 g. All NRM and associated demagnetization measurements were

conducted in the MIT Paleomagnetism Laboratory inside a shielded room (DC field < 200 nT) using a 2G Enterprises Superconducting Rock Magnetometer (SRM) 755 equipped with automated AF demagnetization and remagnetization equipment (Kirschvink et al., 2008). Most sample handling was conducted in a class ~10,000 clean laboratory inside this shielded room. As expected for desert meteorites, most of these stones were found to be essentially completely remagnetized by collectors' hand magnets, as indicated by ratios of NRM to SIRM >10%. However, we found that at least 4 of these stones escaped being touched with a hand magnet as indicated by a lack of high-intensity NRM overprints. We selected one of these stones, NWA 7325 sample B7 (mass of 1.55 g) for more detailed paleomagnetic analyses.

Sample B7 contained no obvious fusion crust and showed few macroscopic signs of weathering. We measured nine mutually oriented subsamples with masses ranging from 70 to 107 mg cut from parent sample B7 using a wire saw lubricated with alcohol in our shielded clean laboratory. Three of these samples (B71-B73) were subjected to static three axis AF demagnetization up to either 85 or 150 mT. These samples were mounted with cyanoacrylate on to GE 124 silica glass slides with moments <10⁻¹¹ Am². The sample magnetic moments were measured after AF application in each of the three orthogonal directions. To reduce the acquisition of spurious remanence, we repeated AF applications 2-4 times for each AF level and averaged the measurements following the Zijderveld-Dunlop method (Tikoo et al., 2014). The remaining six subsamples (B74-B79) were thermally demagnetized up to either 400, 515 or 800°C in a controlled H₂-CO₂ atmosphere (Suavet et al., 2014) under IW-3 conditions in an ASC Scientific Model TD48-SC thermal demagnetizer [see Section 3.2]. Samples B74 and B75 were heated without any disk mounts or adhesives but measured on GE 124 disks using magnetically clean tape (total moment <1×10⁻¹¹ Am²), while B76-B79 were glued to GE 124 disks with potassium silicate for the entire demagnetization and measurement process. Two of the thermally-demagnetized samples (B74 and B75) were pre-treated with AF demagnetization to 5 mT prior to beginning thermal demagnetization to remove any weak IRM overprints (as described in Section 4.2, such overprints were found not to be present). The three-axis moment noise level of the MIT 2G SRM superconducting sensors under typical operating conditions in the MIT Paleomagnetism Laboratory is <1×10⁻¹² Am² (Wang et al., 2017). As such, in this study, the moments of the silica disk mounts are the main factor limiting the sensitivity of the 2G SRM measurements, such that our estimated moment uncertainty here is 0.5-1×10⁻¹¹ Am².

4.2. NRM components

NRM components were identified using principal component analysis (PCA) (Kirschvink, 1980). AF and thermal demagnetization isolated two NRM components that are approximately unidirectional across the parent sample (Figs. 6 and 7 and Table S4). A relatively strong low coercivity (LT), low temperature (LT) component unblocked up to a peak AF of 4-16 mT and a peak temperature of 165-340°C. Two subsamples contained a weak, medium coercivity (MC), medium temperature (MT) component that unblocked from the end of the LC/LT component up to a peak AF of 20.5 mT and temperature of 378°C. After removal of the MC/MT component, further demagnetization produced an approximately spherical cloud of vector endpoints with no demagnetization. The observed moment variations are ~50 times the noise limit imposed by the silica holder moments (see Fig. 6 legend) and so indicate near-total demagnetization of the NRM in combination with spurious remanence acquisition from the AF waveform rather than magnetometer sensor noise (see Section 4.1). PCA fits to this high coercivity (HC) and high temperature (HT) range not constrained to the origin have maximum angular deviation (MAD) values ranging from 30-46°. The high MAD values and lack of decay in magnetization intensity are strong indicators that there is no NRM blocked in the HC/HT range.

5. Paleointensity experiments

5.1. Overview

We conducted paleointensity experiments using both thermal and non-thermal methods. The advantage of thermal methods is that, if alteration can be mitigated, they directly reproduce the acquisition process by which any thermoremanent NRM was acquired and so in principle permit more accurate paleointensities than non-thermal methods. However, the accuracy of thermal methods is fundamentally limited by thermochemical alteration of the samples during heating (i.e., due to oxidation and sulfurization reactions and due to conversion of martensite to taenite). As shown in Section 3.2, our thermal alteration experiments demonstrated that even in our controlled atmosphere, NWA 7325 experiences significant thermochemical alteration above 400°C.

5.2. Methodology

We estimated the paleointensities for the components in NWA 7325 using two nonthermal (ARM- and IRM-based) multicomponent paleointensity techniques and one TRM-based multicomponent paleointensity technique (see supplementary text). All three techniques (ARM, IRM, and thermal) produced broadly similar paleointensities (Fig. 8). The LC/LT components for 9 subsamples have a mean paleointensity of 23 ± 5 μT , while our mean MC paleointensity estimate for subsample B73 was 7.5 ± 2.1 μT [uncertainties are 1 standard deviation (σ) from the mean. Nearly all HC/HT paleointensity estimates were within error of zero. The ARM, IRM, and 800°C total paleointensity experiments provided the most sensitive constraints; their mean HC/HT paleointensity is 1.7 ± 1.4 μT ($\pm 1\sigma$), indicating essentially zero-field conditions during the time of final cooling. Note that these uncertainties do not take into account the systematic uncertainties associated with the unknown efficiency of TRM to ARM and IRM. Taking these into consideration (supplementary text), the mean paleointensities values quoted above are estimated to have a 2- σ error of a factor of 5.

5.3. Paleointensity fidelity tests

Our demagnetization of NRM and associated paleointensity experiments suggest that grains in the HC/HT range do not carry NRM, such that the meteorite last cooled from the martensite ordering temperature in a near-zero field. We now seek to place an upper limit on the field environment at the time of last cooling by (a) determining the minimum paleointensity recoverable by the nonthermal techniques used in Section 5.2 and that (b) could produce a TRM component that exhibits linear, stable demagnetization when thermally demagnetized following our techniques used in Section 4.1. The details of our experiments are described in the supplementary materials. Method (a) shows that we can retrieve paleointensity estimates from NWA 7325 down to TRM-equivalent fields as low as ~ 6 μT (Table S6), but as discussed in Section 5.2, this value has a 2- σ uncertainty of a factor of 5. Method (b) shows that TRM produced by lab fields as weak as ~ 0.3 μT can just marginally be isolated by our thermal demagnetization methods (Fig. 9). Given that our thermal alteration tests (Section 3.2), suggest that heating to 800°C increases ARM by a factor of ~ 10 , these data suggest our thermal demagnetization protocol can isolate ancient TRM produced by paleofields of $0.3 \times 10 = 3$ μT . Importantly, this 3 μT value is not subject to the high uncertainties associated with our non-

thermal paleointensities because it was estimated by direct application of a TRM. The nominal ~6 μT and 3 μT upper limits from methods (a) and (b) are similar, although slightly less restrictive than, the ~1.7 μT upper limit inferred from our paleointensity experiments on the HC/HT magnetization range.

6. Meaning of magnetization in NWA 7325

The preceding experiments make it clear that NWA 7325 does not contain a total TRM. Its NRM essentially completely demagnetizes by peak laboratory temperatures of just 165-378°C (below the temperatures at which significant thermochemical alteration occurs) even though it contains ferromagnetic grains that can readily acquire weak-field ($< \sim 3 \mu\text{T}$) TRM that is stable to thermal demagnetization temperatures of $> 750^\circ\text{C}$. Furthermore its NRM essentially completely AF demagnetizes by just 4-20.5 mT; such low coercivity grains are readily remagnetized by shocks of just 2 GPa that would otherwise leave virtually no petrographic expression (Weiss et al., 2010) and anyway are below the shock pressures known to have been experienced by the meteorite (Section 2). Most of the NRM that is present in NWA 7325 is in the form of the LC/LT component. Its unidirectionality across the parent sample, ~23 μT paleointensity, and low peak unblocking temperature are all consistent with origin as a VRM in the Earth's field following landing on Earth.

The origin of the MC/MT component, unambiguously present in just two of the nine analyzed subsamples is more obscure given its weak paleointensity (about 10% of the Earth's field); it might be another VRM component that has weak (i.e., below that of Earth's field) apparent paleointensities due to changes in the meteorite's position with respect to the Earth's field since landing, or perhaps it is an inefficient crystallization remanent magnetization (CRM) carried by small quantities of secondary ferromagnetic minerals produced during terrestrial weathering. A CRM is broadly consistent with the $\sim 1.8 \times 10^{-10}$ and $\sim 1.3 \times 10^{-10} \text{ Am}^2$ intensities of the MC/MT components in subsamples B73 and B75 given the $\sim 5.7 \times 10^{-5} \text{ Am}^2$ SIRM for magnetite in NWA 7325 inferred from the low-temperature experiments (Section 3.1): for a typical efficiency of CRM produced in an Earth-strength field to SIRM of 0.2-0.02% (Dunlop and Özdemir, 1997), we estimate CRM carried by magnetite in these subsamples should have an intensity of $\sim 0.6\text{--}6 \times 10^{-10}$ and $1\text{--}10 \times 10^{-10} \text{ Am}^2$, respectively.

In any case, after laboratory demagnetization of the LC/LT and MC/MT components, the remaining HC/HT magnetization in NWA 7325 is highly nonunidirectional across the parent sample and has paleointensities within error of zero. The HC/HT magnetization is likely a combination of spurious remanence acquired during the AF and thermal demagnetization process (e.g., due to imperfections in the AF waveform and/or intragrain interaction fields, respectively) (Tikoo et al., 2014) and possibly also spontaneous remanence. We conclude that the field on the parent body of NWA 7325 when it last cooled from its 780°C Curie temperature was most likely $<1.7 \mu\text{T}$. Given that the estimated closure temperature of Mg diffusion in NWA 7325 anorthite exceeds 780°C, we conclude that these zero-field conditions are dated by the meteorite's Al-Mg model age of $4563.09 \pm 0.26 \text{ Ma}$ (supplementary online text).

7. Implications

NWA 7325 is the first recognized small-body achondrite robustly shown to have formed in near-zero field conditions in the early solar system [see original report by Weiss et al. (2013)]. Although numerous basaltic achondrites have been previously been found to have nonunidirectional and highly unstable NRM, most of these samples have very poor magnetic recording properties likely due to the coarse multidomain grain sizes of their constituent metal grains (Weiss et al., 2010). Furthermore, in nearly all such cases, quantitative limits on the maximum intensity and timing of the field experienced by these samples during final cooling were not established. An very recent exception is the volcanic angrites, which during the review period of this paper were also found to have cooled in a near-zero paleofield ($<0.7 \mu\text{T}$ at $4563.5 \pm 0.1 \text{ Ma}$) (Wang et al., 2017). The weakest robustly-recovered paleointensity recovered from an achondrite is the $\sim 12 \mu\text{T}$ value (with a lower bound of $2 \mu\text{T}$ given the uncertainties in the non-thermal paleointensity technique; see above) measured for the eucrite ALHA81001 (Fu et al., 2012). Additionally, two main group pallasites were recently shown to have formed within a minimum field strength of $\sim 1\text{--}7 \mu\text{T}$; this was interpreted as a record of fields from crustal remanence that itself was the product of an earlier thermal convection core dynamo (Nichols et al., 2016).

Although our $<1.7 \mu\text{T}$ paleointensity constraint from NWA 7325 provides no robust record of magnetic fields on the parent body, it is not sufficiently sensitive to exclude the existence of a weak dynamo or crustal remanent fields at the time of final cooling. In particular,

dynamo scaling laws indicate the feasibility of surface fields from planetesimal dynamos weaker than 3 μT (Formisano et al., 2016; Weiss et al., 2010). Furthermore, the crustal fields expected for a body whose crust had been magnetized by even a strong dynamo surface field can also easily be weaker than 3 μT . Following Fu et al. (2012), the maximum possible crustal field at a demagnetized location within the crust of a body carrying a TRM produced by an ancient dynamo field of 100 μT is given by

$$B_{\text{crust}} < \mu_0 M_{\text{TRM}} \rho$$

where $M_{\text{TRM}} \sim 0.03M_{\text{rs}}$ for a TRM in a paleofield of 100 μT , μ_0 is the permeability of free space, and ρ is the crustal density. Assuming the crust is made out of NWA7325-like material (for which $\rho \sim 3000 \text{ kg m}^{-3}$ and $M_{\text{rs}} = 10^{-3} \text{ Am}^2 \text{ kg}^{-1}$; see Table S3), we estimate $B_{\text{crust}} < 0.1 \mu\text{T}$. Therefore, it is entirely possible that the parent body of NWA 7325 generated a strong ($> \sim 100 \mu\text{T}$) dynamo magnetic field prior to the meteorite's period of final cooling time, a weaker dynamo ($< 1.7 \mu\text{T}$) during this cooling period, or an arbitrarily strong field after this period.

Our nominal $< 1.7 \mu\text{T}$ constraint is also consistent with Mercury's $\sim 0.1\text{--}0.7 \mu\text{T}$ present day surface field (Stevenson, 2010). However, this coincidence may have limited significance for the hypothesis that NWA 7325 originates from Mercury given that the age of its magnetization is $\sim 4500 \text{ Ma}$, not the present. Furthermore, NWA 7325's 4563 Ma formation age would make it far older than the oldest large-scale surfaces dated by crater counting (4000–4100 Ma) on Mercury (Koefoed et al., 2016). Therefore, although our magnetic data are formally consistent with formation of NWA 7325 in Mercury's present-day field, we consider a Mercury origin for NWA 7325 unlikely.

Our measurements on NWA 7325 require that all magnetic field sources at the meteorite's location were below $\sim 1.7 \mu\text{T}$ at $4563.09 \pm 0.26 \text{ Ma}$. This includes core dynamo, nebular, and impact-generated fields, which have been predicted under some conditions to well exceed our nominal 1.7 μT constraint. A near-zero nebular field at this time in turn supports the recent conclusion that the solar nebula had likely dissipated by 4 million years after CAIs (Wang et al., 2017). Our field constraint also applies to that of the solar wind, which has been proposed to be the source of the strong middle temperature (MT) component observed in CV chondrites (Tarduno et al., 2016). In fact, our results show that the intensity of the solar wind field was also below the ~ 60 and $\sim 3 \mu\text{T}$ paleointensities measured from the CV chondrites Allende and Kaba, respectively (Carpözen et al., 2011; Gattacceca et al., 2016). On the other hand, the latter high

paleointensities are consistent with a core dynamo origin for this magnetization. Finally, our analysis of NWA 7325 demonstrates that we can successfully and sensitively identify near-zero field constraints from meteorites. This gives greater confidence to the many other meteorite studies that have inferred the existence of past magnetic fields.

8. Summary

- NWA 7325 is an ungrouped achondrite from a differentiated planetesimal
- The meteorite last cooled from above the 780°C in essentially zero-field conditions (nominally $<1.7 \mu\text{T}$, or possibly with an upper limit as high as $<9 \mu\text{T}$ considering uncertainties in non-thermal paleointensity calibration coefficients).
- This requires that any dynamo field as well as any externally generated fields like those from the nebula and solar wind were $<1.7 \mu\text{T}$ at an estimated age $4563.09 \pm 0.26 \text{ Ma}$.
- NWA 7325 is the first small body achondrite recognized to be unmagnetized despite likely having originated from a body that likely formed a metallic core.
- Our paleointensity constraint on NWA 7325 is consistent with the field strength of present-day Mercury, but the meteorite's great age makes a Mercury origin unlikely.
- Our ability to recognize unmagnetized achondrites provides a control that builds confidence in the widespread identification of magnetized achondrites from other bodies.

Acknowledgements. We thank S. Ralew for providing samples of NWA 7325 for study and Neel Chatterjee for assistance with electron microscopy measurements. We gratefully acknowledge the use of the LeRoy Eyring Center for Solid State Science at ASU. BPW thanks the NASA Emerging Worlds program grant #NNX15AH72G, the NASA Solar System Exploration and Research Virtual Institute grant #NNA14AB01A, the U. S. Rosetta program (JPL award #NM0710889), and a generous gift from Thomas F. Peterson, Jr. BD thanks the MIT UROP for support. JG acknowledges funding from the ANER (grant ANR-14-CE33-0012) and the Marie Curie People Programme (FP7/2007-2013, REA grant 29835). We also thank the Rutgers Paleomagnetism Laboratory and the Institute of Rock Magnetism for enabling some of our rock magnetic measurements. Measurements at NSLS, BNL were supported by the U.S. DOE under contract #DE-AC02-98CH10886.

References

- Almeida, T.P., Muxworthy, A.R., Kovács, A., Williams, W., Brown, P.D., Dunin-Borkowski, R.E., 2016. Direct visualization of the thermomagnetic behavior of pseudo–single-domain magnetite particles. *Sci. Adv.* 2, e1501801.
- Archer, G.J., Walker, R.J., Irving, A.J., 2015. Highly siderophile element and ^{187}Re – ^{187}Os isotopic systematics of ungrouped achondrite Northwest Africa 7325. *Lunar Planet Sci. Conf. XLVI*, abstract #1987.
- Baker, D.R., Mancini, L., Polacci, M., Higgins, M.D., Gualda, G.A.R., Hill, R.J., Rivers, M.L., 2012. An introduction to the application of X-ray microtomography to the three-dimensional study of igneous rocks. *Lithos* 148, 262–276.
- Barrat, J.A., Greenwood, R.C., Verchovsky, A.B., Gillet, P., Bollinger, C., Langlade, J.A., Liorzou, C., Franchi, I.A., 2015. Crustal differentiation in the early solar system: Clues from the unique achondrite Northwest Africa 7325 (NWA 7325). *Geochim. Cosmochim. Acta* 168, 280–292.
- Berndt, T., Muxworthy, A.R., Fabian, K., 2016. Does size matter? Statistical limits of paleomagnetic field reconstruction from small rock specimens. *J. Geophys. Res.* 121, doi:10.1002/2015JB012441.
- Bryson, J.F.J., Nichols, C.I.O., Herrero-Albillos, J., Kronast, F., Kasama, T., Alimadadi, H., van der Laan, G., Nimmo, F., Harrison, R.J., 2015. Long-lived magnetism from solidification-driven convection on the pallasite parent body. *Nature* 517, 472–475.
- Carpözen, L., Weiss, B.P., Elkins-Tanton, L.T., Shuster, D.L., Ebel, D.S., Gattacceca, J., 2011. Magnetic evidence for a partially differentiated carbonaceous chondrite parent body. *Proc. Natl. Acad. Sci. USA* 108, 6386–6389.
- Cerantola, V., Walte, N.P., Rubie, D.C., 2015. Deformation of a crystalline olivine aggregate containing two immiscible liquids: Implications for early core–mantle differentiation. *Earth Planet. Sci. Lett.* 417, 67–77.
- Cisowski, S.M., 1991. Remanent magnetic properties of unbrecciated eucrites. *Earth Planet. Sci. Lett.* 107, 173–181.
- Cournède, C., Gattacceca, J., Gounelle, M., Rochette, P., Weiss, B.P., Zanda, B., 2015. An early solar system magnetic field recorded in CM chondrites. *Earth Planet. Sci. Lett.* 410, 62–74.

581 Dunlop, D.J., 2002. Theory and application of the Day plot (M_{rs}/M_s versus H_{cr}/H_c) - 1.
 582 Theoretical curves and tests using titanomagnetite data. J. Geophys. Res. 107,
 583 doi:10.1029/2001JB000486.
 584 Dunlop, D.J., Özdemir, O., 1997. Rock magnetism: Fundamentals and Frontiers. Cambridge
 585 University Press, New York.
 586 Einsle, J.F., Harrison, R.J., Kasama, T., Conbhuí, P.Ó., Fabian, K., Williams, W., Woodland, L.,
 587 Fu, R.R., Weiss, B.P., Midgley, P.A., 2016. Multi-scale three-dimensional characterisation of
 588 iron particles in dusty olivine: implications for paleomagnetism of chondritic meteorites. Am.
 589 Mineral. 101, 2070-2084.
 590 Elkins-Tanton, L.T., Weiss, B.P., Zuber, M.T., 2011. Chondrites as samples of differentiated
 591 planetesimals. Earth Planet. Sci. Lett. 305, 1-10.
 592 Fischer, S.R., Fu, R.R., Weiss, B.P., Li, L., Gattacceca, J., Sonzogni, C., 2013. Paleomagnetic
 593 detection of magnetic fields on a differentiated asteroid during the dynamo epoch. Fall AGU
 594 Meeting, abstract #GP41D-1166.
 595 Formisano, M., Federico, C., De Angelis, S., De Sanctis, M.C., Magni, G., 2016. A core dynamo
 596 in Vesta? Mon. Not. R. Astr. Soc. 458, 695-707.
 597 Fu, R.R., Weiss, B.P., Shuster, D.L., Gattacceca, J., Grove, T.L., Suavet, C., Lima, E.A., Li, L.,
 598 Kuan, A.T., 2012. An ancient core dynamo in asteroid Vesta. Science 338, 238-241.
 599 Gattacceca, J., Weiss, B.P., Gounelle, M., 2016. New constraints on the magnetic history of the
 600 CV parent body and the solar nebula from the Kaba meteorite. Earth Planet. Sci. Lett. 455, 166-
 601 175.
 602 Harrison, R.J., Feinberg, J.M., 2008. FORCinel: An improved algorithm for calculating first-
 603 order reversal curve distributions using locally weighted regression smoothing. Geochem.
 604 Geophys. Geosyst. 9, Q05016.
 605 Irving, A.J., Kuehner, S.M., Bunch, T.E., Ziegler, K., Chen, G., Herd, C.D.K., Conrey, R.M.,
 606 Ralew, S., 2013. Ungrouped mafic achondrite Northwest Africa 7325: A reduced, iron-poor
 607 cumulate olivine gabbro from a differentiated planetary parent body. Lunar Planet Sci. Conf.
 608 XLIV, abstract #2164.
 609 Kirschvink, J.L., 1980. The least-squares line and plane and the analysis of paleomagnetic data:
 610 examples from Siberia and Morocco. Geophys. J. R. Astr. Soc. 62, 699-718.

611 Kirschvink, J.L., Kopp, R.E., Raub, T.D., 2008. Rapid, precise, and high-sensitivity acquisition
 612 of paleomagnetic and rock-magnetic data: Development of a low-noise automatic sample
 613 changing system for superconducting rock magnetometers. *Geochem. Geophys. Geosyst.* 9,
 614 Q05Y01, doi:10.1029/2007GC001856.

615 Koefoed, P., Amelin, Y., Yin, Q.-Z., Wimpenny, J., Sanborn, M.E., Iizuka, T., Irving, A.J., 2016.
 616 U–Pb and Al–Mg systematics of the ungrouped achondrite Northwest Africa 7325. *Geochim.*
 617 *Cosmochim. Acta* 183, 31-45.

618 Krauss, G., 2005. *Steels: Processing, Structure, and Performance*. ASM International, Materials
 619 Park, OH.

620 Kruijer, T.S., Touboul, M., Fischer-Gödde, M., Bermingham, K.R., Walker, R.J., Klein, T.,
 621 2014. Protracted core formation and rapid accretion of protoplanets. *Science* 344, 1150-1154.

622 Moskowitz, B.M., 1981. Methods for estimating Curie temperatures of titanomagnetites from
 623 experimental J_s - T data. *Earth Planet. Sci. Lett.* 53, 84-88.

624 Nichols, C.I.O., Bryson, J.F.J., Herrero-Albillos, J., Kronast, F., Nimmo, F., Harrison, R.J., 2016.
 625 Pallasite paleomagnetism: Quiescence of a core dynamo. *Earth Planet. Sci. Lett.* 441, 103-112.

626 Nimmo, F., 2009. Energetics of asteroid dynamos and the role of compositional convection.
 627 *Geophys. Res. Lett.* 36, doi:10.1029/2009GL037997.

628 Özdemir, Ö., Dunlop, D.J., Moskowitz, B.M., 2002. Changes in remanence, coercivity and
 629 domain state at low temperature in magnetite. *Earth Planet. Sci. Lett.* 194, 343-358.

630 Rückriemen, T., Breuer, D., Spohn, T., 2015. The Fe snow regime in Ganymede's core: A deep-
 631 seated dynamo below a stable snow zone. *J. Geophys. Res.* 120, 1095-1118.

632 Ruzicka, A., 2014. Silicate-bearing iron meteorites and their implications for the evolution of
 633 asteroidal parent bodies. *Chemie der Erde* 74, 3-48.

634 Scheinberg, A., Elkins-Tanton, L.T., Schubert, G., Bercovici, D., 2015a. Core solidification and
 635 dynamo evolution in a mantle-stripped planetesimal. *J. Geophys. Res.* 121,
 636 doi:10.1002/2015JE004843.

637 Scheinberg, A., Fu, R.R., Elkins-Tanton, L.T., Weiss, B.P., 2015b. Asteroid differentiation:
 638 Melting and large-scale structure, in: Michel, P., DeMeo, F., Bottke, W.F. (Eds.), *Asteroids IV*.
 639 University of Arizona Press, Tucson, pp. 533-552.

640 Sterenborg, M.G., Crowley, J.W., 2013. Thermal evolution of early solar system planetesimals
 641 and the possibility of sustained dynamos. *Phys. Earth Planet. Inter.* 214, 53-73.

642 Stevenson, D.J., 1983. Planetary magnetic fields. *Rep. Progr. Phys* 46, 555-620.
 643 Stevenson, D.J., 2010. Planetary magnetic fields: Achievements and prospects. *Space Sci. Rev.*
 644 152, 651-664.
 645 Suavet, C., Weiss, B.P., Grove, T.L., 2014. Controlled-atmosphere thermal demagnetization and
 646 paleointensity analyses of extraterrestrial rocks. *Geochem. Geophys. Geosyst.* 15,
 647 doi:10.1002/2013GC005215.
 648 Sutton, S.R., Wirick, S., Goodrich, C.A., 2014. Ungrouped achondrite NWA 7325: Titanium,
 649 vanadium, and chromium XANES of mafic silicates record highly-reduced origin. *Lunar Planet*
 650 *Sci. Conf. XLV*, abstract #1275.
 651 Swartzendruber, L.J., Itkin, V.P., Alcock, C.B., 1991. The Fe-Ni (iron-nickel) system. *J. Phase*
 652 *Equilib.* 12, 288-312.
 653 Tarduno, J.A., Cottrell, R.D., Nimmo, F., Hopkins, J., Voronov, J., Erickson, A., Blackman, E.,
 654 Scott, E.R.D., McKinley, R., 2012. Evidence for a dynamo in the main group pallasite parent
 655 body. *Science* 338, 939-942.
 656 Tarduno, J.A., O'Brien, T.M., Smirnov, A.V., 2016. Does the magnetization of CV meteorites
 657 record a parent body core dynamo? *Lunar Planet Sci. Conf. XLVII*, abstract #2609.
 658 Tikoo, S.M., Weiss, B.P., Cassata, W.S., Shuster, D.L., Gattacceca, J., Lima, E.A., Suavet, C.,
 659 Nimmo, F., Fuller, M., 2014. Decline of the lunar core dynamo. *Earth Planet. Sci. Lett.* 404, 89-
 660 97.
 661 Wang, H., Weiss, B.P., Bai, X.-N., Downey, B.G., Wang, J., Wang, J., Suavet, C., Fu, R.R.,
 662 Zucolotto, M.E., 2017. Lifetime of the solar nebula constrained by meteorite paleomagnetism.
 663 *Science* 355, 623-627.
 664 Wang, J., Chen, Y.-C.K., Yuan, Q., Tkachuk, A., Erdonmez, C., Hornberger, B., Feser, M.,
 665 2012. Automated markerless full field hard x-ray microscopic tomography at sub-50 nm 3-
 666 dimension spatial resolution. *Appl. Phys. Lett.* 100, 143107.
 667 Weber, I., Morlok, A., Bischoff, A., Hiesinger, H., Ward, D., Joy, K.H., Crowther, S.A.,
 668 Jasterzebski, N.D., Gilmour, J.D., Clay, P.L., Wogelius, R.A., Greenwood, R.C., Franchi, I.A.,
 669 Münker, C., 2016. Cosmochemical and spectroscopic properties of Northwest Africa 7325—A
 670 consortium study. *Meteorit. Planet. Sci.* 51, 3-30.

671 Weisberg, M.K., McCoy, T.J., Krot, A.N., 2006. Systematics and evaluation of meteorite
 672 classification, in: Lauretta, D.S., McSween, H.Y. (Eds.), *Meteorites and the Early Solar System*
 673 II. University of Arizona, Tucson, pp. 19-52.
 674 Weiss, B.P., Gattacceca, J., Stanley, S., Rochette, P., Christensen, U.R., 2010. Paleomagnetic
 675 records of meteorites and early planetesimal differentiation. *Space Sci. Rev.* 152, 341-390.
 676 Weiss, B.P., Gattacceca, J., Suavet, C., Irving, A.J., 2013. Magnetic and nonmagnetic
 677 differentiated planetesimals. Workshop on Planetesimal Formation and Differentiation, abstract
 678 #8047.
 679 Wilson, L., Keil, K., 2012. Volcanic activity on differentiated asteroids: A review and analysis.
 680 *Chemie der Erde* 72, 289-321.
 681 Winklhofer, M., Fabian, K., Heider, F., 1997. Magnetic blocking temperatures of magnetite
 682 calculated with a three-dimensional micromagnetic model. *J. Geophys. Res.* 102, 22695-22709.
 683
 684

686
 687 **Fig. 1.** Electron microscopy analyses of NWA 7325. (A) Overview BSEM image showing
 688 dominant occurrence of FeNi and FeS grains as submicrometer inclusions in plagioclase (plag).
 689 Plagioclase commonly exhibits reaction relationships with adjacent diopside (diop), suggestive
 690 of a partial remelting event after formation. (B) BSEM image of rare CrFeNi metal grains. (C)
 691 BSEM image of rare secondary Fe oxide metal grains associated with an FeNi-FeS assemblage.

(D) BSEM image showing plagioclase (dark gray) with disseminated particles of Fe-Ni metal (white) and troilite (FeS; light gray). The metal and sulfide phases form intergrown assemblages with diameters ranging from tens of nanometers to several micrometers. (E) Higher magnification BSEM of boxed region in (D) showing composite FeS and FeNi particles in plagioclase with associated voids (dark zones adjacent to some metal and sulfide particles). (F) Bright-field (BF) TEM image of an FeS crystal (medium grey) in contact with two FeNi particles (dark grey) and a void (lightest grey) within a polycrystalline aggregate of anorthitic plagioclase (plag). Albite twins in the anorthite are visible as bands of alternating diffraction contrast (arrows). The inset convergent-beam electron diffraction (CBED) pattern, from the lower FeNi particle, is indexed as a $\langle 111 \rangle$ zone axis pattern for a body-centered cubic ($Im\bar{3}m$) α -FeNi. The mottled contrast visible in the central beam and diffraction spots of the CBED pattern illustrates local lattice strain within the FeNi crystal. (G) BF TEM image of the upper FeNi particle from (F) showing a lamellar twin bounded by planar twin boundaries (arrows). The FeNi_a twin domain (dark) is imaged along the $[11\bar{3}]$ zone axis and the FeNi_b domain is imaged nearly along $[11\bar{3}]$. Local variations in diffraction contrast within the twin domains indicate local lattice strain associated with a high density of defects. (H) Selected area electron diffraction (SAED) pattern of the twinned crystal shown in (G). The pattern consists of two $[11\bar{3}]$ zone axis patterns related by a $\sim 28^\circ$ rotation about $[11\bar{3}]$. The patterns were collected with domain FeNi_a aligned along $[11\bar{3}]$ and domain FeNi_b slightly off $[11\bar{3}]$. As shown in Fig. S2, this twin relationship is consistent with the martensitic phase transition from γ -FeNi (fcc) to α_2 -FeNi (bcc).

Fig. 2. Saturation magnetization of NWA 7325 subsample SR1 as a function of temperature during heating up to 800°C and cooling down to 350°C. Fitting the warming data over the interval of 400-750°C with the extrapolation method of Moskowitz (1981) yields a mean martensite-finish temperature (i.e., martensite-to-austenite transition temperature) of 762-796°C. Although this method is designed to quantify Curie temperatures, it should give an accurate estimate of martensite-finish temperature because the latter is very close to martensite's 780°C Curie temperature. These data were acquired on fresh (unheated) subsample SR1 following all other rock magnetic measurements on this subsample (Figs. 4 and S2).

Fig. 3. TXM imaging of NWA 7325. (A) Mosaic TXM image of a silicate grain from NWA 7325 stitched from $40\ \mu\text{m} \times 40\ \mu\text{m}$ images (2×2 pixel binning), front view. (B) Side view of the same NWA 7325 grain. (C, D) Zoom-in to areas containing fine iron-bearing particles. (E-G) Selected frames (0° , 45° , and 90°) from the tomographic three-dimensional reconstruction movie (Movie S1) of a $40\ \mu\text{m}$ cylindrical volume of NWA 7325. In (A)-(D), higher concentrations of iron are darker, whereas in (E) to (G), higher concentrations of iron are brighter.

735

736 **Fig. 4.** Hysteresis properties of NWA 7325. (A) Room temperature hysteresis loop measured
 737 for fresh NWA 7325 subsample SR1. Shown is sample moment as a function of applied field.

738 The black curve represents the hysteresis loop after subtraction of paramagnetic and diamagnetic
 739 contributions. The gray curve represents the raw (uncorrected) data. (B) Dunlop-Day plot

740 (Dunlop, 2002) showing the ratio of coercivity of remanence to coercivity (B_{cr}/B_c) versus the
 741 ratio of saturation remanence to saturation magnetization (M_{rs}/M_s) for NWA 7325 subsamples.

742 Single domain (SD), pseudo single domain (PSD) and multidomain (MD) ranges calculated for
 743 magnetite are indicated, as well as mixing lines between SD and MD and between SD and

744 superparamagnetic (SP) of different sizes are also indicated. Unheated subsamples and
 745 subsamples previously heated are denoted by black and red circles, respectively.

746

Fig. 5. FORC diagrams for NWA 7325. Shown is B_u (a measure of magnetostatic interactions) on the ordinate versus B_c (coercivity) on the abscissa. The color bar shows the probability density of hysterons belonging to a given combination of B_u and B_c . (A) Fresh subsample SR3. (B) Subsample 7B3C, previously heated to 400°C in IW conditions. (C) Subsample SR1a, previously heated to 800°C in IW-3 conditions.

Fig. 6. AF and thermal demagnetization experiments on NRM in ungrouped achondrite NWA 7325. Shown is a two-dimensional projection of the endpoints of the NRM vector during laboratory demagnetization. Solid (open) symbols represent end points of magnetization projected onto the horizontal N-E (vertical Z-E) planes (see left legend). Blue and green arrows denote LC/LT and MC/MT component directions, respectively. Sensor noise of MIT 2G SRM and quartz disk holder moment are shown by two tiny black boxes in right legend. (A) Subsample B71. (B) Subsample B73. (C) Subsample B75. (D) Subsample B79. Peak fields and temperatures for demagnetization steps are labeled in mT and °C, respectively.

Fig. 7. NRM component directions observed in NWA 7325. Shown is an equal area stereographic projection of Fisher mean directions (symbols) and associated maximum angular deviation values (surrounding ellipses) obtained from principal component analyses. (A) LC (circles) and LT (squares) component directions. (B) MC (circles) and MT (squares) component directions. (C) HC (circles) and HT (squares) magnetization directions.

Fig. 8. Paleointensity experiments on selected subsamples of NWA 7325. (A, B) NRM lost as a function of ARM (AF of 200 mT, bias field of 0.2 mT) lost during AF demagnetization for subsample B71 (A) and B73 (B). (C, D) NRM lost as a function of near-saturation IRM (200 mT) lost during AF demagnetization for subsamples B71 (C) and B73 (D). (E, F) NRM lost as a function of TRM (800°C and 30 μ T field) lost during thermal demagnetization for subsamples B78 (E) and B79 (F). Points corresponding to the LC/LT and MC/MT components and HC/HT magnetization are colored blue, green and red, respectively. Selected AF and thermal demagnetization steps are labeled.

780

781 **Fig. 9.** Thermal paleointensity fidelity test using directional criteria. Shown is the fidelity by
 782 which an 800°C total TRM can be isolated by thermal demagnetization. (A, B) Thermal
 783 demagnetization of TRM acquired in a vertical (positive Z) 3 μT field for samples B77 and B78.
 784 (C, D) Thermal demagnetization of TRM acquired in a vertical (positive Z) 0.3 μT field for
 785 samples B77 and B78. Red arrows show where demagnetization of the TRM is visible.

786

787

788

789 **Fig. 10.** Thermal demagnetization of NRM (black curve and points) compared to total TRM
 790 acquired from cooling in a field of 800 $^{\circ}\text{C}$ in fields of 30, 3, and 0.3 μT (blue, red, and green
 791 curves and points, respectively) for sample B79. (A) All data, expressed as moment as a
 792 function of temperature. (B) Zoom into NRM and 0.3 μT TRM data from (A). (C) Moment
 793 normalized to moment at room temperature before starting thermal demagnetization. NRM
 794 loses directional stability by a demagnetization temperature of 300 $^{\circ}\text{C}$; directionally unstable data
 795 at higher demagnetization temperatures are connected with a dashed line.

796

797