1 ENVIRONMENTAL SIGNAL PROPAGATION IN SEDIMENTARY SYSTEMS
2 ACROSS TIMESCALES

3

4 Brian W. Romans¹, Sébastien Castelltort², Jacob A. Covault³, Andrea Fildani⁴, and J.P. Walsh^{5,6}

5 ¹*Department of Geosciences, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061*

6 ²*Section des Sciences de la Terre et de l'Environnement, Université de Genève, 1205 Genève, Suisse*

7 ³*Chevron Energy Technology Company, Houston, TX 77002*

8 ⁴*Statoil RDI, Research Center Austin, TX 78730*

9 ⁵*East Carolina University, Greenville, NC 27858*

10 ⁶*UNC Coastal Studies Institute, Wanchese, NC 27981*

11

12 keywords: source-to-sink; sediment-routing systems; Earth-surface dynamics; stratigraphy; geomorphology; tectonics;
13 paleoclimate

14

15 **Abstract**

16 Earth-surface processes operate across erosionally dominated landscapes and
17 deliver sediment to depositional systems that can be preserved over a range of timescales.
18 The geomorphic and stratigraphic products of this source-to-sink sediment transfer record
19 signals of external environmental forcings, as well as internal, or autogenic, dynamics of
20 the sedimentary system. Here, we evaluate environmental signal propagation across
21 sediment-routing systems with emphasis on sediment supply, Q_s , as the carrier of up-
22 system forcings. We review experimental, numerical, and natural examples of source-to-
23 sink sediment routing and signal propagation during three timescales: (1) Historic, which
24 includes measurement and monitoring of events and processes of landscape change and
25 deposition during decades to centuries; (2) Centuries to several millions of years, referred
26 to as intermediate timescale; and (3) Deep time. We discuss issues related to autogenic

dynamics of sediment transport, transient storage, and release that can introduce noise, lags, and/or completely mask signals of external environmental forcings. We provide a set of conceptual and practical tools for evaluating sediment supply within a source-to-sink context, which can inform interpretations of signals from the sedimentary record. These tools include stratigraphic and sediment-routing system characterization, sediment budget determination, geochronology, detrital mineral analysis (e.g., thermochronology), comparative analog approaches, and modeling techniques to measure, calculate, or estimate the magnitude and frequency of external forcings compared to the characteristic response time of the sediment-routing systems.

1 Introduction

1.1 What is an 'Environmental Signal'?

From the perspective of sedimentary system analysis, signals are changes in sediment production, transport, or deposition, that originate from perturbations of environmental variables such as precipitation, sea level, rock uplift, subsidence, and human modifications. The origin of the perturbations can be 'natural' when they relate to tectonic and climatic processes that have happened over the course of Earth's history, or 'anthropogenic' if they are linked with human actions. Environmental signals occur over many temporal scales, ranging from several hours to millions of years in response to tectonic and climate changes. Signals involve a large range of spatial scales such as localized precipitation affecting small catchments to eustatic sea-level change that affects the globe.

50 An environmental signal can trigger a response of the Earth's surface in the form of
51 erosion, sediment transport, and deposition, and the surface response may be local
52 initially and further afield eventually as it propagates away. A sea-level fall, for example,
53 can create local incision and shoreline regression, but also up-system knickpoint
54 migration and down-system deposition in the deep sea. Similarly, an increase in
55 precipitation can create a wave of incision, alluvial aggradation, and eventually a pulse of
56 sediment discharge to the ocean. The overarching challenge of geomorphology and
57 stratigraphy is to invert the history of environmental signals from landscape and rock
58 records.

59 The transfer, or propagation, of signals is generally examined in the down-system
60 direction, as this is the dominant direction of mass transfer (e.g., [Castelltort and Van Den](#)
61 [Driessche, 2003](#); [Allen, 2008a](#), [Jerolmack and Paola, 2010](#)). However, up-system signal
62 propagation driven by base level change has long been considered in the interpretation of
63 the sedimentary record (e.g., [Fisk, 1944](#)), is important for distributive systems (e.g.,
64 backwater effect in deltas, [Lamb et al., 2012](#)), and is the subject of theoretical work
65 ([Voller et al., 2012](#)).

66 Environmental signals are potentially preserved in the geomorphic expression of
67 landscapes around us, as well as in the stratigraphic record of depositional basins. This
68 review examines how signals propagate within the context of sediment routing systems
69 with emphasis on the nature of sediment supply, or Q_s , as the indicator of up-system
70 forcings (Fig. 1A) ([Allen et al., 2013](#)). We think that reconstructing the rates and
71 magnitudes of signal-generating processes from stratigraphy requires consideration of the
72 nature of system response, and the potential modification of the original signal. It is also

important to recognize that signals can be masked or significantly altered by what can be referred to as ‘noise.’ In the present context, ‘noise’ has the broad meaning of any perturbation of the primary signal of interest, irrespective of its origin, frequency, or magnitude. It is one fundamental goal of stratigraphy to disentangle signal from noise, but what can be considered noise at one timescale may represent a signal at another. One notable type of noise is the result of internal, self-organized, dynamics of a sediment routing system (e.g., [Jerolmack and Paola, 2010](#)), that can potentially ‘shred’ environmental signals as a result of their large magnitude and period relative to the primary signal of interest (e.g., [Jerolmack and Paola, 2010](#); [Wang et al., 2011](#)).

Deciphering signals has obvious implications for the meaning of the sedimentary record of Earth history: what do sediments and rocks tell us about the past? However, understanding the signal-to-noise character of the sedimentary record is also relevant to the prediction of land-to-sea export and burial of terrestrial organic carbon (e.g., [Kao et al., 2014](#); [Leithold et al., this volume](#)), landscape resiliency and hazard management (e.g., [Anthony and Julian, 1999](#)), prediction of depositional systems for natural resource exploration and production ([Bhattacharya et al., this volume](#)), and response of hydrological systems to global climate change (e.g., [Syvitski, 2003](#)). We do not attempt to solve all the outstanding issues related to signal propagation and preservation in this contribution. Our goal is to provide the general Earth scientist a thorough review of the interesting and enigmatic questions and to promote a broader understanding that might attract other researchers to this multidisciplinary field of study.

1.2 *Importance of Timescale of Investigation*

We emphasize the importance of timescale in this review because of its association with the processes of signal generation, propagation, preservation, and analysis. The evaluation of signals requires consideration of the timescale(s) particularly in the context of internally generated ‘noise.’ Also, some signals occur over long durations (e.g., uplift and exhumation of a mountain belt) and, therefore, require a correspondingly long record from which to deduce the signal.

How do we put historical (past few centuries) measurements and observations within the framework of landscapes and stratigraphy constructed over timescales $\geq 10^3$ yr? Put another way, how do we accurately estimate short-term rates from geologic archives that have longer-term temporal resolution? For example, the < centennial stratigraphic record contains information about short-lived events, such as hurricane deposits, which can be reliably dated. The challenge is to extract meaningful insight from such records in the deeper past.

We organize this review of signal propagation and preservation within the context of three important timescales that span a minimum of seven orders of temporal magnitude: (1) Historic, which includes measurement and monitoring of events and processes of landscape change ($< 10^2$ yr); (2) Centennial to several million years, herein referred to as the intermediate timescale (10^2 - 10^6 yr); and (3) Deep time ($\geq 10^7$ yr) (Fig. 2). These timescales are discussed in terms of age of the system as well as duration or period of forcing. The timescale of investigation also influences the application of concepts of steady state, response time, and other system dynamics indicators, which will be discussed in detail in the intermediate timescale Section 3.

1.3 Sediment Routing Systems

Earth-surface processes operate within erosionally dominated landscapes coupled with depositional systems that can be preserved over a range of timescales. A simple and elegant way to consider an integrated sedimentary system was presented by [Schumm \(1977\)](#) wherein he subdivided a system into three spatial zones of dominant mass-flux behavior: denudation/erosion, transfer, and accumulation/deposition. Similar to [Castelltort and Van Den Driessche \(2003\)](#) and [Sadler and Jerolmack \(2014\)](#), we depict a generic sediment routing system in cross section and denote the prominent environmental forcings of interest in this review (Fig. 1B). The ‘transfer zone’ is assumed to be the segment of the sedimentary system that is neither net-denudational nor net-accumulative; rather, it is characterized by the balance between sediment removal/remobilization and sediment storage that feeds or starves down-system accumulation zones. Thus, this zone typically does not produce much sediment via bedrock erosion and, over sufficiently long timescales, it will transfer more mass than it produces or accumulates. We consider the morphology and process history of the transfer zone as an indicator of system response to perturbations, which is important for reconstructing paleo-sediment routing systems. A spatial scale is not shown on Figure 1B because the lengths of these zones vary significantly from system to system (e.g., [Somme et al., 2009](#)). For example, small/high-relief sediment routing systems (10-50 km long) typically have very short transfer zones, which results in negligible transient sediment storage, whereas large, continental-scale systems (100-1000 km long) commonly have long transfer zones containing sediment sinks that can store sediment temporarily or permanently given favorable subsidence

conditions. The magnitudes and timescales of such mass transfer-and-storage behavior, which can be addressed through the estimation of sediment budgets, are fundamental to the propagation of signals. We focus on the down-system mass transfer of inorganic, dominantly siliciclastic, particulates through water-sediment flows and refer the reader to [Leithold et al. \(this volume\)](#) for a review of organic-carbon dynamics of source-to-sink systems. Additionally, we acknowledge the important and unique sediment-supply characteristics of glaciated systems but do not distinguish them here and refer the reader to [Jaeger and Koppes \(this volume\)](#).

2 Sedimentary Process-Response Over Historical ($<10^2$ yr) Timescales

Signals at the historical timescale are the result of individual events that last hours to days (e.g., floods, storms, and earthquakes) to longer-lived changes that occur over decades (e.g., watershed deforestation and other land-use alterations) (Fig. 1). The mechanisms involved in the formation and/or propagation of such signals from source to sink include a range of hillslope (e.g., sheetwash, landsliding), glacial, fluvial, volcanic, oceanic (e.g., tides and storm wave) processes and subaqueous mass movements and flows (e.g., turbidity currents). Data from instruments have provided opportunities to measure and quantify sedimentary dynamics, and the stratigraphic record is also examined to link process to product over longer time. The timescale of this section covers what some consider to be the period of significant anthropogenic influence on Earth surface systems (onset of the Industrial Revolution, or ~250 yr before present; [Crutzen and Stoermer, 2000](#); [Zalasiewicz et al., 2000](#)).

We identify four potential challenges to leveraging historic records to understand millennial-scale and deeper-time geology. First, ancient events might have been non-actualistic; i.e., there is no adequate modern analog regarding process (Myrow and Southard, 1996). For example, globally distributed strata that were produced by the Cretaceous bolide impact (e.g., Bralower et al., 1998). Second, although a recent event may have had profound impact on society (e.g., 2005 Hurricane Katrina, 2011 Mississippi River flood), the geological record produced might be negligible or non-existent, depending on many factors including spatial variation in supply and erosion (Turner et al., 2006; Walsh et al., 2006; Goni et al., 2007; McKee and Cherry, 2009; Reed et al., 2009; Allison et al., 2010; Falcini et al., 2012; Kolker et al., 2014; Xu et al., 2014b). Third, the observation of modern sedimentary processes shows that strata are often destroyed within years after deposition as a result of physical and/or biogenic reworking (Wheatcroft et al., 2007 and references therein). Finally, there is the problem of discontinuous sedimentation and the likelihood of larger gaps in the record (i.e., time recorded as hiatus) as the time interval of sampling increases (Sadler, 1981; Sadler and Jerolmack, 2014), which will be discussed further in Section 4.

Erosional landforms provide a rich record of signals in the annual-to-centennial temporal range (e.g., Viles and Goudie, 2003), but the primary goal of our discussion is to understand signal propagation into the sedimentary record. Many studies at historical timescales are focused on specific processes and segments (e.g., hillslope erosion, shelf sedimentation) and do not strive to directly link source and sink through contemporaneous research. Moreover, simple relationships between event size (e.g., flood magnitude) and strata thickness may be the exception rather than the norm (Corbett

et al., 2014 and references therein). As a result, the source-to-sink stratigraphic connection remains a challenge in many studies despite a wealth of data.

At the shortest end of the signal transfer spectrum (<1 yr), the potential for direct communication of a sediment supply signal to a sink is greatly limited. To produce a measurable signal in the stratigraphic record of the sink, events that drive sediment redistribution must move a relatively large volume of material over a short time. To understand modern system behavior, we recommend consideration of the source signal relative to the sink size; e.g., volume of event-scale Qs versus volume capacity of a sink. Furthermore, system size can impact the timescale of the signal. For example, a flood or earthquake-driven landslide into a confined mountain lake can be captured quickly (hours to days) and potentially with little post-depositional physical and/or biogenic modification (Schillereff et al., 2014) compared to a flood of the vast Mississippi River catchment into the Gulf of Mexico, the effects of which can persist for months (Allison et al.; 2000; Kolker et al., 2014; Xu et al., 2014b). Resolving events occurring in close succession is challenging because the signals might be truncated, overprinted, or commingled (e.g., hurricanes Katrina and Rita; Goni et al., 2007 or the Morokot earthquake and ensuing flood; Carter et al., 2012).

We first discuss key processes and rates of sediment production and transfer over human timescales. We then address the storage in sedimentary sinks and high-resolution dating typical of historical timescales. Finally, we examine two well-studied modern source-to-sink systems and the specificities of stratal preservation and sediment budgets over centennial timescales.

2.1 Sediment Production and Transfer Over Historical Timescales

Sediment production and movement in catchments and river systems is often described in a time-averaged perspective with the timescale of focus related to the measurement tool employed. Annual hillslope erosion rates (in mm/yr or t/ha/yr), sediment loads (t/yr) and yields (t/km²/yr) may be used to compare and contrast systems and help evaluate their overall functioning (Milliman and Syvitski, 1992; Walling and Webb, 1996; Walling, 1999; Syvitski and Saito, 2007; Syvitski and Milliman, 2007; Milliman and Farnsworth, 2011; Covault et al., 2013). Loads and yields are commonly measured with stream gauges (e.g., Milliman and Farnsworth, 2011), which can be used to evaluate catchment erosion rates and/or alluvial storage (e.g., Meade et al., 1990; Walling and Collins, 2008). There are significant challenges to quantifying sediment transfer to the sea by rivers, particularly of large systems because of tidal influence on transport calculations and sediment storage in the lower river (e.g., Milliman et al., 1984; Allison et al., 2012). Historical measurements can be biased as a result of their limited duration or influences of anthropogenic catchment modification, including construction of dams and other land-use activities associated with agriculture, construction, and mining (Wilkinson and McElroy 2007; Milliman and Farnsworth, 2011). Erosion rates also can be measured with cosmogenically derived tracers (e.g., ¹⁰Be; discussed further in Section 3) and radiochemically dated deposits (e.g., ¹⁴C, ¹³⁷Cs or ²¹⁰Pb) from well-defined source areas (e.g., Walling and Collins, 2008). Technological advancements, specifically Light-Detection and Ranging and terrestrial laser scanners, have improved our ability to quantify morphological changes on land. Denudation rates from LiDAR, discharge measurements and ¹⁰Be indicate variability depending on slope and other

factors (commonly <0.5 mm/yr, but locally >3 mm/yr) (e.g., [Hovius et al., 1997](#); [Aalto et al., 2003](#); [Roering et al., 2007](#); [Korup et al., 2014](#)). The contextual and temporal knowledge of precipitation and catchment characteristics usually exceeds what can be measured or inferred in ancient systems, as will be discussed in subsequent sections.

Water-driven transport, especially during intense floods, can generate recognizable sedimentary signals in sink areas. Intense rainfall and associated floods can rapidly (hours to days) move large volumes (>5 Mt [million metric tons]) of sediment through small ($<5,000$ km²) catchments to offshore depositional areas (e.g., [Sommerfield et al., 1999](#); [Hale et al., 2014](#); [Kniskern et al., 2014](#)), and larger catchments ($>50,000$ km²) can generate appreciable sediment supply (>10 s Mt) signals to the sea over the course of days to weeks (e.g., [Palinkas et al., 2005](#); [Kolker et al., 2014](#)). Subaerial and submarine landsliding and other mass movements are related to pre-conditioning factors, such as hillslope soil or rock strength, geomorphology, and short-term conditions (e.g., earthquake and hydrology) ([Dietrich et al., 1995](#); [Roering et al., 2007](#); [Strasser et al., 2006](#); [Goldfinger et al., 2012](#) and references therein).

An earthquake can disturb a catchment by increasing pore pressures and liquefying substrate, among other processes of manipulating gravitational loads on slopes, which can lead to abrupt increases in sediment loads (e.g., [Dadson et al., 2004](#)). Also, earthquake-triggered mass wasting can create conspicuous stratigraphic records in lakes and the deep sea (e.g., [Heezen and Ewing, 1952](#); [Piper and Aksu, 1997](#); [Moernaut et al., 2007](#)). Much research has explored coastal and marine sedimentary records to evaluate the recurrence interval for earthquakes and associated tsunamis (e.g., [Atwater and Hemphill-Haley, 1997](#); [Goldfinger et al., 2003](#); [Strasser et al. 2006](#); [Mournaut et al.,](#)

2007; Goldfinger et al., 2012; Barnes et al., 2013), including some recent detailed research focused on the Sumatra and Tomoko events (e.g., Szczucinski et al., 2012; Patton et al., 2013). There is still vigorous debate regarding deep-sea turbidite deposits as a reliable paleo-seismometer (e.g., Sumner et al., 2013; Atwater et al., 2014).

Over annual to centennial timescales, anthropogenic activities, such as deforestation and pollution, can create signals that become stored in sedimentary sinks (e.g., Paull et al., 2002; Cundy et al., 2003). Many natural and human factors (e.g., land use, dams) have significant influence on sediment yields and loads (Meade et al., 1990; Syvitski et al., 2005; Milliman and Farnsworth, 2011). As a consequence of the potential influence of human activities, Syvitski and Milliman (2007) included an anthropogenic factor in their BQART model that predicts global sediment flux to the oceans. Although intra-system storage might buffer some signals (i.e., low sediment delivery ratios; Phillips, 1991; Walling and Collins; 2008), catchment changes can notably increase Q_s . Damming and leveeing can significantly diminish sediment supply into sink areas (Syvitski et al., 2005; Milliman and Farnsworth, 2011 and references therein), not only precluding new strata development but also yielding land loss in some areas (e.g., Day et al., 2007; Smith and Abdel-Kader, 1988).

2.2 Storage in Sedimentary Sinks Over Historical Timescales

To evaluate the presence of signals, including events, in stratigraphic records over historic timescales, ^{210}Pb and ^{137}Cs are commonly used to date deposits or determine sediment accumulation rates (Fig. 2) (e.g., Sommerfield and Nittrouer, 1999). Bathymetric and sub-bottom observations (i.e., seismic reflection) have revealed the

geomorphic and stratigraphic complexity of subaqueous environments and such data are helpful to strategically position coring sites to obtain desired records (e.g., [Goldfinger et al., 2012](#)) or to inform spatial variability for determining sediment budgets (e.g., [Miller and Kuehl, 2010](#); [Gerber et al., 2010](#)). Recent studies have shown how time-series bathymetric analysis with multibeam may yield new insight into the intermittent nature of fluvial deposition ([Nitttrouer et al., 2008](#)) and subaqueous sediment density flows (e.g., [Smith et al., 2005](#); [Walsh et al., 2006](#); [Paull et al., 2006](#); [Xu et al., 2008](#); [Hughes Clarke et al., 2012](#)). Additionally, researchers are using innovative methods to track sediment transport and deposition, such as short-lived radiochemical tracers (i.e., ^7Be) for catchment and seaward sediment dispersal (e.g., [Sommerfield et al., 1999](#); [Dail et al., 2007](#); [Walling, 2013](#)), and mounted acoustic- and light-based sensors for measuring water and sediment movement (e.g., [Xu et al., 2004](#); [Dinehart and Burau, 2005](#); [Cacchione et al., 2006](#)). Deployed systems have provided flow measurements (i.e., velocity and sediment concentrations), which are essential to modeling sediment transport (e.g., [Traykovski et al., 2007](#); [Moriarty et al., 2014](#)). However, field measurements remain limited especially during extreme and/or rare events when most sediment is moved (e.g., [Ogston et al., 2000](#); [Talling et al., 2013](#); [Hale et al., 2014](#); [Xu et al., 2014a](#); [Stevens et al., 2014](#)).

Sedimentary filling of hollows, ponds, lakes, floodplains, estuaries, and even sinkholes can provide information about individual events or decadal-to-centennial changes in the environment. Evidence for upstream changes includes increased sedimentation rates elevated trace metals, variations in pollen, microfossil organisms and/or assemblages, trace metals, and organic compounds (e.g., estuaries: [Brush et al.,](#)

2001; Cooper et al., 2004; lakes: Noren et al., 2002, Girardclos et al., 2007; floodplains: Aalto et al., 2003; coastal deposits: Sorrel et al., 2012; Lane et al., 2011; shelves: Allison et al., 2012; deep sea: Soutar and Crill, 1977). Gilli et al. (2013) and Schillereff et al. (2014) provide reviews of flooding and climate changes from lake records.

Continental shelves, slopes, and deeper ocean segments are typically viewed as the ultimate depositional sinks, but their records are variably preserved as a result of post-depositional reworking and can be challenging to unravel (Nittrouer et al., 2007). Theory and modeling emphasize that event-layer preservation is a function of the rate of bioturbation, mixing depth, and layer thickness (Wheatcroft et al., 2007 and references therein). However, time-series coring studies of flood-related deposition on continental shelves offshore the Eel, Po, and Waipaoa sediment-routing systems have shown deep (>5 cm) biological reworking over the span of a few years (Wheatcroft et al., 2007; Tesi et al., 2012; Walsh et al., 2014). Areas of rapid sedimentation and physically reworked areas such as topset and foreset regions of clinoforms might have physical stratification preserved at depth, e.g., Amazon delta front (Kuehl et al., 1996; Sommerfield et al., 1999; Walsh et al., 2004; Rose and Kuehl, 2010). However, the presence of discontinuous, heterolithic bedforms can preclude recognition of event-specific beds (Goff et al., 2002; Walsh et al., 2014). Ocean areas with low or no dissolved oxygen inhospitable to benthic organisms (e.g., Soutar and Crill, 1977) are favorable for signal preservation (Allison et al., 2012). Continental margins and basin-margin deep-sea fans capture event records beyond historical timescales. However, during the sea-level highstand of the past several thousand years, off-shelf sediment transport is reduced in some settings (Posamentier and Vail, 1988; Covault and Graham, 2010), with shelf width serving as an important control

(Posamentier et al., 1991; Walsh and Nittrouer, 2003). As a result, limited sediment supply to some deep-sea fans has resulted in condensed sections recording few if any events at historical timescales.

2.3 Modern Sediment Routing System Examples

To further discuss historical ($<10^2$ yr) signal propagation, two differently sized sediment routing systems will be briefly discussed: the Eel River and the Ganges-Brahmaputra-Bengal system. The Eel is a small mountainous river system ($<10^3$ km²) draining northern California, USA that has received intense scrutiny during and since the Office of Naval Research STRATA FORMation on Margins program (STRATAFORM; 1995-2004; Nittrouer et al., 2007). Small mountainous rivers are important for understanding sediment flux to the sea because of the minimal onshore sediment storage (Milliman and Syvitski, 1995; Kuehl et al., 2003; Covault et al., 2011). We contrast this work with the much larger Ganges-Brahmaputra-Bengal sediment-routing system, where abundant sediment is stored onshore, on the shelf, and in the canyon today (Kuehl et al., 2005; Walsh et al., 2013). Collectively, large systems provided potentially a third to a half of the sediment to the sea prior to human alterations (Milliman and Farnsworth, 2011; Walsh et al., 2013).

The Eel River is one of the most comprehensively studied modern sediment routing system over the historical timescale (<500 yr). Its 9,400 km² catchment in a tectonically active setting of outcropping sedimentary rocks is estimated to discharge ~12-16 Mt of sediment to the sea annually (Sommerfield and Nittrouer, 1999; Warrick, 2014; Sommerfield and Nittrouer, 2014) (Fig. 3). Landslides are common in steep portions of the catchment (de la Fuente et al., 2006), but almost 70% of the load comes from the

central portion of the catchment where *mélange* outcrops are more erodible (Brown and Ritter, 1971). The largest recorded flood event occurred in December 1964, a year during which the Eel River is estimated to have discharged more than 160 Mt of sediment (Warrick, 2014). This is >13 times the annual average, with most discharge occurring over a few days. In 1995 (January and March) and 1997 (January), three floods occurred, and STRATAFORM scientists documented the deposition of a widespread layer on the shelf (Fig. 3) (Wheatcroft et al., 1997; Sommerfield and Nittrouer, 1999; Wheatcroft and Borgeld, 2000). The remarkable similarity between the flood deposits and decadal shelf sedimentation patterns demonstrate how important these events are to shelf construction. However, event and decadal sediment budgets indicate most (>50%) of the sediment is exported beyond the shelf (Fig. 3) giving testimony to the effective transport conditions associated with coherent discharge and energetic ocean conditions (Wheatcroft and Borgeld, 2000).

Instrument observations made in winter 1996-1997 revealed that a wave-enhanced sediment gravity flow associated with the floods transported an appreciable amount of sediment to the mid-shelf, exceeding other measured events by two orders of magnitude (Ogston et al., 2000; Traykovski et al., 2000). The widespread and distinctive shelf flood deposit is attributed to this mechanism; however, subsequent examination of the same deposit two years later indicated extensive reworking by physical and biological processes (Wheatcroft et al., 2007 and references therein). Although some shelf core records have stratigraphic and organic carbon evidence suggestive of older events (e.g., 1964 flood) (Sommerfield et al., 1999; Leithold et al., 2005), the documentation of post-event reworking indicates that the Eel shelf does not contain a laterally extensive or high-

fidelity record of flood signals (Goff et al., 2002; Wheatcroft et al., 2007 and references therein).

Subsequent coring and tripod research in the Eel Canyon documented the possibility of more direct sediment gravity flow to deeper water. Resuspension and transport of sediment via waves also were found to have an important control on this off-shelf export (Puig et al., 2003). Cores from the canyon indicate sedimentation is spatially and temporal complex, although export to deeper water is apparent (Mullenbach et al., 2004; Mullenbach and Nittrouer, 2006; Drexler et al., 2006). Nepheloid layers also transport fluvial sediment seaward of the Eel River mouth, allowing hemipelagic sedimentation to accumulate on the slope, but this modest input is easily reworked by the active benthic community precluding event layer formation (Alexander et al., 1999; Walsh and Nittrouer, 1999). These studies demonstrate that unravelling signals from Eel margin stratigraphic records is not straightforward, which a similar story for the Waipaoa Rivers of New Zealand (Kuehl et al., this volume). The apportionment of terrigenous sediment among shelf, slope, and deep-sea segments (Fig. 3D) suggests that shelf records might contain signals of sediment-production events that originated in the catchment, but post-depositional homogenization hampers event-scale determination.

The Ganges-Brahmaputra-Bengal is a large (1,656,000 km² catchment) sediment-routing system fed by tectonically active mountains. Sedimentation on the Bengal Fan (>2,000,000 km² depositional area), the ultimate sink for the system, has varied significantly since the Mesozoic because of plate tectonics (i.e., rifting and then collision in the Eocene) and associated sediment production (Curry, 2014 and references therein). Despite onshore foreland-basin accommodation created by ongoing collisional tectonics,

sediments are moving through most of the system over historical timescales, from the Himalayas (>5000 m elevation) to the Bengal Fan (>4000 m water depth) (Fig. 4; Kuehl et al., 2005). Sediment production in the Ganges-Brahmaputra catchment (including the Meghna River) corresponds to an average catchment denudation rate of 365 mm/kyr, which is over an order of magnitude larger than the global average of 30 mm/kyr (Islam et al., 1999). The sediment load for the integrated catchment is ~1,000 Mt/yr, which equates to a system sediment yield of 556 t/km²/yr. However, sediment yield varies significantly spatially across the catchment. For example, the Brahmaputra River yield is >140% that of the Ganges (Summerfield and Hulton, 1994; Islam et al., 1999), and most of the Brahmaputra sediment is sourced from a smaller portion of the catchment, the High Himalayas (Wesson, 2003).

Gauging stations for rivers are located about 300 km from the coast, and studies indicate ~30% of the sediment is stored landward of the coastline (Fig. 4) (Goodbred and Kuehl, 1999). Sediment sinks include levee, floodplain, and river-bed aggradation, and alluvial accumulation in tectonically subsiding areas (Allison, 1998; Goodbred and Kuehl, 1998). Longer timescale records show that since the middle Holocene (~7 ka) slowdown in sea-level rise, some locations have accumulated >20 m of sediment (Goodbred and Kuehl, 1998), and rates of filling since ~12 ka suggest significant climate forcing on sediment supply (Goodbred and Kuehl, 2000a). Over historical timescales, floodplain areas of the upper delta plain have linear sediment accumulation rates that generally decrease with distance from the river channel (e.g., from 4 cm/yr to <1 cm/yr, Allison, 1998). Sediment dynamics in the lower delta plain are influenced by processes that originate in the marine realm such as sea-level rise, waves, tides, and cyclones

(Allison and Kepple, 2001; Hanebuth et al., 2013). Shoreline areas show a complex pattern of erosion and accretion (Allison, 1998; Shearman et al., 2013), but radiochemical analyses indicate sediment accumulation generally decreases with distance from the coast, reflecting import of fluvial sediment (Allison and Kepple, 2001).

Seismic-reflection profiling has established the presence of a sizable subaqueous delta clinoform on the shelf (Kuehl et al., 1997; Michels et al. 1998). Bathymetric and shoreline changes indicate that ~20% of the fluvial load is building the topset of the clinoform (Fig. 4) (Allison, 1998). Based on core and seismic-reflection data, the foreset region of the clinoform sequesters another 20-31% over historical timescales (Fig. 4) (Michels et al., 1998; Suckow et al., 2001). Transparent layers visible in seismic reflection profiles of the clinoform have been suggested to represent mass flows triggered by earthquakes (Michels et al., 1998). As a result of westward along-shelf currents reworking the delta front, sediment is at present being advected into the head of the Swatch of No Ground submarine canyon and episodically to the Bengal submarine fan (Kuehl et al., 1997; Kuehl et al., 2005). Weber et al. (1997) showed that late Holocene sedimentation occurred on the channel-levee complex (on the middle fan, ~500 km seaward of the shelf), but at a reduced rate compared to latest Pleistocene to early Holocene. Cyclones are hypothesized to be responsible for stratigraphic layering visible on the shelf and in the upper canyon (Fig. 4) (Kudrass et al., 1998; Suckow et al., 2001; Michels et al., 2003). Cyclones also serve as a possible trigger mechanism for episodic mass wasting events (Rogers and Goodbred, 2010). Canyon sedimentation and down-canyon transport, including evidence for turbidite deposition on the Bengal Fan, are hypothesized to account for ~30% of the fluvial load over historical timescales (Fig. 4)

(Goodbred and Kuehl, 1999; Kuehl et al., 2005). A terrestrial erosion-zone signal is being driven down this system, but it has been and continues to be significantly modulated by other processes (e.g., cyclones) along the way. As a result, alluvial storage areas might be the best sites for extracting source forcings over the historical timescale.

The Ganges-Brahmaputra-Bengal and Eel systems research highlight how historic stratigraphic records, accumulation rates, and sediment budgets can inform system functioning and source-to-sink transfer. This work also demonstrates that, although historical timescale records may be data rich and highly temporally resolved relative to intermediate and deep-time records, evaluation of sediment supply signals generated in upland catchments can be difficult. A more detailed and quantitative documentation of processes, rates, and spatial distribution of sedimentation does not necessarily equate to a better understanding of linkages between system segments. Better preserved and potentially more complete records in proximal storage areas, such as lakes, might allow more detailed records to be captured up system, but the localized nature might not reflect broader system functioning (e.g., Orpin et al., 2010). Combining observations from multiple localities will be essential to defining robust regional or broader, global signals (e.g., Noren et al., 2002). Other insights about catchment sediment production can be provided from the geomorphic record of erosional landforms. The sedimentary signature of events, such as floods, earthquakes, and storms, is likely more easily relatable to its forcing if process and response occur within the same or immediately adjacent segment(s) of the sediment-routing system (e.g., coastal overwash fan deposits from landfalling hurricanes; Boldt et al., 2010). The variability in sediment transport and associated deposits generated at $<10^2$ yr timescales is commonly considered noise over

longer timescales as a consequence of combining event-scale and ‘background’ sedimentation into a time-averaged rate. However, the findings from historical timescale studies show that there are signals embedded within the noise.

3 Sediment Routing at Intermediate (10^2 - 10^6 yr) Timescales

The timescale from just beyond historical (several centuries before present; discussed above) to several millions of years is a critical temporal range in Earth surface dynamics because fundamental climate forcings (i.e., Milankovitch cycles) that control the global climate are prominent over this timescale (Hays et al., 1976). Sustained rates of rock uplift and deformation in tectonically active areas lead to exhumation, sediment production, and morphological change at $\geq 10^5$ yr timescales (Burbank and Anderson, 2011). Moreover, it is in this temporal range during which sedimentary deposits can be sufficiently buried to become rock and preserved into the stratigraphic record – durations often referred to as ‘geological timescales’ (e.g., Allen et al., 2013).

We first discuss sedimentary system dynamics and associated signal implications based on numerical and physical models. Unlike the short-term timescales during which an integration of direct observation, monitoring, and modeling informs our understanding of source-to-sink signal propagation, modeling and theory become even more critical for intermediate (10^2 - 10^6 yr) timescales. Examples of recent work on paleo-sediment budgets for sediment-routing systems are also discussed.

3.1 *Model Predictions of Intermediate Timescale Signal Propagation*

We review how tectonic or climatic signals with periods of 10^2 – 10^6 yr are propagated through different portions of the sediment routing system (Fig. 2). We emphasize sediment supply (Q_s) as the principal vector for environmental signal propagation and aim to provide a review on the current state of knowledge with respect to the following important questions: (1) Does the erosion zone produce sediment supply signals in response to climate and tectonic perturbations with periods able to generate stratigraphic patterns? (2) Does the transfer zone faithfully transmit signals to the sedimentary basin, or does it modify signals coming from the erosion zone?

Signal transfer through a system depends on whether its period is smaller or larger than the response time of the system (Paola, 1992; see also Allen, 1974). Moreover, the action of internal, or autogenic, dynamics in any or all of the mass-flux zones can influence Q_s behavior, which affects signal propagation. We first define and review knowledge of response times for the erosion zone of hillslopes and bedrock channels, then focus on the transfer zone of mixed alluvial and bedrock channels to alluvial channels with floodplains, and its linkage to the accumulation zone in sedimentary basins (Fig. 1).

3.1.1 *Q_s Signal Generation and Propagation in the Erosion Zone*

It is beyond our scope to present a comprehensive review of investigations into how climate and/or tectonic forcings are recorded in net-erosional landscapes (e.g., Burbank and Anderson, 2011; and references therein). Rather, we emphasize the propagation of those perturbations out of the erosion zone in the form of sediment supply.

The concept of steady state as applied to landscape evolution (e.g., [Willett and Brandon, 2002](#)) refers to a state in which Earth's surface elevation relative to a datum is broadly constant as a result of a balance between rock uplift and erosion. Thus, the rate of sediment supply out of an area in steady state is, in the simplest case, also constant when averaged beyond timescales of individual events. A perturbation in the form of varying rates of tectonic movement or precipitation induces a response of the landscape system in the form of varying rates of sediment production. A characteristic equilibrium, or response, time for the system is the time that it takes for this transient landscape to respond to this perturbation and then return to a steady state ([Beaumont et al., 2000](#)) (Fig. 5). [Allen \(2008b\)](#) termed landscapes that have a response time shorter than the repeat time of the perturbation as 'reactive' and those with response times longer than perturbation repeat time as 'buffered' landscapes. This equilibrium time is critical to the discussion of signal propagation because it is the *variability* of Q_s out of the erosion zone that can result in recognizable variations in deposit character down system. Here, we focus on the relevance of steady state in terms of denudation because of its close association with sediment production.

The physical laboratory experiments of [Bonnet and Crave \(2003\)](#) highlighted the important observation that climate signals, because they can affect the totality of an area at once, can trigger an immediate response of the landscape. In their case, steady state is characterized by a constant mean elevation ([Montgomery, 2001; Willet and Brandon, 2002](#)) and, thus, a response is a change in mean elevation. This contrasts to rock uplift signals, expressed in the form of baselevel changes (see [Schumm, 1993](#)), which propagate as waves of headward incision and diachronously affect the landscape ([Bonnet](#)

and Crave, 2003). In a study of the response of bedrock channels to tectonic and climate signals using generic stream-power fluvial incision, Whipple (2001) showed response times ranging from 250 kyr to 2.5 Myr to both tectonics and climate. In this study, the response time is the time required for the landscape to return to a steady state defined as a statistically invariant topography (i.e., constant mean elevation; Montgomery, 2001; Willet and Brandon, 2002) and constant denudation rate. When climate and tectonics act jointly, the response of a stream-power fluvial landscape may essentially be immediate (i.e., response time tends to zero; Whipple, 2001). Improvements of the stream-power erosion law produce divergent results as to landscape reactivity. Among these, the consideration of dynamic adjustment of channel width during perturbations induces faster reaction of fluvial landscapes than if channel width is not considered (Attal et al., 2008; Whittaker et al., 2007). Conversely, a series of stream-power inspired models (e.g., Gasparini et al., 2007) including a degree of dependency to saltating bedload tends to suggest longer response times than those predicted by detachment-limited stream power (such as those of Whipple, 2001; see above).

Using a nonlinear 1D diffusive model of catchment erosion, Armitage et al. (2013) showed that small (10-20 km long) catchments, such as those draining normal-fault bounded footwalls, are reactive to single-step, sustained changes of precipitation but tend to temporally buffer cyclic precipitation variations with Milankovitch periodicities (i.e., 100 kyr, 400 kyr, 1.2 Myr). Using a 2D model of landscape evolution including diffusive hillslopes and detachment-limited stream-power-governed bedrock incision, Godard et al. (2013) found that a given landscape possesses a characteristic resonance periodicity for which landscape response to corresponding climatic oscillation is maximized in terms of

sediment supply. For more easily erodible lithologies, landscape response to orbitally controlled climate signals could be a significant increase or decrease of the amplitude of sediment supply variations. Thus, some landscapes respond to, might even amplify, climate and tectonic signals. In landscapes dominated by diffusive hillslopes, however, such as in soil-mantled, low-relief settings, diffusion itself might be very efficient at filtering climatic or tectonic oscillations because of slow signal propagation (e.g., Furbish and Fagherazzi, 2001).

3.1.2 Qs Signal Generation and Propagation in the Transfer Zone and Preservation in the Accumulation Zone

In many instances, the terminal depositional sink is not immediately adjacent to the source area but linked to it by a fluvial system. In such cases, it was recognized that the fundamental problem becomes whether climate and tectonic sediment supply signals that originate in the erosion zone are propagated by the transfer system to the sedimentary basin (Castelltort and Van Den Driessche, 2003). Paola et al. (1992) developed the idea that to understand stratigraphic response to external factors it was fundamental to consider the periodicity of cyclic signals with respect to the characteristic equilibrium, or response, time (T_{eq}) of a sediment-routing system (Fig. 5). They expressed T_{eq} (time unit) for a 1D fluvial profile as a function of characteristic system length (L) and diffusivity (K):

$$T_{eq} \sim L^2/K$$

Thus, the larger the system (i.e., the longer the transfer zone), the longer its response time is whereas the more diffusive the transfer zone, the shorter its response time. A prediction of this model is that cyclic perturbations with periods less than T_{eq} are buffered by the system's response time. In contrast, variations of boundary conditions with periodicities greater than T_{eq} produce stratigraphic patterns in the sedimentary basin, but these patterns might be similar for subsidence and sediment supply variation (Paola et al., 1992; see also Marr et al., 2000 and Allen, 2008b).

On the basis of a comparison between the modern river sediment discharge of some large Asian rivers and the average sediment discharge deduced from sedimentary basins over the last 2 million years, Métiévier and Gaudemer (1999) suggested that large alluvial systems of Asia behave as diffusive entities buffering the high-frequency climate change known for the late Cenozoic (see also Schaller et al., 2001; and Wittmann et al., 2011). Métiévier and Gaudemer (1999) computed equilibrium times of >1 Myr for such rivers using an expression they proposed for the diffusivity (K) of large rivers as a function of sediment discharge (Q_s), river channel or channel-and-floodplain width (W), and slope (S):

$$K=Q_s/(W*S)$$

Following Métiévier and Gaudemer (1999) results, Castelltort and Van Den Driessche (2003) calculated the diffusive response time of 93 of the largest modern rivers to investigate the down-system stratigraphic response to high-frequency (10^4 yr) cycles of sediment supply. Castelltort and Van Den Driessche (2003) find that the characteristic

response times of transfer zones comprising large rivers, which typically include extensive floodplains, are 10^5 - 10^6 yr, exceeding the 10^4 yr climate oscillations. When channel width rather than alluvial valley width is used in this relationship the resulting response times are minimum response times.

These diffusion-based investigations suggest that temporary, and in some cases permanent, storage of sediment in catchment and/or transfer-zone sinks (see also [Allen, 2008a](#); [Wittmann et al., 2011](#); [Covault et al., 2013](#); and references therein) can mask the down-system stratigraphic record of external perturbations to the sediment-routing system. In the case of a large, hinterland-river-continental margin sediment-routing system, this transient storage of sediment can result from deposition in floodplains ([Allen, 2008b](#)). Larger catchments can retain sediment for longer periods as a result of more space available for sediment storage and consequent resistance to complete hinterland-to-continental margin sediment transfer in response to short-term, small-magnitude external perturbations, such as local storms and earthquakes ([Allen, 2008a](#)). [Métivier and Gaudemer \(1999\)](#) suggested that rivers and floodplains proportionally adjust to climate changes and upstream denudation in buffered catchments in which sediment loads are approximately balanced over different timescales. That is, if upstream denudation is reduced, the river will incise its floodplain to keep the sediment load at the outlet constant. Conversely, if climate changes force greater upstream denudation, the river is likely to use that increased sediment load to recharge its previously excavated floodplain. In this way, the steady transfer of reworked floodplain sediment to an outlet can be maintained over a range of timescales ([Métivier and Gaudemer, 1999](#); [Phillips, 2003](#); [Phillips and Slattery, 2006](#); [Covault et al., 2013](#); among many others). The

ubiquitous alluvial terrace fills that ornate many river systems worldwide are witnesses of the residence time of sediments in the transfer zone.

These theoretical results contrast with the sensitivity to Late Quaternary climate change apparently displayed by some large fluvial systems such as the Ganges-Brahmaputra (Goodbred and Kuehl, 1999; 2000a, 2000b; Goodbred, 2003) and suggest that, although alluvial systems may behave diffusively in response to sediment supply variations, they may be sensitive to perturbations of water discharge, which can increase or decrease diffusivity (Simpson and Castelltort, 2012). Using physical laboratory models of river response to water discharge and sediment supply change, Van Den Berg Van Saparoea and Postma (2008) show that experimental rivers respond faster to changes of discharge than to perturbations of up-system sediment supply. Van Den Berg Van Saparoea and Postma (2008) concluded that high-frequency cyclic patterns in marine delta-shelf successions were most likely controlled by high-frequency changes in discharge driven by climate, whereas the low-frequency sequences were likely a result of low-frequency changes in sediment supply driven by tectonic deformation.

We recognize that the results of diffusion-based approaches may be dependent on our current ability to estimate parameters of the diffusion laws. Simpson and Castelltort (2012) explored the response of a 1D alluvial river bed to sediment concentration and water discharge pulses using a physically based numerical model of interacting water flow and sediment transport without *a priori* assumption of diffusive behavior. Consistent with the experiments of Van Den Berg Van Saparoea and Postma (2008), in this model the strong coupling between water discharge and river gradient induces amplified sediment supply variations in response to oscillations of water discharge, whereas

sediment supply oscillations are dampened because of the negative feedback between sediment concentration and channel gradient. In the future, additional constraints on the behavior of sediment transfer will result from other approaches such as computational fluid dynamics (e.g., [Edmonds and Slingerland, 2007](#)) or cellular automata (e.g., [Murray and Paola, 1997](#)).

3.1.3 Potential Influence of Internal Dynamics on Q_s Signal Recognition

In addition to the buffering of signals linked with the processes reviewed in the previous sub-section, perturbations to sedimentary signal propagation arise from sedimentary processes occurring within the river-floodplain and/or river-coastal plain segments that need not be driven by up-system forcings. Such self-organizing processes, referred to as autogenic dynamics ([Paola et al., 2009](#)), and first emphasized by [Beerbower \(1964\)](#), can create organized depositional architecture (e.g., [Hoyal and Sheets, 2009](#)). Critical to this discussion is the potential for climate or tectonic signals that originated in the catchment to be significantly masked, modified, or ‘shredded’ by such autogenic dynamics ([Jerolmack and Paola, 2010](#)). Variability in sediment transport is a result of the following general autogenic cycle: transient storage of sediment, exceedance of some critical threshold, and release of sediment during relaxation following failure. [Jerolmack and Paola \(2010\)](#) likened the threshold behavior of sediment storage and release to morphodynamic turbulence, analogous to turbulence in fluid flows.

Recently, [Ganti et al. \(2014\)](#) developed a quantitative framework to isolate autogenic, morphodynamic processes from external, environmental forcings in the

stratigraphic record. They showed that the calculated advection length (l_a) for settling sediment sets bounds on the scale over which autogenic processes operate:

$$l_a = u h_s / w_s$$

where u is the flow velocity, h_s is the average sediment settling height, and w_s is the settling velocity. The advection length scale is the horizontal length over which an average particle is transported in the flow before falling to the bed. [Ganti et al. \(2014\)](#) argued that morphodynamic feedbacks, or autogenic ‘shredding,’ can only occur if the length scale of interest, e.g., the system size, is larger than l_a .

[Wang et al. \(2011\)](#) recognized the aforementioned work on damping or ‘shredding’ of upstream, external signals by autogenic sediment transport processes, and used numerical and physical experiments, as well as some field data, to gain insight into the timescale of compensational stacking of deposits within a basin. This compensation timescale (T_c) is defined as:

$$T_c = l/r$$

where l is a roughness length scale, equal to the amount of topographic ‘mounding’ due to local channel deposition produced between each avulsion, and r is the basin-wide, long-term sediment accumulation rate. This equation suggests that the geometry of deposits carries the signature of stochastic autogenic dynamics during the time necessary to fill a basin to a depth equal to the amount of surface roughness in a sediment-routing

system. T_c provides an estimate of temporal scales below which stratigraphers should be cautious about interpreting signals. As a case in point, [Wang et al. \(2011\)](#) calculated T_c for the Lower Mississippi Delta in which they consider that the roughness length scale, l , was represented by the mean channel depth for the Lower Mississippi River of 30 m and a sediment accumulation rate of 0.26 m/kyr, estimated for the past 8 Myr ([Straub et al., 2009](#)). T_c is 115 kyr, which is ~100 times larger than the ~1300 yr recurrence of large avulsions of the Lower Mississippi River ([Aslan et al., 2005](#)). However, subsequent field data from the Lower Mississippi River indicate a rapid response to glacio-eustatic variation since Oxygen Isotope Stage 7 (~200 ka) ([Shen et al., 2012](#)). Large amplitude sea-level rise and fall prompted rapid and widespread fluvial aggradation and incision, respectively, the effects of which extended >600 km upstream from the present shoreline ([Shen et al., 2012](#)).

The models and experiments discussed above highlight that signal buffering as a result of sediment storage in up-system segments as well as depositional dynamics in the sink can mask the stratigraphic record of external perturbations to the sediment-routing system, although the quantitative expression of this are still being resolved. In summary, signals of a forcing can be passed to a basin and preserved in the stratigraphic record when their period exceeds the characteristic equilibrium time of the sediment-routing system, but this is valid only if their period is also larger than the characteristic timescale of autogenic sediment transport fluctuation and/or when the magnitude of the forcing is larger than the magnitude of internal oscillations (e.g., on the order of the size of catchment and alluvial accommodation) ([Jerolmack and Paola, 2010](#)). In the next section

we review sediment budgets of natural sedimentary systems, which allow for accounting of our principal vector of relevance, Q_s .

3.2 Paleo-Sediment Budgets of Natural Systems and Implications for Signal Propagation

In this section we will review work on sediment budgets at 10^2 - 10^5 yr timescales and implications for signal propagation via three sediment-routing systems: (1) tectonically active, small systems of southern California; (2) tectonically quiescent, larger systems of the northwestern Gulf of Mexico; and (3) tectonically active, larger systems of southern Asia. By focusing on sediment delivery from onshore catchments to the deep sea, which is the ultimate sink for coarse-grained terrigenous material, we highlight the role of the shelf as a Q_s gateway and filter.

3.2.1 Methods for Paleo-Sediment Budget Reconstruction at Intermediate Timescales

Just as microfossils are the carriers of isotopes used to reconstruct geochemical signals, sediment supply is here considered the carrier of climate and tectonic signals. Thus, determining a paleo-sediment budget, the spatial and temporal partitioning of mass removed, transferred, and deposited within a routing system, is valuable for the interpretation of signal propagation and preservation. For the sake of brevity, we do not present a comprehensive review of the application of sediment budget concepts to timescales beyond direct measurement and instead refer the reader to a recent review by [Hinderer \(2012\)](#). Determining accumulated mass from stratigraphic volumes is straightforward in concept, but can be challenging in practice as a result of lack of

appropriate data (e.g., seismic-reflection with proper chronologic control) and/or uncertainties in post-depositional stratal preservation ([Sadler and Jerolmack, 2014](#)). The geochemistry and mineralogy of sediment is often used to determine routing pathways as well as the relative contributions and potential residence times of terrigenous versus marine-derived material.

Two of the three systems reviewed below combine cosmogenic radionuclide (CRN) analysis for catchment-integrated denudation and radiocarbon dating for continental-margin deposition to reconstruct sediment budgets. Advances in CRN analysis provide catchment-integrated denudation rates and sediment loads at 10^2 – 10^5 yr timescales ([von Blanckenburg, 2005](#)), which are comparably similar to the timescales of deposition measured in offshore basins with radiocarbon ages (generally <50 ka; [Reimer, 2012](#)). CRNs are produced *in situ* as secondary cosmic rays interact with rocks within meters of Earth's surface; longer exposure to secondary cosmic rays as a result of slower denudation produces more nuclides. Sediment can be liberated from these rocks, mixed in the catchment through hillslope and fluvial transport processes, and ultimately deposited near the catchment outlet. Accordingly, the CRN abundance measured in sediment deposited near the catchment outlet can be used to divulge the catchment-wide denudation rate, which is inversely proportional to nuclide abundance ([Brown et al. 1995](#); [Bierman and Steig 1996](#); [Granger et al. 1996](#)).

Regardless of the specific tools used, it is of critical importance that all mass inputs and outputs to the system are considered and accounted for. Attempting to close a sediment budget at timescales beyond direct measurement provides an opportunity to

757 evaluate other inputs and outputs that might not be evident with a qualitative
758 interpretation.

760 **3.2.2 *Small and Tectonically Active Systems of Southern California***

761 Tectonically active southern California is an ideal setting in which to investigate
762 millennial-scale mass balance as a result of close proximity of sediment-routing
763 components: onshore erosion zones are located adjacent to short alluvial-coastal plain
764 depositional environments and offshore, confined sedimentary basins of the California
765 Continental Borderland (Fig. 6A). The confinement of the offshore basins facilitates
766 complete accounting for detrital mass relative to open-ocean basins, such as the Arabian
767 Gulf and Bay of Bengal (Weber et al., 1997; Curray et al., 2002). Furthermore, many of
768 the submarine canyon and fan systems of the California Continental Borderland are
769 consistently linked to the shoreline and maintain connectivity even during Holocene
770 highstand (Normark et al., 2009).

771 Covault et al. (2011) used CRNs from the Peninsular Ranges of southern California to
772 calculate catchment-integrated denudation rates, which varied from 0.07 to 0.24 mm/yr
773 since 10 ka. These denudation rates were calculated to be 1.9-2.4 Mt/yr and integrated
774 across the total area of drainage basins ($>6 \times 10^3 \text{ km}^2$) delivering sediment to the offshore
775 Oceanside littoral cell and the La Jolla submarine canyon and fan system. Based on
776 radiocarbon-constrained seismic-reflection mapping (Covault et al., 2007) the mass
777 accumulation rate of the La Jolla submarine fan was calculated to be 2.6-3.5 Mt/yr since
778 the Last Glacial Maximum. Although the mass of material denuded from Peninsular
779 Ranges catchments is in close agreement, of the same order of magnitude, as the mass of

material deposited in the La Jolla submarine fan, deep-sea deposition exceeds terrestrial denudation by 11%-89%. This additional supply of sediment could be owed to enhanced dispersal of sediment across the shelf caused by sea cliff erosion during postglacial shoreline transgression and initiation of submarine mass wasting.

The terrestrial source to deep-sea sink mass balance does not show orders of magnitude inequalities that might be expected in the wake of major sea-level changes since the Last Glacial Maximum. Thus, sediment-routing processes in a globally significant class of small, tectonically active systems might be fundamentally different from those of larger systems that drain entire orogens, in which sediment storage in coastal plains and wide continental shelves can exceed millions of years ([Milliman and Syvitski, 1992](#)). Furthermore, in such small systems, depositional changes in the deep offshore can reflect onshore changes when viewed over timescales of several thousands of years to more than 10 kyr. For example, [Romans et al. \(2009\)](#) and [Covault et al. \(2010\)](#) examined Holocene deposition of the Hueneme and Newport deep-sea depositional systems offshore of southern California. Integrated datasets of radiocarbon ages from sediment cores and seismic-reflection profiles demonstrated that variability in rates of Holocene deep-sea turbidite deposition is related to complex ocean-atmosphere interactions, including enhanced magnitude and frequency of El Niño-Southern Oscillation (ENSO) cycles, which increased precipitation and fluvial water and sediment discharge in southern California (Fig. 7). Thus, millennial-scale climate forcings are represented as a measureable signal in the stratigraphic record of the deep-sea segment.

3.2.3 *Large and Tectonically Quiescent Systems of the Western Gulf of Mexico*

A sediment budget for the Brazos and Trinity rivers linked to offshore depositional systems in shallow-marine and deep-water environments of the northwestern Gulf of Mexico can be balanced by integrating recent work of [Hidy et al. \(2014\)](#) and [Pirmez et al. \(2012\)](#). In contrast to the small and tectonically active southern California catchments, the Brazos and Trinity rivers drain a large ($\sim 2 \times 10^5 \text{ km}^2$) tectonically quiescent, non-glaciated, low-relief landscape (Fig. 6B). [Hidy et al. \(2014\)](#) evaluated how denudation rates from CRNs responded to climate change during the last glacial cycle ($\sim 15\text{-}45 \text{ ka}$): Brazos River CRNs yielded a mass load of 5.3 Mt/yr since 35 ka; and Trinity River CRNs yielded a mass load of 2-4 Mt/yr. Furthermore, [Hidy et al. \(2014\)](#) analyzed the CRN ratio of $^{26}\text{Al}/^{10}\text{Be}$ in river sediment to evaluate its transient storage in the catchment in route to its final depositional site (see [Wittmann and von Blanckenburg, 2009](#); [Wittmann et al., 2011](#)). Mass storage on the coastal plain was interpreted to have been greater during glacial periods with lower sea level. Denudation rates and mass loads were calculated to be larger during interglacial periods, which suggest that increased weathering rates associated with warmer climates accelerated landscape erosion. Furthermore, increased mass load measured during warm interglacial periods is interpreted to reflect stronger reworking and delivery of sediment to the river mouth than during cooler glacial periods. An implication of this relationship between temperature and mass load is that global sediment and potentially dissolved load delivery to the ocean from analogous, tectonically quiescent, non-glaciated, low-relief landscapes might have been larger during the warm Pliocene than the cooler Quaternary ([Hidy et al., 2014](#)). However, any transient storage of sediment prior to preservation in terrace deposits

would complicate the interpretation of the CRN data as representative of catchment denudation.

[Pirmez et al. \(2012\)](#) developed a robust chronostratigraphic framework from Oxygen Isotope Stage 6, ~120 ka, through the Last Glacial Maximum for the sediment deposited in four deep-water, salt-withdrawal slope basins of the northwestern Gulf of Mexico (see also [Prather et al., 2012](#)) (Fig. 6B). The deep-water depositional systems were linked to the Brazos and Trinity river-deltas only during lowstands of sea level, when the shoreline had regressed >100 km from the present-day beach to the shelf edge ([Mallarino et al., 2006](#); [Anderson et al., this volume](#)). Chronostratigraphy was interpreted from an integrated database of 3D seismic-reflection data, age control from analysis of cores from Integrated Ocean Drilling Program Expedition 308, analysis of proprietary cores from Shell Oil Company, and published literature ([Pirmez et al., 2012](#)). The majority of sediment, ~50 km³, was calculated to have been deposited during a period of relatively low sea level, between ~15-24 ka, yielding a mass accumulation rate during this period of 5.5 Mt/yr, which is within the same order of magnitude of the CRN-derived mass load of the Brazos and Trinity rivers of 7.3-9.3 Mt/yr ([Hidy et al., 2014](#)) during a similar period, generally <35 ka. The imbalance in rates, with diminished deep-water slope basin mass accumulation, is likely a result of Brazos-Trinity river-delta deposition on the subaerially exposed shelf as the shoreline had regressed to the shelf edge between ~15-24 ka. Indeed, [Pirmez et al. \(2012\)](#) estimate a maximum volume of ~25 km³ of deltaic sediment was deposited between ~15-24 ka, which yields a mass accumulation rate of 2.8 Mt/yr. This mass added to the deep-water basin fill yields a total mass accumulation rate of 8.3 Mt/yr,

which is within the range of CRN-derived mass load ([Hidy et al., 2014](#)) of the Brazos and Trinity rivers.

In this system, sea level is interpreted to control the delivery of terrigenous sediment to the deep-water slope basins over 10^5 yr timescales: during periods of relatively high sea level, when the shoreline had transgressed, the slope basins were interpreted to receive only hemipelagic, fine-grained sediment. During periods of relatively low sea level and a subaerially exposed shelf, relatively coarse-grained terrigenous sediment was deposited in the slope basins. However, mass accumulation during periods of relatively low sea level was interpreted to have varied between the four slope basins as a result of a complex interaction between river-delta sediment routing and dynamic, salt-withdrawal slope-basin evolution. The complex history of sediment deposition, storage, and remobilization in the zone between the modern coastline and the shelf edge over the past ~125 kyr ([Anderson et al., this volume](#)) highlights the role of the shelf as an additional filter of signals generated in the catchment. Therefore, in contrast to the southern California examples, offshore depositional records are hypothesized to primarily reflect sea-level-driven accommodation changes as opposed to Q_s variability from the onshore catchments. However, during glacial periods of terrigenous sediment delivery to deep water, depositional records might reflect Q_s variability from rivers (Fig. 6B). This might be common to other sediment-routing systems, where deep-water canyon heads are stranded at the edges of drowned continental shelves during interglacial periods ([Blum and Hattier-Womack, 2009](#); [Covault and Graham, 2010](#)).

3.2.4 Large and Tectonically Active Systems of Southern Asia

Since the Last Glacial Maximum (LGM), the Indus sediment-routing system comprised a steep (total relief of nearly 8 km), tectonically active hinterland and large (~10⁶ km²) catchment (Milliman and Farnsworth, 2011), a delta located on a wide (~120 km) shelf, and a submarine canyon that fed the second largest accumulation of terrigenous sediment in the world, the Indus submarine fan (Fig. 6C). Clift et al. (2014) used seismic-reflection data, radiocarbon ages, and analyzed the geochemistry (a suite of major and trace elements, including Zr/Rb, K/Rb, and Nd; Limmer et al., 2012) and mineralogy of sediment of the Pakistani continental margin to investigate sediment routing from the Indus river-delta to the upper submarine canyon (<1300 m below present sea level) since the LGM. Seismic stratigraphic interpretations of deltaic clinoforms and radiocarbon ages indicate that the majority of Holocene Indus river-delta sediment is stored on the shelf (Giosan et al., 2006; Clift et al., 2014). Clift et al. (2014) interpreted a variety of deltaic clinoform seismic reflections and concluded that sediment used to construct the shelf-edge delta deposits was reworked and dispersed from mid-shelf locations basinward. Neodymium isotopes presented by Limmer et al. (2012) suggests transient storage on the shelf was significant prior to delivery to the canyon and fan system. Neodymium isotope ratios indicate different values compared to those expected from a fluvial source, which points to reworking of marine sediment deposited during the LGM (Clift et al., 2014). Deposition at the head of the Indus Canyon was measured to be rapid during the Holocene, with evidence for annual delivery of Indus river-delta sediment. However, downstream, ~1300 m below present sea level, the

youngest deposits are greater than approximately 7 ka, and no terrigenous sand has reached the upper submarine fan during the Holocene (Clift et al., 2014).

The Indus sedimentary record at the Pakistani continental margin since the LGM indicates reworking and transient storage of sediment on the shelf and within the submarine canyon en route to the deep Arabian Sea. Thus, deep-water deposits of the Indus fan likely do not faithfully record Qs variability related to climatic or tectonic events onshore over timescales of 10^3 - 10^4 years (Clift et al., 2014). This conclusion highlights the role of the shelf segment as a critical process boundary as well as a Qs gateway between land and sea. This is similar to the Brazos-Trinity sediment-routing system (Fig. 6C), where the deep-water depositional record primarily reflects sea-level changes as opposed to Qs variability from the onshore catchments. Moreover, sediment storage in the vast Indus floodplain (Fig. 6C) likely buffers climatic or tectonic signals generated in the up-system headwaters of the Himalayas. Thus, in these large sediment-routing systems (see also the Amazon system, e.g., Wittman et al., 2011), millennial to million-year Qs signals that originated in the erosion zone are potentially recorded in alluvial and floodplain deposits.

3.3 *Synthesis of Intermediate Timescale Signal Propagation*

The theory, models, and natural systems reviewed above inform us about the plausibility of different forcings and how they might generate Qs signals that are then transmitted by the transfer zone to the ultimate depositional sink. These concepts are synthesized in Figure 8, which contains schematic representations of many of the

scenarios reviewed in preceding sections, which can also be viewed as hypotheses about sediment-routing system dynamics worthy of further investigation.

Tectonic signals (e.g., uplift rates) with periods of >50 kyr are likely to produce Qs signals at the outlet of the erosion zone (Fig. 8A). Such signals may be transmitted to the accumulation zone if the transfer zone is short (<300 km) but will be buffered if the transfer zone is long (>300 km), unless their period exceeds 100 kyr. Tectonic signals with periods of <50 kyr are likely to be already buffered by the dynamics of the erosion zone itself; i.e., before they reach the outlet of the erosion zone (Fig. 8B).

Several antithetic results exist with respect to climate signals (e.g., precipitation changes) in the erosion zone. Different models propose that climate signals might be buffered, faithfully transmitted, or even amplified by the erosion zone (Fig. 8C-E). In some natural examples, where sediment budgets have determined the magnitude of sediment supply exiting the erosion zone, there is a relationship between climate (e.g., variation in precipitation) and Qs signals as recorded in the sink (Covault et al., 2009; Romans et al., 2009; Fig. 7). However, it is challenging to test whether climate signals are amplified since no predictive understanding exists between a given climate change and the corresponding amplitude of catchment response in terms of sediment supply. However, as discussed above, numerical models of sediment transport and deposition offer the opportunity to elucidate the question of climate signal amplification (e.g., Armitage et al., 2011). Nonetheless, intermediate climate signals in the form of water discharge (Qw) seem to trigger a strong response of the transfer zone in terms of sediment supply and may thus be transmitted faithfully or even amplified to sedimentary basins (Fig. 8C-E) (Godard et al., 2013; Simpson and Castelltort, 2012).

At the moment, few constraints exist on the characteristic saturation timescales and amplitudes of internally generated Qs fluctuations in natural systems. Current estimates of the typical timescale for channel stacking in large river systems (e.g., Hajek et al., 2010) suggest that autogenic dynamics may be able to completely mask or even destroy sedimentary signals with periods of less than 100 kyr.

4 Deep-Time ($\geq 10^7$ yr) Sediment Routing

4.1 Challenges and Uncertainties in Deep-Time Signal Propagation Analysis

As sedimentary systems age to tens of millions of years and older, the ability to explicitly measure or calculate source-to-sink sediment supply becomes increasingly challenging because: (1) sediment production areas are poorly preserved or not preserved at all, (2) there is increased uncertainty regarding boundary conditions such as tectonic setting and climate regime, (3) of the diminishing resolution of existing chronological tools, and (4) of the completeness of the stratigraphic record (Romans and Graham, 2013).

Reading the sedimentary record in deep time also requires understanding forcings with long periods. The maximum response times resolvable for erosional and depositional processes in most sedimentary systems is commonly $\sim 10^6$ yr (e.g., Paola et al., 1992; Whipple, 2001; Castelltort and Van Den Driessche, 2003; Allen, 2008b). Thus, tectonic and climatic signals with periods of at least several millions of years could induce a measureable equilibrium response of the Earth's surface that is potentially recorded in stratigraphic successions. In other words, long-period stratigraphic archives

959 have more immunity to the internally generated dynamics that plague intermediate
960 timescales. Examples of long-period forcings include the development of orogens and
961 their coupled sedimentary basins (Dickinson, 1974; DeCelles et al., 2009), Phanerozoic
962 changes of Earth's sea level (e.g., Miller et al., 2005), and significant shifts in global
963 climate such as the transition from Cretaceous-Eocene greenhouse to Oligocene-present
964 icehouse conditions (e.g., Zachos et al., 2008).

965 When peering back even further in time ($\geq 10^8$ yr) we lose details about the
966 fundamental boundary conditions of tectonic setting and environmental conditions that
967 are explicitly known for the modern or reliably reconstructed for historical to
968 intermediate timescales. Thus, in many cases, reconstructing those boundary conditions is
969 the primary goal of sedimentary analysis. Linking strata of such old ages to sediment
970 source areas is challenging as a result of major tectonic regime changes (e.g., closing and
971 opening of ocean basins; Wilson, 1966), poorly understood oceanic and atmospheric
972 conditions, and non-actualistic Earth processes.

973 Our ability to reconstruct deep-time Earth surface conditions is based on rock
974 availability: preserved as intact depositional architecture and/or detrital material
975 representative of long-gone source areas. The following sections briefly review
976 methodologies for characterizing the unpreserved sediment-production zones of ancient
977 systems and potential value for interpreting Qs signal propagation.

4.2 *Inferring Catchment Characteristics and Sediment Supply From Stratigraphic Architecture*

Erosion and transfer zones are inherently not preserved in deep time and thus we must rely on preserved stratigraphy to reconstruct their characteristics. In this section we briefly review methods for estimating catchment area from stratigraphic architecture.

The dimensions of modern river channels scale to flood, or bankfull, discharge (Bridge and Demicco, 2008; and references therein), which affords estimation of water discharge from preserved fluvial channel stratigraphic architecture (e.g., Bridge and Tye, 2000; Bhattacharya and Tye, 2004; Adams and Bhattacharya, 2005; Blum et al., 2013; Bhattacharya et al., this volume). Analyses of modern river systems demonstrate a relationship between water discharge and catchment area (e.g., Hack, 1957; Rodier and Roche, 1984; Matthai, 1990) and a global empirical model shows that catchment area and relief are first-order controls on sediment supply (BQART model of Syvitski and Milliman, 2007). These relationships suggest that an estimate of paleo-catchment area can be determined from stratigraphy with the proviso that the channelized strata measured are truly representative of the alluvial system (see Blum et al., 2013 and Bhattacharya et al., this volume for discussion of river channel and alluvial valley scaling with respect to sediment delivery dynamics). However, discharge-to-area relationships as well as sediment load-to-area relationships from modern rivers are shown to vary as a function of precipitation and runoff patterns, vegetation, soil type, and geology (e.g., Milliman and Farnsworth, 2011; Covault et al., 2013 and references therein). Regional hydraulic curves, which capture such characteristics from modern systems (e.g., Leopold and Maddock, 1953), can be used to further constrain the estimate of catchment area if

some aspects of the paleoclimate can be determined. [Davidson and North \(2009\)](#) provide a comprehensive discussion of the values and limitations of the regional hydraulic curve approach, including example applications from the deep-time rock record.

Sediment yield can be approximated with the regional hydraulic curve approach, providing insight about the paleo-sedimentary system. In practice, however, the calculation of any mass supply is only as accurate as the chronologic control available; mass supply averaged over $>10^6$ yr will obviously not capture shorter-period fluctuations. Furthermore, such paleo-hydrologic methods are burdened with uncertainties that are challenging to quantify, which limits accuracy to an order of magnitude ([Holbrook and Wanas, 2014](#)). These methods are also susceptible to aliasing the record of Qs. For example, a single, static paleogeographic reconstruction might be used to inform paleo-hydrology over a large duration of geologic time, which provides average Qs during that time. However, geologic evolution, and especially Qs, is dynamic and influenced by the extreme events. Some applications can be satisfactorily addressed with order-of-magnitude estimates, such as the selection of modern analogs for ancient systems ([Bhattacharya and Tye, 2004](#); [Bhattacharya et al., this volume](#)). How to better link paleo-catchment reconstructions with interpretations of signal propagation remains a challenge. Ultimately, because these methods incorporate information from modern systems, the reliance on an actualistic approach should be acknowledged.

4.3 Source Area Signals From Detrital Material Analysis

Another approach to reconstruct aspects of the erosion zone of deep-time sediment-routing systems is to characterize the detrital material that is preserved in

sedimentary rocks. Provenance analyses focused on detrital products released from the erosion zone has long been used to reconstruct and interpret deep-time paleo-drainage systems and their relationship to tectonic forcings (Dickinson, 1974; Graham et al., 1986; McLennan et al., 1983; among many others). More recently, radioisotope provenance studies have been employed to detect sediment supply signals in deep time by identifying source terranes (Fig. 9) and tracking their evolution through a basin fill (e.g., Dickinson and Gehrels, 2003; Weislogel et al., 2006; Romans et al., 2010; Carrapa, 2010; Blum and Pecha, 2014 and references therein). The common pre-conditions for application of such methods are related to specific characteristic of the source areas, which should be composed of rocks with different tectonic histories, distinctive crystallization and cooling ages, and the presence of the unique mineral(s) (Lawton, 2014).

Combining different thermochronometers on single detrital mineral grains such as zircon, monazite, white mica, and apatite can be used to determine cooling ages associated with different closure temperatures (depths) with which to reconstruct tectono-thermal events (Rahl et al., 2003; Carrapa et al., 2003; Carrapa, 2010; Lawton, 2014). This has potential application for the interpretation of the propagation of a tectonic signal across a paleo-landscape by constraining separate events (e.g., crystallization and cooling) of a single grain such that durations and rates can be determined. These methods provide rates for mountain belt emplacement and exhumation, helping to refine timing and magnitude of tectonic processes.

For example, when zircon U-Pb crystallization ages are coupled with zircon (U-Th)/He exhumation ages, apatite fission-track (AFT), and/or apatite (U-Th)/He methods, we gain critical insight into the timing of rock cooling, inferred exhumation, lag times

between different closure temperature depths, as well as lag times between cooling and deposition (Fig. 10) (e.g., [Rahl et al., 2003](#); [Reiners and Brandon, 2006](#); [Painter et al., 2014](#)). Lag time was originally defined as the difference between the cooling age of a detrital mineral and the depositional age of its host strata ([Brandon and Vance, 1992](#); [Garver et al., 1999](#)). According to theoretical modeling, the variability in cooling age-depositional age lag times through a stratigraphic succession could be used to infer changing erosion rates in an orogenic belt ([Rahl et al., 2007](#)), which is a potentially powerful tool to estimate Q_s in deep time (Fig. 10). Most of these studies focus on synorogenic basins with inferred short transfer zones and correspondingly short duration of transient storage in order to interpret orogenic signals. Minerals yielded by distinct source regions could display overlapping crystallization ages related to different volcanic events but virtually undistinguishable if using a U-Pb dating technique (e.g., [Saylor et al., 2012](#)). Coupling these ages with (U-Th)/He ages can help distinguish older exhumed regions from younger, especially within the context of the omnipresent Grenville U-Pb age populations of the Appalachian orogenic belt ([Rahl et al., 2003](#)).

Tectonic and/or sedimentary burial can reset thermochronometers, complicating a simple exhumation to erosion relationship. In some cases, however, orogenic recycling can be constrained by integrating double dating with time-temperature modeling of burial history ([Fosdick et al., 2014](#)). These techniques will continue to be used to reconstruct paleo-drainage system connectivity and evolution, which will help track changes in sediment supply over $\geq 10^7$ yr timescales ([Carrapa et al., 2003](#)).

4.4 *Sedimentary System Mass Balance in Deep Time*

As reviewed in preceding sections, a full accounting of mass supply (and storage) among erosion, transfer, and accumulation zones of a sediment-routing system can aid interpretation of signal propagation. For example, transient storage of sediment in floodplains and coastal plain segments at intermediate timescales can buffer the transmission of Milankovitch-controlled supply entering the sink (Fig. 6C). These same concepts can be applied to deep-time systems, but with the significant challenges that erosion/transfer zones are typically not morphologically preserved and chronology is more poorly resolved.

Despite these uncertainties, theory and methodologies have been and are being developed with the goal of characterizing mass-balance in ancient systems. Paola and Martin (2012) built on previous work of Paola and Voller (2005) and Strong et al. (2005) to apply mass-balance concepts to quantitative characterization of sedimentary basin fills. This and similar studies (e.g., Whittaker et al., 2011; Carvajal and Steel, 2012; Petter et al. 2013) aim to improve the estimation of Q_s from time-averaged stratigraphy. Sadler and Jerolmack (2014) point out that well-documented 1D measurement-interval effects on rates of denudation (e.g., Gardner et al., 1987) and accumulation (e.g., Sadler, 1981) are eliminated with full spatial averaging because sediment generation and deposition must balance. Closing the sediment budget for an ancient system by accounting for all inputs and outputs is challenging (Hinderer, 2012; Allen et al., 2013). However, Sadler and Jerolmack (2014) make the case that avoiding linear rate measurements (i.e., maximizing volumetric analysis) and avoiding rates of any kind derived from the transfer zone can significantly minimize the measurement-interval bias.

In the same context as box models or other budget diagrams (e.g., [Walling and Collins, 2008](#)), [Hay et al. \(1989\)](#) developed a method for generating mass-balanced paleogeographic maps. These maps aim to depict the source-to-sink redistribution of mass through time by tracking paleotopographic evolution at $\geq 10^7$ yr timescales. Similarly, recent efforts by [Meyers and Peters \(2011\)](#) aim to reconstruct stratigraphic volume and mass distribution through time in relation to long-period ($\geq 10^7$ yr) tectonic and/or sea-level cycles. The utility of such methods to signal propagation has yet to be explicitly addressed. Following [Allen et al. \(2013\)](#), wherein the challenge of estimating sediment supply from strata is termed ‘The Qs problem’, [Michael et al. \(2013\)](#) determine a sediment budget for Late Eocene (~42-34 Ma) foreland basin strata and use resultant grain-size partitioning information to reconstruct tectonic subsidence.

The tectonically active region of southern Asia includes high rates of denudation and mass redistribution, making it an ideal locale to develop longer-term sediment budget concepts. [Johnson \(1994\)](#) accounted for the volume of sediment deposited in foreland basins, deltas, and the Indus and Bengal submarine fans forward of the Himalayas to explore Cenozoic sediment-routing evolution and its relationship to uplift and exhumation. Using information from literature, [Johnson \(1994\)](#) calculated that the Himalayan Main Central Thrust, which is interpreted to have been emplaced ~20 Ma, to be of insufficient volume to balance the depositional systems. Thus, [Johnson \(1994\)](#) posited that regions outside the Himalaya, including Tibet and the Karakoram, might have been significant sediment source areas during the Cenozoic, especially prior to the emplacement of the Himalayan Main Central Thrust.

Clift et al. (2001) used a seismic stratigraphic interpretation to account for the Cenozoic deposition of the Indus Fan. The erosional record on land was constrained from provenance analysis using Pb and Nd isotopic compositions and published thermochronology. Cenozoic land-to-deep sea Indus sediment routing shows a balance of erosion and deposition, with a greater volume of eroded rock during the Neogene, and corresponding greater deposition on the Indus submarine fan during the Neogene. However, the Paleogene was characterized by rapid erosion and coupled accumulation, with deposition focused in the regions of the Katawaz Basin, the Makran accretionary wedge, and the Indus foreland. More recently, Clift et al. (2008) compared published thermochronology from the Himalaya to weathering and climate proxies recorded in Neogene deposits from the South China Sea, Bay of Bengal, and Arabian Sea. Erosion of the Himalaya was interpreted to have intensified ~23-10 Ma, and slowed to ~3.5 Ma, but then began to increase during the Late Pliocene and Pleistocene, which correlates with monsoon intensity interpreted from climate proxies (Clift et al., 2008).

In these studies of sediment budgets of Cenozoic sediment routing systems, the detrital record provides insights into sediment routing evolution, as an indicator of Q_s variability, and insights into the tectonic and climatic controls on erosion and deposition. Similar to the use of modern systems to guide the interpretation of transport and depositional processes from preserved strata, we can use historical-timescale (Figs. 3 and 4) and intermediate-timescale (Fig. 6) sediment-routing systems as analogs of signal-propagation dynamics for deep-time systems. Studies of Cenozoic systems are of particular value to better understand deep-time signal propagation because they combine long-period forcings with relatively well-documented boundary conditions that pre-

Cretaceous systems lack as a result of tectonic reorganization (Romans and Graham, 2013).

5 Discussion and Research Directions

External forcings are initially transformed into an Earth-surface signal through the production of mobile mass that is then redistributed down system as detritus and solutes (Fig. 1) (Allen et al., 2013). The character of that signal and to what extent a signal is preserved in sedimentary records are dependent on the magnitude and frequency of the initial forcing (Fig. 8), on their initial recording or destruction (Wheatcroft et al., 2007), on the responses of the different segments of the sediment-routing system (Castelltort and Van Den Driessche, 2003), on their morphology (e.g., for instance promoting buffered versus reactive sediment and signal transfer) (Covault et al., 2011), and on the ratio of signal to noise, in particular with respect to autogenic dynamics (Jerolmack and Paola, 2010). In this review, we emphasized the importance of sediment supply (Qs) as the main carrier of signals that originate in erosion zones. Thus, approaches for determining Qs by direct observation and measurement (Figs. 3 and 4), calculation from measurement of related process (e.g., denudation via cosmogenic radionuclides; Fig. 6), or estimation from stratigraphy and/or detrital minerals (Figs. 9 and 10) that are reviewed in this paper are critical to understanding how those signals are transmitted through the system. Furthermore, the development of new computational techniques for numerical modeling of source-to-sink sediment transport and deposition promise a deeper understanding of sediment and signal propagation. The benefit of unravelling processes of sediment and signal propagation is an enhanced understanding of the coupling of Earth surface systems

and improved capability to invert stratigraphic and geomorphic records that relate to broader Earth dynamics.

The investigation of signal propagation requires a systems approach, which is provided by the sediment-routing system, or source-to-sink, framework (Allen, 2008a) (Fig. 1). The concept of sediment mass balance is embedded within such a framework and variations in system morphology provide insight into signal propagation and preservation. We emphasized the importance of the transfer zone (Fig. 1) because of its potential role as a ‘buffer’ (e.g., along-system diffusion and temporary floodplain deposition) and, correspondingly, the effect on rates and magnitudes of Qs carried to down-system segments. Such transfer-zone buffering of up-system signals is highly relevant to decoding the meaning of coastal and marine stratigraphic archives (Figs. 4 and 6). The transfer-zone concept at historical timescales ($<10^2$ yr) is elusive because sediment storage can occur across the entire system, including in the erosion zone, leading to potential buffering over short distances. Reconstructing erosion and transfer zones in deep time ($\geq 10^7$ yr) is challenging as a result of incomplete or no preservation of these sediment-routing segments. Characteristics of deep-time sediment production and transfer areas can be interpreted by employing provenance tools, detrital mineral analysis, or application of empirical relationships based on modern systems, and tested with conceptual, analytical, and numerical models.

Implications of signal detection over noise are of paramount importance to interpreting the stratigraphic record of Qs variability. Internal, or autogenic, dynamics of sediment transport, transient storage, and release can introduce noise, lags, and/or completely mask the signal of interest. Experimental and theoretical work has shown that

a Qs signal can be passed to a basin and preserved in the stratigraphic record when its period is similar to or exceeds the characteristic response time of the sediment-routing system, but this is valid only if their periods or magnitudes are also larger than the characteristic timescale of autogenic sediment transport fluctuations (Jerolmack and Paola, 2010). Moreover, autogenic ‘shredding’ is potentially more significant if the length scale of interest, e.g., the system size, is larger than the advection length scale of a particle of sediment (Ganti et al., 2014). Field studies focused on the coarse-grained sediment fraction in small, tectonically active sediment-routing systems of southern California (Fig. 6A) have shown that millennial-scale climate forcings are represented as a measureable signal in the stratigraphic record of the deep-sea segment (Fig. 7) (Romans et al., 2009; Covault et al., 2010). These systems are reactive (*sensu* Allen, 2008b) and comprise sediment-routing segments in close proximity: erosion zones are located adjacent to short transfer zones and offshore confined basins that make up the accumulation zone. Noise over historical timescales can be especially problematic as the observational window is small and the number of signals potentially large (e.g., Sommerfield and Wheatcroft, 2007).

Timescale of observation is fundamentally important for signal analysis in sedimentary systems. Temporal aspects discussed in this review include the duration and period of forcings, the resolution of chronologic tools with which to evaluate Qs (Fig. 2), and preservation into the record. At historical timescales the signal of interest is commonly an individual event, such as an earthquake, flood, or storm. At longer timescales, shifts in the rate and style of sedimentation in the cumulative record can be related to longer-period forcings. We devoted much of our treatment of signal

propagation at the intermediate timescale (10^2 - 10^6 yr) because we consider this to be a critical temporal range in which to understand these dynamics at the scale of entire sediment-routing systems. The shorter-duration end of this range can be linked to direct observation and measurement and the longer, million-year end of this range can serve as a bridge to deep time. The intermediate timescale is the timescale of global climate cycles and the timescale at which climate and tectonic forcings overlap. Moreover, this is the timescale at which meso-scale stratigraphic architecture and high-frequency stratigraphic cycles are created. Our subdivision of timescales in this review is only to aid communication of dominant aspects within each timescale, but we emphasize that sedimentary system research strives to integrate across these timescales.

Our review emphasizes the role of sediment supply, yet we acknowledge the role of accommodation fluctuations in the accumulation zone as an important forcing of stratigraphic patterns (e.g., [Anderson et al., this volume](#)). Deciphering the relative contributions of sediment supply versus accommodation changes to the creation of stratigraphy has been discussed for at least a century (e.g., [Grabau, 1913](#); [Barrell, 1917](#)) and examined via modeling studies for decades (e.g., [Jervy, 1988](#); [Allen and Densmore, 2000](#); [Paola, 2000](#); [Armitage et al., 2013](#)). The nature of signal generation in the sink and potential up-system propagation effects (e.g., [Voller et al., 2012](#)) deserve further attention. Deconvolving the various external forcings from each other and from the products of internal dynamics encoded in the geologic record remains a prime challenge in Earth science ([National Research Council, 2010](#)).

This review provides a set of conceptual and practical tools for reaching informed interpretations of landscape dynamics from the stratigraphic record. These tools include

stratigraphic and sediment-routing system characterization, sediment budget determination, geochronology, detrital mineral analysis (e.g., thermochronology), comparative analog approaches, and modeling techniques to measure, calculate, or estimate the magnitude and frequency of sediment supply signals compared to the characteristic response time of the sediment-routing systems. However, significant research challenges remain, which we distill into four research directions:

1. **Improved documentation of sediment production, transfer, and accumulation rates in natural systems.** The propagation of a signal through a system can be characterized as a phase velocity and, thus, knowledge of time is required. Research aimed at developing new chronometric techniques and studies applying existing techniques in novel ways should continue to be a focus in Earth surface dynamics research. Such work should include the study of and linkage between erosion, transfer, and accumulation zones across a spectrum of system sizes and morphologies as well as across a range of timescales. Within this context, the question of to what degree patterns of stratigraphy (i.e., patterns documented in the absence of absolute chronology) reflect process rates should be further explored.
2. **Grain-size partitioning and signal propagation.** What is the size distribution of sediments exiting the erosion zone as a function of forcings? Our discussion of how sediment supply signals propagate through a system and how they might be preserved in the stratigraphic record is simplistic in that only bulk mass balance is

considered. The variability in transit distances of different grain-size classes for a given forcing might result in the fractionation of catchment-generated signals with certain grain-sizes temporarily stored along the transfer zone. Attempts to capture downstream grain-size fining and to invert it to time-averaged grain-size trends in stratigraphy are promising but still in their embryonic stage (e.g., [Whittaker et al., 2011](#); [Michael et al., 2013](#)). The link between forcings in the erosion zone and the probability density function of the produced grain-size distribution remains a major unknown and constitutes a fundamental input of Qs propagation models at all temporal and spatial scales.

3. **Integration of experimental and modeling approaches with natural systems.**

Many of the modeling efforts reviewed here are based on diffusion assumptions and/or empirical relationships, neither of which truly model sediment transport. Several approaches are currently tackling sediment transfer more explicitly, including using increasingly complete physics of water flow and sediment transport (e.g., Delft3D, broadly labelled Computational Fluid Dynamics, e.g., [Lesser et al., 2004](#)); Reduced Complexity Models (such as cellular automata, e.g. [Murray and Paola, 1997](#); [Liang et al., 2014](#)); and several other modeling approaches as part of CSDMS (Community Surface Dynamics Modeling System) ([Syvitski, 2008](#)). These efforts are complementary and promising, but it is important to maintain a link with natural systems in order to properly assess the appropriate degree of complexity for which model predictions can be compared to data from natural systems. Additionally, scaled-down physical experiments are

contributing valuable insights regarding the timescales of dynamics in depositional landscapes (e.g., autogenic channel avulsion frequency; Paola et al., 2009). However, how such timescales relate to natural-system timescales remains an open question. These experimental approaches must continue to strive to integrate with observation/measurement-based approaches and vice versa.

4. **Integration of particulate transfer dynamics with solute transfer and other geochemical signals.** The denudation of landscapes in erosion zones is the sum of physical and chemical products that are moved down system. Sedimentary archives containing chemical precipitates can be reliable recorders continental weathering as well as atmospheric and oceanic chemistry and have been used to detect climatic and/or oceanographic signals. However, studies that integrate geochemical signal analysis with the concepts and tools for signal propagation as a function of particulate transfer are rare. Additional work combining the particulate and (bio)geochemical perspectives to examine sedimentary system response to environmental change (e.g., Foreman et al., 2013) are necessary to develop a comprehensive understanding of the broader Earth surface system.

Acknowledgements

This review of signal propagation concepts in sedimentary system analysis is purposefully broad in scope and, as a consequence, neither provides a comprehensive treatment of every relevant issue nor does it refer to all studies that have contributed. We hope that readers appreciate the challenge of preparing such a review. BWR, JAC, and

AF would like to acknowledge Steve Graham, George Hilley, and Bill Normark for their mentorship and intellectual guidance. JAC acknowledges intellectual support from Chevron Energy Technology Company scientists Cristian Carvajal, Ashley Harris, Kristy Milliken, Marty Perlmutter, Michael Pyrcz, and Tao Sun. AF would like to thank Angela Hessler and Sam Johnstone for long discussions on deep time. SC acknowledges funding by the Swiss National Science Foundation grant 200021_146822 and inspiring discussion with Guy Simpson, Philip Allen, Jean Van Den Driessche, Sean Willett, Philippe Davy and François Guillocheau. Support for JPW was provided by the National Science Foundation, Marine Geology & Geophysics Program (OCE-0841092), and JPW acknowledges research colleagues for their hard work and valuable insights into signal propagation.

Figure Captions

Figure 1: (A) Schematic portrayal of a sediment supply (Q_s) signal from the erosion zone and how that signal propagates through the system. The leftmost Q_s signal represents as measured at the exit of the erosion zone and for simplicity is the same as the original forcing of interest. The transfer zone Q_s signal is measured within the transfer zone at some distance from exit of erosion zone and the rightmost signal represents that which reaches the accumulation zone and is an input for the stratigraphic record. Dashed lines refer to Q_s signal in up-system segment(s) to illustrate that a signal can be modified during propagation. (B) 2-D profile of a generic sediment-routing system emphasizing erosion, transfer, and accumulation zones (potential for intermediate to deep time

stratigraphic preservation in yellow) and important controls of tectonics (including earthquakes), climate (including storms), base level, and anthropogenic factors. Part B modified from [Castelltort and Van Den Driessche \(2003\)](#).

Figure 2: Overview of three timescales of investigation, some of the chronometric tools with which to constrain process rates, and periods of some of the forcings discussed in this review. Dashed lines at the top emphasize the continuum among the timescales. Temporal range of ‘orogenic cycles’ from [DeCelles et al. \(2009\)](#). Effective dating range of chronometric tools from [Walker \(2005\)](#).

Figure 3: (A) Topography and drainage network of Eel and Mad river catchments, northern California, and bathymetry of the continental margin. Red star denotes location of shelf core x-radiograph shown in (B). (B) X-ray image of shelf reflects bulk density. Light colors (lower bulk density) interpreted as 1995 flood deposit ([Sommerfield and Nittrouer, 1999](#); [Wheatcroft and Borgeld, 2000](#)). (C) Map of Eel-Mad sediment-routing system showing catchment area, areal extent of coastal floodplain, and shelf depocenter (yellow). Red star denotes location of shelf core image shown in (B). (D) Historical timescale sediment budget of the Eel-Mad sediment-routing system showing: 1) there is negligible onshore storage, 2) the shelf stores ~30-50% of the budget, and 3) the remainder moves to the canyon and continental slope. Budget estimations from [Sommerfield and Nittrouer \(1999\)](#) and [Warrick \(2014\)](#).

Figure 4: (A) Topography and drainage network of Ganges, Brahmaputra, and Meghna rivers and bathymetry of shelf, Swatch of No Ground submarine canyon, and part of the

Bengal submarine fan system. Red star denotes location of core record shown in (B). (B) Core from upper canyon showing variation in mean grain size with time compared to storm record from eastern Bengal shelf. Data is from core 96 KL as reported in [Michels et al. \(2003\)](#). (C) Map of Ganges-Brahmaputra-Bengal sediment-routing system showing catchment area, the large delta plain area, shelf depocenter (yellow) and the Bengal submarine channel-levee system. Red star denotes location of core record shown in (B). (D) Historical timescale sediment budget of Ganges-Brahmaputra-Bengal sediment-routing system showing that almost one-third of the budget is stored on the delta plain, ~40% accumulates in the shelf depocenter, split between the topset and foreset regions, and the remaining ~30% is delivered to the canyon and Bengal submarine fan. Budget estimations from [Kuehl et al. \(2005\)](#) and references therein.

Figure 5: The ratio between the timescale of a perturbation (T_p) and the characteristic equilibrium timescale (τ) of the considered system describes the system response to forcing (after [Beaumont et al., 2000](#); see also [Allen, 2008b](#)). A forcing of water discharge (Q_w) and a response of sediment supply (Q_s) are shown for (A) a reactive response when response time is much shorter than timescale of forcing and (B) a buffered response when response time is longer than timescale of perturbation.

Figure 6: Examples of natural sediment routing systems examined at intermediate timescales. (A) Small and tectonically active system, Peninsular Ranges and Continental Borderland of southern California ([Covault et al., 2011](#)); (B) Large and tectonically quiescent system, Texas coastal plain and western Gulf of Mexico ([Hidy et al., 2014](#);

Pirmez et al., 2012); (C) Large and tectonically active system, Indus River and Indus submarine fan (Clift et al., 2014). Indus River floodplain extent from Milliman et al. (1984).

Figure 7: (A) Map showing two southern California sediment-routing systems, each with negligible onshore sediment storage at millennial timescales and correspondingly rapid transfer to offshore submarine fan systems. (B) Alkenone sea surface temperature (SST) proxy for the California coastal region showing a drying trend in the early Holocene followed by the development of the modern El Niño-Southern Oscillation (ENSO), which is known to be sensitive to increased SSTs (Barron et al., 2003). (C) Radiocarbon-constrained weighted-average sediment accumulation rates from Hueneme and Newport submarine fans (Romans et al., 2009; Covault et al., 2011) showing a general correlation of sediment supply to the basin to precipitation regime and, thus, propagation of climatic signal to the stratigraphic record.

Figure 8. Summary of signal propagation from source to sink at intermediate timescales (10^2 - 10^6 yr). The figure considers uplift and climate signals in the erosion zone and their transformation and propagation into sediment supply (Q_s) signals in both the transfer and accumulation zones. The gray shaded regions indicate the cases in which the original forcing has been faithfully transmitted to the accumulation zone. (A) An uplift signal with a period (t) > 50 kyr is transformed into a Q_s signal with same amplitude and period by the erosion zone and transmitted by a short fluvial segment but buffered by a long transfer zone unless $t > 100$ kyr (e.g., Castellort and Van Den Driessche, 2003). (B) An

uplift signal with $t < 50$ kyr is transformed into a buffered Q_s signal by the diffusive catchment dynamics in the erosion zone (Allen and Densmore, 2000) and further buffered by diffusion in the transfer zone (e.g., Castelltort and Van Den Driessche, 2003). (C) A climate signal of water discharge (Q_w) with $t > 100$ kyr in the erosion zone yields a buffered Q_s response signal due to catchment dynamics (Armitage et al., 2013). (D) A climate signal of Q_w with $10 \text{ kyr} < t < 1 \text{ Myr}$ in the erosion zone yields an amplified Q_s response signal due to catchment dynamics (Godard et al., 2013). (E) Climate signal of Q_w with $t > 0$ kyr in erosion zone is transferred to the fluvial system where it is transformed into an amplified Q_s signal by river transport (Simpson and Castelltort, 2012). (F) In the absence of signals coming from the erosion zone, autogenic signals can emerge out of the transfer zone due to the internal dynamics of the fluvial system (e.g., compensational stacking in channelized systems, Hajek et al., 2010), with periodicities and amplitude poorly constrained. Wang et al. (2011) suggest autogenic periodicities of the order of 100 kyr for the Mississippi river delta. (G) Regardless of origin, Q_s signal input to the transfer zone may be ‘shredded’ by the internal dynamics of sediment transport system when their period and amplitude fall within the range of ‘morphodynamic turbulence’ (Jerolmack and Paola, 2010).

Figure 9: Conceptual diagram of crystallization age of detrital minerals (e.g., zircon U-Pb geochronology) as indicator of sediment-routing system connectivity, which can be used to aid reconstruction of sediment supply history from deep-time stratigraphic archives.

Figure 10: Conceptual diagrams of two different types of lag times to be calculated with combinations of cooling ages and depositional age. (A) Schematic cross section depicting trajectory of a particle through cooling via erosional exhumation followed by transport and deposition. (B) Lag time type A is the difference between higher-temperature cooling age (e.g., crystallization age) and lower-temperature cooling age of a single grain (e.g., zircon) and represents exhumation from depth. (C) Lag time type B is the difference between a cooling age and depositional age and represents the time from closure temperature depth to surface exhumation plus transport (and transient storage) in the sediment-routing system prior to deposition (adapted from [Rahl et al., 2007](#)). A consistent Lag time type B calculated through a stratigraphic succession indicates steady erosion whereas departures from that indicate changing erosion rates through time. Note that time within partial annealing/retention zones as well as sedimentary and/or tectonic burial can complicate simple lag time determinations.

References

- Aalto, R., Maurice-Bourgoin, L., Dunne, T., Montgomery, D. R., Nittrover, C. A., and Guyot, J.-L., 2003, Episodic sediment accumulation on Amazonian flood plains influenced by El Nino/Southern Oscillation: *Nature*, v. 425, no. 6957, p. 493-497.
- Adams, M.M. and Bhattacharya, J.P., 2005, No change in fluvial style across a sequence boundary, Cretaceous Blackhawk and Castlegate Formations of central Utah: *Journal of Sedimentary Research*, v., 75, p. 1038-1051.
- Allen, J. R. L., 1974, Reaction, relaxation and lag in natural sedimentary systems: General principles, examples and lessons: *Earth-Science Reviews*, v. 10, p. 263-342.
- Allen, P. A., and Densmore, A. L., 2000, Sediment supply from an uplifting fault block: *Basin Research*, v. 12, p. 367-380.
- Allen, P.A., 2008a, From landscapes into geological history: *Nature*, v. 451, p. 274-276.
- Allen, P.A., 2008b, Time scales of tectonic landscapes and their sediment routing systems: *Geological Society London, Special Publication 296*, p. 7-28.
- Allen, P.A., Armitage, J.J., Carter, A., Duller, R.A., Michael, N.A., Sinclair, H.D., Whitchurch, A.L., and Whittaker, A.C., 2013, The Qs problem: Sediment volumetric balance of proximal foreland basin systems: *Sedimentology*, v. 60, p. 102-130.
- Alexander, C.R. and Simoneau, A.M., 1999, Spatial variability in sedimentary processes on the Eel continental slope, *Marine Geology*: v. 154, p. 243-254.

- 1446 Allison, M.A., 1998, Historical changes in the Ganges-Brahmaputra delta front, *Journal of Coastal*
1447 *Research*, v.14, p.480–490.
- 1448 Allison, M.A., Kineke, G.C., Gordon, E.S., and Goñi, M.A., 2000, Development and reworking
1449 of a seasonal flood deposit on the inner continental shelf off the Atchafalaya River, *Cont.*
1450 *Shelf Res.*, v. 20, p. 2267-2294.
- 1451 Allison, M.A. and Kepple, E.B., 2001, Modern sediment supply to the lower delta plain of the
1452 Ganges–Brahmaputra River in Bangladesh, *Geo-Marine Letters*, v. 21, 66–74.
- 1453 Allison, M.A., Dellapenna, T.M., Gordon, E.S., Mitra, S., and Petsch, S.T., 2010, Impact of
1454 Hurricane Katrina (2005) on shelf organic carbon burial and deltaic evolution, *Geophys.*
1455 *Res. Lett.*, v. 37, p. - L21605.
- 1456 Allison, M.A., Demas, C.R., Ebersole, B.A., Kleiss, B.A., Little, C.D., Meselhe, E.A., Powell,
1457 N.J., Pratt, T.C., and Vosburg, B.M., 2012, A water and sediment budget for the lower
1458 Mississippi-Atchafalaya River in flood years 2008-2010: Implications for sediment
1459 discharge to the oceans and coastal restoration in Louisiana, *Journal of Hydrology*: v.
1460 432-433, p. 84-97.
- 1461 Anderson, J.B., Wallace, D.J., Simms, A.R., Rodriguez, A.B., Weight, R.W.R., and Taha, Z.P.,
1462 this volume, *Recycling sediments between source and sink during a eustatic cycle: Case*
1463 *study of late Quaternary northwestern Gulf of Mexico: Earth-Science Reviews, XXXXX.*
- 1464 Anthony, E.J. and Julian, M., 1999, Source-to-sink sediment transfers, environmental engineering
1465 and hazard mitigation in the steep Var River catchment, French Riviera, southeastern
1466 France: *Geomorphology*, v. 31, p. 337-354.
- 1467 Armitage, J.J., Duller, R.A., Whittaker, A.C., and Allen, P.A., 2011, Transformation of tectonic
1468 and climatic signals from source to sedimentary archive: *Nature Geoscience*, 4, 231-235.
- 1469 Armitage, J.J., Dunkley Jones, T., Duller, R.A., Whittaker, A.C., and Allen, P.A., 2013, Temporal
1470 buffering of climate-driven sediment flux cycles by transient catchment response: *Earth*
1471 *and Planetary Science Letters*, v. 369, p. 200-210.
- 1472 Aslan, A., Autin, W.J., and Blum, M.D., 2005, Causes of river avulsion: Insights from the late
1473 Holocene avulsion of the Mississippi River, USA: *Journal of Sedimentary Research*, v.
1474 75, p 650-664.
- 1475 Attal, M., Tucker, G., Whittaker, A.C., Cowie, P., and Roberts, G.P., 2008, Modeling fluvial
1476 incision and transient landscape evolution: Influence of dynamic channel adjustment:
1477 *Journal of Geophysical Research: Earth Surface*, 113.
- 1478 Atwater, B.F. and Hemphill-Haley, E., 1997, Recurrence intervals for great earthquakes of the
1479 past 3500 years at northeastern Willapa Bay, Washington: U.S. Geological Survey
1480 Professional Paper 1576, 108 p.
- 1481 Atwater, B.F., Carson, B., Griggs, G.B., Johnson, P., and Salmi, M.S., 2014, Rethinking turbidite
1482 paleoseismology along the Cascadia subduction zone, *Geology*: v. 42, p. 827-830.
- 1483 Barnes, P.M., Bostock, H.C., Neil, H.L., Strachan, L.J., and Gosling, M., 2013, A 2300-year
1484 paleoearthquake record of the southern alpine fault and Fiordland subduction zone, New
1485 Zealand, based on stacked turbidites, *Bulletin of the Seismological Society of America*:
1486 v. 103, p. 2424-2446.
- 1487 Barron, J.A., Huesser, L., Herbert, T., and Lyle, M., 2003, High-resolution climatic evolution of
1488 coastal northern California during the past 16,000 years: *Paleoceanography*, 18, 1020.
- 1489 Beerbower, J.R., 1964, Cyclothems and cyclic depositional mechanisms in alluvial plain
1490 sedimentation: Symposium on Cyclic Sedimentation, State Geological Survey of Kansas
1491 Bulletin 169, p. 31-42.
- 1492 Beaumont, C., Kooi, H., and Willett, S., 2000, Coupled tectonic-surface process models with
1493 applications to rifted margins and collisional orogens: In: Summerfield, M.A. (ed.),
1494 *Geomorphology and Global Tectonics*, Wiley, Chichester, pp. 29-55.
- 1495 Barrell, J., 1917, Rhythms and the measurements of geologic time: *Geological Society of*
1496 *America Bulletin*, v. 28, p. 745-904.

- 1497 Bhattacharya, J.P. and Tye, R.S., 2004, Searching for modern Ferron analogs and application to
1498 subsurface interpretation: Regional to wellbore analog for fluvial-deltaic reservoir
1499 modeling: Ferron Sandstone of Utah: AAPG Studies in Geology 50, p 39-57.
- 1500 Bhattacharya, J.P., Copeland, P., Lawton, T.E., and Holbrook, J., this volume, Estimation of
1501 source area, river paleo-discharge, paleoslope, and sediment budgets of linked deep-time
1502 depositional systems and implications for hydrocarbon potential: Earth-Science Reviews,
1503 XXXXX.
- 1504 Bierman, P. and Steig, E.J., 1996, Estimating rates of denudation using cosmogenic isotope
1505 abundances in sediment: Earth Surface Processes and Landforms, v. 21, p. 125-139.
- 1506 Blum, M., and Hattier-Womack, J., 2009, Climate change, sea-level change, and fluvial sediment
1507 supply to deepwater depositional systems: External Controls on Deepwater Depositional
1508 Systems: SEPM Special Publication 92, p. 15-39.
- 1509 Blum, M., Martin, J., Milliken, K., and Garvin, M., 2013, Paleovalley systems: Insights from
1510 Quaternary analogs and experiments: Earth-Science Reviews, v. 116, p. 128-169.
- 1511 Blum, M. and Pecha, M., 2014, Mid-Cretaceous to Paleocene North American drainage
1512 reorganization from detrital zircons: Geology, 10.1130/G35513.1.
- 1513 Boldt, K.V., Lane, P., Woodruff, J.D., and Donnelly, J.P., 2010, Calibrating a sedimentary record
1514 of overwash from Southeastern New England using modeled historic hurricane surges:
1515 Marine Geology, v. 275, p. 127-139.
- 1516 Bonnet, S., and Crave, A., 2003, Landscape response to climate change: Insights from
1517 experimental modeling and implications for tectonic versus climatic uplift of topography:
1518 Geology, v. 31, no. 2, p. 123-126.
- 1519 Bralower, T.J., Paull, C.K., and Mark Leckie, R., 1998, The Cretaceous-Tertiary boundary
1520 cocktail: Chicxulub impact triggers margin collapse and extensive sediment gravity
1521 flows, Geology, v. 26, p. 331-334.
- 1522 Brandon, M.T. and Vance, J.A., 1992, Tectonic evolution of the Cenozoic Olympic subduction
1523 complex, Washington State, as deduced from fission track ages for detrital zircons:
1524 American Journal of Science, v. 292, p. 565-636.
- 1525 Bridge, J.S. and Tye, R.S., 2000, Interpreting the dimensions of ancient fluvial channel bars,
1526 channels, and channel belts from wireline-logs and cores: American Association of
1527 Petroleum Geologists Bulletin, v. 84, p. 1205-1228.
- 1528 Bridge, J.S. and Demicco, R., 2008, Earth Surface Processes, Landforms, and Sediment Deposits:
1529 Cambridge University Press.
- 1530 Brown, W.M., III and Ritter, J.R., 1971, Sediment transport and turbidity in the Eel River basin,
1531 California, U.S. Geological Survey Water-Supply Paper 1986.
- 1532 Brown, E.T., Stallard, R.F., Larsen, M.C., Raisbeck, G.M., and Yio, F., 1995, Denudation rates
1533 determined from the accumulation of in situ-produced ¹⁰Be in the Luquillo experimental
1534 forest, Puerto Rico: Earth and Planetary Science Letters, v. 129 p. 193-202.
- 1535 Brush, G.S., 2001, Natural and Anthropogenic Changes in Chesapeake Bay During the Last 1000
1536 Years, Human and Ecological Risk Assessment: An International Journal: v. 7, p. 1283-
1537 1296.
- 1538 Burbank, D.W. and Anderson, R.S., 2011, Tectonic Geomorphology: John Wiley and Sons.
- 1539 Cacchione, D.A., Sternberg, R.W., and Ogston, A.S., 2006, Bottom instrumented tripods:
1540 History, applications, and impacts, Cont. Shelf Res.: v. 26, p. 2319-2334.
- 1541 Carrapa, B., Wijbrans, J., and Bertotti, G., 2003, Episodic exhumation in the Western Alps:
1542 Geology, v. 31, p. 601-604.
- 1543 Carrapa, B., 2010, Resolving tectonic problems by dating detrital minerals: Geology, v. 38, p.
1544 191-192.
- 1545 Carter, L., Milliman, J.D., Talling, P.J., Gavey, R., and Wynn, R.B., 2012, Near-synchronous and
1546 delayed initiation of long run-out submarine sediment flows from a record-breaking river
1547 flood, offshore Taiwan, Geophys. Res. Lett., v. 39, L12603, p. 1-5.

1548 Carvajal, C. and Steel, R., 2012, Source-to-sink sediment volumes within a tectono-stratigraphic
 1549 model for a Laramide shelf-to-deep-water basin: Methods and results, in: *Tectonics of*
 1550 *Sedimentary Basins: Recent Advances*, C. Busby and A. Azor (Eds.). John Wiley &
 1551 Sons, Ltd, Chichester, UK, pp. 131-151.
 1552 Castelltort, S. and Van Den Driessche, J., 2003, How plausible are high-frequency sediment
 1553 supply-driven cycles in the stratigraphic record?: *Sedimentary Geology*, v. 157, p. 3-13.
 1554 Clift, P., Shimizu, N., Layne, G., Blusztajn, J., Gaedicke, C., Schluter, H.-U., Clark, M., and
 1555 Amjad, S., 2001, Development of the Indus Fan and its significance for the erosional
 1556 history of the Western Himalaya and Karakoram: *Geological Society of America*
 1557 *Bulletin*, v. 113, p. 1039-1051.
 1558 Clift, P.D., Hodges, K.V., Heslop, D., Hannigan, R., Van Long, H., and Calves, G., 2008,
 1559 Correlation of Himalayan exhumation rates and Asian monsoon: *Nature Geoscience*, v. 1,
 1560 p. 875-880.
 1561 Clift, P.D. and Giosan, L., 2014, Sediment fluxes and buffering in the post-glacial Indus Basin:
 1562 *Basin Research*, v. 26, p. 369-386.
 1563 Cooper, S.R., McGlothlin, S.K., Madritch, M., and Jones, D.L., 2004, Paleoecological evidence
 1564 of human impacts on the Neuse and Pamlico estuaries of North Carolina, USA, *Estuaries*:
 1565 v. 27, p. 617-633.
 1566 Corbett, D.R., Walsh, J.P., Harris, C.K., Ogston, A.S, and Orpin, A.R., 2014, Formation and
 1567 preservation of sedimentary strata from coastal events: Insights from measurements and
 1568 modeling: *Continental Shelf Research*, v. 86, p. 1-5.
 1569 Covault, J.A., Normark, W.R., Romans, B.W., and Graham, S.A., 2007, Highstand fans in the
 1570 California Borderland: The overlooked deep-water depositional system: *Geology*, v. 35,
 1571 p. 783-786.
 1572 Covault, J.A. and Graham, S.A., 2010, Submarine fans at all sea-level stands: Tectono-
 1573 morphologic and climatic controls on terrigenous sediment delivery to the deep sea:
 1574 *Geology*, v. 38, p. 939-942.
 1575 Covault, J.A., Romans, B.W., Fildani, A., McGann, M., and Graham, S.A., 2010, Rapid climatic
 1576 signal propagation from source to sink in a southern California sediment-routing system:
 1577 *The Journal of Geology*, v. 118, p. 247-259.
 1578 Covault, J.A., Romans, B.W., Graham, S.A., Fildani, A., and Hilley, G.E., 2011, Terrestrial
 1579 source to deep-sea sink sediment budgets at high and low sea levels: Insights from
 1580 tectonically active southern California: *Geology*, v. 39, p. 619-622.
 1581 Covault, J.A., Craddock W., Romans, B.W., Fildani, A., and Gosai, M., 2013, Spatial and
 1582 temporal variations in landscape evolution: Historic and longer-term sediment flux
 1583 through global catchments: *The Journal of Geology*, v. 121, p. 35-56.
 1584 Crutzen, P.J. and Stoermer, E.F., 2000, The Anthropocene. *Global Change Newsletter*, n. 41, p.
 1585 17-18.
 1586 Cundy, A.B., Croudace, I.W., Cearreta, A., and Irabien, M.J., 2003, Reconstructing historical
 1587 trends in metal input in heavily-disturbed, contaminated estuaries: Studies from Bilbao,
 1588 Southampton Water and Sicily, *Appl. Geochem.*: v. 18, p. 311-325.
 1589 Curray, J.R., Emmel, F.J., and Moore, D.G., 2002, The Bengal Fan: Morphology, geometry,
 1590 stratigraphy, history, and processes: *Marine and Petroleum Geology*, v. 19, p. 1191-1223.
 1591 Curray, J.R., 2014, The Bengal Depositional System: From rift to orogeny, *Mar. Geol.*: v. 352, p.
 1592 59-69.
 1593 Dadson, S.J., Hovius, N., Chen, H., Dade, W.B., Lin, J., Hsu, M., Lin, C., Horng, M., Chen, T.,
 1594 Milliman, J., and Stark, C.P., 2004, Earthquake-triggered increase in sediment delivery
 1595 from an active mountain belt, *Geology*: v. 32, p. 733-736.
 1596 Dail, M.B., Reide Corbett, D., and Walsh, J.P., 2007, Assessing the importance of tropical
 1597 cyclones on continental margin sedimentation in the Mississippi delta region, *Cont. Shelf*
 1598 *Res.*: v. 27, p. 1857-1874.

Davidson, S.K. and North, C.P., 2009, Geomorphological regional curves for prediction of drainage area and screening modern analogues for rivers in the rock record: *Journal of Sedimentary Research*, v. 79, p. 773-792.

Day Jr., J.W., Boesch, D.F., Clairain, E.J., Kemp, G.P., Laska, S.D., Mitsch, W.J., Orth, K., Mashriqui, H., Reed, D.J., Shabman, L., Simenstad, C.A., Streever, B.J., Twilley, R.R., Watson, C.C., Wells, J.T., and Whigham, D.F., 2007, Restoration of the Mississippi Delta: Lessons from Hurricanes Katrina and Rita, *Science*: v. 315, p. 1679-1684.

de la Fuente, J., Miller, A., Elder, D., Faust, R., and Snavely, W., 2006, Landslide sediment production rates in the middle fork and upper Eel River basins, northern California, *Proceedings of the Eighth Federal Interagency Sedimentation Conference (8thFISC)*, April 2-6, 2006, Reno, NV, USA.

DeCelles, P.G., Ducea, M.N., Kapp, P., and Zandt, G., 2009, Cyclicity in Cordilleran orogenic systems: *Nature Geoscience*, v. 2, p. 251-257.

Dickinson, W.R., 1974, Plate tectonics and sedimentation: *SEPM Special Publication* 22.

Dickinson, W.R. and Gehrels, G.E., 2003, U-Pb ages of detrital zircons from Permian and Jurassic eolian sandstones of the Colorado Plateau, USA: Paleogeographic implications: *Sedimentary Geology*, v. 163, p. 29-66.

Dinehart, R.L. and Burau, J.R., 2005, Repeated surveys by acoustic Doppler current profiler for flow and sediment dynamics in a tidal river, *Journal of Hydrology*: v. 314, p. 1-21.

Drexler, T.M., Nittrouer, C.A., and Mullenbach, B.L., 2006, Impact for local morphology on sedimentation in a submarine canyon, ROV studies in Eel Canyon, Northern California, U.S.A, *Journal of Sedimentary Research*: v. 76, p. 839-853.

Edmonds, D. and Slingerland, R., 2007, Mechanics of river mouth bar formation: Implications for the morphodynamics of delta distributary networks: *Journal of Geophysical Research – Earth Surface* 112.

Falcini, F., Khan, N.S., Macelloni, L., Horton, B.P., Lutken, C.B., Mckee, K.L., Santoleri, R., Colella, S., Li, C., Volpe, G., D'emidio, M., Salusti, A., and Jerolmack, D.J., 2012, Linking the historic 2011 Mississippi River flood to coastal wetland sedimentation, *Nature Geoscience*, v. 5, p. 803-807.

Fisk, H.N., 1944, Geological investigation of the alluvial valley of the lower Mississippi River: *Mississippi River Commission*, Vicksburg, Mississippi, p. 1-78.

Fosdick, J.C., Grove, M., Graham, S.A., Hourigan, J.K., Lovera, O., and Romans, B.W., 2014, Detrital thermochronologic record of burial heating and sediment recycling in the Magallanes foreland basin, Patagonian Andes: *Basin Research*, 10.1111/bre.12088.

Furbish, D.J. and Fagherazzi, S., 2001, Stability of creeping soil and implications for hillslope evolution: *Water Resources Research*, v. 37, p. 2607-2618.

Ganti, V., Lamb, M.P., and McElroy, B., 2014, Quantitative bounds on morphodynamics and implications for reading the sedimentary record: *Nature Communications*, v. 5:3298, 10.1038/ncomms4298.

Gardner, T.W., Jorgensen, D.W., Shuman, C., and Lemieux, C.R., 1987, Geomorphic and tectonic process rates: Effects of measured time interval: *Geology*, v. 15, p. 259-261.

Garver, J.I., Brandon, M.T., Roden-Tice, M., and Kamp, P.J., 1999, Exhumation history of orogenic highlands determined by detrital fission-track thermochronology: *Geological Society London Special Publication* 154, p. 283-304.

Gasparini, N.M., Whipple, K.X., and Bras, R.L., 2007, Predictions of steady state and transient landscape morphology using sediment-flux-dependent river incision models: *Journal of Geophysical Research – Earth Surface*, 112.

Gerber, T.P., Pratson, L.F., Kuehl, S., Walsh, J.P., Alexander, C., and Palmer, A., 2010, The influence of sea level and tectonics on Late Pleistocene through Holocene sediment storage along the high-sediment supply Waipaoa continental shelf, *Mar. Geol.*: v. 270, p. 139-159.

1650 Gilli, A., Anselmetti, F., Glur, L., Wirth, S., 2013. Lake Sediments as Archives of Recurrence
1651 Rates and Intensities of Past Flood Events, in: Schneuwly-Bollschweiler, M., Stoffel, M.,
1652 Rudolf-Miklau, F. (Eds.). Springer Netherlands, pp. 225-242.

1653 Giosan, L., Constantinescu, S., Clift, P.D., Tabrez, A.R., Danish, M., and Inam, A., 2006, Recent
1654 morphodynamics of the Indus delta shore and shelf: *Continental Shelf Research*, v. 26, p.
1655 1668-1684.

1656 Girardclos, S., Schmidt, O.T., Sturm, M., Ariztegui, D., Pugin, A., and Anselmetti, F.S., 2007,
1657 The 1996 AD delta collapse and large turbidite in Lake Brienz, *Mar. Geol.*: v. 241, p.
1658 137-154.

1659 Godard, V., Tucker, G. E., Burch Fisher, G., Burbank, D. W., and Bookhagen, B., 2013,
1660 Frequency-dependent landscape response to climatic forcing: *Geophysical Research*
1661 *Letters*, v. 40, no. 5, p. 859-863.

1662 Goff, J.A., Wheatcroft, R.A., Lee, H., Drake, D.E., Swift, D.J.P., and Fan, S., 2002, Spatial
1663 variability of shelf sediments in the STRATAFORM natural laboratory, Northern
1664 California, *Cont. Shelf Res.*: v. 22, p. 1199-1223.

1665 Goldfinger, C., Nelson, C.H., and Johnson, J.E., 2003, Holocene earthquake records from the
1666 Cascadia subduction zone and northern San Andreas fault based on precise dating of
1667 offshore turbidites, *Annu. Rev. Earth Planet. Sci.*: v. 31, p. 555-577.

1668 Goldfinger, C., Nelson, C.H. , Morey, A.E. , Johnson, J.E., Patton, J., Karabanov, E., Gutiérrez-
1669 Pastor, J., Eriksson, A.T., Gràcia, E., Dunhill, G. , Enkin, R.J. , Dallimore, A., and
1670 Vallier, T., 2012, Turbidite event history—methods and implications for Holocene
1671 paleoseismicity of the Cascadia subduction zone, U.S. Geological Survey Professional
1672 Paper 1661-F.

1673 Goni, M.A., Alleau, Y., Corbett, D., Walsh, J.P., Mallinson, D., Allison, M.A., Gordon, E.,
1674 Petsch, S., and Dellapena, T.M., 2007, The effects of Hurricanes Katrina and Rita on the
1675 seabed of the Louisiana shelf, *The Sedimentary Record*, v. 5, p. 5-9.

1676 Goodbred, S.L. and Kuehl, S.A., 1998, Floodplain processes in the Bengal Basin and the storage
1677 of Ganges-Brahmaputra river sediment: an accretion study using Cs- 137 and Pb-210
1678 geochronology, *Sediment. Geol.*: v. 121, p. 239-258.

1679 Goodbred Jr, S.L. and Kuehl, S.A., 1999, Holocene and modern sediment budgets for the
1680 Ganges-Brahmaputra river system: Evidence for highstand dispersal to floodplain, shelf,
1681 and deep-sea depocenters: *Geology*, v. 27, p. 559-562.

1682 Goodbred, S.J. and Kuehl, S.A., 2000a, Enormous Ganges-Brahmaputra sediment discharge
1683 during strengthened early Holocene monsoon, *Geology*: v. 28, p. 1083-1086.

1684 Goodbred Jr, S.L. and Kuehl, S.A., 2000b, The significance of large sediment supply, active
1685 tectonism, and eustasy on margin sequence development: Late Quaternary stratigraphy
1686 and evolution of the Ganges-Brahmaputra delta: *Sedimentary Geology*, v. 133, p. 227-
1687 248.

1688 Goodbred Jr, S. L., 2003, Response of the Ganges dispersal system to climate change: a source-
1689 to-sink view since the last interstade: *Sedimentary Geology*, v. 162, no. 1–2, p. 83-104.

1690 Grabau, A.W., 1913, *Principles of Stratigraphy*: A.G. Seiler and Company, New York.

1691 Graham, S.A., Tolson, R., DeCelles, P., Ingersoll, R., Bargar, E., Caldwell, L., Cavazza, W.,
1692 Edwards, D., Follo, M., and Handschy, J., 1986, Provenance modeling as technique for
1693 analyzing source terrane evolution and controls on foreland sedimentation: *Foreland*
1694 *Basins*, Blackwell Publishing, p. 425-436.

1695 Granger, D.E., Kirchner, J.W., and Finkel, R., 1996, Spatially averaged long-term erosion rates
1696 measured from in situ-produced cosmogenic nuclides in alluvial sediment: *The Journal of*
1697 *Geology*, p. 249-257.

1698 Hack, J.T., 1957, *Studies of longitudinal stream profiles in Virginia and Maryland*: United States
1699 Geological Survey Professional Paper 294-B.

- 1700 Hajek, E.A., Heller, P.L., and Sheets, B.A., 2010, Significance of channel-belt clustering in
1701 alluvial basins: *Geology*, v. 38, p. 535-538.
- 1702 Hale, R.P., Ogston, A.S., Walsh, J.P., and Orpin, A.R., 2014, Sediment transport and event
1703 deposition on the Waipaoa River Shelf, New Zealand, *Cont. Shelf Res.*: v. 86, p. 52-65.
- 1704 Hanebuth, T.J.J., Kudrass, H.R., Linstadter, J., Islam, B., and Zander, A.M., 2013, Rapid coastal
1705 subsidence in the central Ganges-Brahmaputra delta (Bangladesh) since the 17th century
1706 deduced from submerged salt producing kilns, *Geology*: v. 41, p. 987-990.
- 1707 Hays, J.D., Imbrie, J., and Shackleton, N.J., 1976, Variations in the Earth's orbit: Pacemaker of
1708 the ice ages: *Science*, v. 194, p. 1121-1132.
- 1709 Hay, W.W., Wold, C.N., and Shaw, C.A., 1989, Mass-balanced paleogeographic maps:
1710 Background and input requirements: *Quantitative Dynamic Stratigraphy*, Prentice Hall, p.
1711 261-275.
- 1712 Heezen, B. C. and Ewing, M., 1952, Turbidity currents and submarine slumps, and the 1929
1713 Grand Banks earthquake, *Am. J. Sci.*, 250, 849-873.
- 1714 Hidy, A.J., Gosse, J.C., Blum, M.D., and Gibling, M.R., 2014, Glacial-interglacial variation in
1715 denudation rates from interior Texas, USA, established with cosmogenic nuclides: *Earth
1716 and Planetary Science Letters*, v. 390, p. 209-221.
- 1717 Hinderer, M., 2012, From gullies to mountain belts: A review of sediment budgets at various
1718 scales: *Sedimentary Geology*, v. 280, p. 21-59.
- 1719 Holbrook, J. and Wanas, H., 2014, A fulcrum approach to assessing source-to-sink mass balance
1720 using channel paleohydrologic parameters derivable from common fluvial data sets with
1721 an example from the Cretaceous of Egypt: *Journal of Sedimentary Research*, v. 84, p.
1722 349-372.
- 1723 Hovius, N., Stark, C.P., and Allen, P.A., 1997, Sediment flux from a mountain belt derived by
1724 landslide mapping, *Geology*: v. 25, p. 231-234.
- 1725 Hoyal, D. and Sheets, B., 2009, Morphodynamic evolution of experimental cohesive deltas:
1726 *Journal of Geophysical Research – Earth Surface*, 114.
- 1727 Hughes Clarke, J.E., Brucker, S., Muggah, J., Church, I., Cartwright, D., Kuus, P., Hamilton, T.,
1728 Pratomo, D., and Eisan, B., 2012, The Squamish ProDelta: Monitoring Active Landslides
1729 and Turbidity Currents: *Canadian Hydrographic Conference 2012, Proceedings*, 15pp.
- 1730 Jaeger, J.M. and Koppes, M., this volume, *The role of the cryosphere in source to sink systems:*
1731 *Earth-Science Reviews*, XXXXX.
- 1732 Jerolmack, D.J. and Paola, C., 2010, Shredding of environmental signals by sediment transport:
1733 *Geophysical Research Letters*, v. 37, L19401.
- 1734 Jervey, M.T., 1988, Quantitative geological modeling of siliciclastic rock sequences and their
1735 seismic expression, in: Wilgus, Hastings, Kendall, Posamentier, Ross, Van Wagoner,
1736 eds., *Sea Level Changes – an Integrated Approach*.
- 1737 Johnson, M., 1994, Volume balance of erosional loss and sediment deposition related to
1738 Himalayan uplifts: *Journal of the Geological Society*, v. 151, p. 217-220.
- 1739 Kao, S.-J., Hilton, R.G., Selvaraj, K., Dai, M., Zehetner, F., Huang, J.-C., Hsu, S.-C., Sparkes, R.,
1740 Liu, J.T., Lee, T.-Y., Yang, J.-Y.T., Galy, A., Xu, X., and Hovius, N., 2014, Preservation
1741 of terrestrial organic carbon in marine sediments offshore Taiwan: Mountain building and
1742 atmospheric carbon dioxide sequestration: *Earth Surface Dynamics*, v. 2, p. 127-139.
- 1743 Kniskern, T.A., Mitra, S., Orpin, A.R., Harris, C.K., Walsh, J.P., and Corbett, D.R., 2014,
1744 Characterization of a flood-associated deposit on the Waipaoa River shelf using
1745 radioisotopes and terrigenous organic matter abundance and composition, *Cont. Shelf
1746 Res.*: v. 86, p. 66-84.
- 1747 Kolker, A.S., Li, C., Walker, N.D., Pilley, C., Ameen, A.D., Boxer, G., Ramatchandirane, C.,
1748 Ullah, M., and Williams, K.A., 2014, The impacts of the great Mississippi/Atchafalaya
1749 River flood on the oceanography of the Atchafalaya Shelf, *Cont. Shelf Res.*, v. 86, p. 17-
1750 33.

1751 Korup, O., Hayakawa, Y., Codilean, A.T., Matsushi, Y., Saito, H., Oguchi, T., and Matsuzaki, H.,
 1752 2014, Japan's sediment flux to the Pacific Ocean revisited, *Earth-Sci. Rev.*: v. 135, p. 1-
 1753 16.
 1754 Kuehl, S.A., Nittrouer, C.A., Allison, M.A., Faria, L.E.C., Dukat, D.A., Jaeger, J.M., Pacioni,
 1755 T.D., Figueiredo, A.G., and Underkoffler, E.C., 1996, Sediment deposition,
 1756 accumulation, and seabed dynamics in an energetic fine-grained coastal environment,
 1757 *Cont. Shelf Res.*: v. 16, p. 787-815.
 1758 Kuehl, S.A., Levy, B.M., Moore, W.S., and Allison, M.A., 1997, Subaqueous delta of the
 1759 Ganges-Brahmaputra river system, *Mar. Geol.*: v. 144, p. 81-96.
 1760 Kuehl, S.A., Carter, L., Gomez, B. and Trustrum, N., 2003, Holistic approach offers potential to
 1761 quantify mass fluxes across continental margins, *Eos*, v. 84, 379/388.
 1762 Kuehl, S.A., Allison, M.A., Goodbred, S.L. and Kudrass, H., 2005, The Ganges-Brahmaputra
 1763 Delta, in: Giosan, L. and Bhattacharya, J., eds., *River Deltas - Concepts, Models, and*
 1764 *Examples*, SEPM Special Publication No. 83, p. 413-434.
 1765 Kuehl, S.A., Alexander, C.R., Blair, N.E., Harris, C.K., Marsaglia, K.M., Ogston, A.S., Orpin, A.
 1766 R., Roering, J., Bever, A., Bilderback, E.L., Carter, L., Cerovski-Darriau, C., Childress,
 1767 L.B., Corbett, R., Hale, R., Leithold, E.L., Litchfield, N., Moriarty, J.M., Page, M.J.,
 1768 Pierce, L.E., Upton, P., and Walsh, J.P., this volume, A source to sink perspective of the
 1769 Waipaoa River margin: *Earth-Science Reviews*, XXXXX.
 1770 Lamb, M.P., Nittrouer, J.A., Mohrig, D., and Shaw, J., 2012, Backwater and river plume controls
 1771 on scour upstream of river mouths: Implications for fluvio-deltaic morphodynamics:
 1772 *Journal of Geophysical Research*, 117, F01002.
 1773 Lawton, T.F., 2014, Small grains, big rivers, continental concepts: *Geology*, v. 42, p. 639-640.
 1774 Leithold, E.L., Perkey, D.W., Blair, N.E., and Creamer, T.N., 2005, Sedimentation and carbon
 1775 burial on the northern California continental shelf: the signatures of land-use change,
 1776 *Cont. Shelf Res.*: v. 25, p. 349-371.
 1777 Leithold, E.L., Blair, N.E., and Wegmann, K.W., this volume, Source to sink sedimentary
 1778 systems and the global C-cycle: A river runs through it: *Earth-Science Reviews*,
 1779 XXXXX.
 1780 Leopold, L.B. and Maddock, T., 1953, The hydraulic geometry of stream channels and some
 1781 physiographic implications: United States Geological Survey Professional Paper 252.
 1782 Lesser, G.R., Roelvink, J.A., van Kester, J.A.T.M., and Stelling, G., 2004, Development and
 1783 validation of a three-dimensional morphological model: *Journal of Coastal Engineering*,
 1784 v. 51, p. 883-915.
 1785 Liang, M., Voller, V.R., and Paola, C., 2014, A reduced-complexity model for river delta
 1786 formation – Part 1: Modeling deltas with channel dynamics: *Earth Surface Dynamics*
 1787 *Discussions*, v. 2, p. 823-869.
 1788 Limmer, D.R., Kohler, C.M., Hillier, S., Moreton, S.G., Tabrez, A.R., and Clift, P.D., 2012,
 1789 Chemical weathering and provenance evolution of Holocene-Recent sediments from the
 1790 Western Indus Shelf, Northern Arabian Sea, inferred from physical and mineralogical
 1791 properties: *Marine Geology*, v. 326, p. 101-115.
 1792 Mallarino, G., Beaubouef, R.T., Droxler, A.W., Abreu, V., and Labeyrie, L., 2006, Sea level
 1793 influence on the nature and timing of a minibasin sedimentary fill (northwestern slope of
 1794 the Gulf of Mexico): *AAPG Bulletin*, v. 90, p. 1089-1119.
 1795 Marr, J.G., Swenson, J.B., Paola, C., and Voller, R., 2002, A two-diffusion model of fluvial
 1796 stratigraphy in close depositional basins: *Basin Research*, 12, p. 381-398..
 1797 Matthai, H., 1990, *Floods: Surface Water Hydrology*, Geological Society of America, Boulder,
 1798 Colorado, p. 97-120.
 1799 McKee, K.L. and Cherry, J.A., 2009, Hurricane Katrina Sediment Slowed Elevation Loss in
 1800 Subsiding Brackish Marshes of the Mississippi River Delta, *Wetlands*, v. 29, p. 2-15.

1801 McLennan, S.M., Taylor, S., and Eriksson, K.A., 1983, Geochemistry of Archean shales from the
1802 Pilbara Supergroup, western Australia: *Geochimica et Cosmochimica Acta*, v. 47, p.
1803 1211-1222.

1804 Meade, R.H., Yuzyk, T.R., and Day, T.J., 1990, Movement and storage of sediment in rivers of
1805 the United States and Canada. in: M.G. Wolman and H.C. Riggs, eds., *The Geology of*
1806 *North America*, vol. 1, Surface Water Hydrology: Boulder, Geol. Soc. America, p. 255-
1807 280.

1808 Métivier, F. and Gaudemer, Y., 1999, Stability of output fluxes of large rivers in South and East
1809 Asia during the last 2 million years: Implications on floodplain processes: *Basin*
1810 *Research*, v. 11, p. 293-303.

1811 Meyers, S.R. and Peters, S.E., 2011, A 56 million-year rhythm in North American sedimentation
1812 during the Phanerozoic: *Earth and Planetary Science Letters*, v. 303, p. 174-180.

1813 Michael, N.A., Whittaker, A.C., and Allen, P.A., 2013, The functioning of sediment routing
1814 systems using a mass balance approach: Example from the Eocene of the southern
1815 Pyrenees: *The Journal of Geology*, v. 121, p. 581-606.

1816 Michels, K.H., Kudrass, H.R., Hübscher, C., Suckow, A., and Wiedicke, M., 1998, The
1817 submarine delta of the Ganges–Brahmaputra: cyclone-dominated sedimentation patterns,
1818 *Mar. Geol.*: v. 149, p. 133-154.

1819 Michels, K.H., Suckow, A., Breitzke, M., Kudrass, H.R., and Kottke, B., 2003, Sediment
1820 transport in the shelf canyon “Swatch of No Ground” (Bay of Bengal), *Deep Sea*
1821 *Research Part II: Topical Studies in Oceanography*: v. 50, p. 1003-1022.

1822 Miller, K.G., Komins, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E., Sugarman,
1823 P.J., Cramer, B.S., Christie-Blick, N., and Pekar, S.F., 2005, The Phanerozoic record of
1824 global sea-level change: *Science*, v. 310, p. 1293-1298.

1825 Miller, K.G., Komins, M.A., Browning, J.V., Wright, J.D., Mountain, G.S., Katz, M.E.,
1826 Sugarman, P.J., Cramer, B.S., Christie-Blick, N., and Pekar, S.F., 2005, The Phanerozoic
1827 record of global sea-level change: *Science*, 310, 1293-1298.

1828 Miller, A.J. and Kuehl, S.A., 2010, Shelf sedimentation on a tectonically active margin: A
1829 modern sediment budget for Poverty continental shelf, New Zealand, *Mar. Geol.*: v. 270,
1830 p. 175-187.

1831 Milliman, J.D. and Syvitski, J.P., 1992, Geomorphic/tectonic control of
1832 sediment discharge to the ocean: The importance of small mountainous rivers: *The*
1833 *Journal of Geology*, v. 100, p. 525-544.

1834 Milliman, J.D., Quraishie, G.S., and Beg, M.A.A., 1984, Sediment discharge from the Indus
1835 River to the ocean: Past, present, and future: *Marine Geology and Oceanography of the*
1836 *Arabian Sea and Coastal Pakistan*, p. 65-70.

1837 Milliman, J.D. and Syvitski, J.P.M., 1992, Geomorphic/tectonic control of sediment discharge to
1838 the ocean: The importance of small mountainous rivers: *The Journal of Geology*, 100,
1839 525-544.

1840 Milliman, J.D. and Farnsworth, K.L., 2011, River discharge to the coastal ocean: A global
1841 synthesis: Cambridge University Press, 392 p.

1842 Moernaut, J., De Batist, M., Charlet, F., Heirman, K., Chapron, E., Pino, M., Brummer, R., and
1843 Urrutia, R., 2007, Giant earthquakes in South-Central Chile revealed by Holocene mass-
1844 wasting events in Lake Puyehue, *Sediment. Geol.*: v. 195, p. 239-256.

1845 Montgomery, D.R., 2001, Slope distributions, threshold hillslopes, and steady-state topography:
1846 *American Journal of Science*, v. 301, p. 432-454.

1847 Moriarty, J.M., Harris, C.K., and Hadfield, M.G., 2014, A hydrodynamic and sediment transport
1848 model for the Waipaoa Shelf, New Zealand: Sensitivity of fluxes to spatially varying
1849 erodibility and model nesting: *J. Mar. Sci. Eng.*, 2, 336-369.

1850 Mullenbach, B.L., Nittrouer, C.A., Puig, P., and Orange, D.L., 2004, Sediment deposition in a
1851 modern submarine canyon: eel canyon, northern California, *Mar. Geol.*: v. 211, p. 101-
119.

- 1852 Mullenbach, B.L. and Nittrouer, C.A., 2006, Decadal record of sediment export to the deep sea
1853 via Eel Canyon, *Cont. Shelf Res.*: v. 26, p. 2157-2177.
- 1854 Murray, A.B. and Paola, C., 1997, Properties of cellular braided-stream model: *Earth Surface*
1855 *Processes and Landforms*, v. 22, p. 1001-1025.
- 1856 Myrow, P.M. and Southard, J.B., 1996, Tempestite deposition, *Journal of Sedimentary Research*,
1857 v. 66, p. 875-887.
- 1858 National Research Council, 2010, *Landscapes on the Edge: New Horizons for Research on*
1859 *Earth's Surface. Committee on Challenges and Opportunities in Earth Surface Processes*,
1860 National Research Council. ISBN: 0-309-14205-0, 180 p.
- 1861 Nittrouer, C.A., Austin, J.A., Field, M.E., Kravitz, J.H., Syvitski, J.P.M., and P.L. Wiberg. 2007.
1862 *Continental margin sedimentation: from sediment transport to sequence stratigraphy*.
1863 Malden, MA: Blackwell Pub. for the International Association of Sedimentologists. 549
1864 pgs.
- 1865 Nittrouer, J.A., Allison, M.A., and Campanella, R., 2008, Bedform transport rates for the
1866 lowermost Mississippi River, *Journal of Geophysical Research F: Earth Surface*: v. 113.
1867 F03004, p. 1-16.
- 1868 Noren, A.J., Biennan, P.R., Steig, E.J., Lini, A., and Southon, J., 2002, Millennial-scale
1869 storminess variability in the northeastern United States during the Holocene epoch,
1870 *Nature*: v. 419, p. 821-824.
- 1871 Normark, W.R., Piper, D.J., Romans, B.W., Covault, J.A., Dartnell, P., and Sliter, R.W., 2009,
1872 Submarine canyon and fan systems of the California Continental Borderland: *Geological*
1873 *Society of America Special Paper* 454, p. 141-168.
- 1874 Ogston, A.S., Cacchione, D.A., Sternberg, R.W., and Kineke, G.C., 2000, Observations of storm
1875 and river flood-driven sediment transport on the northern California continental shelf,
1876 *Cont. Shelf Res.*: v. 20, p. 2141-2162.
- 1877 Painter, C.S., Carrapa, B., DeCelles, P.G., Gehrels, G.E., and Thomson, S.N., 2014, Exhumation
1878 of the North American Cordillera revealed by multi-dating of Upper Jurassic-Upper
1879 Cretaceous foreland basin deposits: *Geological Society of America Bulletin*,
1880 10.1130/B30999.1.
- 1881 Palinkas, C.M., Nittrouer, C.A., Wheatcroft, R.A., and Langone, L., 2005, The use of ⁷Be to
1882 identify event and seasonal sedimentation near the Po River delta, *Adriatic Sea, Mar.*
1883 *Geol.*: v. 222-223, p. 95-112.
- 1884 Paola, C., Heller, P.L., Angevine, C.L., 1992, The large-scale dynamics of grain-size variation in
1885 alluvial basins, 1: Theory: *Basin Research*, v. 4, p. 73-90.
- 1886 Paola, C., 2000, Quantitative models of sedimentary basin filling: *Sedimentology*, 47, 121-178.
- 1887 Paola, C., Straub, K., Mohrig, D., and Reinhardt, L., 2009, The “unreasonable effectiveness” of
1888 stratigraphic and geomorphic experiments: *Earth-Science Reviews*, v. 97, p. 1-43.
- 1889 Paola, C. and Voller, V., 2005, A generalized Exner equation for sediment mass balance: *Journal*
1890 *of Geophysical Research – Earth Surface*, v. 110.
- 1891 Paola, C. and Martin, J.M., 2012, Mass-balance effects in depositional systems: *Journal of*
1892 *Sedimentary Research*, v. 82, p. 435-450.
- 1893 Patton, J.R., Goldfinger, C., Morey, A.E., Romsos, C., Black, B., Djadjadihardja, Y., and
1894 Udrek., 2013, Seismoturbidite record as preserved at core sites at the Cascadia and
1895 Sumatra-Andaman subduction zones, *Natural Hazards and Earth System Science*: v. 13,
1896 p. 833-867.
- 1897 Paull, C., Greene, H., Ussler, W., and Mitts, P., 2002, Pesticides as tracers of sediment transport
1898 through Monterey Canyon: *Geo-Marine Letters*, 22, pp. 121-126.
- 1899 Petter, A.L., Steel, R.J., Mohrig, D., Kim, W., and Carvajal, C., 2013, Estimation of the paleoflux
1900 of terrestrial-derived solids across ancient basin margins using the stratigraphic record:
1901 *Geological Society of America Bulletin*, v. 125, p. 578-593.

- 1902 Phillips, J.D., 1991, Fluvial sediment delivery to a Coastal Plain estuary in the Atlantic Drainage
1903 of the United States, *Mar. Geol.*: v. 98, p. 121-134.
- 1904 Phillips, J.D., 2003, Sources of nonlinearity and complexity in geomorphic systems: Progress in
1905 Physical Geography, v. 27, p. 1-23.
- 1906 Phillips, J.D. and Slattery, M.C., 2006, Sediment storage, sea level, and sediment delivery to the
1907 ocean by coastal plain rivers: Progress in Physical Geography, v. 30, p. 513-530.
- 1908 Piper, D.W. and Aksu, A., 1987, The source and origin of the 1929 grand banks turbidity current
1909 inferred from sediment budgets, *Geo-Mar. Lett.*: v. 7, p. 177-182.
- 1910 Posamentier, H.W. and Vail, P.R., 1988, Eustatic controls on clastic deposition II9d\sequence and
1911 systems tract models, *Sea-Level Changes: An Integrated Approach*: p. 125-154.
- 1912 Posamentier, H.W., Erskine, R.D., and Mitchum, R.M, Jr., 1991, Submarine fan deposition
1913 within a sequence stratigraphic framework: *Seismic Facies and Sedimentary Processes of*
1914 *Submarine Fans and Turbidite Systems*: Springer-Verlag, p. 127-136..
- 1915 Pirmez, C., Prather, B., Mallarino, G., O'Hayer, W., Droxler, A., and Winker, C., 2012,
1916 Chronostratigraphy of the Brazos-Trinity depositional system, western Gulf of Mexico:
1917 Implications for deepwater depositional models: Application of the Principles of Seismic
1918 Geomorphology to Continental-slope and Base-of-Slope Systems: Case Studies from
1919 Seafloor and Near Seafloor Analogues, *SEPM Special Publication 99*, p. 111-143.
- 1920 Prather, B., Pirmez, C., and Winker, C., 2012, Stratigraphy of linked intraslope basins: Brazos-
1921 Trinity system, western Gulf of Mexico: Application of the Principles of Seismic
1922 Geomorphology to Continental-slope and Base-of-Slope Systems: Case Studies from
1923 Seafloor and Near Seafloor Analogues, *SEPM Special Publication 99*, p. 83-109.
- 1924 Puig, P., Ogston, A.S., Mullenbach, B.L., Nittrouer, C.A., and Sternberg, R.W., 2003, Shelf-to-
1925 canyon sediment-transport processes on the Eel continental margin (northern California),
1926 *Mar. Geol.*: v. 193, p. 129-149.
- 1927 Rahl, J.M., Reiners, P.W., Campbell, I.H., Nicolescu, S., and Allen, C.M., 2003, Combined
1928 single-grain (U-Th)/He and U/Pb dating of detrital zircons from the Navajo Sandstone,
1929 Utah: *Geology*, v. 31, p. 761-764.
- 1930 Rahl, J.M., Ehlers, T.A., and van der Pluijm, B.A., 2007, Quantifying transient erosion of orogens
1931 with detrital thermochronology from syntectonic basin deposits: *Earth and Planetary*
1932 *Science Letters*, v. 256, p. 147-161.
- 1933 Reed, D.J., Commagere, A.M., and Hester, M.W., 2009, Marsh Elevation Response to Hurricanes
1934 Katrina and Rita and the Effect of Altered Nutrient Regimes, *J. Coast. Res.*, p. 166-173.
- 1935 Reimer, P.J., 2012, Refining the radiocarbon time scale: *Science*, v. 338, p. 337-338.
- 1936 Reiners, P.W. and Brandon, M.T., 2006, Using thermochronology to understand orogenic
1937 erosion: *Annual Reviews of Earth and Planetary Science*, v. 34, p. 419-466.
- 1938 Rodier, J. and Roche, M., 1984, World catalogue of maximum observed floods: *IAHS*
1939 *Publication no. 143*.
- 1940 Roering, J.J., Perron, J.T., and Kirchner, J.W., 2007, Functional relationships between denudation
1941 and hillslope form and relief, *Earth Planet. Sci. Lett.*: v. 264, p. 245-258.
- 1942 Rogers, K.G. and Goodbred Jr., S.L., 2010, Mass failures associated with the passage of a large
1943 tropical cyclone over the Swath of No Ground submarine canyon (Bay of Bengal),
1944 *Geology*: v. 38, p. 1051-1054.
- 1945 Romans, B.W., Normark, W.R., McGann, M.M., Covault, J.A., and Graham, S.A., 2009, Coarse-
1946 grained sediment delivery and distribution in the Holocene Santa Monica Basin,
1947 California: Implications for evaluating source-to-sink flux at millennial timescales:
1948 *Geological Society of America Bulletin*, v. 121, p. 1394-1408.
- 1949 Romans, B.W., Fildani, A., Graham, S.A., Hubbard, S.M., and Covault, J.A., 2010, Importance of
1950 predecessor basin history on sedimentary fill of a retroarc foreland basin: Provenance
1951 analysis of the Cretaceous Magallanes basin, Chile (50-52S): *Basin Research*, v. 22, p.
1952 640-658.

- 1953 Romans, B.W. and Graham, S.A., 2013, A deep-time perspective of land-ocean linkages in the
1954 sedimentary record: *Annual Reviews of Marine Science*, v. 5, p. 69-94.
- 1955 Rose, L.E. and Kuehl, S.A., 2010, Recent sedimentation patterns and facies distribution on the
1956 Poverty Shelf, New Zealand, *Mar. Geol.*: v. 270, p. 160-174.
- 1957 Sadler, P.M., 1981, Sediment accumulation rates and the completeness of stratigraphic sections:
1958 *The Journal of Geology*, v. 89, p. 569-584.
- 1959 Sadler, P.M. and Jerolmack, D.J., 2014, Scaling laws for aggradation, denudation, and
1960 progradation rates: The case for time-scale invariance at sediment sources and sinks:
1961 *Geological Society London Special Publication* 404.
- 1962 Saylor, J.E., Stockli, D.F., Horton, B.K., Nie, J., and Mora, A., 2012, Discriminating rapid
1963 exhumation from syndepositional volcanism using detrital zircon double dating:
1964 Implications for the tectonic history of the Eastern Cordillera, Colombia: *Geological*
1965 *Society of America Bulletin*, v. 124, p. 762-779.
- 1966 Schaller, M., von Blanckenburg, F., Hovius, N., and Kubik, P., 2001, Large-scale erosion rates
1967 from in situ-produced cosmogenic nuclides in European river sediments: *Earth and*
1968 *Planetary Science Letters*, v. 188, p. 441-458.
- 1969 Schumm, 1977, *The Fluvial System*, New York, John Wiley & Sons, 338 p.
- 1970 Schumm, S., 1993, River response to baselevel change: Implications for sequence stratigraphy:
1971 *The Journal of Geology*, v. 101, p. 279-294.
- 1972 Schillereff, D.N., Chiverrell, R.C., Macdonald, N., and Hooke, J.M., 2014, Flood stratigraphies in
1973 lake sediments: A review, *Earth-Sci. Rev.*, v. 135, p. 17-37.
- 1974 Shearman, P., Bryan, J., and Walsh, J.P., 2013, Trends in Deltaic Change over Three Decades in
1975 the Asia-Pacific Region, *J. Coast. Res.*: p. 1169-1183.
- 1976 Shen, Z., Tornqvist, T.E., Autin, W.J., Mateo, Z.R.P., Straub, K.M., and Mauz, B., 2012, Rapid
1977 and widespread response of the Lower Mississippi River to eustatic forcing during last
1978 glacial-interglacial cycle: *Geological Society of America Bulletin*, v. 124, p. 690-704.
- 1979 Simpson, G., Castelltort, S., 2012, Model shows that rivers transmit high-frequency climate
1980 cycles to the sedimentary record: *Geology*, v. 40, p. 1131-1134. doi:10.1130/G33451.1
- 1981 Smith, S.E. and Abdel-Kader, A., 1988, Coastal Erosion along the Egyptian Delta, *J. Coast. Res.*:
1982 v. 4, p. 245-255.
- 1983 Smith, D.P., Ruiz, G., Kvitek, R., and Iampietro, P.J., 2005, Semiannual patterns of erosion and
1984 deposition in upper Monterey Canyon from serial multibeam bathymetry: *GSA Bulletin* ,
1985 v. 117, 1123-1133; doi: 10.1130/B25510.1
- 1986 Stevens, T., Paull, C.K., Ussler, W., III, McGann, M., Buylaert, J-P., and Lundsten, E., 2014, The
1987 timing of sediment transport down the Monterey Canyon. *Geological Society of America*
1988 *Bulletin*, v. 126, p. 103-121
- 1989 Straub, K.M., Paola, C., Mohrig, D., Wolinsky, M.A., and George, T., 2009, Compensational
1990 stacking of channelized sedimentary deposits: *Journal of Sedimentary Research*, v. 79, p.
1991 673-688.
- 1992 Strong, N., Sheets, B., Hickson, T., and Paola, C., 2005, A mass-balance framework for
1993 quantifying downstream changes in fluvial architecture: *Fluvial Sedimentology VII*,
1994 *Special Publication* 35, p. 243-253.
- 1995 Sømme, T.O., Helland-Hansen, W., Martinsen, O.J., and Thurmond, J.B., 2009, Relationships
1996 between morphological and sedimentological parameters in source-to-sink systems: A
1997 basis for predicting semi-quantitative characteristics in subsurface systems: *Basin*
1998 *Research*, v. 21, p. 361-387.
- 1999 Sommerfield, C.K. and Nittrouer, C.A., 1999, Modern accumulation rates and a sediment budget
2000 for the Eel shelf: A flood-dominated depositional environment: *Marine Geology*, 154,
2001 227-241.

- 2002 Sommerfield, C.K., Nittrouer, C.A., and Alexander, C.R., 1999, ^7Be as a tracer of flood
2003 sedimentation on the northern Californian continental margin: *Continental Shelf*
2004 *Research*, 19, 335-361.
- 2005 Sommerfield, C.K. and Wheatcroft, R.A., 2007, Late Holocene sediment accumulation on the
2006 northern California shelf: Oceanic, fluvial, and anthropogenic influences, *Bull. Geol. Soc.*
2007 *Am.*: v. 119, p. 1120-1134.
- 2008 Sommerfield, C.K. and Nittrouer, C.A., 2014, Comment on “Eel River margin source-to-sink
2009 sediment budgets: Revisited” by J.A. Warrick [*Marine Geology* 351 (2014) 25–37], *Mar.*
2010 *Geol.*
- 2011 Sorrel, P., Debret, M., Billeaud, I., Jaccard, S.L., McManus, J.F., and Tessier, B., 2012, Persistent
2012 non-solar forcing of Holocene storm dynamics in coastal sedimentary archives, *Nature*
2013 *Geoscience*: v. 5, p. 892-896.
- 2014 Soutar, A. and Crill, P.A., 1977, Sedimentation and climatic patterns in the Santa Barbara Basin
2015 during the 19th and 20th centuries., *Geol. Soc. Am. Bull.*, 88, 1161-1172.
- 2016 Strasser, M., Anselmetti, F.S., Fah, D., Giardini, D., and Schnellmann, M., 2006, Magnitudes and
2017 source areas of large prehistoric northern Alpine earthquakes revealed by slope failures in
2018 lakes, *Geology*: v. 34, p. 1005-1008.
- 2019 Suckow, A., Morgenstern, U., and Kudrass, H.R., 2001, Absolute dating of recent sediments in
2020 the cyclone-influenced shelf area off Bangladesh: comparison of gamma spectrometric
2021 (^{137}Cs , ^{210}Pb , ^{228}Ra), radiocarbon, and ^{32}Si ages, *Radiocarbon*, v. 43, 917- 927.
- 2022 Sumner, E.J., Siti, M.I., McNeill, L.C., Talling, P.J., Henstock, T.J., Wynn, R.B., Djajadihardja,
2023 Y.S., and Permana, H., 2013, Can turbidites be used to reconstruct a paleoearthquake
2024 record for the central Sumatran margin? *Geology*: v. 41, p. 763-766.
- 2025 Szczucinski, W., Kokocinski, M., Rzeszewski, M., Chague-Goff, C., Cachao, M., Goto, K., and
2026 Sugawara, D., 2012, Sediment sources and sedimentation processes of 2011 Tohoku-oki
2027 tsunami deposits on the Sendai Plain, Japan - Insights from diatoms, nannoliths and grain
2028 size distribution, *Sediment. Geol.*: v. 282, p. 40-56
- 2029 Syvitski, J.P., 2003, Supply and flux of sediment along hydrological pathways: Research for the
2030 21st Century: *Global and Planetary Change*, v. 39, p. 1-11.
- 2031 Syvitski, J.P., 2008, Predictive modeling in sediment transport and stratigraphy: *Computers and*
2032 *Geoscience*, 10.1016/j.cageo.2008.02.001.
- 2033 Syvitski, J.P.M., Vorosmarty, C.J., Kettner, A.J., and Green, P., 2005, Impact of humans on the
2034 flux of terrestrial sediment to the global coastal ocean, *Science*, v. 308, p. 376-380.
- 2035 Syvitski, J.P.M. and Milliman, J.D., 2007, Geology, geography, and humans battle for dominance
2036 over the delivery of fluvial sediment to the coastal ocean, *J. Geol.*, v. 115, p. 1-19.
- 2037 Syvitski, J.P.M. and Saito, Y., 2007, Morphodynamics of deltas under the influence of humans,
2038 *Global Planet. Change*, v. 57, p. 261-282.
- 2039 Talling, P.J., Paull, C.K., and Piper, D.J.W., 2013, How are subaqueous sediment density flows
2040 triggered, what is their internal structure and how does it evolve? Direct observations
2041 from monitoring of active flows, *Earth-Sci. Rev.*: v. 125, p. 244-287.
- 2042 Tesi, T., Langone, L., Goñi, M.A., Wheatcroft, R.A., Miserocchi, S., and Bertotti, L., 2012, Early
2043 diagenesis of recently deposited organic matter: A 9-yr time-series study of a flood
2044 deposit, *Geochim. Cosmochim. Acta*: v. 83, p. 19-36.
- 2045 Traykovski, P., Geyer, W.R., Irish, J.D., and Lynch, J.F., 2000, The role of wave-induced
2046 density-driven fluid mud flows for cross-shelf transport on the Eel River continental
2047 shelf, *Cont. Shelf Res.*: v. 20, p. 2113-2140.
- 2048 Traykovski, P., Wiberg, P.L., and Geyer, W.R., 2007, Observations and modeling of wave-
2049 supported sediment gravity flows on the Po prodelta and comparison to prior
2050 observations from the Eel shelf, *Cont. Shelf Res.*: v. 27, p. 375-399.
- 2051 Turner, R.E., Baustian, J.J., Swenson, E.M., and Spicer, J.S., 2006, Wetland sedimentation from
2052 hurricanes Katrina and Rita, *Science*, v. 314, p. 449-452.

2053 Van den Berg van Saparoea, A.-P. and Postma, G., 2008, Control of climate change on the yield
2054 of river systems: Recent Advances in Models of Siliciclastic Shallow-Marine
2055 Stratigraphy: SEPM Special Publication 90, p. 15-33.

2056 Viles, H.A. and Goudie, A.S., 2003, Interannual, decadal and multidecadal scale climatic
2057 variability and geomorphology, *Earth-Sci. Rev.*, v. 61, p. 105-131.

2058 Voller, V.R., Ganti, V., Paola, C., and Foufoula-Georgiou, E., 2012, Does the flow of information
2059 in a landscape have direction?: *Geophysical Research Letters*, 39.

2060 Von Blanckenburg, F., 2005, The control mechanisms of erosion and weathering at basin scale
2061 from cosmogenic nuclides in river sediment: *Earth and Planetary Science Letters*, v. 237,
2062 p. 462-479.

2063 Walling, D.E. and Webb, B.W., 1996, Erosion and sediment yield: a global overview, IAHS
2064 Publication: v. 236, p. 3-19.

2065 Walling, D.E., 1999, Linking land use, erosion and sediment yields in river basins,
2066 *Hydrobiologia*: v. 410, p. 223-240.

2067 Walling, D.E. and Collins, A.L., 2008, The catchment sediment budget as a management tool,
2068 *Environ. Sci. & Policy*: v. 11, p. 136-143.

2069 Walling, D.E., 2013, The evolution of sediment source fingerprinting investigations in fluvial
2070 systems, *Journal of Soils and Sediments*: v. 13, p. 1658-1675.

2071 Walker, M., 2005, Quaternary dating methods: John Wiley & Sons, 286 p.

2072 Walsh, J.P. and Nittrouer, C.A., 1999, Observations of sediment flux to the Eel continental slope,
2073 northern California, *Mar. Geol.*: v. 154, p. 55-68.

2074 Walsh, J.P. and Nittrouer, C.A., 2003, Contrasting styles of off-shelf sediment accumulation in
2075 New Guinea, *Mar. Geol.*: v. 196, p. 105-125.

2076 Walsh, J.P., Nittrouer, C.A., Palinkas, C.M., Ogston, A.S., Sternberg, R.W., and Brunskill, G.J.,
2077 2004, Clinoform mechanics in the Gulf of Papua, New Guinea, *Cont. Shelf Res.*: v. 24, p.
2078 2487-2510.

2079 Walsh, J.P., Corbett, R., Mallinson, D., Goni, M., Dail, M., Loewy, C., Marciniak, K., Ryan, K.,
2080 Smith, C., Stevens, A., Sumners, B., and Tesi, T., 2006, Mississippi delta mudflow
2081 activity and 2005 Gulf hurricanes, *EOS Trans. Am. Geophys. Union*: v. 87, p. 477-479.

2082 Walsh, J.P., Corbett, D.R., Ogston, A.S., Nittrouer, C., Kuehl, S.A., Allison, M.A., and
2083 Goodbred, S.L. Jr., 2013, Shelf and slope sedimentation associated with large deltaic
2084 systems: In: *Biogeochemical Dynamics at Major River-Coastal Interfaces: Linkages with*
2085 *Global Change*, editors: T.S. Bianchi, M.A. Allison, and W.Cai. Cambridge University
2086 Press, New York, NY, p. 86-117.

2087 Walsh, J.P., Corbett, D.R., Kiker, J.M., Orpin, A.R., Hale, R.P., and Ogston, A.S., 2014, Spatial
2088 and temporal variability in sediment deposition and seabed character on the Waipaoa
2089 River margin, New Zealand, *Cont. Shelf Res.*: v. 86, p. 85-102.

2090 Wang, Y., Straub, K.M., and Hajek, E.A., 2011, Scale-dependent compensational stacking: An
2091 estimate of autogenic time scales in channelized sedimentary deposits: *Geology*, v. 39, p.
2092 811-814.

2093 Warrick, J.A., 2014, Eel River margin source-to-sink sediment budgets: Revisited, *Mar. Geol.*: v.
2094 351, p. 25-37.

2095 Weislogel, A.L., Graham, S.A., Chang, E.Z., Wooden, J.L., Gehrels, G.E., and Yang, H., 2006,
2096 Detrital zircon provenance of the Late Triassic Songpan-Ganzi complex: Sedimentary
2097 record of collision of the North and South China Blocks: *Geology*, v. 34, p. 97-100.

2098 Weber, M.E., Wiedicke, M.H., Kudrass, H.R., Hubscher, C., and Erlenkeuser, H., 1997, Active
2099 growth of the Bengal Fan during sea-level rise and highstand: *Geology*, v. 25, p. 315-318.

2100 Wheatcroft, R.A., Sommerfield, C.K., Drake, D.E., Borgeld, J.C., and Nittrouer, C.A., 1997,
2101 Rapid and widespread dispersal of flood sediment on the northern California margin,
2102 *Geology*: v. 25, p. 163-166.

- Wheatcroft, R.A. and Borgeld, J.C., 2000, Oceanic flood deposits on the northern California shelf: Large-scale distribution and small-scale physical properties, *Cont. Shelf Res.*: v. 20, p. 2163-2190.
- Wheatcroft, R.A., Wiberg, P.L., Alexander, C.R., S.J. Bentley, D.E. Drake, C.K. Harris and A.S. Ogston, 2007, in: Nittrouer, C.A., Austin, J.A., Field, M.E., Kravitz, J.H., Syvitski, J.P.M., and P.L. Wiberg, eds., *Continental margin sedimentation: from sediment transport to sequence stratigraphy*. Malden, MA: Blackwell Pub. for the International Association of Sedimentologists. p. 101-155.
- Whipple, K. X., 2001, Fluvial landscape response time: how plausible is steady-state denudation?: *American Journal of Science*, v. 301, p. 313-325.
- Whittaker, A.C., Cowie, P.A., Attal, M., Tucker, G.E., and Roberts, G.P., 2007, Bedrock channel adjustment to tectonic forcing: Implications for predicting river incision rates: *Geology*, v. 35, p. 103-106.
- Whittaker, A.C., Duller, R.A., Springett, J., Smithells, R.A., Whitchurch, A.L., and Allen, P.A., 2011, Decoding downstream trends in stratigraphic grain size as a function of tectonic subsidence and sediment supply: *Geological Society of America Bulletin*, v. 123, p. 1363-1382.
- Willett, S.D. and Brandon, M.T., 2002, On steady state in mountain belts: *Geology*, v. 30, p 175-178.
- Wilkinson, B.H. and McElroy, B.J., 2007, The impact of humans on continental erosion and sedimentation, *Bull. Geol. Soc. Am.*: v. 119, p. 140-156.
- Wilson, J.T., 1966, Did the Atlantic close and then re-open?: *Nature*, v. 211, p. 676-681.
- Wittmann, H. and Von Blanckenburg, F., 2009, Cosmogenic nuclide budgeting of floodplain sediment transfer: *Geomorphology*, v. 109, p. 246-256.
- Wittmann, H., Von Blanckenburg, F., Maurice, L., Guyot, J.-L., and Kubik, P., 2011, Recycling of Amazon floodplain sediment quantified by cosmogenic ²⁶Al and ¹⁰Be: *Geology*, v. 39, p. 467-470.
- Xu, J.P., Sequeiros, O.E., and Noble, M.A., 2014a, Sediment concentrations, flow conditions, and downstream evolution of two turbidity currents, Monterey Canyon, USA, *Deep-Sea Research Part I: Oceanographic Research Papers*: v. 89, p. 11-34.
- Xu, J.P., Wong, F.L., Kvitek, R., Smith, D.P., and Paull, C.K., 2008, Sandwave migration in Monterey Submarine Canyon, Central California, *Mar. Geol.*: v. 248, p. 193-212.
- Xu, J.P., Noble, M.A., and Rosenfeld, L.K., 2004, In-situ measurements of velocity structure within turbidity currents, *Geophys. Res. Lett.*: v. 31, p. L09311 1-4.
- Xu, K., Corbett, D.R., Walsh, J.P., Young, D., Briggs, K.B., Cartwright, G.M., Friedrichs, C.T., Harris, C.K., Mickey, R.C., and Mitra, S., 2014b, Seabed erodibility variations on the Louisiana continental shelf before and after the 2011 Mississippi River flood, *Estuar. Coast. Shelf Sci.*, v. 149, p. 283-293.
- Zachos, J.C., Dickens, G.R., and Zeebe, R.E., 2008, An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics: *Nature*, v. 451, p 279-283.
- Zalasiewicz, J., Williams, M., Smith, A., T. L. Barry, A. L. Coe, P. R. Bown, P. Brenchley, D. Cantrill, A. Gale, P. Gibbard, F. J. Gregory, M. W. Hounslow, A. C. Kerr, P. Pearson, R. Knox, J. Powell, C. Waters, J. Marshall, M. Oates, P. Rawson, and P. Stone. 2008. Are we now living in the Anthropocene? *GSA Today* 18:4-8.

Figure 1

Figure 2

Above is an x-ray image of core. Image intensity reflects the bulk density, where light colors represent low-bulk density sediment from a recently deposited "oceanic flood" layer (Sommerfield and Nittrouer, 1999; Wheatcroft and Borgeld, 2000).

Figure 4

Variation in mean grain size from upper canyon core (location denoted by red star) compared to storm record (Michels et al., 2003). Decadal storm-controlled reworking of shelf depocenter is superimposed on sediment supply signal from catchment.

Figure 6

Figure 7

COLOR EXPLANATION

Romans et al. -- Figure 7

Figure 9

Romans et al. -- Figure 9

Figure 10