The application of diatoms to reconstruct the history of subduction zone earthquakes and tsunamis

Tina Dura^{1, 2 *}

Eileen Hemphill-Haley³

Yuki Sawai⁴

Benjamin P. Horton^{1, 2, 5}

¹ *Sea Level Research, Department of Marine and Coastal Science, Rutgers University, New Brunswick, NJ, USA*

² *Institute of Earth, Ocean, and Atmospheric Sciences, Rutgers University, New Brunswick, NJ, USA*

³ *Department of Geology, Humboldt State University, Arcata, CA, USA*

⁴ *Geological Survey of Japan, National Institute of Advances Industrial Science and Technology (AIST), Tsukuba Central 7, 1-1-1 Higashi, Tsukuba, Ibaraki 305-8567, Japan*

⁵ *Earth Observatory of Singapore and Asian School of the Environment, Nanyang Technological University, Singapore*

*Corresponding Author: Tina Dura
Department of Marine and Coastal Science
Rutgers University
71 Dudley Road, New Brunswick, NJ 08901
Tel: +1 530.990.3468
Email: dura@marine.rutgers.edu

Keywords: Diatoms; earthquakes; tsunamis; subduction zones; sea level

Abstract

Earthquake and tsunami records on centennial and millennial temporal scales are necessary to understanding long-term subduction zone behavior and the occurrences of large, but infrequent events. Microfossils, such as diatoms, incorporated into coastal stratigraphy provide some of the most detailed reconstructions of the history of earthquakes and tsunamis. We explore qualitative and quantitative techniques that employ the relation between diatoms and salinity, tidal elevation, and life form to: (1) reconstruct records of vertical land-level change associated with large earthquakes; and (2) identify anomalous sand and silt beds deposited by tsunamis. A global database shows that diatoms have been successfully employed in the reconstruction of earthquake and tsunami histories in Chile, the Indian Ocean, Japan, New Zealand, the North Sea, the Pacific Northwest of North America, and the South Pacific. We use case studies from some of these locations to highlight advancements in the field and new capabilities that diatoms have enabled. Examples from the Pacific Northwest of North America illustrate the evolution of quantitative diatom-based reconstructions of earthquake-related land-level change. In Alaska and Japan, diatoms have documented land-level changes throughout the earthquake deformation cycle, including possible preseismic land-level change signals and postseismic deformation. Diatoms helped identify coseismic uplift along the central Chile subduction zone coast, and uplift and subsidence along the Alaska-Aleutian megathrust, expanding our knowledge of the variability of slip in megathrust ruptures. In tsunami studies, diatoms help determine the provenance of anomalous sands and silts found in low-energy coastal stratigraphic sequences. In Japan, allochthonous marine and brackish diatoms within sand deposits signaled repeated marine incursions into a coastal lake, helping identify a possible predecessor to the 2011 Tohoku tsunami. In the Pacific Northwest of North America and Chile, diatoms were used to estimate tsunami run-up beyond the landward limit of tsunami sedimentation. Examples from the North Sea, Thailand, and Japan show how the fragmentation and sorting of diatom valves may provide evidence of high-energy transport during the rapid, turbulent flow of a tsunami. To conclude, we emphasize the importance of studying the modern diatom response to changes in land level and/or tsunami inundation to improve diatom-based records of prehistoric earthquakes and tsunamis.

1. Introduction

An incomplete understanding of the earthquake and tsunami hazards associated with the Sunda and Japan subduction zones contributed to the devastating societal impacts of the 2004 Indian Ocean and 2011 Tohoku events (Rhodes et al., 2006; Geller, 2011; Stein and Okal, 2011; Heki, 2011). Instrumental records of previous earthquakes and tsunamis proved too short to estimate the potential magnitude and recurrence interval of such great events that recur centuries to millennia apart (Stein and Okal, 2007). With more than a third of the world's coastlines lying adjacent to active plate boundaries (Lajoie, 1986; Stewart and Vita-Finzi, 1998), we must extend earthquake and tsunami histories to adequately assess hazards from subduction zones (Small et al., 2000; Satake and Atwater, 2007; Stein and Okal, 2011).

Stratigraphic evidence of subduction zone earthquakes and tsunamis has been used to extend records over centuries to millennia. Stratigraphic evidence was first described from the coastal wetlands of Alaska (Plafker, 1965; Plafker, 1972; Bartsch-Winkler and Schmoll, 1987; Combellick, 1991, 1994; Combellick and Reger, 1994), the Pacific Northwest of North America (Atwater, 1987; Darienzo and Peterson, 1990; Atwater 1992; Nelson, 1992; Darienzo et al., 1994; Nelson et al., 1996a), Chile (Wright and Mella, 1963; Bourgeois and Reinhart, 1989; Atwater et al., 1992) and Japan (Kon'no, 1961; Minoura and Nakaya, 1991; Sawai et al., 2002; Nanayama et al., 2003; Sawai et al., 2012, 2015). Subsidence or uplift associated with earthquakes is recorded as a series of sharp stratigraphic contacts that reflect sudden changes in land level (Nelson et al., 1996b; Yeats et al., 1997). Widespread sand beds, rapidly deposited by tsunamis accompanying earthquakes, are often found concomitant with stratigraphic evidence of

sudden land-level change (e.g., Hemphill-Haley et al., 1995a; Cisternas et al., 2005). In addition, trans-oceanic tsunamis produced by the largest subduction zone earthquakes can deposit sediments on coastlines hundreds to thousands of kilometers away from the earthquake source (e.g., Satake et al., 1996; Goff et al., 2006).

Microfossils such as diatoms, foraminifera, and pollen that are incorporated into coastal stratigraphy provide qualitative and quantitative estimates of land-level change associated with subduction zone earthquakes (Horton et al., 2013; Fig. 1). In particular, diatoms and foraminifera have been applied to earthquake and tsunami studies around the world (e.g., Darienzo et al., 1994; Hemphill-Haley, 1995a, 1995b, 1996; Sawai et al., 2004a, b; Shennan and Hamilton, 2006; Hawkes et al., 2011; Engelhart et al., 2013; Grand Pre et al., 2012; Clark et al., 2015; Dura et al., 2015). These microfossils have improved upon early coastal studies of subduction-zone earthquake history that estimated the amount of coseismic land-level change with large errors (>1 m) based on the broad elevational ranges of plant macrofossils (Atwater, 1987, 1992; Clague and Bobrowsky, 1994). Quantitative diatom and foraminiferal-based reconstructions of land-level change (e.g., transfer functions) can produce substantially smaller (<0.3 m) errors (Guilbault et al., 1995, 1996; Zong et al., 2003; Sawai et al., 2004a,b; Shennan and Hamilton, 2006; Nelson et al., 2008; Hawkes et al., 2011; Engelhart et al., 2013; Watcham et al., 2013).

Microfossils are applied to tsunami studies to determine the provenance of tsunami sediments deposited in low-energy coastal environments such as lowlands (Hemphill-Haley, 1995a, 1996; Benson et al., 1997; Nanayama et al. 2007; Sugawara et al., 2009), interdunal depressions (Dawson and Stewart, 2007; Jankaew et al., 2008), lagoons (Minoura and Nakaya, 1991; Nichol et al., 2007; Sawai et al., 2009b; Wilson et al., 2014)

and lakes (Hutchinson et al., 1997; Grauert et al., 2001; Kelsey et al., 2005). Allochthonous marine assemblages or a chaotic mix of marine, brackish and freshwater taxa within terrestrial or brackish settings may be indicative of marine incursions from tsunamis (Tuttle et al., 2004; Cochran et al., 2005; Hawkes et al., 2007; Goff et al., 2012; Clark et al., 2015), and the preservation and distribution of diatoms and foraminifera throughout the deposit can be used to infer high-energy transport and provenance (Hemphill-Haley, 1996; Sawai et al., 2002; Pilarczyk et al., 2012a). Tsunamis of the past two decades have provided an opportunity to characterize the microfossil composition of modern deposits (Dawson, 2007; Sawai et al., 2009a; Chagué-Goff et al., 2011; Szczuciński et al., 2012b), improving our ability to identify such characteristics in the fossil record.

In this review we explore the application of diatoms to earthquake and tsunami studies (Fig. 2; Table 1). We explain the statistical approaches that use diatoms to estimate earthquake related land-level changes, and assess the utility of diatoms in identifying tsunamis in the stratigraphic record. We also examine the knowledge gaps and limitations of diatom analyses in earthquake and tsunami studies, and make recommendations for future research.

2. Reconstructing coastal environments with diatoms

Diatoms are photosynthetic, unicellular algae that inhabit freshwater, brackish, and marine environments (Round et al., 1990; Jones, 2007). Diatoms are a dominant microphyte in coastal wetland environments and their siliceous valves, ranging in size from ~5 μm to ~200 μm , are resistant to taphonomic degradation (e.g., dissolution, abrasion, bioerosion, oxidation, transport, predation; Admiraal, 1984; Palmer and Abbott,

1986; Cooper et al., 2010). As a result, small (1 g wet weight) sample sizes (for example, from narrow-diameter cores 25-50 mm) contain statistically significant (300-600 valves) diatom populations (Birks, 1995; Battarbee et al., 2001; Nelson, 2015).

Diatoms have been described and classified for over 200 years based on their shapes, sizes, and the intricate morphological characteristics of their siliceous valves (Round et al., 1990). Diatom taxa are divided into three main classes: the Coscinodiscophyceae (centric taxa); Fragilariophyceae (araphid pennate taxa); and Bacillariophyceae (raphid pennate taxa). Literature on the identification of diatoms to species level includes van der Werff and Huls (1958-1974), Patrick and Reimer (1966, 1975), Krammer and Lange-Bertalot (1986, 1988, 1991a, b), Hartley et al. (1996), Krammer (2000, 2002, 2003), Lange-Bertalot (2000, 2001, 2011), and Levkov (2009).

Diatoms are a valuable tool in reconstructing paleoenvironmental changes because of their sensitivity to environmental factors including salinity, tidal exposure, substrate, vegetation, pH, nutrient supply, and temperature found in specific coastal wetland environments (e.g., Zong and Horton, 1998). Over time, diatoms become incorporated in coastal sediments, resulting in buried assemblages that represent an environmental history that can span thousands of years. Literature for the ecological classification of diatoms includes Hustedt (1937, 1939, 1953, 1957), Lowe (1974), Patrick and Williams (1990), Denys (1991-1992), Juggins (1992), Vos and de Wolf (1988; 1993), Van Dam et al. (1994), and Lange-Bertalot (2000).

Diatoms' preferences for salinity are valuable for earthquake and tsunami studies. Changes in salinity across the intertidal zone produce vertically zoned diatom assemblages with respect to the tidal frame (Nelson and Kashima, 1993; Hemphill-Haley

1995b; Sherrod et al., 1999; Patterson et al., 2005; Horton and Sawai, 2010; Woodroffe and Long, 2010). Establishing the distribution of diatoms across the intertidal zone is a prerequisite for reconstructing paleoenvironmental changes related to earthquakes and tsunamis (e.g., Zong and Horton, 1998). In the halobian classification scheme, polyhalobous and mesohalobous diatom taxa represent the marine and brackish conditions found in tidal flats and lower tidal marshes and mangroves. Oligohalobous-halophile and oligohalobous-indifferent taxa become dominant through the transition from tidal marsh/mangrove to freshwater environments, and halophobous taxa characterize the most landward freshwater communities above the highest tides. The presence of diatoms in freshwater environments is an obvious advantage compared to foraminifera, which are restricted to areas of marine influence (e.g., Murray, 1991).

The distribution of diatoms along a modern intertidal transect (Fig. 3) typically shows a clear transition from subtidal, open-water marine diatoms (e.g., *Thalassiosira pacifica* and *Odontella aurita*), to marine tidal flat taxa (e.g., *Achnanthes brevipes* and *Tabularia fasciculata*). In the low marsh, where a mixed diatom community is often found, marine-brackish diatoms (e.g., *Planothidium delicatulum* and *Tryblionella granulata*) dominate, followed by freshwater taxa that can tolerate low salinities in the high marsh (e.g., *Caloneis bacillum* and *Cosmioneis pusilla*) and freshwater salt-intolerant taxa in the upland environment (e.g., *Eunotia bilunaris* and *Aulacoseira crassipunctata*).

The distribution of diatoms across the intertidal zone based on their life form is valuable for tsunami studies. The diverse life forms of diatom floras are employed to support the marine provenance of inferred tsunami deposits. Hustedt (1958), Vos (1986),

Vos et al. (1988), and Vos and de Wolf (1993) define diatom life forms based on the substrate—or lack thereof—that particular diatom taxa commonly live on. Benthic diatoms are grouped into epipsammic taxa that live attached to sand grains; epipellic taxa that live on or just below the surface of wet muddy sediment; epiphytic taxa that are attached to larger plants or other surfaces; and aerophile taxa that are able to survive subaerial, temporarily dry conditions. Planktonic diatoms float freely in the water column and do not live attached to any substrate; tycho planktonic diatoms include an array of species that live in the benthos, but are commonly found in the plankton. Based on local conditions, epipsammic, epipellic, planktonic, and tycho planktonic diatoms may comprise tidal flat populations, whereas epiphytic and epipellic forms are more common on tidal marshes/mangroves. Aerophilous forms are most common within the landward communities above the highest tides (Sherrod, 1999).

3. Application of diatoms to subduction zone paleoseismic cycles

3.1 Earthquake deformation cycle

On subduction zone coastlines, the land-level changes associated with earthquakes reflect the strain accumulation and release of the earthquake deformation cycle (Burbank and Anderson, 2001; Nelson, 2013). The nature of the interseismic and coseismic motion of the coastline is determined by its proximity to the trench, the geometry of the subduction zone, and where ruptures stop along strike (Plafker, 1965; Plafker and Savage, 1970; Wang et al., 2012). Some subduction zone coastlines (e.g., Cascadia, the Pacific Northwest of North America) lie within a zone that gradually uplifts in between (interseismic) and abruptly subsides during (coseismic) great earthquakes (Plafker, 1972, Atwater 1987). In coastal wetland stratigraphy, the interseismic period is represented by a

gradual regression from clastic (e.g., tidal flat or shallow subtidal) to organic-rich (e.g., marsh or swamp) sediments reflecting land-level uplift (i.e., relative sea-level fall) and a decrease in marine influence. Abrupt coseismic land subsidence creates a sudden change from organic-rich sediments to clastic sediments, reflecting land-level subsidence (i.e., relative sea-level rise) and an increase in marine influence (Fig. 1).

Conversely, some coastal locations bordering subduction zones (e.g., central Chile; Alaska-Aleutian megathrust) lie within a zone that gradually subsides in between and abruptly uplifts during great earthquakes (Shennan et al., 2009; Dura et al., 2015). During the interseismic period, coastal wetland stratigraphy displays a gradual transgression from organic-rich to clastic sediments as land-level subsides and marine influence increases. Abrupt coseismic uplift creates a sudden change from clastic to organic-rich sediments, reflecting land-level uplift and a decrease in marine influence.

Coastal wetland stratigraphy can record both coseismic uplift and subsidence at one location (e.g., Hayward et al., 2006; Briggs et al., 2014; Ely et al., 2014), depending on the distribution of slip upon the megathrust surface (Savage and Hastie, 1966) or the interaction with upper plate faults. The possibility of a variable uplift-subsidence history illustrates that the paleoseismic record should not be assumed to record exclusively coseismic uplift or subsidence.

Because non-seismic coastal processes can also produce changes in stratigraphy similar to those created by great subduction zone earthquakes, criteria must be considered to support a tectonic origin (Darienzo et al., 1994; Nelson et al., 1996b; Shennan et al., 1996; Dura et al., 2015). The key criteria are the lateral extent of sharp stratigraphic contacts; the suddenness and magnitude of land-level change; the synchronicity of land-

level change among regional sites; and the coincidence of tsunami deposits with sudden changes in stratigraphy. No breaks in sedimentation greater than a few weeks are assumed to occur following earthquake-related land-level change and continuous sedimentation is assumed in the interseismic period. Although many paleoseismic studies have relied on stratigraphic investigations to support an earthquake origin (Atwater, 1987; Dura et al., 2011), microfossils have been particularly useful in providing supporting evidence of earthquake-related land-level change.

3.2 Diatoms and land-level changes related to the earthquake deformation cycle

Much of the early research using diatoms to reconstruct land-level changes related to the earthquake deformation cycle focused on the Pacific Northwest of North America (Cascadia subduction zone). Regional sea-level rise at rates of up to 2 mm/yr along the central Cascadia subduction zone since 6,000 calibrated years BP (Engelhart et al., 2015) resulted in continuous records of tidal sedimentation that contain evidence of coseismic land-level change and tsunami inundation.

Dariento and Peterson (1990) and Dariento et al. (1994) employed qualitative diatom analyses to confirm inferences of sudden and widespread coseismic subsidence based on distinctive stratigraphic contacts in Cascadia coastal wetlands. Dariento et al. (1994) used diatoms to identify broad floral zones (e.g., high marsh, low marsh, tidal flat) in the modern environment and subsequently estimated the magnitude of coseismic subsidence by identifying analogous floral zones from above and below sharp stratigraphic contacts. Because the elevational range of the floral zones was large (0.5-1.0 m), errors for subsidence estimates commonly exceeded 1 m (e.g., Nelson and Kashima,

1993; Darienzo and Peterson, 1990; Darienzo et al., 1994; Kelsey et al., 2002; Witter et al., 2003).

Statistical methods employed the relationship between modern diatom assemblages of known elevation and fossil assemblages to reconstruct land-level changes in stratigraphic sections at Cascadia (Hemphill-Haley, 1995a; Nelson et al., 1996a; Shennan et al., 1996). In southern Washington State, USA, Hemphill-Haley (1995b) used factor analysis to identify the relation of modern diatom distributions to elevation. The result was a list of taxa with dominant occurrences in marsh zones (high marsh, low marsh, and high and low marshes) and three subenvironments of the lower intertidal to shallow subtidal zone (mud flats, sand flats, and *Zostera* (eelgrass) beds). The modern distributions of the same taxa found in the fossil record were used to reconstruct the abrupt changes in environment indicated by stratigraphic contacts spanning the AD 1700 subduction zone earthquake (Fig. 4). Distinct changes in diatom assemblages across these sharp contacts confirmed significant, widespread, and lasting change from an upland environment to a tidal-flat or low marsh environment, consistent with coseismic subsidence of 0.8-1.1 m. Shennan et al. (1996, 1998) and Nelson et al. (1996a) employed similar statistical techniques (i.e., ordination and discriminant function analysis, respectively) to quantitatively compare modern diatom assemblages of known elevation with fossil diatom assemblages. Although the statistical techniques of Hemphill-Haley (1995b), Shennan et al. (1996, 1998), and Nelson et al. (1996a) improved the analysis of diatom data, the reconstructions were still based on differences in elevational ranges between pre-earthquake and postearthquake paleoenvironments, resulting in 0.5-1.0 m errors.

Fully quantitative, microfossil-based transfer functions used to calculate coseismic land-level change at Cascadia have produced substantially smaller errors (<0.5 m; e.g., Guilbault et al., 1995, 1996). The transfer function uses multivariate statistical techniques to formalize the relationship between the relative abundance of microfossil species and the environmental variable of interest (in the case of paleoseismic studies, that variable is elevation; e.g., Horton et al., 1999). The transfer function is applied to microfossil assemblages in coastal wetland stratigraphic sequences to calculate changes in paleo-elevation and can provide a continuous record of land-level changes throughout the earthquake deformation cycle. The performance of transfer functions can be measured using multiple techniques to assess the effect of sample design, the goodness-of-fit between microfossil assemblages and elevation, and the statistical significance of each reconstruction (Telford and Birks, 2011a, b). The technique has since been expanded in Cascadia to include extensive modern diatom (Sherrod, 1999, 2000, 2001), foraminifera (Guilbault et al., 1995, 1996; Hawkes et al., 2010; Engelhart et al., 2013), and pollen (Hughes et al., 2002) datasets, and applied in other geographical locations such as Alaska (e.g., Shennan and Hamilton, 2006), Japan (e.g., Sawai, 2004b), and Chile (e.g., Garrett et al., 2014; Table 1 lists which diatom studies have used quantitative diatom techniques, Fig. 2).

Coastal wetland stratigraphy from eastern Hokkaido, Japan contains evidence of preseismic and postseismic land-level change (Atwater et al., 2004; Sawai et al., 2002, 2004a, b) associated with subduction zone earthquakes originating from the Kuril Trench (Nanayama et al., 2003). Sawai et al. (2004a) used a diatom-based transfer function (Sawai et al., 2001b, 2004b) to document the land-level change associated with a great

17th century earthquake and tsunami (Fig. 6). Fossil diatom assemblages show that tidal flats gradually changed into freshwater upland environments in the decades after the earthquake as a result of up to 1.5 m of postseismic uplift. Sawai et al. (2004a) suggested the large land-level change was the result of a subduction zone earthquake of unusually large size along the Kuril Trench.

Diatom data have helped identify coseismic uplift associated with subduction zone earthquakes in Alaska (e.g., Shennan et al., 2009, 2014) and Chile (e.g., Dura et al., 2015). Along the central Chile coast, Dura et al. (2015) used diatoms to identify six instances of coseismic uplift between 6200 and 3600 calibrated years BP. A repeated influx of freshwater diatoms and other siliceous microfossils above six tsunami sands suggested >1 m of coseismic uplift. The study documented a ~500 year recurrence interval for the earthquakes and accompanying tsunamis in central Chile.

Diatom studies have documented evidence of both coseismic subsidence and uplift at the same site. Briggs et al. (2014) used diatoms and foraminifera to identify five instances of sudden uplift and subsidence in the stratigraphy of a coastal marsh on Sitkinak Island bordering the Alaska-Aleutian megathrust. Diatoms showed uplift accompanying earthquakes in 290-0, 520-300, and 1050-790 calibrated years BP, and subsidence in AD 1964 and 640-510 calibrated years BP (Briggs et al., 2014). Such studies have implications for seismic hazard analysis, as the mixed uplift and subsidence record shows the variability of slip along the megathrust, suggesting that the segmentation of subduction zones is not fixed (Briggs et al., 2014; Ely et al., 2014).

Diatom data have also identified a possible preseismic signal from relatively small amounts of land-level change in the years prior to earthquakes (Bourgeois, 2006;

Shennan and Hamilton, 2006). Zong et al. (2003) used detailed modern and fossil diatom analyses from a coastal wetland along the eastern Alaska-Aleutian megathrust to reconstruct preseismic and coseismic land-level changes associated with the AD 1964 Alaska earthquake. In the 15 years prior to the AD 1964 earthquake, diatoms showed a gradual shift from a freshwater environment dominated by salt-intolerant taxa (e.g., *Eunotia exigua*, *Eunotia pectinales*, and *Achnanthes minutissima*) to salt-tolerant freshwater taxa typical of a high marsh environment (e.g., *Navicula pupula*, *Nitzschia fruticosa*, and *Pinnularia lagerstedtii*). Zong et al. (2003) interpret the shift in diatom assemblages as a phase of gradual land subsidence of ~0.15 m that preceded up to 2 m of coseismic subsidence in the AD 1964 earthquake. If preseismic deformation does occur, the implication is that warning signs may be detectable for several years prior to a great earthquake (Bourgeois, 2006).

Hamilton et al. (2005) explored whether the preseismic signal was a result of mixing of diatom assemblages from biological or physical processes. Hamilton et al. (2005) transplanted a block of marsh peat to a lower elevation in the intertidal zone where it would be buried by tidal mud. The results showed that mixing of diatoms did occur, but only in the top ~1 cm of peat, whereas the preseismic signal observed by Zong et al. (2003) and Hamilton and Shennan (2005b) occurred over 2-5 cm. In addition, Shennan and Hamilton (2006) argued that the preseismic signal was not a result of diatoms filtering down from overlying mud because dominant species that reflect subtle preseismic subsidence, such as *Nitzschia obtusa*, *Navicula begeri*, *Navicula brockmanii*, and *Pinnularia lagerstedtii*, do not occur in the overlying mud (Fig. 5). A similar transplant experiment at Cascadia found that mixing of foraminifera did occur up to 3 cm

below the transplanted marsh peat surface, mimicking a preseismic signal (Engelhart et al., 2013). Englehart et al. (2013) cautioned against interpreting microfossil assemblages in such mixed layers as a preseismic signal.

4. Application of diatoms to tsunami studies

4.1 Paleotsunamis

Sequences of tsunami deposits preserved along subduction zone coastlines can be used to estimate recurrence intervals of tsunamigenic earthquakes over centuries to millennia (Nanayama et al., 2003; Cisternas et al., 2005; Jankaew et al., 2008; Sawai et al., 2012). Stratigraphic sequences, supported by diatom studies, reveal repeated tsunamis in numerous locations including Alaska, the Pacific Northwest of North America, Chile, Japan, New Zealand, and locations bordering the North Sea and the Indian Ocean (Table 1, Fig. 2).

Allochthonous marine and brackish diatoms within tsunami deposits, including benthic and planktonic taxa, can support a seaward provenance of the sediment (e.g., Hemphill-Haley, 1995a; Sawai, 2002; Dura et al., 2015). Diatom analysis helped identify the provenance of three sand beds recovered from a coastal lake at Suijin-numa, on the subduction zone of the Japan Trench (Sawai et al., 2008). The marine and brackish diatoms (*Diploneis smithii*, *Delphineis surirella*) within the sand beds contrasted against the freshwater assemblages (*Aulacoseira granulata*, *A. crassipunctata*, *Eunotia* spp.) in the under and overlying mud. The middle sand bed at Suijin-numa correlates with the Jogan earthquake in AD 869 (Yoshida, 1906), the presumed predecessor to the 2011 Tohoku earthquake. Along the Kuril Trench in northern Japan, Nanayama et al. (2007) used diatoms to identify nine sandy tsunami deposits intercalated with peat. The diatom

assemblage within the peats contained freshwater species (e.g., *Eunotia* spp., *Pinnularia* spp.), whereas the sand beds were dominated by marine taxa (e.g., *Delphineis surirella* and *Odontella aurita*). Based on the record of tsunami deposits, the authors estimated a 365 – 553 year recurrence interval for large Kuril Trench earthquakes. Later, based on over 60 radiocarbon age estimates, Sawai et al. (2009b) found that the interval between tsunamis on the Kuril Trench ranged from 100 to 800 years, with an average recurrence interval of ~400 years.

In the Storegga Slide tsunami¹ deposits in Scotland, Dawson et al. (1996a) found an anomalous marine diatom assemblage with a high occurrence of fragmented diatoms. Up to 90% of valves (60% of which have elongate forms) within the sand bed were fragmented, reflecting the nature of the rapid, turbulent marine incursion of tsunami events (Dawson et al., 1996a; Smith et al., 2004; Fig. 7). Similar preferential fragmentation of diatom valves, in particular elongate forms (>100 µm) has been documented in inferred paleotsunami deposits in the Pacific Northwest of North America (Witter et al., 2009) and Japan (Sawai, 2002). Conversely, anomalously low breakage of diatoms in tsunami deposits has been reported in paleotsunamis from the Pacific coast of Washington State and Puget Sound, USA (Hemphill-Haley, 1996). Hemphill-Haley (1996) suggested that the low breakage of diatoms reflects rapid sedimentation by the tsunamis.

Diatoms can be used to estimate tsunami run-up beyond the landward limit of tsunami deposits. Hemphill-Haley (1996) used the distribution of diatoms to show that the inundation area of the tsunami from the AD 1700 Cascadia subduction zone

¹ The Storegga Slide tsunami (~7900 calibrated years BP) was produced by a submarine landslide (Bondevik et al., 2005).

earthquake was larger than the distribution inferred from the coarser-grained deposit visible in outcrop. Epipsammic tidal flat diatoms were found about 1 km farther upstream from the landward extent mapped in cores and stream channel outcrops.

4.2 Tsunamis of the past two decades

Diatom analysis of recent tsunami deposits provides modern analogues for reconstructing the fossil record. Recent studies characterized the diatom composition of the 1998 Papua New Guinea (Dawson, 2007), 2004 Indian Ocean (Sawai et al., 2009a;), 2009 South Pacific (Chagué-Goff et al., 2011), 2010 Maule Chile (Horton et al., 2011; Garrett et al., 2013), and 2011 Tohoku (Szczuciński et al., 2012b; Sawai et al., 2012) tsunamis. Analyzing the diatom signature of recent tsunamis can be used to infer the sedimentation (e.g., high-energy deposition), sediment provenance, and inundation limits of the events.

The taphonomy of diatom valves may indicate the mode of sedimentation of a tsunami. Dawson (2007) explored the condition of diatom valves in the 1998 Papua New Guinea tsunami deposit and found that linear, sigmoid, and clavate diatoms were more readily fragmented during the high-energy tsunami due to their relatively fragile valve structure. In contrast, high abundances of taphonomically unaltered (i.e., pristine) diatom valves were found in tsunami deposits from the 2004 Indian Ocean tsunami in Thailand (Sawai et al., 2009), and the 2010 Maule tsunami in Chile (Horton et al., 2011).

The upward fining of grain size observed in tsunami sands may be reflected by similar grading of diatom valves, a result of the variable flow speed of a tsunami (Gelfenbaum and Jaffe, 2003). In Thailand, the 2004 Indian Ocean tsunami deposit consisted of a sand bed with a thin mud cap and contained mostly beach and subtidal

344 diatoms (>80%) and very few freshwater species (Fig. 8; Sawai et al., 2009a). The lower
345 section of the deposit was dominated by larger epipsammic marine diatoms, whereas the
346 middle section contained abundant marine planktonic species, and the mud cap was
347 dominated by a mixture of smaller freshwater, brackish, and marine species. Diatom
348 analysis by Chagué-Goff et al. (2011) following the 2009 South Pacific tsunami deposit
349 in Samoa, Horton et al. (2011) following the 2010 Maule tsunami in Chile, and
350 Szczuciński et al. (2012b) following the 2011 Tohoku-Oki tsunami on the Sendai plain
351 also revealed grading of diatom valves.

352 Diatoms recovered from tsunami deposits perpendicular to the coast show variable
353 concentration, ecology, and taphonomy related to the provenance of material.
354 Szczuciński et al. (2012b) found that within 1 km of the coast, the Tohoku-Oki sand beds
355 were derived mainly from the beach and coastal dunes, and because such sediments were
356 devoid of diatoms, the assemblage within the tsunami sediments was composed of a low
357 concentration of fragmented freshwater and few brackish species sourced from the
358 coastal plain. Further inland the diatom concentration within the tsunami sediments
359 increased, and the assemblage and condition of valves in the deposit (% fragmentation of
360 valves) were very similar to the valves in the underlying soil and nearby freshwater canal,
361 suggesting that the sediment was locally sourced and not transported from the coast
362 (Szczuciński et al., 2012b).

363 Chagué-Goff et al. (2015) used the diatom assemblages of the 2010 Maule Chile
364 tsunami deposit to trace tsunami inundation beyond the limit of sedimentological
365 evidence. Marine diatoms could be traced ~100 m beyond the inundation limit identified
366 by the tsunami deposit.

5. Knowledge gaps

Diatoms have improved our understanding of the nature of land-level change associated with past earthquakes and have helped define the impacts of past tsunamis. However, when using diatoms for reconstructing earthquake-related land-level change and tsunami inundation, several matters must be kept in mind.

5.1 Uncertainties in modern diatom distributions

An absence of local to regional modern diatom datasets from a range of coastal environments restricts the application of diatoms to reconstruct the history of subduction zone earthquakes and tsunamis. The relation of diatoms to tidal elevation is unknown in many locations because of a scarcity of modern data (e.g., Dura et al., 2015). However, even modern diatom data sets that include hundreds of samples taken across multiple local coastal wetlands can fail to provide modern analogues for fossil diatom assemblages (Watcham et al., 2013). This non-modern-analogue situation (Birks, 1995) is attributed to local environmental conditions having changed significantly over time (Watcham et al., 2013). If this happens, it is not appropriate to apply quantitative reconstruction techniques (e.g., transfer function) using only local modern diatom assemblages (Watcham et al., 2013). Regional modern training sets compiled from a large range of intertidal coastal environments (e.g. marshes with a variety of vegetation zones, substrates, and elevation gradients) can account for such variation in the distribution of diatoms, providing analogs for diatom assemblages found in fossil cores (Zong et al., 2003; Watcham et al., 2013; Shennan et al., 2014a).

5.2 *Response of diatoms to coseismic land-level change*

When producing diatom-based estimates of earthquake related land-level change across sharp stratigraphic contacts, a possible delay in sediment deposition and diatom response must be considered. If there is a significant (months to years) hiatus in response following an earthquake, estimates of coseismic deformation will include both coseismic and postseismic deformation, thus they will be minimum estimates (Garrett et al., 2013). Shennan et al. (2014a) found that peat formation following the coseismic uplift of an intertidal mudflat along the eastern Alaska-Aleutian megathrust was not instantaneous, resulting in minimum estimates of uplift. The study also found a mixed salinity diatom assemblage in the upper part of the buried silt that formed as a result of ponding and reworking of the uplifted surface before colonization by terrestrial plants, and likely did not represent the pre-earthquake environment and elevation. In Chile, Garrett et al. (2013) found that uplifted marshes had accumulated little to no sediment up to two years after the 2010 earthquake and tsunami, complicating estimates of coseismic uplift. Coseismic subsidence more often results in rapid sedimentation of clastic sediments hours to weeks after an earthquake, due to the accommodation space created in the submerged marsh (Atwater et al., 2001; Hawkes et al., 2011).

5.3 *The production and preservation of diatoms*

Diatom-based reconstructions of earthquake-related land-level change and tsunami inundation are hindered by variable diatom production and preservation. In the Copper River Delta, Shennan et al. (2014a) found low numbers of diatoms in modern samples taken from tidal flat silts, a result of high sediment accretion of the delta environment. Silts in fossil sequences also contained few diatoms, which complicated quantitative

reconstructions of earthquake-related land-level change (Shennan et al., 2014a). In relatively alkaline and warm environments, there are problems with chemical dissolution of diatom valves (Katamani, 1982; Flower, 1993; Barker et al., 1994). Sawai et al. (2009a) found excellent preservation of diatoms in the 2004 Indian Ocean tsunami deposit in Thailand, but three paleotsunami deposits examined at the same site by Jankeaw et al. (2008) contained no fossil diatoms. Dissolution of diatom valves can also make them more susceptible to fragmentation, complicating the use of the fragmentation metric to infer high-energy deposition during a tsunami. Szczuciński et al. (2012b) found a similar degree of fragmentation in diatoms within the soil underlying the 2011 tsunami deposit and within the deposit itself.

5.4 Differentiating autochthonous and allochthonous diatoms

Problems differentiating autochthonous (*in situ*) and allochthonous (transported) diatoms in modern and fossil studies complicate reconstructions. Certain taxa are transported across coastal wetlands and, if included in modern or fossil assemblages, erroneously represent the depositional environment. This problem is common in coastal wetlands where diatom valves are transported from one intertidal floral zone to another by daily tidal currents, or, in colder climates, by ice rafted sediment (Hemphill-Haley et al., 1995a; Hamilton et al., 2005). Hemphill-Haley (1995a) and Sawai (2001b, 2004) suggested several means to distinguish allochthonous diatoms in both modern and fossil assemblages. Planktonic diatoms are considered allochthonous components in modern and fossil coastal wetland assemblages, while benthic taxa (e.g., epiphytic, epipellic and epipsammic types) can be considered as autochthonous (Vos and de Wolf, 1993). Some chain-forming taxa with thickly silicified valves may form prominent allochthonous

assemblages. Hemphill-Haley (1995a) excluded the marine tychoplanktonic diatom *Paralia sulcata* from paleoecological interpretations, because its robust valves and long-chained structure allow its valves to be easily floated and transported by tidal currents and deposited far inland in tidal estuaries (Hemphill-Haley, 1995a). Sawai (2001b, 2004b) recognized allochthonous distributions of certain epiphytic taxa in the intertidal zone. For example, *Cocconeis scutellum* is an epiphytic species commonly found attached to macrophytes in the intertidal zone. After death, the upper (rapheless) valve that is not attached to the macrophyte may detach and be transported by tidal currents (Sawai, 2004b). As a result, rapheless valves of *C. scutellum* may be found in sediment across the entire intertidal zone, although its habitat is limited to the macrophyte zone (Sawai, 2001b).

5.5 Differentiating tsunami and storm surge deposits

Allochthonous marine and brackish diatoms may be found in storm-surge deposits, making it difficult to differentiate from tsunami deposits in the stratigraphic record (e.g. Liu and Fearn, 2000; Parsons, 1998; Tuttle et al., 2004; Horton et al., 2009). However, coseismic land-level change coincident with anomalous silt and sand beds provides a definitive earthquake source for a tsunami. In Washington State, USA Hemphill-Haley (1995a) identified two coarse-grained silt and sand beds dominated by marine diatoms. A tsunami source was inferred from one bed because it was associated with a sudden change in depositional environment from upland soil to intertidal mud, consistent with coseismic submergence. The other bed within a freshwater upland soil was not associated with a change in stratigraphy and was probably deposited by a storm (Hemphill-Haley, 1995a).

6. Conclusions

Using global examples, we illustrated the utility of diatoms in reconstructing land-level change and tsunami inundation histories along subduction zone coastlines on centennial and millennial timescales. We outlined the evolution of fully quantitative, statistical techniques to estimate coseismic land-level change at the Cascadia subduction zone. Examples from the Alaska-Aleutian megathrust and the Japan trench illustrated the expanded application of the transfer function technique to reconstruct preseismic, coseismic, and postseismic land-level change. In central Chile, diatoms were successfully applied to identify coseismic uplift, and in Alaska, diatoms helped reconstruct a mixed uplift and subsidence record preserved in coastal wetland stratigraphy.

We illustrated the utility of diatoms in inferring the sediment provenance, sedimentation, and inundation limit of tsunamis. Allochthonous marine and brackish diatoms within anomalous sand beds along the coasts of Japan, Scotland, and Thailand indicate a seaward provenance of sediments, supporting a tsunami source. The high fragmentation of diatom valves in sand beds deposited by the Storegga tsunami and the 1998 Papua New Guinea tsunami reflect high-energy sedimentation, while the upward fining of diatom valves in the 2004 Indian Ocean tsunami reflect the variable flow speed of a tsunami. Examples from the Cascadia subduction zone and the Chile subduction zone showed diatoms how can be used to estimate tsunami run-up beyond the landward limit of tsunami deposits, a powerful tool in estimating paleotsunami inundation.

We outlined a series of knowledge gaps that should be considered in future research. Many of the knowledge gaps in the study of diatom-based earthquake and tsunami records can be addressed by continuing to explore the modern diatom environments, in

particular exploring diatoms' relation to salinity and substrate, variable production, preservation, and transport. Examining the response of diatoms to recent earthquake-related land-level change and the character of modern tsunami deposit diatom assemblages will continue to provide analogues for identifying events in the fossil record.

Acknowledgements

This work was supported by funding from the National Science Foundation grant (EAR-1144537, 1419824, 1357722) awarded to BPH. We thank Ursula Cochran and Alan Nelson for constructive reviews that substantially improved the manuscript.

References

- Admiraal, W., 1984. The ecology of estuarine sediment-inhabiting diatoms. *Progress in Phycological Research* 3, 269-322.
- Amspoker, M.C., McIntire, C.D., 1978. Distribution of diatoms associated with sediments in Yaquina Estuary, Oregon. *Journal of Phycology*, 14, 387-395.
- Atwater, B. F., 1987. Evidence for great earthquakes along the outer coast of Washington state. *Science*, 236, 942–944. doi:10.1126/science. 236.4804.942.
- Atwater, B. F., Hemphill-Haley, E., 1997. Recurrence intervals for great earthquakes of the past 3,500 years at northeastern Willapa Bay, Washington. U.S. Geological Survey Professional Paper, 1576, 1-108.
- Atwater, B. F., Nunez, H. J., Vita-Finzi, C., 1992. Net late Holocene emergence despite earthquake-induced submergence, south-central Chile, in: *Impacts of Tectonics on Quaternary Coastal Evolution*, Y. Ota, A. R. Nelson and K. R. Berryman (Editors), *Quaternary International*, 15/16, 77–85.
- Atwater, B.F., 1992. Geologic evidence for earthquakes during the past 2000 years along the Copalis River, southern coastal Washington. *Journal of Geophysical Research*, 97, 1901-1919.
- Atwater, B.F., Furukawa, R., Hemphill-Haley, E., Ikeda, Y., Kashima, K., Kawase, K., Kelsey, H.M., Moore, A.L., Nanayama, F., Nishimura, Y., Odagiri, S., Ota, Y., Park, S., Satake, K.,

- Sawai, Y., Shimokawa, K., 2004. Seventeenth-century uplift in eastern Hokkaido, Japan. *The Holocene*, 14, 489-501.
- Barker, P., Fontes, J., Gasse, F., Druart, J., 1994. Experimental dissolution of diatom silica in concentrated salt solutions and implications for paleoenvironmental reconstruction. *Limnology and Oceanography*, 39, 99–110.
- Barlow, N.L.M., Shennan, I., Long, A.J., 2012. Relative sea-level response to Little Ice Age mass change in south central Alaska: reconciling model predictions and geological evidence. *Earth and Planetary Science Letters* 315-316, 62-75.
- Bartsch-Winkler, S., Schmoll, H.R., 1987. Earthquake-caused sedimentary couplets in the upper Cook Inlet region. U.S. Geological Survey, Circular 998, 92-95.
- Battarbee, R. W., Jones, V. J., Flower, R. J., Cameron, N. G., Bennion, H., Carvalho, L., Juggins, S., 2001. *Diatoms* (pp. 155-202). Springer Netherlands.
- Benson, B.E., Grimm, K.A., Clague, J.J., 1997. Tsunami deposits beneath tidal marshes on northwestern Vancouver Island, British Colombia. *Quaternary Research*, 48, 192-204.
- Berkeley, A., Perry, C. T., Smithers, S. G., 2009. Taphonomic signatures and patterns of test degradation on tropical, intertidal benthic foraminifera. *Marine Micropaleontology*, 73(3), 148-163.
- Birks, H.J.B., 1995. Statistical modelling of Quaternary science data. In: Maddy, D., Brew, J.S. (Eds.), *Quantitative Paleoenvironmental Reconstructions*. Quaternary Research Association, Cambridge 5, 161-254.
- Bondevik, S., 2003. Storegga tsunami sand in peat below the Tapes beach ridge at Haroy, western Norway, and its possible relation to an early Stone Age settlement. *Boreas*, 32, 476-483.
- Bondevik, S., Mangerud, J., Dawson, S., Dawson, A., Lohne, O., 2005. Evidence for three North Sea tsunamis at the Shetland Islands between 8000 and 1500 years ago. *Quaternary Science Reviews*, 24, 1757–1775.
- Bourgeois, J., Reinhart, M.A., 1989. Onshore erosion and deposition by the 1960 tsunami at the Rio Lingue estuary, south-central Chile. *American Geophysical Union Transactions*, 70, 1331.
- Bourgeois, J., Petroff, C., Yeh, H., Titov, V., Synolakis, C.E., Benson, B., Kuroiwa, J., Lander, J., Norabuena, E., 1999. Geologic setting, field survey and modeling of the Chimbote, Northern Peru, tsunami of 21 February 1996. *Pure and Applied Geophysics*, 154, 513-540.
- Bourgeois, J., 2006. Earthquakes: a movement in four parts? *Nature*, 440, 430-431.
- Briggs, R. W., Engelhart, S. E., Nelson, A. R., Dura, T., Kemp, A. C., Haeussler, P. J., Corbett, D.R., Angster, S.J., Bradley, L.A., 2014. Uplift and subsidence reveal a non-persistent megathrust rupture boundary (Sitkinak Island, Alaska). *Geophysical Research Letters*, 41, 2289-2296.

- Brill, D., Jankaew, K., Neubauer, N., Kelletat, D., Scheffers, A., Vött, Brückner, H., 2014. Holocene coastal evolution of southwest Thailand – implications for the site-specific preservation of palaeotsunami deposits. *Zeitschrift für Geomorphologie*, 58 (1).
- Chagué-Goff, C., Dawson, S., Goff, J. R., Zachariasen, J., Berryman, K. R., Garnett, D. L., Mildenhall, D.C., 2002. A tsunami (ca. 6300 years BP) and other Holocene environmental changes, northern Hawke's Bay, New Zealand. *Sedimentary Geology*, 150, 89-102.
- Chagué-Goff, C., Schneider, J. L., Goff, J. R., Dominey-Howes, D., Strotz, L., 2011. Expanding the proxy toolkit to help identify past events—lessons from the 2004 Indian Ocean Tsunami and the 2009 South Pacific Tsunami. *Earth-Science Reviews*, 107, 107-122.
- Chagué-Goff, C., Goff, J., Wong, H. K., and Cisternas, M., 2015. Insights from geochemistry and diatoms to characterise a tsunami's deposit and maximum inundation limit. *Marine Geology*, 359, 22-34.
- Chlieh, M., Avouac, J. P., Hjorleifsdottir, V., Song, T. R. A., Ji, C., Sieh, K., Galetzka, J., 2007. Coseismic slip and afterslip of the great Mw 9.15 Sumatra–Andaman earthquake of 2004. *Bulletin of the Seismological Society of America*, 97(1A), S152-S173.
- Cisternas, M., Atwater, B.F., Torrejón, F., Sawai, Y., Machuca, G., Lagos, M., Eipert, A., Youlton, C., Salgado, I., Kamataki, T., Shishikura, M., Rajendran, C.P., Malik, J.K., Rizal, Y., Husni, M., 2005. Predecessors of the giant 1960 Chile earthquake. *Nature* 437, 404-407.
- Clague, J.J., Bobrowsky, P.T., 1994. Evidence for a large earthquake and tsunami 100-400 years ago on western Vancouver Island, British Columbia. *Quaternary Research*, 41, 176-184.
- Clague, J.J., Hutchinson, I., Mathewes, R.W., Patterson, R.T., 1999. Evidence for Late Holocene tsunamis at Catala Lake, British Columbia. *Journal of Coastal Research* 15, 45-60.
- Clark, K. J., Hayward, B. W., Cochran, U. A., Grenfell, H. R., Hemphill–Haley, E., Mildenhall, D. C., Hemphill-Haley, M.A., Wallace, L. M., 2011. Investigating subduction earthquake geology along the southern Hikurangi margin using palaeoenvironmental histories of intertidal inlets. *New Zealand Journal of Geology and Geophysics*, 54(3), 255-271.
- Clark, K. J., Hayward, B. W., Cochran, U. A., Wallace, L. M., Power, W. L., and Sabaa, A. T., 2015. Evidence for Past Subduction Earthquakes at a Plate Boundary with Widespread Upper Plate Faulting: Southern Hikurangi Margin, New Zealand. *Bulletin of the Seismological Society of America*.
- Cleve, P.T., 1894. Synopsis of the naviculoid diatom I and II. *Kongliga Svenska Vetenskaps-Akademien* 26, 1–194.
- Cleve, P.T., 1895. Synopsis of the naviculoid diatom I and II. *Kongliga Svenska Vetenskaps-Akademien* 27, 1–129.
- Cleve-Euler, A., 1922. Om diatomacevegetationen och dess förändringar i Sabysjon, Uppland samt negra damda sjoar i Salatrakten. *Sveriges Geologiska Undersökning C309*, 1–76.

- Cochran, U.A., Berryman, K.R., Mildenhall, D.C., Hayward, B.W., Southall, K., Hollis, C.J., 2005. Towards a record of Holocene tsunami and storms for northern Hawke's Bay, New Zealand. *New Zealand Journal of Geology and Geophysics*, 48(3): 507-515.
- Cochran, U., Berryman, K., Zachariassen, J., Mildenhall, D., Hayward, B., Southall, K., Hollis, K., Wilson, K., 2006. Paleocological insights into subduction zone earthquake occurrence, eastern North Island, New Zealand. *Geological Society of America Bulletin*, 118, 1051-1074.
- Cooper, S., Gaiser, E.E., Wachnicka, A., 2010. Estuarine paleocological reconstructions using diatoms. In: Smol, J.P., Stoermer, E.F. (Eds.), *The Diatoms: Applications for the Environmental and Earth Sciences*. Cambridge University Press, Cambridge, 324-345.
- Combellick, R.A., 1991. Paleoseismicity of the Cook Inlet region, Alaska: Evidence from peat stratigraphy in Turnagain and Knik Arms. *Alaska Division of Geological and Geophysical Surveys Professional Report*, Fairbanks, 52 pp.
- Combellick, R.A., 1994. Investigation of peat stratigraphy in tidal marshes along Cook Inlet, Alaska, to determine the frequency of 1964-style great earthquakes in the Anchorage region. *Alaska Division of Geological and Geophysical Surveys Report of Investigations*, 94-97, Fairbanks, 24 pp.
- Combellick, R., Reger, R.D., 1989. Evaluation of Holocene subsidence events of Cook Inlet estuarine flats near Anchorage, Alaska, as a basis for assessing seismic hazards in south-central Alaska. *USGS Open-File Report 89-453*, 445-446.
- Combellick, R.A., Reger, R.D., 1994. Sedimentological and radiocarbon- age data for tidal marshes along eastern and upper Cook Inlet, Alaska. *Alaska Division of Geological and Geophysical Surveys Report of Investigations*, 94-96, Fairbanks, 60 pp.
- Clark, K.J., Hayward, B.W., Cochran, U.A., Wallace, L.M., Power, W.L., Sabaa, A.T., 2015. Evidence for past subduction earthquakes at a plate boundary with widespread upper plate faulting: southern Hikurangi margin, New Zealand. *Bulletin of the Seismological Society of America*, 105(3): doi: 10.1785/0120140291
- Darlenzo, M.E., Peterson, C.D., 1990. Episodic tectonic subsidence of late Holocene salt marshes, northern Oregon central Cascadia margin. *Tectonics*, 9, 1-22.
- Darlenzo, M.E., Peterson, C.D., Clough, C., 1994. Stratigraphic evidence for great subduction-zone earthquakes at four estuaries in northern Oregon, USA. *Journal of Coastal Research*, 10, 850-876.
- Dawson, S., 2007. Diatom biostratigraphy of tsunami deposits: examples from the 1998 Papua New Guinea tsunami. *Sedimentary Geology*, 200, 328-335.
- Dawson, S., Smith, D.E., 2000. The sedimentology of mid-Holocene tsunami facies in northern Scotland. *Marine Geology*, 170, 69-79.
- Dawson, A.G., Stewart, I., 2007. Tsunami deposits in the geological record. *Sedimentary Geology*, 200, 166-183.

- Dawson, S., Smith, D.E., Ruffman, A., Shi, S., 1996a. The diatom biostratigraphy of tsunami deposits: Examples from recent and middle Holocene events. *Physics and Chemistry of the Earth*, 21, 87-92.
- Dawson, A.G., Shi, S., Dawson, S., Takahashi, T., Shuto, N., 1996b. Coastal sedimentation associated with the June 2nd and 3rd, 1994 tsunami in Rajegwesi, Java. *Quaternary Science Reviews*, 15, 901-912.
- Denys, L., 1991-1992. A check-list of the diatoms in the Holocene deposits of the western Belgian coastal plain with a survey of their apparent ecological requirements. Ministère des affaires économiques, Service Géologique de Belgique.
- Dominey-Howes, D., Cundy, A., Croudace, I., 2000. High energy marine flood deposits on Astypalaea Island, Greece: possible evidence for the AD 1956 southern Aegean tsunami. *Marine Geology*, 163, 303-315.
- Dura, T., Rubin, C.M., Kelsey, H.M., Horton, B.P., Hawkes, A.D., Vane, C.H., Daryono, M., Grand Pre, C., Ladinsky, T., Bradley, S., 2011. Stratigraphic record of Holocene coseismic subsidence, Padang, West Sumatra. *Journal of Geophysical Research*, 116, B11306.
- Dura, T., Cisternas, M., Horton, B.P., Ely, L.L., Nelson, A.R., Wesson, R.L., Pilarczyk, J.E., 2015. Coastal evidence for Holocene subduction-zone earthquakes and tsunamis in central Chile. *Quaternary Science Reviews*, 113, 93-111.
- Ely, L.L., Cisternas, M., Wesson, R.L., Dura, T., 2014. Five centuries of tsunamis and land-level changes in the overlapping rupture area of the 1960 and 2010 Chilean earthquakes. *Geology*. G35830-1.
- Engelhart, S.E., Horton, B.P., Nelson, A.R., Hawkes, A.D., Witter, R.C., Wang, K., Wang, P.-L., Vane, C.H., 2013. Testing the use of microfossils to reconstruct great earthquakes at Cascadia. *Geology*, 41(10), 1067-1070. doi, 10.1130/G34544.1
- Engelhart, S. E., Horton, B. P., Vane, C. H., Nelson, A. R., Witter, R. C., Brody, S. R., Hawkes, A. D., 2013. Modern foraminifera, $\delta^{13}\text{C}$, and bulk geochemistry of central Oregon tidal marshes and their application in paleoseismology. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 377, 13-27.
- Flower, R. J., 1993. Diatom preservation: experiments and observations on dissolution and breakage in modern and fossil material. *Hydrobiologia*, 269/270, 473-84.
- Garrett, E., Shennan, I., Watcham, E.P., Woodroffe, S.A., 2013. Reconstructing paleoseismic deformation, 1: modern analogues from the 1960 and 2010 Chilean great earthquakes. *Quaternary Science Reviews*, 75, 11-21.
- Garrett, E., Shennan, I., Woodroffe, S. A., Cisternas, M., Hocking, E. P., Gulliver, P., 2015. Reconstructing paleoseismic deformation, 2: 1000 years of great earthquakes at Chucalén, south central Chile. *Quaternary Science Reviews*, 113, 112-122.
- Geller, R. J., 2011. Shake-up time for Japanese seismology. *Nature*, 472, 407-409.

- Gelfenbaum, G., Jaffe, B., 2003. Erosion and sedimentation from the 17 July 1998 Papua New Guinea tsunami. *Pure and Applied Geophysics*, 160, 1969-1999.
- Goff, J., McFadgen, B. G., Chagué-Goff, C., 2004. Sedimentary differences between the 2002 Easter storm and the 15th-century Okoropunga tsunami, southeastern North Island, New Zealand. *Marine geology*, 204(1), 235-250.
- Goff, J., Liu, P. L., Higman, B., Morton, R., Jaffe, B. E., Fernando, H., Fernando, S., 2006. Sri Lanka field survey after the December 2004 Indian Ocean tsunami. *Earthquake Spectra*, 22, 155-172.
- Goff, J., Lamarche, G., Pelletier, B., Chagué-Goff, C., Strotz, L., 2011. Predecessors to the 2009 South Pacific tsunami in the Wallis and Futuna archipelago. *Earth-Science Reviews* 107, 91-106.
- Goff, J., Chagué-Goff, Nichol, S., Jaffe, B., Dominey-Howes, D., 2012. Progress in palaeotsunami research. *Sedimentary Geology*, 243, 70-88.
- Goto, T., Satake, K., Sugai, T., Ishibe, T., Harada, T., Murotani, S., 2015. Historical tsunami and storm deposits during the last five centuries on the Sanriku coast, Japan. *Marine Geology*.
- Graehl, N. A., Kelsey, H. M., Witter, R. C., Hemphill-Haley, E., and Engelhart, S. E., 2015. Stratigraphic and microfossil evidence for a 4500-year history of Cascadia subduction zone earthquakes and tsunamis at Yaquina River estuary, Oregon, USA. *Geological Society of America Bulletin*, 127(1-2), 211-226.
- Grand Pre, C.A., Horton, B.P., Kelsey, H.M., Rubin, C.M., Hawkes, A.D., Daryono, M.R., Rosenberg, G., Culver, S.J., 2012. Stratigraphic evidence for an early Holocene earthquake in Aceh, Indonesia. *Quaternary Science Reviews*, 54, 142-151.
- Grauert, M., Björck, S., Bondevik, S., 2001, Storegga tsunami deposits in a coastal lake on Suduroy, the Faroe Islands: *Boreas*, 30, 263-271.
- Guilbault, J.P., Clague, J.J., Lapointe, M., 1995. Amount of subsidence during a late Holocene earthquake – Evidence from fossil tidal marsh foraminifera at Vancouver Island, west coast of Canada. *Palaeogeography Palaeoclimatology Palaeoecology*, 118, 49-71.
- Guilbault, J.-P., Clague, J.J., Lapointe, M., 1996. Foraminiferal evidence for the amount of coseismic subsidence during a late Holocene earthquake on Vancouver Island, west coast of Canada. *Quaternary Science Reviews*, 15, 913-937.
- Hamilton, S., Shennan, I., 2005a. Late Holocene relative sea-level changes and the earthquake deformation cycle around upper Cook Inlet, Alaska. *Quaternary Science Reviews*, 24, 1479-1498.
- Hamilton, S., and Shennan, I., 2005b. Late Holocene great earthquakes and relative sea-level change at Kenai, southern Alaska. *Journal of Quaternary Science*, 20(2), 95-111.

- Hamilton S, Shennan I, Combellick R, Mulholland, J., Noble, C., 2005. Evidence for two great earthquakes at Anchorage, Alaska and implications for multiple great earthquakes through the Holocene. *Quaternary Science Reviews* 24: 2050–2068.
- Hartley, B., Ross, R., Williams, D.M., 1996. A check-list of the freshwater, brackish and marine diatoms of the British Isles and adjoining coastal waters. *Journal of the Marine Biological Association of the United Kingdom* 66, no. 03: 531-610.
- Hawkes, A. D., Bird, M., Cowie, S., Grundy-Warr, C., Horton, B. P., Hwai, A. T. S., Law, L., Macgregor, C., Nott, J., Eong Ong, J., Rigg, J., Robinson, R., Tan-Mullins, M., Tiong Sa, T., Yasin, Z., Aik, L. W., 2007. Sediments deposited by the 2004 Indian Ocean tsunami along the Malaysia–Thailand Peninsula. *Marine Geology*, 242(1), 169-190.
- Hawkes, A. D., Horton, B. P., Nelson, A. R., Hill, D. F., 2010. The application of intertidal foraminifera to reconstruct coastal subsidence during the giant Cascadia earthquake of AD 1700 in Oregon, USA. *Quaternary International*, 221(1), 116-140.
- Hawkes, A.D., Horton, B.P., Nelson, A.R., Vane, C.H., Sawai, Y., 2011. Coastal subsidence in Oregon, USA, during the giant Cascadia earthquake of AD 1700. *Quaternary Science Reviews*, 30, 364-376.
- Hayward, B. W., Grenfell, H. R., Sabaa, A. T., Carter, R., Cochran, U., Lipps, J. H., Shane, P.R., Morley, M. S., 2006. Micropaleontological evidence of large earthquakes in the past 7200 years in southern Hawke's Bay, New Zealand. *Quaternary Science Reviews*, 25(11), 1186-1207.
- Heki, K., 2011. A tale of two earthquakes. *Science*, 332, 1390-1391.
- Hemphill-Haley, E., 1993. Taxonomy of recent and fossil (Holocene) diatoms (Bacillariophyta) from northern Willapa Bay, Washington (No. 93-289). US Geological Survey,.
- Hemphill-Haley, E., 1995a. Diatom evidence for earthquake-induced subsidence and tsunami 300 years ago in southern coastal Washington. *Geological Society of America Bulletin*, 107, 367–378.
- Hemphill-Haley, E., 1995b. Intertidal diatoms from Willapa Bay, Washington: applications to studies of small-scale sea-level changes. *Northwest Science*, 69, 29–45.
- Hemphill-Haley, E., 1996. Diatoms as an aid in identifying late Holocene tsunami deposits. *The Holocene*, 6, 439–448.
- Horton, B.P., 1999. The distribution of contemporary intertidal foraminifera at Cowpen Marsh, Tees Estuary, UK: implications for studies of Holocene sea-level changes. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 149, 127-149.
- Horton, B.P., Edwards, R.J., 2005. The application of local and regional transfer functions to reconstruct former sea levels, North Norfolk, England. *The Holocene*, 15, 216-228.
- Horton, B.P., Edwards, R.J., 2006. Quantifying Holocene sea level change using intertidal foraminifera: lessons from the British Isles. *Cushman Foundation for Foraminiferal Research, Special Publication* 40, 97 pp.

- Horton, B. P., Zong, Y., Hillier, C., Engelhart, S., 2007. Diatoms from Indonesian mangroves and their suitability as sea-level indicators for tropical environments. *Marine Micropaleontology*, 63, 155-168.
- Horton, B.P., Rossi, V., Hawkes, A.D., 2009. The sedimentary record of the 2005 hurricane season from the Mississippi and Alabama coastlines. *Quaternary International*, 195, 15-30.
- Horton, B. P., Sawai, Y., Smol, J. P., Stoermer, E. F., 2010. Diatoms as indicators of former sea levels, earthquakes, tsunamis, and hurricanes. *The diatoms: applications for the environmental and earth sciences*, (Ed. 2), 357-372.
- Horton, B.P., Sawai, Y., Hawkes, A.D., Witter, R.C., 2011. Sedimentology and paleontology of a tsunami deposit accompanying the great Chilean earthquake of February 2010. *Marine Micropaleontology*, 79, 132-138.
- Horton, B.P., Engelhart, S.E., Kemp, A.C., Sawai, Y., 2013. Microfossils in tidal settings as indicators of sea-level change, paleoearthquakes, tsunamis, and tropical cyclones. In: John F. Shroder (ed.) *Treatise on Geomorphology*, vol. 14, San Diego: Academic Press, 292-314.
- Hughes, J.F., Mathewes, R.W., Clague, J.J., 2002. Use of pollen and vascular plants to estimate coseismic subsidence at a tidal marsh near Tofino, British Columbia. *Palaeogeography Palaeoclimatology Palaeoecology*, 185, 145-161.
- Hustedt, F., 1937. Systematische und ökologische Untersuchungen der Diatomeen floren von Java, Bali, Sumatra. *Archiv Hydrobiologie Supplement*, 15, 131–506.
- Hustedt, F., 1939. Systematische und ökologische Untersuchungen der Diatomeen floren von Java, Bali, Sumatra. *Archiv Hydrobiologie Supplement*, 16, 1–394.
- Hustedt, F., 1953. Diatomeen aus dem Naturschutzpark Seeon. *Archiv für Hydrobiologie*, 47, 625–35.
- Hustedt, F., 1957. Die Diatomeenflora des Fluss-systems der Weser im Gebiet der Hansestadt Bremen. *Abhandlungen vom Naturwissenschaftlichen Verein zu Bremen* 34, 181–440.
- Hustedt, F., 1958. Diatomeen aus der Antarktis and dem Siidatlantik. *Deutsche Antarkt. Exped. 1938/39, Wiss. Ergebn.*, 2 103-191.
- Hutchinson, I., Clague, J.J., Mathewes, R.W., 1997. Reconstructing the tsumani record on an emerging coast: A case study of Kanim Lake, Vancouver Island, British Columbia, Canada: *Journal of Coastal Research*, 13, 545–553.
- Hutchinson, I., Guilbault, J. P., Clague, J. J., Bobrowsky, P. T., 2000. Tsunamis and tectonic deformation at the northern Cascadia margin: a 3000-year record from Deserated Lake, Vancouver Island, British Columbia, Canada. *The Holocene*, 10(4), 429-439.
- Iliev, A. Y., Kaistrenko, V. M., Gretskeya, E. V., Tikhonchuk, E. A., Razjigaeva, N. G., Grebennikova, T. A., and Kharlamov, A. A., 2005. Holocene tsunami traces on Kunashir Island, Kurile subduction zone. In *Tsunamis* (pp. 171-192). Springer Netherlands.

- James, T., Gowan, E. J., Hutchinson, I., Clague, J. J., Barrie, J. V., Conway, K. W., 2009. Sea-level change and paleogeographic reconstructions, southern Vancouver Island, British Columbia, Canada. *Quaternary Science Reviews*, 28(13), 1200-1216.
- Jankaew, K., Atwater, B.F., Sawai, Y., Choowong, M., Charoentitirat, T., Martin, M.E., Prendergast, A., 2008. Medieval forewarning of the 2004 Indian Ocean tsunami in Thailand. *Nature*, 455, 1228-1231.
- Jones, V.J., 2007. Diatom Introduction, in: Elias, S., (Ed.) *Encyclopedia of Quaternary Science*. Elsevier, Oxford, pp. 476 - 484.
- Juggins, S., 1992. Diatoms in the Thames estuary, England: ecology, palaeoecology, and salinity transfer function. *Bibliotheca Diatomologica*, 25, 216.
- Kamatani, A., 1982. Dissolution rates of silica from diatoms decomposing at various temperatures. *Marine Biology*, 68, 91-96.
- Kelsey, H. M., Witter, R. C., Hemphill-Haley, E., 2002. Plate-boundary earthquakes and tsunamis of the past 5500 yr, Sixes River estuary, southern Oregon. *Geological Society of America Bulletin*, 114(3), 298-314.
- Kelsey, H. M., Sherrod, B., Johnson, S. Y., Dadisman, S. V., 2004. Land-level changes from a late Holocene earthquake in the northern Puget Lowland, Washington. *Geology*, 32(6), 469-472.
- Kelsey, H.M., Nelson, A.R., Hemphill-Haley, E., Witter, R.C., 2005. Tsunami history of an Oregon coastal lake reveals a 4600 yr record of great earthquakes on the Cascadia subduction zone. *Geological Society of America Bulletin*, 117, 1009-1032.
- Kemp, A.C., Horton, B.P., Culver, S.J., Corbett, D.R., van de Plassche, O., Gehrels, W.R., Douglas, B.C., Parnell, A.C., 2009. Timing and magnitude of recent accelerated sea-level rise (North Carolina, United States). *Geology*, 37, 1035-1038.
- Kokociński, M., Szczuciński, W., Zgrundo, A., and Ibragimow, A., 2009. Diatom assemblages in 26 December 2004 tsunami deposits from coastal zone of Thailand as sediment provenance indicators. *Polish Journal of Environmental Studies*, 18(1), 93-101.
- Kon'no, E., 1961. Geological observations of the Sanriku coastal region damaged by tsunami due to the Chile earthquake in 1960, Contributions to the Institute of Geology and Paleontology of Tohoku University, 52, 1-40. (in Japanese with English abstract)
- Kortekaas, S., Dawson, A. G., 2007. Distinguishing tsunami and storm deposits: an example from Martinhal, SW Portugal. *Sedimentary Geology*, 200, 208-221.
- Krammer, K., Lange-Bertalot, H., 1986. Bacillariophyceae 2/1. Naviculaceae, in: Ettl, H., Gerloff, J., Heynig, H., Mollenhauser (Eds.), *Süßwasserflora von Mitteleuropa*. Gustav Fischer Verlag, Stuttgart, pp. 1-876.
- Krammer, K., Lange-Bertalot, H., 1988. Bacillariophyceae 2/2. Basillariaceae, Epithemiaceae, Surirellaceae, in: Ettl, H., Gerloff, J., Heynig, H., Mollenhauser (Eds.), *Süßwasserflora von Mitteleuropa*. Gustav Fischer Verlag, Stuttgart, pp. 1-600.

- Krammer, K., Lange-Bertalot, H., 1991a. Bacillariophyceae 2/3. Centrales, Fragilariaceae, Eunotiaceae, in: Ettl, H., Gerloff, J., Heynig, H., Mollenhauser (Eds.), Süßwasserflora von Mitteleuropa. Gustav Fischer Verlag, Stuttgart, pp. 1–600.
- Krammer, K., Lange-Bertalot, H., 1991b. Bacillariophyceae 2/4. Achnanthaceae, Kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema, in: Ettl, H. (Ed.), Pascher's Süßwasserflora von Mitteleuropa, vol. 2, part 4. Gustav Fischer Verlag, Stuttgart, pp. 1–437.
- Krammer, K., 2000. Diatoms of Europe: Diatoms of European Inland Water and Comparable Habitats, Vol. 1: The genus Pinnularia. Edited by Horst Lange-Bertalot, Ruggell : Gantner,
- Krammer, K., 2002. Diatoms of Europe: Diatoms of European Inland Water and Comparable Habitats, Vol. 3: The genus Cymbella. Edited by Horst Lange-Bertalot, Ruggell : Gantner,
- Krammer, K., 2003. Diatoms of Europe: Diatoms of European Inland Water and Comparable Habitats, Vol. 4: Cymboplectra, Delicata, Navicymbula, Gomphocymbellopsis, Afrocymbell.. Edited by Horst Lange-Bertalot, Ruggell : Gantner,
- Lajoie, K. R., 1986. Coastal tectonics. Active tectonics, 95-124.
- Lange-Bertalot, H., 2000. Iconographia Diatomologica Annotated Diatom Micrographs – Diatom Flora of Marine Coasts I, Edited: H. Lange-Bertalot; Vol. 7 Pub: Koeltz Scientific Books. Copyright: A.R.G. Gantner Verlag K.G. By: Andrzej Witkowski, Horst Lange- Bertalot and Ditmar Metzeltin.
- Lange-Bertalot, H., 2001. Diatoms of Europe: Diatoms of European Inland Water and Comparable Habitats, Vol. 2: Navicula sensu stricto, 10 Genera Separated from Navicula sensu stricto, Frustulia. Edited by Horst Lange-Bertalot, Ruggell : Gantner.
- Lange-Bertalot, H., Bak, M., Witkowski, A., Tagliaventi, N., 2011. Diatoms of Europe: Diatoms of European Inland Water and Comparable Habitats, Vol. 6: Eunotia and some related genera. Edited by Horst Lange-Bertalot, Ruggell : Gantner.
- Leonard, L.J., Currie, C.A., Mazzotti, S., Hyndman, R.D., 2010. Rupture area and displacement of past Cascadia great earthquakes from coastal coseismic subsidence. Geological Society of America Bulletin, 112, 2079-2096.
- Levkov, Z., 2009. Diatoms of Europe: Diatoms of European Inland Water and Comparable Habitats, Vol. 5: Amphora sensu lato. Edited by Horst Lange-Bertalot, Ruggell : Gantner.
- Liu, K.-b, Fearn, M.L., 2000. Reconstruction of prehistoric landfall frequencies of catastrophic hurricanes in northwestern Florida from lake sediment records. Quat. Res. 54, 238–245.
- Lowe, R. L., 1974. Environmental Requirements and Pollution Tolerance of Freshwater Diatoms. EPA-670/4-74-005. Cincinnati, OH. 344 pp.
- Malik, J. N., Shishikura, M., Echigo, T., Ikeda, Y., Satake, K., Kayanne, H., Dikshit, O., 2011. Geologic evidence for two pre-2004 earthquakes during recent centuries near Port Blair, South Andaman Island, India. Geology, 39(6), 559-562.

- Mamo, B., Strotz, L., Dominey-Howes, D., 2009. Tsunami sediments and their foraminiferal assemblages. *Earth-Science Reviews*, 96, 263-278.
- Minoura, K., and Nakaya, S., 1991. Traces of tsunami preserved in inter-tidal lacustrine and marsh deposits – some examples from northeast Japan. *Journal of Geology*, 99, 265-287.
- Minoura, K., Nakaya, S., and Uchida, M., 1994. Tsunami deposits in a lacustrine sequence of the Sanriku coast, northeast Japan. *Sedimentary Geology*, 89(1), 25-31.
- Minoura, K., Imamura, F., Sugawara, D., Kono, Y., 2001. The 869 Jogan tsunami deposit and recurrence interval of large-scale tsunami on the Pacific coast of northeast Japan. *Journal of Natural Disaster Science*, 23, 83-88.
- Monecke, K., Finger, W., Klarer, D., Kongko, W., McAdoo, B.G., Moore, A.L., Sudrajat, S.U., 2008. A 1,000-year sediment record of tsunami recurrence in northern Sumatra. *Nature*, 455, 1232-1234.
- Morton, R. A., Gelfenbaum, G., Jaffe, B. E., 2007. Physical criteria for distinguishing sandy tsunami and storm deposits using modern examples. *Sedimentary Geology*, 200(3), 184-207.
- Murray, J. W., 1991. Ecology and palaeoecology of benthic foraminifera. Longman Scientific and Technical, Harlow, England. 397 pp.
- Nanayama, F., Shigeno, K., Satake, K., Shimokawa, K., Koitabashi, S., Miyasaka, S., Ishii, M., 2000. Sedimentary differences between the 1993 Hokkaido-nansei-oki tsunami and the 1959 Miyakojima typhoon at Taisei, southwestern Hokkaido, northern Japan. *Sedimentary Geology*, 135(1), 255-264.
- Nanayama, F., Satake, K., Furukawa, R., Shimokawa, K., Atwater, B.F., Shigeno, K., Yamaki, S., 2003. Unusually large earthquakes inferred from tsunami deposits along the Kuril Trench. *Nature*, 424, 660-663.
- Nanayama, F., Furukawa, R., Shigeno, K., Makino, A., Soeda, Y., Igarashi, Y., 2007. Nine unusually large tsunami deposits from the past 4000 years at Kiritappu marsh along the southern Kuril Trench. *Sedimentary Geology*, 200, 275-294.
- Nelson, A. R., 1992. Discordant 14 C ages from buried tidal-marsh soils in the Cascadia subduction zone, southern Oregon coast. *Quaternary Research*, 38(1), 74-90.
- Nelson, A.R., Kashima, K., 1993. Diatom zonation in southern Oregon tidal marshes relative to vascular plants, foraminifera, and sea level. *Journal of Coastal Research*, 9, 673-697.
- Nelson, A.R., Atwater, B.F., Bobrowsky, P.T., Bradley, L.A., Clague, J.J., Carver, G.A., Darienzo, M.E., Grant, W.C., Krueger, H.W., Sparks, R., Stafford, T.W., Stuiver, M., 1995. Radiocarbon evidence for extensive plate-boundary rupture about 300 years ago at the Cascadia subduction zone. *Nature* 378, 371-374.
- Nelson, A.R., Jennings, A.E., Kashima, K., 1996a. Holocene intertidal stratigraphy, microfossils, rapid submergence, and earthquake recurrence at Coos Bay, southern coastal Oregon, USA. *Geological Society of America Bulletin*, 108, 141-154.

1018 Nelson, A.R., Shennan, E., Long, A.J., 1996b. Identifying coseismic subsidence in tidal-wetland
1019 stratigraphic sequences at the Cascadia subduction zone of western North America. *Journal*
1020 *of Geophysical Research* 101, 6115-6135.
1021

1022 Nelson, A. R., Sawai, Y., Jennings, A. E., Bradley, L. A., Gerson, L., Sherrod, B. L., Sabeau, J.,
1023 Horton, B. P., 2008. Great-earthquake paleogeodesy and tsunamis of the past 2000 years at
1024 Alsea Bay, central Oregon coast, USA. *Quaternary Science Reviews*, 27(7), 747-768.
1025

1026 Nelson, A.R., Kashima, K., Bradley, L., 2009. Fragmentary evidence of great - earthquake
1027 subsidence during Holocene emergence, Valdivia Estuary, South Central Chile. *Bulletin of*
1028 *the Seismological Society of America*, 99. 71–86.
1029

1030 Nelson, A. R., 2013. Paleoseismology. *Encyclopedia of Natural Hazards*, 749-750.
1031

1032 Nelson, A.R., 2015. Coastal Sediments. in *Handbook of Sea-level Research*. Shennan, I., Long,
1033 A. J., Horton, B. P. (Eds.). John Wiley and Sons.
1034

1035 Nelson, A. R., Briggs, R. W., Dura, T., Engelhart, S. E., Gelfenbaum, G., Bradley, L. A.,
1036 Formans, S.L., Vane, C.V., Kelley, K. A., 2015. Tsunami recurrence in the eastern Alaska-
1037 Aleutian arc: A Holocene stratigraphic record from Chirikof Island,
1038 Alaska. *Geosphere*, 11(4), 1172-1203.
1039

1040 Nichol, S. L., Goff, J. R., Devoy, R. J. N., Chagué-Goff, C., Hayward, B., James, I., 2007.
1041 Lagoon subsidence and tsunami on the West Coast of New Zealand. *Sedimentary*
1042 *Geology*, 200, 248-262.
1043

1044 Palmer, A.J.M., Abbott, W.H., 1986. Diatoms as indicators of sea-level change. In: van de
1045 Plassche, O. (Ed.), *Sea-level Research: A Manual for the Collection and Evaluation of Data*.
1046 *Geobooks*, Norwich, pp. 457-488.
1047

1048 Pararas-Carayannis, G., 2003. Near and far-field effects of tsunamis generated by the
1049 PAroxyssmal eruptions, explosions, Caldera collapses and massive slope failures of the
1050 Krakatau volcano in Indonesia on August 26-27, 1883. *Science of Tsunami Hazards*.
1051

1052 Parsons, M.L., 1998. Salt marsh sedimentary record of the landfall of Hurricane Andrew on the
1053 Louisiana coast: diatoms and other paleoindicators. *Journal of Coastal Research* 14, 939–950.
1054

1055 Patrick, R. M., Reimer, C.W., 1966. The diatoms of the United States exclusive of Alaska and
1056 Hawaii, Volume 1: Monographs of the Academy of Natural Sciences of Philadelphia, 688.
1057

1058 Patrick, R. M., Reimer, C.W., 1975, The diatoms of the United States exclusive of Alaska and
1059 Hawaii, Volume 2: Monographs of the Academy of Natural Sciences of Philadelphia, 213.
1060

1061 Patrick, R., Williams, D.D., 1990. Aquatic biota in North America. In: Wolman, M.G., Riggs,
1062 H.D., (eds.), *Surface water hydrology: Boulder, Colorado, Geological Society of America,*
1063 *the Geology of North America*, v 0-1.
1064

1065 Patterson, R.T., Dalby, A.P., Roe, H.M., Guilbault, J.-P., Hutchinson, I., Clague, J.J., 2005.
1066 Relative utility of foraminifera, diatoms and macrophytes as high resolution indicators of
1067 paleo-sea level in coastal British Columbia, Canada. *Quaternary Science Reviews* 24, 2002-
1068 2014.

1069
1070 Peterson, C.D., Carver, G.A., Cruikshank, K.M., Abramson, H.F., Garrison-Laney, C.E.,
1071 Dengler, L.A., 2011. Evaluation of the use of paleotsunami deposits to reconstruct inundation
1072 distance and runup heights associated with prehistoric inundation events, Crescent City,
1073 southern Cascadia margin. *Earth Surface Processes and Landforms*, 36, 967-980.
1074
1075 Pilarczyk, J.E., and Reinhardt, E.G., 2012a. Testing foraminiferal taphonomy as a tsunami
1076 indicator in a shallow arid system lagoon: Sur, Sultanate of Oman. *Marine Geology*, 295–
1077 298, 128–136.
1078
1079 Pilarczyk, J.E., Horton, B.P., Witter, R.C., Vane, C.H., Chagué-Goff, C., Goff, J., 2012b.
1080 Sedimentary and foraminiferal evidence of the 2011 Tohoku-oki tsunami on the Sendai
1081 coastal plain, Japan. *Sedimentary Geology*, 282, 78-89.
1082
1083 Plafker, G., 1965. Tectonic deformation associated with the 1964 Alaska earthquake. *Science*,
1084 148, 1675-1687.
1085
1086 Plafker, G., 1969. Tectonics of the March 27, 1964, Alaska earthquake. U.S. Geological Survey
1087 Professional Paper 543-1, 1-74.
1088
1089 Plafker, G., 1972. Alaskan earthquake of 1964 and Chilean earthquake of 1960: Implications for
1090 arc tectonics. *Journal of Geophysical Research*, 77(5), 901-925.
1091
1092 Plafker, G., Savage, J. C., 1970. Mechanism of the Chilean earthquakes of May 21 and 22, 1960.
1093 *Geological Society of America Bulletin* 81, 1001–1030.
1094
1095 Plant, N.G., Stockdon, H.F., Sallenger, A.H., Turco, M.J., East, J.W., Taylor, A.A., Shaffer,
1096 W.A., 2010. Forecasting Hurricane impact on coastal topography. *Eos*, 91, 65-72.
1097
1098 Razjigaeva, N. G., Ganzey, L. A., Grebennikova, T. A., Ivanova, E. D., Kharlamov, A. A.,
1099 Kaistrenko, V. M., Shishkin, A. A., 2013. Coastal sedimentation associated with the Tohoku
1100 tsunami of 11 March 2011 in South Kuril Islands, NW Pacific Ocean. *Pure and Applied*
1101 *Geophysics*, 170(6-8), 1081-1102.
1102
1103 Round, F. E., Crawford, R. M., Mann, D. G., 1990. The diatoms: biology and morphology of the
1104 genera. Cambridge University Press.
1105
1106 Rhodes, B., Tuttle, M., Horton, B., Doner, L., Kelsey, H., Nelson, A., Cisternas, M., 2006.
1107 Paleotsunami research. *Eos*, 87, 205-209.
1108
1109 Riznyk, R. Z., 1973, Interstitial diatoms from two tidal flats in Yaquina Estuary, Oregon USA:
1110 *Botanica Marina*, 16, 113–138.
1111
1112 Sacchi, M., Insinga, D., Milia, A., Molisso, F., Raspini, A., Torrente, M. M., Conforti, A., 2005.
1113 Stratigraphic signature of the Vesuvius 79 AD event off the Sarno prodelta system, Naples
1114 Bay. *Marine Geology*, 222, 443-469.
1115
1116 Satake, K., Shimazaki, K., Tsuji, Y., Ueda, K., 1996. Time and size of a giant earthquake in
1117 Cascadia inferred from Japanese tsunami records of January 1700. *Nature*, 379(6562), 246-
1118 249.

- 1119
1120 Satake, K., 2007. Volcanic origin of the 1741 Oshima-Oshima tsunami in the Japan Sea. *Earth*
1121 *Planets Space*, 59, 381-390.
1122
- 1123 Satake, K., Atwater, B.F., 2007. Long-term perspectives on giant earthquakes and tsunamis at
1124 subduction zones. *Annual Review of Earth and Planetary Sciences*, 35, 349-374.
1125
- 1126 Savage, J. C., L. M. Hastie, 1966. Surface deformation associated with dip-slip faulting, *J.*
1127 *Geophys. Res.*, 71(20), 4897-4904.
1128
- 1129 Sawai, Y., 2001a. Episodic emergence in the past 3000 years at the Akkeshi Estuary, Hokkaido,
1130 Northern Japan. *Quaternary Research*, 56, 231-241.
1131
- 1132 Sawai Y., 2001b. Distribution of living and dead diatoms in tidal wetlands of northern Japan:
1133 relations to taphonomy. *Palaeogeography Palaeoclimatology Palaeoecology*, 173, 125-141.
1134
- 1135 Sawai, Y., Nasu, H., Yasuda, Y., 2002. Fluctuations in relative sea-level during the past 3000 yr
1136 in the Onnetoh estuary, Hokkaido, northern Japan. *Journal of Quaternary Science*, 17, 607-
1137 622.
1138
- 1139 Sawai, Y., Satake, K., Takanobu, K., Nasu, H., Shishikura, M., Atwater, B.F., Horton, B.P.,
1140 Kelsey, H.M., Nagumo, T., Yamaguchi, M., 2004a. Transient uplift after a 17th-century
1141 earthquake along the Kuril subduction zone. *Science* 306, 1918-1920.
1142
- 1143 Sawai, Y., Horton, B.P., Nagumo, T., 2004b. The development of a diatom-based transfer
1144 function along the Pacific coast of eastern Hokkaido, northern Japan – an aid in paleoseismic
1145 studies of the Kuril subduction zone. *Quaternary Science Reviews* 23, 2467-2483.
1146
- 1147 Sawai, Y., Fujii, Y., Fujiwara, O., Kamataki, T., Komatsubara, J., Okamura, Y., Satake, K.,
1148 Shishikura, M., 2008. Marine incursions of the past 1500 years and evidence of tsunamis at
1149 the Suijin-numa, a coastal lake facing the Japan Trench. *The Holocene*, 18, 517-528.
1150
- 1151 Sawai, Y., Jankaew, K., Martin, M.E., Prendergast, A., Choowong, M., Charoentitirat, T., 2009a.
1152 Diatom assemblages in tsunami deposits associated with the 2004 Indian Ocean tsunami at
1153 Phra Thong Island, Thailand. *Marine Micropaleontology*, 73, 70-79.
1154
- 1155 Sawai, Y., Kamataki, T., Shishikura, M., Nasu, H., Okamura, Y., Satake, K., Thomson, K.H.,
1156 Matsumoto, D., Fujii, Y., Komatsubara, J., Aung, T.T., 2009b. Aperiodic recurrence of
1157 geologically recorded tsunamis during the past 5500 years in eastern Hokkaido, Japan .
1158 *Journal of Geophysical Research*, 114, B01319, doi:10.1029/2007JB005503
1159
- 1160 Sawai, Y., Namegaya, Y., Okamura, Y., Satake, K., Shishikura, M., 2012. Challenges of
1161 anticipating the 2011 Tohoku earthquake and tsunami using coastal geology. *Geophysical*
1162 *Research Letters*, 39, L21309, doi: 10.1029/2012GL053692
1163
- 1164 Sawai, Y., Namegaya, Y., Tamura, T., Nakashima, R., Tanigawa, K., 2015. Shorter intervals
1165 between great earthquakes near Sendai: Scour ponds and a sand layer attributable to AD 1454
1166 overwash. *Geophysical Research Letters*, 42(12), 4795-4800.
1167

- Shennan, I., Long, A.J., Rutherford, M.M., Green, F.M., Innes, J.B., Lloyd, J.M., Zong, Y., Walker, K.J., 1996. Tidal marsh stratigraphy, sea level change and large earthquakes: I. A 5000 year record in Washington. *Quaternary Science Reviews*, 15, 1023-1059.
- Shennan, I., Long, A.J., Rutherford, M.M., Innes, J.B., Green, F.M., Walker, K.J., 1998. Tidal marsh stratigraphy, sea-level change and large earthquakes: II. Submergence events during the last 3500 years at Netarts Bay, Oregon. *Quaternary Science Reviews* 17, 365-393.
- Shennan, I., Scott, D.B., Rutherford, M.M., Zong, Y., 1999. Microfossil analysis of sediments representing the 1964 earthquake, exposed at Girdwood Flats, Alaska. *Quaternary International*, 60, 55-73.
- Shennan, I., Hamilton, S., 2006. Coseismic and preseismic subsidence associated with great earthquakes in Alaska. *Quaternary Science Reviews* 25, 1-8.
- Shennan, I., Bruhn, R., Plafker, G., 2009. Multi-segment earthquakes and tsunami potential of the Aleutian megathrust. *Quaternary Science Reviews*, 28, 7-13.
- Shennan, I., Bruhn, R., Barlow, N., Good, K., Hocking, E., 2014a. Late Holocene great earthquakes in the eastern part of the Aleutian megathrust. *Quaternary Science Reviews*, 84, 86-97.
- Shennan, I., Barlow, N., Carver, G., Davies, F., Garrett, E., Hocking, E., 2014b. Great tsunamigenic earthquakes during the past 1000 yr on the Alaska megathrust. *Geology*, 42, 687-690.
- Sherrod, B.L., 1999. Gradient analysis of diatom assemblages in a Puget Sound salt marsh: can such assemblages be used for quantitative paleoecological reconstructions? *Palaeogeography, Palaeoclimatology, Palaeoecology*, 149, 213-226.
- Sherrod, B. L., 2001. Evidence for earthquake-induced subsidence about 1100 yr ago in coastal marshes of southern Puget Sound, Washington. *Geological Society of America Bulletin*, 113(10), 1299-1311.
- Sherrod, B. L., Bucknam, R. C., Leopold, E. B., 2000. Holocene relative sea level changes along the Seattle Fault at Restoration Point, Washington. *Quaternary Research*, 54(3), 384-393.
- Shimazaki, K., Kim, H. Y., Chiba, T., Satake, K., 2011. Geological evidence of recurrent great Kanto earthquakes at the Miura Peninsula, Japan. *Journal of Geophysical Research: Solid Earth* (1978–2012), 116(B12).
- Simons, M., Minson, S.E., Sladen, A., Ortega, F., Jiang, J., Owen, S.E., Meng, L., Ampuero, J-P., Wei, S., Chu, R., Helmberger, D.V., Kanamori, H., Hetland, E., Moore, A.W., Webb, F.H. 2011. The 2011 magnitude 9.0 Tohoku-oki earthquake: mosaicking the megathrust from seconds to centuries. *Science* 332, 1421-1425.
- Small, C., Gornitz, V., Cohen, J.E., 2000. Coastal hazards and the global distribution of human population. *Environmental Geosciences*, 7, 3-12.

- 1217 Smith, D.E., Shi, S., Cullingford, R.A., Dawson, A.G., Dawson, S., Firth, C.R., Foster, I.D.L.,
1218 Fretwell, P.T., Haggart, B.A., Holloway, L.K., Long, D., 2004. The Holocene Storegga slide
1219 tsunami in the United Kingdom. *Quaternary Science Reviews* 23, 2291-2321.
1220
- 1221 Stein, S., Okal, E.A., 2007. Ultralong period seismic study of the December 2004 Indian Ocean
1222 earthquake and implications for regional tectonics and the subduction process. *Bulletin of the*
1223 *Seismological Society of America*, 97, S279-S295.
1224
- 1225 Stein, S., Okal, E. A., 2011. The size of the 2011 Tohoku earthquake need not have been a
1226 surprise. *Eos Transactions of the American Geophysical Union*, 92, 227.
1227
- 1228 Stewart, I. S., Vita-Finzi, C. (Eds.), 1998. *Coastal tectonics*. Geological Society of London.
1229
- 1230 Sugawara, D., Minoura, K., Nemoto, N., Tsukawaki, S., Goto, K., Imamura, F., 2009.
1231 Foraminiferal evidence of submarine sediment transport and deposition by backwash during
1232 the 2004 Indian Ocean tsunami. *Island Arc*, 18(3), 513-525.
1233
- 1234 Switzer, A. D., Jones, B. G., 2008. Large-scale washover sedimentation in a freshwater lagoon
1235 from the southeast Australian coast: sea-level change, tsunami or exceptionally large
1236 storm?. *The Holocene*, 18(5), 787-803.
1237
- 1238 Szczuciński, W., 2012a. The post-depositional changes of the onshore 2004 tsunami deposits on
1239 the Andaman Sea coast of Thailand. *Natural Hazards*, 60, 115-133.
1240
- 1241 Szczuciński, W., M. Kokociński, M. Rzeszewski, C. Chagué-Goff, M. Cachão, K. Goto,
1242 Sugawara, D., 2012b. Sediment sources and sedimentation processes of 2011 Tohoku-oki
1243 tsunami deposits on the Sendai Plain, Japan — Insights from diatoms, nannoliths and grain
1244 size distribution. *Sedimentary Geology*, 282, 40–56.
1245
- 1246 Szkornik, K., Gehrels, W. R., Kirby, J. R., 2006. Using salt-marsh diatoms to reconstruct sea-
1247 level changes in Ho Bugt western Denmark. *Marine Geology*, 235, 137–50.
1248
- 1249 Takashimizu, Y., Urabe, A., Suzuki, K., Sato, Y., 2012. Deposition by the 2011 Tohoku-oki
1250 tsunami on coastal lowland controlled by beach ridges near Sendai, Japan. *Sedimentary*
1251 *Geology*, 282, 124-141.
1252
- 1253 Telford, R.J., Birks, H.J.B., 2011a. A novel method for assessing the statistical significance of
1254 quantitative reconstructions inferred from biotic assemblages. *Quaternary Science*
1255 *Reviews*, 30, 1272-1278.
1256
- 1257 Telford, R.J., Birks, H.J.B., 2011. Effect of uneven sampling along an environmental gradient on
1258 transfer-function performance. *Journal of Paleolimnology*, 46, 99-106.
1259
- 1260 Tuttle, M. P., Ruffman, A., Anderson, T., Jeter, H., 2004. Distinguishing tsunami from storm
1261 deposits in eastern North America: the 1929 Grand Banks tsunami versus the 1991
1262 Halloween storm. *Seismological Research Letters*, 75 (1), 117-131.
1263
- 1264 Van Dam, H., Mertens, A., and Sinkeldam, J., 1994. A coded checklist and ecological indicator
1265 values of freshwater diatoms from the Netherlands. *Netherlands Journal of Aquatic*
1266 *Ecology*, 28(1), 117-133.

1267
1268 van de Plassche, O., Erkens, G., van Vliet, F., Brandsma, J., van der Borg, K., de Jong, A.F.M.,
1269 2006. Salt-marsh erosion associated with hurricane landfall in southern New England in the
1270 fifteenth and seventeenth centuries. *Geology*, 34, 829-832.
1271
1272 van der Werff, H., Huls, H., 1958–1974. Diatomeeenflora van Nederland, 8 parts, published
1273 privately, De Hoef, The Netherlands.
1274
1275 Vos, P.C., 1986. De sediment stabiliserende werking van benthische diatomeeen in het
1276 intergetijde gebied van de Oosterschelde . GEOMOR-nota 86-03, Rijkswaterstaat, Dienst
1277 Getijde-wateren, Middelburg, 156 pp.
1278
1279 Vos, P.C., de Boer, P.L., Misdorp, R., 1988. Sediment stabilization by benthic diatoms in
1280 intertidal sandy shoals, in: P. L. de Boer, A. van Gelder and S . D . Nio (Eds), Tide influenced
1281 sedimentary environments and facies, D.Reidel Publishing Company, Dordrecht, pp. 511-
1282 526.
1283
1284 Vos, P.C., de Wolf, H., 1988. Methodological aspects of paleo-ecological diatom research in
1285 coastal areas of the Netherlands. *Geol Mijnbouw* 67: 31–40.
1286
1287 Vos, P.C., de Wolf, H., 1993. Diatoms as a tool for reconstruction sedimentary environments in
1288 coastal wetlands: methodological aspects. *Hydrobiologia* 269–270, pp.
1289
1290 Wang, K., Hu, Y., and He, J., 2012. Deformation cycles of subduction earthquakes in a
1291 viscoelastic Earth. *Nature*, 484 (7394), 327-332.
1292
1293 Wang, P. L., Engelhart, S. E., Wang, K., Hawkes, A. D., Horton, B. P., Nelson, A. R., Witter, R.
1294 C., 2013. Heterogeneous rupture in the great Cascadia earthquake of 1700 inferred from
1295 coastal subsidence estimates. *Journal of Geophysical Research: Solid Earth*, 118(5), 2460-
1296 2473.
1297
1298 Watcham, E.P., Shennan, I., Barlow, N.L.M., 2013. Scale considerations in using diatoms as
1299 indicators of sea-level change: lessons from Alaska. *Quaternary Science Reviews* 28, 165-
1300 179.
1301
1302 Whiting, M. C., McIntire, C. D., 1985. An investigation of distributional patterns in the diatom
1303 flora of Netarts Bay, Oregon, by correspondence analysis1. *Journal of Phycology*, 21, 655-
1304 661.
1305
1306 Williams, H., Hutchinson, I., 2000. Stratigraphic and microfossil evidence for late Holocene
1307 tsunamis at Swantown marsh, Whidbey Island, Washington. *Quaternary Research*, 54(2),
1308 218-227.
1309
1310 Williams, H.F., Hutchinson, I., Nelson, A.R., 2005. Multiple sources for late-Holocene tsunamis
1311 at Discovery Bay, Washington State, USA. *The Holocene*, 15, 60-73.
1312
1313 Wilson, G. P., Lamb, A. L., 2012. An assessment of the utility of regional diatom-based tidal-
1314 level transfer functions. *Journal of Quaternary Science*, 27(4), 360-370.
1315

- Wilson, R., Hemphill-Haley, E., Jaffe, B., Richmond, B., Peters, R., Graehl, N., Kelsey, H., Leeper, R., Watt, S., McGann, M., Hoirup, D., Chague-Goff, C., Goff, J., Caldwell, D., Looftbourrow, C., 2014. The search for geologic evidence of distant-source tsunamis using new field data in California, chap. C of Ross, S.L., and Jones, L.M., eds., *The SAFRR (Science Application for Risk Reduction) tsunami scenario: U.S. Geological Survey Open-File Report 2013–1170*, 214 p., <http://dx.doi.org/10.3115/20131170/c/>.
- Witter, R.C., Kelsey, H.M., Hemphill-Haley, E., 2001. Pacific storms, El Nino and tsunamis: competing mechanisms for sand deposition in a coastal marsh, Euchre Creek, Oregon. *J. Coast. Res.* 563-583.
- Witter, R.C., Kelsey, H.M., Hemphill-Haley, E., 2003. Great Cascadia earthquakes and tsunamis of the past 6700 years, Coquille River estuary, southern coastal Oregon. *Geological Society of America Bulletin* 115,1289-1306.
- Witter, R. C., Hemphill-Haley, E., Hart, R., Gay, L., 2009. Tracking Prehistoric Cascadia Tsunami Deposits at Nestucca Bay, Oregon. US Geological Survey, National Earthquake Hazards Reduction Program Final Technical Report 08HQGR0076, 92.
- Woodruff, J.D., Donnelly, J.P., Okusu, A., 2009. Exploring typhoon variability over the mid-to-late Holocene: evidence of extreme coastal flooding from Kamikoshiki, Japan. *Quaternary Science Reviews* 28, 1774-1785.
- Woodroffe, S.A., Long, A.J., 2010. Reconstructing recent relative sea-level changes in West Greenland: Local diatom-based transfer functions are superior to regional models. *Quaternary International*, 221, 91-103.
- Wright, C., and Mella, A., 1963. Modifications to the soil pattern of South-Central Chile resulting from seismic and associated phenomena during the period May to August 1960. *Bulletin of the Seismological Society of America*, 53(6), 1367-1402.
- Yeats, R.S., Sieh, K.E., Allen, C.R., 1997. *The geology of earthquakes*. Oxford University Press, Oxford, 568 pp.
- Yoshida, T. Jogan Ju'uichi Nen Mutsu Fujou no Shindou Kouitsu [Quake and flood in 11th year of Jogan era] *Rekishi Chiri* 8, 1033-1040 (1906) [in Japanese]
- Zong, Y.Q., Tooley, M.J., 1996. Holocene sea-level changes and crustal movements in Morecambe Bay, northwest England. *Journal of Quaternary Science* 11, 43-58.
- Zong, Y., 1997. Implications of *Paralia sulcata* abundance in Scottish isolation basins. *Diatom Research*, 12, 125-150.
- Zong, Y., Horton, B. P., 1998. Diatom zones across intertidal flats and coastal saltmarshes in Britain. *Diatom Research*, 13, 375-394.
- Zong, Y.Q., Horton, B.P., 1999. Diatom-based tidal-level transfer functions as an aid in reconstructing Quaternary history of sea-level movements in the UK. *Journal of Quaternary Science*, 14, 153–167.

1366 Zong, Y.Q., Shennan, I., Combellick, R.A., Hamilton, S.L., Rutherford, M.M., 2003. Microfossil
1367 evidence for land movements associated with the 1964 AD Alaska earthquake. *Holocene*, 13,
1368 7-20.
1369

1370 **Figure captions**

1371 **Figure 1:** Schematic drawing of coseismic uplift (a) and subsidence (b) and
1372 accompanying tsunami inundation.

1373

1374 **Figure 2:** Summary of earthquake and tsunami studies applying diatoms that are
1375 referenced in this paper, and listed in Table 1. (a) In: Indian Ocean; (b) Ja: Japan Trench;
1376 (c) Al: Alaska-Aleutian megathrust; (d) Ca: Cascadia subduction zone; (e) Ch: Chilean
1377 Subduction zone; (f) NZ: New Zealand; NS: North Sea (studies marked with an asterisk
1378 because they are not subduction zone studies); SP: South Pacific; PNG: Papua New
1379 Guinea.

1380

1381 **Figure 3:** Qualitative schematic of the distribution of common diatom species along a
1382 modern salt marsh transect. Typical salinity classes (following the halobian classification
1383 scheme of Hemphill-Haley, 1993) for each environment are shown.

1384

1385 **Figure 4:** Diatoms evaluated relative to modern intertidal zones and stratigraphy at the
1386 Niawiakum River in Washington State, USA. (a) Position of the study area relative to the
1387 Cascadia subduction zone (barbed line), the boundary between North America and Juan
1388 de Fuca/Gorda plates that extends from the northern end of the San Andreas Fault (SAF)
1389 to the southern end of the Queen Charlotte Fault (QCF); (b) Location of the Niawiakum
1390 River Valley, on the eastern side of Willapa Bay; (c) Locations of vertical sections

sampled for diatoms in the Niawiakum River Valley. Sites 1-4 are cutbank outcrops exposed during low tide; (d) Changes in diatom assemblages within and above a former upland soil (forming at the transition of extreme high water (EHW) and upland) buried by coseismic subsidence during a Cascadia subduction zone earthquake in AD 1700. Changes in diatom assemblages are consistent with an abrupt change from upland forest to tidal flat or low marsh. (Modified from Hemphill-Haley, 1995a).

Figure 5: Diatom analyses during past earthquake cycles on the Alaska-Aleutian megathrust suggesting preseismic movement. (a) Location of south-central Alaska, USA; (b) area subsided in the AD 1964 earthquake (Plafker, 1969); (c) Relative sea level (RSL) changes reconstructed using a diatom-based transfer function; two short periods of preseismic submergence immediately prior to substantial coseismic subsidence were recognized at the top of peat units and are highlighted with red arrows. (Modified and reprinted from Shennan, I., Hamilton, S., Coseismic and preseismic subsidence associated with great earthquakes in Alaska. *Quaternary Science Reviews* 25, 1-8, 2006, with permission from Elsevier).

Figure 6: Example of land-level reconstructions using diatoms in Hokkaido, northern Japan. (a) Position of study area relative to the Kurile Trench. The solid line with triangles shows the seaward edge of the subduction zone. The volcanoes responsible for tephra layers in c and d are shown and rupture areas of instrumentally recorded earthquakes on the plate boundary off eastern Hokkaido are outlined; (b) Map of Mochirippu Estuary showing location of stratigraphic cross-section; (c) Stratigraphic

cross-section; (d) Photograph and log of stratigraphy. Example of change from tidal-flat mud to lowland-forest peat, punctuated by a tsunami deposit and by volcanic ash layers. (e) Diatom diagram showing the schematic stratigraphy, changes in diatom assemblages, and the results of diatom-based transfer functions showing the seventeenth-century large earthquake. Error bars for height estimates span two standard deviations. (Modified and reprinted from Sawai, Y., Satake, K., Takanobu, K., Nasu, H., Shishikura, M., Atwater, B.F., Horton, B.P., Kelsey, H.M., Nagumo, T., Yamaguchi, M., Transient uplift after a 17th-century earthquake along the Kuril subduction zone. *Science* 306, 1918-1920, 2006, with permission from The American Association for the Advancement of Science).

Figure 7: Diatom analyses of the Storegga Slide tsunami showing fragmentation of diatom valves within the high-energy deposit. (a) Location of the Storegga Slides and sites where evidence for the Holocene Storegga Slide tsunami has been found; (b) Sites in the United Kingdom where evidence for the Holocene Storegga Slide tsunami may be found. Numbers correspond to sites discussed in Smith et al., 2004; (c) Diatom summary diagram from Boreholes 53 and 23 in lower Wick River Valley, Caithness, Scotland (Dawson et al., 1996). Taxa displayed as % of total valves. (Modified and reprinted from Smith, D.E., Shi, S., Cullingford, R.A., Dawson, A.G., Dawson, S., Firth, C.R., Foster, I.D.L., Fretwell, P.T., Haggart, B.A., Holloway, L.K., Long, D., The Holocene Storegga slide tsunami in the United Kingdom. *Quaternary Science Reviews* 23, 2291-2321, 2004, and Dawson, S., Smith, D.E., Ruffman, A., Shi, S., The diatom biostratigraphy of tsunami deposits: Examples from recent and middle Holocene events. *Physics and Chemistry of the Earth*, 21, 87-92, 1996, with permission from Elsevier).

1437

1438 **Figure 8:** Diatom analyses of the 2004 Indian Ocean tsunami deposit showing grading of
1439 diatom valves; (a) Position of study area relative to the Sunda Trench. Fault slip during
1440 the 2004 Sumatra–Andaman earthquake (Chlieh et al., 2007); (b) Phra Thong Island. The
1441 island is isolated from the mainland of Thailand by inlets. Light gray area is grassy beach
1442 ridge plains. Dark gray area is mangrove forests. Landforms traced from 1:50,000-scale
1443 airphotos taken in 1999 and from post-tsunami satellite images at PointAsia.com
1444 (modified from Jankaew et al., 2008); (c) Location of pit and modern samples. Satellite
1445 image is from PointAsia.com; (d) Diatom diagram showing the schematic stratigraphy of
1446 location 6 and changes in diatom assemblages throughout the 2004 tsunami deposit.
1447 Abundant beach and subtidal species are more abundant in the bottom of the sand bed
1448 and freshwater diatoms become more abundant in the top of the sand bed. Diatom valves
1449 are relatively pristine in the tsunami deposit as shown by the high percentage of unbroken
1450 valves. (e) Simplified process of deposition of diatoms and sediment during tsunami; (1)
1451 Fast current. Only beach and subtidal species are incorporated with coarse sediment.
1452 Because turbulent current can keep a substantial amount of sand fraction in the water
1453 column, mixed assemblages of many beach and subtidal, marine plankton are suspended.
1454 Freshwater specimens may be included with eroded soil fractions. (2) Current becomes
1455 slow. Fine fractions fall onto the ground. Eroded, floated, and transported specimens are
1456 also incorporated. (3) Suspension stage (calm current) of tsunami. All floated specimens
1457 are allowed to settle down. Many freshwater species incorporated with their substrata
1458 (plant trash and eroded soil fractions). (Modified and reprinted from Sawai, Y., Jankaew,
1459 K., Martin, M.E., Prendergast, A., Choowong, M., Charoentitirat, T., Diatom

1460 assemblages in tsunami deposits associated with the 2004 Indian Ocean tsunami at Phra
1461 Thong Island, Thailand. *Marine Micropaleontology*, 73, 70–79, 2009, with permission
1462 from Elsevier).

1463

Figure 1

A map of the Pacific Ocean region, including parts of North America, South America, and Australia. The map shows several sampling locations marked with red dots and labeled: NS (North of South America), NS-3°, NS-6°, NS-2,4°, NS-1,4°, NS-1,5°, PNG-1 (near Papua New Guinea), SP-2, and SP-1 (near the Solomon Islands). Six rectangular boxes labeled a through f indicate specific areas of interest: box 'a' is in the western Pacific near Indonesia; box 'b' is in the western Pacific near Japan; box 'c' is in the northern Pacific near Alaska; box 'd' is in the northern Pacific near the US coast; box 'e' is in the southern Pacific near Chile; and box 'f' is in the southern Pacific near New Zealand. Dashed lines represent transects, with one prominent transect running from the western Pacific towards the central Pacific, and another running from the northern Pacific towards the southern Pacific.

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Table 1. A global database of subduction zone earthquake and tsunami studies that have employed diatoms

Map code	Earthquake and tsunami studies employing diatoms	Diatom evidence for coseismic land-level change ¹	Accompanying tsunami ²	Diatom composition of tsunami deposit ³	Preservation of diatom valves ⁴	Concentration of diatom valves ⁵
Alaska						
Al-1	Shennan et al., 1999	Yes (H, Qn)	--	--	--	--
Al-2	Zong et al., 2003	Yes (H, Qn)	--	--	--	--
Al-3	Hamilton et al., 2005	Yes (H, P, Qn)	--	--	--	--
Al-4	Hamilton and Shennan, 2005a	Yes (H, P, Qn)	--	--	--	--
Al-5	Hamilton and Shennan, 2005b	Yes (H, P, Qn)	--	--	--	--
Al-6	Shennan and Hamilton, 2006	Yes (H, P, Qn)	--	--	--	--
Al-7	Watcham et al., 2013	Yes (H, P, Qn)	--	--	--	--
Al-8	Shennan et al., 2009	Yes (H, P, Ql)	--	--	--	--
Al-9	Shennan et al., 2014a	Yes (H, P, Qn)	--	--	--	--
Al-10	Shennan et al., 2014b	Yes (H, P, Qn)	--	--	--	--
Al-11	Briggs et al., 2014	Yes (H, P, Ql)	Yes	Mx	Good	High
Al-12	Nelson et al., 2015	No	Yes (H, P)	Fw	Poor	Low
Cascadia						
Ca-1	Dariento and Peterson, 1990	Yes (P, Ql)	Yes (P)	M&B	--	--
Ca-2	Dariento et al., 1994	Yes (P, Ql)	Yes (P)	M&B	--	--
Ca-3	Hemphill-Haley, 1995a	Yes (P, Ql)	Yes (P)	M&B (Ep)	Very good	--
Ca-4	Hemphill-Haley, 1996	Yes (P, Ql)	Yes (P)	M&B (Ep, Pl)	Very good	High
Ca-5	Shennan et al., 1996	Yes (P, Qn)	Yes (P)	M&B (Ep)	--	--
Ca-6	Nelson et al., 1996b	Yes (P, Qn)	No	--	--	--
Ca-7	Atwater and Hemphill-Haley, 1997	Yes (P, Ql)	Yes (P)	M&B (Ep)	Very good	--
Ca-8	Benson et al., 1997	No	Yes (H, P)	M	--	--
Ca-9	Hutchinson et al., 1997	No	Yes (P)	M&B (Ep)	--	--
Ca-10	Clague et al., 1999	Yes (P, Ql)	Yes (P)	M&B	Good	--
Ca-11	Hutchinson et al., 2000	Yes (P, Ql)	Yes (P)	Mx	Good	Low
Ca-12	Sherrod et al., 2000	Yes (P)	No	--	--	--
Ca-13	Williams and Hutchinson, 2000	No	Yes (P)	M&B	Poor	Low
Ca-14	Sherrod, 2001	Yes (P, Ql)	No	--	--	--
Ca-15	Kelsey et al., 2002	Yes (P, Ql)	Yes (P)	M&B (Ep?)	Good	Low
Ca-16	Kelsey et al., 2004	Yes (P, Ql)	No	--	--	--
Ca-17	Witter et al., 2003	Yes (P, Ql)	Yes (P)	M (Ep)	--	Low
Ca-18	Kelsey et al., 2005	No	Yes (P)	few M	--	Low
Ca-19	Williams et al., 2005	Yes (P, Ql)	Yes (P)	M (Ep, Pl)	Good	High
Ca-20	Nelson et al., 2008	Yes (P, Qn)	Yes (P)	M&B (Ep, Pl)	--	Low
Ca-21	Witter et al., 2009	Yes (P, Ql)	Yes (P)	M (Ep)	Poor	High
Ca-22	Peterson et al., 2011	No	Yes (P)	M&B	--	--
Ca-23	Graehl et al., 2014	Yes (P, Ql)	Yes (P)	M&B (Ep)	Good	Low
Ca-24	Wilson et al., 2014	Yes (P, Ql)	Yes (H, P)	Mx (Pl, G)	Good	High
Chile						
Ch-1	Cisternas et al., 2005	Yes (H, P, Ql)	Yes (H, P)	--	--	--
Ch-2	Nelson et al., 2009	Yes (H, P, Qn)	Yes (P)	Mx	--	--
Ch-3	Horton et al., 2011	No	Yes (H)	M&B (Ep, Pl)	Selective	High
Ch-4	Garrett et al., 2013	Yes (H, Qn)	Yes (H)	Mx (Ep)	--	--
Ch-5	Dura et al., 2014	Yes (P, Ql)	Yes (P)	M&B (Pl, G)	Poor	Low
Ch-6	Garrett et al., 2014	Yes (H, P, Qn)	Yes (H, P)	M&B (Ep)	--	--
Ch-7	Chagué-Goff et al., 2015	No	Yes (H)	M&B	Poor	Low
Indian Ocean						
In-1	Jankaew et al., 2008	No	Yes (H, P)	M&B	Good (H)/Poor (P)	High (H)/Low (P)
In-2	Monecke et al., 2008	No	Yes (H, P)	Fw&B	Good (H)/Poor (P)	High (H)/Low (P)
In-3	Sawai et al., 2009a	No	Yes (H)	Mx (Pl, G)	Good	--
In-4	Kokociński et al., 2009	No	Yes (H)	Mx (Ep, Pl)	Poor	Low