Expanding the Role of Reactive Transport Models in Critical Zone Processes

¹Li Li, ²Kate Maher, ³Alexis Navarre-Sitchler, ⁴Jenny Druhan, ⁵Christof Meile, ⁶Corey Lawrence, ⁷Joel Moore, ⁸Julia Perdrial, ⁹Pamela Sullivan, ¹⁰Aaron Thompson, ¹¹Lixin Jin, ¹²Edward W. Bolton, ¹³Susan L. Brantley, ¹⁴William E. Dietrich, ¹⁵K. Ulrich Mayer, ¹⁶Carl I. Steefel, ¹⁷Albert Valocchi, ¹⁸John Zachara, ¹⁹Benjamin Kocar, ²⁰Jennifer McIntosh, ¹Chen Bao, ²¹Benjamin M. Tutolo, ³Joe Beisman, ²²Mukesh Kumar, ¹⁶Eric Sonnenthal

¹Department of Civil and Environmental Engineering, The Pennsylvania State University, University Park, PA 16802

²Department of Geological & Environmental Sciences, Stanford University, CA 94305-2115

³Department of Geology and Geological Engineering, Colorado School of Mines, Golden, CO 80401

⁴Department of Geology, University of Illinois at Urbana Champaign, Urbana, IL 61801

⁵Department of Marine Sciences, The University of Georgia, Athens, GA 30602

⁶U.S. Geological Survey, MS 980, Box 25046, Federal Center, Denver, CO 80225

⁷Department of Physics, Astronomy, & Geosciences, Towson University, Towson, MD 21252

⁸Department of Geology, University of Vermont, Burlington, VT 05405

⁹Department of Geography, University of Kansas, Lawrence, KS 66045-7613

¹⁰Department of Crop and Soil Sciences, University of Georgia, Athens, GA 30602

¹¹Department of Geological Sciences, University of Texas at El Paso, Texas 79968

¹²Department of Geology and Geophysics, Yale University, New Haven CT 06520-8109

¹³Earth and Environmental Systems Institute and the Department of Geosciences, The Pennsylvania State University, University Park, PA 16802

¹⁴Department of Earth and Planetary Science, University of California Berkeley, Berkeley, CA 94720

¹⁵Department of Earth, Ocean and Atmospheric Sciences, Vancouver, BC Canada V6T 1Z4

¹⁶Earth Sciences Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Mail Stop 90-1116, Berkeley, CA 94720

¹⁷Department of Civil and Environmental Engineering, University of Illinois at Urbana Champaign, Urbana, IL 61801

¹⁸Physical Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99352

¹⁹Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA 0213

²⁰Department of Hydrology and Atmospheric Sciences, University of Arizona, Tucson, Arizona 85716

²¹Department of Earth Sciences, University of Oxford, Oxford, UK OX1 3AN

²²Nicholas School of the Environment, Duke University, Durham, NC 27708-0328

Abstract:

Models test our understanding of processes and can reach beyond the spatial and temporal scales of measurements. Multi-component Reactive Transport Models (RTMs), initially developed more than three decades ago, have been used extensively to explore the interactions of geothermal, hydrologic, geochemical, and geobiological processes in subsurface systems. Driven by extensive data sets now available from intensive measurements efforts, there is a pressing need to couple RTMs with other community models to explore non-linear interactions among the atmosphere, hydrosphere, biosphere, and geosphere. Here we briefly review the history of RTM development, summarize the current state of RTM approaches, and identify new research directions, opportunities, and infrastructure needs to broaden the use of RTMs. In particular, we envision the expanded use of RTMs in advancing process understanding in the Critical Zone, the veneer of the Earth that extends from the top of vegetation to the bottom of groundwater. We argue that, although parsimonious models are essential at larger scales, process-based models are the only way to explore the highly nonlinear coupling that characterizes natural systems. We present seven testable hypotheses that emphasize the unique capabilities of process-based RTMs for (1) elucidating chemical weathering and its physical and biogeochemical drivers; (2) understanding the interactions among roots, micro-organisms, carbon, water, and minerals in the rhizosphere; (3) assessing the effects of heterogeneity across spatial and temporal scales; and (4) integrating the vast quantity of novel data, including “omics” data (genomics, transcriptomics, proteomics, metabolomics), elemental concentration and speciation data, and isotope data into our understanding of complex earth systems. With strong support from data-driven sciences, we are now in an exciting era where integration of RTM framework into other community models will facilitate process understanding across disciplines and across scales.

1. Reactive Transport Models as a Tool for Integration

Reactive transport models (RTMs) quantitatively evaluate coupling of the physical, chemical, and biological processes that determine the formation, operation, and evolution of Earth surface systems. The coupling among processes plays a pivotal role in governing low-temperature biogeochemical systems, including the conversion of rock to soil, the release of nutrients from minerals for ecosystem vitality, the transport and bioremediation of reactive contaminants, and the biogeochemical cycling of important elements. RTMs are process-based models that allow both the integration and differentiation of distinct processes, thus providing an objective way to assess system response to external forcings. RTMs can be applied to optimize sampling campaigns; serve as a repository of knowledge regarding mathematical frameworks; and carry out numerical experiments under conditions where data are not available, thereby expanding forecasting capabilities beyond measurement limitations (assuming the forecast uncertainty can be addressed). From a practical point of view, RTMs could support decision making, risk assessment, and environmental stewardship, especially if data science approaches that address model sensitivity and uncertainty are developed.

Our goal here is to paint a broader picture of how RTMs could be useful to addressing some outstanding biogeochemical questions, rather than discuss issues and methods. We aim to identify opportunities to integrate RTM framework into other relevant disciplines to advance process understanding, with the primary focus on the low-temperature ($< 100^{\circ}\text{C}$) surface and near-surface environment termed the Critical Zone (CZ). The CZ is defined as the zone “from the outer extent of vegetation down to the lower limits of groundwater” (Brantley et al., 2007). We argue that RTMs can play an important role in understanding interactions between water, atmosphere, rock, soil, and life within the CZ. These interactions create complex coupling between chemical, physical, and biological processes that determine Earth system responses to external and anthropogenic forcing (*Figure 1*).

The need for RTMs is in fact also driven by vast growth of observation data documenting CZ processes, and specific hypotheses about solute chemistry and weathering processes suggested by these data (Brantley et al., 2013a; Kim et al., 2014; Perdrial et al., 2015). By integrating processes of interests in different disciplines, RTMs also allow integration between large, diverse, and complex data sets from disparate disciplines and scales, and can thus substantially advance cross-disciplinary research. For example, hydrologic models typically use a subset of observations for calibration, primarily water fluxes and hydraulic heads. Elemental and isotopic concentrations and speciation of dissolved components can also provide constraints on water flow (Jin et al., 2011; Thomas et al., 2013). To use these data requires RTMs. The use of RTMs will enrich our understanding of critical zone systems by eliminating some model behaviors in favor of other model behaviors (in other words some model simulations may reproduce water flux observations but not geochemical observations). RTMs can help discern such incongruities. RTMs therefore provide tools that synthesize understanding at a system level across disciplines while concurrently illuminate operative mechanisms and the contribution of individual processes.

The major advance of modern RTMs was to couple flow and transport within a full geochemical thermodynamic and kinetic framework. RTMs are distinct from geochemical models that primarily calculate geochemical equilibrium, speciation, and thermodynamic state of a system (Wolery et al., 1990). RTMs also differ from reaction path models (Helgeson, 1968; Helgeson et al., 1969) which represent closed or batch systems without either diffusive or advective transport. Multi-component RTMs originated in the 1980s based on the theoretical foundation of the continuum model (Lichtner, 1985; Lichtner, 1988). RTM development advanced substantially in the 1990s with the emergence of widely-used RTM codes and their capabilities (see (Steeffel et al., 2015a) for a review). Several early diagenetic models were developed at a similar time (Soetaert et al., 1996; Van Cappellen and Wang, 1996).

Reactive transport models have since been applied to understand biogeochemical systems for more than three decades (Beaulieu et al., 2011; Bethke, 1996; Brown and Rolston, 1980; Chapman, 1982; Chapman et al., 1982; Lichtner, 1985; Ortoleva, 1987; Regnier et al., 2013; Steefel et al., 2005; Steefel and Lasaga, 1994) in several earth system disciplines, including hydrogeology, geochemistry, and biogeochemistry. RTMs are used across a diverse array of environments involving both porous and fractured media (as reviewed in MacQuarrie and Mayer, 2005; Steefel et al., 2005). They can simulate a wide range of processes, including fluid flow (single or multiphase), solute transport (advection, dispersion, and diffusion), geochemical reactions (e.g., mineral dissolution and

precipitation, ion exchange, surface complexation), and biogeochemical reactions (e.g., microbe-mediated redox reactions, biomass growth and decay). Such models have been applied to understand a wide variety of systems and processes, including chemical weathering and soil formation (Brantley and Lebedeva, 2011; Goddris et al., 2013; Lawrence et al., 2014a; Maher, 2011; Maher et al., 2009a; Moore et al., 2012; Navarre-Sitchler et al., 2011), biogeochemical processes in coastal marine sediments (Dale et al., 2008; Krumins et al., 2013; Regnier et al., 1997), the role of spatial heterogeneity in determining mineral and biogeochemical reactions (Bolton et al., 1996; Bolton et al., 1997; Bolton et al., 1999; Li et al., 2007; Surasani et al., 2013). RTMs have played a key role in understanding and predicting coupled water and chemical mass transfer in environmentally relevant applications, including: bioremediation (Druhan et al., 2012; Li et al., 2009; Steefel et al., 2014; Yabusaki et al., 2011; Zachara et al., 2013); natural attenuation (Chen et al., 2001; Liu et al., 2008; Mayer et al., 2001); geological carbon sequestration (Apps et al., 2010; Atchley et al., 2013; Brunet et al., 2016; Carroll et al., 2016; Navarre-Sitchler et al., 2013; Tutolo et al., 2015; Xu et al., 2003); and nuclear waste storage (e.g., (Saunders and Toran, 1995; Soler and Mader, 2005)).

Microbial processes have been included in RTMs since the mid-1980s. Early applications included the groundwater substrate dynamics and microbial growth (Molz et al., 1986); the transport of bacteria and viruses with surface attachment and detachment (Yates et al., 1987); and multispecies interactions between oxygen, nitrate, organic carbon, and bacteria that regulate denitrification (Kinzelbach et al., 1991). New generation RTM codes have adopted more complex and flexible formulations including the kinetic degradation of microbial biomass and biogeochemical byproducts (Bao et al., 2014; Prommer et al., 2006). With the expansion of the use of isotopes as tracers of mineral-fluid and biologically mediated reactions, recent advances include the explicit treatment of isotopic partitioning due to both kinetic and equilibrium processes (Druhan et al., 2012; Johnson and DePaolo, 1994; Lehmann et al., 2007; Maher et al., 2006b; Van Breukelen et al., 2005; Van Breukelen and Rolle, 2012b; Wanner et al., 2015).

Figure 1. RTMs are being used currently and will be used in the future to cross temporal and spatial scales. The red oval and the red topics indicate disciplines where RTMs have been used most commonly in the past three decades. The blue oval reflects potential target opportunities for further RTM development and application.

Broad ranges of scales have been considered in reactive transport modeling. RTMs have been applied to understand subsurface processes in single pores (Kang et al., 2006; Li et al., 2008; Molins et al., 2012; Molins et al., 2014) and single cells (10^{-6} to 10^{-3} meters) (Fang et al., 2011a; Scheibe et al., 2009), pore networks and columns (10^{-2} to 10^{-1} meters) (Druhan et al., 2014; Knutson et al., 2005; Li et al., 2006; Meile and Tuncay, 2006; Yoon et al., 2012), field scales (10^0 to 10^1 meter) (Li et al., 2011; Liu et al., 2008; Maher et al., 2009a), and more recently the watershed or catchment scales (10^2 to 10^5 meters) (Atchley et al., 2014; Beisman et al., 2015; Navarre-Sitchler et al., 2013). Recent weathering studies have linked regional scale reactive transport models to global vegetation models to investigate the role of climate change in controlling weathering over periods of 10^0 to 10^3 years (Goddris et al., 2013; Godderis et al., 2006; Goddris et al., 2010; Roelandt et al., 2010). Models that

include a full coupling between subsurface reactive transport processes and surface hydrology, land-surface interactions, and meteorological and climatic processes have yet to be developed, although interests in improving process representation in climate models is stimulating these types of efforts.

RTMs are complex, physically based models that represent “pragmatic reality” and are similar to distributed hydrologic models. RTMs therefore suffer similar limitations with the problems “of nonlinearity, of scale, of uniqueness, of equifinality, and of uncertainty” (Beven, 2001a; Beven, 2001b; Ebel and Loague, 2006). Here we are not advocating for that complex models are better than parsimonious models. Such models are defined as models that accomplish a desired level of explanation or prediction power with as few variables as possible (Basu et al., 2010). Both types of models are useful. Parsimonious models can provide good conceptual frameworks. However, to derive and test parsimonious models in nonlinear natural systems, process-based models such as RTM are required to tease apart the importance of individual processes, to test the nonlinear coupling, and to identify controlling processes or parameters that need to be further investigated.

We present seven hypotheses in four different topical areas that highlight important questions and needs in understanding Critical Zone processes that can be uniquely addressed by application of existing RTMs or through the development of new RTM capabilities. We do not provide a comprehensive review of RTM applications, or a detailed discussion of concerns surrounding model use and evaluation. Instead, we aim to identify opportunities to integrate the RTM framework into other relevant disciplines to advance process understanding. The primary focus is in the CZ where interactions between water, atmosphere, rock, soil, and life create complex coupling between chemical, physical, and biological processes that determine system responses to external forcings (*Figure 1*). The hypotheses have the potential to elucidate a broad range of earth surface processes at different spatial/temporal scales and address long-standing questions regarding processes that ultimately sculpt the Earth’s surface and drive shallow subsurface evolution. While our list of hypotheses is by no means inclusive of all outstanding questions (cf. Steefel et al., 2005), these hypotheses focus on a broad cross-section of current research areas and illustrate where transformative scientific advances could be made through RTMs.

2. Testable Hypotheses

2.1 Chemical Weathering

Weathering of fresh bedrock generates soil and releases solutes to streams and groundwater, thus providing nutrients essential for terrestrial ecosystem productivity (e.g., Carey et al., 2005; Huntington, 2000; Moulton et al., 2000) and controlling chemical composition of natural waters (Malmstrom et al., 2000; Williams and Melack, 1997). At the global scale, silicate and carbonate weathering together consume about 0.3 Gt of atmospheric CO₂ per year (Gaillardet et al., 1999). Weathering and CO₂ levels are closely coupled through the dependence of weathering rates on temperature, pH, and water flux. This coupling maintains habitable Earth surface conditions for life over time scales of ~10⁵ – 10⁶ years for silicate weathering and over time scales of ~10³ years for carbonate weathering (Berner et al., 1983; Roelandt et al., 2010; Sundquist and Visser, 2003; Walker et al., 1981).

A process-based understanding of the controls on chemical weathering remains a central challenge for Earth surface studies. These processes include feedbacks between chemical reaction rates and variations in environmental conditions, particularly the availability of water. The introduction of chemically aggressive water in disequilibrium with bedrock drives weathering. Water serves as a medium for transporting weathering reactants and products, setting the mineral-water contact area, and affecting concentrations of reactive gases (e.g., CO₂, O₂). In hydrologically saturated systems with low flow rates, O₂ can be consumed faster than it is replenished via diffusion, resulting in anoxic conditions. In contrast, in unsaturated systems, oxygen diffusion is often sufficient to create oxic conditions and to drive oxidative degradation of organic matter in soils (Bolton et al., 2006). Likewise, rock chemistry can affect the presence or absence of O₂ and CO₂ as a function of depth (Brantley et al., 2013b).

Organisms can control the rates and extent of weathering by transferring water and nutrients among the solid, aqueous, and biological reservoirs in the CZ through root growth and exudation, and through associated fungal and microbial organisms. The availability of reactive gases is tightly coupled to biological processes that are notoriously difficult to resolve. The partial pressure of soil

CO_2 (P_{CO_2}) controls the formation of carbonic acid, which plays a major role setting the acidity of soil pore waters (Drever, 1989; Drever, 1997). The presence of O_2 can drive oxidative weathering of Fe(II)-bearing minerals such as biotite, which may create fractures and porosity that allow water infiltration and weathering of other minerals (Bazilevskaya et al., 2013; Buss et al., 2008; Molins and Mayer, 2007). Oxidation of pyrite can also produce sulfuric acid, which can enhance mineral weathering and create permeable pathways for water and solutes (Brantley et al., 2013b). Much remains to be learned about feedbacks among weathering, transport, and biologically-mediated processes with realistic subsurface structures at spatial scales of micrometers to meters. Despite advances in modeling these processes, their coupling to the long-term evolution of the critical zone and to climate cycles at the global scale have only begun to be explored. In the next sections we present two hypotheses that are ripe for testing with observations and RTM.

Hypothesis 1: Under humid conditions where precipitation (P) exceeds potential evapotranspiration (PET), the rate of chemical weathering for a given mineral assemblage is controlled by the concentration and distribution of reactive gases; in contrast, in arid systems, water is the primary control on chemical weathering rates.

Motivation. We suggest that water availability determines the extent to which reactive gases control the rates of chemical weathering. For example, under humid conditions—such as those in tropical forests ($PET/P < 1$)—water-rock contact area is large and water flow is fast such that reaction products are quickly transported away. The weathering rates are therefore determined by the concentration and spatial distribution of the reactive gases (**Figure 2**). Conversely, under relatively dry conditions—such as those found in semi-arid environments ($PET/P > 1$)—water-rock contact area is small and water flow is slow compared to gas diffusion through the vadose zone, creating water-limiting situations where water controls weathering rates (Perdrial et al., 2014). These fundamentally different controls on the rate of chemical weathering should ultimately result in unique and different physical and chemical architectures (i.e., structure) within the regolith.

Figure 2: conceptual diagram expressing the role of water and gases in driving weathering in different climatic zones: gas (white) transport is likely to be rate-limiting for weathering under humid conditions where water is abundant and precipitation (P) exceeds potential evapotranspiration (PET) ($PET/P < 1$); water availability (blue) is likely to be rate-limiting for weathering in arid ecosystems where water is scarce and gases can rapidly diffuse through unsaturated pore spaces. The blue/green layer indicates the saturated zone.

Current status. The hypothesis about the importance of water availability versus gas concentrations in different systems arises, in part, from previous RTM work on chemical weathering. Modeling of chemical weathering has thus far focused primarily on one-dimensional (1-D) systems at time scales typical of soil development and bedrock weathering. These models emphasize complex reaction networks among primary and secondary minerals, aqueous complexes and adsorbed species (**Sidebar 1**). They have illustrated key controls of Earth's surface evolution, namely fluid transport and reaction thermodynamics and kinetics in regulating both the advance rate and thickness of weathering fronts.

One of the long-standing puzzles in quantifying chemical weathering rates is the five to six orders of magnitude discrepancy between rates of mineral reactions as measured in laboratory versus field contexts (Blum and Stillings, 1995; Drever and Clow, 1995; Richards and Kump, 2003; Sverdrup et al., 1995; White, 1995; White and Brantley, 1995; White and Brantley, 2003). This rate discrepancy has been attributed to two major categories of factors that operate at different spatial scales. One category includes processes and conditions that directly affect the amount and “reactivity” of mineral surfaces. Examples include mineral surface roughness or fractalness (Navarre-Sitchler and Brantley, 2007), passivation or armoring layers as a result of clay coating or secondary mineral precipitation (Helgeson, 1971; Maher et al., 2009b; Nugent et al., 1998), leached layers that act as diffusion barriers (Luce, 1972), and water films between grains that vary under different temperature and pressure conditions (Renard et al., 1997). The second category includes factors that regulate fluid compositions. For example, water flow controls reaction rates by regulating the extent of deviation from equilibrium (Maher, 2010; Salehikhoo et al., 2013). The amount of wetted surface under unsaturated conditions, as well as concentrations of catalyzing or inhibiting aqueous species, also play a significant role (Lawrence et al., 2014a; Oelkers et al., 1994). Both of these categories of factors have been explored using RTMs together with field observations.

Porous medium characteristics—the particular details of pore structure and heterogeneous spatial distribution of porous medium properties (e.g., permeability and mineral abundance)—can lead to large spatial variations in flow rates and deviations from equilibrium, therefore modifying the overall “effective” reaction rates at large scales (Bolton et al., 1996; Bolton et al., 1999; Malmstrom et al., 2004; Meile and Tuncay, 2006; Molins et al., 2012; Salehikhoo and Li, 2015). For example, a mineral that does not have any coating can be bathed in a fluid at equilibrium in the low permeability zone of a heterogeneous medium and therefore is not effectively dissolving. In essence, any factors that influence the reaction kinetics (surface area, concentrations of catalyzing or inhibiting species, and distance from equilibrium) can contribute to the rate discrepancy. In natural systems, all these factors can operate simultaneously resulting in much smaller effectively-dissolving surface areas than the measured Brunauer–Emmett–Teller (BET) or imaged surface area, although the dominance of different factors may vary under different conditions (e.g., (Jin et al., 2011; Nugent et al., 1998; White and Peterson, 1990; Zhu, 2005)).

Modeling studies have also probed the effects of time and climate on weathering rates using chronosequences and climosequences (*i.e.*, soils of different age or soils of similar age and different climate regimes, respectively) (Goddéris et al., 2013; Goddéris et al., 2010; Lawrence et al., 2014a; Maher et al., 2009b; Moore et al., 2012). The advance rate of the weathering front may be limited by the accumulation of weathering products (Maher et al., 2009b; Moore et al., 2012) and by the drainage rate of groundwater from the fresh bedrock (Rempe and Dietrich, 2014). RTMs have also helped place constraints on conversion of rocks to regolith at the weathering front, including basalt clasts and carbonates (Brantley et al., 2013b; Hausrath et al., 2008; Hausrath and Olsen, 2013; Navarre-Sitchler et al., 2011; Navarre-Sitchler et al., 2009). Compared to water availability, the role of reactive gases has been explored to a much lesser extent, although modeling studies have emphasized the importance of including soil CO₂ in quantifying weathering rates (Brantley et al., 2013b).

Future directions. With a process-based framework, RTMs are uniquely suited to evaluate the competition between reactive gases and water availability under otherwise similar weathering conditions. Brantley et al. (2013) hypothesize that the soil O₂ to CO₂ ratio in different landscape positions (ridgetop versus valley) is related to the depths of weathering fronts for different mineral phases, but this has not yet been tested using a fully coupled model framework. Current RTMs include capabilities of multiphase flow that are required to simulate the fluid dynamics under different water availability (wet and dry) conditions, as well as the generation or consumption of reactive gases by microbial processes (Molins and Mayer, 2007). Modeling capabilities need to be expanded to include root processes in order to test hypotheses regarding how plants affect weathering. In addition to the traditional measurement of thermodynamic equilibrium constants and kinetic rate formulations of mineral reactions, coupled measurements of reactive gases and water chemistry at depth are also needed (Hasenmueller et al., 2015). Time-series measurement of mineral and hydrologic properties,

including surface area, porosity, permeability, and unsaturated flow parameters, are rare although they are critical to constrain RTMs (Jin et al., 2010; Navarre-Sitchler et al., 2009; Takagi and Lin, 2011).

Hypothesis 2: *At the global scale, variations in water availability induced by climate change exert more control over chemical weathering than increased CO₂ levels and temperature.*

Motivation. At the global scale, various forcing and feedback loops work together in regulating the ultimate consequences of increasing CO₂ levels and resulting climate change. Insights into critical feedbacks between climate change and chemical weathering, however, are limited (Beerling and Berner, 2005; Berner and Caldeira, 1997; Goddris et al., 2013; Roelandt et al., 2010). There has been debate about whether climate change and global warming will lead to positive or negative feedbacks between atmospheric CO₂ levels, plants, and weathering (Beerling and Berner, 2005) as both rising CO₂ level and temperature can enhance chemical weathering. Climate change can also have profound impacts on the hydrological cycle, including changes in evapotranspiration, rainfall abundance, precipitation patterns, shifts in snow and ice melting, and a change in the frequency of extreme events (Barnett et al., 2005; Bindoff et al., 2013; Chattopadhyay and Hulme, 1997; Jung et al., 2010; Rosenzweig and Neofotis, 2013). As mentioned in Hypothesis 1, water is a key driver for biogeochemical processes (Kump et al., 2000; Maher and Chamberlain, 2014). Changes in water cycles likely trigger shifts in plant response and soil CO₂ levels (Bond-Lamberty and Thomson, 2010), therefore regulating chemical weathering to a more significant extent than increasing atmospheric CO₂ level and temperature alone. In addition, changes in the timing and intensity of precipitation can alter recharge rates, which influence groundwater residence times, flow paths, and mineral weathering fluxes. The need to consider the sensitivity of weathering rates to flow paths, water availability, reactive gases, and temperature presents a classical reactive transport problem. However they have only begun to be explored using RTMs.

Current status. Our limited understanding of the relationship between climate change, CO₂ regulation mechanisms, and chemical weathering exists partly due to the lack of tools that explicitly couple relevant processes at the global scale. Many global models developed to understand the feedbacks between CO₂ level and chemical weathering are zero-dimensional (Berner and Kothavala, 2001; Edmonds and Edmond, 1995) and hence represent global average conditions. The Rock Geochemical Model (RokGeM) was developed for incorporation into the Global ENvironmental and Ecological Simulation of Interactive Systems (GENESIS) to understand the sensitivity of weathering to temperature and run-off (Colbourn et al., 2013). A recent advance has been coupling the rock weathering models with Earth System models to understand the interactions between terrestrial weathering, CO₂ levels, and climate change. For example, the weathering model WITCH has been coupled with the Lund-Potsdam-Jena (LPJ)—a Dynamic Vegetation Model (DVGM)—to examine the role of vegetation in affecting chemical weathering at the continental scale (Roelandt et al., 2010). Sensitivity analysis shows that in tropical environments, plants control weathering by introducing roots with organic acids that increase mineral-water contact and dissolution rates. When atmospheric CO₂ increases, decreases in evapotranspiration enhance infiltration, biosphere productivity, and soil acidity, promoting CO₂ consumption by mineral weathering by an estimated 2.4% - 5% increase in CO₂ consumption levels per 100 ppmv rise in CO₂ (Beaulieu et al., 2010). These estimates however, are highly contingent upon climatic, ecohydrological, lithological, and biological conditions. Most of these models are executed with steady-state CO₂ levels and hydrologic scenarios and do not consider transient precipitation patterns that accompany climate change. In addition, existing models do not consider weathering rate response to human activities such as land use change and fertilizer utilization (Guo et al., 2015; Perrin et al., 2008; Raymond et al., 2008). Only a few researchers have used RTMs to forecast future weathering (Goddris and Brantley, 2013; Goddris et al., 2013).

Future directions. Multi-component RTMs with spatially explicit representations of precipitation patterns and biosphere and geosphere characteristics can advance our predictive understanding of process coupling at continental and global scales (*Sidebar 2*). Model development along these lines, however, faces key challenges. Existing models of continental weathering typically have large grids [e.g., spatial resolution of 0.50.5 latitude-longitude in (Beaulieu et al., 2012)] whereas our understanding and measurements of multicomponent reactive transport processes currently focus on

much smaller scales (micrometers to meters). Translating mechanistic understanding of complex reaction networks at small scales (10^{-3} to 10^1 meters) to a few key reaction mechanisms at continental and global scale models (10^3 to 10^5 meters) presents a daunting task. Such cross-scale understanding, however, is key for advancing not only model development but also conceptual understanding of which processes are important at large scales. Such understanding will reduce computational cost and facilitate explicit incorporation of two-way feedback schemes among processes, instead the current use of cascade, or one-way simulation schemes, which lack feedbacks.

2.2 Carbon Cycling

Soil represents the largest terrestrial pool of organic carbon (C) (Stockmann et al., 2013). Soil C decomposition and stabilization plays a key role in determining the atmospheric CO_2 concentrations and global C cycles (Davidson, 2006). Although the complexity of soil C partitioning has long been recognized (von Lützow et al., 2007), soil C is currently conceptualized and modeled as pools of different decomposition rates and turnover times (Ostle et al., 2009; Thornton et al., 2009). For example, a typical three-pool soil C model includes a readily degradable C pool, with residence times less than 5 years; a slowly degrading pool, with residence times of decades; and a relatively stable pool, with residence times between 10^3 – 10^5 years (Marin-Spiotta et al., 2009; Trumbore, 1993; Trumbore et al., 1995).

One of the grand challenges is to understand mechanisms and quantify rates of C decomposition and stabilization under different environmental conditions (Conant et al., 2011; Lehmann, 2015). Carbon decomposition is primarily biologically driven. Microbes transform C (e.g., leaves, roots) in the presence of multiple electron acceptors including O_2 , nitrate, sulfate, as well as solid phases such as iron-containing and manganese-containing minerals (Dunn et al., 2006; Fierer et al., 2003; Philippot et al., 2009). Carbon stabilization, however, is both biotic and abiotic (Eusterhues et al., 2003; Guggenberger and Kaiser, 2003; Kleber et al., 2015; Lalonde et al., 2012; Mikutta et al., 2006; Mikutta et al., 2009; Rasmussen et al., 2007; Rasmussen et al., 2008; Rumpel et al., 2004). Plant roots and mycorrhizal fungi influence soil properties including physical structure, hydrological and biogeochemical gradients and, as a result, are integral controls on soil structure and water availability (Fierer and Schimel, 2002; Lawrence et al., 2009; Roy et al., 2013). A change in soil properties can limit organic matter degradation through physical separation from microbes and water (Manzoni et al., 2014), or by affecting organic matter-mineral associations (Mikutta et al., 2006; Schmidt et al., 2011). These physiochemical conditions are likely more important controls on stabilized or passive C than biological activities (Schmidt et al., 2011; Torn et al., 1997).

The competing effects of biotic and abiotic factors can be understood through the development of RTMs that simulate water dynamics, physical stabilization, and microbial growth conditions with biological processes and biogeochemical reaction networks (e.g., (Riley et al., 2014)). A detailed mechanistic understanding of the root zone will enable modeling approaches that elucidate the types of processes and the scales required to accurately capture subsurface C cycling, and the derivation of general principles for C stabilization / decomposition. Such mechanism-based principles have yet to be incorporated into earth system models, but hold promise for elucidating the feedbacks that govern the C cycles and climate system at the global scale.

Hypothesis 3: *As soil moisture increases, the fraction of C stabilized through mineral association increases, while the fraction of C stabilized through physical separation decreases.*

Motivation. Soil C stabilization occurs through two general groups of mechanisms: (1) physical separation of the organic carbon from the reactants or enzymes required for mineralization via occlusion, movement of C to depth, isolation of C in zones without suitable electron acceptors; and (2) movement of organic C into an unavailable form, for instance through association with mineral surfaces or through chemical recalcitrance (Conant et al., 2011; Kleber et al., 2015; Lehmann, 2015). Soil moisture governs partitioning between these two general mechanisms (**Figure 3**). As conditions become drier, we hypothesize the physical separation-based mechanisms will become more dominant due to the lack of water. Conversely, an increase in soil moisture and the wetting of new surface area will instead promote additional C-mineral associations, which can come to govern C stabilization.

Soil mineral composition varies significantly with climate (e.g., temperature and rainfall), parent rock, and weathering stage. Weathered soils often have minerals with high surface areas and large cation exchange capacities, which promote soil C stabilization through enhanced mineral-organic matter association and reduce soil C decomposition and CO₂ efflux (Denef et al., 2007; Eusterhues et al., 2003; Guggenberger and Kaiser, 2003; Lalonde et al., 2012; Mikutta et al., 2006; Mikutta et al., 2009; Rasmussen et al., 2007; Rasmussen et al., 2008; Rumpel et al., 2004). Soil mineralogy therefore can play an important role in determining soil C – climate change feedback. Rising CO₂ levels and climate change can in turn impact soil mineralogy by changing chemical weathering rates via shifts in water content, temperature, plant and microbial physiology, and the composition of plant community (Banwart et al., 2009; Berner, 2006) (Hypothesis 2). The codependence of mineral composition, weathering, and CO₂ concentration forms a coupled regulatory system for the C cycle at human-life time scales. Under all conditions, soil mineral and plant composition in the weathering zone will determine the system response and feedback to climate change. Such feedbacks however are not well understood (Trumbore and Czimczik, 2008).

Current status. The dependence of soil C stabilization and decomposition on soil mineral composition has not been considered in models until recently (Sulman et al., 2014). Carbon decomposition is commonly described with first-order dependence on soil moisture and / or temperature (Manzoni and Porporato, 2009; Parton et al., 1987), possibly with an extra Michaelis-Menten term to account for microbial processes (Wieder et al., 2013). The first order dependence on soil moisture can reproduce C behavior under steady-state moisture conditions, but not under dynamic conditions such as pulsed rewetting perturbations (Lawrence et al., 2009). Recent studies have recognized this inadequacy in the representation of complex soil C processes, especially when predicting the soil carbon-climate feedbacks. For example, limited representation of physical and chemical processes under varying hydrological (soil moisture), geochemical (pH, O₂, nutrient supply), and biological (e.g., microbe physiology and efficiency) conditions has been shown to introduce large uncertainties in estimating the temperature sensitivity of soil C decomposition under warming conditions (von Lützow and Kögel-Knabner, 2009). A recent comparison between predictions from 11 ecosystems models and free-air CO₂ enrichments (FACE) data indicate the prediction uncertainty is dominated by the poor representation of the C turnover time (De Kauwe et al., 2014). A model explicitly linking soil C decomposition with microbial and mineral surface sorption has predicted weaker soil carbon-climate feedbacks than models without these processes (Tang and Riley, 2015).

Given the importance of representing soil C process in future models (Georgiou et al., 2015), we need to evaluate the functional relationships between soil C decomposition rates and key soil properties including mineral composition as well as other relevant variables. The recent development of CORPSE (Carbon, Organisms, Rhizosphere, and Protection in the Soil Environment) offers a promising framework for modeling soil C dynamics (Sulman et al., 2014). CORPSE considers the proportion of protected and unprotected soil C pools as a function of clay content, with each pool including a mixture of compounds with different tendency toward decomposition (labile, resistant, and recalcitrant). A dynamic microbial biomass pool controls soil C transformation with rate dependence on the abundance of microbial biomass and concentration of degradable organic carbon through Michaelis-Menten formulation, in addition to the typical temperature and soil moisture dependence. The biomass can grow based on the amount of SOC assimilated, which allows

Figure 3. The primary mechanism for carbon stabilization likely differs in high and low moisture soils. In low moisture soils, physical separation dominates because the lack of water may separate microbial enzymes from readily degradable carbon in particles or micro-clusters. In high-moisture soils, C stabilization may be dominated by organic carbon sorption onto wetted mineral surfaces. High moisture soils also can facilitate microbial degradation because microbial enzymes will more likely be in contact with the C.

simulation of processes such as priming, where simple carbon input greatly increases the overall soil C transformation (Dungait et al., 2012; Finzi et al., 2015; Fontaine et al., 2007; Kuzyakov and Blagodatskaya, 2015). Riley et al. (2014) presents soil cycling model that includes a comprehensive treatment of microbial and fungal activity, stabilization mechanisms and representation of C via polymeric and monomeric carbon substrate groups. Although the model adequately reproduced C turnover times across a variety of sites, there are not yet sufficient measurements to fully evaluate such a complicated model (Riley et al., 2014).

Future directions. Existing RTM capabilities include multiphase flow and gas transport, preferential flow, and microbe-mediated redox reactions by different functional microbial groups. These capabilities can be used to track soil moisture, the evolution of microbial community composition and C decomposition in the presence of roots and mycorrhiza. In particular, multiphase flow dynamics (i.e., air and water in vadose zone) can help us understand transitions between dry and wet conditions, transformations between degradable and stabilized C, as well as important mechanisms for C stabilization under different physiochemical conditions. One challenge in soil C modeling is representing the complex chemistry of organic matter (Allen-King et al., 2002; Chen et al., 2003; Hedges et al., 2000). Molecular-scale representation of soil C chemistry in RTMs is not likely because of the computation cost associated with capturing the diversity of organic components (Tang and Riley, 2015; Tang and Riley, 2013). The advent of more sophisticated tools for characterization of organic matter chemistry offers an opportunity to devise new approaches for representing the functional diversity and reactions of organic matter.

Highly mechanistic representations of soil C have yet to be explicitly included in global scale earth system models such as the Community Earth System Models (CESM) due to observational and computational constraints. Moving from small scale, mechanistic understanding to large-scale earth system modeling presents a major challenge. Simple parameters representing mineralogy (e.g., clay content) and organic matter reactivity (e.g., C pool structure) may not adequately reflect C cycling under changing conditions or across broad spatial scales. Nevertheless, highly resolved models of soil C are necessary to determine the importance of spatial scale and the level of details needed to represent the carbon systematics at larger scales, and could thus be viewed as a critical component in the continued development of earth system models. Overall, RTMs provide a useful framework for assessing rate-limiting processes and system conditions or thresholds.

2.3 Spatial Complexity Across Scales

Natural systems are heterogeneous with large spatial variations in properties and fluxes. The geoscience community has long recognized the importance of spatial heterogeneity in determining earth processes. As an example, the hydrogeology community has studied the role of spatial variation in porosity and permeability in regulating subsurface flow and solute transport since the 1970s. Large-scale tracer experiments have been carried out at field sites with varying degrees of spatial complexity. These include, the low and intermediately heterogeneous aquifers in Cape Cod, Massachusetts (LeBlanc et al., 1991) and the Borden aquifer in Ontario, Canada (Woodbury and Sudicky, 1991); as well as the highly heterogeneous Macrodispersion Experiment (MADE) aquifer in Columbus, Mississippi (Boggs et al., 1992), where the Advection-Dispersion-Equation (ADE) has been significantly challenged (Zheng et al., 2011). The field of stochastic groundwater hydrology was charged with a primary goal of understanding and quantifying the impacts of spatial variability on solute transport and representing this spatial complexity through effective parameters (Adams and Gelhar, 1992; Dagan et al., 2013; Fiori et al., 2010; Gelhar and Axness, 1983; Pedretti et al., 2013; Sudicky, 1986; Sudicky et al., 2010). Effective, upscaled models were developed to capture anomalous behavior (Neuman and Tartakovsky, 2009), including the dual porosity model (Gerke and Van Genuchten, 1993), multi-rate mass transfer model (Haggerty and Gorelick, 1995), non-local methods including Continuous Time Random Walk (Berkowitz et al., 2006), and Fractional Advection Dispersion Equation (fADE) (Benson et al., 2000a; Benson et al., 2000b). Common upscaling and non-local methods for conservative transport, however, do not discriminate between solute spreading and actual mixing (Molz and Widdowson, 1988).

Challenges arising from understanding spatial heterogeneity are universal across all earth disciplines. The past decade has witnessed leaps in understanding the spatial complexity in subsurface biogeochemical processes. The long-standing puzzle in the geochemistry field, i.e., the five to six orders of magnitude lower rates of weathering observed in natural systems compared to those in well-mixed batch reactors, has inspired the use of RTMs in examining the role of spatial heterogeneity in controlling chemical weathering rates (Bolton et al., 1996; Li et al., 2006; Malmstrom et al., 2004; Meile and Tuncay, 2006). RTMs simulating field scale remediation experiments also highlighted the significance of spatial heterogeneity in determining biogeochemical processes (Li et al., 2010; Liu et al., 2013; Tang et al., 2013; Yabusaki et al., 2011; Zachara et al., 2013) along with carefully designed laboratory-scale column or flow cell experiments (Miller et al., 2013; Rolle et al., 2009; Wang and Li, 2015; Willingham et al., 2008).

RTMs have not yet been utilized to understand many systems characterized by extensive and multi-scale heterogeneity. This is an area ripe for advance because geochemical and geobiological data are now being analyzed in well-instrumented field sites such as the Critical Zone Observatories that can constrain RTMs over spatial and temporal scales that have not been possible previously. For example, the root zone represents an example of such extreme heterogeneity. The hyporheic zone, i.e., the interface between river, land and subsurface, is another zone where hydrological and biogeochemical complexities collectively influence biogeochemical processes and reaction rates. The topography, vegetation type, lithological variation, and river network geometry impose spatial heterogeneity that plays a pivotal role in partitioning energy, water, and biogeochemical fluxes across different compartments in watersheds. RTMs are well suited to study how reactions and fluxes scale in these highly heterogeneous systems.

Figure 4: Rhizosphere processes influence the development of biogeochemical gradients at a range of spatial scales including: (a) soil profile scale (b) pore scale and (c) mineral surface scale. Representation of these processes in RTMs requires consideration of how rhizosphere processes vary over the spatial and temporal scales of interest (Figure is adapted from (Lawrence et al., 2014b) and is used with permission).

Hypothesis 4: *Rhizosphere activities shaped by spatial structure of roots dominate soil formation and functioning.*

Motivation. Roots represent structural, biogeochemical and hydrological interfaces of key importance in generation of soil. Root-soil-water-microbe interactions couple abiotic and biotic processes, including chemical weathering and soil C processes discussed in previous hypotheses. The spatial configuration of a root system -- the Root system architecture (RSA) -- anchors and mechanically supports plants, facilitating uptake of water and nutrients (Garré et al., 2012). The RSA varies by plant species, adapts to unevenly distributed resources (Bouma et al., 2001; Lynch, 1995), and is responsive to changing soil conditions (Jung and McCouch, 2013; Pacheco-Villalobos and Hardtke, 2012; Tron et al., 2015).

Root structures also impact soil physical properties and generate macro pores that act as highly permeable conduits for water (Beven and Germann, 1982), profoundly impacting water flow and solute transport. Roots control soil chemistry by modulating soil moisture, pH, sorption capacity, and extent of aqueous complexation (Hinsinger et al., 2011; Hinsinger et al., 2003; Lucas et al., 2011)

(**Figure 4**). Chemicals released from mineral weathering and C decomposition are taken up by roots (Banwart et al., 2011; Brantley and White, 2009; Lopez et al., 2009), while root exudates and other products—often organic acids and CO₂ gas—can accelerate mineral weathering and soil C processes (Hinsinger et al., 2003; Oh et al., 2007). Bacteria and fungi are closely associated with roots and affect soil processes (Drever, 1994; Landeweert et al., 2001; Thorley et al., 2015). Root structures therefore dictate the spatial distribution of water, reactive species, and biogeochemical reaction rates.

Current status. Root models have been developed extensively by plant scientists to describe water and nutrient uptake (e.g., (Diggle, 1988; Lynch et al., 1997; Smithwick et al., 2014; Warren et al., 2014)). Agricultural scientists have developed crop models, which typically simulate soil water dynamics, soil-plant nitrogen and phosphorus cycling, crop canopy and root growth, plant dry matter production, yield, plant residue production and decomposition, and erosion (Bassu et al., 2014; Camargo and Kemanian, 2016; Stöckle et al., 2003; White et al., 2014). However, soil and water biogeochemical dynamics—including soil composition evolution export of dissolved organic carbon—are typically not considered mechanistically in crop models. Early biogeochemical models employed either discrete soil layers with different physical and chemical properties (Veen and Paul, 1981) or continuous descriptions of soil organic matter and nutrient dynamics (O'Brien and Stout, 1978). Most existing models consider multiple soil layers with advection and diffusion processes (Maggi and Porporato, 2007; Miller et al., 2008; Raynaud et al., 2006; Walter et al., 2003), sometime considering the statistical characteristics of root structure (Porporato et al., 2003). Explicit representation of biogeochemical processes associated with spatially heterogeneous roots has been rare, with only a few exceptions (Schnepf et al., 2012; Wu et al., 2007). Effective models have been developed to represent the heterogeneous soil system with averaged and lumped properties (Manzoni et al., 2008). At larger scales, roots are rarely represented explicitly in soil production and landscape evolution models (Lifton and Chase, 1992; Minasny and McBratney, 1999; Minasny and McBratney, 2001; Minasny and McBratney, 2006; Pelletier and Rasmussen, 2009; Yoo and Mudd, 2008). Increasingly, field evidence from some settings suggests that large tree roots could achieve substantial penetration (up to 3 m) into shallow bedrock, therefore accelerating weathering processes (Roering et al., 2010). Key parameters for roots include those relating root depth and root density to macropore permeability and size, all of which are important in distributing water.

Recent incorporation of root functions in large-scale hydrological, ecological, and climate modeling have improved model simulation of latent heat fluxes (Zheng and Wang, 2007), soil moisture dynamics (Ren et al., 2004), and seasonal climate (Barlage and Zeng, 2004). Although vegetation types have been well classified and parameterized in these models, RSA has not been differentiated among different vegetation types. For example, water flow between different soil layers via root systems occurs in different ecosystems and plant species throughout the world (Neumann and Cardon, 2012). This water movement is driven by gradients in water potential at the soil-root interface and is regulated by root structure, which is believed to have significant impacts on water budget, plant transpiration rate, and temperature at both local scale and ecosystem scale (Prieto et al., 2012). To date, land surface models generally consider rooting depth and fine root density in soil layers (Ryel et al., 2002), but do not account for the role of root structure in determining local soil and climate conditions. The lack of representation of root structure in biogeochemical models limits our ability to simulate seasonal water and biogeochemical cycling (e.g., C, N) and potentially increase the model biases relative to dynamic soil moisture, plant transpiration rate, and gross ecosystem productivity data (Zheng and Wang, 2007). Few RTMs have included the chemical effects of roots as a function of depth (Goddéris and Brantley, 2013; Goddéris et al., 2010).

Future directions. Understanding the role of root structure is challenging using experimentation alone, largely because of the complexity of biotic and abiotic interactions, the stochastic and evolving nature of root architecture and the expenses associated with obtaining realistic root structures. RTMs can go beyond measurement limitations and have been used to numerically describe the dynamics of microbial populations, mineral composition, and sorption/desorption processes in heterogeneous, multiphase subsurface systems (Yabusaki et al., 2011). To mechanistically represent intricate interactions among roots, microbes, and minerals, RTMs need to incorporate dynamic root growth as a function of water and/or nutrient concentrations and the effects of soil organisms, root exudates, and the above and below ground linkages that in turn influence soil properties and vegetation composition

(Wardle et al., 2004). Incorporation of these dynamic biological structures and spatial complexities into RTMs will allow prediction of soil biogeochemical processes and ecosystem response to environmental perturbation. The role of root structure on biogeochemical processes could be evaluated at the soil profile scale by comparing measurement data versus modeling output regarding weathering rates and the development of redoximorphic mottling with soil heterogeneity generated from stochastic or resource driven root distributions (e.g., (Schulz et al., 2016)). Such complex models could be used to pinpoint processes of key importance through global sensitivity analysis (Fenwick et al., 2014; Neuman et al., 2012), and ultimately compared with simple models that only include parameters representing root function (e.g., root depth and density). This would allow the distinction between parameters that are critical to the processes of interest and those that are not essential. In other words, RTMs should be used to guide measurement and experimentation by highlighting the most important processes and parameters through sensitivity analysis or data assimilation.

Hypothesis 5: RTMs can be used to identify and quantify critical interfaces that control large-scale biogeochemical reaction rates and ecosystem functioning.

Motivation. Life thrives at interfaces characterized by large gradients in water content, redox, nutrient, and/or pH conditions and organic C availability (Prommer et al., 2006). Here we define critical interfaces as those having distinct hydrological, biogeochemical, and topographic properties where reaction rates are orders of magnitude higher than the rest of the domain. These discrete interfaces can in some cases dominate reactions within an entire watershed. These interfaces include, for example, interfaces defined by internal domains in porous media (IDPM) (Zachara et al., 2016), biofilms on rock surfaces, surface water-ground water mixing zones, discrete reaction fronts within soil pedons (Brantley et al., 2013), and hyporheic zones at river-land-subsurface interfaces and contrasts between low- and high-permeability zones (Hefting et al., 2013; Hester and Gooseff, 2010; McClain et al., 2003; Stonedahl et al., 2010) (**Figure 5**). Biogeochemical processes and biodiversity

Figure 5: Hyporheic processes that occur at various interfaces at different scales. (Figure from (Stonedahl et al., 2010), use with permission).

at these interfaces often outsize the rest of the domain and dictate ecosystem functioning and reaction rates at larger scales. For example, interfaces between different soil zones have been observed to have 2-20 times higher microbe density than the bulk soil (Kuzakov and Blagodatskaya, 2015). In engineered bioremediation systems, microbe-mediated redox reactions dominate at the interface between the high permeability zone—where organic carbon is injected—and the low permeability clay zone—which is rich in electron acceptors (e.g., iron-containing minerals) (Li et al., 2011). Nitrate removal in the shallowest hyporheic zone (<4 cm) can be substantial enough to account for total nitrate removal of the whole stream (Harvey et al., 2013). Respiration rates at the land-river-sediment interface have been found to be 16 times more than those within the alluvial reach (Argerich et al., 2011).

Growing evidence has demonstrated that watershed topography governs the spatial distribution of the flow converging area (Takagi and Lin, 2011), leading to high C decomposition rates (Andrews et al., 2011) and high soil CO₂ concentrations (Hasenmueller et al., 2015) at critical interfaces. Extensive studies have shown that solute transport into and out of these interfaces—often determined by transverse mixing or diffusive fluxes—plays a key role in reaction extent (Cirpka et al., 1999; Cirpka and Valocchi, 2007; Janot et al., 2016; Jose and Cirpka, 2004; Luo and Cirpka, 2008; Werth et

al., 2006). The complexity of the hyporheic zone has continued to inspire many open research questions (Boano et al., 2014; Harvey and Gooseff, 2015). Based on field observations we support the idea that critical interfaces dominate overall behavior and reaction rates at larger scale. In a world with advanced RTM we can explore this idea with the integration of data and numerical simulations. Identification of critical interfaces and quantification of rates at the critical interfaces using RTMs can guide experimental and field campaigns on where, what, and how intensive measurements should occur. We also argue that understanding and quantifying the critical interface is key to upscale from detailed representation at small scales to relatively simplified, large scale representation of salient process characteristics.

Current status. The subsurface community has used multi-component RTMs to understand the role of spatial heterogeneity in determining microbe-mediated redox reactions (Bao et al., 2014; Cirpka et al., 1999; Cirpka and Valocchi, 2007; Scheibe et al., 2011; Zhang et al., 2010b), adsorption-desorption reactions (Deng et al., 2012; Liu et al., 2013; Liu et al., 2008; Sassen et al., 2012; Szecsody et al., 1998a; Szecsody et al., 1998b), and mineral dissolution and precipitation reactions (Atchley et al., 2014; Meile and Tuncay, 2006; Navarre-Sitchler et al., 2013; Werth et al., 2010; Willingham et al., 2008; Zhang et al., 2010a). Various studies have emphasized the importance of connectivity, transverse mixing, and reactive interfaces of heterogeneous porous media in regulating solute transport (Pedretti et al., 2013; Renard and Allard, 2013; Rolle and Kitanidis, 2014) and reactions (Chiogna et al., 2011; Jin et al., 2011; Rolle et al., 2009; Salehikhoo and Li, 2015; Wang and Li, 2015).

Biogeochemical reactions have begun to be integrated with complex flow dynamics to explore heterogeneity in the hyporheic zone (Bardini et al., 2012). At the watershed scale, RTMs need to be coupled with surface hydrological models that have spatially-explicit representations of surface characteristics such as topography, river network, and vegetation types. The surface hydrology community has utilized distributed models for more than five decades to understand processes including rainfall, infiltration, runoff, surface water and subsurface water interactions, and river hydrodynamics (Abbott et al., 1979; Abbott et al., 1986; Beven, 1989; Freeze and Harlan, 1969; Gan et al., 2006; James, 1972; Jarboe and Haan, 1974; Kumar et al., 2009; McDonnell et al., 2007a; Qu and Duffy, 2007b; Quinn et al., 1991; Singh, 1995; Therrien et al., 2010; VanderKwaak and Loague, 2001). Solute and water quality models have been developed as add-ons to hydrological models to understand environmental impacts of contaminants (Arnold and Soil, 1994; Donigian Jr et al., 1995; Santhi et al., 2001). These models use empirical and simplified correlations and do not incorporate physics-based multi-component reaction network and rigorous thermodynamics and kinetics representations (Davison et al., 2014). Model development that integrates surface hydrology model and reactive transport model has recently started (Yeh et al., 2006). Similarly, recent model development has coupled hydrological, land-surface interactions, and reactive transport processes at the watershed scale (Bao et al., 2016; Duffy et al., 2014).

Future directions. Although it is conceptually obvious that critical interfaces regulate biogeochemical reaction rates, it remains unclear how to quantitatively relate critical interfaces to process representation and biogeochemical rates at larger spatial scales. *How large are the critical interfaces? What are the general principles and quantitative function that relate small-scale spatial organization to the formation and fluxes at critical interfaces?* Answers to these questions lie at the heart of developing larger scale predictive models using salient small-scale characteristics. Challenges remain partly due to difficulties and uncertainties in characterizing complex, hierarchical spatial patterns across scales. In this context, process-based models that explicitly incorporate spatial complex present a powerful and cost-effective tool in querying heterogeneity controls of biogeochemical processes.

Models developed for processes across hydrology and biogeochemistry can be used as virtual laboratories for numerical experiments, whether they focus on the hyporheic zone or at the watershed scale. Although computationally expensive, these models can reveal rich information that would allow analysis of ecohydrological and biogeochemical processes at the grid block (local) scale and their key controls, as well as those at large scales through the mass balance of the system. Output from a watershed hydrobiogeochemistry model will reveal the temporal and spatial distribution of variables, including temperature, soil moisture, chemistry of gases, groundwater, and stream flow, and quantities derived from these variables such as rates of soil weathering and carbon decomposition.

Their relationships with key characteristics at the watershed scale, including the dynamic connectivity, critical interface areas, and topographic features, can be explored. Such exercise can potentially “discover” general principles across scales, as well as emergent zones or critical interfaces which are not identified by spatially scarce measurements. As an example, the sediment transport modeling of Chesapeake Bay identified a persistent zone of high particle concentrations formed due to convergent flow, which was later confirmed by field survey that showed maximum abundance of plankton groups in this zone (Hood et al., 1999). Advances along these lines will ultimately allow the fundamental underpinning of dominant processes at different spatial scales and will foster the development of relationships between large-scale rates, reactive interfaces, and key spatial complexity metrics, therefore moving toward predictive understanding across scales.

2.4 Integration of Novel Data

The past decades have witnessed rapid advances in technology and the generation of novel data at scales as small as nanometers to those as large as the globe (i.e., remote sensing from satellites). Advanced observation techniques have expanded our visual capabilities from a relatively narrow range ($10^{-3} \sim 10^1$ meters) to a much wider spectrum ($10^{-9} \sim 10^7$ meters). Our insight into microbial processes has been revolutionized by the ability to broadly query microbial communities through genomics and the development of techniques to measure gene expression, proteins and metabolites (transcriptomics, proteomics, metabolomics) (Sharon and Banfield, 2013)—together referred to as “omics” data. New analytical capabilities have rapidly expanded the use of isotopes to elucidate key processes where multiple reaction and transport pathways may obscure the interpretation of solute concentrations alone (Albarede, 2004; Anbar, 2004; Johnson et al., 2004; Williams and Hervig, 2005). The field of hydrogeophysics has grown enormously with the development of non-destructive imaging to allow glimpses into complex subsurface processes ((Binley et al., 2015; Parsekian et al., 2015; St. Clair et al., 2015). Advances in field characterization methods have provided high-resolution data to identify bioreactive fringes, mixing zones, and “hot spots” in the subsurface (Haberer et al., 2011). Imaging techniques such as X-ray Computed (micro)tomography have enabled unprecedented visualization of pore-scale details, allowing fundamental conceptualization and direct simulation of processes at the scale of microns (Blunt et al., 2013; Bultreys et al., 2016; Molins et al., 2012; Steefel et al., 2015b). In addition, the collection of different data types have started to be coordinated through large research community networks such as the Critical Zone Observatory, allowing for integrated understanding of the interactions among processes that are traditionally done in disparate disciplines (Brantley et al., 2016; Brooks et al., 2015; Clark et al., 2015).

The growth in data type and intensity presents a major advantage for the use of models such as RTMs that typically represent complex processes and require a large number of parameters (Duffy et al., 2014). The incorporation of these novel data into RTM frameworks, however, introduces additional challenges. Many of the new data do not directly relate to solute or solid concentrations that RTMs calculate. Comparison of data and model output therefore requires further conceptualization, insights, and parameterization. For example, electrical geophysical methods typically measure properties such as electrical conductivity and chargeability, which reflect the overall changes induced by aqueous reactions and water-rock interactions such as surface complexation and ion exchange. Incorporation of geophysical data with RTMs has only recently begun (Singha et al., 2011; Wu et al., 2014). In general, the integration of novel data into RTM framework represents a development of broad and interdisciplinary potential.

Hypothesis 6: *The stable isotope partitioning of heterogeneous reactions in natural systems can range from dominantly irreversible to fully reversible, and requires detailed, spatially resolved description of both fluid and solid phase isotopic compositions at fluid-solid interfaces.*

Motivation. Isotopes have been extensively used as tools for hydrologic characterization, analysis of water-rock-life interactions, fluxes, and mixing between hydrologic and chemical reservoirs. Examples include the use of ^{13}C , ^{15}N and ^{14}C in soil and groundwater systems to unravel processes of C and N transformations (e.g. (Dijkstra et al., 2008; Lawrence et al., 2015; Trumbore, 2009)); the characteristic fractionations associated with redox reactions employed to analyze elemental cycling (Cheng et al., 2016; Dale et al., 2009; Druhan et al., 2012; Gibson et al., 2011; Hubbard et al., 2014);

the growth and stability of secondary minerals (2015; Mavromatis et al., 2014; Nielsen et al., 2012; 2015; Oelze et al., 2014; Schott et al., 2014; Watkins et al., 2013); transient isotope partitioning during the initial stages of mineral dissolution (Brantley et al., 2004; Chapman et al., 2009; Crosby et al., 2007; Kiczka et al., 2010; Weiss et al., 2014; Wiederhold et al., 2006; Wimpenny et al., 2010; Ziegler et al., 2005), and application to solute fluxes across landscapes (Dellinger et al., 2014; 2015; Louvat et al., 2014; Noireaux et al., 2014; Rive et al., 2013; 2012a; Tipper et al., 2012b; von Blanckenburg and Bouchez, 2014; von Blanckenburg and Schuessler, 2014).

Current status. Isotope-specific analytical expressions for the combined effects of transport and reactivity have been derived and applied to a variety of stable (Bouchez et al., 2013; Fantle and DePaolo, 2007; Fantle et al., 2010; Jacobson and Holmden, 2008) and radiogenic systems (DePaolo, 2006; Johnson and DePaolo, 1994; Johnson and DePaolo, 1997; Lemarchand et al., 2010; Maher et al., 2006a; Tricca et al., 2001). However, more often approximations using either the Rayleigh distillation model or simple mixing models are applied to interpret stable isotope data in soils and rivers and groundwater systems, although the assumptions of these models are usually violated. Most of these systems could be incorporated into a multi-component reactive transport description that considers a suite of parameters that influence observed reaction rates (Maher et al., 2006b) or fractionation factors (Druhan et al., 2013). As a result, the recent expansion of stable isotope measurements has been accompanied by increased isotope-specific capabilities in RTM. Some models consider transport-based fractionation (Eckert et al., 2012; Rolle et al., 2010; Van Breukelen and Rolle, 2012a), or addition of individual isotopic ‘species’ in existing models that are necessary to describe contaminant degradation and sorption reactions (Hunkeler et al., 2009), including the transformation through multiple reaction pathways (Jin and Rolle, 2014). Others incorporate isotope-specific rate laws for mineral dissolution and precipitation that impact radiogenic and radioactive isotopes (Maher et al., 2006b), stable isotope fractionation associated with mineral growth (Druhan et al., 2013), abiotic redox reactions (Jamieson-Hanes et al., 2012; Wanner et al., 2014) and Monod kinetics (Cheng et al., 2016; Druhan and Maher, 2014; Gibson et al., 2011). A key advantage of these approaches is that the major element compositions are directly associated with the isotopic fractionation, enabling more rigorous evaluation of the transformations than could be obtained by considering the elemental or isotopic data in isolation (Sullivan et al., 2016). Currently, parameterization of existing RTMs is often limited by a lack of information on the processes that lead to differential isotope partitioning.

Future directions. There are two aspects of isotope-enabled RTMs that provide an array of interesting future research directions. The first is the use of isotopes to provide additional constraints on subsurface biogeochemical reactions. In many systems, major elements may participate in multiple reactions whereas the isotopic fractionation may be strongly determined by one reaction. Examples include Ca isotope fractionation due to calcite precipitation (Druhan et al., 2013), Li isotope fractionation during weathering (Bouchez et al., 2013) and metal isotope fractionation due to redox transformations (Druhan et al., 2014). The application of RTMs to interpret multiple isotopic profiles in tandem with the elemental variations may offer new insights into the isotopic system and reactions over space and time.

Nevertheless, to fully exploit the power of isotopes as tracers of biogeochemical processes, much more information about the processes that control fractionation is required. Although decades of research have focused on determining the causes and magnitudes of isotope fractionation for reactions with a continuum of departures from equilibrium (or a mixture of kinetic and equilibrium fractionation), the description of this fractionation remains incomplete. There is thus a pressing need to determine fractionation factors defined as the ratio of either the rate constants (kinetic fractionation) or the equilibrium constants (equilibrium fractionation) for two isotopologues of the same species. This definition is compatible with the existing structure of RTMs and is in stark contrast to the empirical, condition-specific fractionation factors. Ultimately, the rigorous mass balance provided by RTMs may enable us to determine fractionation factors in the field using multi-element approaches to constrain the isotope partitioning. In addition, as diffusive fractionation and apparent fractionation due to mixing processes are widespread in subsurface systems, better understanding of the kinetic and equilibrium fractionation factors would enable RTM applications to separate the effects of transport (*i.e.*, diffusion and mixing) from biogeochemical processes.

An additional challenge involves treatment of the solid phase during heterogeneous reactions. Continuum-scale RTMs describe the mineralogical composition of each grid block as a collection of mineral phases. The stable isotope composition of each mineral must then be incorporated and secondary minerals formed in the simulation must be described as a mixture of isotopic end members with temporally variable or spatially zoned isotopic compositions. A consequence of describing heterogeneous isotope fractionation in current continuum-scale RTMs is that the growth of a mineral phase requires the use of a solid solution, while dissolution of a solid is accomplished by the use of a pre-specified isotopic composition (Druhan et al., 2013). As a result, this approach lacks the capability to describe stable isotopes across reversible saturation states. Current research using lattice-Boltzmann (Huber et al., 2013) and surface kinetic (Maher et al., 2016) or ion-by-ion (Nielsen et al., 2012) models offer more accurate descriptions of isotopic incorporation into solids. These methods have yet to be reconciled with continuum scale approaches. Future research on the reversibility of stable isotope fractionation factors needs to be developed at spatial scales appropriate to describe fluid-solid interfaces in order to support and validate model development. Grain-scale models with moving boundaries within a continuum grid cell could be adapted for isotopic composition evolution as has been done for trace element mineral compositions during recrystallization (Neogi et al., 2014).

Hypothesis 7: *The incorporation of “omics” data into RTMs will enable predictive understanding of the emergent behavior of dynamic biogeochemical systems.*

Motivation. Biogeochemical systems are complex, internally interactive, and fundamentally nonlinear. Dynamically evolving microbial communities shape the distribution of biogeochemical processes in earth surface environments (**Figure 6**). The advent of omics data in recent decades had motivated the rapid advances in developing metabolic models to understand bacterial cell signal-transduction proteins, regulators, and metabolic and respiratory subsystems (Fredrickson et al., 2008; Reed and Palsson, 2003). The integration of omics data into RTM development, however, has not kept pace (Fang et al., 2011b; King et al., 2009; Vallino, 2003; Zhuang et al., 2011). The development of omics-enabled RTMs is challenging for several reasons. First, it requires expertise in microbial ecology and geochemical modeling along with omics methods and bioinformatics (Feist et al., 2009). Only recently have we begun to train scientists to effectively bridge these fields. Second, the nature of omics data differs from concentration, geochemical speciation, or direct rate measurements that traditionally have been integrated with RTMs. Such data is not *directly* comparable to the output of the current generation of RTMs. Third, and most importantly, the complexity of the microbial community revealed by omics data needs to be reduced to a level that is practical and can be parameterized, but yet inclusive of system-determining biogeochemical processes for integration into RTMs.

Current status. The incorporation of microbial activity in RTMs has mostly been limited to bulk biomass growth (Barry et al., 2002; Tartakovsky et al., 2009) and the indirect effects of microbial activity on electron acceptors and organic C concentrations using Monod or Michaelis-Menten rate laws (Li et al., 2010; Tartakovsky et al., 2013). These model formulations aim at simulating the impact of environmental conditions (e.g., substrate availability) on microbial activity, which then feedback on the environment (by community growth, for example). With careful simulation construction, numerical models have been used successfully to discriminate between various microbial functions and to predict the microbial community response to biostimulation conditions where microbial populations respond to an overwhelming abundance of a known C source (e.g. (Druhan et al., 2012; Li et al., 2011; Yabusaki et al., 2011). Conversely, there are functional genes that code for specific microbial activities. One example is nitrogen cycling where genes are known for each enzymatic step. The transcription of these functional genes therefore can tell us the limiting steps / processes in nitrogen cycling in soils, even though the data do not yield rates directly (Wallenstein and Vitgalys, 2005; Yan et al., 2003). A key challenge is to predict feedbacks between the microbial community and environmental conditions (Bouskill et al., 2012). For example, nutrient limitation is likely to affect the emergent behavior and is difficult to predict. To date, the effects of nutrient limitation are only coarsely accounted for and only within the context of a generally rather limited, *a priori* defined reaction network (Bethke et al., 2008; Jin et al., 2012; King et al., 2010).

911

912

Figure 6. Reaction network and metabolic pathway using natural organic matter (OM) and a wide range of available electron acceptors. The two blue cells schematically represent microbial communities (MC) consisting of many different organisms. The two red arrows expand the MCs, leading to the oval region schematically describing enzymatic pathways. The oval region describes collective enzymatic pathways, or reaction networks, for the microbial communities that may be elucidated by genomic or proteomic analyses. The effective concentrations of involved enzymes along with other parameters descriptive of enzyme kinetics may be used to drive reaction-based calculations in RTM (courtesy of Chongxuan Liu, Pacific Northwestern National Laboratory).

Future directions. It is timely to consider how omics information can be used to develop functional-gene or protein-based models that include key enzymatic reactions (metabolic groups) controlling system behavior and response to environmental or geochemical change (**Figure 6**). To keep computational burden manageable, a limited number of microbial actors capable of carrying out biogeochemically significant reactions could be selected as the model core with steady state concentrations assumed for intracellular metabolites. Such a model would explicitly consider the dynamics of biomass, genes, enzymes, substrates, as well as the community control strategy (e.g., maximizing energy yield and/or biomass growth; (Jin et al., 2012)), and could be used to explore hypotheses on the role of dynamic microbial community membership and interactions on the functioning and resilience of biogeochemical systems. In addition, the integration of omics data into RTMs can take place through the establishment of quantitative relationships between the fundamentally measurable biological characteristics and model parameters, such as growth yields under varying environmental conditions (e.g., (Vieira-Silva and Rocha, 2010)). Complementary to the use of molecular biology data to inform the model structure or parameterization *a priori*, rich data sets also offer strong constraints on RTMs. This includes the insights offered on active pathways through, e.g., transcriptomics and proteomics, or the use of concentrations of a wide range of chemicals (e.g., metabolomics) in the calculation of bioenergetics differentiating between thermodynamically feasible and infeasible processes.

As a new generation of molecular-biology enabled RTMs are developed, there is potential to describe and predict isotope partitioning as a function of the evolving relationship between multiple enzymatic reactions within an individual cell and their expression as a function of dynamic microbial community membership and interaction. Elements that are strongly influenced by biological cycling, such as iron and sulfur, record long-term variations in isotopic fractionation through the rock record that preserves information about climate, redox conditions and microbial dynamics (**Sidebar 3**). Stable isotope distributions therefore offer key means of validating a new generation of molecular-biology enabled RTMs in complement to direct metagenomic data, particularly where acquisition of these data is limited or precluded. This will be relevant particularly in applications beyond modern day systems, such as paleosols and chronosequences where direct metagenomic observables have

been largely erased, but the characteristic partitioning of stable isotopes associated with past microbial communities is preserved. The combination of omics and isotope data will therefore enhance our interpretation of processes with relevance to short-term response (e.g. contamination or biostimulation) and in natural systems over geologic timescales. In fact, because of the long time scales involved in natural systems, we may be able to infer fractionation factors and reaction rate constants that are more accurate than those from laboratory systems, especially for those systems with small fractionation and rate constants (Goddéris et al., 2010).

3. LOOKING FORWARD

Over the past three decades, researchers have demonstrated the usefulness of RTMs in understanding and predicting coupled physical, chemical, and biological processes in natural and engineered subsurface environments. We recognize that the use of RTMs can be expanded to understand various earth surface processes and to address some of the pressing research needs of other disciplines. However, RTMs alone are insufficient (Duffy et al., 2014). Looking forward, we see many opportunities where the RTM framework can be integrated into existing models in related disciplines (*Figure 1*). This becomes particularly important as the science community increasingly recognizes the significance of subsurface processes in influencing earth system functioning and formation. For instance, groundwater has been found to profoundly impact wetland formation and vegetation gradients (Fan, 2015; Fan et al., 2013). Soil C decomposition and stabilization influence global C cycle and climate change (Schmidt et al., 2011; Trumbore and Czimczik, 2008). The seven hypotheses outlined here are examples that can be tested with current capabilities or future development of RTMs. In particular, we envision advanced understanding by integrating RTMs into disciplines including hydrology, geochemistry, soil sciences, terrestrial ecosystem sciences, and geomorphology. Although coupling models across disciplines presents numerous challenges including upscaling, evaluating and interrogating complex models, and data availability (Duffy et al., 2014), we believe this should be attempted. Otherwise, we will miss very important linkages and feedbacks among processes that regulate different parts of the earth surface systems. The current lack of models describing process-level coupling will significantly limit the use of large-scale models for forecasting global hydrology and biogeochemistry.

Challenges. RTMs are similar to distributed hydrologic models and therefore share their problems and limitations. Whereas hydrological models solve for water, hydraulic or pressure head and saturation, RTMs solve for a suite of chemical species, often in the order of tens. That leads to an additional level of reaction complexity with nonlinearly laws of thermodynamic relationships and kinetics. The RTM approach therefore imposes a higher data demand for defining model domains, conditions, and constraints. For instance, a distributed hydrologic model domain needs information on topography, soil maps, and soil hydrological properties as a function of depth. Many of these data are available from national databases, including the USGS National Elevation Dataset (NED), the Soil Survey Geographic (SSURGO) database, and the National Land Cover Database (NLCD). An RTM will need an additional suite of biogeochemical properties including spatial distribution of mineralogy, specific surface area, ion exchange capacity, organic matter content, and microbial biomass. In constraining the model output, it would need not only water fluxes but also concentrations of reaction products including, major cations (e.g., Ca, Mg, Na, K), major anions (e.g., SO₄, NO₃, Cl), dissolved organic and inorganic carbon, nutrients (N, P), and pH. The use of these additional data in constraining RTM however will enrich our understanding of complex earth systems.

As models become more complex, new tools to evaluate them will need to be developed. Uncertain parameters may range from categorical (e.g., boundary conditions), continuous (e.g., mean permeability or rate constants) to spatial (e.g., uncertainty on facies models). The problem of uncertainty in distributed models has been routinely addressed with the Generalized Likelihood Uncertainty Estimation (GLUE) in the hydrology community (Beven and Binley, 1992). Other approaches include distance-based global sensitivity analysis (Fenwick et al., 2014) and Maximum Likelihood Bayesian Model Averaging (BMA) framework (Neuman et al., 2012; Ye et al., 2004), can aid in evaluating parameter sensitivity across multiple categories and assist in model reduction. Unfortunately, robust approaches are not yet routinely applied to RTM-based studies. Data assimilation, originated from numerical weather prediction (Daley, 1991), is another technique that has been used widely in recent decades to address the issue of parameter non-uniqueness or

equifinality in atmospheric, geographic, oceanic, and hydrologic sciences (Beven, 2001b; Navon, 2009; Rabier, 2005). Data assimilation, however, has not been used to date with RTMs.

Upscaling represents another grand challenge in integrating RTMs into large scale hydrologic and earth system models that operate at much larger spatial scales than traditional RTMs. A similar challenge remains in upscaling from the hydrologic time of months and years to the landscape evolution timescale of millennia (Duffy et al., 2014). This challenge lies at the heart of Critical Zone science where one major goal is to understand measurements across systems operating at different time scales. As spatial and temporal scales increase, dominant processes might change due to the “averaging” and “aggregation”. This would require different mathematical formulation in representing large-scale processes, parameters, as well as boundary and initial conditions. Although theoretical upscaling frameworks such as volume averaging exist (Whitaker, 1999; Wood and Whitaker, 1998), these approaches fall short when dealing with complex biogeochemical systems that have a large number of species and highly nonlinear reaction thermodynamics and kinetics. A range of multi-scale hybrid modeling approaches have been reviewed and analyzed recently (Scheibe et al., 2015; Scheibe and Smith, 2015). Simple averaging however is not possible due to the high nonlinearity and complex process coupling. Massive numbers of process-based model simulations can be run to derive general principles and performance patterns and can be used to translate small-scale processes and characteristics into large scale representations (McDonnell et al., 2007b).

Directions. The integration of RTM framework into the broader hydrologic sciences will be a powerful approach toward understanding complexity at large spatial scales. Hydrologic models at the catchment scale typically include surface hydrological processes (Brunner and Simmons, 2012; Kollet and Maxwell, 2006; Kumar et al., 2009; Maxwell, 2013; Qu and Duffy, 2007a) and recently have been coupled with atmospheric drivers and land-surface interactions (Clark et al., 2015; Maxwell and Miller, 2005; Shi et al., 2013). Indeed, most hydrologic models are underconstrained: the incorporation of RTM modules into hydrologic models offers the potential to constrain simulations by incorporating expressions for the wide array of solutes moving in surface and groundwaters. Coupling hydrologic models with RTMs will advance our understanding of watershed hydrobiogeochemistry with the potential to resolve long-standing puzzles including, for example, the observed concentration-discharge relationship at the watershed scale (e.g., (Godsey et al., 2009; Kirchner, 2003; NRC, 2012)). RTMs incorporating complex physical settings of hyporheic zones and water corridors can offer fundamental insights into intriguing interactions among flow, transport, and biogeochemical reactions, likely demystifying observations at these critical reactive interfaces (Harvey and Gooseff, 2015).

The root zone represents another focal point where mechanistic understanding can be obtained by extending current RTMs capabilities. Such understanding is of interest to an array of disciplines (biogeochemistry, soil sciences, microbiology, hydrology, and terrestrial ecosystem sciences). Integration of spatially-explicit RTMs into existing Earth System Models (ESMs) will capture interactions among atmosphere, biosphere, and geosphere under changing climate. Expansion of temporal scales into geological time scales will allow the integration of geochemical processes with geomorphic frameworks to understand chemical, physical, and biological coupling in landscape evolution. Landscape evolution models have been developed that can explain topography (e.g. slope, relief, channel density) (e.g., (Chen et al., 2014; Goren et al., 2014)). Whereas soil production and thickness has been predicted in landscape evolution models (Dietrich et al., 1995), existing RTMs of chemical weathering have been developed mostly from the geochemical perspective. Initial steps have been taken to couple the modeling of chemical weathering and erosion (Lebedeva and Brantley, 2013; Maher, 2010). Such models predict the depth to fresh bedrock across hillslopes on the basis of simplifying assumptions, but other such models connecting hillslope development to the landscape scale are only now emerging (Anderson et al., 2013; Lebedeva and Brantley, 2013; Rempe and Dietrich, 2014; St. Clair et al., 2015). The incorporation of geochemical processes into long-timescale models of the spatial patterns of soil and weathered bedrock will increase our understanding of the co-evolution of climate, ecosystems, and landscapes (Duffy et al., 2014; Dupré et al., 2003; Goddérís and Brantley, 2013; National Research Council, 2010).

Much like the fractured nature of data collection within separate disciplines (Brooks et al., 2015), model development also reflect fragmentation among disciplines. As data collection has been coordinated within networks of communities through joint platforms such as the Critical Zone

Observatory, Long Term Ecological Research (LTER) network, and United States Geological Survey (USGS), we also emphasize the need for interdisciplinary model development and the integration of RTM capabilities into existing models in other communities (Duffy et al., 2014; Goddris and Brantley, 2013). For example, to understand the role of biotic processes in chemical weathering, root models from the plant community can be coupled with RTMs (DOE, 2015). In this context, a flexible modular approach is needed to advance modeling capabilities across disciplines. Several geoscience research communities have worked toward that direction, including the Community Surface Dynamics Modeling Systems (CSDMS) (<http://csdms.colorado.edu/>), where modeling infrastructure has been developed to minimize overlapping efforts. The Department of Energy (DOE) Advanced Scientific Computing Research (ASCR) and Basic Energy Research (BER) have embarked on a pilot effort to re-factor existing High Performance Computing (HPC) software and develop new computational tools that can be scaled up to the next generation ‘extreme-scale’ platforms. Efforts are needed to identify common interests to leverage resources and push synergistic model development for multiple scientific communities.

We are now in an exciting era characterized by an unprecedented luxury of rich data (Fan et al., 2015). Comparisons between data and models will offer insights about the key data that are critical to our improved understanding. Collaborative science is dissolving the boundaries that separate traditional disciplines. RTMs are acting as frameworks that quantitatively integrate across the fields of hydrology, geochemistry, geomorphology, plant physiology, and microbiology. Such interfacing will lead to greater understanding of the formation, evolution, and functioning of the Critical Zone (*Figure I*). While the era of numerical weather prediction by computers began about half a century ago, earth process forecasting will become possible with increasing observations and computational capabilities in the coming decades. Although large uncertainties and challenges will remain, we are ready to build numerical integrators for processes, data, disciplines, and scales to explore the evolution and functions of the critical zone.

ACKNOWLEDGEMENT

This manuscript grew out of the workshop “Expanding the role of reactive transport modeling in biogeochemical sciences”, held in Alexandria, Virginia on April 13 – 15, 2014 (Li et al., 2014) with support from the NSF Low Temperature Geochemistry and Geobiology (EAR 14-14558). EWB received support from NASA Astrobiology Institute’s Virtual Planetary Laboratory under Cooperative Agreement number NNA13AA93A. SLB acknowledges funding from DOE OBES DE-FG02-OSER15675. We acknowledge insightful feedbacks and careful editing from Jon Chorover and David L. Parkhurst, as well as stimulating and constructive comments from Olaf Cirpka and two anonymous reviewers that have significantly improved this paper. We appreciate Dr. Joan-Albert Sanchez-Cabeza for handling this manuscript.

Sidebars:

Sidebar 1: A One-dimensional RTM for chemical weathering

In many soil profiles, primary minerals such as plagioclase are depleted over time via chemical weathering. The relative depletion of minerals in relation to their original concentrations in parent materials is characterized by τ values. The τ values represent a fractional loss (or accumulation of minerals) where 0 is equivalent to the parent, -1 is 100% loss relative to the parent and 0.25 is 25% gain relative to the parent (Anderson et al., 2002; Brimhall and Dietrich, 1987). RTMs offer a valuable tool for understanding processes controlling chemical weathering. RTMs require information on the parent rock such as mineralogy, porosity, and mineral surface areas; climate information such as rainfall and evapotranspiration (to determine average flow rates through the soil profile) and atmospheric CO_2 concentrations over geological time. Reaction thermodynamics (equilibrium constants) and kinetics are also needed for such simulations, where mineral dissolution and precipitation rates is often represented by the Transition State Theory (TST) rate laws with rate dependence on aqueous chemistry. Figure S1 shows the use of an 1D RTM (FLOTRAN) to understand and quantify the role of environmental conditions (flow, pCO_2) and mineral properties (e.g., reactive surface area) in controlling plagioclase weathering, e.g., plagioclase τ values for a 250 ka granitic

alluvium soil profile from Merced, CA, USA (circles, Figure S1) (Moore et al., 2012). Such RTMs can be developed with constraints from field data and can be extrapolated to different geochemical conditions to help elucidate reaction mechanisms and identify key controls of chemical weathering. RTMs can also be used to quantify weathering rates, casting the evolution of relevant properties over geological time scales (Navarre-Sitchler et al., 2011; Navarre-Sitchler et al., 2009).

Figure S1: Field (circles) and modeled plagioclase reaction fronts (lines) as τ values versus depth for different (A) flow rates or $p\text{CO}_2$, and (B) reactive surface areas for a Merced chronosequence profile (250 ka). The best fit model (thicker black solid line) compares well with the plagioclase field data (dots). The figure shows that the reaction front advance rate could be altered by flow and $p\text{CO}_2$ (A) while the thickness of the front could only be changed by reducing the reactive surface area (B) relative to BET surface area (modified from (Moore et al., 2012)).

Sidebar 2: CO_2 consumption by chemical weathering at the regional scale

According to predicted future anthropogenic emission scenarios, atmospheric CO_2 concentration is expected to double before the end of the 21st century, which will promote changes in hydrological cycles and vegetation cover. Although typically considered unimportant at the century scale, recent studies have demonstrated that chemical weathering can be highly sensitive to climate changes. A recent modeling study demonstrates that doubling CO_2 level will likely alter CO_2 consumption rates significantly for the Mackenzie River Basin, one of the most important Arctic watersheds (Beaulieu et al., 2012).

Figure S2. A. B-WITCH LPJ takes climate, lithology, and soil data to calculate the land cover and water and carbon cycles, the output of which is further fed into WITCH, a geochemical model that simulates mineral weathering and CO_2 consumption through transition state theory rate laws (Roelandt et al., 2010); B. Predicted spatial distribution of differences in CO_2 consumption rates when atmospheric CO_2 rises from 355 to 560 ppmv (Beaulieu et al., 2012).

The prediction was made using B-WITCH, a code that couples the dynamic global vegetation model LPJ with a process-based continental weathering model (Figure S2). Using spatially explicit

data on lithology, vegetation cover, soil properties, and climate conditions, the model reproduced field data of river chemistry (Beaulieu et al., 2010). Two simulations were carried out at the CO₂ levels of 355 and 560 ppmv to quantify the effects of carbon doubling. With an average CO₂ consumption rates of 0.06 mol/m²/yr, the predicted CO₂ consumption rates in the south (red area) that sits on carbonaceous shale increase by more than three times responding to the carbon doubling. In contrast, the northeastern watershed sees no change or up to a maximum of one time decrease. This emphasizes the importance of including chemical weathering in understanding the evolution of the global carbon cycle under changing climate conditions.

Sidebar 3: Using isotopes to identify key biogeochemical processes in marine sediments

Here a 1D RTM constrained by a comprehensive dataset (C, ³²S, ³⁴S, N, Fe) allows the identification of the complex reaction network associated with sulfur cycling in the organic-rich mud underlying the Namibian shelf waters (Dale et al., 2009). The sulfur dynamics are influenced by the bacterium *Thiomargarita namibiensis* that oxidizes H₂S to SO₄²⁻ using nitrate as the electron acceptor. The oxidized sulfate can further oxidize organic carbon (G) or methane (Figure S3). Detailed quantification of reaction rates indicated sulfate reduction (SR) and H₂S oxidation by nitrate (> 6.0 mol/S/m²/yr). As a result, only a small fraction (~ 0.3 wt.%) of total H₂S production is trapped

as sedimentary pyrite (FeS₂) and organic sulfur (S_{org}). A fractionation approaching 100‰ was observed, far higher than laboratory-based estimates for these processes. Mass balance calculations argue against the commonly believed view that this is caused by disproportionation of autochthonous elemental sulfur. Instead, the model indicates that repeated multi-step sulfide oxidation and intracellular disproportionation by *Thiomargarita* could, in principle, generate the measured isotopic fractionation.

References:

- Abbott, M., Warren, I., Jensen, K.H., Jønhc-Clausen, T. and Havnø, K., 1979. Coupling of unsaturated and saturated zone models, 19th Congress of the IAHR.
- Abbott, M.B., Bathurst, J.C., Cunge, J.A., O'Connell, P.E. and Rasmussen, J., 1986. An introduction to the European Hydrological System—Systeme Hydrologique European, "SHE", 1: History and philosophy of a physically-based, distributed modelling system. Journal of hydrology, 87(1): 45-59.
- Adams, E.E. and Gelhar, L.W., 1992. Field study of dispersion in a heterogeneous aquifer: 2. Spatial moments analysis. Water Resources Research, 28(12): 3293-3307.

- Albarede, F., 2004. The stable isotope geochemistry of copper and zinc, *Geochemistry of Non-Traditional Stable Isotopes. Reviews in Mineralogy & Geochemistry*, pp. 409-427.
- Allen-King, R.M., Grathwohl, P. and Ball, W.P., 2002. New modeling paradigms for the sorption of hydrophobic organic chemicals to heterogeneous carbonaceous matter in soils, sediments, and rocks. *Advances in Water Resources*, 25(8-12): 985-1016.
- Anbar, A.D., 2004. Molybdenum stable isotopes: Observations, interpretations and directions, *Geochemistry of Non-Traditional Stable Isotopes. Reviews in Mineralogy & Geochemistry*, pp. 429-454.
- Anderson, R.S., Anderson, S.P. and Tucker, G.E., 2013. Rock damage and regolith transport by frost: an example of climate modulation of the geomorphology of the critical zone. *Earth Surface Processes and Landforms*, 38(3): 299-316.
- Andrews, D.M., Lin, H., Zhu, Q., Jin, L. and Brantley, S.L., 2011. Hot Spots and Hot Moments of Dissolved Organic Carbon Export and Soil Organic Carbon Storage in the Shale Hills Catchment All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher. *Vadose Zone Journal*, 10(3).
- Apps, J.A., Zheng, L., Zhang, Y., Xu, T. and Birkholzer, J.T., 2010. Evaluation of Potential Changes in Groundwater Quality in Response to CO₂ Leakage from Deep Geologic Storage. *Transport in Porous Media*, 82(1): 215-246.
- Argerich, A., Haggerty, R., Martí, E., Sabater, F. and Zarnetske, J., 2011. Quantification of metabolically active transient storage (MATS) in two reaches with contrasting transient storage and ecosystem respiration. *Journal of Geophysical Research: Biogeosciences*, 116(G3): n/a-n/a.
- Arnold, J. and Soil, G., 1994. SWAT (Soil and Water Assessment Tool). Grassland, Soil and Water Research Laboratory, USDA, Agricultural Research Service.
- Atchley, A.L., Maxwell, R.M. and Navarre-Sitchler, A.K., 2013. Using streamlines to simulate stochastic reactive transport in heterogeneous aquifers: Kinetic metal release and transport in CO₂ impacted drinking water aquifers. *Advances in Water Resources*, 52(0): 93-106.
- Atchley, A.L., Navarre-Sitchler, A.K. and Maxwell, R.M., 2014. The effects of physical and geochemical heterogeneities on hydro-geochemical transport and effective reaction rates. *Journal of Contaminant Hydrology*, 165(0): 53-64.
- Banwart, S., Bernasconi, S.M., Bloem, J., Blum, W., Brandao, M., Brantley, S., Chabaux, F., Duffy, C., Kram, P., Lair, G., Lundin, L., Nikolaidis, N., Novak, M., Panagos, P., Ragnarsdottir, K.V., Reynolds, B., Rousseva, S., de Ruiter, P., van Gaans, P., van Riemsdijk, W., White, T. and Zhang, B., 2011. Soil Processes and Functions in Critical Zone Observatories: Hypotheses and Experimental Design. *Vadose Zone Journal*, 10(3): 974-987.
- Banwart, S.A., Berg, A. and Beerling, D.J., 2009. Process-based modeling of silicate mineral weathering responses to increasing atmospheric CO₂ and climate change. *Global Biogeochemical Cycles*, 23(4): n/a-n/a.
- Bao, C., Li, L., Shi, Y. and Duffy, C., 2016. Understanding Hydrogeochemical Processes at the Watershed Scale: 1. the development of process-based model RT-Flux-PIHM. *Water Resources Research*: (in revision).
- Bao, C., Wu, H., Li, L., Newcomer, D., Long, P.E. and Williams, K.H., 2014. Uranium Bioreduction Rates across Scales: Biogeochemical Hot Moments and Hot Spots during a Biostimulation Experiment at Rifle, Colorado. *Environmental Science & Technology*, 48(17): 10116-10127.

1215 Bardini, L., Boano, F., Cardenas, M.B., Revelli, R. and Ridolfi, L., 2012. Nutrient cycling in
1216 bedform induced hyporheic zones. *Geochimica et Cosmochimica Acta*, 84: 47-61.

1217 Barlage, M. and Zeng, X., 2004. Impact of observed vegetation root distribution on seasonal
1218 global simulations of land surface processes. *Journal of Geophysical Research:*
1219 *Atmospheres*, 109(D9): n/a-n/a.

1220 Barnett, T.P., Adam, J.C. and Lettenmaier, D.P., 2005. Potential impacts of a warming
1221 climate on water availability in snow-dominated regions. *Nature*, 438(7066): 303-309.

1222 Barry, D.A., Prommer, H., Miller, C.T., Engesgaard, P., Brun, A. and Zheng, C., 2002.
1223 Modelling the fate of oxidisable organic contaminants in groundwater. *Advances in*
1224 *Water Resources*, 25(8-12): 945-983.

1225 Bassu, S., Brisson, N., Durand, J.L., Boote, K., Lizaso, J., Jones, J.W., Rosenzweig, C.,
1226 Ruane, A.C., Adam, M. and Baron, C., 2014. How do various maize crop models vary
1227 in their responses to climate change factors? *Global Change Biology*, 20(7): 2301-
1228 2320.

1229 Basu, N.B., Rao, P.S.C., Winzeler, H.E., Kumar, S., Owens, P. and Merwade, V., 2010.
1230 Parsimonious modeling of hydrologic responses in engineered watersheds: Structural
1231 heterogeneity versus functional homogeneity. *Water Resources Research*, 46(4): n/a-
1232 n/a.

1233 Bazilevskaya, E., Lebedeva, M., Pavich, M., Rother, G., Parkinson, D.Y., Cole, D. and
1234 Brantley, S.L., 2013. Where fast weathering creates thin regolith and slow weathering
1235 creates thick regolith. *Earth Surface Processes and Landforms*, 38(8): 847-858.

1236 Beaulieu, E., Godderis, Y., Donnadieu, Y., Labat, D. and Roelandt, C., 2012. High sensitivity
1237 of the continental-weathering carbon dioxide sink to future climate change. *Nature*
1238 *Clim. Change*, 2(5): 346-349.

1239 Beaulieu, E., Goddris, Y., Labat, D., Roelandt, C., Calmels, D. and Gaillardet, J., 2011.
1240 Modeling of water-rock interaction in the Mackenzie basin: Competition between
1241 sulfuric and carbonic acids. *Chemical Geology*, 289(1-2): 114-123.

1242 Beaulieu, E., Goddris, Y., Labat, D., Roelandt, C., Oliva, P. and Guerrero, B., 2010. Impact
1243 of atmospheric CO₂ levels on continental silicate weathering. *Geochemistry,*
1244 *Geophysics, Geosystems*, 11(7): Q07007.

1245 Beerling, D.J. and Berner, R.A., 2005. Feedbacks and the coevolution of plants and
1246 atmospheric CO₂. *Proceedings of the National Academy of Sciences of the United*
1247 *States of America*, 102(5): 1302-1305.

1248 Beisman, J., Maxwell, R., Navarre-Sitchler, A., Steefel, C. and Molins, S., 2015.
1249 ParCrunchFlow: an efficient, parallel reactive transport simulation tool for physically
1250 and chemically heterogeneous saturated subsurface environments. *Computational*
1251 *Geosciences*: 1-20.

1252 Benson, D.A., Wheatcraft, S.W. and Meerschaert, M.M., 2000a. Application of a fractional
1253 advection-dispersion equation. *Water Resources Research*, 36(6): 1403-1412.

1254 Benson, D.A., Wheatcraft, S.W. and Meerschaert, M.M., 2000b. The fractional-order
1255 governing equation of Lvy Motion. *Water Resources Research*, 36(6): 1413-1423.

1256 Berkowitz, B., Cortis, A., Dentz, M. and Scher, H., 2006. Modeling non-Fickian transport in
1257 geological formations as a continuous time random walk. *Reviews of Geophysics*,
1258 44(2): RG2003.

1259 Berner, R.A., 2006. GEOCARBSULF: A combined model for Phanerozoic atmospheric O₂
1260 and CO₂. *Geochimica et Cosmochimica Acta*, 70(23): 5653-5664.

1261 Berner, R.A. and Caldeira, K., 1997. The need for mass balance and feedback in the
1262 geochemical carbon cycle. *Geology*, 25(10): 955-956.

1263 Berner, R.A. and Kothavala, Z., 2001. Geocarb III: A Revised Model of Atmospheric CO₂
1264 over Phanerozoic Time. *American Journal of Science*, 301(2): 182-204.

1265 Berner, R.A., Lasaga, A.C. and Garrels, R.M., 1983. The carbonate-silicate geochemical
 1266 cycle and its effect on atmospheric carbon-dioxide over the past 100 million years.
 1267 American Journal of Science, 283(7): 641-683.
 1268 Bethke, C., 1996. Geochemical reaction modeling : concepts and applications. Oxford
 1269 University Press, New York, 397 pp.
 1270 Bethke, C.M., Ding, D., Jin, Q.S. and Sanford, R.A., 2008. Origin of microbiological zoning
 1271 in groundwater flows. Geology, 36(9): 739-742.
 1272 Beven, K., 1989. Changing ideas in hydrology—the case of physically-based models. Journal
 1273 of hydrology, 105(1): 157-172.
 1274 Beven, K., 2001a. How far can we go in distributed hydrological modelling? Hydrology and
 1275 Earth System Sciences, 5(1): 1-12.
 1276 Beven, K., 2001b. On stochastic models and the single realization. Hydrological Processes,
 1277 15(5): 895-896.
 1278 Beven, K. and Binley, A., 1992. THE FUTURE OF DISTRIBUTED MODELS - MODEL
 1279 CALIBRATION AND UNCERTAINTY PREDICTION. Hydrological Processes,
 1280 6(3): 279-298.
 1281 Beven, K. and Germann, P., 1982. MACROPORES AND WATER-FLOW IN SOILS. Water
 1282 Resources Research, 18(5): 1311-1325.
 1283 Bindoff, N.L., Stott, P.A., AchutaRao, K.M., Allen, M.R., Gillett, N., Gutzler, D., Hansingo,
 1284 K., Hegerl, G., Hu, Y., Jain, S., Mokhov, I.I., Overland, J., Perlwitz, J., Sebbari, R.
 1285 and Zhang, X., 2013. Detection and Attribution of Climate Change: from Global to
 1286 Regional. In: T.F. Stocker, D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J.
 1287 Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (Editors), Climate Change
 1288 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth
 1289 Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge
 1290 University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 867–
 1291 952.
 1292 Binley, A., Hubbard, S.S., Huisman, J.A., Revil, A., Robinson, D.A., Singha, K. and Slater,
 1293 L.D., 2015. The emergence of hydrogeophysics for improved understanding of
 1294 subsurface processes over multiple scales. Water Resources Research, 51(6): 3837-
 1295 3866.
 1296 Blum, A.E. and Stillings, L.L., 1995. Feldspar dissolution kinetics. Reviews in Mineralogy
 1297 and Geochemistry, 31(1): 291-351.
 1298 Blunt, M.J., Bijeljic, B., Dong, H., Gharbi, O., Iglauer, S., Mostaghimi, P., Paluszny, A. and
 1299 Pentland, C., 2013. Pore-scale imaging and modelling. Advances in Water Resources,
 1300 51: 197-216.
 1301 Boano, F., Harvey, J.W., Marion, A., Packman, A.I., Revelli, R., Ridolfi, L. and Wörman, A.,
 1302 2014. Hyporheic flow and transport processes: Mechanisms, models, and
 1303 biogeochemical implications. Reviews of Geophysics, 52(4): 603-679.
 1304 Boggs, J.M., Young, S.C., Beard, L.M., Gelhar, L.W., Rehfeldt, K.R. and Adams, E.E., 1992.
 1305 Field study of dispersion in a heterogeneous aquifer: 1. Overview and site description.
 1306 Water Resources Research, 28(12): 3281-3291.
 1307 Bolton, E.W., Berner, R.A. and Petsch, S.T., 2006. The Weathering of Sedimentary Organic
 1308 Matter as a Control on Atmospheric O₂: II. Theoretical Modeling. American Journal
 1309 of Science, 306(8): 575-615.
 1310 Bolton, E.W., Lasaga, A.C. and Rye, D.M., 1996. A model for the kinetic control of quartz
 1311 dissolution and precipitation in porous media flow with spatially variable
 1312 permeability: Formulation and examples of thermal convection. Journal of
 1313 Geophysical Research: Solid Earth, 101(B10): 22157-22187.

1314 Bolton, E.W., Lasaga, A.C. and Rye, D.M., 1997. Dissolution and precipitation via forced-
1315 flux injection in a porous medium with spatially variable permeability: Kinetic control
1316 in two dimensions. *Journal of Geophysical Research: Solid Earth*, 102(B6): 12159-
1317 12171.

1318 Bolton, E.W., Lasaga, A.C. and Rye, D.M., 1999. Long-term flow/chemistry feedback in a
1319 porous medium with heterogeneous permeability: kinetic control of dissolution and
1320 precipitation. *Am. J. Sci.*, 299: 1-68.

1321 Bond-Lamberty, B. and Thomson, A., 2010. Temperature-associated increases in the global
1322 soil respiration record. *Nature*, 464(7288): 579-582.

1323 Bouchez, J., von Blanckenburg, F. and Schuessler, J.A., 2013. Modeling novel stable isotope
1324 ratios in the weathering zone. *American Journal of Science*, 313(4): 267-308.

1325 Bouma, T., Nielsen, K.L., Van Hal, J. and Koutstaal, B., 2001. Root system topology and
1326 diameter distribution of species from habitats differing in inundation frequency.
1327 *Functional Ecology*, 15(3): 360-369.

1328 Bouskill, N.J., Tang, J., Riley, W.J. and Brodie, E.L., 2012. Trait-based representation of
1329 biological nitrification: model development, testing, and predicted community
1330 composition. *The microbial nitrogen cycle*: 116.

1331 Brantley, S., DiBiase, R., Russo, T., Shi, Y., Lin, H., Davis, K., Kaye, M., Hill, L., Kaye, J.,
1332 Eissenstat, D., Hoagland, B., Dere, A., Neal, A., Brubaker, K. and Arthur, D., 2016.
1333 Designing a suite of measurements to understand the critical zone. *Earth Surface*
1334 *Dynamics*, 4: 1-25.

1335 Brantley, S.L., Goldhaber, M.B. and Ragnarsdottir, K.V., 2007. Crossing Disciplines and
1336 Scales to Understand the Critical Zone. *Elements*, 3(5): 307-314.

1337 Brantley, S.L., Holleran, M.E., Jin, L. and Bazilevskaya, E., 2013a. Probing deep weathering
1338 in the Shale Hills Critical Zone Observatory, Pennsylvania (USA): the hypothesis of
1339 nested chemical reaction fronts in the subsurface. *Earth Surface Processes and*
1340 *Landforms*, 38(11): 1280-1298.

1341 Brantley, S.L. and Lebedeva, M., 2011. Learning to Read the Chemistry of Regolith to
1342 Understand the Critical Zone. In: R. Jeanloz and K.H. Freeman (Editors), *Annual*
1343 *Review of Earth and Planetary Sciences*, Vol 39. *Annual Review of Earth and*
1344 *Planetary Sciences*, pp. 387-416.

1345 Brantley, S.L., Lebedeva, M. and Bazilevskaya, E., 2013b. Relating weathering fronts for
1346 acid neutralization and oxidation to pCO₂ and pO₂. In: J. Farquhar, Kasting, J.,
1347 Canfield, D. (Editor), *Treatise of Geochemistry. The Atmosphere -- History*. Elsevier,
1348 Amsterdam, The Netherlands.

1349 Brantley, S.L., Liermann, L.J., Gynn, R.L., Anbar, A., Icopini, G.A. and Barling, J., 2004.
1350 Fe isotopic fractionation during mineral dissolution with and without bacteria.
1351 *Geochimica Et Cosmochimica Acta*, 68(15): 3189-3204.

1352 Brantley, S.L. and White, A.F., 2009. Approaches to Modeling Weathered Regolith.
1353 *Thermodynamics and Kinetics of Water-Rock Interaction*, 70: 435-484.

1354 Brooks, P.D., Chorover, J., Fan, Y., Godsey, S.E., Maxwell, R.M., McNamara, J.P. and
1355 Tague, C., 2015. Hydrological partitioning in the critical zone: Recent advances and
1356 opportunities for developing transferrable understanding of water cycle dynamics.
1357 *Water Resources Research*: n/a-n/a.

1358 Brown, B. and Rolston, D., 1980. Transport and transformation of methyl bromide in soils.
1359 *Soil science*, 130(2): 68-75.

1360 Brunet, J.-P.L., Li, L., Karpyn, Z.T. and Huerta, N.J., 2016. Fracture opening or self-sealing:
1361 Critical residence time as a unifying parameter for cement-CO₂-brine interactions.
1362 *International Journal of Greenhouse Gas Control*, 47: 25-37.

1363 Brunner, P. and Simmons, C.T., 2012. HydroGeoSphere: A Fully Integrated, Physically
1364 Based Hydrological Model. *Ground Water*, 50(2): 170-176.

1365 Bultreys, T., De Boever, W. and Cnudde, V., 2016. Imaging and image-based fluid transport
1366 modeling at the pore scale in geological materials: A practical introduction to the
1367 current state-of-the-art. *Earth-Science Reviews*, 155: 93-128.

1368 Buss, H.L., Sak, P.B., Webb, S.M. and Brantley, S.L., 2008. Weathering of the Rio Blanco
1369 quartz diorite, Luquillo Mountains, Puerto Rico: Coupling oxidation, dissolution, and
1370 fracturing. *Geochimica et Cosmochimica Acta*, 72(18): 4488-4507.

1371 Camargo, G. and Kemanian, A., 2016. Six crop models differ in their simulation of water
1372 uptake. *Agricultural and Forest Meteorology*, 220: 116-129.

1373 Carey, A.E., Lyons, W.B. and Owen, J.S., 2005. Significance of landscape age, uplift, and
1374 weathering rates to ecosystem development. *Aquatic Geochemistry*, 11: 215-239.

1375 Carroll, S., Carey, J.W., Dzombak, D., Huerta, N.J., Li, L., Richard, T., Um, W., Walsh,
1376 S.D.C. and Zhang, L., 2016. Review: Role of chemistry, mechanics, and transport on
1377 well integrity in CO₂ storage environments. *International Journal of Greenhouse Gas*
1378 *Control*, 49: 149-160.

1379 Chapman, B.M., 1982. Numerical simulation of the transport and speciation of
1380 nonconservative chemical reactants in rivers. *Water Resources Research*, 18(1): 155-
1381 167.

1382 Chapman, B.M., James, R.O., Jung, R.F. and Washington, H.G., 1982. Modelling the
1383 transport of reacting chemical contaminants in natural streams. *Marine and*
1384 *Freshwater Research*, 33(4): 617-628.

1385 Chapman, J.B., Weiss, D.J., Shan, Y. and Lemburger, M., 2009. Iron isotope fractionation
1386 during leaching of granite and basalt by hydrochloric and oxalic acids. *Geochimica Et*
1387 *Cosmochimica Acta*, 73(5): 1312-1324.

1388 Chattopadhyay, N. and Hulme, M., 1997. Evaporation and potential evapotranspiration in
1389 India under conditions of recent and future climate change. *Agricultural and Forest*
1390 *Meteorology*, 87(1): 55-73.

1391 Chen, A., Darbon, J. and Morel, J.-M., 2014. Landscape evolution models: A review of their
1392 fundamental equations. *Geomorphology*, 219: 68-86.

1393 Chen, B.H., Wu, F.H. and Liu, H.H., 2001. Study on microbes and their effects on rare earth
1394 extraction in weathering crust of granite. *Journal of Rare Earths*, 19(3): 161-167.

1395 Chen, J., Gu, B.H., Royer, R.A. and Burgos, W.D., 2003. The roles of natural organic matter
1396 in chemical and microbial reduction of ferric iron. *Science of the Total Environment*,
1397 307(1-3): 167-178.

1398 Cheng, Y., Hubbard, C.G., Li, L., Bouskill, N., Molins, S., Zheng, L., Sonnenthal, E.,
1399 Conrad, M.E., Engelbrektson, A., Coates, J.D. and Ajo-Franklin, J.B., 2016. Reactive
1400 Transport Model of Sulfur Cycling as Impacted by Perchlorate and Nitrate
1401 Treatments. *Environmental Science & Technology*, 50(13): 7010–7018.

1402 Chiogna, G., Cirpka, O.A., Grathwohl, P. and Rolle, M., 2011. Transverse mixing of
1403 conservative and reactive tracers in porous media: Quantification through the
1404 concepts of flux-related and critical dilution indices. *Water Resources Research*, 47.

1405 Cirpka, O.A., Frind, E.O. and Helmig, R., 1999. Numerical simulation of biodegradation
1406 controlled by transverse mixing. *Journal of Contaminant Hydrology*, 40(2): 159-182.

1407 Cirpka, O.A. and Valocchi, A.J., 2007. Two-dimensional concentration distribution for
1408 mixing-controlled bioreactive transport in steady state. *Advances in Water Resources*,
1409 30(6-7): 1668-1679.

1410 Clark, M.P., Fan, Y., Lawrence, D.M., Adam, J.C., Bolster, D., Gochis, D.J., Hooper, R.P.,
1411 Kumar, M., Leung, L.R., Mackay, D.S., Maxwell, R.M., Shen, C., Swenson, S.C. and

- Zeng, X., 2015. Improving the representation of hydrologic processes in Earth System Models. *Water Resources Research*: n/a-n/a.
- Colbourn, G., Ridgwell, A. and Lenton, T.M., 2013. The Rock Geochemical Model (RokGeM) v0.9. *Geosci. Model Dev.*, 6(5): 1543-1573.
- Conant, R.T., Ryan, M.G., Ågren, G.I., Birge, H.E., Davidson, E.A., Eliasson, P.E., Evans, S.E., Frey, S.D., Giardina, C.P., Hopkins, F.M., Hyvönen, R., Kirschbaum, M.U.F., Lavallee, J.M., Leifeld, J., Parton, W.J., Megan Steinweg, J., Wallenstein, M.D., Martin Wetterstedt, J.Å. and Bradford, M.A., 2011. Temperature and soil organic matter decomposition rates – synthesis of current knowledge and a way forward. *Global Change Biology*, 17(11): 3392-3404.
- Crosby, H.A., Roden, E.E., Johnson, C.M. and Beard, B.L., 2007. The mechanisms of iron isotope fractionation produced during dissimilatory Fe(III) reduction by *Shewanella putrefaciens* and *Geobacter sulfurreducens*. *Geobiology*, 5(2): 169-189.
- Dagan, G., Fiori, A. and Jankovic, I., 2013. Upscaling of flow in heterogeneous porous formations: Critical examination and issues of principle. *Advances in Water Resources*, 51: 67-85.
- Dale, A.W., Brüchert, V., Alperin, M. and Regnier, P., 2009. An integrated sulfur isotope model for Namibian shelf sediments. *Geochimica et Cosmochimica Acta*, 73(7): 1924-1944.
- Dale, A.W., Regnier, P., Knab, N.J., Jorgensen, B.B. and Van Cappellen, P., 2008. Anaerobic oxidation of methane (AOM) in marine sediments from the Skagerrak (Denmark): II. Reaction-transport modeling. *Geochimica Et Cosmochimica Acta*, 72(12): 2880-2894.
- Daley, R., 1991. *Atmospheric Data Analysis*. Cambridge Atmospheric and Space Science Series. Cambridge University Press.
- Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature*, 440(7081): 165-173.
- Davison, J., Hwang, H., Sudicky, E. and Lin, J., 2014. Development of a fully integrated water cycle model: HydroGeoSphere-Weather Research and Forecasting (HGS-WRF), AGU Fall Meeting Abstracts, pp. 0915.
- De Kauwe, M.G., Medlyn, B.E., Zaehle, S., Walker, A.P., Dietze, M.C., Wang, Y.-P., Luo, Y., Jain, A.K., El-Masri, B., Hickler, T., Wårlind, D., Weng, E., Parton, W.J., Thornton, P.E., Wang, S., Prentice, I.C., Asao, S., Smith, B., McCarthy, H.R., Iversen, C.M., Hanson, P.J., Warren, J.M., Oren, R. and Norby, R.J., 2014. Where does the carbon go? A model–data intercomparison of vegetation carbon allocation and turnover processes at two temperate forest free-air CO₂ enrichment sites. *New Phytologist*, 203(3): 883-899.
- Dellinger, M., Gaillardet, J., Bouchez, J., Calmels, D., Galy, V., Hilton, R.G., Louvat, P. and France-Lanord, C., 2014. Lithium isotopes in large rivers reveal the cannibalistic nature of modern continental weathering and erosion. *Earth and Planetary Science Letters*, 401: 359-372.
- Dellinger, M., Gaillardet, J., Bouchez, J., Calmels, D., Louvat, P., Dosseto, A., Gorge, C., Alanoca, L. and Maurice, L., 2015. Riverine Li isotope fractionation in the Amazon River basin controlled by the weathering regimes. *Geochimica Et Cosmochimica Acta*, 164: 71-93.
- Denef, K., Zotarelli, L., Boddey, R.M. and Six, J., 2007. Microaggregate-associated carbon as a diagnostic fraction for management-induced changes in soil organic carbon in two Oxisols. *Soil Biology and Biochemistry*, 39(5): 1165-1172.
- Deng, H., Dai, Z., Wolfsberg, A.V., Ye, M., Stauffer, P.H., Lu, Z. and Kwicklis, E., 2012. Upscaling retardation factor in hierarchical porous media with multimodal reactive mineral facies. *Chemosphere*.

1462 DePaolo, D., 2006. Isotopic effects in fracture-dominated reactive fluid-rock systems.
 1463 *Geochimica et Cosmochimica Acta*, 70: 1077-1096.
 1464 Dietrich, W.E., Reiss, R., Hsu, M.L. and Montgomery, D.R., 1995. A process-based model
 1465 for colluvial soil depth and shallow landsliding using digital elevation data.
 1466 *Hydrological processes*, 9(3-4): 383-400.
 1467 Diggle, A.J., 1988. ROOTMAP—a model in three-dimensional coordinates of the growth and
 1468 structure of fibrous root systems. *Plant and Soil*, 105(2): 169-178.
 1469 Dijkstra, P., LaViolette, C.M., Coyle, J.S., Doucett, R.R., Schwartz, E., Hart, S.C. and
 1470 Hungate, B.A., 2008. N-15 enrichment as an integrator of the effects of C and N on
 1471 microbial metabolism and ecosystem function. *Ecology Letters*, 11(4): 389-397.
 1472 DOE, U.S., 2015. Building Virtual Ecosystems: Computational Challenges for Mechanistic
 1473 Modeling of Terrestrial Environments: Workshop Report. DOE/SC-0171. U.S.
 1474 Department of Energy Office of Science.
 1475 <http://doesbr.org/BuildingVirtualEcosystems/>.
 1476 Donigian Jr, A., Bicknell, B., Imhoff, J. and Singh, V., 1995. Hydrological Simulation
 1477 Program-Fortran (HSPF). Computer models of watershed hydrology.: 395-442.
 1478 Drever, J., 1994. The effect of land plants on weathering rates of silicate minerals.
 1479 *Geochimica et Cosmochimica Acta*, 58(10): 2325-2332.
 1480 Drever, J. and Clow, D., 1995. Weathering rates in catchments. *Reviews in Mineralogy and*
 1481 *Geochemistry*, 31(1): 463-483.
 1482 Drever, J.I., 1989. *The Geochemistry of Natural Waters*. Prentice-Hall, Inc., Englewood
 1483 Cliffs, NJ, 388 pp.
 1484 Drever, J.I., 1997. Weathering Processes. In: O.M. Saether and d.C. P. (Editors),
 1485 *Geochemical processes, weathering and groundwater recharge in catchments*. A.A.
 1486 Balkema, pp. 400.
 1487 Druhan, J.L. and Maher, K., 2014. A Model Linking Stable Isotope Fractionation to Water
 1488 Flux and Transit Times in Heterogeneous Porous Media. *Procedia Earth and Planetary*
 1489 *Science*, 10(0): 179-188.
 1490 Druhan, J.L., Steefel, C.I., Conrad, M.E. and DePaolo, D.J., 2014. A large column analog
 1491 experiment of stable isotope variations during reactive transport: I. A comprehensive
 1492 model of sulfur cycling and delta S-34 fractionation. *Geochimica Et Cosmochimica*
 1493 *Acta*, 124: 366-393.
 1494 Druhan, J.L., Steefel, C.I., Molins, S., Williams, K.H., Conrad, M.E. and DePaolo, D.J.,
 1495 2012. Timing the Onset of Sulfate Reduction over Multiple Subsurface Acetate
 1496 Amendments by Measurement and Modeling of Sulfur Isotope Fractionation.
 1497 *Environmental Science & Technology*, 46(16): 8895-8902.
 1498 Druhan, J.L., Steefel, C.I., Williams, K.H. and DePaolo, D.J., 2013. Calcium isotope
 1499 fractionation in groundwater: Molecular scale processes influencing field scale
 1500 behavior. *Geochimica Et Cosmochimica Acta*, 119: 93-116.
 1501 Duffy, C., Shi, Y., Davis, K., Slingerland, R., Li, L., Sullivan, P., Goddris, Y. and Brantley,
 1502 S.L., 2014. Designing a Suite of Models to Explore Critical Zone Function. *Procedia*
 1503 *Earth and Planetary Science*, 10: 7-15.
 1504 Dungait, J.A.J., Hopkins, D.W., Gregory, A.S. and Whitmore, A.P., 2012. Soil organic matter
 1505 turnover is governed by accessibility not recalcitrance. *Global Change Biology*, 18(6):
 1506 1781-1796.
 1507 Dunn, R.M., Mikola, J., Bol, R. and Bardgett, R.D., 2006. Influence of microbial activity on
 1508 plant-microbial competition for organic and inorganic nitrogen. *Plant and Soil*, 289(1-
 1509 2): 321-334.

1510 Dupré, B., Dessert, C., Oliva, P., Goddérès, Y., Viers, J., François, L., Millot, R. and
 1511 Gaillardet, J., 2003. Rivers, chemical weathering and Earth's climate. *Comptes*
 1512 *Rendus Geoscience*, 335(16): 1141-1160.
 1513 Ebel, B.A. and Loague, K., 2006. Physics-based hydrologic-response simulation: Seeing
 1514 through the fog of equifinality. *Hydrological Processes*, 20(13): 2887-2900.
 1515 Eckert, D., Rolle, M. and Cirpka, O.A., 2012. Numerical simulation of isotope fractionation
 1516 in steady-state bioreactive transport controlled by transverse mixing. *Journal of*
 1517 *Contaminant Hydrology*, 140: 95-106.
 1518 Edmonds, H.N. and Edmond, J.M., 1995. A 3-Component Mixing Model for Ridge-Crest
 1519 Hydrothermal Fluids. *Earth and Planetary Science Letters*, 134(1-2): 53-67.
 1520 Eusterhues, K., Rumpel, C., Kleber, M. and Kogel-Knabner, I., 2003. Stabilisation of soil
 1521 organic matter by interactions with minerals as revealed by mineral dissolution and
 1522 oxidative degradation. *Organic Geochemistry*, 34(12): 1591-1600.
 1523 Fan, Y., 2015. Groundwater in the Earth's critical zone: Relevance to large-scale patterns and
 1524 processes. *Water Resources Research*, 51(5): 3052-3069.
 1525 Fan, Y., Li, H. and Miguez-Macho, G., 2013. Global Patterns of Groundwater Table Depth.
 1526 *Science*, 339(6122): 940-943.
 1527 Fan, Y., Richard, S., Bristol, R.S., Peters, S.E., Ingebritsen, S.E., Moosdorf, N., Packman, A.,
 1528 Gleeson, T., Zaslavsky, I., Peckham, S., Murdoch, L., Fienen, M., Cardiff, M.,
 1529 Tarboton, D., Jones, N., Hooper, R., Arrigo, J., Gochis, D., Olson, J. and Wolock, D.,
 1530 2015. DigitalCrust – a 4D data system of material properties for transforming research
 1531 on crustal fluid flow. *Geofluids*, 15(1-2): 372-379.
 1532 Fang, Y., Scheibe, T.D., Mahadevan, R., Garg, S., Long, P.E. and Lovley, D.R., 2011a.
 1533 Direct coupling of a genome-scale microbial in silico model and a groundwater
 1534 reactive transport model. *Journal of Contaminant Hydrology*, 122(1-4): 96-103.
 1535 Fang, Y., Scheibe, T.D., Mahadevan, R., Garg, S., Long, P.E. and Lovley, D.R., 2011b.
 1536 Direct coupling of a genome-scale microbial in silico model and a groundwater
 1537 reactive transport model. *Journal of contaminant hydrology*, 122(1): 96-103.
 1538 Fantle, M.S. and DePaolo, D.J., 2007. Ca isotopes in carbonate sediment and pore fluid from
 1539 ODP Site 807A: The $\text{Ca}^{2+}(\text{aq})$ -calcite equilibrium fractionation factor and calcite
 1540 recrystallization rates in Pleistocene sediments. *Geochimica Et Cosmochimica Acta*,
 1541 71(10): 2524-2546.
 1542 Fantle, M.S., Maher, K.M. and DePaolo, D.J., 2010. Isotopic approaches for quantifying the
 1543 rates of marine burial diagenesis. . *Reviews of Geophysics*, 48: RG3002.
 1544 Feist, A.M., Herrgard, M.J., Thiele, I., Reed, J.L. and Palsson, B.O., 2009. Reconstruction of
 1545 biochemical networks in microorganisms. *Nat Rev Micro*, 7(2): 129-143.
 1546 Fenwick, D., Scheidt, C. and Caers, J., 2014. Quantifying Asymmetric Parameter Interactions
 1547 in Sensitivity Analysis: Application to Reservoir Modeling. *Mathematical*
 1548 *Geosciences*, 46(4): 493-511.
 1549 Fierer, N. and Schimel, J.P., 2002. Effects of drying-rewetting frequency on soil carbon and
 1550 nitrogen transformations. *Soil Biology & Biochemistry*, 34(6): 777-787.
 1551 Fierer, N., Schimel, J.P. and Holden, P.A., 2003. Variations in microbial community
 1552 composition through two soil depth profiles. *Soil Biology and Biochemistry*, 35(1):
 1553 167-176.
 1554 Finzi, A.C., Abramoff, R.Z., Spiller, K.S., Brzostek, E.R., Darby, B.A., Kramer, M.A. and
 1555 Phillips, R.P., 2015. Rhizosphere processes are quantitatively important components
 1556 of terrestrial carbon and nutrient cycles. *Global Change Biology*, 21(5): 2082-2094.
 1557 Fiori, A., Boso, F., de Barros, F.P.J., De Bartolo, S., Frampton, A., Severino, G., Suweis, S.
 1558 and Dagan, G., 2010. An indirect assessment on the impact of connectivity of

conductivity classes upon longitudinal asymptotic macrodispersivity. *Water Resources Research*, 46(8): W08601.

Fontaine, S., Barot, S., Barre, P., Bdioui, N., Mary, B. and Rumpel, C., 2007. Stability of organic carbon in deep soil layers controlled by fresh carbon supply. *Nature*, 450(7167): 277-280.

Fredrickson, J.K., Romine, M.F., Beliaev, A.S., Auchtung, J.M., Driscoll, M.E., Gardner, T.S., Nealson, K.H., Osterman, A.L., Pinchuk, G., Reed, J.L., Rodionov, D.A., Rodrigues, J.L.M., Saffarini, D.A., Serres, M.H., Spormann, A.M., Zhulin, I.B. and Tiedje, J.M., 2008. Towards environmental systems biology of *Shewanella*. *Nat Rev Micro*, 6(8): 592-603.

Freeze, R.A. and Harlan, R., 1969. Blueprint for a physically-based, digitally-simulated hydrologic response model. *Journal of Hydrology*, 9(3): 237-258.

Gaillardet, J., Dupré, B., Louvat, P. and Allègre, C.J., 1999. Global silicate weathering and CO₂ consumption rates deduced from the chemistry of large rivers. *Chemical Geology*, 159(1-4): 3-30.

Gan, T.Y., Gusev, Y., Burges, S.J., Nasonova, O., Andréassian, V., Hall, A., Chahinian, N. and Schaake, J., 2006. Performance comparison of a complex physics-based land surface model and a conceptual, lumped-parameter hydrological model at the basin-scale. Large sample basin experiments for hydrological model parameterization: results of the model parameter experiment (MOPEX): 196-208.

Garré, S., Pagès, L., Laloy, E., Javaux, M., Vanderborght*, J. and Vereecken, H., 2012. Parameterizing a Dynamic Architectural Model of the Root System of Spring Barley from Minirhizotron Data. *Vadose Zone Journal*, 11(4).

Gelhar, L.W. and Axness, C.L., 1983. Three-dimensional stochastic analysis of macrodispersion in aquifers. *Water Resources Research*, 19(1): 161-180.

Georgiou, K., Koven, C.D., Riley, W.J. and Torn, M.S., 2015. Towards improved model structures for analyzing priming: potential pitfalls of using bulk turnover time. *Global Change Biology*: n/a-n/a.

Gerke, H.H. and Van Genuchten, M.T., 1993. A dual-porosity model for simulating the preferential movement of water and solutes in structured porous media. *Water Resources Research*, 29(2): 305-319.

Gibson, B.D., Amos, R.T. and Blowes, D.W., 2011. S-34/S-32 Fractionation during Sulfate Reduction in Groundwater Treatment Systems: Reactive Transport Modeling. *Environmental Science & Technology*, 45(7): 2863-2870.

Goddéris, Y. and Brantley, S.L., 2013. Earthcasting the future Critical Zone. *Elem. Sci. Anth.*, 1(1): 000019.

Goddéris, Y., Brantley, S.L., Francois, L.M., Schott, J., Pollard, D., Deque, M. and Dury, M., 2013. Rates of consumption of atmospheric CO₂ through the weathering of loess during the next 100 yr of climate change. *Biogeosciences*, 10(1): 135-148.

Goddéris, Y., Francois, L.M., Probst, A., Schott, J., Moncoulon, D., Labat, D. and Viville, D., 2006. Modelling weathering processes at the catchment scale: The WITCH numerical model. *Geochimica et Cosmochimica Acta*, 70: 1128-1147.

Goddéris, Y., Williams, J.Z., Schott, J., Pollard, D. and Brantley, S.L., 2010. Time evolution of the mineralogical composition of Mississippi Valley loess over the last 10 kyr: Climate and geochemical modeling. *Geochimica Et Cosmochimica Acta*, 74(22): 6357-6374.

Godsey, S.E., Kirchner, J.W. and Clow, D.W., 2009. Concentration–discharge relationships reflect chemostatic characteristics of US catchments. *Hydrological Processes*, 23(13): 1844-1864.

1608 Goren, L., Willett, S.D., Herman, F. and Braun, J., 2014. Coupled numerical–analytical
1609 approach to landscape evolution modeling. *Earth Surface Processes and Landforms*,
1610 39(4): 522-545.

1611 Guggenberger, G. and Kaiser, K., 2003. Dissolved organic matter in soil: challenging the
1612 paradigm of sorptive preservation. *Geoderma*, 113(3–4): 293-310.

1613 Guo, J., Wang, F., Vogt, R.D., Zhang, Y. and Liu, C.-Q., 2015. Anthropogenically enhanced
1614 chemical weathering and carbon evasion in the Yangtze Basin. *Scientific Reports*, 5:
1615 11941.

1616 Haberer, C.M., Rolle, M., Liu, S., Cirpka, O.A. and Grathwohl, P., 2011. A high-resolution
1617 non-invasive approach to quantify oxygen transport across the capillary fringe and
1618 within the underlying groundwater. *Journal of Contaminant Hydrology*, 122(1–4): 26-
1619 39.

1620 Haggerty, R. and Gorelick, S.M., 1995. multiple-rate mass transfer for modeling diffusion
1621 and surface reactions in media with pore-scale heterogeneity. *Water Resources*
1622 *Research*, 31(10): 2383-2400.

1623 Harvey, J. and Gooseff, M., 2015. River corridor science: Hydrologic exchange and
1624 ecological consequences from bedforms to basins. *Water Resources Research*: n/a-
1625 n/a.

1626 Harvey, J.W., Böhlke, J.K., Voytek, M.A., Scott, D. and Tobias, C.R., 2013. Hyporheic zone
1627 denitrification: Controls on effective reaction depth and contribution to whole-stream
1628 mass balance. *Water Resources Research*, 49(10): 6298-6316.

1629 Hasenmueller, E.A., Jin, L., Stinchcomb, G.E., Lin, H., Brantley, S.L. and Kaye, J.P., 2015.
1630 Topographic controls on the depth distribution of soil CO₂ in a small temperate
1631 watershed. *Applied Geochemistry*, 63: 58-69.

1632 Hausrath, E.M., Navarre-Sitchler, A.K., Sak, P., Steefel, C.I. and Brantley, S.L., 2008. Basalt
1633 weathering rates on Earth and the duration of liquid water on the plains of Gusev
1634 Crater, Mars. *Geology*, 36: 67-70.

1635 Hausrath, E.M. and Olsen, A.A., 2013. Using the chemical composition of carbonate rocks
1636 on Mars as a record of secondary interaction with liquid water. *American*
1637 *Mineralogist*, 98(5-6): 897-906.

1638 Hedges, J.I., Eglinton, G., Hatcher, P.G., Kirchman, D.L., Arnosti, C., Derenne, S., Evershed,
1639 R.P., Kögel-Knabner, I., de Leeuw, J.W., Littke, R., Michaelis, W. and Rullkötter, J.,
1640 2000. The molecularly-uncharacterized component of nonliving organic matter in
1641 natural environments. *Organic Geochemistry*, 31(10): 945-958.

1642 Hefting, M.M., van den Heuvel, R.N. and Verhoeven, J.T.A., 2013. Wetlands in agricultural
1643 landscapes for nitrogen attenuation and biodiversity enhancement: Opportunities and
1644 limitations. *Ecological Engineering*, 56: 5-13.

1645 Helgeson, H.C., 1971. Kinetics of mass transfer among silicates and aqueous solutions.
1646 *Geochimica et Cosmochimica Acta*, 35(5): 421-469.

1647 Helgeson, R.C., 1968. Evaluation of irreversible reactions in geochemical processes
1648 involving minerals and aqueous solutions: I. Thermodynamic relations. *Geochim.*
1649 *Cosmochim. Acta*, 32: 853–877.

1650 Helgeson, R.C., R.M. Garrels and MacKenzie, F.T., 1969. Evaluation of irreversible
1651 reactions in geochemical processes involving minerals and aqueous solutions: II.
1652 Applications,. *Geochim. Cosmochim. Acta*, 33: 455– 482.

1653 Hester, E.T. and Gooseff, M.N., 2010. Moving Beyond the Banks: Hyporheic Restoration Is
1654 Fundamental to Restoring Ecological Services and Functions of Streams.
1655 *Environmental Science & Technology*, 44(5): 1521-1525.

1656 Hinsinger, P., Brauman, A., Devau, N., Gérard, F., Jourdan, C., Laclau, J.-P., Le Cadre, E.,
1657 Jaillard, B. and Plassard, C., 2011. Acquisition of phosphorus and other poorly mobile

1658 nutrients by roots. Where do plant nutrition models fail? *Plant and Soil*, 348(1-2): 29-
1659 61.

1660 Hinsinger, P., Plassard, C., Tang, C. and Jaillard, B., 2003. Origins of root-mediated pH
1661 changes in the rhizosphere and their responses to environmental constraints: A
1662 review. *Plant and Soil*, 248(1-2): 43-59.

1663 Hood, R.R., Wang, H.V., Purcell, J.E., Houde, E.D. and Harding, L.W., 1999. Modeling
1664 particles and pelagic organisms in Chesapeake Bay: Convergent features control
1665 plankton distributions. *Journal of Geophysical Research: Oceans*, 104(C1): 1223-
1666 1243.

1667 Hubbard, C.G., West, L.J., Rodriguez-Blanco, J.D. and Shaw, S., 2014. Laboratory study of
1668 spectral induced polarization responses of magnetite - Fe²⁺ redox reactions in porous
1669 media. *Geophysics*, 79(1): D21-D30.

1670 Huber, C., Parmigiani, A., Latt, J. and Dufek, J., 2013. Channelization of buoyant nonwetting
1671 fluids in saturated porous media. *Water Resources Research*, 49(10): 6371-6380.

1672 Hunkeler, D., Van Breukelen, B.M. and Elsner, M., 2009. Modeling Chlorine Isotope Trends
1673 during Sequential Transformation of Chlorinated Ethenes. *Environmental Science &
1674 Technology*, 43(17): 6750-6756.

1675 Huntington, T.G., 2000. The potential for calcium depletion in forest ecosystems of
1676 southeastern United States: Review and analysis. *Global Biogeochemical Cycles*,
1677 14(2): 623-638.

1678 Jacobson, A.D. and Holmden, C., 2008. delta Ca-44 evolution in a carbonate aquifer and its
1679 bearing on the equilibrium isotope fractionation factor for calcite. *Earth and Planetary
1680 Science Letters*, 270(3-4): 349-353.

1681 James, L.D., 1972. Hydrologic modeling, parameter estimation, and watershed
1682 characteristics. *Journal of Hydrology*, 17(4): 283-307.

1683 Jamieson-Hanes, J.H., Amos, R.T. and Blowes, D.W., 2012. Reactive Transport Modeling of
1684 Chromium Isotope Fractionation during Cr(VI) Reduction. *Environmental Science &
1685 Technology*, 46(24): 13311-13316.

1686 Janot, N., Lezama Pacheco, J.S., Pham, D.Q., O'Brien, T.M., Hausladen, D., Noël, V.,
1687 Lallier, F., Maher, K., Fendorf, S., Williams, K.H., Long, P.E. and Bargar, J.R., 2016.
1688 Physico-Chemical Heterogeneity of Organic-Rich Sediments in the Rifle Aquifer,
1689 CO: Impact on Uranium Biogeochemistry. *Environmental Science & Technology*,
1690 50(1): 46-53.

1691 Jarboe, J.E. and Haan, C., 1974. Calibrating a water yield model for small ungaged
1692 watersheds. *Water Resources Research*, 10(2): 256-262.

1693 Jin, B. and Rolle, M., 2014. Mechanistic approach to multi-element isotope modeling of
1694 organic contaminant degradation. *Chemosphere*, 95: 131-139.

1695 Jin, L., Andrews, D.M., Holmes, G.H., Lin, H. and Brantley, S.L., 2011. Opening the "Black
1696 Box": Water Chemistry Reveals Hydrological Controls on Weathering in the
1697 Susquehanna Shale Hills Critical Zone Observatory. *Vadose Zone Journal*, 10(3):
1698 928-942.

1699 Jin, L., Ravella, R., Ketchum, B., Bierman, P.R., Heaney, P., White, T. and Brantley, S.L.,
1700 2010. Mineral weathering and elemental transport during hillslope evolution at the
1701 Susquehanna/Shale Hills Critical Zone Observatory. *Geochimica Et Cosmochimica
1702 Acta*, 74(13): 3669-3691.

1703 Jin, Q., Roden, E.E. and Giska, J.R., 2012. Geomicrobial Kinetics: Extrapolating Laboratory
1704 Studies to Natural Environments. *Geomicrobiology Journal*, 30(2): 173-185.

1705 Johnson, C.M., Beard, B.L. and Albarede, F., 2004. Overview and general concepts,
1706 Geochemistry of Non-Traditional Stable Isotopes. *Reviews in Mineralogy &
1707 Geochemistry*, pp. 1-24.

Johnson, T.M. and DePaolo, D.J., 1994. Interpretation of Isotopic Data in Groundwater-Rock Systems - Model Development and Application to Sr Isotope Data from Yucca Mountain. *Water Resources Research*, 30(5): 1571-1587.

Johnson, T.M. and DePaolo, D.J., 1997. Rapid exchange effects on isotope ratios in groundwater systems .2. Flow investigation using Sr isotope ratios. *Water Resources Research*, 33(1): 197-209.

Jose, S.C. and Cirpka, O.A., 2004. Measurement of mixing-controlled reactive transport in homogeneous porous media and its prediction from conservative tracer test data. *Environmental Science & Technology*, 38(7): 2089-2096.

Jung, J.K. and McCouch, S., 2013. Getting to the roots of it: genetic and hormonal control of root architecture. *Frontiers in plant science*, 4.

Jung, M., Reichstein, M., Ciais, P., Seneviratne, S.I., Sheffield, J., Goulden, M.L., Bonan, G., Cescatti, A., Chen, J., de Jeu, R., Dolman, A.J., Eugster, W., Gerten, D., Gianelle, D., Gobron, N., Heinke, J., Kimball, J., Law, B.E., Montagnani, L., Mu, Q., Mueller, B., Oleson, K., Papale, D., Richardson, A.D., Rouspard, O., Running, S., Tomelleri, E., Viovy, N., Weber, U., Williams, C., Wood, E., Zaehle, S. and Zhang, K., 2010. Recent decline in the global land evapotranspiration trend due to limited moisture supply. *Nature*, 467(7318): 951-954.

Kang, Q.J., Lichtner, P.C. and Zhang, D.X., 2006. Lattice Boltzmann pore-scale model for multicomponent reactive transport in porous media. *Journal of Geophysical Research-Solid Earth*, 111.

Kiczka, M., Wiederhold, J.G., Frommer, J., Kraemer, S.M., Bourdon, B. and Kretzschmar, R., 2010. Iron isotope fractionation during proton- and ligand-promoted dissolution of primary phyllosilicates. *Geochimica Et Cosmochimica Acta*, 74(11): 3112-3128.

Kim, H., Bishop, J.K.B., Dietrich, W.E. and Fung, I.Y., 2014. Process dominance shift in solute chemistry as revealed by long-term high-frequency water chemistry observations of groundwater flowing through weathered argillite underlying a steep forested hillslope. *Geochimica et Cosmochimica Acta*, 140: 1-19.

King, E., Tuncay, K., Ortoleva, P. and Meile, C., 2009. In silico *Geobacter sulfurreducens* metabolism and its representation in reactive transport models. *Applied and environmental microbiology*, 75(1): 83-92.

King, E.L., Tuncay, K., Ortoleva, P. and Meile, C., 2010. Modeling biogeochemical dynamics in porous media: Practical considerations of pore scale variability, reaction networks, and microbial population dynamics in a sandy aquifer. *Journal of Contaminant Hydrology*, 112(1-4): 130-140.

Kinzelbach, W., Schafer, W. and Herzer, J., 1991. Numerical Modeling of Natural and Enhanced Denitrification Processes in Aquifers. *Water Resources Research*, 27(6): 1123-1135.

Kirchner, J.W., 2003. A double paradox in catchment hydrology and geochemistry. *Hydrological Processes*, 17(4): 871-874.

Kleber, M., Eusterhues, K., Keiluweit, M., Mikutta, C., Mikutta, R. and Nico, P.S., 2015. Chapter One - Mineral–Organic Associations: Formation, Properties, and Relevance in Soil Environments. In: L.S. Donald (Editor), *Advances in Agronomy*. Academic Press, pp. 1-140.

Knutson, C.E., Werth, C.J. and Valocchi, A.J., 2005. Pore-scale simulation of biomass growth along the transverse mixing zone of a model two-dimensional porous medium. *Water Resources Research*, 41(7).

Kollet, S.J. and Maxwell, R.M., 2006. Integrated surface-groundwater flow modeling: A free-surface overland flow boundary condition in a parallel groundwater flow model. *Advances in Water Resources*, 29(7): 945-958.

1758 Krumins, V., Gehlen, M., Arndt, S., Van Cappellen, P. and Regnier, P., 2013. Dissolved
1759 inorganic carbon and alkalinity fluxes from coastal marine sediments: model
1760 estimates for different shelf environments and sensitivity to global change.
1761 *Biogeosciences*, 10(1): 371-398.

1762 Kumar, M., Duffy, C.J. and Salvage, K.M., 2009. A Second-Order Accurate, Finite Volume–
1763 Based, Integrated Hydrologic Modeling (FIHM) Framework for Simulation of
1764 Surface and Subsurface Flow All rights reserved. No part of this periodical may be
1765 reproduced or transmitted in any form or by any means, electronic or mechanical,
1766 including photocopying, recording, or any information storage and retrieval system,
1767 without permission in writing from the publisher. *Vadose Zone J.*, 8(4): 873-890.

1768 Kump, L.R., Brantley, S.L. and Arthur, M.A., 2000. Chemical weathering, atmospheric CO₂,
1769 and climate. *Annual Review of Earth and Planet Science*, 28: 611 - 667.

1770 Kuzyakov, Y. and Blagodatskaya, E., 2015. Microbial hotspots and hot moments in soil:
1771 Concept & review. *Soil Biology and Biochemistry*, 83: 184-199.

1772 Lalonde, K., Mucci, A., Ouellet, A. and Gelin, Y., 2012. Preservation of organic matter in
1773 sediments promoted by iron. *Nature*, 483(7388): 198-200.

1774 Landeweert, R., Hoffland, E., Finlay, R.D., Kuiper, T.W. and van Breemen, N., 2001.
1775 Linking plants to rocks: ectomycorrhizal fungi mobilize nutrients from minerals.
1776 *Trends in Ecology & Evolution*, 16(5): 248-254.

1777 Lawrence, C., Harden, J. and Maher, K., 2014a. Modeling the influence of organic acids on
1778 soil weathering. *Geochimica et Cosmochimica Acta*, 139(0): 487-507.

1779 Lawrence, C., Steefel, C. and Maher, K., 2014b. Abiotic/Biotic Coupling in the Rhizosphere:
1780 A Reactive Transport Modeling Analysis. *Procedia Earth and Planetary Science*, 10:
1781 104-108.

1782 Lawrence, C.R., Harden, J.W., Xu, X.M., Schulz, M.S. and Trumbore, S.E., 2015. Long-term
1783 controls on soil organic carbon with depth and time: A case study from the Cowlitz
1784 River Chronosequence, WA USA. *Geoderma*, 247: 73-87.

1785 Lawrence, C.R., Neff, J.C. and Schimel, J.P., 2009. Does adding microbial mechanisms of
1786 decomposition improve soil organic matter models? A comparison of four models
1787 using data from a pulsed rewetting experiment. *Soil Biology and Biochemistry*, 41(9):
1788 1923-1934.

1789 Lebedeva, M.I. and Brantley, S.L., 2013. Exploring geochemical controls on weathering and
1790 erosion of convex hillslopes: beyond the empirical regolith production function. *Earth
1791 Surface Processes and Landforms*, 38(15): 1793-1807.

1792 LeBlanc, D.R., Garabedian, S.P., Hess, K.M., Gelhar, L.W., Quadri, R.D., Stollenwerk, K.G.
1793 and Wood, W.W., 1991. Large-scale natural gradient tracer test in sand and gravel,
1794 Cape Cod, Massachusetts: 1. Experimental design and observed tracer movement.
1795 *Water Resources Research*, 27(5): 895-910.

1796 Lehmann, J., Kleber, M., 2015. The contentious nature of soil organic matter. *Nature*,
1797 528(7580): 60-68.

1798 Lehmann, M.F., Sigman, D.M., McCorkle, D.C., Granger, J., Hoffmann, S., Cane, G. and
1799 Brunelle, B.G., 2007. The distribution of nitrate ¹⁵N/¹⁴N in marine sediments and
1800 the impact of benthic nitrogen loss on the isotopic composition of oceanic nitrate.
1801 *Geochimica et Cosmochimica Acta*, 71(22): 5384-5404.

1802 Lemarchand, E., Chabaux, F., Vigier, N., Millot, R. and Pierret, M.-C., 2010. Lithium isotope
1803 systematics in a forested granitic catchment (Strengbach, Vosges Mountains, France).
1804 *Geochimica et Cosmochimica Acta*, 74(16): 4612-4628.

1805 Li, L., Gawande, N., Kowalsky, M.B., Steefel, C.I. and Hubbard, S.S., 2011.
1806 Physicochemical heterogeneity controls on uranium bioreduction rates at the field
1807 scale. *Environmental Science and Technology*, 45(23): 9959-66.

1808 Li, L., Maher, K.M. and Navarre-Sitchler, A., 2014. Expanding the Role of Reactive
 1809 Transport Modeling in Biogeochemical Sciences. *Eos, Transactions American*
 1810 *Geophysical Union*, 95(35): 316-316.
 1811 Li, L., Peters, C.A. and Celia, M.A., 2006. Upscaling geochemical reaction rates using pore-
 1812 scale network modeling. *Advances in Water Resources* 29(9): 1351-1370.
 1813 Li, L., Peters, C.A. and Celia, M.A., 2007. Effects of mineral spatial distribution on reaction
 1814 rates in porous media. *Water Resources Research*, 43(1):
 1815 doi:10.1029/2005WR004848.
 1816 Li, L., Steefel, C., Kowalsky, M., Englert, A. and Hubbard, S., 2010. Effects of physical and
 1817 geochemical heterogeneities on mineral transformation and biomass accumulation
 1818 during a biostimulation experiment at Rifle, Colorado. *J Contamin Hydrol*, 112: 45 -
 1819 63.
 1820 Li, L., Steefel, C.I., Williams, K.H., Wilkins, M.J. and Hubbard, S.S., 2009. Mineral
 1821 transformation and biomass accumulation during uranium bioremediation at Rifle,
 1822 Colorado. *Environ. Sci. Technol.*, 43(14): 5429-5435. doi: 10.1021/es900016v.
 1823 Li, L., Steefel, C.I. and Yang, L., 2008. Scale dependence of mineral dissolution rates within
 1824 single pores and fractures. *Geochimica Et Cosmochimica Acta*, 72(2): 360-377.
 1825 Lichtner, P.C., 1985. Continuum Model for Simultaneous Chemical-Reactions and Mass-
 1826 Transport in Hydrothermal Systems. *Geochimica et Cosmochimica Acta*, 49(3): 779-
 1827 800.
 1828 Lichtner, P.C., 1988. The Quasi-Stationary State Approximation to Coupled Mass- Transport
 1829 and Fluid-Rock Interaction in a Porous-Medium. *Geochim. Cosmochim. Acta.*, 52(1):
 1830 143-165.
 1831 Lifton, N.A. and Chase, C.G., 1992. Fractals in Geomorphology Tectonic, climatic and
 1832 lithologic influences on landscape fractal dimension and hypsometry: implications for
 1833 landscape evolution in the San Gabriel Mountains, California. *Geomorphology*, 5(1):
 1834 77-114.
 1835 Liu, C., Shang, J., Shan, H. and Zachara, J.M., 2013. Effect of Subgrid Heterogeneity on
 1836 Scaling Geochemical and Biogeochemical Reactions: A Case of U(VI) Desorption.
 1837 *Environmental Science & Technology*, 48(3): 1745-1752.
 1838 Liu, C., Zachara, J.M., Qafoku, N.P. and Wang, Z., 2008. Scale-dependent desorption of
 1839 uranium from contaminated subsurface sediments. *Water Resources Research*, 44(8).
 1840 Lopez, B., Bashan, Y., Bacilio, M. and De la Cruz-Agüero, G., 2009. Rock-colonizing plants:
 1841 abundance of the endemic cactus *Mammillaria fraileana* related to rock type in the
 1842 southern Sonoran Desert. *Plant Ecology*, 201(2): 575-588.
 1843 Louvat, P., Gayer, E. and Gaillardet, J., 2014. Boron behavior in the rivers of Reunion island,
 1844 inferred from boron isotope ratios and concentrations of major and trace elements. In:
 1845 J. Gaillardet (Editor), *Geochemistry of the Earth's Surface Ges-10. Procedia Earth and*
 1846 *Planetary Science*, pp. 231-237.
 1847 Lucas, R.W., Klaminder, J., Futter, M.N., Bishop, K.H., Egnell, G., Laudon, H. and Hogberg,
 1848 P., 2011. A meta-analysis of the effects of nitrogen additions on base cations:
 1849 Implications for plants, soils, and streams. *Forest Ecology and Management*, 262(2):
 1850 95-104.
 1851 Luce, R.W., Bartlett, R.W., Parks, G.A., 1972. Dissolution kinetics of magnesium silicates.
 1852 *Geochimica et Cosmochimica Acta*, 36: 35-50.
 1853 Luo, J. and Cirpka, O.A., 2008. Traveltime-based descriptions of transport and mixing in
 1854 heterogeneous domains. *Water Resources Research*, 44(9).
 1855 Lynch, J., Nielsen, K., Davis, R. and Jabllokow, A., 1997. SimRoot: Modelling and
 1856 visualization of root systems. *Plant and Soil*, 188(1): 139-151.
 1857 Lynch, J.P., 1995. Root architecture and plant productivity. *Plant physiology*, 109(1): 7.

1858 MacQuarrie, K.T.B. and Mayer, K.U., 2005. Reactive transport modeling in fractured rock: A
1859 state-of-the-science review. *Earth-Science Reviews*, 72(3-4): 189-227.

1860 Maggi, F. and Porporato, A., 2007. Coupled moisture and microbial dynamics in unsaturated
1861 soils. *Water Resources Research*, 43(7): W07444.

1862 Maher, K., 2010. The dependence of chemical weathering rates on fluid residence time. *Earth
1863 and Planetary Science Letters*, 294(1-2): 101-110.

1864 Maher, K., 2011. The role of fluid residence time and topographic scales in determining
1865 chemical fluxes from landscapes. *Earth and Planetary Science Letters*, 312(1-2): 48-
1866 58.

1867 Maher, K. and Chamberlain, C.P., 2014. Hydrologic Regulation of Chemical Weathering and
1868 the Geologic Carbon Cycle. *Science*, 343(6178): 1502-1504.

1869 Maher, K., DePaolo, D.J. and Christensen, J.N., 2006a. U-Sr isotopic speedometer: flow and
1870 chemical weathering in aquifers. *Geochimica Et Cosmochimica Acta*, 70(17): 4417-
1871 4435.

1872 Maher, K., Johnson, N.C., Jackson, A., Lammers, L.N., Torchinsky, A.B., Weaver, K.L.,
1873 Bird, D.K. and Brown Jr, G.E., 2016. A spatially resolved surface kinetic model for
1874 forsterite dissolution. *Geochimica et Cosmochimica Acta*, 174: 313-334.

1875 Maher, K., Steefel, C.I., DePaolo, D.J. and Viani, B.E., 2006b. The mineral dissolution rate
1876 conundrum: Insights from reactive transport modeling of U isotopes and pore fluid
1877 chemistry in marine sediments. *Geochimica et Cosmochimica Acta*, 70(2): 337-363.

1878 Maher, K., Steefel, C.I., White, A.F. and Stonestrom, D.A., 2009a. The role of reaction
1879 affinity and secondary minerals in regulating chemical weathering rates at the Santa
1880 Cruz Soil Chronosequence, California. *Geochimica et Cosmochimica Acta*, 73: 2804-
1881 2831.

1882 Maher, K., Steefel, C.I., White, A.F. and Stonestrom, D.A., 2009b. The role of reaction
1883 affinity and secondary minerals in regulating chemical weathering rates at the Santa
1884 Cruz Soil Chronosequence, California. *Geochimica Et Cosmochimica Acta*, 73(10):
1885 2804-2831.

1886 Malmstrom, M.E., Destouni, G., Banwart, S.A. and Stromberg, B.H.E., 2000. Resolving the
1887 Scale-Dependence of Mineral Weathering Rates. *Environmental Science &
1888 Technology*, 34(7): 1375-1378.

1889 Malmstrom, M.E., Destouni, G. and Martinet, P., 2004. Modeling expected solute
1890 concentration in randomly heterogeneous flow systems with multicomponent
1891 reactions. *Environ. Sci. Technol.*, 38: 2673-2679.

1892 Manzoni, S. and Porporato, A., 2009. Soil carbon and nitrogen mineralization: Theory and
1893 models across scales. *Soil Biology and Biochemistry*, 41(7): 1355-1379.

1894 Manzoni, S., Porporato, A. and Schimel, J.P., 2008. Soil heterogeneity in lumped
1895 mineralization-immobilization models. *Soil Biology and Biochemistry*, 40(5): 1137-
1896 1148.

1897 Manzoni, S., Schaeffer, S.M., Katul, G., Porporato, A. and Schimel, J.P., 2014. A theoretical
1898 analysis of microbial eco-physiological and diffusion limitations to carbon cycling in
1899 drying soils. *Soil Biology and Biochemistry*, 73: 69-83.

1900 Marin-Spiotta, E., Silver, W.L., Swanston, C.W. and Ostertag, R., 2009. Soil organic matter
1901 dynamics during 80 years of reforestation of tropical pastures. *Global Change
1902 Biology*, 15(6): 1584-1597.

1903 Mavromatis, V., Bundeleva, I.A., Shirokova, L.S., Millo, C., Pokrovsky, O.S., Benezeth, P.,
1904 Ader, M. and Oelkers, E.H., 2015. The continuous re-equilibration of carbon isotope
1905 compositions of hydrous Mg carbonates in the presence of cyanobacteria. *Chemical
1906 Geology*, 404: 41-51.

1907 Mavromatis, V., Meister, P. and Oelkers, E.H., 2014. Using stable Mg isotopes to distinguish
1908 dolomite formation mechanisms: A case study from the Peru Margin. *Chemical*
1909 *Geology*, 385: 84-91.

1910 Maxwell, R.M., 2013. A terrain-following grid transform and preconditioner for parallel,
1911 large-scale, integrated hydrologic modeling. *Advances in Water Resources*, 53: 109-
1912 117.

1913 Maxwell, R.M. and Miller, N.L., 2005. Development of a coupled land surface and
1914 groundwater model. *Journal of Hydrometeorology*, 6(3): 233-247.

1915 Mayer, K.U., Benner, S.G., Frind, E.O., Thornton, S.F. and Lerner, D.N., 2001. Reactive
1916 transport modeling of processes controlling the distribution and natural attenuation of
1917 phenolic compounds in a deep sandstone aquifer. *Journal of Contaminant Hydrology*,
1918 53(3-4): 341-368.

1919 McClain, M.E., Boyer, E.W., Dent, C.L., Gergel, S.E., Grimm, N.B., Groffman, P.M., Hart,
1920 S.C., Harvey, J.W., Johnston, C.A., Mayorga, E., McDowell, W.H. and Pinay, G.,
1921 2003. Biogeochemical hot spots and hot moments at the interface of terrestrial and
1922 aquatic ecosystems. *Ecosystems*, 6(4): 301-312.

1923 McDonnell, J., Sivapalan, M., Vaché, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C.,
1924 Hooper, R., Kirchner, J. and Roderick, M., 2007a. Moving beyond heterogeneity and
1925 process complexity: A new vision for watershed hydrology. *Water Resources*
1926 *Research*, 43(7).

1927 McDonnell, J.J., Sivapalan, M., Vache, K., Dunn, S., Grant, G., Haggerty, R., Hinz, C.,
1928 Hooper, R., Kirchner, J., Roderick, M.L., Selker, J. and Weiler, M., 2007b. Moving
1929 beyond heterogeneity and process complexity: A new vision for watershed hydrology.
1930 *Water Resources Research*, 43(7).

1931 Meile, C. and Tuncay, K., 2006. Scale dependence of reaction rates in porous media.
1932 *Advances in Water Resources*, 29(1): 62-71.

1933 Mikutta, R., Kleber, M., Torn, M.S. and Jahn, R., 2006. Stabilization of soil organic matter:
1934 Association with minerals or chemical recalcitrance? *Biogeochemistry*, 77(1): 25-56.

1935 Mikutta, R., Schaumann, G.E., Gildemeister, D., Bonneville, S., Kramer, M.G., Chorover, J.,
1936 Chadwick, O.A. and Guggenberger, G., 2009. Biogeochemistry of mineral-organic
1937 associations across a long-term mineralogical soil gradient (0.3-4100 kyr), Hawaiian
1938 Islands. *Geochimica et Cosmochimica Acta*, 73(7): 2034-2060.

1939 Miller, A.W., Rodriguez, D.R. and Honeyman, B.D., 2013. Simplified behaviors from
1940 increased heterogeneity: II. 3-D Uranium transport at the decimeter scale and
1941 intertank comparisons. *Journal of Contaminant Hydrology*, 148: 51-66.

1942 Miller, G.R., Rubin, Y., Mayer, K.U. and Benito, P.H., 2008. Modeling Vadose Zone
1943 Processes during Land Application of Food-Processing Waste Water in California's
1944 Central Valley. *Journal of Environmental Quality*, 37(5_Supplement): S-43-S-57.

1945 Minasny, B. and McBratney, A.B., 1999. A rudimentary mechanistic model for soil
1946 production and landscape development. *Geoderma*, 90(1-2): 3-21.

1947 Minasny, B. and McBratney, A.B., 2001. A rudimentary mechanistic model for soil
1948 formation and landscape development: II. A two-dimensional model incorporating
1949 chemical weathering. *Geoderma*, 103(1-2): 161-179.

1950 Minasny, B. and McBratney, A.B., 2006. Mechanistic soil-landscape modelling as an
1951 approach to developing pedogenetic classifications. *Geoderma*, 133(1-2): 138-149.

1952 Molins, S. and Mayer, K.U., 2007. Coupling between geochemical reactions and
1953 multicomponent gas and solute transport in unsaturated media: A reactive transport
1954 modeling study. *Water Resources Research*, 43(5): n/a-n/a.

1955 Molins, S., Trebotich, D., Steefel, C.I. and Shen, C., 2012. An investigation of the effect of
1956 pore scale flow on average geochemical reaction rates using direct numerical
1957 simulation. *Water Resources Research*, 48.

1958 Molins, S., Trebotich, D., Yang, L., Ajo-Franklin, J.B., Ligocki, T.J., Shen, C. and Steefel,
1959 C.I., 2014. Pore-scale controls on calcite dissolution rates from flow-through
1960 laboratory and numerical experiments. *Environmental science & technology*, 48(13):
1961 7453-7460.

1962 Molz, F.J. and Widdowson, M.A., 1988. Internal inconsistencies in dispersion-dominated
1963 models that incorporate chemical and microbial kinetics. *Water Resources Research*,
1964 24(4): 615-619.

1965 Molz, F.J., Widdowson, M.A. and Benefield, L.D., 1986. Simulation of microbial-growth
1966 dynamics coupled to nutrient and oxygen-transport in porous media. *Water Resources*
1967 *Research*, 22(8): 1207-1216.

1968 Moore, J., Lichtner, P.C., White, A.F. and Brantley, S.L., 2012. Using a reactive transport
1969 model to elucidate differences between laboratory and field dissolution rates in
1970 regolith. *Geochimica Et Cosmochimica Acta*, 93: 235-261.

1971 Moulton, K.L., West, J. and Berner, R.A., 2000. Solute flux and mineral mass balance
1972 approaches to the quantification of plant effects on silicate weathering. *American*
1973 *Journal of Science*, 300(7): 539-570.

1974 National Research Council, A.B., 2010. *Landscapes on the Edge: New Horizons for Research*
1975 *on Earth's Surface*. The National Academies Press, Washington, DC, 180 pp.

1976 Navarre-Sitchler, A. and Brantley, S., 2007. Basalt weathering across scales. *Earth and*
1977 *Planetary Science Letters*, 261: 321-334.

1978 Navarre-Sitchler, A., Steefel, C.I., Sak, P.B. and Brantley, S.L., 2011. A reactive-transport
1979 model for weathering rind formation on basalt. *Geochimica Et Cosmochimica Acta*,
1980 75(23): 7644-7667.

1981 Navarre-Sitchler, A., Steefel, C.I., Yang, L., Tomutsa, L. and Brantley, S.L., 2009. Evolution
1982 of porosity and diffusivity associated with chemical weathering of a basalt clast.
1983 *Journal of Geophysical Research-Earth Surface*, 114.

1984 Navarre-Sitchler, A.K., Maxwell, R.M., Siirila, E.R., Hammond, G.E. and Lichtner, P.C.,
1985 2013. Elucidating geochemical response of shallow heterogeneous aquifers to CO₂
1986 leakage using high-performance computing: Implications for monitoring of CO₂
1987 sequestration. *Advances in Water Resources*, 53(0): 45-55.

1988 Navon, I., 2009. Data assimilation for numerical weather prediction: a review. In: S. Park and
1989 X. L (Editors), *Data assimilation for atmospheric, oceanic, hydrologic applications*.
1990 Springer, Berlin.

1991 Neogi, S., Bolton, E. and Chakraborty, S., 2014. Timescales of disequilibrium melting in the
1992 crust: constraints from modelling the distribution of multiple trace elements and a
1993 case study from the Lesser Himalayan rocks of Sikkim. *Contributions to Mineralogy*
1994 *and Petrology*, 168(2): 1-22.

1995 Neuman, S.P. and Tartakovsky, D.M., 2009. Perspective on theories of non-Fickian transport
1996 in heterogeneous media. *Advances in Water Resources*, 32(5): 670-680.

1997 Neuman, S.P., Xue, L., Ye, M. and Lu, D., 2012. Bayesian analysis of data-worth considering
1998 model and parameter uncertainties. *Advances in Water Resources*, 36(0): 75-85.

1999 Neumann, R.B. and Cardon, Z.G., 2012. The magnitude of hydraulic redistribution by plant
2000 roots: a review and synthesis of empirical and modeling studies. *New Phytologist*,
2001 194(2): 337-352.

2002 Nielsen, L.C., DePaolo, D.J. and De Yoreo, J.J., 2012. Self-consistent ion-by-ion growth
2003 model for kinetic isotopic fractionation during calcite precipitation. *Geochimica Et*
2004 *Cosmochimica Acta*, 86: 166-181.

2005 Noireaux, J., Gaillardet, J., Sullivan, P.L. and Brantley, S.L., 2014. Boron isotope
2006 fractionation in soils at Shale Hills CZO. In: J. Gaillardet (Editor), *Geochemistry of*
2007 *the Earth's Surface Ges-10. Procedia Earth and Planetary Science*, pp. 218-222.
2008 NRC, 2012. *Challenges and opportunities in the hydrologic sciences*. The National
2009 Academies Press, Washington, D. C. .
2010 Nugent, M.A., Brantley, S.L., Pantano, C.G. and Maurice, P.A., 1998. The influence of
2011 natural mineral coatings on feldspar weathering. *Nature*, 395(6702): 588-591.
2012 O'Brien, B.J. and Stout, J.D., 1978. Movement and turnover of soil organic matter as
2013 indicated by carbon isotope measurements. *Soil Biology and Biochemistry*, 10(4):
2014 309-317.
2015 Oelkers, E.H., Schott, J. and Devidal, J.-L., 1994. The effect of aluminum, pH, and chemical
2016 affinity on the rates of aluminosilicate dissolution reactions. *Geochimica et*
2017 *Cosmochimica Acta*, 58(9): 2011-2024.
2018 Oelze, M., von Blanckenburg, F., Bouchez, J., Hoellen, D. and Dietzel, M., 2015. The effect
2019 of Al on Si isotope fractionation investigated by silica precipitation experiments.
2020 *Chemical Geology*, 397: 94-105.
2021 Oelze, M., von Blanckenburg, F., Hoellen, D., Dietzel, M. and Bouchez, J., 2014. Si stable
2022 isotope fractionation during adsorption and the competition between kinetic and
2023 equilibrium isotope fractionation: Implications for weathering systems. *Chemical*
2024 *Geology*, 380: 161-171.
2025 Oh, N.-H., Hofmockel, M., Lavine, M.L. and Richter, D.D., 2007. Did elevated atmospheric
2026 CO₂ alter soil mineral weathering?: an analysis of 5-year soil water chemistry data at
2027 Duke FACE study. *Global Change Biology*, 13(12): 2626-2641.
2028 Ortoleva, P., Merino, E., Moore, C. and Chadam, J. , 1987. Geochemical self organization, I.
2029 Feedbacks, quantitative modeling. *Amer. J. Science*, 287: 979–1007.
2030 Ostle, N.J., Smith, P., Fisher, R., Woodward, F.I., Fisher, J.B., Smith, J.U., Galbraith, D.,
2031 Levy, P., Meir, P., McNamara, N.P. and Bardgett, R.D., 2009. Integrating plant–soil
2032 interactions into global carbon cycle models. *Journal of Ecology*, 97(5): 851-863.
2033 Pacheco-Villalobos, D. and Hardtke, C.S., 2012. Natural genetic variation of root system
2034 architecture from *Arabidopsis* to *Brachypodium*: towards adaptive value.
2035 *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1595):
2036 1552-1558.
2037 Parsekian, A.D., Singha, K., Minsley, B.J., Holbrook, W.S. and Slater, L., 2015. Multiscale
2038 geophysical imaging of the critical zone. *Reviews of Geophysics*, 53(1): 1-26.
2039 Parton, W.J., Schimel, D.S., Cole, C.V. and Ojima, D.S., 1987. Analysis of Factors
2040 Controlling Soil Organic Matter Levels in Great Plains Grasslands. *Soil Science*
2041 *Society of America Journal*, 51(5): 1173-1179.
2042 Pedretti, D., Fernández-García, D., Bolster, D. and Sanchez-Vila, X., 2013. On the formation
2043 of breakthrough curves tailing during convergent flow tracer tests in three-
2044 dimensional heterogeneous aquifers. *Water Resources Research*, 49(7): 4157-4173.
2045 Pelletier, J.D. and Rasmussen, C., 2009. Quantifying the climatic and tectonic controls on
2046 hillslope steepness and erosion rate. *Lithosphere*, 1(2): 73-80.
2047 Perdrial, J., Thompson, A. and Chorover, J., 2015. Chapter 6 - Soil Geochemistry in the
2048 Critical Zone: Influence on Atmosphere, Surface- and Groundwater Composition. In:
2049 R.G. John and H. Chris (Editors), *Developments in Earth Surface Processes*. Elsevier,
2050 pp. 173-201.
2051 Perdrial, J.N., McIntosh, J.C., Harpold, A., Brooks, P.D., Zapata-Rios, X., Ray, J., Meixner,
2052 T., Kanduc, T., Litvak, M., Troch, P. and Chorover, J., 2014. Stream water carbon
2053 controls in seasonally snow-covered mountain catchments: impact of inter annual

2054 variability of water fluxes, catchment aspect and seasonal processes.
 2055 Biogeochemistry, 118(1-3): 273-290.
 2056 Perrin, A.S., Probst, A. and Probst, J.L., 2008. Impact of nitrogenous fertilizers on carbonate
 2057 dissolution in small agricultural catchments : implications for weathering CO₂, uptake
 2058 at regional and global scales. *Geochimica et Cosmochimica Acta*, 72: 3105-3123.
 2059 Philippot, L., Bru, D., Saby, N.P.A., Čuhel, J., Arrouays, D., Šimek, M. and Hallin, S., 2009.
 2060 Spatial patterns of bacterial taxa in nature reflect ecological traits of deep branches of
 2061 the 16S rRNA bacterial tree. *Environmental Microbiology*, 11(12): 3096-3104.
 2062 Porporato, A., D'Odorico, P., Laio, F. and Rodriguez-Iturbe, I., 2003. Hydrologic controls on
 2063 soil carbon and nitrogen cycles. I. Modeling scheme. *Advances in Water Resources*,
 2064 26(1): 45-58.
 2065 Prieto, I., Armas, C. and Pugnaire, F.I., 2012. Water release through plant roots: new insights
 2066 into its consequences at the plant and ecosystem level. *New Phytologist*, 193(4): 830-
 2067 841.
 2068 Prommer, H., Tuxen, N. and Bjerg, P.L., 2006. Fringe-controlled natural attenuation of
 2069 phenoxy acids in a landfill plume: Integration of field-scale processes by reactive
 2070 transport modeling. *Environmental Science & Technology*, 40(15): 4732-4738.
 2071 Qu, Y. and Duffy, C.J., 2007a. A semidiscrete finite volume formulation for multiprocess
 2072 watershed simulation. *Water Resources Research*, 43(8).
 2073 Qu, Y. and Duffy, C.J., 2007b. A semidiscrete finite volume formulation for multiprocess
 2074 watershed simulation. *Water Resources Research*, 43(8): W08419.
 2075 Quinn, P., Beven, K., Chevallier, P. and Planchon, O., 1991. The prediction of hillslope flow
 2076 paths for distributed hydrological modelling using digital terrain models.
 2077 *Hydrological processes*, 5(1): 59-79.
 2078 Rabier, F., 2005. Overview of global data assimilation developments in numerical weather-
 2079 prediction centres. *Quarterly Journal of the Royal Meteorological Society*, 131(613):
 2080 3215-3233.
 2081 Rasmussen, C., Southard, R.J. and Horwath, W.R., 2007. Soil mineralogy affects conifer
 2082 forest soil carbon source utilization and microbial priming. *Soil Science Society of
 2083 America Journal*, 71(4): 1141-1150.
 2084 Rasmussen, C., Southard, R.J. and Horwath, W.R., 2008. Litter type and soil minerals control
 2085 temperate forest soil carbon response to climate change. *Global Change Biology*,
 2086 14(9): 2064-2080.
 2087 Raymond, P.A., Oh, N.-H., Turner, R.E. and Broussard, W., 2008. Anthropogenically
 2088 enhanced fluxes of water and carbon from the Mississippi River. *Nature*, 451(7177):
 2089 449-452.
 2090 Raynaud, X., Lata, J.-C. and Leadley, P., 2006. Soil microbial loop and nutrient uptake by
 2091 plants: a test using a coupled C:N model of plant-microbial interactions. *Plant and
 2092 Soil*, 287(1-2): 95-116.
 2093 Reed, J.L. and Palsson, B.Ø., 2003. Thirteen Years of Building Constraint-Based In Silico
 2094 Models of *Escherichia coli*. *Journal of Bacteriology*, 185(9): 2692-2699.
 2095 Regnier, P., Arndt, S., Goossens, N., Volta, C., Laruelle, G.G., Lauerwald, R. and Hartmann,
 2096 J., 2013. Modelling Estuarine Biogeochemical Dynamics: From the Local to the
 2097 Global Scale. *Aquatic geochemistry*, 19(5-6): 591-626.
 2098 Regnier, P., Wollast, R. and Steefel, C.I., 1997. Long-term fluxes of reactive species in
 2099 macrotidal estuaries: Estimates from a fully transient, multicomponent reaction-
 2100 transport model. *Marine Chemistry*, 58(1-2): 127-145.
 2101 Rempe, D.M. and Dietrich, W.E., 2014. A bottom-up control on fresh-bedrock topography
 2102 under landscapes. *Proceedings of the National Academy of Sciences*, 111(18): 6576-
 2103 6581.

- Ren, D., Xue, M. and Henderson-Sellers, A., 2004. Incorporating hydraulic lift into a land surface model and its effects on surface soil moisture prediction. *Journal of Hydrometeorology*, 5(6): 1181-1191.
- Renard, F., Ortoleva, P. and Gratier, J.P., 1997. Pressure solution in sandstones: influence of clays and dependence on temperature and stress. *Tectonophysics*, 280(3–4): 257-266.
- Renard, P. and Allard, D., 2013. Connectivity metrics for subsurface flow and transport. *Advances in Water Resources*, 51(0): 168-196.
- Richards, P.L. and Kump, L.R., 2003. Soil pore-water distributions and the temperature feedback of weathering in soils. *Geochimica et cosmochimica acta*, 67(20): 3803-3815.
- Riley, W.J., Maggi, F., Kleber, M., Torn, M.S., Tang, J.Y., Dwivedi, D. and Guerry, N., 2014. Long residence times of rapidly decomposable soil organic matter: application of a multi-phase, multi-component, and vertically resolved model (BAMS1) to soil carbon dynamics. *Geosci. Model Dev.*, 7(4): 1335-1355.
- Rive, K., Gaillardet, J., Agrinier, P. and Rad, S., 2013. Carbon isotopes in the rivers from the Lesser Antilles: origin of the carbonic acid consumed by weathering reactions in the Lesser Antilles. *Earth Surface Processes and Landforms*, 38(9): 1020-1035.
- Roelandt, C., Goddris, Y., Bonnet, M.P. and Sondag, F., 2010. Coupled modeling of biospheric and chemical weathering processes at the continental scale. *Global Biogeochemical Cycles*, 24(2): GB2004.
- Roering, J.J., Marshall, J., Booth, A.M., Mort, M. and Jin, Q., 2010. Evidence for biotic controls on topography and soil production. *Earth and Planetary Science Letters*, 298(1): 183-190.
- Rolle, M., Chiogna, G., Bauer, R., Griebler, C. and Grathwohl, P., 2010. Isotopic Fractionation by Transverse Dispersion: Flow-through Microcosms and Reactive Transport Modeling Study. *Environmental Science & Technology*, 44(16): 6167-6173.
- Rolle, M., Eberhardt, C., Chiogna, G., Cirpka, O.A. and Grathwohl, P., 2009. Enhancement of dilution and transverse reactive mixing in porous media: Experiments and model-based interpretation. *Journal of Contaminant Hydrology*, 110(3-4): 130-142.
- Rolle, M. and Kitanidis, P.K., 2014. Effects of compound-specific dilution on transient transport and solute breakthrough: A pore-scale analysis. *Advances in Water Resources*, 71: 186-199.
- Rosenzweig, C. and Neofotis, P., 2013. Detection and attribution of anthropogenic climate change impacts. *Wiley Interdisciplinary Reviews: Climate Change*, 4(2): 121-150.
- Roy, M., Martin, J.B., Cable, J.E. and Smith, C.G., 2013. Variations of iron flux and organic carbon remineralization in a subterranean estuary caused by inter-annual variations in recharge. *Geochimica et Cosmochimica Acta*, 103(0): 301-315.
- Rumpel, C., Eusterhues, K. and Kogel-Knabner, I., 2004. Location and chemical composition of stabilized organic carbon in topsoil and subsoil horizons of two acid forest soils. *Soil Biology & Biochemistry*, 36(1): 177-190.
- Ryel, R., Caldwell, M., Yoder, C., Or, D. and Leffler, A., 2002. Hydraulic redistribution in a stand of *Artemisia tridentata*: evaluation of benefits to transpiration assessed with a simulation model. *Oecologia*, 130(2): 173-184.
- Salehikhoo, F. and Li, L., 2015. The role of magnesite spatial distribution patterns in determining dissolution rates: When do they matter? *Geochimica et Cosmochimica Acta*, 155(0): 107-121.
- Salehikhoo, F., Li, L. and Brantley, S.L., 2013. Magnesite dissolution rates at different spatial scales: The role of mineral spatial distribution and flow velocity. *Geochimica Et Cosmochimica Acta*, 108(0): 91-106.

2154 Santhi, C., Arnold, J.G., Williams, J.R., Dugas, W.A., Srinivasan, R. and Hauck, L.M., 2001.
2155 VALIDATION OF THE SWAT MODEL ON A LARGE RWER BASIN WITH
2156 POINT AND NONPOINT SOURCES¹. Wiley Online Library.

2157 Sassen, D.S., Hubbard, S.S., Bea, S.A., Chen, J., Spycher, N. and Denham, M.E., 2012.
2158 Reactive facies: An approach for parameterizing field-scale reactive transport models
2159 using geophysical methods. *Water Resources Research*, 48(10): W10526.

2160 Saunders, J.A. and Toran, L.E., 1995. Modeling of radionuclide and heavy metal sorption
2161 around low- and high- $\{pH\}$ waste disposal sites at $\{Oak Ridge, Tennessee\}$. *Appl.*
2162 *Geochem.*, 10(6): 673-684.

2163 Scheibe, T.D., Hubbard, S.S., Onstott, T.C. and DeFlaun, M.F., 2011. Lessons Learned from
2164 Bacterial Transport Research at the South Oyster Site. *Ground Water*, 49(5): 745-763.

2165 Scheibe, T.D., Mahadevan, R., Fang, Y., Garg, S., Long, P.E. and Lovley, D.R., 2009.
2166 Coupling a genome-scale metabolic model with a reactive transport model to describe
2167 in situ uranium bioremediation. *Microbial Biotechnology*, 2(2): 274-286.

2168 Scheibe, T.D., Murphy, E.M., Chen, X., Rice, A.K., Carroll, K.C., Palmer, B.J., Tartakovsky,
2169 A.M., Battiato, I. and Wood, B.D., 2015. An analysis platform for multiscale
2170 hydrogeologic modeling with emphasis on hybrid multiscale methods. *Groundwater*,
2171 53(1): 38-56.

2172 Scheibe, T.D. and Smith, J.C., 2015. Multiscale computation: Needs and opportunities for
2173 BER science.

2174 Schmidt, M.W.I., Torn, M.S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I.A.,
2175 Kleber, M., Kogel-Knabner, I., Lehmann, J., Manning, D.A.C., Nannipieri, P., Rasse,
2176 D.P., Weiner, S. and Trumbore, S.E., 2011. Persistence of soil organic matter as an
2177 ecosystem property. *Nature*, 478(7367): 49-56.

2178 Schnepf, A., Leitner, D. and Klepsch, S., 2012. Modeling Phosphorus Uptake by a Growing
2179 and Exuding Root System. *Vadose Zone Journal*, 11(3).

2180 Schott, J., Mavromatis, V., Gonzalez-Gonzalez, A. and Oelkers, E.H., 2014. Kinetic and
2181 thermodynamic controls of divalent metals isotope composition in carbonate:
2182 Experimental investigations and applications. In: J. Gaillardet (Editor), *Geochemistry*
2183 *of the Earth's Surface Ges-10. Procedia Earth and Planetary Science*, pp. 168-172.

2184 Schulz, M., Stonestrom, D., Lawrence, C., Bullen, T., Fitzpatrick, J., Kyker-Snowman, E.,
2185 Manning, J. and Mnich, M., 2016. Structured heterogeneity in a marine terrace
2186 chronosequence: Upland mottling. *Vadose Zone Journal*, 15(2).

2187 Sharon, I. and Banfield, J.F., 2013. Genomes from Metagenomics. *Science*, 342(6162): 1057-
2188 1058.

2189 Shi, Y., Davis, K.J., Duffy, C.J. and Yu, X., 2013. Development of a Coupled Land Surface
2190 Hydrologic Model and Evaluation at a Critical Zone Observatory. *Journal of*
2191 *Hydrometeorology*, 14(5): 1401-1420.

2192 Singh, V.P., 1995. Computer models of watershed hydrology. *Water Resources Publications*.

2193 Singha, K., Li, L., Day-Lewis, F.D. and Regberg, A.B., 2011. Quantifying solute transport
2194 processes: Are chemically "conservative" tracers electrically conservative?
2195 *Geophysics*, 76(1): F53-F63.

2196 Smithwick, E.A.H., Lucash, M.S., McCormack, M.L. and Sivandran, G., 2014. Improving
2197 the representation of roots in terrestrial models. *Ecological Modelling*, 291: 193-204.

2198 Soetaert, K., Herman, P.M.J. and Middelburg, J.J., 1996. A model of early diagenetic
2199 processes from the shelf to abyssal depths. *Geochimica et Cosmochimica Acta*, 60(6):
2200 1019-1040.

2201 Soler, J.M. and Mader, U.K., 2005. Interaction between hyperalkaline fluids and rocks
2202 hosting repositories for radioactive waste: Reactive transport simulations. *Nuclear*
2203 *Science and Engineering*, 151(1): 128-133.

2204 St. Clair, J., Moon, S., Holbrook, W.S., Perron, J.T., Riebe, C.S., Martel, S.J., Carr, B.,
 2205 Harman, C., Singha, K. and Richter, D.d., 2015. Geophysical imaging reveals
 2206 topographic stress control of bedrock weathering. *Science*, 350(6260): 534-538.
 2207 Steefel, C.I., Appelo, C.A.J., Arora, B., Jacques, D., Kalbacher, T., Kolditz, O., Lagneau, V.,
 2208 Lichtner, P.C., Mayer, K.U., Meeussen, J.C.L., Molins, S., Moulton, D., Shao, H.,
 2209 Šimůnek, J., Spycher, N., Yabusaki, S.B. and Yeh, G.T., 2014. Reactive transport
 2210 codes for subsurface environmental simulation. *Computational Geosciences*: 1-34.
 2211 Steefel, C.I., Appelo, C.A.J., Arora, B., Jacques, D., Kalbacher, T., Kolditz, O., Lagneau, V.,
 2212 Lichtner, P.C., Mayer, K.U., Meeussen, J.C.L., Molins, S., Moulton, D., Shao, H.,
 2213 Šimůnek, J., Spycher, N., Yabusaki, S.B. and Yeh, G.T., 2015a. Reactive transport
 2214 codes for subsurface environmental simulation. *Computational Geosciences*, 19(3):
 2215 445-478.
 2216 Steefel, C.I., DePaolo, D.J. and Lichtner, P.C., 2005. Reactive transport modeling: An
 2217 essential tool and a new research approach for the Earth Sciences. *Earth and Planetary
 2218 Science Letters*, 240: 539-558.
 2219 Steefel, C.I., Emmanuel, S. and Anovitz, L.M., 2015b. PREFACE. *Reviews in Mineralogy
 2220 and Geochemistry*, 80(1): iii-v.
 2221 Steefel, C.I. and Lasaga, A.C., 1994. A coupled model for transport of multiple chemical
 2222 species and kinetic precipitation/dissolution reactions with application to reactive
 2223 flow in single phase hydrothermal systems. *Am. J. Sci.*, 294: 529-592.
 2224 Stöckle, C.O., Donatelli, M. and Nelson, R., 2003. CropSyst, a cropping systems simulation
 2225 model. *European Journal of Agronomy*, 18(3-4): 289-307.
 2226 Stockmann, U., Adams, M.A., Crawford, J.W., Field, D.J., Henakaarchchi, N., Jenkins, M.,
 2227 Minasny, B., McBratney, A.B., Courcelles, V.d.R.d., Singh, K., Wheeler, I., Abbott,
 2228 L., Angers, D.A., Baldock, J., Bird, M., Brookes, P.C., Chenu, C., Jastrow, J.D., Lal,
 2229 R., Lehmann, J., O'Donnell, A.G., Parton, W.J., Whitehead, D. and Zimmermann, M.,
 2230 2013. The knowns, known unknowns and unknowns of sequestration of soil organic
 2231 carbon. *Agriculture, Ecosystems & Environment*, 164: 80-99.
 2232 Stonedahl, S.H., Harvey, J.W., Wörman, A., Salehin, M. and Packman, A.I., 2010. A
 2233 multiscale model for integrating hyporheic exchange from ripples to meanders. *Water
 2234 Resources Research*, 46(12): n/a-n/a.
 2235 Sudicky, E.A., 1986. A natural gradient experiment on solute transport in a sand aquifer:
 2236 spatial variability of hydraulic conductivity and its role in the dispersion process.
 2237 *Water Resources Research*, 22(13): 2069-2086.
 2238 Sudicky, E.A., Illman, W.A., Goltz, I.K., Adams, J.J. and McLaren, R.G., 2010.
 2239 Heterogeneity in hydraulic conductivity and its role on the macroscale transport of a
 2240 solute plume: From measurements to a practical application of stochastic flow and
 2241 transport theory. *Water Resources Research*, 46.
 2242 Sullivan, P.L., Ma, L., West, N., Jin, L., Karwan, D.L., Noireaux, J., Steinhoefel, G., Gaines,
 2243 K.P., Eissenstat, D.M., Gaillardet, J., Derry, L.A., Meek, K., Hynek, S. and Brantley,
 2244 S.L., 2016. CZ-tope at Susquehanna Shale Hills CZO: Synthesizing multiple isotope
 2245 proxies to elucidate Critical Zone processes across timescales in a temperate forested
 2246 landscape. *Chemical Geology*.
 2247 Sulman, B.N., Phillips, R.P., Oishi, A.C., Shevliakova, E. and Pacala, S.W., 2014. Microbe-
 2248 driven turnover offsets mineral-mediated storage of soil carbon under elevated CO₂.
 2249 *Nature Clim. Change*, 4(12): 1099-1102.
 2250 Sundquist, E.T. and Visser, K., 2003. The Geologic History of the Carbon Cycle. In: H.D.
 2251 Holland and K.K. Turekian (Editors), *Treatise on Geochemistry*. Pergamon, Oxford,
 2252 pp. 425-472.

2253 Surasani, V.K., Li, L., Ajo-Franklin, J.B., Hubbard, C., Hubbard, S.S. and Wu, Y., 2013.
 2254 Bioclogging and Permeability Alteration by *L. mesenteroides* in a Sandstone
 2255 Reservoir: A Reactive Transport Modeling Study. *Energy & Fuels*, 27(11): 6538-
 2256 6551.

2257 Sverdrup, H., Warfvinge, P., Blake, L. and Goulding, K., 1995. Modelling recent and historic
 2258 soil data from the Rothamsted Experimental Station, UK using SAFE. *Agriculture,*
 2259 *ecosystems & environment*, 53(2): 161-177.

2260 Szecsody, J.E., Chilakapati, A., Zachara, J.M. and Garvin, A.L., 1998a. Influence of iron
 2261 oxide inclusion shape on Co(II/III)EDTA reactive transport through spatially
 2262 heterogeneous sediment. *Water Resources Research*, 34(10): 2501-2514.

2263 Szecsody, J.E., Zachara, J.M., Chilakapati, A., Jardine, P.M. and Ferency, A.S., 1998b.
 2264 Importance of flow and particle-scale heterogeneity on {CoII/III}EDTA reactive
 2265 transport. *J. Hydro.*, 209(1-4): 112-136.

2266 Takagi, K. and Lin, H.S., 2011. Temporal Dynamics of Soil Moisture Spatial Variability in
 2267 the Shale Hills Critical Zone Observatory. *Vadose Zone Journal*, 10(3): 832-842.

2268 Tang, G., Wu, W.-M., Watson, D.B., Parker, J.C., Schadt, C.W., Shi, X. and Brooks, S.C.,
 2269 2013. U(VI) Bioreduction with Emulsified Vegetable Oil as the Electron Donor –
 2270 Microcosm Tests and Model Development. *Environmental Science & Technology*,
 2271 47(7): 3209-3217.

2272 Tang, J. and Riley, W.J., 2015. Weaker soil carbon-climate feedbacks resulting from
 2273 microbial and abiotic interactions. *Nature Clim. Change*, 5(1): 56-60.

2274 Tang, J.Y. and Riley, W.J., 2013. A total quasi-steady-state formulation of substrate uptake
 2275 kinetics in complex networks and an example application to microbial litter
 2276 decomposition. *Biogeosciences*, 10(12): 8329-8351.

2277 Tartakovsky, A.M., Scheibe, T.D. and Meakin, P., 2009. Pore-Scale Model for Reactive
 2278 Transport and Biomass Growth. *Journal of Porous Media*, 12(5): 417-434.

2279 Tartakovsky, G.D., Tartakovsky, A.M., Scheibe, T.D., Fang, Y., Mahadevan, R. and Lovley,
 2280 D.R., 2013. Pore-scale simulation of microbial growth using a genome-scale
 2281 metabolic model: Implications for Darcy-scale reactive transport. *Advances in Water*
 2282 *Resources*, 59: 256-270.

2283 Therrien, R., McLaren, R., Sudicky, E. and Panday, S., 2010. HydroGeoSphere: A three-
 2284 dimensional numerical model describing fully-integrated subsurface and surface flow
 2285 and solute transport. Groundwater Simulations Group, University of Waterloo,
 2286 Waterloo, ON.

2287 Thomas, E.M., Lin, H., Duffy, C.J., Sullivan, P.L., Holmes, G.H., Brantley, S.L. and Jin, L.,
 2288 2013. Spatiotemporal Patterns of Water Stable Isotope Compositions at the Shale
 2289 Hills Critical Zone Observatory: Linkages to Subsurface Hydrologic Processes.
 2290 *Vadose Zone Journal*, 12(4).

2291 Thorley, R.M., Taylor, L.L., Banwart, S.A., Leake, J.R. and Beerling, D.J., 2015. The role of
 2292 forest trees and their mycorrhizal fungi in carbonate rock weathering and its
 2293 significance for global carbon cycling. *Plant Cell Environ*, 38(9): 1947-61.

2294 Thornton, P.E., Doney, S.C., Lindsay, K., Moore, J.K., Mahowald, N., Randerson, J.T., Fung,
 2295 I., Lamarque, J.F., Feddes, J.J. and Lee, Y.H., 2009. Carbon-nitrogen interactions
 2296 regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general
 2297 circulation model. *Biogeosciences*, 6(10): 2099-2120.

2298 Tipper, E.T., Calmels, D., Gaillardet, J., Louvat, P., Capmas, F. and Dubacq, B., 2012a.
 2299 Positive correlation between Li and Mg isotope ratios in the river waters of the
 2300 Mackenzie Basin challenges the interpretation of apparent isotopic fractionation
 2301 during weathering. *Earth and Planetary Science Letters*, 333: 35-45.

2302 Tipper, E.T., Lemarchand, E., Hindshaw, R.S., Reynolds, B.C. and Bourdon, B., 2012b.
 2303 Seasonal sensitivity of weathering processes: Hints from magnesium isotopes in a
 2304 glacial stream. *Chemical Geology*, 312: 80-92.

2305 Torn, M.S., Trumbore, S.E., Chadwick, O.A., Vitousek, P.M. and Hendricks, D.M., 1997.
 2306 Mineral control of soil organic carbon storage and turnover. *Nature*, 389(6647): 170-
 2307 173.

2308 Tricca, A., Wasserburg, G.J., Porcelli, D. and Baskaran, M., 2001. The transport of U- and
 2309 Th-series nuclides in a sandy unconfined aquifer. *Geochimica Et Cosmochimica Acta*,
 2310 65(8): 1187-1210.

2311 Tron, S., Bodner, G., Laio, F., Ridolfi, L. and Leitner, D., 2015. Can diversity in root
 2312 architecture explain plant water use efficiency? A modeling study. *Ecological*
 2313 *Modelling*, 312(0): 200-210.

2314 Trumbore, S., 2009. Radiocarbon and Soil Carbon Dynamics, *Annual Review of Earth and*
 2315 *Planetary Sciences. Annual Review of Earth and Planetary Sciences*, pp. 47-66.

2316 Trumbore, S.E., 1993. Comparison of carbon dynamics in tropical and temperate soils using
 2317 radiocarbon measurements. *Global Biogeochemical Cycles*, 7(2): 275-290.

2318 Trumbore, S.E. and Czimczik, C.I., 2008. An Uncertain Future for Soil Carbon. *Science*,
 2319 321(5895): 1455-1456.

2320 Trumbore, S.E., Davidson, E.A., Barbosa de Camargo, P.n., Nepstad, D.C. and Martinelli,
 2321 L.A., 1995. Belowground cycling of carbon in forests and pastures of eastern
 2322 Amazonia.

2323 Tutolo, B.M., Luhmann, A.J., Kong, X.-Z., Saar, M.O. and Seyfried Jr, W.E., 2015. CO2
 2324 sequestration in feldspar-rich sandstone: Coupled evolution of fluid chemistry,
 2325 mineral reaction rates, and hydrogeochemical properties. *Geochimica et*
 2326 *Cosmochimica Acta*, 160: 132-154.

2327 Vallino, J.J., 2003. Modeling Microbial Consortia as Distributed Metabolic Networks.
 2328 *Biological Bulletin*, 204(2): 174-179.

2329 Van Breukelen, B.M., Hunkeler, D. and Volkerling, F., 2005. Quantification of sequential
 2330 chlorinated ethene degradation by use of a reactive transport model incorporating
 2331 isotope fractionation. *Environmental science & technology*, 39(11): 4189-4197.

2332 Van Breukelen, B.M. and Rolle, M., 2012a. Transverse Hydrodynamic Dispersion Effects on
 2333 Isotope Signals in Groundwater Chlorinated Solvents' Plumes. *Environmental Science*
 2334 *& Technology*, 46(14): 7700-7708.

2335 Van Breukelen, B.M. and Rolle, M., 2012b. Transverse Hydrodynamic Dispersion Effects on
 2336 Isotope Signals in Groundwater Chlorinated Solvents' Plumes. *Environmental*
 2337 *Science & Technology*, 46(14): 7700-7708.

2338 Van Cappellen, P. and Wang, Y.F., 1996. Cycling of iron and manganese in surface
 2339 sediments: a general theory for the coupled transport and reaction of carbon, oxygen,
 2340 nitrogen, sulfur, iron, and manganese. *Am J Sci*, 296: 197-243.

2341 VanderKwaak, J.E. and Loague, K., 2001. Hydrologic-response simulations for the R-5
 2342 catchment with a comprehensive physics-based model. *Water Resources Research*,
 2343 37(4): 999-1013.

2344 Veen, J.A.V. and Paul, E.A., 1981. ORGANIC CARBON DYNAMICS IN GRASSLAND
 2345 SOILS. 1. BACKGROUND INFORMATION AND COMPUTER SIMULATION.
 2346 *Canadian Journal of Soil Science*, 61(2): 185-201.

2347 Vieira-Silva, S. and Rocha, E.P., 2010. The systemic imprint of growth and its uses in
 2348 ecological (meta)genomics. *PLoS Genet*, 6(1): e1000808.

2349 von Blanckenburg, F. and Bouchez, J., 2014. River fluxes to the sea from the ocean's Be-
 2350 10/Be-9 ratio. *Earth and Planetary Science Letters*, 387: 34-43.

- von Blanckenburg, F. and Schuessler, J.A., 2014. Element cycling in the Critical Zone as viewed by new isotope tools. In: J. Gaillardet (Editor), *Geochemistry of the Earth's Surface* Ges-10. *Procedia Earth and Planetary Science*, pp. 173-178.
- von Lützow, M. and Kögel-Knabner, I., 2009. Temperature sensitivity of soil organic matter decomposition—what do we know? *Biology and Fertility of Soils*, 46(1): 1-15.
- von Lützow, M., Kögel-Knabner, I., Ekschmitt, K., Flessa, H., Guggenberger, G., Matzner, E. and Marschner, B., 2007. SOM fractionation methods: Relevance to functional pools and to stabilization mechanisms. *Soil Biology and Biochemistry*, 39(9): 2183-2207.
- Walker, J.C.G., Hays, P.B. and Kasting, J.F., 1981. A negative feedback mechanism for the long-term stabilization of Earth's surface temperature. *Journal of Geophysical Research-Oceans and Atmospheres*, 86(C10): 9776-9782.
- Wallenstein, M.D. and Vilgalys, R.J., 2005. Quantitative analyses of nitrogen cycling genes in soils. *Pedobiologia*, 49(6): 665-672.
- Walter, C., Rossel, R.A.V. and McBratney, A.B., 2003. Spatio-Temporal Simulation of the Field-Scale Evolution of Organic Carbon over the Landscape. *Soil Sci. Soc. Am. J.*, 67(5): 1477-1486.
- Wang, L. and Li, L., 2015. Illite Spatial Distribution Patterns Dictate Cr(VI) Sorption Macrocapacity and Macrokinetics. *Environmental Science & Technology*, 49(3): 1374-1383.
- Wanner, C., Druhan, J., Amos, R., Alt-Epping, P. and Steefel, C., 2014. Benchmarking the simulation of Cr isotope fractionation. *Computational Geosciences*: 1-25.
- Wanner, C., Druhan, J., Amos, R., Alt-Epping, P. and Steefel, C., 2015. Benchmarking the simulation of Cr isotope fractionation. *Computational Geosciences*, 19(3): 497-521.
- Wardle, D.A., Bardgett, R.D., Klironomos, J.N., Setälä, H., Van Der Putten, W.H. and Wall, D.H., 2004. Ecological linkages between aboveground and belowground biota. *Science*, 304(5677): 1629-1633.
- Warren, J.M., Hanson, P.J., Iversen, C.M., Kumar, J., Walker, A.P. and Wulfschleger, S.D., 2014. Root structural and functional dynamics in terrestrial biosphere models – evaluation and recommendations. *New Phytologist*: n/a-n/a.
- Watkins, J.M., Nielsen, L.C., Ryerson, F.J. and DePaolo, D.J., 2013. The influence of kinetics on the oxygen isotope composition of calcium carbonate. *Earth and Planetary Science Letters*, 375: 349-360.
- Weiss, D.J., Boye, K., Caldeas, C. and Fendorf, S., 2014. Zinc Isotope Fractionation during Early Dissolution of Biotite Granite. *Soil Science Society of America Journal*, 78(1): 171-179.
- Werth, C.J., Cirpka, O.A. and Grathwohl, P., 2006. Enhanced mixing and reaction through flow focusing in heterogeneous porous media. *Water Resources Research*, 42(12).
- Werth, C.J., Zhang, C.Y., Brusseau, M.L., Oostrom, M. and Baumann, T., 2010. A review of non-invasive imaging methods and applications in contaminant hydrogeology research. *Journal of Contaminant Hydrology*, 113(1-4): 1-24.
- Whitaker, S., 1999. *The method of volume averaging*. Springer Netherlands.
- White, A.F., 1995. Chemical weathering rates of silicate minerals in soils. *Reviews in Mineralogy and Geochemistry*, 31(1): 407-461.
- White, A.F. and Brantley, S.L., 1995. Chemical weathering rates of silicate minerals: an overview. *Chemical Weathering Rates of Silicate Minerals*, 31: 1-22.
- White, A.F. and Brantley, S.L., 2003. The effect of time on the weathering of silicate minerals: why do weathering rates differ in the laboratory and field? *Chemical Geology*, 202(3-4): 479-506.

- White, A.F. and Peterson, M.L., 1990. Role of Reactive-Surface-Area Characterization in Geochemical Kinetic-Models. *Acs Symposium Series*, 416: 461-475.
- White, C.M., Kemanian, A.R. and Kaye, J.P., 2014. Implications of carbon saturation model structure for simulated nitrogen mineralization dynamics. *Biogeosciences Discussions*, 11(6): 9667-9695.
- Wieder, W.R., Bonan, G.B. and Allison, S.D., 2013. Global soil carbon projections are improved by modelling microbial processes. *Nature Clim. Change*, 3(10): 909-912.
- Wiederhold, J.G., Kraemer, S.M., Teutsch, N., Borer, P.M., Halliday, A.N. and Kretzschmar, R., 2006. Iron isotope fractionation during proton-promoted, ligand-controlled, and reductive dissolution of goethite. *Environmental Science & Technology*, 40(12): 3787-3793.
- Williams, L.B. and Hervig, F.L., 2005. Probing crystal growth mechanisms with non-traditional isotopes. *Geochimica Et Cosmochimica Acta*, 69(10): A211-A211.
- Williams, M.R. and Melack, J.M., 1997. Atmospheric deposition, mass balances, and processes regulating streamwater solute concentration in mixed-conifer catchments of the Sierra Nevada, California. *Biogeochemistry*, 37(2): 111-144.
- Willingham, T.W., Werth, C.J. and Valocchi, A.J., 2008. Evaluation of the effects of porous media structure on mixing-controlled reactions using pore-scale modeling and micromodel experiments. *Environmental Science & Technology*, 42(9): 3185-3193.
- Wimpenny, J., Gislason, S.R., James, R.H., Gannoun, A., Pogge Von Strandmann, P.A.E. and Burton, K.W., 2010. The behaviour of Li and Mg isotopes during primary phase dissolution and secondary mineral formation in basalt. *Geochimica Et Cosmochimica Acta*, 74(18): 5259-5279.
- Wolery, T.J., Jackson, K.J., Bourcier, W.L., Bruton, C.J., Viani, B.E., Knauss, K.G. and Delany, J.M., 1990. CURRENT STATUS OF THE EQ3/6 SOFTWARE PACKAGE FOR GEOCHEMICAL MODELING. *Acs Symposium Series*, 416: 104-116.
- Wood, B.D. and Whitaker, S., 1998. Diffusion and reaction in biofilms. *Chemical Engineering Science*, 53(3): 397-425.
- Woodbury, A.D. and Sudicky, E.A., 1991. The geostatistical characteristics of the borden aquifer. *Water Resources Research*, 27(4): 533-546.
- Wu, L., McGechan, M.B., McRoberts, N., Baddeley, J.A. and Watson, C.A., 2007. SPACSYS: Integration of a 3D root architecture component to carbon, nitrogen and water cycling—Model description. *Ecological Modelling*, 200(3-4): 343-359.
- Wu, Y., Surasani, V.K., Li, L. and Hubbard, S.S., 2014. Geophysical monitoring and reactive transport simulations of bioclogging processes induced by *Leuconostoc mesenteroides*. *Geophysics*, 79(1): E61-E73.
- Xu, T.F., Apps, J.A. and Pruess, K., 2003. Reactive geochemical transport simulation to study mineral trapping for CO₂ disposal in deep arenaceous formations. *Journal of Geophysical Research-Solid Earth*, 108(B2).
- Yabusaki, S.B., Fang, Y., Williams, K.H., Murray, C.J., Ward, A.L., Dayvault, R.D., Waichler, S.R., Newcomer, D.R., Spane, F.A. and Long, P.E., 2011. Variably saturated flow and multicomponent biogeochemical reactive transport modeling of a uranium bioremediation field experiment. *Journal of Contaminant Hydrology*, 126(3-4): 271-290.
- Yan, T., Fields, M.W., Wu, L., Zu, Y., Tiedje, J.M. and Zhou, J., 2003. Molecular diversity and characterization of nitrite reductase gene fragments (*nirK* and *nirS*) from nitrate- and uranium-contaminated groundwater. *Environmental Microbiology*, 5(1): 13-24.
- Yates, M.V., Yates, S.R., Wagner, J. and Gerba, C.P., 1987. Modeling virus survival and transport in the subsurface. *Journal of Contaminant Hydrology*, 1(3): 329-345.

- Ye, M., Neuman, S.P. and Meyer, P.D., 2004. Maximum likelihood Bayesian averaging of spatial variability models in unsaturated fractured tuff. *Water Resources Research*, 40(5).
- Yeh, G.-T., Huang, G., Cheng, H.-p., Zhang, F., Lin, H.-c., Edris, E. and Richards, D., 2006. A first principle, physics based watershed model: WASH123D. In: V.P. Singh and D.K. Frevert (Editors), *Watershed models*. CRC, Boca Raton.
- Yoo, K. and Mudd, S.M., 2008. Toward process-based modeling of geochemical soil formation across diverse landforms: A new mathematical framework. *Geoderma*, 146(1–2): 248-260.
- Yoon, H., Valocchi, A.J., Werth, C.J. and Dewers, T., 2012. Pore-scale simulation of mixing-induced calcium carbonate precipitation and dissolution in a microfluidic pore network. *Water Resources Research*, 48.
- Zachara, J., Brantley, S., Chorover, J., Ewing, R., Kerisit, S., Liu, C., Perfect, E., Rother, G. and Stack, A., 2016. Internal Domains of Natural Porous Media Revealed: Critical Locations for Transport, Storage, and Chemical Reaction. *Environmental Science & Technology*.
- Zachara, J.M., Long, P.E., Bargar, J., Davis, J.A., Fox, P., Fredrickson, J.K., Freshley, M.D., Konopka, A.E., Liu, C., McKinley, J.P., Rockhold, M.L., Williams, K.H. and Yabusaki, S.B., 2013. Persistence of uranium groundwater plumes: Contrasting mechanisms at two DOE sites in the groundwater–river interaction zone. *Journal of Contaminant Hydrology*, 147(0): 45-72.
- Zhang, C., Dehoff, K., Hess, N., Oostrom, M., Wietsma, T.W., Valocchi, A.J., Fouke, B.W. and Werth, C.J., 2010a. Pore-Scale Study of Transverse Mixing Induced CaCO₃ Precipitation and Permeability Reduction in a Model Subsurface Sedimentary System. *Environmental Science & Technology*, 44(20): 7833-7838.
- Zhang, C., Kang, Q., Wang, X., Zilles, J.L., Mueller, R.H. and Werth, C.J., 2010b. Effects of Pore-Scale Heterogeneity and Transverse Mixing on Bacterial Growth in Porous Media. *Environmental Science & Technology*, 44(8): 3085-3092.
- Zheng, C., Bianchi, M. and Gorelick, S.M., 2011. Lessons Learned from 25 Years of Research at the MADE Site. *Ground Water*, 49(5): 649-662.
- Zheng, Z. and Wang, G., 2007. Modeling the dynamic root water uptake and its hydrological impact at the Reserva Jaru site in Amazonia. *Journal of Geophysical Research: Biogeosciences* (2005–2012), 112(G4).
- Zhu, C., 2005. In situ feldspar dissolution rates in an aquifer. *Geochimica Et Cosmochimica Acta*, 69(6): 1435-1453.
- Zhuang, K., Izallalen, M., Mouser, P., Richter, H., Risso, C., Mahadevan, R. and Lovley, D.R., 2011. Genome-scale dynamic modeling of the competition between *Rhodoferax* and *Geobacter* in anoxic subsurface environments. *ISME J*, 5(2): 305-316.
- Ziegler, K., Chadwick, O.A., Brzezinski, M.A. and Kelly, E.F., 2005. Natural variations of delta Si-30 ratios during progressive basalt weathering, Hawaiian Islands. *Geochimica Et Cosmochimica Acta*, 69(19): 4597-4610.