Omega-3 fatty acids, lipid rafts, and T cell signaling

Tim Y. Hou^{a,b,c}, David N. McMurray^{b,c,e,f,g}, and Robert S. Chapkin^{a,b,c,d,e,f}

^aDepartment of Biochemistry and Biophysics; ^bDepartment of Nutrition and Food Science;

^cProgram in Integrative Nutrition and Complex Diseases; ^dCenter for Translational

Environmental Health Research; ^eDepartment of Veterinary Integrative Biosciences, Texas

A&M University, College Station, Texas, USA; ^fDepartment of Microbial Pathogenesis and

Immunology, Texas A&M University System Health Science Center, College Station, Texas,

77843, USA.

^gAddress correspondence to: Robert S. Chapkin, 111 Cater-Mattil, MS 2253, Texas A&M

University, College Station, TX, 77843. Phone: 1-979-845-0419. E-mail address: r-

chapkin@tamu.edu.

Abstract:

n-3 PUFA have been shown in many clinical studies to attenuate inflammatory responses. Although inflammatory responses are orchestrated by a wide spectrum of cells, CD4⁺ T cells play an important role in the etiology of many chronic inflammatory diseases such as inflammatory bowel disease and obesity. In light of recent concerns over the safety profiles of non-steroidal anti-inflammatory drugs (NSAIDs), alternatives such as bioactive nutraceuticals are becoming more attractive. In order for these agents to be accepted into mainstream medicine, however, the mechanisms by which nutraceuticals such as n-3 polyunsaturated fatty acids (PUFA) exert their anti-inflammatory effects must be fully elucidated. Lipid rafts are nanoscale, dynamic domains in the plasma membrane that are formed through favorable lipid-lipid (cholesterol, sphingolipids, and saturated fatty acids) and lipid-protein (membrane-actin cytoskeleton) interactions. These domains optimize the clustering of signaling proteins at the membrane to facilitate efficient cell signaling which is required for CD4⁺ T cell activation and differentiation. This review summarizes novel emerging data documenting the ability of n-3 PUFA to perturb membrane-cytoskeletal structure and function in CD4⁺ T cells. An understanding of these underlying mechanisms will provide a rationale for the use of n-3 PUFA in the treatment of chronic inflammation.

Keywords:

Omega-3 fatty acids

Lipid rafts

T cell activation

T cell differentiation

Contents:

36	1. Introduction
37	2. n-3 PUFA and lipid rafts in the CD4 ⁺ T cell plasma membrane.
38	2.1. Lipid rafts.
39	2.2. Effects of n-3 PUFA on lipid rafts in CD4 ⁺ T cell plasma membrane.
40	3. CD4 ⁺ T cell activation.
41	3.1. Role of lipid rafts in CD4 ⁺ T cell activation.
42	3.2. Effects of n-3 PUFA on CD4 ⁺ T cell activation.
43	4. CD4 ⁺ T cell differentiation.
44	4.1. Role of lipid rafts in CD4 ⁺ T cell differentiation.
45	4.2. Effects of n-3 PUFA on CD4 ⁺ T cell differentiation.
46	4.3. Effects of n-3 PUFA on Th17 CD4 ⁺ T cell differentiation.
47	5. Future Directions
48	6. Acknowledgements
49	7. References

50 **1. Introduction**

51 The mammalian immune system is critical in defending the host against foreign
52 pathogens and malignant cells. Under normal conditions, T lymphocytes circulate throughout
53 the body to survey for foreign antigens and transformed cells and target them for destruction.
54 However, under certain pathophysiological conditions, the adaptive immune system may lose the
55 ability to differentiate between self and foreign antigens, resulting in self-reactive T lymphocyte
56 activation and effector function. The result of the loss of self-tolerance could be autoimmune
57 diseases such as inflammatory bowel disease (IBD), e.g., Crohn's disease and ulcerative colitis
58 (Zenewicz et al., 2009). Alternatively, the adaptive immune system may become over-reactive

against self-antigens and unable to resolve appropriately, resulting in chronic inflammatory diseases such as rheumatoid arthritis. The mammalian immune system is comprised of the innate and the adaptive system; this review will focus on the adaptive arm, specifically CD4⁺ T lymphocytes. These cells typically further differentiate into other effector cell types (T_H1, T_H2, T_{reg}, T_H17), which have opposing roles in autoimmune diseases such as IBD (Zenewicz et al., 2009).

The cell membrane, composed of a phospholipid bilayer and a myriad of proteins, constitutes the outer boundary of the cell. Not only does the cell membrane control molecular transport, but it also regulates communication between the cell and its environment by transducing signals. . The first model of the plasma membrane, the fluid mosaic model, was proposed in 1972 by Singer and Nicolson (Singer and Nicolson, 1972). In this model, the phospholipid bilayer is thought of as a fluid, dynamic, passive solvent in which proteins are either embedded in and span the membrane (i.e. integral proteins), or loosely associate (i.e. peripheral proteins), with the phospholipid bilayer. The plasma membrane contains three classes of amphiphilic lipids: phospholipids, glycolipids, and sterols. Phospholipids and glycolipids are further subdivided into various fatty acids and headgroups at the *sn*-1, *sn*-2, and *sn*-3 positions (Fahy et al., 2009). For example, a major species of phosphatidylinositol-(4,5)-bisphosphate [PI(4,5)P₂] is composed of a saturated C18:0 fatty acid at the *sn*-1 position, an unsaturated C20:4^{Δ5,8,11,14} fatty acid at the *sn*-2 position, and *myo*-inositol 4,5-bisphosphate at the *sn*-3 position. The heterogeneity of the lipids in the plasma membrane is not well studied, with the potential to generate 9,000 – 100,000 different molecular species (Shevchenko and Simons, 2010; van Meer, 2005; Yetukuri et al., 2008). With all these layers of complexity, could lipids in the plasma membrane form local structures that can function to regulate cell signaling?

2. n-3 PUFA and lipid rafts in the CD4⁺ T cell plasma membrane.

2.1 *Lipid rafts*

In a simple model system where two lipids (one high melting temperature, one low melting temperature) and cholesterol are mixed together, micron-scale domains phase separate and are easily visualized using conventional fluorescence microscopy (Nicolau et al., 2006). These micron-sized microdomains, one example of local structures in the plasma membrane, can be observed in epithelial cells, where the apical plasma membrane is enriched in sphingolipids, while the basolateral plasma membrane is enriched in phosphatidylcholine (Zidovetzki and Levitan, 2007). Small invaginations in the plasma membrane, enriched with cholesterol, sphingolipids, and the protein caveolin, can also be found in many cells such as endothelial and intestinal epithelial cells and adipocytes (Ma et al., 2004; Toulmay and Prinz, 2013). Smaller, highly dynamic, nanoscale lipid rafts enriched in sphingolipids, cholesterol, and saturated fatty acids, have been proposed to play a role in signal transduction (**Fig. 1**). In fact, stable nanodomains can be visualized in yeast vacuole membranes in response to various stresses such as nutrient deprivation and pH change; proteins that sort to these vacuolar membranes also segregate to one of two domains, similar to what would be predicted by the simple system composed of two lipids and cholesterol (Toulmay and Prinz, 2013). These nanodomains are thought to organize select proteins to optimize their signaling capacity upon ligand engagement. Computer simulations suggest that in order for lipid rafts to promote protein-protein interactions, these nanoscale domains must be small (6 to 14 nm in diameter) in order to operate as protein concentrators in the plasma membrane (Nicolau et al., 2006).

Although lipid rafts can associate and dissociate as a mechanism to regulate the formation of raft phases in the plasma membrane, one way to achieve a stabilized raft phase (i.e.,

stabilize the size and/or lifetime of the raft) is the presence of the actin cytoskeleton. Monomeric actin (G-actin) protein is capable of polymerization to form long, complex filamentous actin (F-actin) that can provide the force required for organelle movement, and the scaffold required for stabilization of membrane raft phases. F-actin is connected to the plasma membrane by interacting with integral and membrane-associated proteins; e.g., various protein-actin cytoskeleton interactions in erythrocytes (Luna and Hitt, 1992). One current model is that these “membrane skeletons” form the “fences” in the plasma membrane, impeding the diffusion of membrane proteins and lipids (Kusumi et al., 2012).

The participation of the actin cytoskeleton in the formation of nanoscale domains was first postulated when it was observed that the coefficient of diffusion of phospholipid probes were significantly lower in live cells (Lee et al., 1993; Swaisgood and Schindler, 1989), than those estimated in artificial membranes (Ladha et al., 1996; Sonnleitner et al., 1999). This difference was also observed for transmembrane protein markers and glycosylphosphatidylinositol-anchored protein markers (Kusumi et al., 2012). One suggestion to explain the difference in the diffusion coefficient was the presence of the membrane cytoskeleton in live cells compared to artificial membranes. The requirement for the actin cytoskeleton in maintaining the properties of nanoscale domains was also observed using secondary ion mass spectrometry, in which treatment of NIH 3T3 mouse fibroblasts with latrunculin A (prevents G-actin polymerization) resulted in a random distribution of sphingomyelin instead of sphingomyelin-enriched microdomains in control fibroblasts (Frisz et al., 2013).

The actin cytoskeleton is bridged to membranes through interactions with integral proteins, or through proteins recruited to the plasma membrane by specific phospholipids. One such phospholipid is phosphatidylinositol-(4,5)-bisphosphate [PI(4,5)P₂], which comprises 5% of

the total phosphatidylinositol species, and 1% of the total plasma membrane phospholipid in mammalian cells (Di Paolo and De Camilli, 2006; Kwiatkowska, 2010). Classically, PI(4,5)P₂ is important in signal transduction because of the hydrolysis products diacylglycerol (DAG), and inositol(1,4,5)-trisphosphate (IP₃) produced by phospholipase C. However, an additional function of PI(4,5)P₂ is to recruit actin remodeling proteins to localized sites at the plasma membrane. PI(4,5)P₂ is thought to activate actin-regulatory proteins that induce actin polymerization, while concurrently inhibiting those proteins that promote actin disassembly. Indeed, the increase in local PI(4,5)P₂ results in stress-fiber formation, indicative of increased actin filament formation (Kwiatkowska, 2010). Due to multiple acidic phosphate groups on PI(4,5)P₂, basic protein domains are thought to play a role in recruiting proteins to the plasma membrane. For example, the Wiskott-Aldrich syndrome protein (WASP) is released from its autoinhibitory conformation when the basic domain binds to PI(4,5)P₂ (Padrick et al., 2008). Similarly, PI(4,5)P₂ can recruit and activate Rho family GTPase through its polybasic region. Some small GTPases play a role in cytoskeletal remodeling (Saarikangas et al., 2010).

A third interaction to consider in the formation and organization of optimal nanoscale lipid rafts is the role of higher-ordered protein complexes (i.e., protein-protein interactions), as proposed by Kai Simons and Akihiro Kusumi (Kusumi et al., 2011; Simons and Sampaio, 2011). As an example, epidermal growth factor receptor (EGFR) undergoes conformational change and dimerization upon binding to its ligand (EGF), resulting in downstream signaling (Turk et al., 2012; Turk and Chapkin, 2013). Optimal EGFR signaling, however, takes place in lipid rafts (Turk and Chapkin, 2013; 2015), highlighting the role of protein-protein interactions in driving the formation and organization of optimal nanoscale domains for cellular signaling.

2.2 *Effects of n-3 PUFA on lipid rafts in CD4⁺ T cell plasma membrane.*

Eicosapentaenoic acid (EPA, 20:5^{Δ5,8,11,14,17}) and docosaheptaenoic acid (DHA, 22:6^{Δ4,7,10,13,16,19}) are thought to be the major bioactive n-3 PUFA in fish oil. EPA and DHA are incorporated into the two major classes of phospholipids in the plasma membrane, phosphatidylethanolamine and phosphatidylcholine at the *sn*-2 position (Fan et al., 2004; Fan et al., 2003), consistent with the observation that saturated fatty acids occupy the *sn*-1 position of phospholipids, while unsaturated fatty acids are inserted at the *sn*-2 position. DHA is highly disordered and adopts various conformations on the subnanosecond time scale (Gawrisch and Soubias, 2008; Soubias and Gawrisch, 2007). By eliminating the double bond at the n-3 position to generate docosapentaenoic acid (DPA, 22:5^{Δ4,7,10,13,16}), chain dynamics of the fatty acid are reduced (Gawrisch and Soubias, 2008), suggesting that the unsaturation at the n-3 position affords unique properties to EPA and DHA in the plasma membrane.

Studies on ion channels have demonstrated that EPA and DHA can modulate membrane protein properties. The incorporation of DHA into the lipid bilayer resulted in a decrease in bilayer stiffness without changing the negative curvature of the bilayer (Bruno et al., 2007). DHA increased the appearance and lifetime of gramicidin channels, and decreased the free energy for channel formation. Oleic acid (OA, 18:1^{Δ9}) that intercalated into the plasma membrane at a greater rate, had no effect on gramicidin channel formation, suggesting that the effect of DHA was due to changes in bilayer properties and not due to specific binding (Bruno et al., 2007; Bruno et al., 2013). EPA and DHA inhibit cardiac Na⁺ and L-type Ca²⁺ channels (Xiao et al., 1997; Xiao et al., 1995), activate TRAAK-1 and TRPV1 channels (Fink et al., 1998; Matta et al., 2007), and increase desensitization of nAChR and GABA_A channels (Bouzat and Barrantes, 1993; Nabekura et al., 1998).

DHA is thought to play a major role in altering the size and/or stability of nanoscale lipid rafts in the plasma membrane. As a result of its highly disordered nature, DHA may modify the lateral organization of the plasma membrane by forming a distinct non-raft DHA domain (Shaikh et al., 2014; Wassall and Stillwell, 2008). This DHA domain is distinct from the sphingolipid and cholesterol enriched lipid rafts due to the highly flexible DHA, which is incompatible with the rigidity of sphingolipids and cholesterol. Incorporation of DHA into the plasma membrane results in the distinct non-raft DHA domain inserting itself into lipid rafts, increasing the size of lipid rafts. This mechanism would explain nuclear magnetic resonance (NMR) data which indicated that EPA and DHA are incorporated into both raft and non-raft domains in a phosphatidylethanolamine/sphingomyelin/cholesterol membrane mixture (Williams et al., 2012). Similar results have also been observed *in vivo* by analyzing phospholipids of CD4⁺ T cells after isolating the detergent-resistant and -soluble membrane fractions (Fan et al., 2004; Fan et al., 2003), as well as utilizing lipid-sensitive fluorescent probes (Kim et al., 2014; Kim et al., 2008).

An alternative mechanism proposed to explain the effects of n-3 PUFA on lipid raft organization comes from the flexibility of n-3 PUFA which is incompatible with the rigid cholesterol. This effect promotes the aggregation of cholesterol in the plasma membrane, resulting in the coalescence of sphingolipid and cholesterol lipid rafts from the bulk membrane (Wassall and Stillwell, 2009). Indeed, it has been shown that EPA and DHA can increase the size of lipid rafts in HEK cells, CD4⁺ T cells, and B cells (Chapkin et al., 2008; Kim et al., 2008; Rockett et al., 2012). The resulting alteration in the optimal size of nanoscale lipid rafts could perturb the cellular signaling required for CD4⁺ T cell activation and differentiation.

3. CD4⁺ T cell activation.

The engagement of the CD4⁺ T cell receptor (TCR) by an antigenic peptide presented by the major histocompatibility complex II (MHCII) molecule results in the formation of the immunological synapse (IS, **Fig. 2**). Proteins essential for propagating the signal are enriched at the IS to form the central supramolecular activation cluster (cSMAC). Adhesion molecules necessary to stabilize the IS form the peripheral supramolecular activation cluster (pSMAC) leading to the “bull’s eye” pattern of the IS (Lee et al., 2003; Monks et al., 1998). Proteins such as phosphatases that can abrogate signaling are excluded from the cSMAC and pSMAC in order for T cell activation to take place. At the IS, tyrosine kinases Lck and ZAP70 are activated and subsequently phosphorylate the adaptor protein linker for activation of T cells (LAT), leading to the assembly of the signalsome comprised of many proteins, including GADS, SLP76, NCK, ITK, VAV1, PAK, and PLC- γ 1 (Tybulewicz and Henderson, 2009). The proper formation of the IS is required for sustained T cell activation and is stabilized by the actin cytoskeleton (Gomez and Billadeau, 2008; Huang and Burkhardt, 2007; Meiri, 2005). When the adhesion proteins coalesce to form the pSMAC, the actin cytoskeleton is connected to the IS by proteins such as talin, vinculin, and WASp. The dynamics of the actin cytoskeleton, however, are also important for sustained T cell activation. It is thought that the centripetal flow of F-actin to the initial engagement site between TCR and MHCII is important for the strength and duration of T cell activation. Indeed, T cell activation can be disrupted by inhibiting actin polymerization (Campi et al., 2005; DeMond et al., 2008; Kaizuka et al., 2007; Yokosuka et al., 2005).

3.1 Role of lipid rafts in CD4⁺ T cell activation.

Lipid rafts in the plasma membrane are thought to be critical in CD4⁺ T cell activation. The interaction between Ag:MHCII and TCR results in a major reorganization of the nanoscale lipid rafts and signaling proteins to form the IS (Jury et al., 2007; Lillemeier et al., 2010; Meiri,

2005). IS formation involves changes in lipid-lipid interactions in the membranes. Liquid ordered (Lo) lipids such as cholesterol and sphingolipids accumulate at the IS (Burack et al., 2002; Fan et al., 2003; Kim et al., 2008; Zech et al., 2009). The involvement of lipid rafts in CD4⁺ T cell activation is conclusive; disruption of lipid rafts with various agents such as methyl- β -cyclodextrin (Xavier et al., 1998), 7-ketocholesterol (Rentero et al., 2008), or n-3 PUFA such as EPA (discussed below, Zech et al., 2009) resulted in suppressed T cell activation. Importantly, these data suggest that an alteration in the size of rafts can impact IS formation, and thus regulate T cell activation (Rentero et al., 2008; Zech et al., 2009).

3.2 *Effects of n-3 PUFA on CD4⁺ T cell activation.*

Since it has been shown that n-3 PUFA can alter the size and/or stability of lipid rafts in CD4⁺ T cells (Fan et al., 2004; Fan et al., 2003; Kim et al., 2014; Kim et al., 2008; Zech et al., 2009), it is not surprising that many studies have shown downstream perturbations in T cell activation when the CD4⁺ T cell plasma membrane is enriched with n-3 PUFA. Studies using the immortalized Jurkat T cell line demonstrated that n-3 PUFA displaced many of the signaling proteins necessary for T cell activation, including the Src family kinases Lck and Fyn (Stulnig et al., 1998) and LAT (Stulnig et al., 2001) from detergent-resistant membrane fractions. Recruitment and activation of signaling proteins such as PKC θ , LAT, Fas, PLC- γ 1, and F-actin were also altered (Fan et al., 2004; Kim et al., 2008; Sanderson and Calder, 1998). n-3 PUFA, however, did not universally inhibit protein localization; one study observed an enhancement of surface CTLA-4 expression, a negative regulator of T cell activation, in CD4⁺ T cells isolated from mice fed an EPA-enriched diet (Ly et al., 2006). It is now appreciated that n-3 PUFA broadly suppress downstream activation signaling in CD4⁺ T cells, including mitochondrial translocation (Yog et al., 2010), IL-2 secretion (Arrington et al., 2001; Chapkin et al., 2002; Fan

et al., 2004; Jolly et al., 1997; Ly et al., 2006; McMurray et al., 2000), and lymphoproliferation (Fan et al., 2008; Kim et al., 2008; McMurray et al., 2000).

Both lipid-lipid interactions and lipid-protein interactions (i.e., membrane-actin cytoskeleton interactions) are important for the formation of the IS. In a breakthrough finding, we recently demonstrated that n-3 PUFA directly perturb the membrane-actin cytoskeleton interactions by depleting the overall level of PI(4,5)P₂ in CD4⁺ T cells, resulting in the suppression/normalization of actin cytoskeletal rearrangement upon T cell activation (Hou et al., 2012). This effect was rescued by exogenous incubation of n-3 PUFA-enriched CD4⁺ T cells with PI(4, 5)P₂, demonstrating the direct perturbation of membrane-actin cytoskeleton interactions by n-3 PUFA. Perturbation of membrane-actin cytoskeleton interactions by n-3 PUFA may occur in other cell types. Colonocytes enriched with n-3 PUFA exhibited suppressed activation of cytoskeletal remodeling proteins such as PLC-γ1, Rac1, and Cdc42 (Turk et al., 2013).

4. CD4⁺ T cell differentiation.

Activated CD4⁺ T cells differentiate into pro-inflammatory effector subsets (T_H1, T_H17) in the presence of the cytokines such as interferon-γ (IFN-γ) and interleukin-12 (IL-12) for T_H1 cells, and Transforming growth factor-β (TGFβ), interleukin-6 (IL-6), and interleukin 21 (IL-21) for T_H17 cells (Zhu et al., 2010). CD4⁺ T cells can differentiate into anti-inflammatory T_H2 cells in the presence of IL-4 and IL-2, and into regulatory Treg cells (Zhu et al., 2010). Typically these CD4⁺ T effector cell subsets are defined by their transcriptional signatures. However, the role of the plasma membrane in directing the differentiation of CD4⁺ T cells is underappreciated. Although the anti-inflammatory effects of EPA and DHA have been well documented, their precise biochemical mechanisms of action with respect to cell

polarization/differentiation have yet to be clearly defined. We propose that the ability of n-3 PUFA to alter plasma membrane and cytoskeleton-dependent signaling may impact CD4⁺ T cell differentiation. These molecular targets may provide a novel therapeutic strategy for modulating the pathologic effector T cell subsets in autoimmune and chronic inflammatory diseases.

4.1 Role of lipid rafts in CD4⁺ T cell differentiation.

Early reports suggested that different CD4⁺ effector T cell populations depend upon distinct plasma membrane characteristics. For example, treatment of T cells with methyl- β -cyclodextrin to disturb lipid rafts demonstrated that T_H1, but not T_H2, cells were more sensitive as detected by suppressed Ca²⁺ influx upon antigen stimulation (Balamuth et al., 2001). Indeed, later studies demonstrated that T_H1 and T_H2 cells have distinct IS. The T_H1 cell IS is characterized by the bull's eye pattern where the cSMAC, comprised of Ag:MHCII/TCR interactions, is surrounded by the pSMAC populated with adhesion molecules. In contrast, the T_H2 cell IS is described as multifocal and dependent on the concentration of antigens (Thauland et al., 2008). Furthermore, individual human CD4⁺ T cells can be categorized into three classes based on the lipid order of their plasma membrane (low, intermediate, and high), as determined by the generalized polarization of di-4-ANEPPDHQ, a probe that changes fluorescence intensities at 570 nm and 620 nm based on the lipid order of its surrounding microenvironment upon incorporation into the plasma membrane (Miguel et al., 2011). CD4⁺ T cells with intermediate membrane order were associated with IFN- γ production (i.e. T_H1 phenotype), while high membrane order was associated with IL-4 production (i.e. T_H2 phenotype); low membrane order CD4⁺ T cells were associated with increased apoptosis. When CD4⁺ T cells were cultured in either T_H1 or T_H2-polarizing conditions, the membrane order of the CD4⁺ T cells changed to the corresponding T cell subset; intermediate membrane order was observed under T_H1

conditions, while high membrane order was observed under T_H2 conditions. To further link the role of membrane order with CD4⁺ T cell differentiation, the reduction of membrane order with 7-ketoxycholesterol resulted in an increase in the number of CD4⁺ T cells producing IFN- γ , indicative of a switch to an intermediate membrane order associated with the T_H1 phenotype. Clinically, CD4⁺ T cells isolated from human patients with systemic lupus erythematosus, Sjogrens Syndrome, or rheumatoid arthritis have an increased population of intermediate membrane order CD4⁺ T cells, associated with IFN- γ production and pro-inflammatory phenotype (McDonald et al., 2014; Miguel et al., 2011). T_H17 cells are also known to be affected by membrane order. Decreasing levels of glycosphingolipid, a lipid known to be associated with lipid rafts in CD4⁺ T cells, resulted in a reduction in T_H17 differentiation (Zhu et al., 2011). These studies demonstrate the structural importance of membrane nanoscale domains in directing CD4⁺ T cell differentiation.

4.2 *Effects of n-3 PUFA on CD4⁺ T cell differentiation.*

n-3 PUFA suppress CD4⁺ T cell activation and differentiation. Only T_H1-like cells exhibited enhanced activation-induced cell death (AICD) upon enrichment with n-3 PUFA, whereas T_H2-like cells were unaffected by n-3 PUFA (Switzer et al., 2004; Switzer et al., 2003). In addition, CD4⁺ T cells enriched with n-3 PUFA failed to polarize into T_H1 or T_H17 cells as efficiently when compared to control CD4⁺ T cells (Monk et al., 2013; Monk et al., 2012a; Zhang et al., 2005). These phenotypes were accompanied by changes in downstream signaling, such as reduced activation of T_H17-related STAT3, reduced expression of T_H17 signature cytokines (i.e., IL-17A), reduced expression of the T_H17 transcription factor (i.e., ROR γ t), and reduced surface expressions of critical surface receptors for T_H17 polarization (i.e., IL-6R and IL-23R) (Monk et al., 2013; Monk et al., 2012a; Zhang et al., 2005).

Interestingly, n-3 PUFA do not seem to affect the polarization of CD4⁺ T cells into T_H2 or T_{reg} subsets (Monk et al., 2013; Monk et al., 2012a; Monk et al., 2012b). This may be due to the different membrane properties optimized for polarization into these subsets (Miguel et al., 2011; Thauland et al., 2008). However, a recent study reported a decrease in CD4⁺ T cell differentiation under T_H2 polarization conditions in cells isolated from *Fat-1* mice, which incorporate n-3 PUFA into cell membranes endogenously (Jang et al., 2014). This highlights the need for more study into how n-3 PUFA affect CD4⁺ T cell differentiation, such as examining the dose of n-3 PUFA, and the differentiation environment the CD4⁺ T cells are selected under.

One mechanism by which long chain n-3 PUFA (i.e., EPA and DHA) suppress CD4⁺ T cell polarization into T_H17 cells involves the IL-6-gp-130-STAT3 signaling axis which regulates the earliest events in T_H17 cell differentiation (Bettelli et al., 2006; Jones et al., 2013; Jones et al., 2010; Nishihara et al., 2007; Veldhoen et al., 2006). A hexameric signaling complex composed of two IL-6 binding to two membrane bound IL-6 receptors (mIL-6R) and two molecules of glycoprotein 130 (gp130) is formed, leading to the phosphorylation of STAT3, translocation to the nucleus, and subsequent activation of RORγt, the master regulator of T_H17 transcription (Briso et al., 2008; Jones et al., 2010). As mentioned above, phosphorylation of STAT3 and activation of RORγt are both suppressed by n-3 PUFA in CD4⁺ T cells (Monk et al., 2013; Monk et al., 2012a). While the IL-6 receptor (IL-6R) is an 80 kDa surface protein expressed on naive CD4⁺ T cells (Betz and Muller, 1998), gp130 is ubiquitously expressed and mediates signal transduction for IL-6 and other cytokines (Silver and Hunter, 2010). Both IL-6R and gp130 are associated with lipid rafts; the localization of IL-6R at the plasma membrane is regulated by cholesterol (Matthews et al., 2003), while gp130 has been found to reside in lipid

rafts of kidney (Buk et al., 2004) and neuroepithelial (Yanagisawa et al., 2004) cell plasma membranes. Based upon the essential nature of the IL-6-gp130-STAT3 signaling axis in T_H17 differentiation (Veldhoen et al., 2006) and the fact that membrane modulation disrupts T_H17 development (Zhu et al., 2011), Allen and colleagues hypothesized that n-3 PUFA reduce T_H17 differentiation by interfering with IL-6 signaling in a lipid raft-dependent fashion (Allen et al., 2014). The presence of n-3 PUFA decreased surface expression of IL-6R and the association of gp130 with lipid rafts (**Fig. 3**). Specifically, the homodimerization of gp130 was reduced in activated CD4⁺ T cells enriched with n-3 PUFA. The lipid raft perturbations caused by n-3 PUFA led to downstream STAT3 phosphorylation. These results demonstrate that membrane alterations induced by n-3 PUFA suppress T_H17 differentiation by altering the IL6/gp130/Stat3 pathway.

5. Future Directions

A third potential target for n-3 PUFA in the formation and organization of optimal mesoscale lipid rafts are higher-ordered protein complexes (i.e., protein-protein interactions), as proposed by Kai Simons and Akihiro Kusumi (Kusumi et al., 2011; Simons and Sampaio, 2011). Similar to EGFR, Fas receptor (FasR) localizes in lipid rafts, and upon engagement with its ligand, Fas ligand (FasL), FasR undergoes oligomerization to induce favorable interactions between lipid rafts, again highlighting how protein-protein interactions can mediate lipid raft formation and organization (Muppidi and Siegel, 2004; Wang et al., 2010; Wilson et al., 2009). The TCR itself forms nanoclusters in the plasma membrane in a cholesterol-dependent fashion, requiring both cholesterol and sphingomyelin for the formation of TCR dimers (Molnar et al., 2012). Thus, the hierarchical organization of the plasma membrane is aided by lipid-lipid, lipid-protein, and protein-protein interactions. With advances in super-resolution microscopy to

visualize nanoclusters at the plasma membrane, it will be possible to determine whether n-3 PUFA can also affect protein-protein interactions in forming TCR nanoclusters at the plasma membrane of CD4⁺ T cells.

Not all n-3 PUFA are effective at perturbing plasma membrane lipid rafts. Phenotypes are not interchangeable when cells are treated with EPA or DHA (Corsetto et al., 2012; Turk et al., 2013), and biophysical studies have demonstrated that EPA and DHA do not disrupt lipid rafts with equal efficiency (Williams et al., 2012). Adding to the complication, 1-palmitoyl-2-docosahexaenoylphosphatidylethanolamine preferentially segregates into the non-raft plasma membrane, while 1-palmitoyl-2-docosahexaenoylphosphatidylcholine prefers the mesoscale raft domains of the plasma membrane (Shaikh et al., 2014), demonstrating that the headgroups of phospholipids can also affect the mechanisms by which these n-3 PUFA modulate plasma membrane properties.

Although it is appreciated that n-3 PUFA are pleotropic and can act upon targets outside of the plasma membrane, cogent data demonstrate that incorporation of n-3 PUFA can alter the biochemical and biophysical properties of CD4⁺ T cell plasma membranes, favorably modulating cytoskeletal dependent CD4⁺ T cell activation and differentiation (**Fig. 1**). In conclusion, n-3 PUFA target the plasma membrane of immune cells, and may contribute in the attenuation of autoimmune and chronic inflammatory diseases. However, in order to provide further rationale for their potential use as therapeutics, additional research is required to further elucidate how n-3 PUFA affect the nanoscale organization of the plasma membrane, and how this mediates the suppression of CD4⁺ T cell activation and differentiation into pathologic effector subsets.

Acknowledgements:

T.Y.H. was supported by pre-doctoral fellowships from the National Science and Engineering

379 Research Council (NSERC) of Canada and Texas A&M University System Health Science
380 Center Microbial Pathogenesis Training Grant Faculty. We also acknowledge support from U.S.
381 Department of Agriculture CSREES (Cooperative State Research, Education, and Extension
382 Service) Special Grant, “Designing Foods for Health” [2008-34402-17121] and NIH grants
383 CA129444 and P30ES023512.

384 **References:**

385 Allen, M.J., Fan, Y.Y., Monk, J.M., Hou, T.Y., Barhoumi, R., McMurray, D.N., Chapkin, R.S.,
386 2014. n-3 PUFAs reduce T-helper 17 cell differentiation by decreasing responsiveness to
387 interleukin-6 in isolated mouse splenic CD4(+) T cells. *J. Nutr.* 144, 1306-1313.

388 Arrington, J.L., McMurray, D.N., Switzer, K.C., Fan, Y.Y., Chapkin, R.S., 2001.
389 Docosahexaenoic acid suppresses function of the CD28 costimulatory membrane receptor in
390 primary murine and Jurkat T cells. *J. Nutr.* 131, 1147-1153.

391 Balamuth, F., Leitenberg, D., Unternaehrer, J., Mellman, I., Bottomly, K., 2001. Distinct patterns
392 of membrane microdomain partitioning in Th1 and th2 cells. *Immunity* 15, 729-738.

393 Bettelli, E., Carrier, Y., Gao, W., Korn, T., Strom, T.B., Oukka, M., Weiner, H.L., Kuchroo,
394 V.K., 2006. Reciprocal developmental pathways for the generation of pathogenic effector TH17
395 and regulatory T cells. *Nature* 441, 235-238.

396 Betz, U.A., Muller, W., 1998. Regulated expression of gp130 and IL-6 receptor alpha chain in T
397 cell maturation and activation. *Int. Immunol.* 10, 1175-1184.

398 Bouzat, C.B., Barrantes, F.J., 1993. Effects of long-chain fatty acids on the channel activity of
399 the nicotinic acetylcholine receptor. *Receptors Channels* 1, 251-258.

400 Briso, E.M., Dienz, O., Rincon, M., 2008. Cutting edge: soluble IL-6R is produced by IL-6R
401 ectodomain shedding in activated CD4 T cells. *J. Immunol.* 180, 7102-7106.

402 Bruno, M.J., Koeppe, R.E., 2nd, Andersen, O.S., 2007. Docosahexaenoic acid alters bilayer
 403 elastic properties. *Proc. Natl. Acad. Sci. U. S. A.* 104, 9638-9643.

404 Bruno, M.J., Rusinova, R., Gleason, N.J., Koeppe, R.E., 2nd, Andersen, O.S., 2013. Interactions
 405 of drugs and amphiphiles with membranes: modulation of lipid bilayer elastic properties by
 406 changes in acyl chain unsaturation and protonation. *Faraday Discuss.* 161, 461-480; discussion
 407 563-489.

408 Buk, D.M., Waibel, M., Braig, C., Martens, A.S., Heinrich, P.C., Graeve, L., 2004. Polarity and
 409 lipid raft association of the components of the ciliary neurotrophic factor receptor complex in
 410 Madin-Darby canine kidney cells. *J. Cell. Sci.* 117, 2063-2075.

411 Burack, W.R., Lee, K.H., Holdorf, A.D., Dustin, M.L., Shaw, A.S., 2002. Cutting edge:
 412 quantitative imaging of raft accumulation in the immunological synapse. *J. Immunol.* 169, 2837-
 413 2841.

414 Campi, G., Varma, R., Dustin, M.L., 2005. Actin and agonist MHC-peptide complex-dependent
 415 T cell receptor microclusters as scaffolds for signaling. *J. Exp. Med.* 202, 1031-1036.

416 Chapkin, R.S., Arrington, J.L., Apanasovich, T.V., Carroll, R.J., McMurray, D.N., 2002. Dietary
 417 n-3 PUFA affect TcR-mediated activation of purified murine T cells and accessory cell function
 418 in co-cultures. *Clin. Exp. Immunol.* 130, 12-18.

419 Chapkin, R.S., Wang, N., Fan, Y.Y., Lupton, J.R., Prior, I.A., 2008. Docosahexaenoic acid alters
 420 the size and distribution of cell surface microdomains. *Biochim. Biophys. Acta* 1778, 466-471.

421 Corsetto, P.A., Cremona, A., Montorfano, G., Jovenitti, I.E., Orsini, F., Arosio, P., Rizzo, A.M.,
 422 2012. Chemical-physical changes in cell membrane microdomains of breast cancer cells after
 423 omega-3 PUFA incorporation. *Cell. Biochem. Biophys.* 64, 45-59.

424 DeMond, A.L., Mossman, K.D., Starr, T., Dustin, M.L., Groves, J.T., 2008. T cell receptor
 425 microcluster transport through molecular mazes reveals mechanism of translocation. *Biophys. J.*
 426 94, 3286-3292.

427 Di Paolo, G., De Camilli, P., 2006. Phosphoinositides in cell regulation and membrane
 428 dynamics. *Nature* 443, 651-657.

429 Fahy, E., Subramaniam, S., Murphy, R.C., Nishijima, M., Raetz, C.R., Shimizu, T., Spener, F.,
 430 van Meer, G., Wakelam, M.J., Dennis, E.A., 2009. Update of the LIPID MAPS comprehensive
 431 classification system for lipids. *J. Lipid Res.* 50 Suppl, S9-14.

432 Fan, Y.Y., Kim, W., Callaway, E., Smith, R., Jia, Q., Zhou, L., McMurray, D.N., Chapkin, R.S.,
 433 2008. fat-1 transgene expression prevents cell culture-induced loss of membrane n-3 fatty acids
 434 in activated CD4⁺ T-cells. *Prostaglandins Leukot. Essent. Fatty Acids* 79, 209-214.

435 Fan, Y.Y., Ly, L.H., Barhoumi, R., McMurray, D.N., Chapkin, R.S., 2004. Dietary
 436 docosahexaenoic acid suppresses T cell protein kinase C θ lipid raft recruitment and IL-2
 437 production. *J. Immunol.* 173, 6151-6160.

438 Fan, Y.Y., McMurray, D.N., Ly, L.H., Chapkin, R.S., 2003. Dietary (n-3) polyunsaturated fatty
 439 acids remodel mouse T-cell lipid rafts. *J. Nutr.* 133, 1913-1920.

440 Fink, M., Lesage, F., Duprat, F., Heurteaux, C., Reyes, R., Fosset, M., Lazdunski, M., 1998. A
 441 neuronal two P domain K⁺ channel stimulated by arachidonic acid and polyunsaturated fatty
 442 acids. *EMBO J.* 17, 3297-3308.

443 Frisz, J.F., Lou, K., Klitzing, H.A., Hanafin, W.P., Lizunov, V., Wilson, R.L., Carpenter, K.J.,
 444 Kim, R., Hutcheon, I.D., Zimmerberg, J., Weber, P.K., Kraft, M.L., 2013. Direct chemical
 445 evidence for sphingolipid domains in the plasma membranes of fibroblasts. *Proc. Natl. Acad.*
 446 *Sci. U. S. A.* 110, E613-622.

447 Gawrisch, K., Soubias, O., 2008. Structure and dynamics of polyunsaturated hydrocarbon chains
 448 in lipid bilayers-significance for GPCR function. *Chem. Phys. Lipids* 153, 64-75.

449 Gomez, T.S., Billadeau, D.D., 2008. T cell activation and the cytoskeleton: you can't have one
 450 without the other. *Adv. Immunol.* 97, 1-64.

451 Hou, T.Y., Monk, J.M., Fan, Y.Y., Barhoumi, R., Chen, Y.Q., Rivera, G.M., McMurray, D.N.,
 452 Chapkin, R.S., 2012. n-3 polyunsaturated fatty acids suppress phosphatidylinositol 4,5-
 453 biphosphate-dependent actin remodelling during CD4+ T-cell activation. *Biochem. J.* 443, 27-
 454 37.

455 Huang, Y., Burkhardt, J.K., 2007. T-cell-receptor-dependent actin regulatory mechanisms. *J.*
 456 *Cell Sci.* 120, 723-730.

457 Jang, H.Y., Lim, K., Lee, S.M., Park, B.H., 2014. Effects of n-3 PUFA on the CD4(+) type 2
 458 helper T-cell-mediated immune responses in Fat-1 mice. *Mol. Nutr. Food Res.* 58, 365-375.

459 Jolly, C.A., Jiang, Y.H., Chapkin, R.S., McMurray, D.N., 1997. Dietary (n-3) polyunsaturated
 460 fatty acids suppress murine lymphoproliferation, interleukin-2 secretion, and the formation of
 461 diacylglycerol and ceramide. *J. Nutr.* 127, 37-43.

462 Jones, G.W., Greenhill, C.J., Williams, J.O., Nowell, M.A., Williams, A.S., Jenkins, B.J., Jones,
 463 S.A., 2013. Exacerbated inflammatory arthritis in response to hyperactive gp130 signalling is
 464 independent of IL-17A. *Ann. Rheum. Dis.* 72, 1738-1742.

465 Jones, G.W., McLoughlin, R.M., Hammond, V.J., Parker, C.R., Williams, J.D., Malhotra, R.,
 466 Scheller, J., Williams, A.S., Rose-John, S., Topley, N., Jones, S.A., 2010. Loss of CD4+ T cell
 467 IL-6R expression during inflammation underlines a role for IL-6 trans signaling in the local
 468 maintenance of Th17 cells. *J. Immunol.* 184, 2130-2139.

469 Jury, E.C., Flores-Borja, F., Kabouridis, P.S., 2007. Lipid rafts in T cell signalling and disease.
 470 *Semin. Cell. Dev. Biol.* 18, 608-615.

471 Kaizuka, Y., Douglass, A.D., Varma, R., Dustin, M.L., Vale, R.D., 2007. Mechanisms for
 472 segregating T cell receptor and adhesion molecules during immunological synapse formation in
 473 Jurkat T cells. *Proc. Natl. Acad. Sci. U. S. A.* 104, 20296-20301.

474 Kim, W., Barhoumi, R., McMurray, D.N., Chapkin, R.S., 2014. Dietary fish oil and DHA down-
 475 regulate antigen-activated CD4+ T-cells while promoting the formation of liquid-ordered
 476 mesodomains. *Br. J. Nutr.* 111, 254-260.

477 Kim, W., Fan, Y.Y., Barhoumi, R., Smith, R., McMurray, D.N., Chapkin, R.S., 2008. n-3
 478 polyunsaturated fatty acids suppress the localization and activation of signaling proteins at the
 479 immunological synapse in murine CD4+ T cells by affecting lipid raft formation. *J. Immunol.*
 480 181, 6236-6243.

481 Kusumi, A., Fujiwara, T.K., Chadda, R., Xie, M., Tsunoyama, T.A., Kalay, Z., Kasai, R.S.,
 482 Suzuki, K.G., 2012. Dynamic organizing principles of the plasma membrane that regulate signal
 483 transduction: commemorating the fortieth anniversary of Singer and Nicolson's fluid-mosaic
 484 model. *Annu. Rev. Cell. Dev. Biol.* 28, 215-250.

485 Kusumi, A., Suzuki, K.G., Kasai, R.S., Ritchie, K., Fujiwara, T.K., 2011. Hierarchical mesoscale
 486 domain organization of the plasma membrane. *Trends Biochem. Sci.* 36, 604-615.

487 Kwiatkowska, K., 2010. One lipid, multiple functions: how various pools of PI(4,5)P(2) are
 488 created in the plasma membrane. *Cell. Mol. Life Sci.* 67, 3927-3946.

489 Ladha, S., Mackie, A.R., Harvey, L.J., Clark, D.C., Lea, E.J., Brullemans, M., Duclohier, H.,
 490 1996. Lateral diffusion in planar lipid bilayers: a fluorescence recovery after photobleaching

491 investigation of its modulation by lipid composition, cholesterol, or alamethicin content and
 492 divalent cations. *Biophys. J.* 71, 1364-1373.

493 Lee, G.M., Zhang, F., Ishihara, A., McNeil, C.L., Jacobson, K.A., 1993. Unconfined lateral
 494 diffusion and an estimate of pericellular matrix viscosity revealed by measuring the mobility of
 495 gold-tagged lipids. *J. Cell Biol.* 120, 25-35.

496 Lee, K.H., Dinner, A.R., Tu, C., Campi, G., Raychaudhuri, S., Varma, R., Sims, T.N., Burack,
 497 W.R., Wu, H., Wang, J., Kanagawa, O., Markiewicz, M., Allen, P.M., Dustin, M.L.,
 498 Chakraborty, A.K., Shaw, A.S., 2003. The immunological synapse balances T cell receptor
 499 signaling and degradation. *Science* 302, 1218-1222.

500 Lillemeier, B.F., Mortelmaier, M.A., Forstner, M.B., Huppa, J.B., Groves, J.T., Davis, M.M.,
 501 2010. TCR and Lat are expressed on separate protein islands on T cell membranes and
 502 concatenate during activation. *Nat. Immunol.* 11, 90-96.

503 Luna, E.J., Hitt, A.L., 1992. Cytoskeleton--plasma membrane interactions. *Science* 258, 955-
 504 964.

505 Ly, L.H., Smith, R., Switzer, K.C., Chapkin, R.S., McMurray, D.N., 2006. Dietary
 506 eicosapentaenoic acid modulates CTLA-4 expression in murine CD4+ T-cells. *Prostaglandins*
 507 *Leukot. Essent. Fatty Acids* 74, 29-37.

508 Ma, D.W., Seo, J., Davidson, L.A., Callaway, E.S., Fan, Y.Y., Lupton, J.R., Chapkin, R.S., 2004.
 509 n-3 PUFA alter caveolae lipid composition and resident protein localization in mouse colon.
 510 *FASEB J.* 18, 1040-1042.

511 Matta, J.A., Miyares, R.L., Ahern, G.P., 2007. TRPV1 is a novel target for omega-3
 512 polyunsaturated fatty acids. *J. Physiol.* 578, 397-411.

513 Matthews, V., Schuster, B., Schutze, S., Bussmeyer, I., Ludwig, A., Hundhausen, C., Sadowski,
 514 T., Saftig, P., Hartmann, D., Kallen, K.J., Rose-John, S., 2003. Cellular cholesterol depletion
 515 triggers shedding of the human interleukin-6 receptor by ADAM10 and ADAM17 (TACE). *J.*
 516 *Biol. Chem.* 278, 38829-38839.

517 McDonald, G., Deepak, S., Miguel, L., Hall, C.J., Isenberg, D.A., Magee, A.I., Butters, T., Jury,
 518 E.C., 2014. Normalizing glycosphingolipids restores function in CD4⁺ T cells from lupus
 519 patients. *J. Clin. Invest.* 124, 712-724.

520 McMurray, D.N., Jolly, C.A., Chapkin, R.S., 2000. Effects of dietary n-3 fatty acids on T cell
 521 activation and T cell receptor-mediated signaling in a murine model. *J. Infect. Dis.* 182 Suppl 1,
 522 S103-107.

523 Meiri, K.F., 2005. Lipid rafts and regulation of the cytoskeleton during T cell activation. *Philos.*
 524 *Trans. R. Soc. Lond. B Biol. Sci.* 360, 1663-1672.

525 Miguel, L., Owen, D.M., Lim, C., Liebig, C., Evans, J., Magee, A.I., Jury, E.C., 2011. Primary
 526 human CD4⁺ T cells have diverse levels of membrane lipid order that correlate with their
 527 function. *J. Immunol.* 186, 3505-3516.

528 Molnar, E., Swamy, M., Holzer, M., Beck-Garcia, K., Worch, R., Thiele, C., Guigas, G., Boye,
 529 K., Luescher, I.F., Schwille, P., Schubert, R., Schamel, W.W., 2012. Cholesterol and
 530 sphingomyelin drive ligand-independent T-cell antigen receptor nanoclustering. *J. Biol. Chem.*
 531 287, 42664-42674.

532 Monk, J.M., Hou, T.Y., Turk, H.F., McMurray, D.N., Chapkin, R.S., 2013. n3 PUFAs reduce
 533 mouse CD4⁺ T-cell ex vivo polarization into Th17 cells. *J. Nutr.* 143, 1501-1508.

534 Monk, J.M., Hou, T.Y., Turk, H.F., Weeks, B., Wu, C., McMurray, D.N., Chapkin, R.S., 2012a.
 535 Dietary n-3 polyunsaturated fatty acids (PUFA) decrease obesity-associated Th17 cell-mediated
 536 inflammation during colitis. *PLoS One* 7, e49739.

537 Monk, J.M., Jia, Q., Callaway, E., Weeks, B., Alaniz, R.C., McMurray, D.N., Chapkin, R.S.,
 538 2012b. Th17 cell accumulation is decreased during chronic experimental colitis by (n-3) PUFA
 539 in Fat-1 mice. *J. Nutr.* 142, 117-124.

540 Monks, C.R., Freiberg, B.A., Kupfer, H., Sciaky, N., Kupfer, A., 1998. Three-dimensional
 541 segregation of supramolecular activation clusters in T cells. *Nature* 395, 82-86.

542 Muppidi, J.R., Siegel, R.M., 2004. Ligand-independent redistribution of Fas (CD95) into lipid
 543 rafts mediates clonotypic T cell death. *Nat. Immunol.* 5, 182-189.

544 Nabekura, J., Noguchi, K., Witt, M.R., Nielsen, M., Akaike, N., 1998. Functional modulation of
 545 human recombinant gamma-aminobutyric acid type A receptor by docosahexaenoic acid. *J. Biol.*
 546 *Chem.* 273, 11056-11061.

547 Nicolau, D.V., Jr., Burrage, K., Parton, R.G., Hancock, J.F., 2006. Identifying optimal lipid raft
 548 characteristics required to promote nanoscale protein-protein interactions on the plasma
 549 membrane. *Mol. Cell. Biol.* 26, 313-323.

550 Nishihara, M., Ogura, H., Ueda, N., Tsuruoka, M., Kitabayashi, C., Tsuji, F., Aono, H., Ishihara,
 551 K., Huseby, E., Betz, U.A., Murakami, M., Hirano, T., 2007. IL-6-gp130-STAT3 in T cells
 552 directs the development of IL-17+ Th with a minimum effect on that of Treg in the steady state.
 553 *Int. Immunol.* 19, 695-702.

554 Padrick, S.B., Cheng, H.C., Ismail, A.M., Panchal, S.C., Doolittle, L.K., Kim, S., Skehan, B.M.,
 555 Umetani, J., Brautigam, C.A., Leong, J.M., Rosen, M.K., 2008. Hierarchical regulation of
 556 WASP/WAVE proteins. *Mol. Cell* 32, 426-438.

557 Rentero, C., Zech, T., Quinn, C.M., Engelhardt, K., Williamson, D., Grewal, T., Jessup, W.,
 558 Harder, T., Gaus, K., 2008. Functional implications of plasma membrane condensation for T cell
 559 activation. *PLoS One* 3, e2262.

560 Rockett, B.D., Teague, H., Harris, M., Melton, M., Williams, J., Wassall, S.R., Shaikh, S.R.,
 561 2012. Fish oil increases raft size and membrane order of B cells accompanied by differential
 562 effects on function. *J. Lipid Res.* 53, 674-685.

563 Saarikangas, J., Zhao, H., Lappalainen, P., 2010. Regulation of the actin cytoskeleton-plasma
 564 membrane interplay by phosphoinositides. *Physiol. Rev.* 90, 259-289.

565 Sanderson, P., Calder, P.C., 1998. Dietary fish oil appears to prevent the activation of
 566 phospholipase C-gamma in lymphocytes. *Biochim. Biophys. Acta* 1392, 300-308.

567 Shaikh, S.R., Kinnun, J.J., Leng, X., Williams, J.A., Wassall, S.R., 2014. How polyunsaturated
 568 fatty acids modify molecular organization in membranes: Insight from NMR studies of model
 569 systems. *Biochim. Biophys. Acta* 1848, 211-219.

570 Shevchenko, A., Simons, K., 2010. Lipidomics: coming to grips with lipid diversity. *Nat Rev*
 571 *Mol. Cell. Biol.* 11, 593-598.

572 Silver, J.S., Hunter, C.A., 2010. gp130 at the nexus of inflammation, autoimmunity, and cancer.
 573 *J. Leukoc. Biol.* 88, 1145-1156.

574 Simons, K., Sampaio, J.L., 2011. Membrane organization and lipid rafts. *Cold Spring Harb.*
 575 *Perspect. Biol.* 3, a004697.

576 Singer, S.J., Nicolson, G.L., 1972. The fluid mosaic model of the structure of cell membranes.
 577 *Science* 175, 720-731.

578 Sonnleitner, A., Schutz, G.J., Schmidt, T., 1999. Free Brownian motion of individual lipid
 579 molecules in biomembranes. *Biophys. J.* 77, 2638-2642.

580 Soubias, O., Gawrisch, K., 2007. Docosahexaenoyl chains isomerize on the sub-nanosecond time
 581 scale. *J. Am. Chem. Soc.* 129, 6678-6679.

582 Stulnig, T.M., Berger, M., Sigmund, T., Raederstorff, D., Stockinger, H., Waldhausl, W., 1998.
 583 Polyunsaturated fatty acids inhibit T cell signal transduction by modification of detergent-
 584 insoluble membrane domains. *J. Cell Biol.* 143, 637-644.

585 Stulnig, T.M., Huber, J., Leitinger, N., Imre, E.M., Angelisova, P., Nowotny, P., Waldhausl, W.,
 586 2001. Polyunsaturated eicosapentaenoic acid displaces proteins from membrane rafts by altering
 587 raft lipid composition. *J. Biol. Chem.* 276, 37335-37340.

588 Swaisgood, M., Schindler, M., 1989. Lateral diffusion of lectin receptors in fibroblast
 589 membranes as a function of cell shape. *Exp. Cell. Res.* 180, 515-528.

590 Switzer, K.C., Fan, Y.Y., Wang, N., McMurray, D.N., Chapkin, R.S., 2004. Dietary n-3
 591 polyunsaturated fatty acids promote activation-induced cell death in Th1-polarized murine CD4+
 592 T-cells. *J. Lipid Res.* 45, 1482-1492.

593 Switzer, K.C., McMurray, D.N., Morris, J.S., Chapkin, R.S., 2003. (n-3) Polyunsaturated fatty
 594 acids promote activation-induced cell death in murine T lymphocytes. *J. Nutr.* 133, 496-503.

595 Thauland, T.J., Koguchi, Y., Wetzel, S.A., Dustin, M.L., Parker, D.C., 2008. Th1 and Th2 cells
 596 form morphologically distinct immunological synapses. *J. Immunol.* 181, 393-399.

597 Toulmay, A., Prinz, W.A., 2013. Direct imaging reveals stable, micrometer-scale lipid domains
 598 that segregate proteins in live cells. *J. Cell Biol.* 202, 35-44.

599 Turk, H.F., Barhoumi, R., Chapkin, R.S., 2012. Alteration of EGFR spatiotemporal dynamics
 600 suppresses signal transduction. *PLoS One* 7, e39682.

601 Turk, H.F., Chapkin, R.S., 2013. Membrane lipid raft organization is uniquely modified by n-3
 602 polyunsaturated fatty acids. *Prostaglandins Leukot. Essent. Fatty Acids* 88, 43-47.

603 Turk, H.F., Chapkin, R.S., 2015. Analysis of epidermal growth factor receptor dimerization by
604 BS(3) cross-linking. *Methods Mol. Biol.* 1233, 25-34.

605 Turk, H.F., Monk, J.M., Fan, Y.Y., Callaway, E.S., Weeks, B., Chapkin, R.S., 2013. Inhibitory
606 effects of omega-3 fatty acids on injury-induced epidermal growth factor receptor transactivation
607 contribute to delayed wound healing. *Am. J. Physiol. Cell Physiol.* 304, C905-917.

608 Tybulewicz, V.L., Henderson, R.B., 2009. Rho family GTPases and their regulators in
609 lymphocytes. *Nat. Rev. Immunol.* 9, 630-644.

610 van Meer, G., 2005. Cellular lipidomics. *EMBO J.* 24, 3159-3165.

611 Veldhoen, M., Hocking, R.J., Atkins, C.J., Locksley, R.M., Stockinger, B., 2006. TGFbeta in the
612 context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing
613 T cells. *Immunity* 24, 179-189.

614 Wang, L., Yang, J.K., Kabaleeswaran, V., Rice, A.J., Cruz, A.C., Park, A.Y., Yin, Q., Damko,
615 E., Jang, S.B., Raunser, S., Robinson, C.V., Siegel, R.M., Walz, T., Wu, H., 2010. The Fas-
616 FADD death domain complex structure reveals the basis of DISC assembly and disease
617 mutations. *Nat. Struct. Mol. Biol.* 17, 1324-1329.

618 Wassall, S.R., Stillwell, W., 2008. Docosahexaenoic acid domains: the ultimate non-raft
619 membrane domain. *Chem. Phys. Lipids* 153, 57-63.

620 Wassall, S.R., Stillwell, W., 2009. Polyunsaturated fatty acid-cholesterol interactions: domain
621 formation in membranes. *Biochim. Biophys. Acta* 1788, 24-32.

622 Williams, J.A., Batten, S.E., Harris, M., Rockett, B.D., Shaikh, S.R., Stillwell, W., Wassall, S.R.,
623 2012. Docosahexaenoic and eicosapentaenoic acids segregate differently between raft and
624 nonraft domains. *Biophys. J.* 103, 228-237.

625 Wilson, N.S., Dixit, V., Ashkenazi, A., 2009. Death receptor signal transducers: nodes of
 626 coordination in immune signaling networks. *Nat. Immunol.* 10, 348-355.

627 Xavier, R., Brennan, T., Li, Q., McCormack, C., Seed, B., 1998. Membrane compartmentation is
 628 required for efficient T cell activation. *Immunity* 8, 723-732.

629 Xiao, Y.F., Gomez, A.M., Morgan, J.P., Lederer, W.J., Leaf, A., 1997. Suppression of voltage-
 630 gated L-type Ca^{2+} currents by polyunsaturated fatty acids in adult and neonatal rat ventricular
 631 myocytes. *Proc. Natl. Acad. Sci. U. S. A.* 94, 4182-4187.

632 Xiao, Y.F., Kang, J.X., Morgan, J.P., Leaf, A., 1995. Blocking effects of polyunsaturated fatty
 633 acids on Na^{+} channels of neonatal rat ventricular myocytes. *Proc. Natl. Acad. Sci. U. S. A.* 92,
 634 11000-11004.

635 Yanagisawa, M., Nakamura, K., Taga, T., 2004. Roles of lipid rafts in integrin-dependent
 636 adhesion and gp130 signalling pathway in mouse embryonic neural precursor cells. *Genes Cells*
 637 9, 801-809.

638 Yetukuri, L., Ekroos, K., Vidal-Puig, A., Oresic, M., 2008. Informatics and computational
 639 strategies for the study of lipids. *Mol. Biosyst.* 4, 121-127.

640 Yog, R., Barhoumi, R., McMurray, D.N., Chapkin, R.S., 2010. n-3 polyunsaturated fatty acids
 641 suppress mitochondrial translocation to the immunologic synapse and modulate calcium
 642 signaling in T cells. *J. Immunol.* 184, 5865-5873.

643 Yokosuka, T., Sakata-Sogawa, K., Kobayashi, W., Hiroshima, M., Hashimoto-Tane, A.,
 644 Tokunaga, M., Dustin, M.L., Saito, T., 2005. Newly generated T cell receptor microclusters
 645 initiate and sustain T cell activation by recruitment of Zap70 and SLP-76. *Nat. Immunol.* 6,
 646 1253-1262.

647 Zech, T., Ejsing, C.S., Gaus, K., de Wet, B., Shevchenko, A., Simons, K., Harder, T., 2009.
 648 Accumulation of raft lipids in T-cell plasma membrane domains engaged in TCR signalling.
 649 EMBO J. 28, 466-476.
 650 Zenewicz, L.A., Antov, A., Flavell, R.A., 2009. CD4 T-cell differentiation and inflammatory
 651 bowel disease. Trends Mol. Med. 15, 199-207.
 652 Zhang, P., Smith, R., Chapkin, R.S., McMurray, D.N., 2005. Dietary (n-3) polyunsaturated fatty
 653 acids modulate murine Th1/Th2 balance toward the Th2 pole by suppression of Th1
 654 development. J. Nutr. 135, 1745-1751.
 655 Zhu, J., Yamane, H., Paul, W.E., 2010. Differentiation of effector CD4 T cell populations. Annu.
 656 Rev. Immunol. 28, 445-489.
 657 Zhu, Y., Gumlaw, N., Karman, J., Zhao, H., Zhang, J., Jiang, J.L., Maniatis, P., Edling, A.,
 658 Chuang, W.L., Siegel, C., Shayman, J.A., Kaplan, J., Jiang, C., Cheng, S.H., 2011. Lowering
 659 glycosphingolipid levels in CD4+ T cells attenuates T cell receptor signaling, cytokine
 660 production, and differentiation to the Th17 lineage. J. Biol. Chem. 286, 14787-14794.
 661 Zidovetzki, R., Levitan, I., 2007. Use of cyclodextrins to manipulate plasma membrane
 662 cholesterol content: evidence, misconceptions and control strategies. Biochim. Biophys. Acta
 663 1768, 1311-1324.
 664
 665

Figure Legends:

Figure 1. Proposed mechanisms by which n-3 PUFA modulate adaptive immune responses by

A) modulating lipid-lipid interactions in the plasma membrane; and **B)** altering plasma

membrane lipid-protein interactions by decreasing PI(4,5)P₂ level, thereby lowering the

recruitment of actin-binding proteins and suppressing actin cytoskeleton remodeling.

Consequently, incorporation of n-3 PUFA into the plasma membrane increase the size and/or

stability of the mesoscale lipid rafts and physiologically. This translates into suppressed CD4⁺ T

cell activation and differentiation. Red highlight indicates liquid ordered lipid rafts; Blue

indicates bulk membrane.

Figure 2. Immunological synapse at the **A)** cellular; and **B)** biochemical level. When CD4⁺ T

cells recognize cognate antigen presented by the antigen presenting cell, an immunological

synapse is formed at the cellular level. At the biochemical level, major lipidomic and proteomic

rearrangement occurs to propagate T cell activation. At the lipidomic level, lipids associated

with lipid rafts are found to coalesce at the immunological synapse (represented in red).

Adhesion molecules, such as LFA-1, are important for the stabilization of the immunological

synapse, while CD45, which is required to terminate T cell activation, is excluded from the

immunological synapse. Additionally, the actin cytoskeleton is important for T cell activation by

stabilizing the immunological synapse.

Figure 3. Proposed mechanism by which n-3 PUFA suppress T_H17 cell differentiation. n-3

PUFA directly modulate IL-6/gp130 signaling at the plasma membrane, thus decreasing T_H17

differentiation. **A)** n-3 PUFA decrease colocalization of gp130 in lipid rafts, as assessed by

colocalization with cholera toxin (CTxB), a marker of lipid rafts. Red highlight indicates liquid

ordered lipid rafts; Blue indicates bulk membrane. **B)** n-3 PUFA decrease the dimerization of

689 gp130 upon IL-6 stimulation, suppressing downstream T_H17 differentiation (Allen et al., 2014).

Figure 1

A) n-3 PUFA increase the size and/or stability of lipid rafts in the plasma membrane

Effect on lipid-lipid interactions

B) n-3 PUFA suppress phosphatidylinositol-(4,5)-bisphosphate dependent actin remodeling

Effect on lipid-protein interactions

n-3 PUFA containing phospholipids

Figure 2

A) Cellular

B) Biochemical

Figure 3

A) n-3 PUFA decrease the colocalization of gp130 with the lipid raft marker GM1

B) n-3 PUFA decrease gp130 dimerization upon IL-6 stimulation