1 **Hypothalamic-pituitary-adrenal axis activity is not elevated in a songbird (*Junco***
2 ***hyemalis*) preparing for migration**

3

4 Carolyn M. Bauer ^{a,*}, Katie B. Needham ^a, Chuong N. Le ^a, Emily C. Stewart ^a, Jessica L.
5 Graham ^a, Ellen D. Ketterson ^b and Timothy J. Greives ^a

6

7 ^a *Department of Biological Sciences, North Dakota State University, Fargo, ND, USA*

8 ^b *Department of Biology, Indiana University, Bloomington, IN, USA*

9

10 * Corresponding author. Fax +1 701 231 7149.

11 *E-mail address:* carolyn.marie.bauer@gmail.com (C.M. Bauer).

12

13 **Abstract**

14

15 During spring, increasing daylengths stimulate gonadal development in migratory birds.
16 However, late-stage reproductive development is typically postponed until migration has
17 been completed. The hypothalamic-pituitary-adrenal (HPA) axis regulates the secretion
18 of glucocorticoids, which have been associated with pre-migratory hyperphagia and
19 fattening. The HPA-axis is also known to suppress the hypothalamic-pituitary-gonadal
20 (HPG) axis, suggesting the possibility that final transition into the breeding life history
21 stage may be slowed by glucocorticoids. We hypothesized that greater HPA-axis activity
22 in individuals preparing for migration may foster preparation for migration while
23 simultaneously acting as a “brake” on the development of the HPG-axis. To test this

hypothesis, we sampled baseline corticosterone (CORT), stress-induced CORT, and negative feedback efficacy of Dark-eyed Juncos (*Junco hyemalis*) in an overwintering population that included both migratory (*J.h. hyemalis*) and resident (*J.h. carolinensis*) individuals. We predicted that compared to residents, migrants would have higher baseline CORT, higher stress-induced CORT, and weaker negative feedback. Juncos were sampled in western Virginia in early March, which was about 2–4 wk before migratory departure for migrants and 4–5 wk before first clutch initiation for residents. Contrary to our predictions, we found that migrants had lower baseline and stress-induced CORT and similar negative feedback efficacy compared with residents, which suggests that delayed breeding in migrants is influenced by other physiological mechanisms. Our findings also suggest that baseline CORT is not elevated during pre-migratory fattening, as migrants had lower baseline CORT and were fatter than residents.

Keywords: corticosterone, fattening, migration, negative feedback, reproductive timing, stress.

1. Introduction

By migrating, temperate-zone birds take advantage of seasonally abundant food resources during the summer months while improving their likelihood of survival by departing from locales that are harsh and resource-poor during winter (Lack, 1968). In preparation for mid- and long-distance migration, birds undergo a migratory developmental phase that includes several physiological changes such as flight muscle

hypertrophy, red blood cell proliferation (erythropoiesis), and fattening (Cornelius et al., 2013; Piersma et al., 1999; Ramenofsky, 2011). Fattening, driven by hyperphagia, is especially important for migrants because fat is the main fuel utilized during extended periods of flight (Ramenofsky, 1990). Initiation of the migratory developmental phase in spring is regulated by both endogenous circannual rhythms and increasing photoperiod or daylength (Gwinner, 1996a; Ramenofsky and Wingfield, 2007). Exposure to lengthening days causes increased secretion of gonadotropin-releasing hormone (GnRH), which plays a critical role in male spring migratory development through its stimulatory effects on androgen production and release (Deviche, 1995; Owen et al., 2014; Ramenofsky and Nemeth, 2014). Increasing daylengths also stimulate secretion of other neuropeptides, whose specific roles in migratory preparation are still largely unknown (Cornelius et al., 2013; Ramenofsky, 2011).

In addition to stimulating migratory development, increasing daylengths also activate gonadal development and other physiological and behavioral changes essential for breeding (Gwinner, 1996b; Perrins, 1970; Rowan, 1926). Androgens in particular are known to have a permissive effect on pre-migratory fattening in male birds (Mattocks, 1976; Ramenofsky and Nemeth, 2014; Tonra et al., 2011). Partial reproductive development before migratory departure allows migrants to commence breeding shortly after arrival to the breeding grounds. Nevertheless, migrants typically delay complete reproductive maturity until migration has been completed, likely because full gonadal development could interfere with migratory performance owing to increased body mass, metabolic pathway constraints, increased lipoprotein use, and decreased hematocrit (Williams, 2012).

Final transition into the breeding life history stage could be mediated by glucocorticoid production and release, which is controlled by the hypothalamic-pituitary-adrenal (HPA) axis. Corticosterone (CORT), the main glucocorticoid in birds, has numerous physiological effects and is generally considered important for migratory development due to its permissive effects on hyperphagia and fattening (Landys et al., 2004a; Landys et al., 2004b). Additionally, CORT has well-known suppressive effects on the hypothalamic-pituitary-gonadal (HPG) axis (Wingfield and Sapolsky, 2003). Therefore, CORT could be an important regulator for promoting migratory preparation and delaying full reproductive maturity until arrival at the breeding grounds. Indeed, in most bird species CORT levels increase during spring migratory preparation (Holberton et al., 2008; Romero, 2002). However, we also know that in many bird species gonadal recrudescence increases during migration (Bauchinger et al., 2007), apparently despite high CORT levels. One possible way to reconcile these observations is to suggest that gonadal recrudescence occurs during stopovers when CORT decreases (Eikenaar et al., 2014) from elevated levels during migratory flight (Falsone et al., 2009).

To determine whether increased CORT exposure during migratory preparation might act as a “brake” on the HPG-axis, we compared CORT profiles of migrant and resident Dark-eyed Juncos (*Junco hyemalis hyemalis* and *J.h. carolinensis*, respectively) that share the same overwintering site. Juncos were sampled for baseline CORT, stress-induced CORT, and negative feedback efficacy. Baseline CORT concentrations represent levels needed for everyday, predictable activities, and generally reflect current energy use (Sapolsky et al., 2000). Because baseline CORT concentrations represent CORT levels only when birds are undisturbed, we also measured stress-induced CORT to determine

relative sensitivity of migrants and residents to stressors (Sapolsky et al., 2000). Negative feedback is another important component of the endocrine stress response, as strong negative feedback can quickly reduce CORT secretion, whereas weak negative feedback prolongs CORT exposure. Compared to other components of the HPA-axis, weak negative feedback is the strongest indicator of chronic stress (Dickens and Romero, 2013). By measuring baseline CORT, stress-induced CORT, and negative feedback strength, we estimated each individual's potential *total exposure* to CORT. We predicted that if CORT plays a stimulatory role in migratory preparation, then prior to migratory departure migrant birds would have higher baseline CORT, higher stress-induced CORT, weaker negative feedback, and thus higher total CORT exposure as compared to resident birds.

2. Methods

2.1. Study animals

We studied an overwintering population of Dark-eyed Juncos (*Junco hyemalis*) at Mountain Lake Biological Station (37°22'N, 80°32'W), located in Giles County, western Virginia and managed by University of Virginia. We used *J. hyemalis* in this study because both resident (*J.h. carolinensis*: hereafter called "residents") and migrant (*J.h. hyemalis*: hereafter called "migrants") subspecies overwinter at Mountain Lake. On average, residents weigh more and are slightly larger than migrants, and males weigh more and are slightly larger than females (Nolan et al., 2002). Migrant males, migrant females, resident males, and resident females can be identified based on bill color, wing

chord length, and plumage characteristics (Ketterson and Nolan, 1976; Nolan et al., 2002). Plumage characteristics can also be used to age juncos as young (less than 1 year old) or old (greater than 1 year old) (Ketterson, 1979). This study was carried out March 12–19, 2015, which is about 4–5 wk before first clutch initiation for residents (April 16) and 2–4 wk before migratory departure for migrants. We estimated migrant departure date based on evidence that there were very few migrants (< 3% of junco population) when we returned to Mountain Lake in mid-April, and because migrants from this population showed very low levels of Zugunruhe throughout the month of March in a captive study (A. Fudickar, personal communication). During this study, both resident and migrant juncos were observed foraging in winter flocks and showed little evidence of territorial behavior.

2.2. Trapping and blood sampling

Birds were captured in baited Potter traps and mist-nets. Traps and nets were opened 1 hr after sunrise and closed at least 2.5 h before sunset (0830–1700h). While many vertebrates possess diel CORT rhythms (Rich and Romero, 2001; Romero and Remage-Healey, 2000), we found no relationships between baseline CORT concentrations and time of capture in our study (regression analysis, data not shown). While traps were open, 2–4 observers would scan with binoculars in order to determine time of capture. Birds were bled within 3 min of capture to obtain a baseline CORT sample (~60 µL) as CORT levels typically increase within 3 min of disturbance (Romero and Reed, 2005). Birds for which capture time could not be assigned were excluded from

all CORT analyses. For a sub-set of male resident and migrant juncos, we took an additional 60 μ L of blood within 5 min of capture so we could measure baseline testosterone, which would give us an estimate of relative reproductive readiness in resident and migrant males (Fudickar et al., in press). All blood samples were taken from the alar vein and were collected with heparinized, microhematocrit capillary tubes. Birds were kept in opaque, cloth bags between blood sampling time-points.

After baseline blood sample collection, birds were identified as residents or migrants, sexed, aged, banded if needed, and weighed to the nearest 0.25 g. Birds were given both abdominal and furcular subcutaneous fat scores, with a score of 0 meaning no visible fat and 5 meaning visible fat bulging (Nolan and Ketterson, 1983; O'Neal et al., 2011). Stress-induced blood samples (~30 μ L) were then taken 30 min after capture, as CORT peaks around this time-point in Dark-eyed Juncos (Schoech et al., 1999). Birds were then immediately given an intramuscular injection of dexamethasone (DEX) into the pectoral muscle. DEX is a synthetic glucocorticoid and is commonly used for determining negative feedback efficacy (Dickens et al., 2009; Romero, 2004; Sapolsky and Altmann, 1991). We used a 1 mg/kg of body weight dose, as previous studies have shown this dosage to effectively suppress endogenous CORT production in songbirds (Rich and Romero, 2001). Through a validation study, we determined that juncos maximally decrease CORT production 90 min after DEX injection. Therefore, we assessed negative feedback 90 min after DEX injection in our study (**Fig. 1**). After this last blood sample (~ 60 μ L), birds were then released at their site of capture.

Blood samples were kept in a cooler with icepacks for no more than 9 h before centrifugation. After centrifugation, plasma was drawn off and stored at -80° C until

further analysis. Sample sizes for full stress series were: migrant males ($n = 17$), migrant females ($n = 3$), resident males ($n = 24$), and resident females ($n = 6$). Sample sizes for testosterone were: migrant males ($n = 11$) and resident males ($n = 11$). Sample sizes for body weight and fat were: migrant males ($n = 29$), migrant females ($n = 12$), resident males ($n = 32$), and resident females ($n = 9$). All procedures were approved by the North Dakota State University Institutional Animal Care and Use Committee.

2.3. DEX validation study

Briefly, *J.h. hyemalis* individuals were trapped in Fargo, ND, USA during migration in November 2014. Stress-induced samples were taken 30 min after capture, and birds were then injected with DEX ($n = 8$) or a control phosphate buffer solution (PBS) ($n = 8$). Subsequent blood samples ($\sim 60 \mu\text{L}$) were taken 60 and 90 min after injection. While CORT levels did not significantly change over time after injection with PBS (**Fig. 1**; one-way repeated measures ANOVA, $F_{2,7} = 2.65$, $p = 0.11$); CORT levels significantly decreased both 60 and 90 min after injection with DEX (**Fig. 1**; one-way repeated measures ANOVA $F_{2,7} = 12.16$, $p < 0.01$, LSM post-hoc tests all $ps < 0.05$). However, compared to PBS-injected birds, DEX-injected birds had significantly stronger negative feedback only at the 90 min time-point (**Fig. 1**; t-tests with Bonferroni corrections, 60 min $t_{(1,14)} = 2.24$, $p = 0.084$; 90 min $t_{(1,15)} = 2.80$, $p = 0.03$).

2.4. CORT enzyme immunoassay

We measured CORT concentrations with an enzyme immunoassay kit (#KO14, Arbor Assays, Ann Arbor, MI, USA) that had low, reported cross-reactivity with DEX (0.12%). While previous studies have successfully used this kit in songbirds (DeVries and Jawor, 2013; Pryor and Casto, 2015), we further validated the kit for use in the Dark-eyed Junco. To test for parallelism, we compared slopes between our standard curve (created from kit-provided reference standards) and serially diluted pooled junco plasma. Parallelism assumptions were met ($F_{1,10} = 5.5$, $p = 0.96$) as residual variance did not significantly increase when the two lines shared a common slope (Grotjan and Keel, 1996). Briefly, we added 20 μ L plasma to 200 μ L ultra-pure water. Samples were then extracted two times with diethyl ether in a methanol/dry ice bath, dried with N_2 , and reconstituted with 200 μ L assay buffer. Samples were then plated in duplicate and processed according to the manufacturer's instructions. Samples were spread across plates to balance migrant/resident status, sex, and age. Reported assay sensitivity was 18.6 pg/mL and intra- and inter-assay variation were 4.5% and 9.3%, respectively. For samples with undetectable CORT levels, we assigned them a value of 62 pg/mL, as this was the lowest level of CORT we detected among our samples.

2.5. Testosterone enzyme immunoassay

We measured testosterone concentrations with an enzyme immunoassay kit (#901-065, Enzo Life Sciences Inc., Farmingdale, NY, USA) that had previously been validated for use in the Dark-eyed Junco (Clotfelter et al., 2004; Rosvall et al., 2012). Briefly, we added 30 μ L plasma to 200 μ L ultra-pure water. Samples were then extracted

two times with diethyl ether in a methanol/dry ice bath, dried with N₂, and reconstituted with 300 µL assay buffer. Samples were then plated in duplicate and processed according to the manufacturer's instructions. All samples were run on a single plate. Reported assay sensitivity was 5.67 pg/mL and intra-assay variation was 2.3%. For samples with undetectable testosterone levels, we assigned them a value of 10.8 pg/mL, as this was the lowest level of testosterone we detected among our samples.

2.6. Statistical analyses

Negative feedback efficacy was measured as the percent decrease in CORT from stress-induced levels (collected 30 min after capture) to those 90 min after DEX injection: $(\text{stress-induced CORT} - \text{post-DEX CORT}) / (\text{stress-induced CORT}) * 100$ (Lattin et al., 2012). Since baseline CORT, stress-induced CORT, and negative feedback have been shown to be regulated independently, we analyzed each CORT variable separately (Landys et al., 2006; Romero, 2006). Baseline CORT, stress-induced CORT, negative feedback efficacy, baseline testosterone, body weight, abdominal fat, and furcular fat were analyzed using mixed model ANOVAs. For all mixed models except testosterone, fixed effects included migrant/resident status, sex, and age (first-year/adult young or older than first-year/adult old). For testosterone, fixed effects include migrant/resident status and age. For baseline CORT, stress-induced CORT, and negative feedback analyses, EIA plate number was included as a random effect. Cohen's f^2 , a commonly used effect size measure for mixed models (Cohen, 1988; Selya et al., 2012), was reported for all significant main effects. Cohen's f^2 values of 0.02, 0.15, and 0.35 are

considered to represent small, medium, and large magnitudes of effect, respectively (Cohen, 1988). All mixed models were performed using the PROC MIXED procedure in SAS (Version 9.4).

3. Results

3.1. CORT and testosterone

Migrants had significantly lower baseline CORT levels compared with residents (**Fig. 2a**; mean $\pm$ SE, 0.2 ± 0.1 vs. 0.7 ± 0.2 ng/mL, $F_{1,42} = 4.87$, $p = 0.03$, Cohen's $f^2 = 0.09$). Baseline CORT levels were undetectable for 60% of migrants ($n = 30$) and 27% of residents ($n = 20$). However, baseline CORT did not significantly differ with sex ($F_{1,42} = 1.33$, $p = 0.25$) or age ($F_{1,42} = 2.15$, $p = 0.15$). Migrants also had significantly lower stress-induced CORT levels compared with residents (**Fig. 2b**; mean $\pm$ SE, 7.7 ± 1.0 vs. 20.6 ± 1.6 ng/mL, $F_{1,41} = 40.07$, $p < 0.01$, Cohen's $f^2 = 0.91$), and stress-induced CORT also did not significantly differ with sex ($F_{1,41} = 0.02$, $p = 0.88$) or age ($F_{1,41} = 3.39$, $p = 0.07$). Negative feedback efficacy did not significantly differ between migrant and resident (**Fig. 2c**; $F_{1,41} = 0.62$, $p = 0.44$), male and female ($F_{1,41} = 0.06$, $p = 0.44$), and adult young and adult old ($F_{1,41} = 0.00$, $p = 0.97$) birds. Resident males had significantly higher baseline testosterone levels than migrant males (**Fig. 3**; mean $\pm$ SE, 1.2 ± 0.4 vs. 0.1 ± 0.03 ng/mL, $F_{1,19} = 9.25$, $p < 0.01$, Cohen's $f^2 = 0.41$), although testosterone did not significantly differ with age ($F_{1,19} = 1.58$, $p = 0.22$). Baseline testosterone levels were undetectable for 18% of migrant males ($n = 11$) and 0% of resident males ($n = 11$).

3.2. Body condition

As expected based on established sub-species and sex differences in size, residents weighed significantly more than migrants (**Fig. 4**; $F_{1,78} = 27.07$, $p < 0.01$, Cohen's $f^2 = 0.33$), and males weighed significantly more than females (**Fig. 4**; mean $\pm$ SE, resident males: 22.7 ± 0.3 g, resident females: 22.3 ± 0.4 g, migrant males: 21.7 ± 0.5 g, migrant females: 20.3 ± 0.5 g, $F_{1,78} = 8.43$, $p < 0.01$, Cohen's $f^2 = 0.09$). Adult young and old birds did not significantly differ in weight ($F_{1,78} = 0.22$, $p = 0.64$). Compared with residents, migrants had significantly more abdominal and furcular fat (**Fig. 5a**; mean $\pm$ SE, 2.8 ± 0.4 vs. 1.2 ± 0.3 , abdominal fat: $F_{1,78} = 11.52$, $p < 0.01$, Cohen's $f^2 = 0.13$; **Fig. 5b**; mean $\pm$ SE, 2.9 ± 0.3 vs. 1.3 ± 0.3 , furcular fat: $F_{1,78} = 15.56$, $p < 0.01$, Cohen's $f^2 = 0.18$). Neither abdominal nor furcular fat differed significantly with sex (abdominal fat: $F_{1,78} = 0.45$, $p = 0.51$; furcular fat: $F_{1,78} = 1.84$, $p = 0.18$) or age (abdominal fat: $F_{1,78} = 0.03$, $p = 0.86$; furcular fat: ($F_{1,78} = 0.00$, $p = 0.96$).

4. Discussion

The purpose of this comparative study was to determine whether heightened HPA-axis activity during spring migratory development acts as a “brake” on the HPG-axis in birds preparing to migrate. Counter to expectation, we found that within an overwintering population of *J. hyemalis*, migrant individuals had less-reactive HPA-axes than resident individuals prior to migratory departure. Therefore, our results suggest that

the HPA-axis is not responsible for delayed reproductive maturation during spring migratory development.

Migrants were observed to have significantly greater fat stores compared with residents, thus suggesting that migratory development was well under way. Despite apparent evidence that these individuals were undergoing migratory preparation, their baseline and stress-induced CORT levels were lower than resident individuals. One possibility may be that we sampled migrant juncos too early during migratory development (2–4 wk before departure) to see heightened HPA-axis activity, although Holberton et al. (2008) found that captive Dark-eyed Juncos increased baseline CORT in concert with fat scores. Furthermore, a similar migrant vs. resident junco study in captivity found that prior to anticipated migratory departure, migrants had higher fat scores but lower baseline CORT concentrations compared to residents (Fudickar et al., in press). It is also possible that migrant and resident juncos have inherently different HPA-axes; future studies should measure winter CORT profiles of resident and migrant juncos to assess the relative increase in CORT from winter to pre-migration, which may be greater in migrants than residents. While this alternative is possible, it seems unlikely given how low baseline CORT levels already were in migrants.

Lower baseline and stress-induced CORT levels in migrants versus residents could also be caused by up-regulation of the HPA-axis prior to breeding (Eikenaar et al., 2015), as residents were at least a few weeks closer to breeding than migrants, and previous studies have found baseline CORT levels significantly increase from winter to pre-breeding (Hegner and Wingfield, 1990 but see Lattin et al., 2012). However, past studies have also shown that testosterone levels and testes size start to increase 1–2 mo

before CORT levels start to increase from winter levels (Hegner and Wingfield, 1990). Given that all juncos were still in their winter flocks and showed little signs of territorial behavior, we doubt that pre-breeding up-regulation of baseline and stress-induced CORT levels explain higher CORT levels in residents vs. migrants. Future studies should also better determine pre-migratory CORT profiles of migrant and resident female juncos. While our study found no significant CORT or negative feedback differences between sexes, our female sample size was very small ($n = 3$ and 6 for migrant and resident females, respectively).

Additionally, while migrant juncos did not have higher circulating levels of CORT or weaker negative feedback compared to resident juncos, downstream components of the migrant HPA-axis could have caused migrants to be more sensitive to stress. For example, altered concentrations of glucocorticoid receptors, corticosteroid binding globulin (CBG), and 11β -Hydroxysteroid dehydrogenase (11β -HSD) could all affect stress reactivity (Sapolsky et al., 2000). There is increasing evidence that seasonal changes in downstream components of the HPA-axis may contribute to increased stress sensitivity before breeding (Lattin et al., 2012; Lattin et al., 2015), thus helping birds optimally time breeding initiation.

Several other physiological mechanisms may mediate delayed reproductive development in migrants. One possibility is that migrants may require longer daylengths than residents to initiate reproductive development, as other studies have found population differences in photoperiodic thresholds (Lambrechts et al., 1997; Lambrechts et al., 1996; Silverin et al., 1993). An increased photoperiodic threshold in migrants could cause them to start gonadal recrudescence and reproductive development later than

residents, which is supported by our finding that resident males had significantly higher testosterone levels than migrant males during this study. Alternatively, migrants and residents may share the same photoperiodic threshold but differ in the rate of later-stage gonadal recrudescence and reproductive maturation (Caro et al., 2005; Caro et al., 2006). Later-stage reproductive development could also be contingent on the relative rate of increasing photoperiod, as migrants will experience rapidly increasing daylengths as they migrate toward higher latitudes. However, evidence suggests that gonadal recrudescence rates in birds are driven by current photoperiod, rather than the rate or change in direction of photoperiod (Dawson, 2015). Delayed reproductive development in migrants could also be mediated by gonadotropin-inhibiting hormone (GnIH) either through increased GnIH secretion or GnIH receptor concentrations (Kriegsfeld et al., 2015). Thyroid hormones could also play a role in delayed reproductive development, as thyroid hormone levels have been positively associated with fattening, flight muscle hypertrophy, and migratory restlessness (Bishop et al., 2000; Pant and Chandolasaklani, 1993). However, studies have reported both suppressive and permissive effects of the thyroid hormones on reproductive development in seasonally breeding birds (Dawson et al., 2001).

Reported baseline CORT levels in this study were surprisingly low, as many sample concentrations were less than 1 ng/mL. Only one other avian study has measured baseline CORT using this same kit (DetectX Corticosterone EIA Kit, Arbor Assays, Ann Arbor, MI, USA), but their lowest reported CORT concentration was 3.08 ng/mL (Pryor and Casto, 2015), which is more typical of other studies. However, our results may differ from Pryor and Casto (2015) because they used the kit-supplied dissociation reagent

whereas we did a double-extraction with diethyl ether. Therefore, lower-calculated
CORT concentrations may be specific to the EIA kit we used, as antibody and protocol
differences can significantly alter measured hormone concentrations (Brown et al., 2003;
Lattin et al., 2011), which is why different studies should not directly compare hormone
concentrations (Brown et al., 2003). Lower measures of CORT caused by kit differences
are also supported by the fact that our stress-induced CORT measurements are in the
lower range of other studies in wild birds. Regardless, these “lower” CORT
measurements should not matter since migrant/resident status, sex, and age were evenly
distributed among plates.

Contrary to previous studies, our results found that baseline CORT is not
elevated during migratory preparation since migrants had lower baseline CORT and
higher fat scores compared with residents. However, studies that have reported positive
relationships between baseline CORT and pre-migratory hyperphagia and fattening have
predominately been correlative (Fudickar et al., 2013; Holberton, 1999; Holberton et al.,
2008; Holberton et al., 1996; Long and Holberton, 2004), while the majority of
manipulative studies have found that neither CORT nor CORT-agonist administration
increases hyperphagia in birds (Astheimer et al., 1992; Gray et al., 1990; Holberton et al.,
2007; Saldanha et al., 2000; Simon, 1989 but see Landys et al., 2004b; Lohmus et al.,
2006). Furthermore, studies in Northern wheatears (*Oenanthe oenanthe*) during
migratory stopovers found negative correlations between baseline CORT levels and
refueling rates (Eikenaar et al., 2014; Eikenaar et al., 2013). Eikenaar et al. (2013)
suggested that since CORT levels rapidly decrease after arrival to stopover sites and then
steadily increase as birds put on more fat, CORT acts more as a departure signal. This

hypothesis fits well with similar studies in Bar-tailed Godwits (*Limosa lapponica*) (Landys-Ciannelli et al., 2002; Ramenofsky et al., 1995), Red Knots (*Calidris canutus*) (Piersma et al., 2000), and Gray Catbirds (*Dummetella carolinensis*) (Holberton et al., 1996), which also found that fatter birds had higher baseline CORT levels. Therefore, low baseline CORT levels may be necessary during hyperphagia and fuel deposition, while high baseline CORT levels may signal sufficient fat stores for departure. Future studies should determine whether *J.h. hyemalis* baseline CORT levels continue to rise during fattening and peak immediately before departure.

5. Conclusions

By comparing resident and migrant Dark-eyed Juncos that share the same overwintering site, this study examined whether up-regulated HPA-axis activity in migrants could potentially suppress later-stage reproductive development during spring migratory preparation. Contrary to our predictions, we found that migrants actually had lower circulating CORT levels and similar negative feedback efficacy compared with residents, thus suggesting that systemic glucocorticoids do not act as a brake on reproductive maturation prior to migratory departure. Future studies should examine the relationship between individual CORT levels and fueling rates during pre-migratory development, as our study design did not allow us to test how CORT changed over the migratory preparation stage. Future studies should also examine when residents start to increase CORT from winter to breeding levels, as it's possible we sampled residents too close to breeding. These studies would also verify whether migrants and residents have

inherently different HPA-axes. Regardless, our results strongly indicated that delayed reproductive development in migrants is not being driven by increased HPA-axis activity, therefore future studies should examine whether higher photoperiodic thresholds or other, physiological differences mediate delayed reproductive development in migrants. Finally, our data lead us to propose that high CORT levels may not be required for pre-migratory hyperphagia and fattening.

Acknowledgements

We thank the University of Virginia for use of their field site and facilities and J. Kittilson for assistance in the lab. We thank M. Abolins-Abols, J. Atwell, A. Fudickar, B. Heidinger, C. Lattin, K. Quinn, M. Ramenofsky, and L.M. Romero for their assistance and comments. We also thank C. Eikenaar and an anonymous reviewer for their comments on an earlier draft of this manuscript. This work was funded by NSF Grant IOS-1257527 (T.J.G) and IOS- 1257474 (E.D.K.). C.M.B. was funded by ND EPSCoR, a Wilson Ornithological Society Louis Agassiz Fuertes Grant, and an American Ornithologists' Union Hesse Research Award.

References

Astheimer, L.B., Buttemer, W.A., Wingfield, J.C., 1992. Interactions of Corticosterone with Feeding, Activity and Metabolism in Passerine Birds. *Ornis Scandinavica* 23, 355-365.

415 Bauchinger, U., Van't Hof, T., Biebach, H., 2007. Testicular development during long-
 416 distance spring migration. *Hormones and Behavior* 51, 295-305.

417 Bishop, C.M., McCabe, C.J., Gittoes, N.J.L., Butler, P.J., Franklyn, J.A., 2000. Tissue-
 418 specific regulation of thyroid hormone receptor mRNA isoforms and target gene
 419 proteins in domestic ducks. *Journal of Endocrinology* 165, 607-615.

420 Brown, J., Walker, S., Steinman, K., 2003. Endocrine manual for reproductive
 421 assessment of domestic and non-domestic species. Smithsonian National
 422 Zoological Park, Conservation and Research Center, Front Royal, Virginia.

423 Caro, S.P., Balthazart, J., Thomas, D.W., Lacroix, A., Chastel, O., Lambrechts, M.M.,
 424 2005. Endocrine correlates of the breeding asynchrony between two corsican
 425 populations of blue tits (*Parus caeruleus*). *General and Comparative*
 426 *Endocrinology* 140, 52-60.

427 Caro, S.P., Lambrechts, M.M., Chastel, O., Sharp, P.J., Thomas, D.W., Balthazart, J.,
 428 2006. Simultaneous pituitary-gonadal recrudescence in two Corsican populations
 429 of male blue tits with asynchronous breeding dates. *Hormones and Behavior* 50,
 430 347-360.

431 Clotfelter, E.D., O'Neal, D.M., Gaudioso, J.M., Casto, J.M., Parker-Renga, I.M., Snajdr,
 432 E.A., Duffy, D.L., Nolan, V., Ketterson, E.D., 2004. Consequences of elevating
 433 plasma testosterone in females of a socially monogamous songbird: evidence of
 434 constraints on male evolution? *Hormones and Behavior* 46, 171-178.

435 Cohen, J., 1988. *Statistical Power Analysis for the Behavioral Sciences*. Academic Press,
 436 Inc., New York.

437 Cornelius, J.M., Boswell, T., Jenni-Eiermann, S., Breuner, C.W., Ramenofsky, M., 2013.
 438 Contributions of endocrinology to the migration life history of birds. *General and*
 439 *Comparative Endocrinology* 190, 47-60.
 440 Dawson, A., 2015. Annual gonadal cycles in birds: Modeling the effects of photoperiod
 441 on seasonal changes in GnRH-1 secretion. *Frontiers in Neuroendocrinology* 37,
 442 52-64.
 443 Dawson, A., King, V.M., Bentley, G.E., Ball, G.F., 2001. Photoperiodic control of
 444 seasonality in birds. *Journal of Biological Rhythms* 16, 365-380.
 445 Deviche, P., 1995. Androgen regulation of avian premigratory hyperphagia and fattening-
 446 from eco-physiology to neuroendocrinology. *American Zoologist* 35, 234-245.
 447 DeVries, M.S., Jawor, J.M., 2013. Natural variation in circulating testosterone does not
 448 predict nestling provisioning rates in the northern cardinal, *Cardinalis cardinalis*.
 449 *Animal Behaviour* 85, 957-965.
 450 Dickens, M.J., Earle, K.A., Romero, L.M., 2009. Initial transference of wild birds to
 451 captivity alters stress physiology. *General and comparative endocrinology* 160,
 452 76-83.
 453 Dickens, M.J., Romero, L.M., 2013. A consensus endocrine profile for chronically
 454 stressed wild animals does not exist. *General and comparative endocrinology* 191,
 455 177-189.
 456 Eikenaar, C., Bairlein, F., Stowe, M., Jenni-Eiermann, S., 2014. Corticosterone, food
 457 intake and refueling in a long-distance migrant. *Hormones and Behavior* 65, 480-
 458 487.

459 Eikenaar, C., Frittsch, A., Bairlein, F., 2013. Corticosterone and migratory fueling in
 460 Northern wheatears facing different barrier crossings. *General and Comparative*
 461 *Endocrinology* 186, 181-186.
 462 Eikenaar, C., Müller, F., Klinner, T., Bairlein, F., 2015. Baseline corticosterone levels are
 463 higher in migrating than sedentary common blackbirds in autumn, but not in
 464 spring. *General and Comparative Endocrinology*.
 465 doi:10.1016/j.ygcen.2015.07.003
 466 Falsone, K., Jenni-Eiermann, S., Jenni, L., 2009. Corticosterone in migrating songbirds
 467 during endurance flight. *Hormones and behavior* 56, 548-556.
 468 Fudickar, A.M., Grieves, T.J., Atwell, J.W., Stricker, C., Ketterson, E.D., in press.
 469 Reproductive allochrony in seasonally sympatric populations maintained by
 470 differential response to photoperiod: implications for population divergence and
 471 response to climate change. *American Naturalist*.
 472 Fudickar, A.M., Schmidt, A., Hau, M., Quetting, M., Partecke, J., 2013. Female-biased
 473 obligate strategies in a partially migratory population. *Journal of Animal Ecology*
 474 82, 863-871.
 475 Gray, J.M., Yarian, D., Ramenofsky, M., 1990. Corticosterone, Foraging Behavior, and
 476 Metabolism in Dark-Eyed Juncos, *Junco-Hyemalis*. *General and comparative*
 477 *endocrinology* 79, 375-384.
 478 Grotjan, H.E., Keel, B.A., 1996. Data interpretation and quality control., in: Diamandis,
 479 E.P., Christopoulos, T.K. (Eds.), *Immunoassay*. Academic Press, New York, pp.
 480 51-93.

481 Gwinner, E., 1996a. Circadian and circannual programmes in avian migration. *Journal of*
 482 *Experimental Biology* 199, 39-48.
 483 Gwinner, E., 1996b. Circannual clocks in avian reproduction and migration. *Ibis* 138, 47-
 484 63.
 485 Hegner, R.E., Wingfield, J.C., 1990. Annual cycle of gonad size, reproductive hormones
 486 and breeding activity of free-living house sparrows (*P. domesticus*), in: Pinowski,
 487 J., Summer-Smith, J.D. (Eds.), *General Meetings of the Working Group on*
 488 *Granivorous Birds*, INTECOL. Warszawa: Polish Scientific Publishers, Ottawa,
 489 CA, pp. 123-135.
 490 Holberton, R.L., 1999. Changes in patterns of corticosterone secretion concurrent with
 491 migratory fattening in a neotropical migratory bird. *General and Comparative*
 492 *Endocrinology* 116, 49-58.
 493 Holberton, R.L., Boswell, T., Hunter, M.J., 2008. Circulating prolactin and corticosterone
 494 concentrations during the development of migratory condition in the Dark-eyed
 495 Junco, *Junco hyemalis*. *General and Comparative Endocrinology* 155, 641-649.
 496 Holberton, R.L., Parrish, J.D., Wingfield, J.C., 1996. Modulation of the adrenocortical
 497 stress response in neotropical migrants during autumn migration. *Auk* 113, 558-
 498 564.
 499 Holberton, R.L., Wilson, C.M., Hunter, M.J., Cash, W.B., Sims, C.G., 2007. The role of
 500 corticosterone in supporting migratory lipogenesis in the dark-eyed junco, *Junco*
 501 *hyemalis*: A model for central and peripheral regulation. *Physiological and*
 502 *Biochemical Zoology* 80, 125-137.
 503 Ketterson, E.D., 1979. Status Signaling in Dark-Eyed Juncos. *Auk* 96, 94-99.

504 Ketterson, E.D., Nolan, V., 1976. Geographic Variation and its Climatic Correlates in
 505 Sex-Ratio of Eastern-Wintering Dark-Eyed Juncos (*Junco-Hyemalis Hyemalis*).
 506 Ecology 57, 679-693.

507 Kriegsfeld, L.J., Ubuka, T., Bentley, G.E., Tsutsui, K., 2015. Seasonal control of
 508 gonadotropin-inhibitory hormone (GnIH) in birds and mammals. Frontiers in
 509 Neuroendocrinology 37, 65-75.

510 Lack, D., 1968. Bird Migration and Natural Selection. Oikos 19, 1-9.

511 Lambrechts, M.M., Blondel, J., Maistre, M., Perret, P., 1997. A single response
 512 mechanism is responsible for evolutionary adaptive variation in a bird's laying
 513 date. Proceedings of the National Academy of Sciences of the United States of
 514 America 94, 5153-5155.

515 Lambrechts, M.M., Perret, P., Blondel, J., 1996. Adaptive differences in the timing of egg
 516 laying between different populations of birds result from variation in
 517 photore sponsiveness. Proceedings of the Royal Society B-Biological Sciences
 518 263, 19-22.

519 Landys, M.M., Piersma, T., Ramenofsky, M., Wingfield, J.C., 2004a. Role of the low-
 520 affinity glucocorticoid receptor in the regulation of behavior and energy
 521 metabolism in the migratory red knot *Calidris canutus islandica*. Physiological
 522 and Biochemical Zoology 77, 658-668.

523 Landys, M.M., Ramenofsky, M., Guglielmo, C.G., Wingfield, J.C., 2004b. The low-
 524 affinity glucocorticoid receptor regulates feeding and lipid breakdown in the
 525 migratory Gambel's white-crowned sparrow *Zonotrichia leucophrys gambelii*.
 526 Journal of Experimental Biology 207, 143-154.

527 Landys, M.M., Ramenofsky, M., Wingfield, J.C., 2006. Actions of glucocorticoids at a
 528 seasonal baseline as compared to stress-related levels in the regulation of periodic
 529 life processes. *General and comparative endocrinology* 148, 132-149.

530 Landys-Ciannelli, M.M., Ramenofsky, M., Piersma, T., Jukema, J., Wingfield, J.C.,
 531 Castricum Ringing, G., 2002. Baseline and stress-induced plasma corticosterone
 532 during long-distance migration in the bar-tailed godwit, *Limosa lapponica*.
 533 *Physiological and Biochemical Zoology* 75, 101-110.

534 Lattin, C.R., Bauer, C.M., de Bruijn, R., Romero, L.M., 2012. Hypothalamus-pituitary-
 535 adrenal axis activity and the subsequent response to chronic stress differ
 536 depending upon life history stage. *General and comparative endocrinology* 178,
 537 494-501.

538 Lattin, C.R., Breuner, C.W., Romero, L.M., 2015. Does corticosterone regulate the onset
 539 of breeding in free-living birds?: The CORT-Flexibility Hypothesis and six
 540 potential mechanisms for priming corticosteroid function. *Hormones and*
 541 *Behavior*. doi:10.1016/j.ygcen.2015.07.003

542 Lattin, C.R., Reed, J.M., DesRochers, D.W., Romero, L.M., 2011. Elevated
 543 corticosterone in feathers correlates with corticosterone-induced decreased feather
 544 quality: a validation study. *Journal of Avian Biology* 42, 247-252.

545 Lohmus, M., Sundstrom, F., Moore, F.R., 2006. Non-invasive corticosterone treatment
 546 changes foraging intensity in red-eyed vireos *Vireo olivaceus*. *Journal of Avian*
 547 *Biology* 37, 523-526.

548 Long, J.A., Holberton, R.L., 2004. Corticosterone secretion, energetic condition, and a
 549 test of the migration modulation hypothesis in the hermit thrush (*Catharus*
 550 *guttatus*), a short-distance migrant. *Auk* 121, 1094-1102.

551 Mattocks, P.W., Jr., 1976. The role of gonadal hormones in the regulation of the
 552 premigratory fat deposition in the white-crowned sparrow, *Zonotrichia leucophrys*
 553 *gambelii*. University of Washington, Seattle.

554 Nolan, V., Jr., Ketterson, E.D., Cristol, D.A., Rogers, C.M., Clotfelter, E.D., Titus, R.C.,
 555 Schoech, S.J., Snajdr, E., 2002. Dark-eyed junco: *Junco hyemalis*. *Birds of North*
 556 *America* 716, 1-44.

557 Nolan, V., Ketterson, E.D., 1983. An analysis of body-mass, wing length, and visible fat
 558 deposits of Dark-eyed Juncos wintering at different latitudes. *Wilson Bulletin* 95,
 559 603-620.

560 O'Neal, D.M., Kiley, R.P., Ketterson, E.D., 2011. The effect of winter sex ratio on
 561 immune function and condition in a differential migrant. *Physiology & Behavior*
 562 102, 406-413.

563 Owen, J.C., Garvin, M.C., Moore, F.R., 2014. Elevated testosterone advances onset of
 564 migratory restlessness in a nearctic-neotropical landbird. *Behavioral Ecology and*
 565 *Sociobiology* 68, 561-569.

566 Pant, K., Chandolasaklani, A., 1993. A role for thyroid-hormones in the development of
 567 premigratory disposition in redheaded bunting, *Emberiza bruniceps*. *Journal of*
 568 *Comparative Physiology B-Biochemical Systemic and Environmental Physiology*
 569 163, 389-394.

570 Perrins, C.M., 1970. Timing of Birds Breeding Seasons. *Ibis* 112, 242-&.

571 Piersma, T., Gudmundsson, G.A., Lilliendahl, K., 1999. Rapid changes in the size of
 572 different functional organ and muscle groups during refueling in a long-distance
 573 migrating shorebird. *Physiological and Biochemical Zoology* 72, 405-415.
 574 Piersma, T., Reneerkens, J., Ramenofsky, M., 2000. Baseline corticosterone peaks in
 575 shorebirds with maximal energy stores for migration: A general preparatory
 576 mechanism for rapid behavioral and metabolic transitions? *General and*
 577 *Comparative Endocrinology* 120, 118-126.
 578 Pryor, L.J.E., Casto, J.M., 2015. Blood-feeding ectoparasites as developmental stressors:
 579 Does corticosterone mediate effects of mite infestation on nestling growth,
 580 immunity and energy availability? *Journal of Experimental Zoology Part A:*
 581 *Ecological Genetics and Physiology* 323, 466-477.
 582 Ramenofsky, M., 1990. Fat storage and fat metabolism in relation to migration., in:
 583 Gwinner, E. (Ed.), *Bird Migration*. Springer-Verlag, Berlin, Germany, pp. 214-
 584 231.
 585 Ramenofsky, M., 2011. Hormones in Migration and Reproductive Cycles of Birds, in:
 586 Norris, D.O., Lopez, K.H. (Eds.), *Hormones and Reproduction of Vertebrates*,
 587 Vol 4: Birds. Elsevier Academic Press Inc, San Diego, pp. 205-237.
 588 Ramenofsky, M., Nemeth, Z., 2014. Regulatory mechanisms for the development of the
 589 migratory phenotype: Roles for photoperiod and the gonad. *Hormones and*
 590 *Behavior* 66, 148-158.
 591 Ramenofsky, M., Piersma, T., Jukema, J., 1995. Plasma-corticosterone in Bar-tailed
 592 Godwits at a major stop-over site during spring migration. *Condor* 97, 580-585.
 593 Ramenofsky, M., Wingfield, J.C., 2007. Regulation of migration. *Bioscience* 57, 135-143.

594 Rich, E.L., Romero, L.M., 2001. Daily and photoperiod variations of basal and stress-
 595 induced corticosterone concentrations in house sparrows (*Passer domesticus*).
 596 Journal of Comparative Physiology B-Biochemical Systemic and Environmental
 597 Physiology 171, 543-547.

598 Romero, L.M., 2002. Seasonal changes in plasma glucocorticoid concentrations in free-
 599 living vertebrates. General and comparative endocrinology 128, 1-24.

600 Romero, L.M., 2004. Physiological stress in ecology: lessons from biomedical research.
 601 Trends in Ecology & Evolution 19, 249-255.

602 Romero, L.M., 2006. Seasonal changes in hypothalamic-pituitary-adrenal axis sensitivity
 603 in free-living house sparrows (*Passer domesticus*). General and comparative
 604 endocrinology 149, 66-71.

605 Romero, L.M., Reed, J.M., 2005. Collecting baseline corticosterone samples in the field:
 606 is under 3 min good enough? Comparative Biochemistry and Physiology A-
 607 Molecular & Integrative Physiology 140, 73-79.

608 Romero, L.M., Remage-Healey, L., 2000. Daily and seasonal variation in response to
 609 stress in captive starlings (*Sturnus vulgaris*): Corticosterone. General and
 610 comparative endocrinology 119, 52-59.

611 Rosvall, K.A., Reichard, D.G., Ferguson, S.M., Whittaker, D.J., Ketterson, E.D., 2012.
 612 Robust behavioral effects of song playback in the absence of testosterone or
 613 corticosterone release. Hormones and Behavior 62, 418-425.

614 Rowan, W., 1926. On photoperiodism, reproductive periodicity, and the annual
 615 migrations of birds and certain fishes. Proc Boston Soc Nat Hist 38, 147-189.

616 Saldanha, C.J., Schlinger, B.A., Clayton, N.S., 2000. Rapid effects of corticosterone on
617 cache recovery in mountain chickadees (*Parus gambeli*). *Hormones and Behavior*
618 37, 109-115.

619 Sapolsky, R.M., Altmann, J., 1991. Incidence of Hypercortisolism and Dexamethasone
620 Resistance Increases with Age among Wild Baboons. *Biological psychiatry* 30,
621 1008-1016.

622 Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence
623 stress responses? Integrating permissive, suppressive, stimulatory, and preparative
624 actions. *Endocrine reviews* 21, 55-89.

625 Schoech, S.J., Ketterson, E.D., Nolan, V., 1999. Exogenous testosterone and the
626 adrenocortical response in Dark-eyed Juncos. *Auk* 116, 64-72.

627 Selya, A.S., Rose, J.S., Dierker, L.C., Hedeker, D., Mermelstein, R.J., 2012. A practical
628 guide to calculating Cohen's $f(2)$, a measure of local effect size, from PROC
629 MIXED. *Frontiers in Psychology* 3, 6.

630 Silverin, B., Massa, R., Stokkan, K.A., 1993. Photoperiodic adaptation to breeding at
631 different latitudes in Great Tits. *General and Comparative Endocrinology* 90, 14-
632 22.

633 Simon, J., 1989. Chicken as a useful species for the comprehension of insulin action.
634 *Critical Reviews in Poultry Biology* 2, 121-148.

635 Tonra, C.M., Marra, P.P., Holberton, R.L., 2011. Early elevation of testosterone advances
636 migratory preparation in a songbird. *Journal of Experimental Biology* 214, 2761-
637 2767.

638 Williams, T.D., 2012. Physiological Adaptions for Breeding in Birds. Princeton
639 University Press, Princeton, NJ.
640 Wingfield, J.C., Sapolsky, R.M., 2003. Reproduction and resistance to stress: When and
641 how. Journal of neuroendocrinology 15, 711-724.
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660

Figure Captions

Figure 1. Mean $\pm$ SE negative feedback efficacy 60 and 90 min after dexamethasone (DEX) ($n = 8$) or PBS ($n = 8$) injection in migratory Dark-eyed Juncos (*Junco hyemalis* captured during autumn in North Dakota, USA (45°53'N, 96°47'W). Negative feedback efficacy was calculated as the percent decrease from CORT levels 30 min after capture (stress-induced levels) to those 60 and 90 min after injection with DEX or PBS. Asterisks indicate significant differences ($p < 0.05$) between DEX and PBS-injected birds.

Figure 2. Mean $\pm$ SE **a)** baseline corticosterone (CORT) ($n = 20$ and 30), **b)** stress-induced CORT ($n = 18$ and 31), and **c)** negative feedback efficacy levels ($n = 18$ and 31) measured in Dark-eyed Junco residents (*Junco hyemalis carolinensis*) and migrants (*J.h. hyemalis*). Birds were captured during spring in VA, USA (37°22'N, 80°31'W). Negative feedback efficacy was calculated as the percent decrease from stress-induced CORT levels to those 90 min after dexamethasone (DEX) injection. Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 3. Mean $\pm$ SE baseline plasma testosterone concentrations of male Dark-eyed Junco residents (*Junco hyemalis carolinensis*) ($n = 11$) and migrants (*J.h. hyemalis*) ($n = 11$), respectively. Birds were captured during spring in VA, USA (37°22'N, 80°31'W). Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 4. Mean $\pm$ SE body weight of male and female Dark-eyed Junco residents (*Junco hyemalis carolinensis*) ($n = 32$ and 9) and migrants (*J.h. hyemalis*) ($n = 29$ and 12), respectively. Birds were captured during spring in VA, USA ($37^{\circ}22'N$, $80^{\circ}31'W$). Males weighed significantly more than females. Migrants are known to be smaller-bodied than residents (Nolan et al., 2002). Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 5. Mean $\pm$ SE **a)** abdominal and **b)** furcular fat scores of Dark-eyed Junco residents (*Junco hyemalis carolinensis*) ($n = 41$) and migrants (*J.h. hyemalis*) ($n = 41$). Fats scores ranged from 0 (no visible fat) to 5 (visible fat bulging). Birds were captured during spring in VA, USA ($37^{\circ}22'N$, $80^{\circ}31'W$). Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 1. Mean $\pm$ SE negative feedback efficacy 60 and 90 min after dexamethasone (DEX) ($n = 8$) or PBS ($n = 8$) injection in migratory Dark-eyed Juncos (*Junco hyemalis hyemalis*) captured during autumn in North Dakota, USA (45°53'N, 96°47'W). Negative feedback efficacy was calculated as the percent decrease from CORT levels 30 min after capture (stress-induced levels) to those 60 and 90 min after injection with DEX or PBS. Asterisks indicate significant differences ($p < 0.05$) between DEX and PBS-injected birds.

Figure 2. Mean $\pm$ SE **a)** baseline corticosterone (CORT) ($n = 20$ and 30), **b)** stress-induced CORT ($n = 18$ and 31), and **c)** negative feedback efficacy levels ($n = 18$ and 31) measured in Dark-eyed Junco residents (*Junco hyemalis carolinensis*) and migrants (*J.h. hyemalis*). Birds were captured during spring in VA, USA ($37^{\circ}22'N$, $80^{\circ}31'W$). Negative feedback efficacy was calculated as the percent decrease from stress-induced CORT levels to those 90 min after dexamethasone (DEX) injection. Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 3. Mean $\pm$ SE baseline plasma testosterone concentrations of male Dark-eyed Junco residents (*Junco hyemalis carolinensis*) ($n = 11$) and migrants (*J.h. hyemalis*) ($n = 11$), respectively. Birds were captured during spring in VA, USA (37°22'N, 80°31'W). Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 4. Mean $\pm$ SE body weight of male and female Dark-eyed Junco residents (*Junco hyemalis carolinensis*) ($n = 32$ and 9) and migrants (*J.h. hyemalis*) ($n = 29$ and 12), respectively. Birds were captured during spring in VA, USA ($37^{\circ}22'N$, $80^{\circ}31'W$). Males weighed significantly more than females. Migrants are known to be smaller-bodied than residents (Nolan et al., 2002). Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.

Figure 5. Mean $\pm$ SE **a)** abdominal and **b)** furcular fat scores of Dark-eyed Junco residents (*Junco hyemalis carolinensis*) ($n = 41$) and migrants (*J.h. hyemalis*) ($n = 41$). Fats scores ranged from 0 (no visible fat) to 5 (visible fat bulging). Birds were captured during spring in VA, USA (37°22'N, 80°31'W). Asterisks indicate significant differences ($p < 0.05$) between residents and migrants.