

Sex steroid profiles in zebra finches: effects of reproductive state and domestication

Nora H. Prior^{1*}, Kang Nian Yap², Mark C. Mainwaring⁵, Hans H. Adomat⁶, Ondi L. Crino^{5,7}, Chunqi Ma², Emma S. Guns⁶, Simon C. Griffith⁵, Katherine L. Buchanan⁷⁺, Kiran K. Soma^{1,2,3,4+}

Abstract

The zebra finch is a common model organism in neuroscience, endocrinology, and ethology. Zebra finches are generally considered opportunistic breeders, but the extent of their opportunism depends on the predictability of their habitat. This plasticity in the timing of breeding raises the question of how domestication, a process that increases environmental predictability, has affected their reproductive physiology. Here, we compared circulating steroid levels in various “strains” of zebra finches. In Study 1, using radioimmunoassay, we examined circulating testosterone levels in several strains of zebra finches (males and females). Subjects were wild or captive (Captive Wild-Caught, Wild-Derived, or Domesticated). In Study 2, using liquid chromatography-tandem mass spectrometry (LC-MS/MS), we examined circulating sex steroid profiles in wild and domesticated zebra finches (males and females). In Study 1, circulating testosterone levels in males differed across strains. In Study 2, six steroids were detectable in plasma from wild zebra finches (pregnenolone, progesterone, dehydroepiandrosterone (DHEA), testosterone, androsterone, and 5 α -dihydrotestosterone (5 α -DHT)). Only pregnenolone and progesterone levels changed across reproductive states in wild finches. Compared to wild zebra finches, domesticated zebra finches had elevated levels of circulating pregnenolone, progesterone, DHEA, testosterone, androstenedione, and androsterone. These data suggest that domestication has profoundly altered the endocrinology of this common model organism. These results have implications for interpreting studies of domesticated zebra finches, as well as studies of other domesticated species.

Keywords: Australia, domestic, opportunistic breeding, songbird, steroid profiling

1. Introduction

Natural selection shapes both the behavior of wild animals and its underlying physiological mechanisms. The process of domestication alters natural selection and introduces artificial selection pressures. Some effects of domestication are similar across species (Zeder, 2012; Wilkins, et al., 2014). For example, domesticated animals show changes to the seasonal regulation of reproduction and molt (Belyaev, 1979; Lincoln et al., 1990; Setchell, 1992). Furthermore, domesticated animals show decreased hypothalamic-pituitary-adrenal (HPA) axis reactivity (Martin, 1978; Woodward and Strange, 1987; Künzl and Sascher, 1999; Albert et al., 2008; Suzuki et al., 2012; Homberger et al., 2013; Ericsson et al., 2014). However, other effects of domestication are different across species and depend on the characteristics of the original population and the specifics of the domestication process (Zeder, 2012; Adkins-Regan, 2009).

Zebra finches are an extensively used model organism in a wide range of behavioral, evolutionary, and neuroscience research (Griffith and Buchanan, 2010). Initial breeding stocks of zebra finches were removed from the wild in Australia during the 19th century and were established into a number of independent breeding populations across the globe (Zann, 1996). Given the ease with which they can be studied in the laboratory, it is not surprising that most of the research on the physiology and neuroscience of this species has been conducted on domesticated zebra finches (Griffith and Buchanan, 2010). The laboratory environment itself may have caused persistent differences to arise between wild and domesticated individuals with respect to both their physiology and behavior (Calisi and Bentley, 2009; Dickens and Bentley, 2014). Furthermore, the process of domestication may have led to genetic differentiation

between wild and domesticated zebra finch populations, even in the few hundred generations for which they have been separated (Forstmeier et al., 2007). It is currently unclear to what extent there may be population differences in functional traits relating to behavior and physiology, between wild and domesticated finches and also across different domesticated populations. The demonstration of such effects will provide insight into the mechanisms underlying domestication, but may also necessitate caution when comparing across populations of zebra finches that differ in their domestication history.

For some traits, wild and domesticated zebra finches are extremely similar. For example, both wild and domesticated zebra finches form long-term pair bonds, provide biparental care, and breed opportunistically (Zann, 1996). With regard to morphology, zebra finches with wild-type plumage are common in domesticated populations, and beak coloration is similar in wild and domesticated zebra finches (Burley et al., 1992). However, other traits differ between wild and domesticated zebra finches. Domesticated zebra finches are generally larger and express many different plumage colorations (Sossinka, 1970; Carr and Zann, 1986; Zann, 1996; Forstmeier et al., 2007). Furthermore, wild zebra finches have lower rates of extra-pair copulations (Griffith et al., 2010) and differ subtly in nesting behavior (Mainwaring et al., 2010; Gilby et al., 2013). In mate choice tests, captive wild-caught females are choosier than domesticated females, and domesticated males are more active courters than wild-caught males (Rutstein et al., 2007). However, it is currently unclear whether there are fundamental differences between wild and domesticated zebra finches in their endocrine profiles.

Zebra finches are opportunistic breeders; however, the extent of opportunism varies with habitat and environmental conditions (Zann, 1996; Perfito et al., 2007). Overall, the regulation of breeding in zebra finches is not as well understood as it is in seasonally-breeding species. However, studies have suggested that the HPG axis functions similarly in wild and domesticated zebra finches (Perfito et al., 2007; 2011; Prior and Soma, 2015). For example, the effects of natural droughts on wild finches and the effects of experimental water restriction on domesticated finches are similar (Perfito, 2010; Prior and Soma, 2015). Both cause zebra finches to stop breeding and inhibit parts of the HPG axis (Prior et al., 2013). Therefore, it is likely that (1) wild zebra finches have sex steroid profiles that change across a breeding cycle; and that (2) domestication, a process that reduces environmental variability, has affected sex steroid profiles.

We began to [investigate the above hypotheses](#) in two studies. In Study 1, we used a radioimmunoassay (RIA) to quantify circulating testosterone levels in males and females from four “strains” of zebra finches: Wild, Captive Wild-Caught, Wild-Derived (males only), and Domesticated. In Study 2, we used liquid chromatography-tandem mass spectrometry (LC-MS/MS) to quantify multiple sex steroids and precursors in wild and domesticated zebra finches. We also [described circulating steroid profiles of wild zebra finches at different reproductive states across](#) a breeding cycle.

2. Materials and Methods

2.1 Ethics statement

These studies were carried out under a UBC Animal Care Committee protocol and under ethics licenses from the Animal Ethics Committee at Deakin University (B24- 2012) and

Macquarie University (ARA 2010/053). Experiments followed the guidelines of the Canadian Council on Animal Care and the Australian and New Zealand Council for the Care of Animals in Research and Teaching.

Study 1: Radioimmunoassay (RIA) to measure testosterone levels in plasma

We used a radioimmunoassay to compare circulating testosterone levels among four strains of zebra finches (1) Wild (n = 20 females and 11 males), (2) Captive Wild-Caught (n = 4 females and 4 males), (3) Wild Derived (n = 0 females and 42 males), and (4) Domesticated (n = 15 females and 15 males).

Wild finches were sampled while “not nesting” or while actively nesting (incubating or feeding chicks) (females: n = 8 Not Nesting and 12 Nesting; males: n = 5 Not Nesting and 6 Nesting). Subjects in other strains were not nesting at the time of blood sampling, allowing us to compare circulating testosterone levels across all four strains, in subjects that were not nesting. Wild-Derived males were housed in same-sex aviaries, and all other subjects were paired. See below for more details.

2.3 Subjects

2.3.1 Wild zebra finches

Free-roaming, wild zebra finch pairs were captured for blood sampling at Fowler’s Gap Arid Zone Research Station, in the Western region of New South Wales, Australia. More specifically, pairs were from nest box colonies within Gap Hills (Griffith et al., 2008). Blood samples were collected between September-November 2012. At this location, October is often a peak breeding month for zebra finches (Griffith et al,

2008), and in this season of 2012, there was continuous breeding activity in the study area throughout the time of sampling. However, in other areas of the Fowler's Gap study site, environmental conditions were different and there was no breeding.

At the time of blood sampling, subjects were either not nesting or nesting. Breeding activity in nest box populations was assessed every 1-3 days. For active nests, the dates that eggs were laid and hatched were recorded, allowing us to determine the days of incubation and feeding chicks. Breeding females and males (either incubating or feeding chicks) were caught in the nest box during the early morning. Subjects that were not nesting were caught in a walk-in feeder trap from an area where no birds were nesting in nest boxes (Mariette et al., 2011; McCowan et al., 2015). While there was no nesting at these sites, because the overall environmental conditions were appropriate for breeding, it is difficult to be certain that these individuals were not maintaining a breeding nest elsewhere. Therefore, we classified these individuals as not nesting, instead of non-breeding, because they may have been physiologically capable of breeding (Prior et al., 2013; Perfito et al., 2007). Note that in zebra finches, only females develop a brood patch (Zann, 1996). Additionally, all individuals examined in these colonies had a brood patch score of 1-2, making it difficult to identify breeding state from brood patch score.

2.3.2 Captive Wild-Caught zebra finches

Wild adult female and male zebra finches were caught at the Fowler's Gap study site (in September 2012) and housed in open-air aviaries under natural photoperiod for 1-2 months prior to collecting blood samples. Subjects had access to *ad libitum* seed, water, cuttlefish bone, and grit. These subjects were housed in several aviaries, and each

aviary held 1-2 pairs. All subjects were pair bonded; however, they were not nesting at the time of blood sampling (Prior et al., 2016).

2.3.3 *Wild-Derived zebra finches*

Wild-Derived adult male zebra finches were also examined. These subjects were largely first- or second-generation and born in captivity at Macquarie University. Some of these birds might be the original individuals taken from the wild in either 2007 or 2009 (from Fowler's Gap and another population in Far-Western New South Wales). These subjects were housed in a same-sex open-air aviary (8 X 10 X 2 m). All individuals had *ad libitum* access to seed, water, cuttlefish bone, and grit. Blood was collected in the morning. Prior to blood collection, individuals were housed in the aviary for one month without contact with females. Therefore, these males were not nesting.

2.3.4 *Domesticated zebra finches*

Adult zebra finches (>120 days old) housed at University of British Columbia (UBC) were used as the domesticated sample. This colony was derived from birds from Eastern Birds Supplies (Quebec, Canada) and birds from Simon Fraser University (Burnaby, Canada) (Forstmeier et al. 2007). At UBC, subjects were housed indoors at ~22 °C, ~35% relative humidity, and on a 14:10 h light:dark cycle. All individuals had *ad libitum* access to seed (50/50, panicum millet/white millet, Just For Birds, Langley BC), water, cuttlefish bone, and grit. These birds were housed in pairs, which were not nesting.

2.4 Blood collection

We collected 100 to 150 μ L of blood from the brachial vein. Time to bleed after capture by Strain: Wild = within 15 min (7.3 ± 0.3 min); Captive Wild-Caught = within 13 min (6.9 ± 0.4 min); Wild Derived = within 7 min (3.1 ± 0.2 min); Domesticated = within 10 min (4.0 ± 0.1 min). For Wild zebra finches, blood samples were kept on wet ice until centrifugation to obtain plasma. For all other strains, blood samples were immediately centrifuged. All plasma samples were frozen until radioimmunoassay.

2.5 RIA to measure plasma testosterone levels

Steroids were extracted from plasma samples (10 – 20 μ L) using solid phase extraction (Newman et al., 2008; Taves et al., 2010; 2011). Testosterone was quantified using a sensitive and specific radioimmunoassay (RIA) (MP Biomedicals, 07189102). Validations and modifications of this RIA have been described (Overk et al., 2013). Samples were analyzed in 4 assays. Intra-assay variation was 10.76% and inter-assay variation was 12.75%. Samples were corrected for recovery of testosterone from zebra finch plasma (102.5%).

2.6 Study 2: Liquid chromatography-tandem mass spectrometry (LC-MS/MS) to measure sex steroid profiles in plasma

We used a sensitive and specific liquid chromatography-tandem mass spectrometry (LC-MS/MS) method to quantify circulating steroids. Wild zebra finches samples were collected as described above during September-November 2012. Here samples were categorized into 3 breeding stages: Not Nesting (n = 11 females and 10

males), Incubating (n = 17 females and 7 males), and Feeding Chicks (n = 9 females and 7 males). Some of these individuals are the same as the wild zebra finches in Study 1. In addition, steroid profiles were compared between wild zebra finches (n = 37 females and 24 males) and domesticated zebra finches (n = 8 females and 5 males). Again, UBC domesticated zebra finches were sampled. These are different individuals than those sampled for Study 1, although all birds were housed under the same conditions (see above). For this comparison, both wild and domesticated finches were not nesting. Blood sampling was the same as above.

2.7 LC-MS/MS to measure plasma steroid profiles

Steroid extraction and LC-MS/MS protocols were adapted from Kalhorn et al., 2007. This LC-MS/MS assay is optimized to measure 11 androgens and progestins: 6 steroids from the “traditional” pathway to 5 α -dihydrotestosterone (5 α -DHT) [pregnenolone; progesterone; 17-OH-pregnenolone; dehydroepiandrosterone (DHEA); androstenedione and testosterone], 4 steroids from the “backdoor” pathway to 5 α -DHT [pregnan-3,20-dione; pregnan-3 α ,ol-20-one; pregnan-3,17-diol-20-one and androsterone], and 5 α -DHT (Auchus, 2004; Fokidis et al., 2015; Prior et al., 2016). Internal standards (deuterated testosterone, 5 α -DHT and androstanediol, C/D/N Isotopes Inc, Quebec Canada) were added to plasma samples (~30 μ l). Steroids were then extracted using 2000 μ l of 60:40 (v/v) hexane:ethyl acetate for 30 min, and the upper layer was collected. Next, two additional extractions were done and pooled with the initial extract to maximize extraction efficiency. Pooled extracts were dried in a centrifugal vacuum evaporator (Centrivap, Labconco).

The resulting residues were derivatized using hydroxylamine (HA, Fluka, Sigma, Oakville Ontario Canada), which enhances sensitivity for ketosteroids. Extracts were dissolved in 50 μ L of 50 mM HA in 50% methanol, centrifuged at 20,000g for 2 min, transferred to an LC insert, and then heated to 65°C for 30 min prior to LC-MS/MS analysis.

Analysis was carried out with the analytical system of a Waters Acquity UPLC Separations Module coupled to a Waters Quattro Premier XE Tandem Mass Spectrometer (Waters Corporation, Massachusetts, USA). A 2.1x100mm BEH 1.7 μ M C18 column was used (Water Corporation, Massachusetts, USA). The two mobile phases were water and acetonitrile, both containing 0.1% formic acid, using the following gradient: 0 min, 25%; 0.2 min, 25%; 8 min, 70%; 9 min, 100%; 10 min, 100%; 10.2 min, 25% (% acetonitrile) with a total run length of 12 min. Column temperature was 35 °C and injection volume was 15 μ L. The MS was set at unit resolution, capillary was 3 kV, source and desolvation temperatures were 120 °C and 300 °C respectively, desolvation and cone gas flows were 1000 L/h and 50 L/h and the collision cell pressure was held at 4.6×10^{-3} mbar. All data were collected in electrospray (positive mode) (ES+) by multiple reaction monitoring (MRM) with instrument parameters optimized for the m/z ratios and corresponding fragments of the oxime-steroids monitored for each MRM.

Data processing was conducted with Quanlynx (Waters Corporation Massachusetts, USA) using area under curve of analyte/IS. Calibration samples consisted of neat standards and equivalently extracted spiked charcoal-stripped serum (6 standards ranging from 0.02 to 10ng/ml) and normalized to sample volume. The limit of detection (LOD) ranged from approximately 0.01-0.02 ng/mL (3x background). Recoveries and

conversions to derivatized steroid species were greater than 80% for each steroid.

Statistics were conducted for steroids where some samples were above the LOD; for

these steroids, all samples below the LOD were set to zero.

2.8 Statistics

All statistics were conducted in R v3.1.2 (R core team, 2014). Time to bleed did not significantly impact circulating steroid levels, and thus no such correction was made in the models. ANOVAs were run using glm models, and data were transformed as necessary to meet the assumptions of the test. All models are reported as ANOVA summaries with the resulting F statistic.

2.8.1 Study 1: RIA to measure testosterone levels in plasma

In order to test whether Sex or Nesting Activity (Not Nesting vs. Nesting) affected circulating testosterone levels of Wild zebra finches, we used an ANOVA with Sex and Nesting Activity as between-subject factors.

In zebra finches that were not nesting, in order to test whether Strain (Wild, Captive Wild-Caught, Wild-Derived, and Domesticated) affected circulating testosterone levels, we used a non-parametric one-way ANOVA (Kruskal-Wallis test), with Strain as a between-subject factor. Note that no Wild-Derived females were sampled. Separate Kruskal-Wallis tests were conducted on females and males. Pair-wise comparisons were conducted using Wilcoxon rank sum tests.

2.8.2 Study 2: LC-MS/MS to measure sex steroid profiles in plasma

In order to test whether sex steroid levels varied with Reproductive State in wild finches, we conducted an ANOVA for each steroid. Sex and Reproductive State (Not Nesting, Incubating, or Feeding Chicks) were between-subject factors. Significant main effects of Reproductive State were followed up with pair-wise Wilcoxon rank sum tests.

In zebra finches that were not nesting, in order to test whether sex steroids differed between wild and domesticated finches, we conducted an ANOVA for each steroid. Sex and Strain (Wild or Domesticated) were both between-subject factors. Significant interactions were followed up Wilcoxon rank sum tests (Strain) for females and males separately.

3. Results

3.1 Study 1: RIA to measure testosterone levels in plasma

In wild zebra finches, there was no effect of Nesting Activity (Not Nesting vs. Nesting) on female or male circulating testosterone levels (Figure 1A; $F_{1,28} = 0.24$, $P = 0.63$). There was also no Sex \times Nesting Activity interaction (Figure 1A; $F_{1,27} = 0.35$, $P = 0.56$). However, there was a main effect of Sex, with males having significantly higher circulating testosterone levels than females (Figure 1A; $F_{1,29} = 8.30$, $P < 0.01$).

We then focused on subjects that were not nesting in 4 “strains” (Wild, Captive Wild-Caught, Wild Derived, and Domesticated). The effect of Strain was examined in females and males separately. Overall, circulating testosterone varied significantly by Strain in females (Figure 1B; Kruskal-Wallis, $\chi^2 = 15.22$, $P < 0.001$) and males (Figure 1C; $\chi^2 = 30.71$, $P < 0.001$). In females, Wild females had higher circulating testosterone

than Captive Wild-Caught females and Domesticated females (Figure 1B; $P \leq 0.01$ in both cases). In males, Wild and Domesticated males had higher circulating testosterone than Wild-Derived males (Figure 1C; $P < 0.001$ in both cases).

3.2 Study 2: LC-MS/MS to measure sex steroid profiles in plasma

Six steroids were detectable in the circulation of wild zebra finches: pregnenolone, progesterone, DHEA, testosterone, androsterone, and 5α -DHT (Figure 2; Table 1). However, androsterone and 5α -DHT were only detectable in a few plasma samples (Table 1). Androstenedione was not detectable in plasma samples from wild zebra finches.

Levels of pregnenolone, progesterone, and DHEA were not significantly different between females and males (pregnenolone: $F_{1,57} = 2.58$, $P = 0.11$; progesterone: $F_{1,57} = 0.16$, $P = 0.69$; DHEA: $F_{1,57} = 2.63$, $P = 0.11$; Figure 3A-C). Testosterone levels were significantly higher in males than females ($F_{1,57} = 8.31$, $P < 0.01$; Figure 3D).

Pregnenolone and progesterone levels differed across Reproductive State (pregnenolone: $F_{2,58} = 4.59$, $P = 0.01$; progesterone: $F_{2,58} = 10.67$, $P < 0.001$; Figure 3A,B). Pregnenolone levels were highest during Feeding Chicks (Figure 3A; Not Nesting vs. Feeding Chicks, $P = 0.03$; Incubating vs. Feeding Chicks, $P = 0.04$; Not Nesting vs. Incubating, $P = 0.94$). Similarly, progesterone levels were highest during Feeding Chicks and lowest during Not Nesting (Figure 3B; Not Nesting vs. Feeding Chicks, $P < 0.001$; Incubating vs. Feeding Chicks, $P < 0.01$; Not Nesting vs. Incubating, $P < 0.01$). Circulating DHEA levels (Figure 3C) and testosterone levels (Figure 3D) were not affected by Reproductive State in males or females (DHEA: $F_{2,58} = 1.77$, $P = 0.18$;

testosterone: $F_{2,58} = 0.93$, $P = 0.40$). There were no significant Sex \times Reproductive State interactions (all P values > 0.05). Because the residuals of the GLM model for testosterone were not normally distributed, we confirmed there was no effect of Reproductive State using Kruskal-Wallis tests (Figure 3D; Females: $\chi^2 = 1.39$, $P = 0.50$. Males: $\chi^2 = 0.65$, $P = 0.72$).

Independent of reproductive state, domesticated zebra finches had a greater number of steroids that were detectable in plasma, compared to wild zebra finches. First, 7 steroids were detectable in the circulation of domesticated zebra finches (Table 1; Figure 2). Androstenedione, which was not detectable in plasma from wild zebra finches, was detectable in plasma from domesticated males and females. Second, a greater proportion of domesticated zebra finch samples had detectable levels of progesterone, testosterone, androsterone, and 5α -DHT (Table 1).

Next, we focused on subjects that were not nesting and compared wild and domesticated zebra finches. For the 6 steroids that were detectable in the circulation of both wild and domesticated zebra finches, steroid levels were generally higher in domesticated finches than wild finches (Table 2, Figure 4A-G). Circulating levels of testosterone were higher in males than females, regardless of Strain (Table 2, Figure 4D).

We further tested the effect of Strain on androsterone and 5α -DHT separately in males and females (Figure 4F-G). Domesticated males had higher circulating androsterone than wild males (Figure 4F; $W = 49.0$; $P < 0.01$), but this pattern was not significant for females (Figure 4F; $W = 55.5$, $P = 0.24$). Circulating 5α -DHT was not significantly different between wild and domesticated zebra finches (Figure 4G; males, $W = 38.0$; $P = 0.07$; females, $W = 49.5$, $P = 0.29$).

4. Discussion

Our results highlight the effects of both strain (level of domestication) and reproductive state on circulating steroid profiles of male and female zebra finches. In Study 1, we found significant main effects of level of domestication on circulating testosterone levels of males and females that were not nesting. Domesticated males had significantly higher circulating testosterone levels relative to Wild-Derived males. While the direction of this effect was not consistent across the sexes in Study 1, in Study 2, both male and female circulating testosterone levels were significantly elevated in domesticated zebra finches, relative to wild zebra finches. Furthermore, in Study 2, we found that circulating levels of several steroids were consistently elevated in domesticated zebra finches. During domestication, selection may have acted on traits relating to mate acquisition and resource defense in a captive environment, which could have contributed to this result. In Study 2, we also found that circulating pregnenolone and progesterone levels varied across the reproductive states in wild zebra finches. To our knowledge, this is the first characterization of steroid profiles in wild zebra finches. Finally, from our literature review, we conclude that large variation across strains and reproductive states is a phenomenon seen across multiple laboratories that have studied zebra finches.

4.1 Steroid profiles in the circulation of wild and domesticated zebra finches

In Study 2, the observed elevations of both circulating steroid hormones and their precursors in domesticated zebra finches may signify a general elevation of all, or many, sex steroids in circulation and tissues. Such a general elevation could be caused by differences in the availability of the steroid precursor, cholesterol, or by differences in

proteins involved in early stages of gonadal and adrenal steroidogenesis (e.g., TSPO, StAR, CYP11A1). Future studies should examine levels of cholesterol in the circulation and levels of steroidogenic proteins in the adrenals and gonads of wild and domesticated zebra finches.

In addition, the data indicate variation in circulating steroid levels across the breeding cycle. Many songbirds show peaks in circulating testosterone during courtship and/or egg laying (e.g., Vleck and Priedkalns, 1985; Hegner and Wingfield, 1986; Ball and Wingfield, 1987; Wingfield and Farner, 1978). Here, we found no changes in circulating testosterone levels across three reproductive states in wild zebra finches; however, all of our samples were collected after egg laying and we might have missed an earlier peak in testosterone. Interestingly, we did see elevations of circulating pregnenolone and progesterone during Feeding Chicks. To our knowledge, this pattern has not been previously reported in songbirds. The functions of pregnenolone and progesterone during Feeding Chicks are unclear. These steroids might regulate egg laying (Furr et al., 1973), and circulating levels of pregnenolone in Greylag geese (*Anser anser*) are lowest during incubation (Hirschenhauser et al., 1999).

Furthermore, we compiled circulating testosterone levels in male zebra finches across reproductive states from previously published studies (Table 3). Consistent with the present results, there is a general pattern of elevated plasma testosterone levels in males from domesticated populations of zebra finches compared to wild stocks, across reproductive conditions (Table 3). Additionally, there is pronounced variability in circulating testosterone levels across studies of domesticated males. Some of this variability likely reflects different assay methods, but some of the variability might

reflect genetic and environmental differences among domesticated populations. Taken together, these results highlight the importance of reporting and controlling for reproductive state and population in studies of zebra finches. Future studies should examine circulating testosterone levels across a full breeding cycle (including non-breeding condition) for both wild and domesticated zebra finches.

4.2 Environmental predictability and breeding in zebra finches

While zebra finches have long been cited as the classic example of an opportunistically-breeding bird species (Serventy, 1971; Perfito et al., 2007; Hahn et al., 2008; Griffith and Buchanan 2010), the degree of this opportunism depends on the predictability of the habitat. Zebra finches breed more seasonally along the coast of Australia, where rain and weather conditions are seasonally predictable (Perfito, 2010). Perfito et al. (2007) compared the hypothalamic-pituitary-gonadal (HPG) axis of non-breeding (winter) zebra finches from the coast and non-breeding (drought) zebra finches from central Australia. Non-breeding zebra finches from the predictable coastal habitat had lower circulating levels of luteinizing hormone (LH) and smaller gonads than non-breeding zebra finches from the unpredictable central habitat (Perfito et al., 2007). It is possible that in unpredictable environments, zebra finches maintain intermediate levels of circulating testosterone year round.

Assuming captivity increases environmental predictability, we might expect that domesticated male zebra finches would have elevated testosterone relative to wild male zebra finches. Whereas wild zebra finches are nomadic (Zann, 1996) and likely travel through high- and low-quality habitats, domesticated zebra finches have

experienced generations of high-quality conditions for breeding. This prolonged change in environmental conditions could select for further elevations in circulating testosterone of domesticated males. Indeed our survey of the literature suggests that circulating testosterone levels may be elevated in domesticated male zebra finches across reproductive states (including non-breeding condition) (Table 3). Interestingly, our data also show that domesticated female zebra finches may have elevated plasma steroid levels, suggesting there may have been selection for elevated steroids in domesticated female zebra finches as well as males.

4.3 Additional potential constraints on circulating sex steroid levels in wild zebra finches

While unpredictable environmental conditions may be one constraint on circulating steroid levels in wild zebra finches, there are other potential constraints. Elevated circulating testosterone has many “costs” (Wingfield et al., 2001), including (1) suppression of the immune system (Folstad and Karter, 1992, Roberts et al., 2004), (2) reduction of fat stores, and (3) elevation of basal metabolic rate (Buchanan et al 2001).

Testosterone can also have costs for monogamous pair bonding. Monogamous avian species generally have lower circulating testosterone levels than polygynous avian species (Garamszegi et al., 2005). In fact, elevated circulating testosterone in males is associated with an increase in extra-pair paternity across species (Garamszegi et al., 2005). Further, testosterone administration increases extra-pair copulations in sparrows (Wingfield, 1984). While domesticated zebra finches are highly sexually monogamous compared to many songbirds, they do have much higher levels of extra-pair paternity than wild zebra finches (Griffith et al., 2010). This relationship raises the possibility that

selection for a different mating strategy in domesticated zebra finches could have had a concomitant influence on circulating testosterone levels.

Decreased female mate choice in domesticated zebra finches could result in an increase in extra-pair courtship in males, if females choose mates based on their perceived commitment to the partnership. Indeed, females appear “less choosy” during mate choice tests in some domesticated zebra finches (Rustein et al., 2007). Alternatively, in domesticated populations, there may be greater selection for testosterone-dependent mating behaviors by males that secure additional mating opportunities. Forstmeier et al. (2011) found higher levels of extra-pair paternity and male competition in domesticated zebra finches, relative to wild-caught zebra finches. Taken together, these results raise interesting questions about the effect of domestication on mating systems and social behavior in zebra finches.

4.5 Conclusions

Overall, the present results from LC-MS/MS analyses suggest that domesticated zebra finches generally have higher levels of circulating steroids, including testosterone and other androgens and progestins, than wild zebra finches. Increased environmental predictability or other factors during domestication might have selected for higher circulating androgen levels in domesticated zebra finches. These results highlight the importance of considering the potential effects of strain and reproductive state in research on domesticated zebra finches, and suggest that extrapolations from domesticated to wild birds (or vice versa) need to be qualified and tested. Furthermore, these results raise

interesting questions about the effect of domestication on mating systems and social behavior in zebra finches.

Acknowledgements

We thank Keith Leggett, Garry Dowling, Mark Tilley and Vicki Dowling for logistical support at the University of New South Wales Fowler's Gap Arid Zone Research Station. For assistance in the field, we thank Luke S.C. McCowan. Funding sources included: Canadian Institutes of Health Research (CIHR) Operating Grant 133606 to KKS; National Science Foundation graduate research fellowship (NSF GRFP) to NHP; and Australian Research Council grant DP0881019 to SCG, and DP130100417 to SCG and KLB.

References

- Adkins-Regan, E. 2009. Neuroendocrinology of social behavior. *ILAR Journal*, 50, 5-14.
- Adkins-Regan, E., Abdelnabi, M., Mobarak, M. Ottinger, M. A. 1990. Sex steroid levels in developing and adult male and female zebra finches (*Poephila guttata*). *Gen. Comp. Endocrinol.*, 78, 93-109.
- Albert, F. W., O. Shchepina, C. Winter, H. Römler, D. Teupser et al., 2008. Phenotypic differences in behavior, physiology and neurochemistry between rats selected for tameness and for defensive aggression towards humans. *Horm. Behav.* 53: 413–421.
- Alonso-Alvarez, C., Bertrand, S., Faivre, B., Chastel, O., Sorci, G. 2007. Testosterone and oxidative stress: the oxidation handicap hypothesis. *Proc. R. Soc. B* 274, 819–825
- Auchus, R.J. 2004. The backdoor pathway to dihydrotestosterone. *Trends Endocrinol. Metabol.*, 15, 432–438.
- Ball, G.F. and Wingfield, J.C. 1987. Changes in Plasma Levels of Luteinizing Hormone and Sex Steroid Hormones in Relation to Multiple-Broodedness and Nest-Site Density in Male Starlings. *Physiol. Zool.*, 60, 191-199.
- Belyaev, D.K., 1979. Destabilizing selection as a factor in domestication. *J. Heredity*, 70:301-308.
- Black, J. M., 1996. *Partnerships in Birds: The Study of Monogamy*. Oxford University Press, UK.
- Buchanan, K.L., Evans, M.R., Goldsmith, A.R., Bryant, D.M., Rowe, L.V. 2001. Testosterone influences basal metabolic rate in male house sparrows: a new cost of dominance signalling? *Proc. R. Soc. B.*, 268, 1337-1344.
- Burley, N.T., Price, D.K., Zann, R.A. 1992. Bill color, reproduction and condition effects in wild and domesticated zebra finches. *Auk*, 109, 13-23.
- Calisi, R.M., and Bentley, G.E. 2009. Lab and field experiments: are they the same animal? *Horm. Behav.*, 56, 1-10.
- Carr, R.A. and Zann, R.A. 1986. The morphological identification of domesticated zebra finches, *Poephila guttata* (Passeriformes:Estrilidae), in Australia. *Aust. J. Zool.*, 34: 439-48.

- Charlier, T.D., Po, D.W.L., Newman, A.E., Shah, A.H., Saldanha, C.J., Soma, K.K.,
2010. 17β -Estradiol levels in male zebra finch brain: Combining Palkovits punch
and an ultrasensitive radioimmunoassay. *Gen. Comp. Endocrinol.*, 167:18-26.
- Dickens, M.J., and Bentley, G.E. 2014. Stress, captivity, and reproduction in a wild bird
species. *Horm Behav*, 66, 685-693.
- Ericsson, M., Fallahsharoudi, A., Bergquist, J., Kushnir, M.M., Jensen, P. 2014.
Domestication effects on behavioural and hormonal responses to acute stress in
chickens. *Physiol. Behav.*, 133: 161-169.
- Fokidis, H.B., Adomat, H.H., Kharmate, G., Hosseini-Beheshti, E., Guns, E.S., and Soma,
K.K. 2015. Regulation of local steroidogenesis in the brain and in prostate cancer:
Lessons learned from interdisciplinary collaboration. *Front Neuroendocrinol*, 36,
108-129.
- Folstad, I., and Karter, A.J. 1992. Parasites, bright males, and the immunocompetence
handicap. *Am. Nat.*, 139, 603-622,
- Forstmeier, W., Segelbacher, G., Mueller, J.C., Kempenaers, B. 2007. Genetic variation
and differentiation in captive and wild zebra finches (*Taeniopygia guttata*) *Mol.*
Ecol., 16: 4039–4050 doi: 10.1111/j.1365-294X.2007.03444.x
- Forstmeier, W., Martin, K., Bolund, E., Schielzeth, H., Kempenaers, B. 2011. Female
extrapair mating behavior can evolve via indirect selection on males. *PNAS*, 108,
10608-10613.
- Furr, B. J.A., Bonney, R.C., England, R.J., Cunningham, F.J. 1973. Luteinizing hormone
and progesterone in peripheral blood during the ovulatory cycle of the hen *Gallus*
domesticus. *J. Endocrinol.*, 57,159-169.
- Garamszegi, L.Z., Eens, M., Hurtrez-Bousses, S., Møller, A.P. 2005. Testosterone, testes
size, and mating success in birds: a comparative study. *Horm. Behav.*, 47, 389-
409.
- Griffith, S.C., and Buchanan, K.L., 2010. The Zebra Finch: the ultimate Australian
supermodel. *Emu*. 110, v–xii
- Griffith, S.C., Pryke, S.R., Mariette, M. 2008. Nest box use by the zebra finch
Taeniopygia guttata: implications for reproductive success and research. *Emu*.
108,311-319.
- Griffith, S. C., Holleley, C. E., Mariette, M. M., Pryke, S. R., Svedin, N., 2010. Low level
of extrapair parentage in wild zebra finches. *Anim. Behav.* 79, 261–264.

- Gilby, A.J., Mainwaring, M.C., Griffith, S.C. 2013. Incubation behaviour and hatching synchrony differ in wild and captive populations of the zebra finch. *Anim Behav*, 85, 1329-1334.
- Hahn, T.P., Cornelius, J.M., Sewall, K.B., Kelsey, T.R., Hau, M., Perfito, N. 2008. Environmental regulation of annual schedules in opportunistically-breeding songbirds: adaptive specializations or variations on a theme of white-crowned sparrow? *Gen. Comp. Endocrinol.*, 157, 217-226.
- Hegner, R.E., and Wingfield, J.C. 1986. Behavioral and endocrine correlates of multiple brooding in the semicolonial house sparrow *Passer domesticus* I. Males. *Horm. Behav.*, 20, 294-312.
- Homberger, B., Jenni-Eiermann, S., Roulin, A., Jenni, L. 2013. The impact of pre- and post-natal contexts on immunity, glucocorticoids and oxidative stress resistance in wild and domesticated grey partridges. *Funct. Ecol.*, 27, 1042-1054.
- Hirschenhauser, K. Möstl, E., Kotrschal, K. 1999. Seasonal Patterns of Sex Steroids Determined from Feces in Different Social Categories of Greylag Geese (*Anser anser*). *Gen. Comp. Endocrinol.*, 144, 67-79
- Hutchison, J.B., Wingfield, J.C., Hutchison, R.E. 1984. Sex differences in plasma concentrations of steroids during the sensitive period for brain differentiation in the zebra finch. *J. Endocrinol.* 103:363-369.
- Kabelik, D., Schrock, S.E., Ayres, L.C., Goodson, J.L. 2011 Estrogenic regulation of dopaminergic neurons in the opportunistically breeding zebra finch. *Gen. Comp. Endocrinol.*, 173, 96-104
- Kalhorn, T.F., Page, S.T., Howald, W.N., Mostaghel, E.A., Nelson, P.S. 2007. Analysis of testosterone and dihydrotestosterone from biological fluids as the oxime derivatives using high-performance liquid chromatography/tandem mass spectrometry. *Rapid Commun. Mass Spectrom.*, 21, 3200-3206.
- Koren, L., Ng, E.S.M., Soma, K.K., Wynne-Edwards, K.E. 2012. Sample Preparation and Liquid Chromatography-Tandem Mass Spectrometry for Multiple Steroids in Mammalian and Avian Circulation. *PLoS ONE*. 7(2): e32496.
- Korsia, S., and Bottjer, S.W. 1991. Chronic Testosterone Treatment Impairs Vocal Learning in Male Zebra Finches during a Restricted Period of Development. *J. Neurosci.*, 11, 2362-237.
- Künzl, C., and Sachser, N. 1999. The behavioral endocrinology of domestication: a comparison between the domestic Guinea Pig (*Cavia aperea* f. *porcellus*) and Its Wild Ancestor, the Cavy (*Cavia aperea*). *Horm. Behav.*, 35,28-37.

- Lincoln, G.A., Lincoln, C.E., McNeilly, A.S. 1990. Seasonal cycles in the blood plasma concentration of FSH, inhibin and testosterone, and testicular size in rams of wild, feral and domesticated breeds of sheep. *J. Reprod. Fertility*, 88, 623-633.
- Luine, V., Nottebohm, F., Harding C., McEwen, B.S. 1980. Androgen affects cholinergic enzymes in syringeal motor neurons and muscle. *Brain Res.* 192,89-107.
- Lynn, S.E., Perfito, N., Guardado, D., Bentley, G.E. 2015. Food, stress, and circulating testosterone: Cue integration by the testes, not the brain, in male zebra finches (*Taeniopygia guttata*). *Gen. Comp. Endocrinol.*, 215, 1–9.
- Mainwaring, M.C., Hartley, I.R., Gilby, A.J., Griffith, S.C. 2010. Hatching asynchrony and growth tradeoffs within domesticated and wild zebra finch, *Taeniopygia guttata*, broods. *Biol J Linnean Soc*, 100, 763-773.
- Mariette, M.M., Pariser, E.C., Gilby, A.J., Magrath, M.J., Pryke, S.R., Griffith, S.C. 2011. Using an electronic monitoring system to link offspring provisioning and foraging behavior of a wild passerine. *Auk*, 128, 26-35.
- Martin, J.T. 1978. Embryonic pituitary adrenal axis, behavior development, and domestication in birds. *Amer. Zool.* 18: 489-499.
- McCowan, L. S., Mainwaring, M. C., Prior, N. H., Griffith, S. C. 2015. Personality in the wild zebra finch: exploration, sociality, and reproduction. *Behav. Ecol. aru239*.
- Newman, A.E.M., Chin, E.H., Schmidt, K.L., Bond, L., Wynne-Edwards, K.E., Soma, K.K., 2008. Analysis of steroids in songbird plasma and brain by coupling solid phase extraction to radioimmunoassay. *Gen. Comp. Endocrinol.*, 155,503–510.
- Overk, C.R., Perez, S.E., Ma, C., Taves, M.D., Soma, K.K., Mufson, E.J. 2013. Sex steroid levels and AD-like pathology in 3xTgAD mice. *J. Neuroendocrinol.*, 25:131-144.
- Perfito, N., 2010. The reproductive and stress physiology of zebra finches in context: integrating field and laboratory studies. *Emu* 110:199-208.
- Perfito, N., Zann, R. A., Bentley, G. E., Hau, M., 2007. Opportunism at work: habitat predictability affects reproductive readiness in free-living zebra finches. *Funct. Ecology.* 21:291–301.
- Perfito, N., Zann, R., Ubuka, T., Bentley, G., Hau, M. 2011. Potential roles for GnIH and GNRH-II in reproductive axis regulation of an opportunistically breeding songbird. *Gen. Comp. Endocrinol.*, 173, 20-26.

- Priedkalns, J., Oksche, A., Vleck, C.M., Bennett, R.K. 1984. The response of the hypothalamo-gonadal system to environmental factors in the zebra finch, *Poephila guttata castanotis*. *Cell Tissue. Res.*, 238, 23–35. doi:10.1007/BF00215140
- Pröve, E., 1983. Hormonal correlates of behavioural development in male zebra finches, in: Balthazart, J., Pröve, E., Gilles, R. (Eds.), *Hormones and Behaviour in Higher Vertebrates*, Springer-Verlag, Berlin/Heidelberg, pp. 368–374.
- Prior, N.H., and Soma, K.K. 2015. Neuroendocrine regulation of long-term pair maintenance in the monogamous zebra finch. *Horm Behav.*, in press. doi:10.1016/j.yhbeh.2015.04.014
- Prior, N.H., Heimovics, S.A., Soma, K. K., 2013. Effects of water restriction on reproductive physiology and affiliative behavior in an opportunistically-breeding and monogamous songbird, the zebra finch. *Horm. Behav.* 63, 462-474.
- Prior, N.H., Yap, K.N., Adomat, H.H., Mainwaring, M.C., Fokidis, H.B., Guns, E.S., Buchanan, K.L., Griffith, S.C., Soma, K.K. 2016. Sex steroid profiles and pair-maintenance behavior of captive wild-caught zebra finches (*Taeniopygia guttata*). *J. Comp. Physiol. A.* 202, 35-44.
- R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Roberts, M.L., Buchanan, K.L., Evans, M.R. 2004. Testing the immunocompetence handicap hypothesis: a review of the evidence. *Anim. Behav.*, 68, 227-239.
- Rutstein, A.N., Brazill-Boast, J., Griffith, S.C. 2007. Evaluating mate choice in the zebra finch. *Anim. Behav.* 74, 1277-1284.
- Schlinger, B.A., Ira Lan, N., Grisham, W., Thompson, L. 1999. Androgen Synthesis in a Songbird: A Study of Cyp17 (17 α -Hydroxylase/C17,20-Lyase) Activity in the Zebra Finch. *Gen. Comp. Endocrinol.* 113, 46-58.
- Setchell, B. P. 1992. Domestication and reproduction. *Anim. Reprod. Sci.*, 28, 195- 202.
- Serventy, D.L., 1971. Biology of desert birds. In: Farner, D.S., King, J.L. (Eds.), *Avian Biology 1*. Academic Press, New York, pp. 287–33
- Sossinka, R., 1970. Domestikationsercheinungen beim Zebrafinken *Taeniopygia guttata castanotis* (GOULD). *Jahrb. Abt. Syst. Okol. Geogr. Tiere* 97, 455–521.
- Suzuki, K., Yamada, H., Kobayashi, T., Okanoya, K., 2012. Decreased fecal corticosterone levels due to domestication: a comparison between the white-backed munia (*Lonchura striata*) and its domesticated strain, the Bengalese finch

- (*Lonchura striata* var. *domestica*) with a suggestion for complex song evolution. *J. Exp. Zool. A.*, 317, 561–570.
- Taves, M.D., Schmidt, K.L., Ruhr, I.M., Kapusta, K., Prior, N.H., Soma, K.K., 2010. Steroid concentrations in plasma, whole blood and brain: effects of saline perfusion to remove blood contamination from brain. *PLoS One* 5:e15727
- Taves, M.D., Ma, C., Heimovics, S.A., Saldanha, C.J., Soma, K.K., 2011. Measurement of steroid concentrations in brain tissue: methodological considerations. *Front Endocrinol* 2, 39.
- Wade, J., Springer, M.L., Wingfield, J.C., Arnold, A.P. 1996. Neither testicular androgens nor embryonic aromatase activity alters morphology of the neural song system in zebra finches. *Biol. Reprod.*, 55, 1126-1132.
- Wilbrecht, L., Williams, H., Gangadhar, N., Nottebohm, F. 2006. High Levels of New Neuron Addition Persist When the Sensitive Period for Song Learning Is Experimentally Prolonged. *J. Neurosci.*, 26, 9135–9141.
- Wilkins, A.S., Wrangham, R.W., Tecumseh, W.F. 2014. The “Domestication Syndrome” in mammals: A unified explanation based on neural crest cell behavior and genetics. *Genetics*. 197, 795-808.
- Wingfield, J.C. 1984. Androgens and mating systems: Testosterone-induced polygyny in normally monogamous birds. *Auk*. 101: 665-671.
- Wingfield, J.C. and Farner, D.S. 1975. The determination of five steroids in avian plasma by radioimmunoassay and competitive protein binding. *Steroids* 26, 311-327.
- Wingfield, J.C., Lynn, S., Soma, K.K. 2001. Avoiding the “costs” of testosterone: ecological bases of hormone-behavior interactions. *Brain Behav. Evol.*, 57, 239-251.
- Woodward, C.C., and Strange, R.J. 1987. Physiological stress responses in wild and hatchery-reared rainbow trout. *Trans. Am. Fish Soc.* 1987;116:574–9.
- Vleck, C., and Friedkalns, J., 1985. Reproduction in zebra finches: Hormone levels and effect of dehydration. *Condor*. 87, 37–46.
- Vockel, A. Prove, E. Balthazart, J. 1990. Effects of castration and testosterone treatment on the activity of testosterone-metabolizing enzymes in the brain of male and female zebra finches. *J. Neurobiol.* 21, 808-825.
- Zann, R. A., 1996. *The Zebra Finch: A Synthesis of Laboratory and Field Studies*. Oxford University Press, USA.

740 Zeder, M.A. 2012 Pathways to animal domestication, In: Biodiversity in Agriculture:
741 Domestication, Evolution, and Sustainability, edited by P. Gepts, T.R. Famula,
742 R.L. Bettinger et al. Published by Cambridge University Press, USA.
743

744 Table 1: Percent of plasma samples from wild and domesticated zebra finches that are non-detectable (ND), 3-10X above background,
745 or >10X above background using a LC-MS/MS method to simultaneously examine multiple steroids.

		Pregnenolone	Progesterone	DHEA	Androsterone	Androstenedione	Testosterone	5 α -DHT
Wild N = 61	ND	0	13	0	61	0	54	93
	3-10X	0	41	0	37	0	26	0
	>10X	100	46	100	2	0	20	7
Domesticated N = 13	ND	0	0	0	38	46	0	70
	3-10X	0	30	0	54	38	30	30
	>10X	100	70	100	8	15	70	0

746

747 Note: Both males and females are included here. Wild zebra finches were in several reproductive states, and domesticated zebra
748 finches were sampled while not actively nesting.

749

Table 2: Effects of Strain (Wild or Domesticated) and Sex on circulating steroid levels, as measured by a LC/MS-MS method.

	Strain		Sex		Strain \times Sex	
	F _(1,32)	P	F _(1,31)	P	F _(1,30)	P
Pregnenolone	96.18	< 0.001	< 0.01	0.95	0.47	0.50
Progesterone	31.86	< 0.001	0.19	0.67	1.84	0.19
DHEA	44.05	< 0.001	1.53	0.23	3.35	0.08
Testosterone	22.15	< 0.001	13.97	< 0.001	0.11	0.75
Androsterone*	20.84	< 0.001	12.92	0.001	11.51	< 0.01
5 α -DHT*	4.33	< 0.05	0.26	0.61	0.06	0.80

Note: ANOVAs were run using glm models, and data were transformed as necessary to meet the assumptions of the test. All models are reported as ANOVA summaries with the resulting F statistic. All subjects were not nesting. Bolded values indicate significant effects. *For androsterone and 5 α -DHT, residuals of the GLM models were not normally distributed, and Wilcoxon rank sum tests were used to determine the effect of Strain on females and males separately.

765 Table 3: Studies reporting circulating testosterone levels in wild or domesticated adult male zebra finches.

766

	Reference	Wild vs. Domesticated	Housing conditions	Strain/Location	Testosterone (ng/mL)	Method	n
Non-Breeding	Vleck & Priedkalns, 1985	Wild-Caught	Water Restricted	Barmera, S. Aus., AUS	< 0.28*	RIA	16 (1 pooled)
	Priedkalns et al., 1984	Wild-Caught	Water Restricted	S. Australia, AUS	< 0.45*	RIA	16 (4 pooled)
	Perfito et al., 2007	Wild	NA	Alice Springs, AUS	0.20 ± 0.02	RIA	12 (2/11 samples were detectable)
	Prior et al., 2013	Domesticated	Water Restricted	Vancouver, BC, CAN	0.70 ± 0.2	RIA	11
Not Nesting	Hutchinson et al., 1984	Domesticated	Individually housed	Cambridge, UK	0.29	RIA	9
	Wade et al., 1996	Domesticated	Same Sex housed	Los Angeles, CA, USA	0.52 ± 0.1	RIA	7
	Alonso-Alvarez et al., 2007	Domesticated	Individually housed	Station Biologique de Foljuif, FR	1.58 ± 0.4	RIA	12
	Prior et al., 2013	Domesticated	Not Nesting	Vancouver, BC, CAN	1.90 ± 0.4	RIA	9
	Charlier et al., 2010	Domesticated	Same Sex housed	Vancouver, BC, CAN	2.50	RIA	8
	Lynn et al., 2015	Domesticated	Individually housed	Berkeley, CA, USA	3.50	EIA	8
Breeding	Perfito et al., 2007	Wild	NA	Victoria, AUS	0.51 ± 0.24	RIA	6 (6/8 samples were detectable)
	Vleck & Priedkalns, 1985	Domesticated	Late Incubating	Adelaide, S. Aus., AUS	0.24-1.06*	RIA	5 (1 pooled)
	Vleck &	Domesticated	Feeding	Adelaide, S. Aus.,	0.25*	RIA	4

	Priedkalns, 1985		chicks	AUS			(1 pooled)
	Koren et al., 2012	Domesticated	?	Vancouver, BC, CAN	0.51	LC-MS/MS	7
	Vleck & Priedkalns, 1985	Domesticated	Courting	Adelaide, S. Aus., AUS	0.80*	RIA	9 (1 pooled)
	Vleck & Priedkalns, 1985	Domesticated	Early Incubating	Adelaide, S. Aus., AUS	1.06*	RIA	4 (1 pooled)
Unknown	Vleck & Priedkalns, 1985	Wild-Caught	?	Barmera, S. Aus., AUS	0.30*	RIA	5 (1 pooled)
	Wilbrecht et al., 2006	Domesticated	?	Millbrook, NY, USA	0.16 ± 0.16	RIA	4
	Korsia and Bottjer, 1991	Domesticated	?	Los Angeles, CA, USA	0.17 ± 0.13	RIA	13
	Wilbrecht et al., 2006	Domesticated	?	Millbrook, NY, USA	0.25 ± 0.1	RIA	6
	Schlenger et al., 1999	Domesticated	?	Los Angeles, CA, USA	0.31	RIA	?
	Pröve, 1983	Domesticated	?	Bielfeld, Germany	0.33	RIA	7
	Luine et al., 1980	Domesticated	?	Deer Park, NY, USA	0.65 ± 0.1	RIA	9
	Wilbrecht et al., 2006	Domesticated	?	Millbrook, NY, USA	0.68 ± 0.37	RIA	5
	Adkins-Regan et al., 1990	Domesticated	?	College Park, MD, USA	1.90	RIA	16
	Vockel et al., 1990	Domesticated	?	Liege, Belgium	3.20	RIA	16
	Kabelik et al., 2011	Domesticated	?	Bloomington, IN, USA	3.47 ± 0.67	RIA	6

767

768

769 Note: If we estimated testosterone levels from graphs, then we present only the estimated mean. If the mean and SEM were reported in
770 the text of the original reference, then we provide those here. The strain/location of the last author is reported when no specific details
771 are listed in the text. NA = not applicable, EIA = enzyme immunoassay (ELISA), RIA = radioimmunoassay, LC-MS/MS = liquid
772 chromatography-tandem mass spectrometry.

773 *Total androgens measured, not a specific assay for testosterone.

Figure Legends

Figure 1: In Study 1, plasma testosterone levels were measured via radioimmunoassay (RIA).

(A) Plasma testosterone levels in wild zebra finches did not vary with nesting activity but were higher in males than females. Here, nesting zebra finches includes subjects that were incubating eggs or feeding chicks. (B) In female zebra finches that were not nesting, plasma testosterone levels differed across “strains” (Wild, Captive Wild-Caught, and Domesticated) and were highest in wild females. Wild-derived females were not assessed (NA). (C) In male zebra finches that were not nesting, plasma testosterone levels also differed across strains and were lowest in wild-derived males in captivity. In (A), $** p \leq 0.01$. In (B) and (C), letters indicate significant differences.

Figure 2: A simplified steroidogenic pathway indicating steroids in the “traditional” pathway to 5α -DHT and the “backdoor” pathway to 5α -DHT. Steroids in the backdoor pathway are in italics. Boxed steroids indicate those that were detectable in the plasma of zebra finches (solid line = detectable in wild and domesticated finches; dotted line = detectable in domesticated finches only).

Figure 3: In Study 2, a panel of steroids was measured via liquid chromatography-tandem mass spectrometry (LC-MS/MS) in plasma from wild zebra finches. Wild female and male zebra finches were examined across reproductive states. (A) Pregnenolone, (B) Progesterone, (C) DHEA, and (D) Testosterone. Plasma pregnenolone and progesterone levels were higher during the Feeding Chicks stage, and plasma testosterone levels were higher in males than females. In (A) and (B), letters indicate significant differences.

797

798 Figure 4: In Study 2, a panel of steroids was measured via LC-MS/MS in plasma from wild and
799 domesticated zebra finches that were not nesting. Overall, domesticated zebra finches have
800 higher circulating steroid levels than wild finches. (A) Pregnenolone, (B) Progesterone, (C)
801 DHEA, (D) Testosterone, (E) Androstenedione, (F) Androsterone, and (G) 5 α -DHT. ND = non-
802 detectable. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$.


803

804

Figure 1


Study 1: RIA

A



B

Not Nesting Females



C

Not Nesting Males




Figure 2




Figure 3

Study 2: LC-MS/MS

Wild Zebra Finches




Figure 4

Study 2: LC-MS/MS

