Lipid signaling and fat storage in the dark-eyed junco

Jacqueline M. Ho^{a,b,*}, Christine M. Bergeon Burns^{a,b}, Nikki M. Rendon^{a,b}, Kimberly A. Rosvall^{a,b,c}, Heather B. Bradshaw^{b,c,d}, Ellen D. Ketterson^{a,b} and Gregory E. Demas^{a,b,c}

Abstract

Seasonal hyperphagia and fattening promote survivorship in migratory and wintering birds, but reduced adiposity may be more advantageous during the breeding season. Factors such as photoperiod, temperature, and food predictability are known environmental determinants of fat storage, but the underlying neuroendocrine mechanisms are less clear. Endocannabinoids and other lipid signaling molecules regulate multiple aspects of energy balance including appetite and lipid metabolism. However, these functions have been established primarily in mammals; thus the role of lipid signals in avian fat storage remains largely undefined. Here we examined relationships between endocannabinoid signaling and individual variation in fat storage in captive white-winged juncos (*Junco hyemalis aikenii*) following a transition to long-day photoperiods. We report that levels of the endocannabinoid 2-arachidonoylglycerol (2-AG), but not anandamide (AEA), in furcular and abdominal fat depots correlate negatively with fat mass. Hindbrain mRNA expression of CB₁ endocannabinoid receptors also correlates negatively with levels of fat, demonstrating that fatter animals experience less central and peripheral endocannabinoid signaling when in breeding condition. Concentrations of the anorexigenic lipid, oleoylethanolamide (OEA), also inversely relate to adiposity. These findings demonstrate unique and significant relationships between adiposity and lipid signaling molecules in the brain and periphery, thereby suggesting a potential role for lipid signals in mediating adaptive levels of fat storage.

Key words: fattening, adiposity, endocannabinoid, CB₁, oleoylethanolamide, palmitoylethanolamide

1. Introduction

Many species accumulate energy stores in anticipation of prolonged periods of increased energy demand or decreased food availability. In passerines, this primarily occurs prior to autumn or vernal migration and during winter (Baldwin and Kendeigh, 1938; Berthold, 1993). Benefits conferred by greater fat storage (e.g., fasting capacity) are tempered by costs that include increased energetic demands of flight (Freed, 1981), loss of agility (Blem, 1975), and greater predation risk with higher foraging rates (Lima, 1986). Thus, individuals must effectively balance this trade-off to maximize survivorship benefits at minimal cost (Rogers, 2015). As a result, fat stores are reduced as birds complete gonadal development and enter the breeding life-history stage (Williams, 2012). The timing and extent of winter fattening are driven by environmental factors such as temperature (Rogers, 1995), food abundance and predictability (Rogers and Smith, 1993; Witter et al., 1995; Biebach, 1996), and climate (Nolan and Ketterson, 1983), whereas migratory fattening follows a circannual rhythm predominantly synchronized by changes in photoperiod (Gwinner, 1986; Berthold, 1996).

The neuroendocrine mechanisms that mediate seasonal changes in fat are less clear. Studies have typically focused on few physiological systems, and results often suggest complex relationships that may vary by species as well as life-history stage. For example, corticosterone has been suggested to drive changes in feeding and fat deposition during winter in the migratory slate-colored race of the dark-eyed junco (*Junco hyemalis hyemalis*), a common North American songbird (Gray et al., 1990; Rogers et al., 1993). In contrast, evidence for relationships between corticosterone and pre-migratory fattening in this subspecies have been mixed, and may further be complicated by differences between migratory and resident subspecies (Ramenofsky et al., 1999; Holberton et al., 2007; Holberton et al., 2008; Bauer et al., 2015; Fudickar et al., 2016). In

addition to glucocorticoids, hormones such as testosterone (Lofts and Marshall, 1961; Morton and Mewalt, 1962; Stetson and Erickson, 1972; Schwabl and Farner, 1989; Deviche, 1995) and prolactin (Meier and Farner, 1964; Buntin and Tesch, 1985; Boswell et al., 1995; Holberton and Dufty, 2005) have been associated with feeding and fat regulation in birds. However, neuroendocrine regulators classically implicated in mammalian energy balance such as neuropeptide Y, insulin, and glucagon have been examined to a lesser extent (Totzke et al., 1997; Holberton and Dufty, 2005; Ramakrishnan et al., 2007).

Similarly, little is known of energy regulatory functions of endocannabinoids in birds, despite well-established roles of these ubiquitous neuromodulators in mammalian food intake (Pagotto et al., 2006), lipogenesis (Osei-Hyiaman et al., 2005; Vettor and Pagano, 2009), and energy expenditure (Cluny et al., 2010; Quarta et al., 2010; Tam et al., 2010). Endocannabinoids are produced locally and on demand in the brain and periphery. They are capable of fine-tuning synaptic transmission by activating CB₁ receptors on presynaptic axons to inhibit transmitter release (Di Marzo et al., 2005). The appetite-stimulating effects of endocannabinoids are in part due to such retrograde inhibition of classical hypothalamic circuits (Jo et al., 2005). Moreover, *in vivo* and *in vitro* models suggest that peripheral endocannabinoid signaling promotes fat deposition by stimulating glucose uptake and lipogenesis via adipocyte CB₁ receptors (Cota et al., 2003; Jbilo et al., 2005; Pagano et al., 2007). Although relatively few studies have examined endocannabinoid signaling in birds, existing reports suggest similar functions to those observed in mammals, including promotion of food intake in chickens (Novoseletsky et al., 2011) and inhibition of hypothalamo-pituitary-adrenal (HPA) activity in European starlings (Dickens et al., 2015). Because hyperphagia is the primary means to increase fat stores in birds (King et al., 1963; King, 1972; Baggott, 1975; Blem, 1980; Piersma and Jukema, 1990), the endocannabinoid

system is a prime candidate to maintain plasticity and seasonally appropriate levels of food intake and fat storage.

In the present study, we examined the endocannabinoid system in relation to fat stores in captive dark-eyed juncos acclimating to long days. Specifically, we determined whether individual variation in adiposity of two major fat depots relate to local concentrations of the endocannabinoids 2-arachidonoylglycerol (2-AG) and anandamide (AEA). To determine whether central endocannabinoid signaling contributes to fat levels, we quantified mRNA expression of CB₁ receptors in the hindbrain, a brain region that regulates feeding in birds (Dubbeldam, 1984; Boswell et al., 1995; Moghaddam et al., 2010) and mammals (Berthoud, 2004). Given CB₁ receptors' ability to promote food intake in birds (Novoseletsky et al., 2011), we predicted that central (CB₁) and peripheral (2-AG; AEA) endocannabinoid signaling would relate positively to fat levels in the present study. We also predicted that concentrations of oleoylethanolamide (OEA) and palmitoylethanolamide (PEA), two structurally-related lipids with opposing effects on energy balance from those of endocannabinoids (Rodriguez de Fonseca et al., 2001), would correlate negatively with adiposity. However, if the relationship between lipid signaling and adiposity is dependent on current energetic need, then an alternative prediction is that fat levels would negatively correlate with orexigenic lipid signaling in contexts of fat loss, such as during preparation for breeding.

2. Methods

2.1 Animals and housing

Thirty-five male and female dark-eyed juncos (subspecies: white-winged junco, *Junco hyemalis aikenii*) were captured as juveniles in the Black Hills National Forest near Custer, South Dakota (43° 46' N, 103° 36' W) at the end of the 2009 breeding season. Juncos were caught using baited

mist-nets and potter traps. The birds were banded, measured, bled from the wing vein upon capture, and group-housed in a temporary outdoor aviary in the field. In early August, they were transported to Indiana University (39° 09' N, 86° 31' W) where they were group-housed in an indoor aviary held around 16°C and maintained on a diet of *ad libitum* millet, sunflower seeds, orange slices, mealworms (*Tenebrio molitor*), and a mixture of puppy chow, hardboiled eggs, and carrots. From August through February, the free-flying birds were captured approximately every 2-3 weeks (17±1 days) and subcutaneous fat was visually scored using established criteria (Rogers, 1991) as outlined in Table 1. Photoperiod was adjusted every two weeks to match the naturally changing photoperiod at their capture site. This allowed birds to molt and become photosensitive during spring 2010. In the wild, this subspecies overwinters at lower elevations extending from the breeding range in the Black Hills into the foothills of the nearby Rocky Mountains (Whitney, 1968; eBird, 2016). Thus, the duration of this study tracked a period of time during which birds naturally fatten over winter and lose fat as they enter breeding condition.

During the first week of March 2010, before natural photoperiod reached a stimulatory day length, biweekly photoperiod adjustments were discontinued and photoperiod was increased by 1 h every two days until a 16:8 h light:dark cycle was achieved (March 10). Birds remained on this long-day photoperiod for the duration of this study (through mid-April). During this time, birds were individually housed across four similarly-sized, mixed-sex housing rooms. This rapid increase in photoperiod is sufficient to initiate gonadal growth and loss of winter fat (Dawson et al., 2001; Bergeon Burns et al., 2014; Rosvall et al., 2016), but allows for variation in the rate at which birds adjust fat levels. For this study, we capitalized on this individual variation to examine birds that vary in degree of fat storage and pre-breeding fat loss. Juncos had high survivorship and maintained excellent physical condition in captivity. Thirty-two individuals (19

males, 13 females) were used in this study. All procedures were approved by the Bloomington Institutional Animal Care and Use Committee (BIACUC) at Indiana University.

2.2 Tissue collection

Prior to tissue collection for the present study, all birds participated in a separate experiment in which they received a series of three hormone challenges (two of chicken gonadotropin-releasing hormone, one of ovine luteinizing hormone) separated by five-day intervals. Small blood samples (100 µl) were collected before and shortly after hormone administration to assess transient hormone responses (Rosvall et al., 2013; Bergeon Burns et al., 2014). All birds were exposed to the same manipulations and these hormone challenges evoke only a short-term hormonal response that returns to basal condition within hours (Jawor et al., 2006; Rosvall et al., 2016). Basal hormone levels do not differ significantly between challenges separated by five-day intervals (Bergeon Burns et al., 2014), suggesting that five days is of sufficient duration for birds to return to basal physiological condition. Variation in the present measures of interest at the time of sample collection can therefore be reasonably attributed to natural variation among individuals.

Five days after the final hormone challenge (in mid-April; 38d after reaching 16L:8D), birds were overdosed with isoflurane inhalation and decapitated. Brains and gonads were quickly extracted under RNase-free conditions and immediately frozen on crushed dry ice. Visual analyses of gonads revealed that all birds were in reproductive condition and had testosterone concentrations comparable to free-living birds in the early breeding season (Rosvall et al., 2013; Bergeon Burns et al., 2014). Each bird was assigned a fat score by visual assessment, as before. Fat depots from the furculum and abdomen were then excised, weighed, and frozen on dry ice. Samples were stored at -80°C until further microdissection (brains) or processing (fat).

2.3 Compound extraction from fat samples

Tissues were processed based on a protocol previously described (Bradshaw et al., 2006). Briefly, fat samples (~0.08 g) were incubated on ice with 50 volumes of 100% HPLC-grade methanol for 1 h. One hundred pmol [²H₈]-anandamide (Cayman Chemical, Ann Arbor, MI) was added to the methanol/tissue sample and used as an internal standard to track recovery of the test compounds. Following incubation, samples were maintained on ice and homogenized (Polytron PT 10-35) for 1-2 min. Samples were centrifuged at 42,858 x g and 24°C for 20 min. Supernatants were collected and diluted to a 25% organic solution with HPLC-grade water. Compounds were partially purified from the supernatant using 500 mg C18 solid phase extraction columns (Varian #12113027, Harbor City, CA), conditioned with 5 ml 100% HPLC-grade methanol and 2.5 ml HPLC water prior to addition of the water/supernatant solution. Columns were washed with 2.5 ml of HPLC-grade water and three elutions were collected (1.5 ml each of 65, 80, and 100% HPLC-grade methanol). Elutions were stored at -20°C until mass spectrometric analysis. Individual furcular and abdominal fat samples were processed separately with both sexes analyzed within each batch.

2.4 HPLC/MS/MS analysis and quantification of lipid signalers

Each elution was warmed to room temperature and vortexed prior to mass spectrometric analysis. Rapid separation was obtained with 10-20 µl injections of analyte (CBM-20A prominence controller and SIL-20AC prominence autosampler; Shimadzu, Columbia, MD; maintained at 24°C) onto a Zorbax eclipse XDB 2.1x50 mm reversed phase column. Gradient elution (methanol and water; 200 µl/min) was formed under pressure on a pair of 10AdVP pumps (Shimadzu). Mass spectrometric analysis was performed using electrospray ionization

with a triple quadrupole mass spectrometer (API3000, Applied Biosystems/MDS Sciex, Foster City, CA). Levels of each compound were analyzed by multiple reactions monitoring on the HPLC/MS/MS system. In this mode, detection of each compound is based on fragmentation of a precursor ion $[M+H]^+$ or $[M-H]^+$ to yield a prominent product ion. Mass spectrometric conditions were optimized for each compound using direct flow injection of synthetic standards of each compound (Cayman Chemical). Test compounds of interest were the endocannabinoids 2-AG and AEA as well as non-cannabinoid but structurally-related *N*-acyl ethanolamides, OEA and PEA.

2.5 Hindbrain RNA isolation and cDNA synthesis

Brains were microdissected into functional regions following Soma et al. (1999), and hindbrain was reserved for this study. Hindbrain tissue was homogenized for 30-45 s to isolate total RNA using TRIzol reagent (Invitrogen, Carlsbad, CA) according to manufacturer instructions. RNA concentration was determined using a spectrophotometer and RNA integrity was verified by gel electrophoresis. All RNA samples were stored at -80°C until further analysis. Following DNase treatment (Invitrogen), 1 μg RNA was reverse transcribed into cDNA in 20 μl total volume according to manufacturer instructions using oligo dT primers and Superscript III Reverse Transcriptase (Invitrogen), yielding a stock concentration of 50 ng/ μl .

2.6 Primer Design

To ensure high specificity of primers, a portion of the gene for CB_1 , the primary endocannabinoid receptor in the brain, was sequenced for this species. Initial primers were designed using PrimerQuest (Integrated DNA Technologies, Coralville, IA) from zebra finch (*Taeniopygia guttata*) CB_1 (GenBank accession no. AF255388.1): forward, 5'-

AGTGGGACTTCCTCCGATGCATTT-3'; reverse, 5'-
ACGATAGCGATGGTCCACATCACA-3'. A 10 µl PCR reaction was performed using 1X
GoTaq Flexi DNA polymerase (Promega, Madison, WI), 2.5 mM MgCl₂, 0.4 µM of each primer,
0.2 mM of each deoxynucleotide triphosphate (dNTP), 1 U taq, distilled water, and cDNA as a
template at a concentration of 5 ng/µl. The reaction was carried out at 95°C for 2 min, then 30
cycles of 95°C for 30 sec/54°C for 30 sec/72°C for 1.5 min, followed by 72°C for 5 min. The
reaction amplified an 809-bp product which was confirmed on a 1% TBE agarose gel containing
ethidium bromide. PCR products were purified using a QIAquick PCR purification kit (Qiagen,
Valencia, CA) according to manufacturer instructions, sequenced at the Indiana Molecular
Biology Institute (Bloomington, IN), and deposited in GenBank (accession no. KX574508).
Product sequences were confirmed to be highly homologous to the CB₁ sequences for zebra
finch (97%) and the closely-related white-throated sparrow (*Zonotrichia albicollis*; 99%) using
NCBI BLAST. Junco-specific qPCR primers for CB₁ were then designed based on this sequence
using PrimerQuest: forward (300 nM), 5'-TTACGTGGGCTCCAACGATATCCA-3'; reverse
(300 nM), 5'-TGATGTTGACCTGCTCTGAGGGAA-3' (amplifying a 179-bp product).
Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) served as a reference gene and its
primers were based from zebra finch sequence (GenBank accession no. AF255390): forward
(300 nM), 5'-TGACCTGCCGTCTGGAAAA-3'; reverse (300 nM), 5'-
CCATCAGCAGCAGCCTTCA-3' (amplifying a 70-bp product). Standard curves with serial
dilutions of cDNA resulted in >0.99 correlation coefficients and amplification efficiency at
100.1% and 109.3% for CB₁ and GAPDH, respectively.

2.7 Quantitative PCR

Quantitative PCR was performed on a Stratagene MX3000P qPCR machine by using PerfeCta

SYBR Green SuperMix with Low ROX (Quanta BioSciences, Inc., Gaithersburg, MD). Twenty-five μ l reactions were carried out containing half the reaction volume (12.5 μ l) of SYBR, 12.5 ng cDNA, and forward and reverse primers at a concentration of 300 nM. Samples were run in duplicate at 95°C for 5 min, 40 cycles of 95°C for 30 sec/60°C for 1 min/70°C for 30 sec, followed by dissociation curves of PCR products to confirm primer specificity.

Samples were run across two plates and a common calibrator run on each confirmed low inter-plate variability (% CV = 8.25). No template controls were also run in duplicate on each plate and demonstrated no amplification. To quantify CB₁ mRNA expression, mean C_T values were determined between reaction duplicates for each sample. Mean CB₁ C_T values were then normalized to those of an internal reference gene (GAPDH) using the $2^{-\Delta\Delta C_T}$ method (Schmittgen and Livak, 2008).

2.8 Statistical analyses

Statistical analyses were conducted using SPSS 17.0 (Chicago, IL) and JMP 12.0 (Cary, NC). Pearson correlation coefficients were calculated for all correlational data except for the correlation between fat score and combined fat mass, for which Spearman's rho was calculated instead due to the ordinal nature of fat score. Outliers were determined by Grubbs' test (GraphPad Software, La Jolla, CA) and excluded from analyses. Data from one individual for which lipid concentrations were not detectable in furcular fat were also removed from analyses, resulting in final sample sizes of 30-32. Changes in adiposity over time were quantified using a repeated-measures MANOVA. A post-hoc Welch's ANOVA was used to contrast fat scores in the last score before light advancement (Feb 24) and the final score at the end of the study (April 17). Because males and females did not differ in any outcome measure after light advancement,

data were collapsed across sex for all end-point analyses. Findings were considered significant when $p < 0.05$, and we report means $\pm$ SEM unless otherwise noted.

3. Results

3.1 Adiposity

There was a strong positive correlation between abdominal and furcular fat mass (Fig. 1A; $r = 0.936$, $p < 0.001$) and between combined fat mass (i.e., abdominal + furcular fat mass) and visual fat score (Fig. 1B; $r = 0.912$, $p < 0.001$) thereby validating the use of this measure in estimating fat levels. White-winged juncos responded to light advancement with significant reductions in adiposity (Welch's ANOVA: $F = 6.24$, $p = 0.01$; Fig. 2). Just prior to light advancement, mean fat score was 3.72 ± 0.30 and differed significantly between males and females (Repeated measures MANOVA, Date: $F = 33.05$, $p < 0.001$; Sex: $F = 7.38$, $p = 0.012$; Sex*Date: $F = 2.9256$, $p = 0.008$). After 38 d in a long-day photoperiod, mean fat score was 2.76 ± 0.16 and did not differ between sexes ($t = 1.12$, $p = 0.27$). Based on the relationship between fat score and mass (Fig. 1B), this decline in fat score amounts a loss of almost one gram of fat (average: 0.87 g), with some birds gaining up to 0.35 g and others losing as much as 1.54 g.

3.2 Lipid signals in adipose tissue

In general, concentrations of lipid signals corresponded significantly with adiposity levels but did not differ between sexes (all $p > 0.1$). Individuals with lower amounts of fat tended to have higher levels of 2-AG and OEA. Specifically, furcular 2-AG levels correlated negatively with all three measures of adiposity (i.e., furcular, abdominal, and combined fat mass) ($r = -0.442$, $p = 0.014$ for combined fat mass; Fig. 3A), and similar correlations were observed with OEA ($r = -0.462$, $p = 0.010$ for combined fat mass; Fig. 3C). In contrast, furcular PEA trended towards a

positive correlation with combined fat mass (Fig. 3D; $r = 0.322$, $p = 0.082$), whereas no associations were observed between furculum-derived AEA and any adiposity measure (Fig. 3B; $p > 0.05$). Concentrations of lipid signals in abdominal fat demonstrated somewhat similar patterns as in the furculum: abdominal 2-AG and OEA correlated negatively with all adiposity measures ($r = -0.557$, $p = 0.001$; 2-AG and combined fat mass; Fig. 4A), ($r = -0.535$, $p = 0.002$; OEA and combined fat mass; Fig. 4C) whereas abdominal AEA and PEA showed little relation to adiposity (Fig. 4B and D; $p > 0.05$). In both fat depots, 2-AG and OEA were the only compounds that correlated significantly with each other (furculum: $r = 0.371$, $p = 0.048$; abdominal fat: $r = 0.509$, $p = 0.004$; all others: $p > 0.09$, $r < |0.326|$).

3.3 CB₁ expression in hindbrain

Hindbrain CB₁ mRNA expression did not differ between sexes ($p > 0.05$) and correlated significantly and negatively with abdominal fat mass ($r = -0.364$, $p = 0.044$). A similar pattern was observed between CB₁ mRNA and furcular fat levels ($r = -0.338$, $p = 0.063$), resulting in a significant negative correlation between hindbrain CB₁ mRNA expression and overall adiposity (Fig. 5; $r = -0.357$, $p = 0.048$).

4. Discussion

Seasonally appropriate fat storage has important consequences for survival and reproductive success. Given its role in mammals to promote food intake and lipogenesis (Osei-Hyiaman et al., 2005; Pagotto et al., 2006; Vettor and Pagano, 2009), endocannabinoid signaling may be one mechanism by which rapid modulation of feeding and fattening is achieved in seasonally-breeding birds. In the present study, we used photostimulation of captive birds to induce physiological readiness to reproduce and concurrent fat reduction. We report negative

correlations between adiposity and local 2-AG and OEA concentrations, as well as with hindbrain mRNA expression of CB₁ receptors. Our results suggest that different lipid signals serve different functions that may depend on the context under which fat storage occurs.

The birds in this study exhibited highly variable and in some cases extremely pronounced fat while under winter-like photoperiods with ad libitum access to food (Fig. 2). The experimental increase in photoperiod was sufficient to trigger an overall reduction in adiposity, as well as gonadal growth and marked steroidogenic capabilities demonstrating that birds were in reproductive condition (Rosvall et al., 2013; Bergeon Burns et al., 2014; Rosvall et al., 2016). Mean fat score of birds at study end remained elevated relative to initial measures in August when birds were on natural long-day photoperiods, suggesting that birds may still have been losing fat at the time of terminal measures. At the time of collection, there remained high variation in adiposity among individuals (CV=31.9%).

Such variation in fatness revealed negative relationships between 2-AG concentrations and furcular and abdominal fat mass, indicating that fatter birds have less 2-AG content in fat depots. These findings are consistent with a previous study of seasonal adiposity in which lower 2-AG was observed in subcutaneous white adipose tissue of fat, summer-acclimated Siberian hamsters compared with lean, winter-acclimated counterparts (Ho et al., 2012). However, these findings sit in contrast to our predictions and reports of elevated 2-AG concentrations in adipose tissue of obese humans and diet-induced obese mice (Matias et al., 2006). One possible explanation for the discrepancy between studies may be the context in which adiposity occurs. In the context of seasonal adaptation, the negative relationship between endocannabinoid signaling and adiposity may reflect a drive towards fat loss or maintenance once sufficient stores are achieved. Although food intake was not measured in the present study, low 2-AG levels may

reflect an appropriate response for individuals with sufficient fat stores by inhibiting further food intake and lipogenesis, and perhaps even promoting continued fat loss in preparation for breeding. In contrast, high 2-AG levels in clinical obesity may result from dysregulation of regulatory systems that include endocannabinoid signaling. Because our findings provide only a snapshot of dynamic changes in lipid signaling, studies that compare signaling across life-history stages (such as pre- and post- migratory fattening, winter fattening, and/or breeding) are needed to determine how 2-AG concentrations relate to adiposity in natural conditions.

No correlations were observed between the other major endocannabinoid ligand, AEA, and any measure of adiposity, indicating a functional separation between endocannabinoid signals. This finding is not surprising, as dissociations between 2-AG and AEA have been reported across a variety of physiological contexts (Di Marzo and Maccarrone, 2008) and concentrations of 2-AG correlated significantly with OEA, but not AEA, in this study. Indeed, concentrations of 2-AG, but not AEA, are significantly elevated in epididymal fat of diet-induced obese mice and in visceral fat of obese humans in comparison with their respective lean controls (Matias et al., 2006). Moreover, each ligand uses a different set of enzymatic machinery (Dinh et al., 2002; Sugiura, 2008), and AEA also interacts with non-cannabinoid receptors (Zygmunt et al., 1999). Our findings suggest that as with mammals, avian 2-AG and AEA are independent from one another in form and function.

The patterns we observed in the brain mimic those seen for 2-AG in the periphery: individual juncos demonstrating greater adiposity had less hindbrain CB₁ mRNA expression. In mammals, this brain region includes the dorsal vagal complex, which sits at the interface of the gut-brain axis and integrates meal-related satiety signals (Berthoud, 2004). To the extent that mRNA abundance is predictive of receptor expression, less hindbrain CB₁ mRNA in fatter birds

is suggestive of less endocannabinoid suppression of satiety signals that emerge from this brain region, thereby promoting satiety. These results may once again reflect a drive to maintain or reduce energy intake in fat birds that have already achieved sufficient energy stores. These results stand in contrast to previous studies in Siberian hamsters that demonstrate endocannabinoid signaling fluctuates seasonally in the hindbrain, such that high summertime adiposity is associated with higher protein levels of hindbrain CB₁ and 2-AG relative to lean, winter-like states (Ho, 2011; Ho et al., 2012). Again, further studies that address how hindbrain CB₁ signaling relates to appetite and adiposity with greater temporal resolution will help clarify its role in regulating adiposity.

Because endocannabinoids are part of a family of biologically active lipids, we examined two additional lipids found in the fat depots that are structurally related to AEA and relevant to energy regulation, although with very different functions than endocannabinoids. OEA and PEA do not act at cannabinoid receptors, but instead activate peroxisome proliferator-activated receptors (PPARs) that regulate lipid metabolism. OEA attenuates body mass gains and stimulates breakdown of fat through PPAR- α (Fu et al., 2003), but its peripherally-based anorectic effects (Rodriguez de Fonseca et al., 2001) may be in part due to its ability to activate GPR119 and elicit glucagon-like peptide-1 (GLP-1) secretion (Lauffer et al., 2009; Moss et al., 2016), which itself has anorectic effects in the hindbrain (Hayes et al., 2008). Thus we predicted that OEA would function in a direction opposite to that of endocannabinoids with respect to fat regulation; instead, we found that OEA concentrations correlated negatively with adiposity, much like our 2-AG findings. Interestingly, OEA concentrations were notably higher than any other lipid examined. While it is not yet known how changes in OEA signaling alter adiposity in birds, these patterns suggest that this compound may be important and should be studied further.

PEA also activates PPAR- α (Fu et al., 2003; LoVerme et al., 2006) and inhibits food intake in rodents, albeit to a lesser extent than OEA (Rodriguez de Fonseca et al., 2001). In our study, furcular PEA correlated somewhat positively with combined fat mass, but no relationship was observed between abdominal PEA and fat mass. It is presently unclear why patterns of PEA expression contrast with those of OEA given their similar effects on appetite, but these patterns may relate to PEA's involvement in other functions such as inflammation (Solorzano et al., 2009) and analgesia (Calignano et al., 1998). Moreover, relationships between adiposity and the other lipids examined in this study were generally consistent between furcular and abdominal depots, thus the functional significance of these particular findings are unclear.

Conclusions

Considered together, our findings reveal several significant relationships between lipid signaling and adiposity that collectively suggest a role for peripheral and central endocannabinoid signaling in the regulation of fat storage in birds. The patterns we have identified lay the groundwork for future studies that are essential to understanding relationships between fat storage and lipid signaling, including those that ultimately determine how lipid signaling relates to temporal changes in adiposity. Further studies that experimentally manipulate lipid signaling, examine food intake and energy expenditure, and define temporal changes across periods of fat gain and loss will help determine the functional effects of these lipid signals and whether they causally relate to seasonal changes in adiposity. Our study is the first to report OEA and PEA concentrations in birds and demonstrates significant associations between lipid signaling and adiposity. As a consequence, these results add to the wealth of knowledge on the adaptive function of fat regulation in birds by shedding new light on the lesser known, proximate

mechanisms involved in this process. The present study takes an important step towards establishing a potential role for lipid signaling in regulating adiposity in birds.

Acknowledgments

We are grateful to the Black Hills National Forest for access to field site, and to J and C Gorsuch, M Boser, EM Schultz, and C Wood for assistance in South Dakota. B Duncan, R Kiley, MP Peterson, and R Stewart provided valuable assistance. This research was supported by: NIH F32HD068222 to KAR; NIH DA006668 to HBB; NIH T32HD049336; NSF IOS-0919911 to GED; NSF IOS-0909834 to CMBB; NSF IOS-0820055 to EDK; NSF DBI-0851607; and James P. Holland Graduate Fellowship (Indiana University) to NMR. The authors have no conflicts of interest to disclose, and funding sources had no involvement in any aspect of the study design, execution or publication.

405 **Literature cited**

- 406
- 407 Baggott, G.K., 1975. Molt, flight-muscle hypertrophy and premigratory lipid deposition of
408 juvenile willow warbler, *Phylloscopus-Trochilus*. J. Zool. 175, 299-314.
- 409 Baldwin, S.P., Kendeigh, S.C., 1938. Variations in the weight of birds. Auk 55, 416-467.
- 410 Bauer, C.M., Needham, K.B., Le, C.N., Stewart, E.C., Graham, J.L., Ketterson, E.D., Greives,
411 T.J., 2016. Hypothalamic-pituitary-adrenal axis activity is not elevated in a songbird
412 (*Junco hyemalis*) preparing for migration. Gen. Comp. Endocrinol. 232, 60-66.
- 413 Bergeon Burns, C.M., Rosvall, K.A., Hahn, T.P., Demas, G.E., Ketterson, E.D., 2014.
414 Examining sources of variation in HPG axis function among individuals and populations
415 of the dark-eyed junco. Horm. Behav. 65:179-187.
- 416 Berthold, P., 1993. Bird Migration - A General Survey. Oxford University Press, Oxford.
- 417 Berthold, P., 1996. Control of Bird Migration. Chapman and Hall, London.
- 418 Berthoud, H.R., 2004. The caudal brainstem and the control of food intake and energy balance,
419 in: Stricker, E.M., Woods, S.C. (Eds.), Neurobiology of Food and Fluid Intake, vol. 14.
420 Springer, New York, pp. 195-240.
- 421 Biebach, H., 1996. Energetics of winter and migratory fattening, in: Carey, C. (Ed.), Avian
422 energetics and nutritional ecology. Chapman and Hall, New York, pp. 280-323.
- 423 Blem, C.R., 1975. Geographic variation in wing-loading of the House Sparrow. Wilson Bull. 87,
424 543-549.
- 425 Blem, C.R., 1980. The energetics of migration, in: Gauthreaux, S.A. (Ed.), Animal Migration,
426 Orientation, and Navigation. Academic Press, London, pp. 175-224.
- 427 Boswell, T., Richardson, R.D., Seeley, R.J., Ramenofsky, M., Wingfield, J.C., Friedman, M.I.,
428 Woods, S.C., 1995. Regulation of food intake by metabolic fuels in white-crowned
429 sparrows. Am. J. Physiol. Regul. Integr. Comp. Physiol. 269, R1462-R1468.
- 430 Bradshaw, H.B., Rimmerman, N., Krey, J.F., Walker, J.M., 2006. Sex and hormonal cycle
431 differences in rat brain levels of pain-related cannabimimetic lipid mediators. Am. J.
432 Physiol. Regul. Integr. Comp. Physiol. 291, R349-R358.
- 433 Buntin, J.D., Tesch, D., 1985. Effects of intracranial prolactin administration on maintenance of
434 incubation readiness, ingestive behavior, and gonadal condition in ring doves. Horm.
435 Behav. 19, 188-203.
- 436 Calignano, A., La Rana, G., Giuffrida, A., Piomelli, D., 1998. Control of pain initiation by
437 endogenous cannabinoids. Nature 394, 277-281.
- 438 Cluny, N.L., Vemuri, V.K., Chambers, A.P., Limebeer, C.L., Bedard, H., Wood, J.T., Lutz, B.,
439 Zimmer, A., Parker, L.A., Makriyannis, A., Sharkey, K.A., 2010. A novel peripherally
440 restricted cannabinoid receptor antagonist, AM6545, reduces food intake and body
441 weight, but does not cause malaise, in rodents. Br. J. Pharmacol. 161, 629-642.
- 442 Cota, D., Marsicano, G., Tschoop, M., Grubler, Y., Flachskamm, C., Schubert, M., Auer, D.,
443 Yassouridis, A., Thone-Reineke, C., Ortmann, S., Tomassoni, F., Cervino, C., Nisoli, E.,
444 Linthorst, A.C., Pasquali, R., Lutz, B., Stalla, G.K., Pagotto, U., 2003. The endogenous
445 cannabinoid system affects energy balance via central orexigenic drive and peripheral
446 lipogenesis. J. Clin. Invest. 112, 423-431.
- 447 Dawson, A., King, V.M., Bentley, G.E., Ball, G.F., 2001. Photoperiodic control of seasonality in
448 birds. J. Biol. Rhythms 16, 365-380.

- Deviche, P.E., 1995. Androgen regulation of avian premigratory hyperphagia and fattening: from eco-physiology to neuroendocrinology. *Am. Zool.* 35, 234-245.
- Di Marzo, V., De Petrocellis, L., Bisogno, T., 2005. The biosynthesis, fate and pharmacological properties of endocannabinoids, in: Pertwee, R.G. (Ed.), *Handbook of Experimental Pharmacology*, Springer-Verlag, Heidelberg, Germany, pp. 147-185.
- Di Marzo, V., Maccarrone, M., 2008. FAAH and anandamide: is 2-AG really the odd one out? *Trends Pharmacol. Sci.* 29, 229-233.
- Dickens, M.J., Vecchiarelli, H.A., Hill, M.N., Bentley, G.E., 2015. Endocannabinoid Signaling in the stress response of male and female songbirds. *Endocrinology* 156, 4649-4659.
- Dinh, T.P., Freund, T.F., Piomelli, D., 2002. A role for monoglyceride lipase in 2-arachidonoylglycerol inactivation. *Chem. Phys. Lipids* 121, 149-158.
- eBird, 2016. An online database of bird distribution and abundance. <http://www.ebird.org/content/ebird/> (accessed 28.06.16)
- Freed, L.A., 1981. Loss of mass in breeding wrens: stress or adaptation? *Ecology* 62, 1179-1186.
- Fu, J., Gaetani, S., Oveisi, F., Lo Verme, J., Serrano, A., de Fonseca, F.R., Rosengarth, A., Luecke, H., Di Giacomo, B., Tarzia, G., Piomelli, D., 2003. Oleyethanolamide regulates feeding and body weight through activation of the nuclear receptor PPAR- α . *Nature* 425, 90-93.
- Fudickar, A.M., Greives, T.J., Atwell, J.W., Stricker, C.A., Ketterson, E.D., 2016. Reproductive allochrony in seasonally sympatric populations maintained by differential response to photoperiod: implications for population divergence and response to climate change. *Am. Nat.* 187, 436-446.
- Gray, J.M., Yarian, D., Ramenofsky, M., 1990. Corticosterone, foraging behavior, and metabolism in dark-eyed juncos, *Junco hyemalis*. *Gen. Comp. Endocrinol.* 79, 375-384.
- Gwinner, E., 1986. Circannual rhythms endogenous circadian clocks in the organization of seasonal processes. Springer, Berlin Heidelberg New York.
- Hayes, M.R., Skibicka, K.P., Grill, H.J., 2008. Caudal brainstem processing is sufficient for behavioral, sympathetic, and parasympathetic responses driven by peripheral and hindbrain glucagon-like-peptide-1 receptor stimulation. *Endocrinology* 149, 4059-4068.
- Ho, J.M., 2011. Seasonal changes in endocannabinoid signaling and energy regulation. Indiana University, Bloomington.
- Ho, J.M., Smith, N.S., Adams, S.A., Bradshaw, H.B., Demas, G.E., 2012. Photoperiodic changes in endocannabinoid levels and energetic responses to altered signalling at CB1 receptors in Siberian hamsters. *J. Neuroendocrinol.* 24, 1030-1039.
- Holberton, R.L., Boswell, T., Hunter, M.J., 2008. Circulating prolactin and corticosterone concentrations during the development of migratory condition in the dark-eyed junco, *Junco hyemalis*. *Gen. Comp. Endocrinol.* 155, 641-649.
- Holberton, R.L., Dufty, A.M., Jr., 2005. Hormone patterns and variation in life history strategies of migratory and non-migratory birds, in: Marra, P.P., Greenberg, R., (Eds.), *Birds of Two Worlds: The Ecology and Evolution of Migratory Birds*. Johns Hopkins Press, Baltimore, pp. 90-302.
- Holberton, R.L., Wilson, C.M., Hunter, M.J., Cash, W.B., Sims, C.G., 2007. The role of corticosterone in supporting migratory lipogenesis in the dark-eyed junco, *Junco hyemalis*: a model for central and peripheral regulation. *Physiol. Biochem. Zool.* 80, 125-137.

494 Jawor J.M., McGlothlin, J.W., Casto, J.M., Greives, T.J., Snajdr, E.A., Bentley, G.E., Ketterson,
 495 E.D., 2006. Seasonal and individual variation in response to GnRH challenge in male
 496 dark-eyed juncos (*Junco hyemalis*). Gen. Comp. Endocrinol. 149, 182-189.
 497 Jbilo, O., Ravinet-Trillou, C., Arnone, M., Buisson, I., Bribes, E., Peleraux, A., Penarier, G.,
 498 Soubrie, P., Le Fur, G., Galiegue, S., Casellas, P., 2005. The CB1 receptor antagonist
 499 rimonabant reverses the diet-induced obesity phenotype through the regulation of
 500 lipolysis and energy balance. FASEB J 19, 1567-1569.
 501 Jo, Y.H., Chen, Y.J., Chua, S.C., Jr., Talmage, D.A., Role, L.W., 2005. Integration of
 502 endocannabinoid and leptin signaling in an appetite-related neural circuit. Neuron 48,
 503 1055-1066.
 504 King, J.R., 1972. Adaptive periodic fat storage by birds. Proc. Int. Ornithol. Congr. 15, 200-217.
 505 King, J.R., Barker, S., Farner, D.S., 1963. A comparison of energy reserves during autumnal and
 506 vernal migratory periods in white-crowned sparrow, *Zonotrichia leucophrys gambelii*.
 507 Ecology 44, 513-521.
 508 Lauffer, L.M., Iakoubov, R., Brubaker, P.L., 2009. GPR119 is essential for oleoylethanolamide-
 509 induced glucagon-like peptide-1 secretion from the intestinal enteroendocrine L-cell.
 510 Diabetes 58, 1058-1066.
 511 Lima, S.L., 1986. Predation risk and unpredictable feeding conditions: determinants of body
 512 mass in birds. Ecology 67, 366-376.
 513 Lofts, B., Marshall, A.J., 1961. Zugunruhe activity in castrated bramblings (*Fringilla*
 514 *montifringilla*). Ibis 103, 189-194.
 515 LoVerme, J., Russo, R., La Rana, G., Fu, J., Farthing, J., Mattace-Raso, G., Meli, R., Hohmann,
 516 A., Calignano, A., Piomelli, D., 2006. Rapid broad-spectrum analgesia through activation
 517 of peroxisome proliferator-activated receptor- α . J. Pharm. Exp. Ther. 319, 1051-
 518 1061.
 519 Matias, I., Gonthier, M.P., Orlando, P., Martiadis, V., De Petrocellis, L., Cervino, C., Petrosino,
 520 S., Hoareau, L., Festy, F., Pasquali, R., Roche, R., Maj, M., Pagotto, U., Monteleone, P.,
 521 Di Marzo, V., 2006. Regulation, function, and dysregulation of endocannabinoids in
 522 models of adipose and beta-pancreatic cells and in obesity and hyperglycemia. J. Clin.
 523 Endocrinol. Metab. 91, 3171-3180.
 524 Meier, A.H., Farner, D.S., 1964. A possible endocrine basis for premigratory fattening in the
 525 white-crowned sparrow, *Zonotrichia leucophrys gambelii* (Nuttall). Gen. Comp.
 526 Endocrinol. 4, 584-595.
 527 Morton, M.L., Mewalt, L.R., 1962. Some effects of castration on a migratory sparrow
 528 (*Zonotrichia atricapilla*). Physiol. Zool. 35, 237-247.
 529 Moss, C.E., Glass, L.L., Diakogiannaki, E., Pais, R., Lenaghan, C., Smith, D.M., Wedin, M.,
 530 Bohlooly, Y.M., Gribble, F.M., Reimann, F., 2016. Lipid derivatives activate GPR119
 531 and trigger GLP-1 secretion in primary murine L-cells. Peptides 77, 16-20.
 532 Nolan, V., Ketterson, E.D., 1983. An analysis of body mass, wing length, and visible fat deposits
 533 of dark-eyed juncos wintering at different latitudes. Wilson Bull. 95, 603-620.
 534 Novoseletsky, N., Nussinovitch, A., Friedman-Einat, M., 2011. Attenuation of food intake in
 535 chicks by an inverse agonist of cannabinoid receptor 1 administered by either injection or
 536 ingestion in hydrocolloid carriers. Gen. Comp. Endocrinol. 170, 522-527.
 537 Osei-Hyiaman, D., DePetrillo, M., Pacher, P., Liu, J., Radaeva, S., Batkai, S., Harvey-White, J.,
 538 Mackie, K., Offertaler, L., Wang, L., Kunos, G., 2005. Endocannabinoid activation at

- hepatic CB1 receptors stimulates fatty acid synthesis and contributes to diet-induced obesity. *J. Clin. Invest.* 115, 1298-1305.
- Pagano, C., Pilon, C., Calcagno, A., Urbanet, R., Rossato, M., Milan, G., Bianchi, K., Rizzuto, R., Bernante, P., Federspil, G., Vettor, R., 2007. The endogenous cannabinoid system stimulates glucose uptake in human fat cells via phosphatidylinositol 3-kinase and calcium-dependent mechanisms. *J. Clin. Endocrinol. Metab.* 92, 4810-4819.
- Pagotto, U., Marsicano, G., Cota, D., Lutz, B., Pasquali, R., 2006. The emerging role of the endocannabinoid system in endocrine regulation and energy balance. *Endocr. Rev.* 27, 73-100.
- Piersma, T., Jukema, J., 1990. Budgeting the flight of a long-distance migrant - changes in nutrient reserve levels of bar-tailed godwits at successive spring staging sites. *Ardea* 78, 315-337.
- Quarta, C., Bellocchio, L., Mancini, G., Mazza, R., Cervino, C., Bräulke, L.J., Fekete, C., Latorre, R., Nanni, C., Bucci, M., Clemens, L.E., Heldmaier, G., Watanabe, M., Leste-Lassere, T., Maitre, M., Tedesco, L., Fanelli, F., Reuss, S., Klaus, S., Srivastava, R.K., Monory, K., Valerio, A., Grandis, A., De Giorgio, R., Pasquali, R., Nisoli, E., Cota, D., Lutz, B., Marsicano, G., Pagotto, U., 2010. CB(1) signaling in forebrain and sympathetic neurons is a key determinant of endocannabinoid actions on energy balance. *Cell. Metab.* 11, 273-285.
- Ramakrishnan, S., Strader, A.D., Wimpee, B., Chen, P., Smith, M.S., Buntin, J.D., 2007. Evidence for increased neuropeptide Y synthesis in mediobasal hypothalamus in relation to parental hyperphagia and gonadal activation in breeding ring doves. *J. Neuroendocrinol.* 19, 163-171.
- Ramenofsky, M., Savard, R., Greenwood, M.R.C., 1999. Seasonal and diel transitions in physiology and behavior in the migratory dark-eyed junco. *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* 122, 385-397.
- Rodriguez de Fonseca, F., Navarro, M., Gomez, R., Escuredo, L., Nava, F., Fu, J., Murillo-Rodriguez, E., Giuffrida, A., LoVerme, J., Gaetani, S., Kathuria, S., Gall, C., Piomelli, D., 2001. An anorexic lipid mediator regulated by feeding. *Nature* 414, 209-212.
- Rogers, C.M., 1995. Experimental evidence for temperature-dependent winter lipid storage in the dark-eyed junco (*Junco hyemalis oreganus*) and song sparrow (*Melospiza melodia morphna*). *Physiol. Zool.* 68, 277-289.
- Rogers, C.M., 1995. Testing optimal body mass theory: Evidence for cost of fat in wintering birds. *Ecosphere* 6(4): 55.
- Rogers, C.M., Ramenofsky, M., Ketterson, E.D., Nolan, V., Wingfield, J.C., 1993. Plasma corticosterone, adrenal mass, winter weather, and season in nonbreeding populations of dark-eyed juncos (*Junco hyemalis hyemalis*). *Auk* 110, 279-285.
- Rogers, C.M., Smith, J.N.M., 1993. Life-history theory in the nonbreeding period: trade-offs in avian fat reserves? *Ecology* 74, 419-426.
- Rosvall, K.A., Bergeon Burns, C.M., Hahn, T.P., Ketterson, E.D., 2013. Sources of variation in HPG axis reactivity and individually consistent elevation of sex steroids in a female songbird. *Gen. Comp. Endocrinol.* 194, 230-239.
- Rosvall, K.A., Bergeon Burns, C.M., Jayaratna, S.P., Ketterson, E.D., 2016. Divergence along the gonadal steroidogenic pathway: Implications for hormone-mediated phenotypic evolution. *Horm. Behav.* 84, 1-8.

584 Schmittgen, T.D., Livak, K.J., 2008. Analyzing real-time PCR data by the comparative C(T)
 585 method. Nat. Protoc. 3, 1101-1108.
 586 Schwabl, H., Farner, D.S., 1989. Endocrine and environmental-control of vernal migration in
 587 male white-crowned sparrows, *Zonotrichia-leucophrys-gambelii*. Physiol. Zool. 62, 1-10.
 588 Solorzano, C., Zhu, C., Battista, N., Astarita, G., Lodola, A., Rivara, S., Mor, M., Russo, R.,
 589 Maccarrone, M., Antonietti, F., Duranti, A., Tontini, A., Cuzzocrea, S., Tarzia, G.,
 590 Piomelli, D., 2009. Selective N-acylethanolamine-hydrolyzing acid amidase inhibition
 591 reveals a key role for endogenous palmitoylethanolamide in inflammation. Proc. Natl.
 592 Acad. Sci. USA 106, 20966-20971.
 593 Soma, K.K., Bindra, R.K., Gee, J., Wingfield, J.C., Schlinger, B.A., 1999. Androgen-
 594 metabolizing enzymes show region-specific changes across the breeding season in the
 595 brain of a wild songbird. J. Neurobiol. 41, 176-188.
 596 Stetson, M.H., Erickson, J.E., 1972. Hormonal-control of photoperiodically induced fat
 597 deposition in white-crowned sparrows. Gen. Comp. Endocrinol. 19, 355-362.
 598 Sugiura, T., 2008. Biosynthesis of anandamide and 2-arachidonylglycerol, in: Köfalvi, A., (Ed.)
 599 Cannabinoids and the Brain. Springer Science + Business Media LLC, Coimbra, Portugal
 600 pp. 15-30.
 601 Tam, J., Vemuri, V.K., Liu, J., Batkai, S., Mukhopadhyay, B., Godlewski, G., Osei-Hyiaman, D.,
 602 Ohnuma, S., Ambudkar, S.V., Pickel, J., Makriyannis, A., Kunos, G., 2010. Peripheral
 603 CB1 cannabinoid receptor blockade improves cardiometabolic risk in mouse models of
 604 obesity. J. Clin. Invest. 120, 2953-2966.
 605 Totzke, U., Hubinger, A., Bairlein, F., 1997. A role for pancreatic hormones in the regulation of
 606 autumnal fat deposition of the garden warbler (*Sylvia borin*)? Gen. Comp. Endocrinol.
 607 107, 166-171.
 608 Vettor, R., Pagano, C., 2009. The role of the endocannabinoid system in lipogenesis and fatty
 609 acid metabolism. Best Pract. Res. Clin. Endocrinol. Metab. 23, 51-63.
 610 Whitney, N.R., 1968. White-winged junco, in: Austin, O.L. (Ed.), Life Histories of North
 611 American Cardinals, Grosbeaks, Buntings, Towhees, Finches, Sparrows, and Their Allies
 612 Pt 2 (Vol. 237) U.S. Natl. Mus. Bull., pp. 1021-1029.
 613 Williams, T.D., 2012. Physiological Adaptations for Breeding in Birds. Princeton University
 614 Press, Princeton, NJ.
 615 Witter, M.S., Swaddle, J.P., Cuthill, I.C., 1995. Periodic food availability and strategic regulation
 616 of body-mass in the European starling, *Sturnus-vulgaris*. Funct. Ecol. 9, 568-574.
 617 Zygmunt, P.M., Petersson, J., Andersson, D.A., Chuang, H., Sorgard, M., Di Marzo, V., Julius,
 618 D., Hogestatt, E.D., 1999. Vanilloid receptors on sensory nerves mediate the vasodilator
 619 action of anandamide. Nature 400, 452-457.

Table 1. Criteria used to assign fat scores. *some scores were assigned an additional + or – 0.25 for greater accuracy

Fat score*	Criterion
1	Fat present in abdomen or furculum (one place)
2	Fat present in both abdomen and furculum
3	Fat flush with body curvature in abdomen or furculum (one place)
4	Fat bulging in abdomen or furculum (one place)
5	Fat bulging in both abdomen and furculum

Figure legends

Figure 1. Relationships between visual fat score and combined mass of abdominal and furcular fat depots (A) and between abdominal and furcular fat mass (B). $p < 0.05$, Pearson's correlation coefficient and Spearman's rho.

Figure 2. Relationship between adiposity and date, as birds gain fat over-winter and lose fat as they enter breeding condition. Error bars represent one standard error from the mean. The shaded bar highlights the period of light advancement, and the dashed line indicates when birds reached a long-day photoperiod (16L:8D). Although the sexes differed in overwinter fat levels, males (gray) and females (black), they did not differ after light advancement. Repeated measures MANOVA, Date: $F = 33.05$, $p < 0.001$; Sex: $F = 7.38$, $p = 0.012$; Sex*Date: $F = 2.926$, $p = 0.008$. Post-hoc t -test compared sexes in April: $t = 1.12$, $p = 0.27$.

Figure 3. Relationships between combined abdominal and furcular fat mass and furcular levels of 2-arachidonoylglycerol (2-AG) (A), anandamide (AEA) (B), oleoylethanolamide (OEA) (C), and palmitoylethanolamide (PEA) (D). Pearson's correlation coefficient.

Figure 4. Relationships between combined abdominal and furcular fat mass and abdominal levels of 2-arachidonoylglycerol (2-AG) (A), anandamide (AEA) (B), oleoylethanolamide (OEA) (C), and palmitoylethanolamide (PEA) (D). Pearson's correlation coefficient.

Figure 5. Relationship between combined abdominal and furcular fat mass and hindbrain CB₁ mRNA expression levels. Expression levels were calculated relative to those of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) and normalized to calibrator values. $p < 0.05$, Pearson's correlation coefficient.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5