Interaction between L-aspartate and the brucite $[\text{Mg}(\text{OH})_2]$ -water interface

Charlene F. Estrada^{a,b,*}, Dimitri A. Sverjensky^a, Manuel Pelletier^{c,d}, Angéline Razafitianamaharavo^{c,d}, and Robert M. Hazen^b

^a Johns Hopkins University, Dept. of Earth & Planetary Sciences, 3400 N Charles St, Baltimore MD, 21218 USA

^b Geophysical Laboratory, Carnegie Institution of Washington, Broad Branch Rd NW, Washington DC 20015 USA

^c Université de Lorraine, Laboratoire Interdisciplinaire des Environnements Continentaux, 15 avenue du Charmois, 54500 Vandœuvre les Nancy, France

^d CNRS, UMR 7360, Laboratoire Interdisciplinaire des Environnements Continentaux, 15 avenue du Charmois, 54500 Vandœuvre les Nancy, France.

*Corresponding author at: Geophysical Laboratory, Carnegie Institution of Washington, 5251 Broad Branch Rd NW, Washington DC, 20015, USA. Tel: +1 202 478 8900

Email address: cestrada@jhu.edu (C. Estrada).

Abstract

The interaction of biomolecules at the mineral-water interface could have played a prominent role in the emergence of more complex organic species in life's origins. Serpentine-hosted hydrothermal vents may have acted as a suitable environment for this process to occur, although little is known about biomolecule-mineral interactions in this system. We used batch adsorption experiments and surface complexation modeling to study the interaction of L-aspartate onto a thermodynamically stable product of serpentinization, brucite $[\text{Mg}(\text{OH})_2]$, over a wide range of initial aspartate concentrations at four ionic strengths governed by $[\text{Mg}^{2+}]$ and $[\text{Ca}^{2+}]$. We observed that up to $1.0 \mu\text{mol}$ of aspartate adsorbed per m^2 of brucite at $\text{pH} \sim 10.2$ and low Mg^{2+} concentrations ($0.7 \times 10^{-3} \text{ M}$), but surface adsorption decreased at high Mg^{2+} concentrations ($5.8 \times 10^{-3} \text{ M}$). At high Ca^{2+} concentrations ($4.0 \times 10^{-3} \text{ M}$), aspartate surface adsorption doubled (to $2.0 \mu\text{mol} \cdot \text{m}^{-2}$), with Ca^{2+} adsorption at $29.6 \mu\text{mol} \cdot \text{m}^{-2}$. We used the extended triple-layer model (ETLM) to construct a quantitative thermodynamic model of the adsorption data. We proposed three surface reactions involving the adsorption of aspartate (HAsp^-) and/or Ca^{2+} onto brucite:

We used the ETLM to predict that brucite particle surface charge becomes more negative with increasing $[\text{Mg}^{2+}]$, creating an unfavorable electrostatic environment for a negatively-charged aspartate molecule to adsorb. In contrast, our addition of Ca^{2+} to the system resulted in Ca^{2+} adsorption and development of positive surface charge. Our prediction of surface speciation of aspartate on brucite with Ca^{2+} revealed that the calcium-aspartate complex is the predominant surface aspartate species, which suggests that the increase in aspartate adsorption with Ca^{2+} is primarily driven by calcium adsorption. The cooperative effect of Ca^{2+} and the inhibitive effect of Mg^{2+} on aspartate adsorption onto brucite indicate that serpentine-hosted hydrothermal fluids provide an ideal environment for these interactions to take place.

1. Introduction

The origin of life on early Earth was a geochemical sequence of emergent steps involving increasing complexity (Hazen, 2005). Among the first of these steps was the synthesis of prebiotic organic molecules in the midst of the Archean atmosphere, ocean, and lithologies. The emergence of simple molecules such as amino acids and sugars may have occurred under a variety of plausible Archean environments (Ponnamperuma and Pering, 1966; Deamer and Pashley, 1989; Hennet et al., 1992; Holm and Andersson, 2005; Jalbout et al., 2007; Cleaves et al., 2008; Kim et al., 2011). Far less is known about the selection, concentration, and self-organization of these biomolecules into macromolecular systems from dilute solutions.

Mineral surfaces may provide the means by which simple biomolecules self-organize. The mineral-water interface is known to concentrate biomolecules through adsorption (Jonsson et al., 2009; Jonsson et al., 2010; Cleaves et al., 2012). Mineral surfaces change significantly with shifts in pH owing to surface protonation reactions. Accordingly, mineral surfaces may have potentially played a key role in the origin of life, whereby the mineral-water interface provided sites for the selection, concentration, and self-organization of biomolecules (Goldschmidt, 1952; Hazen, 2006; Cleaves et al., 2012).

Hydrothermal vent systems have been discussed as plausible environments for the origin of life by providing templates for macromolecular assembly (Baross and Hoffman, 1985; Shock and Canovas, 2010; Sleep et al., 2011). The disequilibria established by mixing hot, hydrothermal fluids (<350 °C) with the overlying cold seawater (2 °C) may drive the formation of some prebiotic molecules (Shock and Schulte, 1998; Shock and Canovas, 2010). A different type of hydrothermal system known as the Lost City hydrothermal field was discovered off the Mid-Ocean Ridge Axis (Kelley et al., 2001). The Lost City hydrothermal field belongs to a class

of hydrothermal systems that undergo serpentinization, which involves the hydration of magnesium-rich olivine to serpentine and ferro-brucite (Bach et al., 2006), according to the reaction:

Serpentinite-hosted hydrothermal vents such as the Lost City hydrothermal field release fluids that contain up to 19×10^{-3} M and 4×10^{-3} M of H_2 and CH_4 , respectively (Kelley et al., 2005; Proskurowski et al., 2006). The mixing of these hydrothermal fluids with seawater creates a strong redox gradient that may facilitate the formation of prebiotic molecules (Holm et al., 2006; Shock and Canovas, 2010). Furthermore, greenstone sequences discovered at the Isua Formation (3.8 Ga) and the Pilbara Formation (3.5 Ga) suggest that serpentinite-hosted hydrothermal activity may have been present on early Earth (Dymek et al., 1988; Van Kranendonk et al., 2007; Shibuya et al., 2010). These localities and the presence of Archean komatiites provide evidence that the oceanic crust on early Earth was more ultramafic compared to today's mafic, basalt-hosted seafloor (Herzberg et al., 2010; Van Kranendonk, 2010). Depending on the prevalence of the ultramafic host rock, serpentinite-hosted hydrothermal fields may have been prolific on the Archean seafloor prior to the emergence of the first cellular lifeform at 3.48 Ga (Noffke et al. 2013).

Ferro-brucite [$\text{Mg}_{0.75}\text{Fe}_{0.25}(\text{OH})_2$] is a thermodynamically stable mineral of the serpentinization reaction in Eq. (1). The ferrous iron in ferro-brucite controls the amount of $\text{H}_{2(\text{aq})}$ produced in the hydrothermal field as magnetite is subsequently precipitated:

As this reaction proceeds to the right, the amount of $\text{H}_{2(\text{aq})}$ flowing through the system becomes limited, and the brucite in the serpentinite-hosted hydrothermal field approaches the magnesium

endmember (Bach et al., 2006; McCollom and Bach, 2009). Holm and coworkers (2006; 2012) have suggested that magnesium plays a role in stabilizing prebiotic biomolecules such as proteins and nucleic acids. As the primary constituent of brucite, magnesium may promote the formation of more complex molecules at the mineral-water interface within a serpentinite-hosted hydrothermal system.

Previous studies of brucite surface chemistry have focused on its precipitation and dissolution kinetics (Hostetler, 1963; Schott, 1981; Jordan and Rammensee, 1996; Pokrovsky and Schott, 2004; Pokrovsky et al., 2005; Kudoh et al., 2006). Nevertheless, few studies exist on the adsorption of organic species onto the brucite surface over a wide range of conditions. Pokrovsky and coworkers (2005) have studied the effect of oxalate, citrate, glycine, xylose, and catechol on brucite dissolution and precipitation. These biomolecules affected brucite dissolution at concentrations greater than 1×10^{-2} M and at neutral to weakly alkaline pH conditions. Citrate adsorbed in amounts as high as $0.4 \mu\text{mol} \cdot \text{m}^{-2}$, and it and the other biomolecules may have attached onto the brucite surface as an inner-sphere species (Pokrovsky et al., 2005). Pokrovsky and coworkers (2005) also observed that concentrations of at least 1×10^{-2} M Ca^{2+} increase the brucite dissolution rate at pH= 4.9, and they assumed that calcium might adsorb onto brucite as a bidentate or monodentate mononuclear surface species.

In this investigation, we characterized the adsorption of the amino acid L-aspartate ($\text{C}_4\text{H}_7\text{NO}_4$) onto powdered brucite. Aspartate is a polar amino acid with three protonation sites: two carboxyl groups and one amine group. Hennet and coworkers (1992) synthesized aspartate under mineral-buffered, simulated hydrothermal conditions. The adsorption of aspartate has been previously well-characterized on rutile (Giacomelli et al., 1995; Roddick-Lanzilotta and McQuillan, 2000; Jonsson et al., 2010), calcite (Hazen et al., 2001), kaolinite (Ikhsan et al.,

2004), and alumina (Greiner et al., 2014). However, there are no studies on the adsorption of aspartate on brucite under the effect of variable divalent cation concentration, molecular concentration, and/or pH conditions.

The aim of this study is to evaluate the potential role that minerals within serpentinite-hosted hydrothermal fields play in concentrating and selecting prebiotic molecules from a dilute aqueous environment. In this framework, we characterized the extent and possible importance of this interaction with a combination of batch adsorption experiments at 25 °C and 1 bar and surface complexation modeling. Furthermore, we investigated the effects that common dissolved ions, including Mg^{2+} and Ca^{2+} , have on the interaction between aspartate and the brucite-water interface so that we may predict mineral surface reactions in plausible prebiotic aqueous environments.

2. Materials and Methods

2.1. Brucite Synthesis

We synthesized a microcrystalline brucite powder with a precipitation method previously reported by Henrist and coworkers (2003) and Lu and coworkers (2004). With 18.2 MΩcm Milli-Q water (Millipore), we prepared a 2.0 M $MgCl_2$ solution (99.99 % Alfa Aesar dry ampoules under Ar gas) and a solution containing 2.0 M NH_4OH (Sigma-Aldrich, 25 wt. % solution). We added the $MgCl_2$ dropwise to the NH_4OH solution at a rate of approximately 2.5 mL•min⁻¹ using a graduated buret while we kept the latter solution vigorously stirred at a constant temperature of 45 ± 3 °C. Lu and colleagues (2004) have proposed that the precipitation reaction

occurred during the addition of MgCl_2 to NH_4OH . After brucite precipitated, we set the sealed reaction vessel aside to cool at room temperature for 3 days. We thoroughly washed the soluble NH_4Cl byproduct from the precipitate with Milli-Q water. We then dried the precipitate on a Corning PC-420D hot plate at 135 °C for 1 hour.

2.1.1 Hydrothermal Treatment

Henrist and coworkers (2003) and Lu and colleagues (2004) observed that the crystallinity and morphology of synthetic brucite improved when it was sealed with water in a reaction vessel heated above 100 °C over an extended period of time. We carried out a hydrothermal aging treatment on our synthetic brucite by sealing an aqueous suspension of the synthetic brucite in Milli-Q water within a PTFE-lined stainless steel reaction vessel (Col-Int-Tech). We heated the precipitate in a furnace at 150 °C for 3 days, and after cooling the brucite-water solution to room temperature, we dried the powder at 135 °C with a hotplate. We then stored the hydrothermally-treated brucite powder in a glass vial under an Ar gas atmosphere at room temperature.

SEM imaging (JEOL 8500F) revealed that the untreated brucite has a crystal habit characterized by rosettes of rough, circular, and oblong platelets (Fig. 1a), whereas the hydrothermally-treated brucite is distinguished by clusters of lamellar, hexagonal platelets (Fig. 1b). Additionally, the powder XRD (Bruker D2 Phaser) pattern of the hydrothermally-treated brucite demonstrates higher intensity peaks in the case of reflection (001) and reduced intensity for the (100), (101) and (111) reflections (Fig. 2). This pattern is consistent with powder XRD patterns collected under ambient conditions that have been archived at the International Centre for Diffraction Data (<http://icdd.com>). Our SEM and powder XRD observations may be

interpreted as evidence that the hydrothermal treatment process enhanced the crystallinity in the basal direction by reducing terraces and kink sites and improving hexagonal morphology.

2.2. *Brucite Specific Surface Area*

We characterized the surface area of the synthetic brucite powder with the multi-point BET N₂ adsorption method (Micromeritics, Norcross, GA, USA) to obtain a specific surface area of $17.6 \pm 0.1 \text{ m}^2 \cdot \text{g}^{-1}$. A high proportion of this measured surface area most likely represents the prominent (001) basal surface, although the distribution between the basal and lateral (edge) surfaces cannot be determined with the classical N₂ gas adsorption method. This distinction is important because previous adsorption studies involving mineral surfaces similar to brucite, such as gibbsite, goethite, illite, and kaolinite, conclude that only the lateral surfaces are active in the protonation and deprotonation reactions that lead to surface adsorption (Prélot et al., 2003; Sayed-Hassan et al., 2006; Liu et al., 2013). If the lateral surfaces primarily adsorb aspartate, it is necessary to determine the surface area specific to these edge surfaces.

We assayed the edge-specific surface area (ESA) with low-pressure Ar gas adsorption at the Laboratoire Interdisciplinaire des Environnements Continentaux (Vandœuvre les Nancy, France). Consistent basal- and edge-specific surface areas were previously obtained with this technique for several lamellar minerals including talc, kaolinite, illite, lamellar silica, saponite, and smectite clays (Villieras et al., 2002; Michot and Villieras, 2002; Eypert-Blaison et al., 2002; Tournassat et al., 2003; Sayed-Hassan et al., 2005; Sayed-Hassan et al., 2006; Perronnet et al., 2007). Other types of minerals have also been investigated with this method, including goethite, anatase, and manganese dioxide (Prélot et al., 2003; Stevanovic et al., 2010; Ali Ahmad et al., 2012).

We determined a total surface area of $19.8 \pm 0.2 \text{ m}^2 \cdot \text{g}^{-1}$ with low-pressure Ar gas adsorption, which is comparable to the BET determined by N₂ gas ($17.6 \pm 0.1 \text{ m}^2 \cdot \text{g}^{-1}$). Using the methods detailed in previous studies (Villieras et al., 1992; Villieras et al., 1997; Villieras et al., 2002; Tournassat et al., 2003; Sayed-Hassan et al., 2006), we have estimated that 89.5 % of the surface area can be represented by basal brucite surfaces, whereas 10.5 % are represented by lateral brucite surfaces (see Appendix A). We adopted a basal surface area of $15.7 \text{ m}^2 \cdot \text{g}^{-1}$ and a lateral surface area of $1.9 \text{ m}^2 \cdot \text{g}^{-1}$ by assuming that the total surface area was $17.6 \text{ m}^2 \cdot \text{g}^{-1}$. We tested these surface areas by deriving the brucite particle thickness (t) and length (l) with the equations

$$l = \frac{4}{\rho \cdot S_l} \quad (4)$$

and

$$t = \frac{2}{\rho \cdot S_b} \quad (5)$$

where ρ is the density of brucite ($2.39 \text{ g} \cdot \text{cm}^{-3}$) and S_l and S_b are the lateral and basal surface areas, respectively. We calculated an average brucite plate length of $0.9 \text{ } \mu\text{m}$ and thickness of 53 nm , which were consistent with the dimensions we observed with SEM ($l = 0.1$ to $1 \text{ } \mu\text{m}$ and $t = 30$ to 50 nm , see Fig. 1b). We therefore consider $1.9 \text{ m}^2 \cdot \text{g}^{-1}$ to accurately represent the ESA of the lateral surface of our brucite sample.

2.3. Batch Adsorption Experiments

In a single batch sample, we added between 2 and $300 \text{ } \mu\text{M}$ aspartate to $10.0 \pm 0.5 \text{ g} \cdot \text{L}^{-1}$ brucite in a 15 mL sterile Falcon tube. The pH of each sample (10.2 to 10.3 without added MgCl_2) was strongly buffered by the brucite powder. We studied four ionic strengths that resulted from the dissolution of Mg^{2+} from brucite and our addition of either MgCl_2 or CaCl_2 to

the aqueous phase. We conducted low-Mg²⁺, high-Mg²⁺, low-Ca²⁺, and high-Ca²⁺ experiments, which we describe in Table 1.

We conducted preliminary adsorption experiments over multiple periods of time and determined that aspartate adsorption on brucite reaches a steady state within 16 hours. In this study, we placed the batch samples on a test tube rotator at 40 rpm (25 ± 1 °C, 1 bar) for 22 hours to ensure that the brucite-aspartate suspensions had reached a steady state. We then measured the pH of the brucite-aspartate suspensions with a combination glass electrode (Thermo-Electron Orion 8103BNUWP) that we calibrated prior to the experiment with NIST-standardized buffers that referred to pH values of 4, 7, and 10 (Fischer Scientific). As we obtained pH measurements, we flushed each sample with Ar gas to avoid contamination by CO₂. We centrifuged the samples for 10 minutes with a relative centrifugal force of 1073g. In each sample, we then separated the liquid supernatant from the solid brucite.

We analyzed the concentration of the aspartate, Mg²⁺ and Ca²⁺ in the supernatant with ion chromatography (IC). We used a Dionex ICS-5000 DP dual pump system equipped with an AminoPac10 column, IonPac CS12A column, and Chromeleon 6.8 software (Dionex Corporation, Sunnyvale, CA, USA). We analyzed the samples in triplicate and determined the mean concentration of aspartate remaining in the liquid supernatant within ±1 standard error (σ_M). It was necessary to analyze the aspartate samples in triplicate because we initially determined that low amounts of aspartate adsorbed onto the brucite surface (<10 %), and collecting data in triplicate better constrained accurate aspartate adsorption values. It was not necessary for us to analyze Ca²⁺ adsorption data in triplicate because we initially observed that Ca²⁺ adsorbed onto brucite in large amounts (>10 %), and we calculated that ±1 standard deviation (σ) was equal to about 8 % of the adsorption values. From the analytical concentrations

we determined the surface adsorption, (Γ_{ads} , $\mu\text{mol}\cdot\text{m}^{-2}$) of aspartate or Ca^{2+} on brucite, according to

$$\Gamma_{\text{ads}} = \frac{[\text{X}]_0 - [\text{X}]_{\text{eq}}}{C_s A_s} \quad (6)$$

in which $[\text{X}]_0$ is the initial concentration of aspartate or Ca^{2+} , $[\text{X}]_{\text{eq}}$ is the concentration of aspartate or Ca^{2+} remaining in the liquid supernatant, C_s is the solid concentration, and A_s is the edge-specific surface area of brucite. We report Γ_{ads} as an isotherm as a function of $[\text{X}]_{\text{eq}}$.

2.4. Surface Complexation Modeling Approach

We modeled the data from our batch adsorption experiments with the extended triple-layer model or ETLM (Sverjensky and Sahai, 1996; Sahai and Sverjensky, 1997; Sverjensky, 2003; Sverjensky, 2005; Sverjensky and Fukushima, 2006). Unlike other solid-water interface models, the ETLM includes the electrical work resulting from the release of water dipoles during inner-sphere adsorption. We iteratively fit the adsorption data in a manner described in previous studies (Bahri et al., 2011; Jonsson et al., 2009) and obtained reaction stoichiometries that represent our best fit to the adsorption data. When it has been possible to test such surface complexation modeling with spectroscopy and quantum chemical modeling (Parikh et al., 2011; Lee et al., 2012), there has been excellent agreement as to the nature of the adsorbed species. We carried out calculations for the ETLM with the computer code GEOSURF (Sahai and Sverjensky, 1998). We estimated a $\text{pH}_{\text{PZC}} = 10.5$ from measurements of the pH after only aspartate or water had been added to brucite during our batch adsorption experiments. We previously calculated speciation reactions involving aqueous aspartate and found them to be consistent with those determined with potentiometric titrations (De Robertis et al., 1991; Jonsson et al., 2010). We obtained both protonation and electrolyte

adsorption parameters for the brucite surface from a regression of surface titration data measured by Pokrovsky and Schott (2004). As illustrated by Fig. 3, we fit these data with the ETLM using parameters that were derived from a constant capacitance model ($C = 500 \mu\text{F}\cdot\text{cm}^{-2}$), where $\text{pH}_{\text{PZC}} = 11.0$, site density (N_s) = $10 \text{ sites}\cdot\text{nm}^{-2}$, and BET surface area (A_s) = $9.2 \text{ m}^2\cdot\text{g}^{-1}$ were reported for the titration study at $1.0 \times 10^{-2} \text{ M NaCl}$ and 1.0 M NaCl . We recalculated the protonation and electrolyte adsorption parameters for our estimated pH_{PZC} , lateral A_s , and N_s using the methods detailed by Sverjensky (2003; 2005). These parameters are displayed in Table 2.

Site densities can be estimated from the number of broken bonds at the surface, and these estimations are usually consistent with measurements by tritium exchange experiments (Koretsky et al., 1998). Accordingly, we calculated site densities for the lateral brucite surfaces (100), (010), (110), and (111) by examining the broken bonds of each idealized surface with the program CrystalMaker (Palmer, 2009). We estimated a minimum possible site density of $9.7 \text{ sites}\cdot\text{nm}^{-2}$ and a maximum possible site density of $20.0 \text{ sites}\cdot\text{nm}^{-2}$ (Table 3). However, to fit both the Ca^{2+} and aspartate adsorption data we observed for the low- Ca^{2+} and high- Ca^{2+} experiments, we required a site density of $38 \text{ sites}\cdot\text{nm}^{-2}$. This value is greater than what we estimated for the idealized surfaces, but it is consistent with site densities estimated ($40.8 \text{ sites}\cdot\text{nm}^{-2}$) and measured ($36 \text{ sites}\cdot\text{nm}^{-2}$) on periclase [MgO], a mineral that is structurally similar to brucite and may well be covered by a thin layer of brucite in water (Anderson et al., 1965; Koretsky et al., 1998). We can also use our measurements of aspartate and Ca^{2+} adsorption to estimate site density (e.g., $1.0 \mu\text{mol}\cdot\text{m}^{-2} = 0.6 \text{ sites}\cdot\text{nm}^{-2}$). However, due to possible intermolecular forces between surface species, it may be unlikely that every active site within one nm^2 would interact with Ca^{2+} or aspartate. It may also be possible that surface imperfections primarily caused by the

dissolution of Mg^{2+} from brucite in the aqueous phase explain why it is necessary for us to adopt a higher site density. Our estimated site density of $38 \text{ sites}\cdot\text{nm}^{-2}$ could be tested by future adsorption studies involving a variety of chemically different ions as probes of the surface.

With the parameters we established in Table 2, we carried out iterative fits of the adsorption data to establish the stoichiometry of surface reactions that could account for the data. We collected surface adsorption data at four ionic strengths with different dissolved ions present and a variable pH range; thus, we thoroughly tested our surface complexation model to permit the selection of a unique set of surface reactions that successfully predict the adsorption of aspartate and Ca^{2+} onto brucite.

3. Results and Discussion

3.1. Aspartate Adsorption onto Brucite at Low Mg^{2+} Concentrations

The adsorption isotherm in Fig. 4 shows the surface adsorption of aspartate on brucite as a function of the remaining aspartate concentration in the aqueous phase ($[\text{Asp}]_{\text{eq}}$) when low concentrations of Mg^{2+} dissolved from the surface. We report all aspartate adsorption data in Table 4. Aspartate surface adsorption (Γ_{ads}) at a pH of 10.3 ± 0.1 increased steadily as a function of $[\text{Asp}]_{\text{eq}}$. The error in Fig. 4 is $\pm 1 \text{ }\sigma$ of each Γ_{ads} and $[\text{Asp}]_{\text{eq}}$ value. Aspartate surface adsorption was between 8.4×10^{-2} and $1.0 \text{ }\mu\text{mol}\cdot\text{m}^{-2}$. This range of Γ_{ads} corresponds to 16.4 to 3.6 % of the initial aspartate added, where the percentage of adsorbed aspartate decreases with increasing Γ_{ads} . As a consequence, the reported error substantially increases with Γ_{ads} and $[\text{Asp}]_{\text{eq}}$.

As previously discussed, we assumed that only the lateral brucite surfaces were active in adsorption reactions; however, we were unable to distinguish whether there was one lateral surface that more predominantly adsorbed aspartate. Our crystallographic examination of the

idealized lateral surfaces (100), (110), (111), and (010) revealed two distinct terminations on the three-coordinated brucite hydroxyl group: a geminal site ($>\text{Mg}(\text{OH})_2$) with two bonds broken and a bridging site ($>\text{MgOH}$) with a single bond broken at the surface. We could not determine whether aspartate attached at a geminal or bridging reaction site with the ETLM because our proposed surface reactions in Table 2 resulted in the same amount of electrical work at both sites. Therefore to avoid ambiguity, we display surface reactions occurring at the (100) brucite plane as an example surface only, and henceforth regard the neutral brucite surface as “ $>\text{SOH}$ ”.

We used the ETLM to carry out a series of iterative calculations to establish possible surface reactions for the aspartate. We proposed a surface reaction represented by the stoichiometry

This reaction resulted in a close fit to the adsorption data, within analytical uncertainties, as illustrated by Fig. 4. We subsequently interpreted this surface reaction as involving a bidentate mononuclear, or “leaning” configuration with two points of attachment (one that is hydrogen-bonded and one that is inner-sphere) onto the brucite surface. A possible schematic illustration of the “leaning” surface species is shown in Fig. 5. Pokrovsky and coworkers (2005) also concluded that it was likely that biomolecules attached onto brucite through an inner-sphere attachment. Our observed range of aspartate adsorption on the brucite surface is consistent with the citrate adsorption ($0.4 \mu\text{mol}\cdot\text{m}^{-2}$) measurements by Pokrovsky and colleagues (2005). Furthermore, we previously measured similar amounts of aspartate adsorption ($1 \mu\text{mol}\cdot\text{m}^{-2}$) on rutile [TiO_2], and our calculations with the ETLM also predicted an inner-sphere attachment of the aspartate molecule at the mineral-water interface (Jonsson et al., 2010).

The reaction in Eq. (7) corresponds to the equilibrium constant

$${}^*K_{>\text{SOH}_2^+>\text{SAsp}^-}^0 = \frac{a_{>\text{SOH}_2^+>\text{SAsp}^-} a_{>\text{H}_2\text{O}}}{a_{>\text{SOH}}^2 a_{\text{HAsp}^-} a_{\text{H}^+}} 10^{\frac{F\Delta\psi_{r,7}}{2.303RT}} \quad (8)$$

where the superscript “*” indicates the reaction is expressed relative to the >SOH surface, and “0” refers to a hypothetical 1.0 molal standard state (Sverjensky, 2003). The value $\Delta\psi_{r,7}$ represents the electrical work involved in Eq. (7). The electrical work includes a contribution from the movement of water dipoles off the brucite surface according to $\Delta\psi_{r,7} = -n_{\text{H}_2\text{O}}(\psi_0 - \psi_\beta)$, where $n_{\text{H}_2\text{O}}$ are the number of water molecules on the right-hand side of the reaction. In Eq. (7), $n_{\text{H}_2\text{O}} = 1$, which resulted in $\Delta\psi_{r,7} = \psi_0 - \psi_\beta$.

3.2. Aspartate Adsorption onto Brucite at High Mg^{2+} Concentrations

When brucite is immersed in water, the surface dissolves in aqueous solution according to

where the equilibrium constant of Eq. (9) is

$$\log K = \frac{a_{\text{Mg}^{2+}}}{a_{\text{H}^+}^2} \quad (10)$$

We calculated equilibrium constants of $\log K = 17.3 \pm 0.01$ and $\log K = 17.3 \pm 0.06$ for batch experiments in which we added only water or aqueous aspartate, respectively, to our synthetic brucite powder. These equilibrium constants are consistent with Pokrovsky and Schott’s measured value of $\log K = 17.6 \pm 0.5$ for a ground, natural sample of brucite. We calculated the steady-state activity ratio from the activity coefficient of Mg^{2+} with GEOSURF and our measured pH values from the batch adsorption experiments. We observed that the activity ratio in Eq. (10) did not change as a function of pH (Fig. 6). Therefore, we assumed that brucite was in equilibrium with the aqueous phase during our batch adsorption experiments. If brucite was in equilibrium with the aqueous phase, we estimated that our addition of MgCl_2 to an aspartate

batch experiment would decrease solution pH while maintaining the equilibrium constant. We therefore calculated the activity ratio with data from the high-Mg²⁺ experiment. As we predicted, pH decreased (pH= 9.9 ± 0.1), and the average activity ratio of logK= 17.2 ± 0.04 was consistent within the range of uncertainty of the logK we calculated for the low-Mg²⁺ experiment.

The adsorption isotherm in Fig. 7 shows aspartate adsorption on brucite at high Mg²⁺ concentrations along with data from the low-Mg²⁺ experiment. We had no indication from our IC analysis that the MgCl₂ we added adsorbed onto the brucite surface. Aspartate adsorption at high Mg²⁺ concentrations was between 1.7×10⁻² and 0.6 μmol•m⁻². In comparison with adsorption data we obtained under low Mg²⁺ concentrations, Γ_{ads} decreased by a factor of about 1.5.

Using the ETLM and the surface reaction proposed in Eq. (7), we predicted adsorption of aspartate for the high-Mg²⁺ data at pH= 9.9. As shown in Fig. 8, there is close agreement between the curve predicted by the “leaning” aspartate species from Eq. (7) and our data at two distinct Mg²⁺ concentrations. This fit lends support to our use of the surface reaction in Eq. (7).

We also tested and dismissed dozens of other surface reactions with the ETLM. For instance, we considered the reaction stoichiometry

which resulted in an excellent fit to the adsorption data at low Mg²⁺ concentrations. However, this reaction also predicted substantially higher amounts of aspartate adsorption (up to 3.5 μmol•m⁻²) at high Mg²⁺ concentrations. This strong disagreement with the experimental data led us to dismiss this surface reaction. Through this process of elimination, we determined that only the surface reaction in Eq. (11) could predict aspartate adsorption at both Mg²⁺ concentrations.

According to the reaction in Eq. (7), Asp²⁻ attaches onto brucite. However, when the pH decreased from 10.2 in the low-Mg²⁺ experiment to 9.9 in the high-Mg²⁺ experiment, the

abundance of the doubly-deprotonated aspartate species (pKa=10.01) decreased in the aqueous phase. Therefore, the decrease in pH may have caused a decrease in aspartate adsorption.

3.3. Aspartate Adsorption onto Brucite in the Presence of Ca^{2+}

The adsorption isotherms in Fig. 8 illustrate aspartate adsorption onto brucite in the presence of low and high Ca^{2+} concentrations. The low- Ca^{2+} and high- Ca^{2+} experiments were conducted at a pH of 10.2 ± 0.02 and 10.2 ± 0.05 , respectively. We observed that surface adsorption increased with Ca^{2+} concentration, where Γ_{ads} was between 4.3×10^{-2} and $1.8 \mu\text{mol} \cdot \text{m}^{-2}$ for the low- Ca^{2+} experiment and between 0.1 and $2.1 \mu\text{mol} \cdot \text{m}^{-2}$ for the high- Ca^{2+} experiment. The range of Γ_{ads} in the high- Ca^{2+} experiment corresponded to between 56.4 and 17.1 % aspartate adsorption, respectively.

Aspartate surface adsorption was significantly greater in the presence of Ca^{2+} compared with the low- Mg^{2+} experiment (see Fig. 4). In particular, when aspartate adsorbed onto brucite at high Ca^{2+} concentrations, Γ_{ads} increased two-fold. In addition to this increase in aspartate adsorption, we observed that a significant amount of Ca^{2+} adsorbed onto brucite as illustrated in Fig. 9. The amount of Ca^{2+} surface adsorption averaged $21.6 \pm 1.8 \mu\text{mol} \cdot \text{m}^{-2}$ (41.8 %) for the low- Ca^{2+} experiment and $29.6 \pm 2.3 \mu\text{mol} \cdot \text{m}^{-2}$ (14.4 %) for the high- Ca^{2+} experiment. We report all calcium adsorption data in Table 5.

We developed a model of the surface adsorption of aspartate and Ca^{2+} by iteratively fitting the adsorption data. We discovered that in addition to the surface reaction in Eq. (7), two surface reactions involving Ca^{2+} were needed to fit the experimental data. The first surface reaction involved the formation of a calcium- aspartate complex

that may attach as a monodentate outer-sphere species on brucite. Franchi and colleagues (2003) observed enhanced adsorption of nucleic acids on the clay minerals montmorillonite and kaolinite in the presence of Ca^{2+} . The authors suggested a configuration involving calcium-nucleic acid complexes where the Ca^{2+} ion would act as a “bridge” between the nucleic acid and the clay-water interface (Franchi et al., 2003). We estimated a similar configuration of our predicted calcium-aspartate species at the brucite surface in Fig. 10a. We proposed that the calcium-aspartate species adsorbs in addition to the “leaning” aspartate species (Fig. 5). The equilibrium constant of the surface reaction in Eq. (12) is

$$^*K_{>\text{SO}^-\text{Ca}(\text{HAsp})^+}^0 = \frac{a_{>\text{SO}^-\text{(HAsp)}^+} a_{\text{H}^+}}{a_{>\text{SOH}} a_{\text{HAsp}^-} a_{\text{Ca}^{2+}}} 10^{\frac{F\Delta\psi_{r,12}}{2.303RT}} \quad (13)$$

where the value $\Delta\psi_{r,12} = -\psi_0 + \psi_\beta$.

The second reaction we proposed required the adsorption of a hydrated calcium complex

that can be interpreted as a monodentate outer-sphere surface species as illustrated in Fig. 10b.

The equilibrium constant of the surface reaction in Eq. (14) is

$$^*K_{>\text{SOH}_2^+\text{Ca}(\text{OH})_2}^0 = \frac{a_{>\text{SOH}_2^+\text{Ca}(\text{OH})_2} a_{\text{H}^+}}{a_{>\text{SOH}} a_{\text{Ca}^{2+}} a_{\text{H}_2\text{O}}^2} 10^{\frac{F\Delta\psi_{r,14}}{2.303RT}} \quad (15)$$

where $\Delta\psi_{r,14} = \psi_0$.

Our proposed species differs from the prediction by Pokrovsky and coworkers (2005) that Ca^{2+} may be attaching onto brucite as a monodentate or bidentate inner-sphere species. We tested surface reactions containing these species with the ETLM. However, they failed to adequately fit the Ca^{2+} and aspartate adsorption data. For example, the reaction stoichiometry

which we can alternatively express as

437 might result in a monodentate inner-sphere calcium species at the brucite surface. This reaction
 438 underestimated the amount of Ca^{2+} adsorption for the high- Ca^{2+} experiment by approximately
 439 $5.7 \mu\text{mol}\cdot\text{m}^{-2}$. Furthermore, the addition of this surface reaction in combination with Eqs. (7) and
 440 (12) suppressed the amount of aspartate adsorption to less than $0.1 \mu\text{mol}\cdot\text{m}^{-2}$ at both Ca^{2+}
 441 concentrations. We therefore could not consider Eq. (16) as a surface reaction representing Ca^{2+}
 442 adsorption onto brucite.

443 We combined the surface reactions in Eqs. (7), (12), and (14) to fit the aspartate
 444 adsorption data at low and high Ca^{2+} concentrations (Fig. 9). The calculated curve for the low-
 445 Ca^{2+} experimental data fit within analytical uncertainty. Although in overall agreement with the
 446 isotherm shape, the curve calculated at high Ca^{2+} concentrations is a bit low compared with all of
 447 the data points except for the highest surface coverage, but is consistent with the data within ± 2.5
 448 σ_M , or approximately 7 % uncertainty.

449 We predicted curves for Ca^{2+} adsorption as a function of initial aspartate concentration,
 450 $[\text{Asp}]_0$. As shown in Fig. 9, there are two data points from the high- Ca^{2+} experiment that are
 451 elevated above the predicted curve. However, we determined that the average predicted surface
 452 adsorption of Ca^{2+} was $28.1 \mu\text{mol}\cdot\text{m}^{-2}$, which is well within the analytical uncertainty of the
 453 average experimental surface adsorption of Ca^{2+} , $29.6 \pm 2.3 \mu\text{mol}\cdot\text{m}^{-2}$. Therefore, our prediction
 454 of Ca^{2+} surface adsorption is in acceptable agreement with the data collected at low and high
 455 Ca^{2+} concentrations. We thus regard the surface reactions proposed by Eqs. (7), (12), and (14) as
 456 sufficiently characterizing the aspartate-brucite and Ca^{2+} -brucite adsorption systems with respect
 457 to different ionic strengths, pH conditions, divalent cation concentrations and initial aspartate
 458 concentrations.

3.4. Prediction of Brucite Particle Surface Charge

The significant increase in aspartate adsorption with Ca^{2+} and its decrease with MgCl_2 addition suggest that the particle surface charge of brucite shifted relative to the experimental conditions we investigated (Table 1). We therefore calculated the particle surface charge by predicting the ζ -potential of the brucite surface with the ETLM as a function of $[\text{Asp}]_0$ for each type of batch experiment. In these calculations, we assumed that the ζ -potential was equal to the potential at the start of the diffuse layer (Ψ_d) in the ETLM. As shown in Fig. 11, we predicted that ζ -potential was neutral to slightly negative for the low- Mg^{2+} experiment, and it decreased with increasing $[\text{Mg}^{2+}]$. This could be a result of decreased pH with added MgCl_2 , which may directly affect the brucite surface charge. Moreover, we did not observe Mg^{2+} adsorption on brucite, although we added MgCl_2 to the system. In Eq. (7), we also predicted the doubly-deprotonated aspartate molecule, Asp^{2-} , adsorbed onto brucite. A decrease in surface particle charge could create an unfavorable electrostatic environment for Asp^{2-} to adsorb onto brucite, which reflects our observation of decreased aspartate adsorption with added MgCl_2 . In Fig. 11, we modeled a reversal of the ζ -potential with Ca^{2+} . This calculation of positive particle surface charge reflects the ETLM reactions in Eqs. (12) and (14) where calcium adsorbs onto the brucite surface resulting in a net positive surface charge.

The cooperative calcium-aspartate complex, CaHAsp^+ , may have enabled the aspartate molecule to adsorb in greater amounts. We predicted the distribution of the two adsorbing aspartate species in Eqs. (7) and (12) with the ETLM at both low and high Ca^{2+} concentrations as a function of $[\text{Asp}]_0$ in Fig. 12. At both Ca^{2+} concentrations, we observed that the calcium-aspartate outer-sphere species overwhelmingly predominates, whereas very little of the surface

adsorption can be represented by the “leaning” aspartate species. This result may provide evidence that the formation of a calcium-aspartate complex enhances the adsorption of aspartate.

We predicted that a calcium-aspartate surface species forms at the brucite-water interface (Eq. 12). We predicted the distribution of aqueous aspartate species that can be represented by calcium-aspartate complexes (Fig. 13), and we determined that up to 25.8 % and 6.5 % of aqueous aspartate species are represented by CaAsp^0 at high Ca^{2+} and low Ca^{2+} concentrations, respectively. This prediction indicates that the calcium-aspartate aqueous complex could easily form during our batch adsorption experiments. Whereas CaAsp^0 predominates in aqueous solution, CaHAsp^+ does not exceed 0.5 % at low Ca^{2+} concentrations and 2 % at high Ca^{2+} concentrations. Nevertheless, we predicted that the CaHAsp^+ species predominates as a surface species, which illustrates the tendency for the mineral-water interface to adsorb organic species that may not always predominate in bulk water (Geiger, 2009; Malin et al., 2009).

It is possible that because aspartate is a polar, negatively-charged amino acid, it may readily form a cooperative complex with calcium that leads to advantageous surface adsorption on brucite. A similar amino acid, such as glutamate ($\text{C}_5\text{H}_9\text{NO}_4$), may also form a complex with Ca^{2+} and follow a similar pattern of increased surface adsorption. Likewise, Ca^{2+} may not strongly bond to a positively-charged amino acid, such as lysine ($\text{C}_6\text{H}_{14}\text{N}_2\text{O}_2$), and competitively adsorb with the amino acid on the brucite surface. Whether Ca^{2+} may either cooperatively or competitively affect the surface adsorption of a particular biomolecule onto brucite is worth future investigation.

3.5. Prediction of Aspartate Adsorption in Lost City Fluids

We propose that an ideal natural environment for aspartate to adsorb onto brucite would have elevated concentrations of Ca^{2+} and little or no Mg^{2+} present. Serpentinite-hosted hydrothermal systems provide such an environment with brucite and possibly other biomolecules such as aspartate (Kelley et al., 2001; Shock and Canovas, 2010). Kelley and coworkers (2002) report that vent fluids at the Lost City hydrothermal field have high concentrations of Ca^{2+} (23.3×10^{-3} M) relative to a 0 M Mg^{2+} endmember composition over a pH range between 9.0 and 9.8. We invoked the surface reactions in Eqs. (7), (12), and (14) with the major ion (Na^+ , Cl^- , Ca^{2+} and SO_4^{2-}) concentrations reported by Kelley and coworkers (2002) to predict the surface adsorption of aspartate on brucite in Lost City vent fluids at 25 °C and 1 bar. We predicted surface adsorption at pH values of 9.8 and 10.2. We found that surface adsorption steadily increases with a decrease in pH. For instance, we predicted that surface adsorption at a pH of 9.8 is approximately double the adsorption at a pH of 10.2 at a wide range (2-300 μM) of aspartate concentrations. Although some of the aspartate concentrations in our predictions might exceed prebiotically-plausible concentrations of amino acids at hydrothermal vents, our model could possibly predict changes in amino acid adsorption as pH fluctuates as a consequence of an evolving hydrothermal system.

Our predictions for aspartate adsorption in the presence of the Lost City vent fluids are at ambient conditions. However, it is possible that the ETLM predictions could be extrapolated to higher temperatures in future studies if experiments are conducted under more plausible hydrothermal temperatures and pressures. It also may be that vent fluids in a Lost City-type hydrothermal field may mix with cold seawater to approximate ambient temperatures. This low-temperature environment may hold an ideal $\text{Ca}^{2+}/\text{Mg}^{2+}$ ratio that could promote favorable aspartate adsorption onto brucite. Therefore, the results we both observed and predicted in this

study appear to be highly relevant for natural serpentinite-hosted hydrothermal systems. Within this plausible prebiotic environment we have observed a cooperative surface adsorption effect between Ca^{2+} and aspartate on brucite. Our combination of batch adsorption data and surface complexation modeling contributes to a fundamental understanding of the role of the brucite-water interface in the emergence of complex biochemistry and the geochemical origin of life.

Acknowledgements

We would like to thank Cécile Feuillie, Namhey Lee, Alyssa K. Adcock, Timothy Strobel, Dionysis Foustoukos, Paul Goldey, John Armstrong, Adrian Villegas-Jimenez, Stephen Hodge and Steven Coley for their invaluable expertise and advice throughout this project. We thank the National Science Foundation, the NASA Astrobiology Institute, the Deep Carbon Observatory, Johns Hopkins University, and the Carnegie Institution for Science for support of this research.

References

- Anderson, P. J., Horlock, R. F., and Oliver, J. F. (1965) Interaction of water with the magnesium oxide surface. *Trans. Faraday Soc.* **61**, 2754-2762.
- Ali Ahmad M., Prélôt B., Razafitianamaharavo A., Douillard J.-M., Zajac J., Dufour F., Durupthy O., Chanéac C. and Villiéras F. (2012) Influence of morphology and crystallinity on surface reactivity of nanosized anatase TiO_2 studied by adsorption techniques. 1. The use of gaseous molecular probes. *J. Phys. Chem., C* **116**, 24596–24606.
- Bach W., Paulick H., Garrido C. J., Ildefonse B., Meurer W. P. and Humphris S. E. (2006) Unraveling the sequence of serpentinization reactions: Petrography, mineral chemistry, and petrophysics of serpentinites from MAR 15°N (ODP Leg 209, Site 1274). *Geophys. Res. Lett.* **3**, 1-4.
- Bahri, S., Jonsson, C. M., Jonsson, C. L., Azzolini, D., Sverjensky, D. A. and Hazen, R. M. (2011) Adsorption and surface complexation study of L-DOPA on rutile ($\alpha\text{-TiO}_2$) in NaCl solutions. *Environ. Sci. Technol.* **45**, 3959-3966.
- Baross J. A. and Hoffman S. E. (1985) Submarine hydrothermal vents and associated gradient environments as sites for the origin and evolution of life. *Orig. Life Evol. Biosph.* **15**, 327–345.

- 557 Cleaves H. J., Chalmers J. H., Lazcano A., Miller S. L. and Bada J. L. (2008) A reassessment of
558 prebiotic organic synthesis in neutral planetary atmospheres. *Orig. Life Evol. Biosph.* **38**,
559 105–115.
- 560 Cleaves H. J., Michalkova Scott A., Hill F. C., Leszczynski J., Sahai N. and Hazen R. M. (2012)
561 Mineral–organic interfacial processes: Potential roles in the origins of life. *Chem. Soc. Rev.*
562 **41**, 5502–5525.
- 563 De Robertis A., De Stefano C. and Gianguzza A. (1991) Salt effects on the protonation of L-
564 histidine and L-aspartic acid: A complex formation model. *Thermochim. Acta* **177**, 39–57.
- 565 Deamer D. and Pashley R. M. (1989) Amphiphilic Components of the Murchison Carbonaceous
566 Chondrite - Surface-Properties and Membrane Formation. *Orig. Life Evol. Biosph.* **19**, 21–38.
- 567 Dymek R. F., Brothers S. C. and Schiffries C. M. (1988) Petrogenesis of ultramafic metamorphic
568 rocks from the 3800 Ma Isua Supracrustal Belt, West Greenland. *J. Petrol.* **29**, 1353–1397.
- 569 Eypert-Blaison C., Villiéras F., Michot L. J., Pelletier M., Humbert B., Ghanbaja J. and Yvon J.
570 (2002) Surface heterogeneity of kanemite, magadiite and kenyaite: a high-resolution gas
571 adsorption study. *Clay Miner.* **37**, 531–542.
- 572 Franchi M., Ferris J. P. and Gallori E. (2003) Cations as mediators of the adsorption of nucleic
573 acids on clay surfaces in prebiotic environments. *Orig. Life Evol. Biosph.* **33**, 1–16.
- 574 Geiger F. M. (2009) Second Harmonic Generation, Sum Frequency Generation, and χ (3):
575 Dissecting environmental interfaces with a nonlinear optical swiss army knife. *Annu. Rev.*
576 *Phys. Chem.* **60**, 61–83.
- 577 Giacomelli C. E., Avena M. J. and De Pauli C. P. (1995) Aspartic acid adsorption onto TiO₂
578 particles surface. Experimental data and model calculations. *Langmuir* **11**, 3483–3490.
- 579 Goldschmidt V. M. (1952) Geochemical aspects of the origin of complex organic molecules on
580 the Earth, as precursors to organic life. *New Biol.* **12**, 97–105.
- 581 Greiner E., Kumar K., Sumit M., Giuffre A., Zhao W., Pedersen J. and Sahai N. (2014)
582 Adsorption of L-glutamic acid and L-aspartic acid to γ -Al₂O₃. *Geochim. Cosmochim. Acta*
583 **133**, 142–155.
- 584 Hazen R. M. (2005) *Genesis: The Scientific Quest for Life's Origin*, Joseph Henry Press,
585 Washington, DC.
- 586 Hazen R. M. (2006) Presidential address to the Mineralogical Society of America, Salt Lake City,
587 October 18, 2005: Mineral surfaces and the prebiotic selection and organization of
588 biomolecules. *Am. Mineral.* **91**, 1715–1729.
- 589 Hazen R. M., Filley T. and Goodfriend G. (2001) Selective adsorption of L- and D-amino acids
590 on calcite: Implications for biochemical homochirality. *PNAS* **98**, 5487–5490.

- 591 Hennes R., Holm N. G. and Engel M. H. (1992) Abiotic synthesis of amino-acids under
592 hydrothermal conditions and the origin of life - a perpetual phenomenon.
593 *Naturwissenschaften* **79**, 361–365.
- 594 Henrist C., Mathieu J., Vogels C., Rulmont A. and Cloots R. (2003) Morphological study of
595 magnesium hydroxide nanoparticles precipitated in dilute aqueous solution. *J. Cryst. Growth*
596 **249**, 321–330.
- 597 Herzberg C., Condie K. and Korenaga J. (2010) Thermal history of the Earth and its petrological
598 expression. *Earth Planet. Sci. Lett.* **292**, 79–88.
- 599 Holm N. G. (2012) The significance of Mg in prebiotic geochemistry. *Geobiology* **10**, 269–279.
- 600 Holm N. G. and Andersson E. (2005) Hydrothermal simulation experiments as a tool for studies
601 of the origin of life on earth and other terrestrial planets: A review. *Astrobiology* **5**, 444–460.
- 602 Holm N. G., Dumont M., Ivarsson M. and Konn C. (2006) Alkaline fluid circulation in
603 ultramafic rocks and formation of nucleotide constituents: a hypothesis. *Geochem. Trans.* **7**,
604 1–13.
- 605 Hostetler P. B. (1963) The stability and surface energy of brucite in water at 25 degrees C. *Am. J.*
606 *Sci.* **261**, 238–258.
- 607 Ikhsan J., Johnson B. B., Wells J. D. and Angove M. J. (2004) Adsorption of aspartic acid on
608 kaolinite. *J. Colloid Interface Sci.* **273**, 1–5.
- 609 Jalbout A. F., Abrell L., Adamowicz L., Polt R., Apponi A. J. and Ziurys L. M. (2007) Sugar
610 synthesis from a gas-phase formose reaction. *Astrobiology* **7**, 433–442.
- 611 Jonsson C. M., Jonsson C. L., Estrada C. F., Sverjensky D. A., Cleaves H. J. II and Hazen R. M.
612 (2010) Adsorption of L-aspartate to rutile (α -TiO₂): Experimental and theoretical surface
613 complexation studies. *Geochim. Cosmochim. Acta* **74**, 2356–2367.
- 614 Jonsson C. M., Jonsson C. L., Sverjensky D. A., Cleaves H. J. and Hazen R. M. (2009)
615 Attachment of L-glutamate to rutile (α -TiO₂): A potentiometric, adsorption, and surface
616 complexation Study. *Langmuir* **25**, 12127–12135.
- 617 Jordan G. and Rammensee W. (1996) Dissolution rates and activation energy for dissolution of
618 brucite (001): A new method based on the microtopography of crystal surfaces. *Geochim.*
619 *Cosmochim. Acta* **60**, 5055–5062.
- 620 Kelley D. S., Baross J. A. and Delaney J. R. (2002) Volcanoes, fluids, and life at mid-ocean
621 ridge spreading centers. *Annu. Rev. Earth Planet. Sci.* **30**, 385–491.
- 622 Kelley D. S., Karson J. A., Fruh-Green G. L., Yoerger D. R., Shank T. M., Butterfield D. A.,
623 Hayes J. M., Schrenk M. O., Olson E. J., Proskurowski G., Jakuba M., Bradley A., Larson B.,
624 Ludwig K., Glickson D., Buckman K., Bradley A.S., Brazelton W.J., Roe K., Elend M.J.,
625 Delacour A., Bernasconi S.M., Lilley M.D., Baross J.A., Summons R.T., and Sylva S.P.

626 (2005) A serpentinite-hosted ecosystem: the Lost City hydrothermal field. *Science* **307**,
627 1428-1434.

628 Kelley D. S., Karson J., Blackman D., Frueh-Green G., Butterfield D., Lilley M., Olson E.,
629 Schrenk M., Roe K., Lebon G., Rivizzigno P., and the AT3-60 Shipboard Party (2001) An
630 off-axis hydrothermal vent field near the Mid-Atlantic Ridge at 30 degrees N. *Nature* **412**,
631 145–149.

632 Kim H.-J., Ricardo A., Illangkoon H. I., Kim M. J., Carrigan M. A., Frye F. and Benner S. A.
633 (2011) Synthesis of carbohydrates in mineral-guided prebiotic cycles. *J. Am. Chem. Soc.* **133**,
634 9457–9468.

635 Koretsky C. M., Sverjensky D. A. and Sahai N. (1998) A model of surface site types on oxide
636 and silicate minerals based on crystal chemistry; implications for site types and densities,
637 multi-site adsorption, surface infrared spectroscopy, and dissolution kinetics. *Am. J. Sci.* **298**,
638 349–438.

639 Kudoh Y., Kameda J. and Kogure T. (2006) Dissolution of brucite on the (001) surface at neutral
640 pH: In situ atomic force microscopy observations. *Clay Clay Miner.* **54**, 598–604.

641 Lee, N., Hummer, D. R., Sverjensky, D. A., Rajh, T., Hazen, R. M., Steele, A. and Cody, G. D.
642 (2012) Speciation of L-DOPA on nanorutile as a function of pH and surface coverage using
643 surface-enhanced Raman spectroscopy (SERS). *Langmuir* **28**, 17322-17330.

644 Liu X., Cheng J., Sprik M., Lu X. and Wang R. (2013) Understanding surface acidity of gibbsite
645 with first principles molecular dynamics simulations. *Geochim. Cosmochim. Acta* **120**, 487–
646 495.

647 Lu J., Qiu L. and Qu B. (2004) Controlled growth of three morphological structures of
648 magnesium hydroxide nanoparticles by wet precipitation method. *J. Cryst. Growth* **267**,
649 676–684.

650 Malin J. N., Holland J. G. and Geiger F. M. (2009) Free energy relationships in the electric
651 double layer and alkali earth speciation at the fused silica/water interface. *J. Phys. Chem., C*
652 **113**, 17795–17802.

653 McCollom T. M. and Bach W. (2009) Thermodynamic constraints on hydrogen generation
654 during serpentinization of ultramafic rocks. *Geochim. Cosmochim. Acta* **73**, 856–875.

655 Michot L. J. and Villiéras F. (2002) Assessment of surface energetic heterogeneity of synthetic
656 Na-saponites. The role of layer charge. *Clay Miner.* **37**, 39–57.

657 Noffke N., Christian D., Wacey D. and Hazen R. M. (2013) Microbially induced sedimentary
658 structures recording an ancient ecosystem in the ca. 3.48 billion-year-old Dresser Formation,
659 Pilbara, Western Australia. *Astrobiology* **13**, 1103–1124.

660 Parikh, S. J., Kubicki, J. D., Jonsson, C. M., Jonsson, C. J., Hazen, R. M., Sverjensky, R. M.,
661 Sverjensky, D. A., and Sparks, D. L. (2011) Evaluating glutamate and aspartate binding

662 mechanisms to rutile (α -TiO₂) via ATR-FTIR spectroscopy and quantum chemical
 663 calculations. *Langmuir* **27**, 1778-1787.

664 Palmer D. (2009) CrystalMaker for Mac OS X, 8.2. 2. *CrystalMaker Software Limited:*
 665 *Oxfordshire*.

666 Perronnet M., Villiéras F., Jullien M., Razafitianamaharavo A., Raynal J. and Bonnin D. (2007)
 667 Towards a link between the energetic heterogeneities of the edge faces of smectites and their
 668 stability in the context of metallic corrosion. *Geochim. Cosmochim. Acta* **71**, 1463–1479.

669 Pokrovsky O. S. and Schott J. (2004) Experimental study of brucite dissolution and precipitation
 670 in aqueous solutions: surface speciation and chemical affinity control. *Geochim. Cosmochim.*
 671 *Acta* **68**, 31–45.

672 Pokrovsky O. S., Schott J. and Castillo A. (2005) Kinetics of brucite dissolution at 25°C in the
 673 presence of organic and inorganic ligands and divalent metals. *Geochim. Cosmochim. Acta*
 674 **69**, 905–918.

675 Ponnamperna C. and Pering K. (1966) Possible abiogenic origin of some naturally occurring
 676 hydrocarbons. *Nature* **209**, 979–982.

677 Prélôt B., Villiéras F., Pelletier M., Gérard G., Gaboriaud F., Ehrhardt J.-J., Perrone J., Fedoroff
 678 M., Jeanjean J., Lefèvre G., Mazerolles L., Pastol J.-L., Rouchaud J.-C. and Lindecker C.
 679 (2003) Morphology and surface heterogeneities in synthetic goethites. *J. Colloid Interface*
 680 *Sci.* **261**, 244–254.

681 Proskurowski G., Lilley M. D., Kelley D. S. and Olson E. J. (2006) Low temperature volatile
 682 production at the Lost City Hydrothermal Field, evidence from a hydrogen stable isotope
 683 geothermometer. *Chem. Geol.* **229**, 331–343.

684 Roddick-Lanzilotta A. D. and McQuillan A. J. (2000) An in situ Infrared spectroscopic study of
 685 glutamic acid and of aspartic acid adsorbed on TiO₂: Implications for the biocompatibility of
 686 titanium. *J. Colloid Interface Sci.* **227**, 48–54.

687 Sahai N. and Sverjensky D. A. (1998) GEOSURF: A computer program for modeling adsorption
 688 on mineral surfaces from aqueous solution. *Comput. Geosci.* **24**, 853–873.

689 Sahai N. and Sverjensky D. A. (1997) Solvation and electrostatic model for specific electrolyte
 690 adsorption. *Geochim. Cosmochim. Acta* **61**, 2827–2848.

691 Sayed-Hassan M., Villiéras F., Gaboriaud F. and Razafitianamaharavo A. (2006) AFM and low-
 692 pressure argon adsorption analysis of geometrical properties of phyllosilicates. *J. Colloid*
 693 *Interface Sci.* **296**, 614–623.

694 Sayed-Hassan M., Villiéras F., Razafitianamaharavo A. and Michot L. J. (2005) Role of
 695 exchangeable cations on geometrical and energetic surface heterogeneity of kaolinites.
 696 *Langmuir* **21**, 12283–12289.

- 697 Schott H. (1981) Electrokinetic studies of magnesium hydroxide. *J. Pharm. Sci.* **70**, 486–489.
- 698 Shibuya T., Komiya T., Nakamura K., Takai K. and Maruyama S. (2010) Highly alkaline, high-
699 temperature hydrothermal fluids in the early Archean ocean. *Precambrian Res.* **182**, 230–
700 238.
- 701 Shock E. L. and Canovas P. (2010) The potential for abiotic organic synthesis and biosynthesis
702 at seafloor hydrothermal systems. *Geofluids* **10**, 161–192.
- 703 Shock E. L. and Schulte M. D. (1998) Organic synthesis during fluid mixing in hydrothermal
704 systems. *J. Geophys. Res.* **103**, 28513.
- 705 Sleep N. H., Bird D. K. and Pope E. C. (2011) Serpentinite and the dawn of life. *Philos. Trans. R.*
706 *Soc. Lond., B* **366**, 2857–2869.
- 707 Stevanovic S., AliAhmad M., Razafitianamaharavo A., Villiras F., Odile B., Prlot B., Zajac J.,
708 Douillard J.-M. and Chanac C. (2010) Evidences for the relationship between surface
709 structure and reactivity of goethite nanoparticles based on advanced molecular-probe
710 methods. *Adsorption* **16**, 185–195.
- 711 Sverjensky D. A. (2005) Prediction of surface charge on oxides in salt solutions: Revisions for
712 1:1 (M+L) electrolytes. *Geochim. Cosmochim. Acta* **69**, 225–257.
- 713 Sverjensky D. A. (2003) Standard states for the activities of mineral surface sites and species.
714 *Geochim. Cosmochim. Acta* **67**, 17–28.
- 715 Sverjensky D. A. and Fukushi K. (2006) A predictive model (ETLM) for As(III) adsorption and
716 surface speciation on oxides consistent with spectroscopic data. *Geochim. Cosmochim. Acta*
717 **70**, 3778–3802.
- 718 Sverjensky D. A. and Sahai N. (1996) Theoretical prediction of single-site surface-protonation
719 equilibrium constants for oxides and silicates in water. *Geochim. Cosmochim. Acta* **60**,
720 3773–3797.
- 721 Tournassat C., Neaman A., Villiras F., Bosbach D. and Charlet L. (2003) Nanomorphology of
722 montmorillonite particles: Estimation of the clay edge sorption site density by low-pressure
723 gas adsorption and AFM observations. *Am. Mineral.* **88**, 1989–1995.
- 724 Van Kranendonk M. J. (2010) Two types of Archean continental crust: Plume and plate tectonics
725 on early Earth. *Am. J. Sci.* **310**, 1187–1209.
- 726 Van Kranendonk M. J., Hugh Smithies R., Hickman A. H. and Champion D. C. (2007) Review:
727 secular tectonic evolution of Archean continental crust: Interplay between horizontal and
728 vertical processes in the formation of the Pilbara Craton, Australia. *Terra Nova* **19**, 1–38.
- 729 Villiras F., Cases J. M., Franois M., Michot L. J. and Thomas F. (1992) Texture and surface
730 energetic heterogeneity of solids from modeling of low pressure gas adsorption isotherms.
731 *Langmuir* **8**, 1789–1795.

- 732 Villiéras F., Michot L. J., Bardot F., Cases J. M., François M. and Rudziński W. (1997) An
733 improved derivative isotherm summation method to study surface heterogeneity of clay
734 minerals. *Langmuir* **13**, 1104–1117.
- 735 Villiéras F., Michot L. J., Bardot F., Chamerois M., Eypert-Blaison C., François M., Gérard G.
736 and Cases J.-M. (2002) Surface heterogeneity of minerals. *C. R. Geosci.* **334**, 597–609.

Table 1. Description of the low-Mg²⁺, high-Mg²⁺, low-Ca²⁺, and high-Ca²⁺ experiments. Dissolved Mg²⁺ concentrations from brucite are present in each experiment in addition to added MgCl₂ and CaCl₂. All concentrations are expressed in mol/L.

Experiment	Dissolved Mg ²⁺	Added MgCl ₂	Added CaCl ₂	Ionic Strength
Low-Mg ²⁺	0.7x10 ⁻³	N/A	N/A	2.1x10 ⁻³
High-Mg ²⁺	0.6x10 ⁻³	5.2x10 ⁻³	N/A	17.4x10 ⁻³
Low-Ca ²⁺	0.9x10 ⁻³	N/A	1.0x10 ⁻³	5.8x10 ⁻³
High-Ca ²⁺	1.0x10 ⁻³	N/A	4.0x10 ⁻³	15.2x10 ⁻³

Table 2. Aqueous aspartate properties^a, characteristics of brucite^b, and ETLM parameters for proton, electrolyte, and aspartate surface adsorption.

Reaction Type	Reaction	log K
Aqueous aspartate equilibria	$\text{Asp}^{2-} + \text{H}^+ = \text{HAsp}^-$	10.01
	$\text{HAsp}^- + \text{H}^+ = \text{H}_2\text{Asp}$	3.88
	$\text{H}_2\text{Asp} + \text{H}^+ = \text{H}_3\text{Asp}^+$	1.92
	$\text{HAsp}^- + \text{Cl}^- + 2\text{H}^+ = \text{H}_3\text{AspCl}$	5.3
	$\text{HAsp}^- + \text{Na}^+ = \text{NaHAsp}$	-0.3
	$\text{HAsp}^- + \text{Na}^+ = \text{NaAsp}^- + \text{H}^+$	-9.6
	$\text{HAsp}^- + \text{Ca}^{2+} + \text{H}^+ = \text{CaH}_2\text{Asp}^{2+}$	-4.43
	$\text{HAsp}^- + \text{Ca}^{2+} = \text{CaHAsp}^+$	1.43
	$\text{HAsp}^- + \text{Ca}^{2+} = \text{CaAsp} + \text{H}^+$	-7.49
Surface equilibria	Hypothetical 1.0 m standard state	
$\log K_1^0$	$>\text{SOH} + \text{H}^+ = >\text{SOH}_2^+$	9.60
$\log K_2^0$	$>\text{SO}^- + \text{H}^+ = >\text{SOH}$	11.40
$\log^* K_{\text{Na}^+}^0$	$>\text{SOH} + \text{Na}^+ = >\text{SO}^- \text{Na}^+ + \text{H}^+$	-9.00
$\log^* K_{\text{Cl}^-}^0$	$>\text{SOH} + \text{Cl}^- + \text{H}^+ = >\text{SOH}_2^+ \text{Cl}^-$	11.80
$\log^* K_{>\text{SOH}_2^+ >\text{SAsp}^-}^0$	$2>\text{SOH} + \text{H}^+ + \text{HAsp}^- = >\text{SOH}_2^+ >\text{SAsp}^- + \text{H}_2\text{O}$	16.08
$\log^* K_{>\text{SO}^- \text{CaHAsp}^+}^0$	$>\text{SOH} + \text{Ca}^{2+} + \text{HAsp}^- = >\text{SO}^- \text{CaHAsp}^+ + \text{H}^+$	-3.73
$\log^* K_{>\text{SOH}_2^+ \text{Ca(OH)}_2}^0$	$>\text{SOH} + \text{Ca}^{2+} + 2\text{H}_2\text{O} = >\text{SOH}_2^+ \text{Ca(OH)}_2 + \text{H}^+$	-5.20
Surface equilibria ^c	Site-occupancy standard states	
$\log K_{>\text{SOH}_2^+ >\text{SAsp}^-}^\theta$	$2>\text{SOH}_2^+ + \text{HAsp}^- = >\text{SOH}_2^+ >\text{SAsp}^- + \text{H}_2\text{O} + \text{H}^+$	18.80
$\log K_{>\text{SO}^- \text{CaHAsp}^+}^\theta$	$>\text{SO}^- + \text{Ca}^{2+} + \text{HAsp}^- = >\text{SO}^- \text{CaHAsp}^+$	-5.34
$\log K_{>\text{SOH}_2^+ \text{Ca(OH)}_2}^\theta$	$>\text{SOH}_2^+ + \text{Ca}^{2+} + 2\text{H}_2\text{O} = >\text{SOH}_2^+ \text{Ca(OH)}_2 + \text{H}^+$	-3.87

^a Protonation constants and electrolyte ion pair constants from (De Robertis et al., 1991; Jonsson et al., 2010)

^b Brucite properties are $N_s = 38 \text{ sites} \cdot \text{nm}^{-2}$, $A_s = 1.9 \text{ m}^2 \cdot \text{g}^{-1}$, $C_1 = 190 \text{ } \mu\text{F} \cdot \text{cm}^{-2}$, $C_2 = 190 \text{ } \mu\text{F} \cdot \text{cm}^{-2}$, $\text{pH}_{\text{PPZC}} = 10.5$, $\Delta \text{p}K_n^0 = 2.08$, $\log K_1^0 = 9.46$, $\log K_2^0 = 11.54$, $\log^0 K_{\text{Na}^+} = -9.14$, $\log^0 K_{\text{Cl}^-} = 11.66$

^c Equilibrium constants relative to site occupancy standard states can be written relative to charged surface sites calculated using the equations: $\log K_{>\text{SOH}_2^+ >\text{SAsp}^-}^\theta = \log^* K_{>\text{SOH}_2^+ >\text{SAsp}^-}^0 + \log(N_s A_s C_s)/100$; $\log K_{>\text{SO}^- \text{CaHAsp}^+}^\theta = \log^* K_{>\text{SO}^- \text{CaHAsp}^+}^0 + \log(N_s A_s)/100$; $\log K_{>\text{SOH}_2^+ \text{Ca(OH)}_2}^\theta = \log^* K_{>\text{SOH}_2^+ \text{Ca(OH)}_2}^0 + \log(N_s A_s)/100$, where N_s is site density, A_s is BET edge surface area $\text{m}^2 \cdot \text{g}^{-1}$, and C_s is solid concentration $\text{g} \cdot \text{L}^{-1}$.

Table 3. A summary of site densities calculated from idealized lateral brucite surfaces using the estimation procedures described by Koretsky and coworkers (1998), where a broken bond at a surface is considered one site per unit area. We estimated multiple site densities for every possible termination of the surfaces (100), (010), (110), and (111).

Surface	N_s (Broken Bonds, sites•nm ⁻²)
(100)	20.0
	13.4
	20.0
(010)	20.0
	13.4
	20.0
(110)	15.4
	11.6
(111)	12.9
	12.9
	9.7

Table 4a. Aspartate surface adsorption data observed for the low-Mg²⁺ and high-Mg²⁺ batch experiments.

Batch Experiment	[Asp] _{aq} (μM)	Γ _{ads} (μmol•m ⁻²)	% Ads
<i>Low Mg²⁺</i>	4.5	0.03	10.2
	17.6	0.14	12.9
	37.3	0.16	7.8
	45.8	0.24	9.0
	70.0	0.28	7.4
	110.7	0.52	8.3
	134.7	0.85	10.8
	188.2	0.70	6.7
	1.9	0.01	3.6
	8.5	0.08	16.4
	27.3	0.15	11.6
	52.1	0.43	13.8
	77.4	0.43	9.8
	87.4	0.40	8.5
	99.2	0.57	10.3
	277.4	1.00	8.1
<i>High Mg²⁺</i>	4.8	0.02	4.0
	9.8	0.02	4.5
	18.3	0.09	9.4
	46.1	0.20	8.1
	70.4	0.23	6.1
	95.0	0.34	6.6
	144.5	0.42	3.8
	190.4	0.52	5.0
	53.6	0.28	9.5
	36.4	0.16	7.7
	28.1	0.08	5.5
	238.4	0.63	4.3
	116.0	0.19	2.5
	79.8	0.27	5.4

Table 4b. Aspartate surface adsorption data observed for the low- Ca^{2+} and high- Ca^{2+} batch experiments.

Batch Experiment	[Asp] _{aq} (μM)	Γ_{ads} ($\mu\text{mol}\cdot\text{m}^{-2}$)	% Ads
<i>Low Ca^{2+}</i>	217.2	1.77	13.6
	127.2	1.20	15.3
	84.6	0.76	14.9
	61.1	0.71	18.8
	40.4	0.48	19.3
	20.3	0.25	18.8
	7.9	0.11	21.5
	4.2	0.04	17.3
<i>High Ca^{2+}</i>	90.4	1.50	24.8
	208.0	2.10	17.1
	115.9	1.73	22.8
	73.2	1.36	26.6
	51.9	1.19	30.9
	33.6	0.86	32.8
	13.4	0.59	46.4
	4.4	0.29	56.4
	2.4	0.13	52.7

Table 5. Ca^{2+} surface adsorption data observed for the low- Ca^{2+} and high- Ca^{2+} batch experiments.

Batch Experiment	$[\text{Ca}^{2+}] (\mu\text{M})$	$\Gamma_{\text{ads}} (\mu\text{mol}\cdot\text{m}^{-2})$	% Ads
<i>Low Ca^{2+}</i>	614.1	21.88	40.7
	591.2	23.25	42.9
	605.0	22.21	41.6
	586.2	22.32	43.4
	597.4	21.75	42.3
	622.9	21.66	39.8
	597.3	22.80	42.3
	601.7	21.65	41.9
<i>High Ca^{2+}</i>	3416.5	31.75	15.6
	3519.4	26.04	13.1
	3479.0	28.83	14.1
	3411.7	32.59	15.7
	3451.2	30.59	14.8
	3473.5	30.24	14.2
	3482.4	28.94	14.0
	3505.0	28.05	13.4
	3453.3	29.79	14.7

Fig. 1. SEM image of synthetic brucite (a) before hydrothermal treatment where oblong platelets are clustered in rosettes and (b) after hydrothermal treatment where platelets are hexagonal and in lamellar clusters (scale= 1 μ M).

Fig. 2. Synthetic brucite powder XRD patterns, both untreated (blue) and hydrothermally treated (black). (*hkl*) reflections are indexed by the red symbols.

Fig. 3. Surface titration data for brucite at 25 °C determined by Pokrovsky and Schott (2004) as function of solution pH at 1.0 M NaCl (red) and 1.0×10^{-2} M NaCl (blue). Dashed curves are modeled fits to the experimental data using the ETLM and the initial parameters $\log K_1 = 10.0$, $\log K_2 = 12.0$, $\log K_{\text{Na}^+}^* = -9.6$, $\log K_{\text{Cl}^-}^* = -12.2$, $A_s = 9.2 \text{ m}^2 \cdot \text{g}^{-1}$, and $N_s = 10 \text{ sites} \cdot \text{nm}^{-2}$.

Fig. 4. Adsorption of aspartate on brucite as a function of aspartate concentration remaining in solution for the low- Mg^{2+} experiments. . Dashed curve represents the modeled fit to the experimental data calculated with the parameters in Table 2. Symbols represent experimental data that are the average of a triplicate run and horizontal and vertical error bars are ± 1 standard error from the average.

Fig. 5. Possible representation of the surface species predicted by surface complexation calculations using the parameters in Table 2 where aspartate attaches onto brucite as bidentate mononuclear, or “leaning” species. Red spheres represent oxygen, yellow spheres are magnesium, tan spheres are hydrogen, black spheres are carbon, and pale blue spheres are nitrogen atoms.

Fig. 6. Equilibrium constant for Eq. (10), or $\log(a_{\text{Mg}^{2+}}/a_{\text{H}^+})^2$, as a function of solution pH for low- Mg^{2+} , water, and high- Mg^{2+} experiments. The vertical error bars are ± 1 standard deviation of each dataset.

Fig. 7. Adsorption of aspartate onto brucite as a function of aspartate concentration remaining in solution for the low- Mg^{2+} (red) and high- Mg^{2+} (blue) experiments. Dashed curves represent the modeled fits to the experimental data calculated with the parameters in Table 2. Symbols represent experimental data that are the average of a triplicate run and horizontal and vertical error bars are ± 1 standard error from the average

Fig. 8. Adsorption of aspartate onto brucite as a function of aspartate concentration remaining in solution for the low- Ca^{2+} (purple) and high- Ca^{2+} (green) experiments. Dashed curves represent the modeled fits to the experimental data calculated with the parameters in Table 2. Symbols represent experimental data that are the average of a triplicate run and horizontal and vertical error bars are ± 1 standard error from the average.

Fig. 9. Adsorption of Ca^{2+} onto brucite as a function of initial aspartate concentration determined for the low- Ca^{2+} (purple) and high- Ca^{2+} (green) experiments. Dashed curves represent the modeled fits to the experimental data calculated with the parameters in Table 2. Symbols represent experimental data and vertical error bars represent an error of 8 %.

Fig. 10. Possible representation of surface species predicted by surface complexation calculations using the parameters in Table 2. (a) Attachment of a calcium-aspartate complex to the (100) surface forming a monodentate outer-sphere species through an electrostatic interaction between a surface oxygen atom and the calcium atom. (b) Attachment of a hydrated calcium molecule to the brucite surface forming a monodentate outer-sphere species. Red spheres represent oxygen, yellow spheres are magnesium, tan spheres are hydrogen, large light blue spheres are calcium, small pale blue spheres are nitrogen and black spheres are carbon atoms.

Fig. 11. ζ -potential (mv), or particle surface charge, predicted by surface complexation modeling with the parameters in Table 2 as a function of initial aspartate concentration for the low- Mg^{2+} (red), high- Mg^{2+} (blue), low- Ca^{2+} (purple), and high- Ca^{2+} (green) experiments.

Fig. 12. The predicted surface speciation of aspartate on brucite as a function of initial aspartate concentration for the (a) low- Ca^{2+} and (b) high- Ca^{2+} experiments. The “leaning” and outer-sphere species refer to Fig. 5 and Fig. 10a and reaction stoichiometries in Eqs. (7) and (11), respectively. Dashed curves represent the individual species and the solid curve represents the total aspartate adsorption.

Fig. 13. The predicted distribution (%) of the aspartate aqueous species that can be represented by calcium-aspartate complexes, including $\text{CaH}_2\text{Asp}^{2+}$, CaHAsp^+ , and CaAsp^0 as a function of initial aspartate concentration in the presence for the a) low- Ca^{2+} and b) high- Ca^{2+} experiments.

Fig. 14. Predicted aspartate surface adsorption on brucite as a function of aspartate concentration remaining in solution in the presence of Lost City-type fluids at 25 °C and 1 bar. Aspartate surface adsorption is calculated with the parameters in Table 2 in the presence of $[\text{Na}^+]=0.49$ M, $[\text{Cl}^-]=0.55$ M, $[\text{Ca}^{2+}]=23.3\times 10^{-3}$ M, and $[\text{SO}_4^{2-}]=12.9\times 10^{-3}$ M at a pH of 9.8 (Kelley et al., 2002) and a pH of 10.2.