Compositional evolution of the upper continental crust through time, as constrained by ancient glacial diamictites

Richard M. Gaschnig^{1*}, Roberta L. Rudnick^{1**}, William F. McDonough¹, Alan J. Kaufman², John Valley³, Zhaochu Hu⁴, Shan Gao⁴, Michelle L. Beck⁵

1. Department of Geology, University of Maryland, College Park, MD 20742

2. Department of Geology and Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD 20742

3. Department of Geoscience, University of Wisconsin - Madison, Madison, WI 53706

4. State Key Laboratory of Geological Processes and Mineral Resources, China University of Geosciences, Wuhan, China 430074

5. Department of Biological Sciences, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061

* Now at School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332

** Now at Department of Earth Science, University of California - Santa Barbara, Santa Barbara, CA 93106

ABSTRACT

The composition of the fine-grained matrix of glacial diamictites from the Mesoproterozoic, Paleoproterozoic, Neoproterozoic, and Paleozoic, collected from four modern continents, reflect the secular evolution of the average composition of the upper continental crust (UCC). The effects of localized provenance are present in some cases, but distinctive geochemical signatures exist in diamictites of the same age from different localities, suggesting that these are global

signatures. Archean UCC, dominated by greenstone basalts and less so komatiites, was more mafic, based on major elements and transition metal trace elements. Temporal changes in oxygen isotope ratios, rare earth elements, and high field strength elements indicate that the UCC became more differentiated and that tonalite-trondhjemite-granodiorite suites became less important with time, findings consistent with previous studies. We also document the concentrations of siderophile and chalcophile elements (Ga, Ge, Cd, In, Sn, Sb, W, Tl, Bi) and lithophile Be in the UCC through time, and use the data for the younger diamictites to construct a new estimate of average UCC along with associated uncertainties.

1. Introduction

The Earth is unique amongst other terrestrial planets in our solar system in that it has continents, which are composed of silicic crust distinct from the basalts that dominate Earth's ocean basins and the crust of other planets. Continental crust reflects differentiation associated with plate tectonics on Earth (Taylor and McLennan, 1985; Rudnick, 1995; Condie and Kröner, 2013). How continents form and how they have grown through Earth's history remain salient first-order questions in Earth science (Committee on Grand Research Questions in the Solid-Earth Sciences, National Research Council, 2008). A necessary starting point to deciphering how continents form is knowledge of their bulk composition. The continental crust, especially the upper crust, is enriched in incompatible elements, by as much as a thousand fold relative to the mantle, and thus it exerts a large influence on the planetary budget of these elements, despite comprising a minute proportion (0.5%) of the mass of the silicate Earth. These elements include the heat producing elements (K, Th, and U) that partially power mantle convection and drive plate tectonics. Thus, quantifying the composition of the upper continental crust, the associated

uncertainties on that composition, and how this composition may have changed through time is an essential step in constraining the Earth's evolving power budget and, from that, mantle dynamics.

Previous attempts to quantify the average composition of the upper continental crust (UCC) have generally followed one of two approaches: systematic grid-based bedrock sampling, or analysis of fine-grained sediments (or sedimentary rocks) such as shales and loess. The former approach typically involves the collection and analysis of thousands of samples and has been carried out in the Canadian Shield (Eade and Fahrig, 1973; Shaw et al., 1976; Shaw et al., 1967), Russia (Borodin, 1998; Ronov and Yaroshevsky, 1967, 1976), China (Gao et al., 1998), and Japan (Togashi et al., 2000). The latter approach assumes that natural geological (sedimentary) processes provide a robust average of insoluble elements in the provenance of the sediments, and has been used in many studies aimed at tracking the compositional evolution of the UCC (McLennan et al., 1979, 1980, 1983; Taylor and McLennan, 1985, 1995; Nance and Taylor, 1976, 1977; Wronkiewicz and Condie, 1987, 1989, 1990; Peucker-Ehrenbrink and Jahn, 2001; Kamber et al., 2005; Chauvel et al., 2014; Sauzeat et al., 2015). Most of these studies use either shale, glacially-derived loess, or desert loess to infer the average composition of the UCC, which works well for insoluble elements, but not for soluble elements. Although the cold temperatures and largely physical weathering that produce the glacial loess limits loss of the soluble elements compared with shales, such loess still records a weathering signature (Gallet et al., 1998), and both types of loess are also affected by wind-driven particle sorting that fractionates heavy minerals, which may dominate the budgets of certain elements (e.g., Ti, Zr, Hf, Sn) (Barth et al., 2000; Liu et al., 1993; McLennan, 2001; Taylor et al., 1983, Chauvel et al., 2014; Sauzeat et al., 2015).

By contrast, use of till or its lithified equivalent, glacial diamictite, may avoid many of the above problems. Like loess, till is produced by physical weathering and thus, may be expected to provide useful information on the soluble element concentrations of the UCC. Unlike loess, there is minimal sorting, as tills are not reworked by wind (or water). Goldschmidt (1933; 1958) first suggested the utility of related glacial loam as a proxy for the average composition of the UCC and the use of till for this purpose was recently explored by Canil and Lacourse (2011) in a study of Quaternary valley till in the Canadian Cordillera, who showed that the tills provided robust averages of the bedrock in the valleys the glaciers sampled.

Here, we report the major, trace element and oxygen isotopic compositions of pre-Cenozoic diamictites deposited by continental ice sheets, and use these diamictites to evaluate the compositional evolution of the UCC, for both soluble and insoluble elements, as well as elements for which few data existed previously (e.g., Be, Ga, Ge, Cd, In, Sn, Sb, W, Tl, Bi). The diamictites studied here were identified by previous workers as being of glacial origin based on the presence of faceted and striated clasts, the presence of dropstones in associated laminated sediments, as well as aspects of stratigraphic facies architecture (e.g., Hambrey and Harland, 1981). We hereafter refer to these rocks as diamictites, with the understanding that all samples investigated here have been proposed to be of glacial origin.

2. Methods

2.1 Samples and their preparation

Glacial diamictites deposited by continental ice sheets are recognized around the world from four pre-Cenozoic intervals (~2.9, 2.4-2.2, 0.75-0.57, and 0.30 Ga); we sampled diamictites

from all four intervals on several modern continents. Stratigraphic units, ages, localities, and references to the original investigations of these units are given in Table 1.

The samples investigated here comprise two rock types: (1) most samples comprise massive (i.e., un-stratified) diamictite, which consists of muddy to sandy matrix material with clasts ranging from pebble to cobble size, and (2) a few samples are dropstone-bearing argillites, which are fine-grained stratified sedimentary rocks spatially associated with the diamictites. In both cases, our analytical campaign focused on the fine-grained matrix material. Pieces with fewer clasts were selected from each sample and crushed to chips in an alumina jaw crusher. Any chips containing clasts larger than ~5 mm were then removed by hand. A fraction of the remaining material was then ground to a fine powder in an alumina swing mill. We endeavored to produce 100 grams of powder for each sample, although this was not possible with some clast-rich varieties.

In addition to preparing individual samples for analysis, we also prepared composite samples of each stratigraphic unit, with the exceptions of the Paleoproterozoic Pecors, Neoproterozoic Blasskranz, and Ordovician Pakhuis formations, for which we have no more than two samples each. Each composite sample consists of equal weights of powder from each individual sample of a stratigraphic unit that was thoroughly mixed together. These composites are intended to serve as a new upper crustal reference suite of samples, comparable to PAAS (post-Archean Australian shale; Nance and Taylor, 1976) and NASC (North American shale composite; Haskin et al., 1966), and represent average compositions of the glacially sampled upper crust found in each stratigraphic unit. The composites are available for further investigations upon request to the second author.

2.2 X-ray Fluorescence Analysis

Major element concentrations and a limited number of trace elements were determined by XRF at Franklin and Marshall College. Loss of ignition (LOI) was first determined at 900°C. From the resulting anhydrous powder, 0.4 g were mixed with 3.6 g of lithium tetraborate flux and fused in a platinum crucible. The fusion disc was used for measurement of major elements, along with Sr, Zr, V, and Cr. Other trace elements were measured on a powder briquette made with copolywax binder. Analyses were conducted on a PANalytical 2404 X-ray XRF vacuum spectrometer with a 4 kW super sharp X-ray tube. Major element results for the composite samples are new, whereas results for the individual samples were previously reported in Gaschnig et al. (2014) and are provided here for completeness (Table 2). Trace elements by XRF are reported and discussed in the Supplemental Material.

2.3 ICP-MS analysis

2.3.1 Sample digestion

Most samples were digested by conventional high pressure HF/HNO₃ dissolution in Teflon Parr bombs, closely following the methods of Zhang et al. (2010b). For each sample, fifty milligrams of powder was weighed into a bomb along with 1 mL and 3 mL of concentrated HNO₃ and HF, respectively. Bombs were then capped and placed in an oven for at least 4 days at ~180°C. Bombs were then opened and acids were evaporated on a hot plate. Two mL of concentrated HNO₃ were added to the salt and evaporated, followed by 1 mL of HNO₃ with 2 mL of H₂O. Bombs were then capped and returned to the oven for one day at 150°C. Samples were transferred to centrifuge capsules and diluted to 15 mL using 2% HNO₃ containing a trace of HF.

A subset of samples (Paleoproterozoic samples from Wyoming and Ontario and Paleozoic samples from Bolivia) was digested using the NH_4HF_2 method developed by Zhang et al. (2010a). In this method, ~ 0.2 g of purified NH_4HF_2 was added to 0.05 g of sample powder with a drop of water in a Teflon screw-top Savillex beaker. Beakers were placed in an oven for 12 hours at ~230°C. Upon returning to room temperature, 2 mL of HNO_3 were added to the solid salt and the beakers were capped and placed on the hot plate at 160°C for at least an hour. Beakers were then uncapped and solutions were evaporated. One mL of HNO_3 and 2mL of H_2O were added to the salt and the beaker was capped and placed on the hot plate for at least 6 hours. Samples were transferred to centrifuge capsules and diluted to 15 mL with 2% HNO_3 containing a trace of HF.

Due to the large range in concentration of the elements analyzed (spanning five orders of magnitude, with Fe and Bi as end members) two dilutions were prepared for analysis from the master sample aliquot. The first dilution was prepared using 0.5 mL of master solution, adding 0.5 mL of an indium standard solution (as an internal drift corrector), and diluting to 15 mL with 2% HNO_3 containing a trace of HF to prevent precipitation of the highly insoluble high field strength elements. The second dilution was prepared by taking 1 mL of the first solution and diluting this to 7.5 mL with 2% HNO_3 . The dilutions were analyzed within 24 hours of preparation in order to avoid precipitation of elements having low solubilities.

2.3.2 ICP-MS protocols

Most samples were analyzed for trace elements on a Thermo-Finnigan Element 2 high resolution-ICP-MS in the Department of Geology at the University of Maryland. Instrumental parameters are given in Supplemental Table A.1. Samples were analyzed with aliquots of USGS whole-rock standards AGV-2, BHVO-1, GSP-1, and W-2 that were prepared at the same time as

the samples. Elemental concentrations in samples were determined by constructing calibration curves with the four USGS standards. Precision and accuracy were evaluated by repeated analysis of separately dissolved AGV-2 aliquots treated as unknowns and are reported in Table 3. Precision was better than 4% relative standard deviation (1σ) for the majority of elements.

A small subset of samples (those from the Konnarock, Nantuo and Gucheng formations) were analyzed on an Agilent 7700x quadrupole ICP-MS in the State Key Laboratory for Geological Processes and Mineral Resources at the China University of Geoscience in Wuhan, China and these are shown in italics in Table 2. Instrumental parameters are given in Supplemental Table A.1. Calibration was conducted using multi-element standard solutions to create a calibration line, and then adjusting this calibration after analysis of dissolved whole-rock standards. Precision and accuracy are discussed in Hu and Gao (2008).

2.3.3 Standard addition analysis

A subset of samples was analyzed by standard addition ICP-MS for Ga, Ge, Mo, Cd, In, Sn, Sb, W, Tl, and Bi, since the concentrations of these elements are relatively poorly constrained in the whole-rock standards used for calibration. Two spikes were prepared from single-element standard solutions, one containing Ga, Ge, Mo, Sn, and W, and one containing Cd, In, Sb, Tl, and Bi. These two spikes were prepared due to the large difference in concentrations found in typical crustal rocks and the potential for polyatomic interferences between the groups (e.g., Mo-oxides on Cd). Four solutions were analyzed per sample (for a given spike) consisting of: 1) a total analytical blank; 2) an unspiked aliquot of sample; 3) a spiked aliquot of sample; and 4) an aliquot of a sample spiked with twice the volume of spike as the previous one. All aliquots were also spiked with a Rh solution, which was used for internal drift correction. Precision and accuracy of this method based on analyses of AGV-2 as an

unknown are presented in Table 3. Further details of this approach, including data for other standard reference materials, are given in Gaschnig et al (2015).

2.4 Oxygen isotope analysis

Whole-rock powders were analyzed for oxygen isotope ratios at the University of Wisconsin-Madison by laser fluorination (Valley et al., 1995) using an air-lock sample chamber (Spicuzza et al., 1998) with BrF₅ as a reagent and a 25W CO₂ laser ($\lambda = 10.2 \mu\text{m}$). By this means, 2-3 mg samples of powder were outgassed at vacuum overnight and fluorinated individually so as to prevent cross-contamination. Evolved oxygen was cryogenically cleaned, passed through heated Hg vapor, and converted to CO₂ over hot graphite for analysis. Isotope ratios were measured on a Finnigan MAT 251 dual inlet mass spectrometer and are reported in standard permil notation relative to VSMOW (Vienna Standard Mean Ocean Water). Measured values were standardized using the UWG-2 garnet standard ($\delta^{18}\text{O} = 5.80\text{‰}$ VSMOW; Valley et al., 1995). Overall reproducibility of the standard during the analytical sessions was $\pm 0.18\text{‰}$ (2 SD).

3. Results

3.1 Major elements

The major element compositions of siliciclastic sediments such as diamictites reflect the relative proportions of quartz to feldspars and clays/phyllosilicates, and the majority of diamictites falls along the expected mixing trend in a plot of SiO₂ vs. Al₂O₃ (Fig 1a). Several samples deviate from this trend and are characterized by either unusually high CaO or Fe₂O₃(T) (Figs. 1b and 1c). Although these outlier samples encompass all four age groups, they all share a common origin in southern Africa. The high CaO samples also have high LOI values (see

supplemental figure A.2), consistent with high levels of carbonate. The high levels of carbonate and Fe have a dilution effect on the other major elements.

3.2 Trace elements

Trace element concentrations are plotted in Figures 2-4, organized by element groups. In order to account for the effects of dilution from quartz, carbonate, or Fe on absolute abundances of trace elements (see above), samples are first normalized to the average UCC abundance of Al_2O_3 (15.26%) of Rudnick and Gao (2003). Aluminum is used for this purpose because it is an immobile element with a relatively restricted range in crustal igneous rocks (i.e., both basalts and granites commonly have ~15 wt% Al_2O_3). This normalization is done by dividing 15.26 by the measured Al_2O_3 in each sample, which results in a value greater than one for this “adjustment factor” in samples that are affected by quartz, carbonate, or Fe dilution. We then multiply the concentrations for each trace element by the adjustment factor. Data for individual elements are then normalized to the average UCC of Rudnick and Gao (2003) (hereafter referred to as UCC_{RG}) (with the exception of the chondrite normalized left half of Fig. 2), which provides a frame of reference. For the sake of clarity, only composites are plotted in Figures 2-4. This is justified by the close comparison between the calculated medians of individual samples for a given formation and their composites. Further comparison of the individual sample results to the composites is shown in supplemental Figure A.1.

3.2.1 Rare earth elements (REE)

The REE are incompatible, insoluble lithophile elements that show a systematic increase in compatibility with increasing atomic number, with the exception of Eu, which mirrors the behavior of Ca and Sr when in a 2+ oxidation state. REE patterns, including Y (which behaves

like a heavy REE and is plotted between Ho and Er), are shown in Figure 2. The diamictites all show chondrite-normalized REE patterns (Figs. 2a) that are characterized by light REE (LREE)-enrichment and negative Eu anomalies (Eu anomaly or $\text{Eu}/\text{Eu}^* = (\text{Eu}/(\text{Gd} \cdot \text{Sm})^{0.5})$, with all concentrations normalized to chondrite), save one Mesoarchean sample that has a positive Eu anomaly. Differences are more apparent when the results are normalized to UCC_{RG} (Figs. 2b). Three of the Mesoarchean composites have normalized REE abundances and LREE to heavy rare earth element (HREE) fractionation similar to that of UCC_{RG} , whereas the fourth is depleted in the LREE but not the HREE relative to UCC_{RG} . All Archean composites are enriched in Eu relative to UCC_{RG} . The majority of the Paleoproterozoic composites show normalized concentrations and LREE to HREE fractionation comparable to UCC_{RG} , whereas Eu may be slightly enriched or depleted relative to UCC_{RG} . The Neoproterozoic composites differ from the older ones in that many show a positive slope when normalized to UCC_{RG} , indicating less fractionation between the LREE and HREE, and all are Eu depleted. In addition, the Neoproterozoic Idaho composite is distinctly more enriched in all REE than the other composites of all ages. The Paleozoic composites are indistinguishable from the Neoproterozoic composites, although Dwyka West has lower overall abundances and is not depleted in Eu.

3.2.2 Transition and post-transition metals and metalloids

The transition metals are generally compatible, may exhibit multiple valence states, and may show lithophile, chalcophile, and/or siderophile characteristics. The post-transition metals and metalloids exhibit both compatible and incompatible behavior and are primarily chalcophile. The transition metal concentrations change as a function of the age of the sediment (Fig. 3a). Mesoarchean composites are generally enriched in all transition metals relative to UCC_{RG} but the enrichment is significantly greater for Cr and Ni relative to the other metals. Paleoproterozoic

composites show mild enrichment in Cr, but otherwise are similar to UCC_{RG}. Neoproterozoic composites are generally depleted in Cr, Co, and Ni but show similar Sc and V concentrations to UCC_{RG}. Of the Paleozoic composites, the western Dwyka shows significant enrichment in all transition metals, comparable to the Mesoarchean composites. The other Paleozoic composites have transition metal levels overlapping those of the Neoproterozoic composites and similar to UCC_{RG}.

The post-transition metals and metalloids are plotted, along with transition metals Cu and Zn, in Figure 3b. Copper concentrations vary more than those of all other elements (roughly two orders of magnitude). Although the Mesoarchean composites are all enriched in Cu, the younger composites show both enrichment and depletion relative to UCC_{RG}. Paleoproterozoic composites are particularly notable for the large range of normalized Cu concentrations observed. Zinc is also quite variable in the Paleoproterozoic composites but composites from other periods show a narrower range of concentrations and are consistently enriched relative to UCC_{RG}. Cadmium also shows significant variability but tends to be enriched in most composites relative to UCC_{RG}. Compared to all other post-transition metals, concentrations of Ga and In, which belong to the same group as Al on the Periodic Table, show little variability and match UCC_{RG} estimates. Germanium concentrations vary widely, especially in the Paleoproterozoic and Mesoarchean composites, and are both enriched and depleted relative to UCC_{RG}; there is no correlation between Ge and SiO₂ contents. Tin is depleted relative to UCC_{RG} in the Mesoarchean and some of the Paleoproterozoic composites, but is enriched in all other composites. Antimony concentrations are highly variable in diamictites of all time periods, and show no obvious trend with age, but tend to be enriched in the diamictites relative to UCC_{RG}.

3.2.3 Alkali and alkaline earth elements

The alkali and alkaline earth elements are incompatible and relatively soluble lithophile elements; their normalized concentrations in the diamictite composites are shown in Fig. 4a. Lithium is enriched relative to UCC_{RG} in nearly all composites, and Rb is generally enriched in all post-Archean samples. Cesium is highly variable, showing both large enrichments and depletions in different composites with no obvious trend with age. Beryllium concentrations appear to increase with age. Strontium is depleted relative to UCC_{RG} in all composites save the carbonate-rich Neoproterozoic Ghaub Formation. Barium concentrations show a relatively narrow range in the diamictites, with some suggestion of an increasing concentration over time with respect to UCC_{RG} .

3.2.4 High field strength elements (HFSE)

The HFSE are incompatible and primarily lithophile elements, although Mo and W may show chalcophile and siderophile characteristics, respectively. Normalized HFSE concentrations for the composites are shown in Fig. 4b. Zirconium and Hf concentrations show limited variability and are close to UCC_{RG} , with the exception of the Neoproterozoic Pocatello (Idaho) and Blaubeker formations that are enriched in Zr and Hf. Niobium and Ta concentrations vary more widely and show an increase with time relative to UCC_{RG} . Several Neoproterozoic composites show significant enrichment in Nb and Ta, and the enrichment is especially large (~10 times UCC_{RG}) in the Pocatello composite. Molybdenum concentrations show significant variability within and between the different time periods, but are generally higher in the Mesoarchean and Paleoproterozoic than the Neoproterozoic and Paleozoic, with the exception of the Mo-enriched western Dwyka composite. Tungsten concentrations are depleted in the Mesoarchean and Neoproterozoic composites relative to UCC_{RG} , but are similar to or enriched relative to UCC_{RG} in the Paleoproterozoic and two of the three Paleozoic composites.

3.2.5 Heavy post-transition metals and actinides

The heavy post-transition metals (Tl, Pb, and Bi) (Fig. 4c) are incompatible elements with both lithophile and chalcophile characteristics. The actinides (Th and U) are incompatible lithophile elements. Concentrations of all five elements vary widely in the composites of all periods except the Paleozoic. Thallium is either similar to or depleted relative to UCC_{RG} in all periods. Lead concentrations are either higher or lower than UCC_{RG} and show no temporal trend. Bismuth concentrations are enriched relative to UCC_{RG} for most composites. Thorium and U concentrations are depleted in the Mesoarchean but enriched in the other periods. Th/U increases systematically from the Mesoarchean, which has lower Th/U than UCC_{RG} , to the Neoproterozoic, which has higher Th/U than UCC_{RG} . By contrast, Th/U of the Paleozoic samples are similar to UCC_{RG} .

3.3 Oxygen isotopes

Figure 5a shows the $\delta^{18}O$ values for individual glacial diamictite powders as a function of age, after filtering the data set to exclude samples with LOI values greater than 7%. (We exclude these samples to eliminate the effects of carbonate, which leads to abnormally high $\delta^{18}O$ values. See supplemental figure A.2 for justification of this cut-off criterion). The oxygen isotope ratios of the diamictites tend to become heavier and more variable with time. Values of $\delta^{18}O$ are the lowest in the Archean diamictites, which overlap with the modern mantle. Most of the Paleoproterozoic diamictites have $\delta^{18}O$ values that overlap with those of the higher $\delta^{18}O$ Archean diamictites, but diamictites from the Makganyene and Duitschland formations from South Africa have considerably higher $\delta^{18}O$, ranging from 13 to 18‰. The majority of Neoproterozoic samples have values between 10 and 15‰. Several of the Paleozoic samples overlap with the main Neoproterozoic group but a few samples have lower $\delta^{18}O$ values.

Figure 5b shows $\delta^{18}\text{O}$ of the diamictites versus their chemical index of alteration values (CIA = molar Al_2O_3 / ($\text{Al}_2\text{O}_3 + \text{CaO}^* + \text{K}_2\text{O} + \text{Na}_2\text{O}$), where CaO^* is corrected for apatite and carbonate; carbonate correction using approach of McLennan, 1993) (Nesbitt and Young, 1982), which are a measure of chemical weathering intensity (see below). While a positive correlation is present for the Neoproterozoic and Paleozoic diamictites, $\delta^{18}\text{O}$ and CIA are largely decoupled in the older diamictites. In particular, the Mesoarchean samples have very high CIA but mantle-like $\delta^{18}\text{O}$.

4. Discussion

The efficacy of using glacial diamictites to track changes in the average UCC composition is contingent upon two main factors. First, the diamictites must preserve the original composition of the eroded crustal material without being significantly modified by syn- or post-depositional processes such as chemical weathering. Second, the diamictites must consist of material pulverized by ice sheets from large continental areas, as opposed to preferentially sampling local sources near the ice sheet terminus. We consider these two factors below, and then discuss evidence in the diamictites for secular changes in the average UCC composition and finally, use our new data to derive an average UCC composition, along with uncertainties, and compare that with previous results.

4.1 Identifying and seeing through the effects of chemical weathering

Sediments and sedimentary rocks frequently show the effects of chemical weathering, which obscures information about the bulk composition of their sources. Several geochemical indices have been used to recognize chemical weathering effects, the most popular of which is the chemical index of alteration. Fresh igneous rocks have CIA values ranging from 40 to 55

and pure kaolinite and chlorite have a value of 100, owing to their lack of Ca, Na, and K. Gaschnig et al. (2014) noted that the majority of diamictites have CIA values elevated above those of igneous rocks; they also attributed the pervasive Sr depletion seen in the diamictites to the influence of chemical weathering. This weathering signature appears to be inherited from the sediment source in most cases, rather than reflecting post-depositional changes, or weathering during glacial erosion and transport (Gaschnig et al., 2014; Li et al., 2016).

A weakness of using the CIA value to assess chemical weathering effects is its inability to take into account the original bulk composition of the protolith. This is especially problematic at low CIA because unweathered igneous rocks can show a range of CIA values: basalts tend to have the lowest CIA, around 40, whereas granites have higher CIA, up to 55. One way of extracting information on source lithologies from major element data of sedimentary rocks, which also show the influence of chemical weathering, is to pair the CIA value with another index called the ICV (index of chemical variability) (Cox et al., 1995), which is defined as $ICV = (TiO_2 + Fe_2O_3(T) + MnO + MgO + CaO + K_2O + Na_2O) / Al_2O_3$ (with all oxides as weight percent). ICV values are higher for more mafic source lithologies and when plotted against CIA values (Figure 6) (Potter et al., 2005), straight lines connecting a completely weathered end member (pure kaolinite/chlorite) to average igneous compositions reflect weathering trends. The majority of the diamictite data plot on a broad weathering trend consistent with an average andesitic source region, which is close to the expected granodioritic average composition of the UCC. However, many Mesoarchean and Paleoproterozoic samples fall along trends suggestive of a basaltic or komatiitic average source composition, indicating that the UCC from these earlier time periods was more mafic.

Another method used for separating source composition from weathering affects in sedimentary rocks was developed by Ohta and Arai (2007). This approach uses principle component analysis to assign and weight all major element oxides into mafic (M), felsic (F), and weathered (W) end members. Individual diamictites are plotted in this framework in Figure 7. As in Figure 6, the majority of samples plot in a broad field reflecting a weathering trend projecting to intermediate composition igneous protoliths, except for a select group of Paleozoic Dwyka samples (the west Dwyka) and nearly all of the Archean samples. These samples form a weathering trend that projects to a mafic igneous protolith. Thus, although the diamictites all bear chemical weathering signatures of various intensities, it is possible to use their major elements to identify a major change in the nature of the crust being weathered and eroded between the Archean and Paleoproterozoic.

4.2 Identifying and seeing through the effects localized or non-representative provenance

Research on modern and Pleistocene glacial erosion and sedimentation has shown that large ice sheets are capable of transporting sediment great distances, but that sediment from upstream sources may become diluted by material from nearby sources with increasing downstream distance (e.g., Clark, 1987; Klassen, 1999; Hildes et al., 2004; Farmer et al., 2006). Dispersal distance appears to be controlled primarily by basal ice velocity and bedrock topography (Clark, 1987). The effects of localized provenance are seen on the geochemistry of some of the diamictites. The two most prominent examples are found in the Pocatello Formation and Dwyka Group. Diamictites from the Neoproterozoic Pocatello Formation are unusually enriched in Nb and Ta (by an order of magnitude relative to UCC_{RG} , Fig. 4b), and the REEs, especially the LREEs (enriched by a factor of two relative to UCC_{RG} , Fig. 2). These enrichments reflect the contribution from local Neoproterozoic bimodal alkaline volcanic rocks formed

during the rifting of Rodinia, which are strongly enriched in these elements (Yonkee et al., 2014). The influence of these alkaline rocks is supported by the large number of detrital zircons of the appropriate age found in this diamictite (Yonkee et al., 2014), and the high concentrations of Zr and Hf (Fig. 4b). Despite this clear local signature, the Pocatello Formation also shares distinctive chemical characteristics that are common to all Neoproterozoic diamictites, such as the depletion of V, Sc, Mo, and especially Cr and Ni relative to REE.

Another example of the influence of localized provenance is found in the Dwyka Group, where major geochemical differences exist between diamictites collected in separate parts of the Karoo Basin. Samples from the western Dwyka have significantly higher concentrations of the compatible transition metals (Sc, V, Cr, Co, Ni, Cu, Zn, Mo, Cd) and lower concentrations of the REE than samples from the southeastern Dwyka. These chemical characteristics appear to reflect differences in provenance governed by ice flow directions. Detrital zircons from the west Dwyka samples are 2.0 Ga and older (Gaschnig et al., 2015), indicating that the ice in this region was exclusively sampling the Archean interior of the Kaapvaal craton (plus the Paleoproterozoic Bushveld Complex). By contrast, a large diversity of detrital zircon ages, ranging from Early Paleozoic to Archean, are found in the southeastern portion of the Dwyka Group (Vorster, 2013), indicating both a greater diversity of sediment sources and a dominance of Mesoproterozoic and younger ages. Thus, the transition metal enrichments (Fig. 3a) and the major element evidence for a much more mafic source rock (Fig. 7) for the west Dwyka diamictites relative to the remainder of the Dwyka samples both reflect sampling of Archean crust by the Paleozoic glaciers in the west.

Despite the influence of localized provenance seen in the geochemistry of certain samples discussed above, there are common geochemical features seen globally amongst similarly aged

samples. This is especially apparent for elements such as the REE, which show similar fractionation patterns and abundances for each time period, and the transition metals, which show relatively uniform levels of enrichment or depletion in each time period. Furthermore, many of the other diamictites for which detrital zircons have been dated show age spectra with a wide range of ages, indicating a diversity of sediment sources (e.g., Liu et al., 2008; Hofmann et al., 2015). Consequently, the majority of the diamictites appear to consist of sediments drawn from large continental areas, making them reliable proxies for the average UCC for many elements. We next explore the key geochemical features seen amongst diamictites of different ages.

4.3 Secular changes in diamictite compositions

Systematic secular changes in the compositions of the diamictites are observed that reflect changes in the average composition of the UCC through time. We explore these changes below in plots of elemental ratios versus age for insoluble elements in the composite samples (Figs. 8 through 12), and measure the robustness of these changes through statistical analysis. We used Kolmogorov-Smirnov tests and normality plots to determine if variables met the assumptions of a parametric test and log transformed the ratios when necessary to improve normality. We used MANOVA (multivariate analysis of the variance) to compare normally distributed ratios among the four time periods and used post-hoc univariate ANOVAs and Tukey tests to identify significant differences among periods. All tests were two-tailed and we set $\alpha = 0.05$. For ratios that were not normally distributed, we used a Kruskal-Wallis test to compare ratios among the different time periods. Analyses were performed in PASW 18 (SPSS). Results are provided in the Supplemental Tables A2 and A3. All of the element ratios examined differed significantly among the four time periods (all $p < 0.05$), with the exception of Gd/Lu.

4.3.1 Evidence for increasingly differentiated and felsic UCC with time

As noted above, techniques using major element composition to separate sediment source from weathering effects reveal that the Mesoarchean and some of the Paleoproterozoic diamictites have distinctly more mafic provenances that include komatiites (Figs. 6 and 7). Moreover, from the Archean to the present, diamictites show a secular decrease in the abundance of the compatible first row transition metals (Sc, V, Cr, Co, and Ni, Fig. 3), a feature also documented in Figure 4 of Gaschnig et al. (2014). These transition metals are contained primarily in mafic minerals, and the depletion of these elements with decreasing age is consistent with the UCC becoming less mafic with time. We can summarize this observation using the Ni/Lu ratio plotted against deposition age (Fig. 8a), where Lu is used as a reference element having limited absolute variation (in the same way that Al is used in the multi-element plots in Figs. 2-4). Ni/Lu ratios in Mesoarchean diamictites are distinctly higher than all other composites, demonstrating that these diamictites were sourced from a UCC with high Ni content. Diamictites from the post-Archean glaciations show successively lower Ni/Lu ratios (with the exception of the western Dwyka samples previously discussed). These observations are consistent with the UCC being more mafic in the Archean and becoming progressively more differentiated and felsic in composition with time (Tang et al., 2016).

The Th/Sc ratio is used to differentiate between mafic and felsic provenance of sediments, given the compatibility of Sc and the incompatibility of Th in igneous systems (e.g., Taylor and McLennan, 1985). Sediments from granitic source rocks thus have higher Th/Sc ratios than those from basaltic source rocks. The Mesoarchean diamictite composites show the lowest Th/Sc ratios, which is consistent with the other lines of evidence indicating a predominantly mafic source (Fig. 8b). In contrast to the continuous change with time seen in the

Ni/Lu ratio, the Th/Sc ratio increases between the Mesoarchean and Paleoproterozoic but does not show a continuous secular trend.

The Eu/Eu* ratio is another measure of intracrustal differentiation. This is because Eu²⁺ enters feldspar relative to Eu³⁺, which leads to the development of a negative Eu anomaly (Eu/Eu* < 1) in evolved magmas derived from melting or differentiation within the crust of typical thickness (≤35 km) (e.g., Taylor and McLennan, 1985; Rudnick and Gao, 2003; Tang et al., 2015). As with the Ni/Lu ratios, the diamictites show a continuous decrease in Eu/Eu* with time (Fig. 8c). This suggests that intracrustal differentiation became increasingly important through time and is consistent with a greater proportion of mafic, undifferentiated magmas making up the bulk of the Archean UCC.

The oxygen isotope compositions of the diamictites likely reflect the original δ¹⁸O of the provenance, which, in turn, reflects its chemical weathering or alteration history. Low temperature interactions with water at or near the Earth's surface drives the oxygen isotopic compositions of rocks towards higher δ¹⁸O values, starting from the mantle δ¹⁸O value of 5.6 ± 0.3 ‰ (Mattey et al., 1994; Valley et al., 1998; Eiler, 2001; Grimes et al., 2011). As noted above, the δ¹⁸O of the diamictites generally increase with time (Fig. 5a). The oxygen isotope composition of Mesoarchean diamictites have both relatively low δ¹⁸O values and a limited range (Fig. 5a). These samples overlap in part with the δ¹⁸O composition of the modern mantle and are lower, on average, than the published values for Archean siliciclastic sediments compiled by Payne et al. (2015) and Bindeman et al. (2016). The δ¹⁸O values of the Archean diamictites are also decoupled from the CIA values, which are high (~70 to 90) (Fig. 5b). The persistence of the low Archean δ¹⁸O values suggests either that weathering processes did not significantly contribute to their oxygen isotope compositions, or that Archean surface waters had a

significantly different oxygen isotope composition than today (Knauth and Lowe, 2003; Muehlenbachs, 1998).

The main exceptions to the increase in $\delta^{18}\text{O}$ with time are the very high values from the Paleoproterozoic Makganyene and Duitschland formations. These units tend to have high CIAs and are dominated by marine sedimentary clasts (carbonate and chert), suggesting that their heavy oxygen isotope compositions are influenced by incorporation of a greater proportion of preexisting sediments compared to other diamictites that contain an assortment of clast types and are often rich in igneous and meta-igneous clasts.

The increase in the average $\delta^{18}\text{O}$ values of the diamictites with time (Fig. 5a) and the increase in the range of $\delta^{18}\text{O}$ values after the Mesoarchean is broadly consistent with both the igneous $\delta^{18}\text{O}$ record provided by zircons (Valley et al., 2005) and the compilation of existing whole-rock siliciclastic sedimentary data of Payne et al. (2015) and Bindeman et al. (2016). Both studies attributed the parallel changes in the igneous and sedimentary oxygen isotope record of continents to the increased production of clay and incorporation of it into sedimentary and igneous rocks. This increase in clay production may have been driven by a combination of crustal thickening and crustal growth, both of which led to greater amounts of emergent continental material available for weathering (Arndt, 1999; Rey and Coltice, 2008; Gaillard et al., 2011).

In summary, the above observations from major and trace elements indicate that the crust sampled by the ice sheets in the Archean was more mafic and less differentiated compared to later time periods. The similarity of these results to previous studies of Archean UCC geochemistry based on shales and map-based methods (e.g., Taylor and McLennan, 1985; 2010; Condie, 1993) strongly suggest that the diamictites are faithful proxies for the UCC composition.

The presence of a negative Eu anomaly (albeit a small one) in all but one of the oldest diamictite units is consistent with Condie's (1993) contention that the average Archean UCC had a negative Eu anomaly, and its growth with time supports the hypothesis that intracrustal differentiation increasingly became more important from late Archean onward (Taylor and McLennan, 1985). The increase in $\delta^{18}\text{O}$ values with time is also consistent (at least indirectly) with increasing recycling of continental crust from the Mesoarchean onwards, possibly the result of a thicker and more voluminous emergent crust (Payne et al., 2015).

4.3.2 Changes in the concentrations of heat producing elements

Thorium and U are two of the three major heat-producing elements in the Earth. While these elements are concentrated in the continental crust and depleted in the mantle, the amount remaining in the mantle after crust extraction plays an important role in driving convection and plate tectonics. It is for this reason, and their role as progenitors of several radiogenic Pb isotopes, that constraining the concentrations of Th and U in the upper crust is of great importance to understanding Earth differentiation and evolution.

Thorium and U are similarly incompatible during mantle melting, so the Th/U ratio in juvenile crust is expected to match the mantle ratio. While Rudnick and Gao (2003) inferred an average Th/U of 3.8 for the UCC, there are reasons to believe that the abundance of these heat-producing elements have changed through time. McLennan and Taylor (1980) and Taylor and McLennan (1985) noted a secular increase of Th/U for fine-grained sedimentary rocks, with average values between 3.5 and 4.0 for the Archean while Phanerozoic values were ~5. They attributed this trend to increased recycling of sedimentary material under oxidized atmospheric conditions after the Great Oxidation Event (GOE) around 2.4 Ga (e.g., Holland et al., 2006), leading to conversion of U^{+4} to soluble U^{+6} and its loss from the UCC to the oceans through

weathering. Similarly, Collerson and Kamber (1999) documented a complementary decrease in the Th/U of depleted mantle through the same interval, which they attributed to increased oxidation of Earth's surface and preferential recycling of U.

The normalized concentration of Th in the diamictites shows a step-wise increase between the Mesoarchean and Paleoproterozoic (Fig. 9a), which (as mentioned above) is consistent with a more felsic and differentiated UCC after the Archean. On the other hand, the concentration of U shows a more subtle increase after the Archean and generally shows more scatter (Fig. 9b). The Th/U ratios of the diamictites are lower in Archean and Paleoproterozoic diamictites compared to those of the Neoproterozoic and Paleozoic (Fig 9c) but they do not show a smooth temporal change. The Neoproterozoic samples are distinct in that they show both the greatest variation and the highest average Th/U. The temporal changes in Th/U reflect a combination of the step-wise increase in Th after the Archean and the depletion of U in most of the Neoproterozoic, Paleozoic, and one of the Paleoproterozoic (Deutschland) diamictites. The U depletion is most likely the result of oxidative weathering. Overall, the diamictite results are consistent with other UCC proxies in showing an upward redistribution of Th and U into the UCC due to enhanced intracrustal differentiation after the Archean and then loss of U to the oceans via oxidative weathering.

4.3.3 Changes in the fractionation of REEs

With the exception of Eu (discussed above), the REEs display coherent behavior during geochemical processes due to their identical oxidation states (under igneous conditions) and progressively decreasing ionic radii with increasing atomic number, which results in more incompatible behavior of the LREEs compared to the HREEs. The pressure at which partial melting and/or crystal fractionation occurs is one of the main processes controlling the

fractionation of REEs, as the presence of garnet at higher pressures results in high LREE/HREE ratios in melts in equilibrium with these minerals (Drummond and Defant, 1990). Such TTGs (tonalite-trondhjemite-granodiorite associations) are a major constituent of Archean crust (e.g., Moyen and Martin, 2012). Ratios reflecting the fractionation between light, medium, and heavy REEs as a function of time are shown in Figures 10a, c, and e. The Neoproterozoic Pocatello (Idaho) diamictites and one of the Mesoproterozoic units (the Afrikander Formation of the Witwatersrand Supergroup) are prominent outliers, which are attributable to the influence of local incompatible element-enriched rift-related igneous rocks in the case of the former (Yonkee et al., 2014) and a more mafic/ultramafic provenance of the latter (relative to the other Archean units). Even excluding these outliers, La/Lu, representing overall LREE/HREE fractionation, shows significant overlap amongst the four time periods (Fig. 10a). The LREEs show more resolvable differences amongst the four periods, with a peak in La/Sm in the Paleoproterozoic followed by a drop to lower values in the Neoproterozoic and Paleozoic (Fig. 10c). Decreasing fractionation of LREE with time is consistent with the shift from a granitic component dominated by TTG magmatism in the earlier Archean and Hadean, to “modern-style” K-rich granitic magmatism in the Neoproterozoic during a period of global cratonization (e.g., Whalen et al., 2004; Condie and O’Neill, 2011; Arndt, 2013; Laurent et al., 2015), a shift that is observed in a statistical analysis of the geochemistry of granites through time (Keller and Schoene, 2012). The HREEs do not show significant differences in fractionation amongst the four time periods, as seen in the Gd/Lu (Fig. 10e) (the latter of which is supported by a statistical test indicating data from the four time periods belong to the same population). While Archean TTGs have high Gd/Lu, they also have low concentrations of HREEs relative to Archean basalts, which tend to have flat HREE patterns (e.g., Condie, 1993). Thus, if the Gd/Lu of Archean diamictites reflects

a mixture of basalt and TTG, this can explain the lack of fractionation between the Archean and post-Archean UCC.

In addition to secular changes in the fractionation of LREEs, the concentrations of the REEs (normalized to Al) change with time (Figs. 10b, d, and f). Although there is significant overlap in La concentrations in the different time periods, Gd and especially Lu concentrations increase in younger units. This temporal increase is likely a reflection of the increasingly felsic and differentiated composition of the UCC, with the particularly marked changes in Lu (and other HREEs) likely reflecting an increasingly smaller proportion of the UCC comprised of HREE-depleted TTGs, as described above.

4.3.4 Sm-Nd and Lu-Hf

The Sm/Nd and Lu/Hf ratio are important in the study of planetary differentiation and crust formation due to their radiogenic isotopic systems (i.e., decay of ^{147}Sm to ^{143}Nd , ^{146}Sm to ^{142}Nd , and ^{176}Lu to ^{176}Hf). The Sm/Nd ratios of the diamictites show an increase between the Paleoproterozoic and Neoproterozoic (Fig. 11a), but this is an expected outcome given the change in LREE fractionation seen in Figure 10c, and is probably attributable to the same causes. The Sm/Nd values of the diamictites are higher than in the UCC model of Rudnick and Gao (2003), and references therein; however the diamictites do not exceed the range of other published models, particularly the higher values of Kamber et al. (2005) and Chauvel et al. (2014). The diamictites show a small increase in Lu/Hf with time, but also show a much greater range for each period compared to Sm/Nd, probably due to the significantly different geochemical behavior of Lu-Hf compared to the Sm-Nd pair (Fig. 11b).

4.3.5 HFSEs

The high field strength elements (HFSE), including Zr, Hf, Nb, and Ta, are highly incompatible and enriched in the continental crust relative to the rest of the silicate Earth, but they are also insoluble in aqueous fluids and consequently are fractionated from other similarly incompatible elements (e.g., alkali and alkaline earth) during subduction zone processes (e.g., Pearce and Peate, 1995). In addition, the pairs Zr-Hf and Nb-Ta share oxidation states and near identical ionic radii (4+, 715 pm; 5+, 640 pm, respectively) and are therefore difficult to fractionate from one another. This coherent behavior has led to their use in studies of Earth differentiation and crustal growth (e.g., Green et al., 1995; Barth et al., 2000; Pfander et al., 2007; Hoffmann et al., 2011).

Zr/Hf shows no secular variations, whereas Nb/Ta and La/Nb ratios of the diamictites show a jump between the Paleoproterozoic and Neoproterozoic, with older diamictites having lower Nb/Ta and higher La/Nb (Fig. 12). The latter observation is particularly interesting in light of the temporal increase in La concentrations (Fig. 10b), as it requires an even greater increase in Nb concentration in order to explain the increase in the La/Nb ratio. The lower Nb seen in the Paleoproterozoic and Mesoarchean diamictites is consistent with the other lines of evidence listed above for a greater TTG influence on the Mesoarchean and Paleoproterozoic diamictites. Although TTGs have a wide range of Nb/Ta ratios, they generally are skewed towards significantly lower ratios (e.g., Condie, 2005; Moyen and Martin, 2012). The secular changes in La/Nb and Nb/Ta ratios point to differentiation processes driving the element fractionation. The evidence from Hoffmann et al. (2011) and Marschall et al. (2013) is suggestive of a role for residual rutile in TTG genesis (McDonough 1990; Rudnick et al., 2000).

4.4 Reexamining the integrated average UCC

The results presented here indicate that the UCC experienced a major change in composition between the Mesoarchean and Paleoproterozoic glaciations, with more subdued changes between the Paleoproterozoic and Neoproterozoic glaciations. There is no evidence from the diamictites to indicate that the UCC changed significantly in composition between the Neoproterozoic and Paleozoic, consistent with the lack of evidence for changes in crust formation and modification processes in the last billion years. Therefore, the Neoproterozoic and Paleozoic diamictites together should provide information on the average global UCC today. Below, we compare the results obtained from the Neoproterozoic and Paleozoic diamictites to previous estimates of the average UCC.

4.4.1 The major element composition of the UCC

Using the Neoproterozoic and Paleozoic diamictite samples as representative of the present-day UCC, we calculate its average major element composition. Table 4 compares UCC_{RG} to two new estimates (unfiltered and filtered) for the average UCC based on two appraisals of the Neoproterozoic and Paleozoic diamictite data. We report the median and the median absolute deviation (MAD, defined as the median of the absolute deviations of the measurements from the population median) rather than the arithmetic mean and standard deviation because the former is less influenced by outliers. Both medians are normalized to 100% anhydrous (omitting the LOI) in order to properly compare them to existing major element models. The first estimate (unfiltered) is calculated based on all Neoproterozoic and Paleozoic diamictite samples. However, a weakness of this approach is that some Namibian Neoproterozoic and the South African Paleozoic diamictites contain significant carbonate, leading to high CaO (and MgO) and unusually low values for the other major element oxides, making them clear outliers, as exemplified in Figure 1. The second estimate (filtered) corrects

for carbonate enrichment, with the median major element values providing a more robust estimate of the average UCC composition.

Major element oxide medians for the Neoproterozoic and Paleozoic are shown normalized to UCC_{RG} in Figure 13. The filtered dataset has slightly lower MgO and CaO than the unfiltered dataset but the other major element oxides are nearly identical and all are within error of one another, indicating that carbonate-rich samples do not significantly skew the results. Major differences between the Rudnick and Gao (2003) model and the diamictite-derived model are the significantly lower MgO, CaO, and Na_2O , and marginally higher K_2O of the latter. The low MgO, CaO, and Na_2O values are almost certainly the result of chemical weathering, as these major elements are highly soluble and unlikely to be fixed in secondary minerals formed during incongruent weathering (whereas K_2O can be retained if kaolinite reacts with K^+ in pore water to form illite, Fedo et al., 1995). The depletion of these elements relative to UCC_{RG} is not surprising given that this and other models cited therein are based largely on sampling crystalline bedrock of continental shields. Use of crystalline rock compositions to determine the average UCC ignores the fact that some fraction of the UCC consists, by definition, of sediments and some fraction is chemically weathered. Accordingly, the diamictite-based model may provide a more realistic estimate of the major element composition of the UCC. However, an important question is how deeply this weathering signature extends into the crust. The thickness of UCC is typically assumed to be no more than about 10 km (e.g., Taylor and McLennan, 1985), or the approximate upper one third of the continental crust (e.g., Rudnick and Fountain, 1995). It is likely that the signature of chemical weathering that is so obvious in the diamictites attenuates with depth, but the shape of this attenuation profile is unknown and is difficult to determine.

4.4.2 The trace element composition of the UCC

A new diamictite-based estimate for the trace element composition of the UCC is presented in Table 4 based on the median values of the Neoproterozoic and Paleozoic samples. Trace element medians are shown normalized to UCC_{RG} in Figure 14. The filtered and unfiltered dataset yield median values that are within the unfiltered median absolute deviations, although the filtered dataset medians have smaller deviations. Compared to UCC_{RG} , the diamictite-based model yields significantly higher Li, Rb, Y, MREE, and HREE, all of which are 20 to 60% higher. Transitional metals Cr, Co, and Ni, along with W and Tl, are roughly 40% lower, and Sr and Mo are depleted by roughly 70% relative to UCC_{RG} . The depletions in Sr and Mo have already been discussed and are attributed to the effects of chemical weathering (Gaschnig et al., 2014) and the other deviations from UCC_{RG} are still within the range of other UCC models (Fig 14b). In particular, the diamictite medians for Li, Rb, Y, MREE, HREE, W, and Tl are within uncertainty of the Australian alluvium-based model of Kamber et al. (2005), and the median for Li is also similar to that of Teng et al. (2004) (35 ppm), as well as a newly proposed average UCC value based on loess analyses (Sauzeat et al., 2015) (30.5 ppm). The diamictite medians for Cr, Co, and Ni, while lower than UCC_{RG} , are higher than the Shaw et al. (1967; 1976) and Eade and Fahrig (1973) models based on bedrock sampling, but are comparable to the Kamber et al (2005) model for Cr and Ni. The diamictite median for Bi is comparable to the previous estimate of Gao et al. (1998).

4.4.2.1 Concentrations of chalcophile and siderophile elements and Be in the UCC

Several infrequently analyzed chalcophile and siderophile elements (Ga, Ge, Cd, In, Sn, Sb, W, Tl, and Bi) were analyzed (in most cases by standard addition) in this study in order to better characterize their abundance in the UCC. We include the lithophile element Be in this discussion because it is also infrequently analyzed and its abundance in the UCC is likewise less

constrained. Molybdenum was also analyzed but shows strong effects of mobilization during oxidative weathering in younger diamictites (Gaschnig et al., 2014), and consequently, it is only discussed here in the context of its relationship with Cd.

Of these elements, only Bi shows no statistical difference in concentration amongst the four glacial epochs. Be and Sn show the only evidence for secular changes, and both have higher concentrations in the Neoproterozoic and Paleozoic than Mesoarchean and Paleoproterozoic (red and green symbols vs. blue and black symbols, respectively, in Fig 15 a, b). This observation is consistent with greater differentiation of the crust through time, as both elements are incompatible and concentrated in melt during partial melting and/or fractional crystallization (e.g., Jochum et al., 1993; Ryan, 2002). Both elements correlate well with REEs, particularly Sm in the case of Be, and Yb in the case of Sn (Fig. 15a, b). Gallium correlates strongly with the other Group 13 elements, Al and In, and also with Ti, Sn, and Be (Fig. 15c). Germanium does not correlate with its Group 14 neighbors, Si and Sn, nor other trace elements, but it correlates with Fe (Fig. 15d) in older diamictite samples. Mesoarchean and Paleoproterozoic samples show correlations between Ge and Fe with differing slopes. This may indicate that Ge (which is siderophile) is following reduced iron prior to the GOE, but that this relationship does not hold for oxidized iron. Cadmium shows a complex but positive correlation with Mo (Fig. 15e) with separate diamictite units showing different Cd vs. Mo slopes. The presence or absence of oxidative weathering is the major process controlling Mo distribution in the diamictites (Gaschnig et al., 2014) and Cd may be affected by the same processes, as it is found in sulfides that are readily broken down by oxidative weathering and will remain in solution in sulfide-poor waters (e.g., Calvert and Pedersen, 1993). Indium correlates well with the MREE to HREEs and Y (Fig. 15f) and moderately well with Nb and Ta. Antimony does not

correlate with other trace elements but shows a broad positive correlation with the CIA (Fig. 15g), suggesting that it may be adhering to clays during chemical weathering, although the lack of correlation with other elements with a similar tendency (e.g., Rb) suggests several processes may be at work. Tungsten correlates weakly with Th and Tl (Fig. 15g). Tl correlates well with Rb, and to a lesser extent with Th, Cs, and K (Fig. 15h). Bi does not correlate well with any other element.

Using these correlations we calculate new average UCC concentrations for chalcophile and siderophile elements. This follows the approach of McLennan (2001), where the existing average UCC concentration for a well-constrained element (the “pinning” element), along with associated uncertainties, is input into the regression equation ($y = mx + b$) to calculate a new average value for the chalcophile or siderophile element of interest. For this purpose we have used both UCC_{RG} and the median values derived from the diamictites for the concentrations of the “pinning” elements. Propagating the errors on the slope and intercept of the regression, we obtain the results shown in Table 5. An obvious weakness of this general approach is that the newly-calculated concentrations are only as accurate as those for the pinning elements, and the approach can only be used in cases where correlations exist. In general, the concentrations derived from the regressions using different values for the pinning elements agree within uncertainty, and also agree with those of UCC_{RG} , except for Tl, which is significantly higher in the latter.

4.4.3 Towards a UCC consensus

An emerging consensus on the average abundance for many major and trace elements in the UCC can be seen in Figs. 13 and 14. All estimates of the average SiO_2 , Al_2O_3 , TiO_2 , and K_2O in the UCC are within $\pm 15\%$. The same is true of Ga, Ge, In, La, Ce, Nd, Pb, and Th.

While much of the dispersion seen amongst UCC models for other elements is attributable to chemical weathering effects, major differences are present amongst the insoluble first row transition metals, as well as the MREE and HREE. These differences may be the result of changing proportions of Archean to post-Archean UCC types in the sampled bedrock areas, in the case of the grid-based models, and sediment sources in the case of the sediment-based models.

5. Conclusions

Major and trace element and oxygen isotope compositions of ancient glacial diamictites from the Mesoarchean, Paleoproterozoic, Neoproterozoic, and Paleozoic paint a portrait of secular changes in the average composition of the upper continental crust with time. The diamictites indicate that the Archean UCC was dominated by greenstone belt basalt, and komatiite, with subordinate TTG and that the average UCC became more felsic and differentiated with time. These changes are consistent with previous studies that have utilized the geochemistry of shales, map-based reconstructions, and other approaches (e.g., Condie, 1993; Taylor and McLennan, 1985, 1995; Keller and Schoene, 2012), validating the utility of glacial diamictites as UCC proxies.

We use the diamictite data to develop a new model for the modern UCC composition based on the glacially sampled crust from the Neoproterozoic and Paleozoic eras. The major elements trends show depletions in Ca, Na, and Mg and enrichment in K compared other UCC models, which we attribute to the effects of chemical weathering, an intrinsic characteristic of the UCC. How deep this signature extends into the crust is an important, unresolved question. The diamictite-derived trace element estimates of UCC all fall within the range of existing models, with the exceptions of large depletions in Sr and Mo that result from chemical weathering.

Of the siderophile and chalcophile elements (and Be) analyzed, only Be and Sn concentrations show evidence for secular changes in the UCC, becoming more enriched with time. Gallium behaves most similarly to Al, Ge follows Fe (for pre-Great Oxidation Event samples), Cd follows Mo, In and Sn follow the MREEs, W follows Th, and Tl follows Rb. Antimony correlates only with the CIA, and Bi does not correlate with any other element or compositional parameter. New estimates for the UCC abundances of a subset of these elements (Be, Ga, In, Sn, Tl) derived from correlations with better-known elements and from median values for Paleozoic and Neoproterozoic diamictites are within uncertainty of UCC_{RG}, except for Tl, which is significantly higher in the latter.

Acknowledgements

This research was supported by NSF EAR-1321954, NSF EAR-1144454, NSF FESD EAR-1338810, and a grant from the State Key Laboratory of Geological Processes and Mineral Resources at China University of Geosciences in Wuhan. We thank Nic Beukes, Paul Link, Lian Zhou, Charlie Hoffmann, and Guy Narbonne for help with sampling and geologic interpretation; Heidi Anderson for providing the Bolivian samples; Richard Ash, Igor Puchtel, Zhang Wen, and Lin Lin for help with analytical work; Mike Spicuzza for oxygen isotope analysis; Mike Ream and Will Junkin for help with sample processing; and Lew Ashwal and Sue Webb for providing accommodations and libations in Johannesburg during our South African field work. We thank Mark Fornace for his help in initially developing the standard addition method that we now use in the Plasma Lab. We are especially grateful to Richard Ash for his endless support, insights, and positive spirit in keeping the Plasma Lab operational. This paper was improved by detailed and thoughtful reviews by Nick Arndt and Lang Farmer.

772

773 **Figure 1.** Diamictite major element compositions (and sediments in general) are chiefly
774 controlled by the ratio of quartz to feldspars and clays/phyllsilicates, which forms the primary
775 trend. Departures from this trend are caused by unusually high levels of either CaCO_3 or iron, as
776 seen in (b) and (c), which cause relative dilutions in the other major elements. The “carbonate
777 cut-off” line in (b) separates the carbonate-rich samples on the left from those used to calculate
778 the filtered medians in Figs. 13-14.

779 **Figure 2.** Rare earth element (REE) variations in diamictite composites ordered by depositional
780 age. Measured concentrations are double-normalized, with each analysis corrected for quartz
781 dilution by normalizing it to the average UCC_{RG} Al_2O_3 content and then normalizing individual
782 elements to chondrite (McDonough and Sun, 1995) in panel a, and average UCC_{RG} in panel b.

783 **Figure 3.** Transition metals (a) and post-transition metals and metalloids (b) in diamictite
784 composites, double normalized to UCC_{RG} , as in Fig. 2.

785 **Figure 4.** Alkali and alkaline earth elements (a), high field strength elements (HFSE; b), and
786 actinides (c) in diamictite composites, double normalized to UCC_{RG} , as in Fig. 2.

787 **Figure 5.** (a) Whole-rock oxygen isotope compositions of individual diamictite samples versus
788 depositional age. Diamictites with LOI values greater than 7% were excluded in order to
789 eliminate the skewing effects of high carbonate content on $\delta^{18}\text{O}$ (see supplemental material for
790 discussion). Large colored boxes are median values for each of the four time periods. The
791 median absolute error is shown as a vertical line for the Paleozoic but is smaller than the symbols
792 for the other three periods. Mantle composition is from Valley et al. (1998). Shaded field is
793 range of clastic sediment/sedimentary rock analyses compiled by Payne et al. (2015), filtered

here to exclude samples metamorphosed at medium and high-grade conditions, with additional data from Bindeman et al. (2016). Mesoarchean diamictites extend to distinctly lower $\delta^{18}\text{O}$ values compared to post-Archean diamictites. Although the other three glacial periods contain outliers, the average values become progressively higher with time. (b) $\delta^{18}\text{O}$ versus CIA (chemical index of alteration, a measure of chemical weathering intensity). If $\delta^{18}\text{O}$ values are being controlled by chemical weathering, a positive correlation should be present in the data. While this is true of the Neoproterozoic and Paleozoic diamictites, the $\delta^{18}\text{O}$ and CIA values are largely decoupled for the Mesoarchean and Paleoproterozoic diamictites.

Figure 6. Upper: Chemical index of alteration (CIA) versus index of chemical variability (ICV). Lower: an expanded view of the dashed box in the upper panel. The ICV increases as the composition becomes more mafic. See text for formal definitions. Average igneous compositions shown as purple triangles are from Condie (1993) and tie lines to CIA=100 and ICV=0 are weathering trends. UCC_{RG} is also shown as a purple triangle and plots with a primary igneous CIA because it is largely derived from grid sampling studies of igneous basement. Samples with >7% LOI have been excluded, as their high carbonate content results in extreme ICV values. Most diamictites fall along a weathering trend leading back to an andesitic composition but many of the Mesoarchean and some of the Paleoproterozoic samples fall near tie lines more consistent with a komatiitic source.

Figure 7. Ternary plots separating the effects of weathering from source rock composition, after Ohta and Arai (2007). This approach is based on principal component analysis of the major elements and divides compositions amongst “mafic” (M), “felsic” (F), and “weathered” (W) end-members. Unweathered igneous rocks plot along the arc connecting the M and F apices whereas weathering will draw compositions toward the W apex. Most diamictites plot in positions

consistent with a weathered intermediate source composition but Mesoarchean samples clearly derive from a more mafic source.

Figure 8. Trace element ratios sensitive to differentiation and/or mafic versus felsic sources plotted against depositional age for the composites, which represent average compositions for each diamictite unit. The decreasing (a) Ni/Lu, increasing (b) Th/Sc, and increasingly negative Eu anomaly (c) are all consistent with a progressively less mafic, more differentiated UCC with time. Large boxes and vertical bars represent median values and median absolute deviations for all individual samples from each time period, with the exception of the western Dwyka samples, which are dominated by >2.0 Ga sources (see text for discussion). In this and subsequent figures, the UCC_{RG} line represents the average UCC composition of Rudnick and Gao (2003) and shaded area represents the range of other published UCC models (Shaw et al, 1967, 1976; Eade and Fahrig, 1973; Taylor and McLennan, 1985; Condie, 1993; Wedepohl, 1995; Gao et al., 1998; Kamber et al., 2005; Hu and Gao, 2008; Chauvel et al., 2014).

Figure 9. Changes in Th and U with time for diamictite composites. Th and U are divided by Al (in ppm) in (a) and (b) to correct for quartz dilution effects. (a) Th shows a clear increase between the Mesoarchean and Paleoproterozoic. (b) A clear trend with age is less clear for U, but this is at least partially due to the effects of U mobility during oxidative weathering. (c) Th/U shows a steady increase with time and is highly variable in the Neoproterozoic. Both the increase and variability are likely driven by oxidative weathering, which became increasingly important after the Paleoproterozoic glaciations. UCC_{RG} and range of other UCC estimates are shown as a solid line and shaded area, respectively, as in Figure 8.

Figure 10. Changes in REE fractionation (a, c, e) and normalized abundances (b, d, f) with time for diamictite composites. (a, c, e) La/Lu represents the fractionation between the LREE and

HREEs and shows a large range for the different periods but appears to decrease slightly with time. La/Sm represents the fractionation within the LREEs and shows a significant decrease with time. Gd/Lu represents the fractionation within the HREEs and does not vary time. The changes seen in La/Lu and La/Sm may reflect a decrease in relative abundance of TTGs as a percent of the UCC and the increase in abundance of potassic “true” granites. (b, d, f) La, Gd, and Lu are given as representatives of the LREEs, MREEs, and HREEs, respectively. All elements are divided by Al (in ppm) to correct of quartz dilution effects. Increases in abundance are seen in all four elements with time and are especially marked for the MREEs and HREEs. This likely reflects both the increasingly felsic and differentiated nature of the UCC with time and the decrease in the relative proportion of TTGs to potassic granites, as the latter are more enriched in MREEs and HREEs. UCC_{RG} and range of other UCC estimates are shown as a solid line and shaded area, respectively, as in Figure 8.

Figure 11. (a) Sm-Nd and (b) Lu-Hf ratios in diamictite composites versus age. Significant overlap exists for all periods. Sm-Nd ratios are consistently higher than the Rudnick and Gao (2003) UCC estimate but do not exceed the full range of values in the literature. UCC_{RG} and range of other UCC estimates are shown as a solid line and shaded area, respectively, as in Figure 8.

Figure 12. HFSE ratios in diamictite composites versus age. (a) Nb/Ta shows a prominent increase between the Paleoproterozoic and Neoproterozoic, and (c) La/Nb shows an analogous decrease. This may reflect a decrease in the relative abundance of TTGs with time in the UCC. UCC_{RG} and range of other UCC estimates are shown as a solid line and shaded area, respectively, as in Figure 8.

Figure 13. (a) Comparison of major elements in versions of the diamictite-based UCC model, normalized to the Rudnick and Gao (2003) UCC. The squares are derived from the median of all individual Neoproterozoic and Paleozoic samples and the diamonds are based on the median of individual samples to the right of the carbonate cut-off boundary in Fig. 1b. Error bars are median absolute deviations. (b) Major elements from other UCC models, normalized to Rudnick and Gao (2003). References: Shaw et al. (1967, 1976), Eade and Fahrig (1973), Condie (1993), Taylor and McLennan (1985), Ronov and Yaroshevskiy (1967, 1976), and Borodin (1998).

Figure 14. Comparison of trace elements in the (a) two versions of the diamictite-based UCC model, as in Fig. 13. (b) Other UCC models are also shown. References: Shaw et al. (1967, 1976), Eade and Fahrig (1973), Condie (1993), Gao et al. (1998), Kamber et al. (2005), Taylor and McLennan (1985) including additional data from McLennan (2001), Wedepohl (1995), Hu and Gao (2008), and Chauvel et al. (2014) incorporating the Li concentration from Sauzeat et al. (2015).

Figure 15. Correlations seen amongst the chalcophile and siderophile trace elements and Be in the diamictites.

References

Arndt, N. (1999) Why was flood volcanism on submerged continental platforms so common in the Precambrian? *Precambrian Research* **97**, 155-164.

883 Arndt, N. (2013) The formation and evolution of the continental crust: *Geochemical*
884 *Perspectives*, **2**, 405-533.

885

886 Barth, M.G., McDonough, W.F., Rudnick, R.L. (2000) Tracking the budget of Nb and Ta in the
887 continental crust. *Chemical Geology* **165**, 197-213.

888

889 Bindeman, I.N., Bekker, A., Zakharov, D.O. (2016) Oxygen isotope perspective on crustal
890 evolution on early Earth: A record of Precambrian shales with emphasis on Paleoproterozoic
891 glaciations and Great Oxygenation Event. *Earth and Planetary Science Letters* **437**, 101-113.

892

893 Borodin, L.S. (1998) Estimated chemical composition and petrochemical evolution of the upper
894 continental crust. *Geochemistry International* **37**, 723-734.

895

896 Calvert S.E., Pedersen T.F. (1993) Geochemistry of Recent oxic and anoxic marine sediments:
897 Implications for the geological record. *Marine Geology* **113**, 67-88.

898

899 Canil, D., Lacourse, T. (2011) An estimate for the bulk composition of juvenile upper
900 continental crust derived from glacial till in the North American Cordillera. *Chemical Geology*
901 **284**, 229-239.

902

903 Carto, S.L., Eyles, N. (2011) The deep-marine glaciogenic Gaskiers Formation, Newfoundland,
904 Canada. In *The Geological Record of Neoproterozoic Glaciations* (eds. Arnaud, E., Halverson,
905 G.P., Shields-Zhou, G.), The Geological Society, London. pp. 467-473.

906

907 Chauvel, C., Garcon, M., Bureau, S., Besnault, A., Jahn, B.-M., Ding, Z. (2014) Constraints from
 908 loess on the Hf-Nd isotopic composition of the upper continental crust. *Earth and Planetary*
 909 *Science Letters* **388**, 48-58.
 910
 911 Clark, P.U. (1987) Subglacial sediment dispersal and till composition. *The Journal of Geology*
 912 **95**, 527-541.
 913
 914 Collerson, K.D., Kamber, B.S. (1999) Evolution of the continents and the atmosphere inferred
 915 from Th-U-Nb systematics of the depleted mantle. *Science* **283**, 1519-1522.
 916
 917 Committee on Grand Research Questions in the Solid-Earth Sciences (2008) Origin and
 918 Evolution of Earth: Research Questions for a Changing Planet, National Research Council, 152
 919 p. ISBN: 978-0-309-11717-3.
 920
 921 Condie, K.C. (1993) Chemical composition and evolution of the upper continental crust:
 922 Contrasting results from surface samples and shales. *Chemical Geology* **104**, 1-37.
 923
 924 Condie K.C. (2005) TTGs and adakites: are they both slab melts? *Lithos* **80**, 33-44.
 925
 926 Cox, R.N., Lowe, D.R., Cullers, R.L. (1995) The influence of sediment recycling and basement
 927 composition on evolution of mudrock chemistry in the southwestern United States. *Geochimica*
 928 *et Cosmochimica Acta* **59**, 2919-2940.
 929

930 Drummond, M.S., Defant, M.J. (1990) A model for trondhjemite-tonalite-dacite genesis and
 931 crustal growth via slab melting: Archean to modern comparisons. *Journal of Geophysical*
 932 *Research* **95**, 21503-21521.

933

934 Eade, K.E., Fahrig, W.F. (1973) Regional, lithological, and temporal variation in the abundances
 935 of some trace elements in the Canadian Shield. *Geological Survey of Canada Paper* **72**.

936 Eiler, J.M. (2001) Oxygen isotope variations of basaltic lavas and upper mantle rocks. *Rev. in*
 937 *Mineral. Geochem* **43**, 319-364.

938

939 Farmer, G.L., Licht, K., Swope, R.J. and Andrews, J. (2006) Isotopic constraints on the
 940 provenance of fine-grained sediment in LGM tills from the Ross Embayment, Antarctica. *Earth*
 941 *and Planetary Science Letters* **249**, 90-107.

942

943 Fedo, C.M., Wayne Nesbitt, H. and Young, G.M. (1995) Unraveling the effects of potassium
 944 metasomatism in sedimentary rocks and paleosols, with implications for paleoweathering
 945 conditions and provenance. *Geology* **23**, 921-924.

946

947 Frimmel, H.E. (2011) The Kaigas and Numees formations, Port Nolloth Group, in South Africa
 948 and Namibia. In *The Geological Record of Neoproterozoic Glaciations* (eds. Arnaud, E.,
 949 Halverson, G.P., Shields-Zhou, G.). The Geological Society, London, pp. 223-231.

950

951 Gaillard, F., Scaillet, B., Arndt, N.T. (2011) Atmospheric oxygenation caused by a change in
 952 volcanic degassing pressure. *Nature* **478**, 229-232.

953

954 Gao, S., Luo, T.-C., Zhang, B.-R., Zhang, H.-F., Han, Y.-w., Zhao, Z.-D., Hu, Y.-K. (1998)

955 Chemical composition of the continental crust as revealed by studies in East China. *Geochimica*

956 *et Cosmochimica Acta* **62**, 1959-1975.

957

958 Gaschnig, R.M., Rudnick, R.L., McDonough, W.F. (2015) Determination of Ga, Ge, Mo, Ag,

959 Cd, In, Sn, Sb, W, Tl, and Bi in USGS whole-rock reference materials by standard addition ICP-

960 MS. *Geostandards and Geoanalytical Research* **39**, 371-379.

961

962 Gaschnig, R.M., Rudnick, R.L., McDonough, W.F., Kaufman, A.J., Vervoort, J.D., Fisher, C.

963 (2015) Insights into crustal growth from detrital zircons in ancient glacial deposits: *EOS*,

964 *Transactions, American Geophysical Union*.

965

966 Gaschnig, R.M., Rudnick, R.L., McDonough, W.F., Kaufman, A.J., Hu, Z., Gao, S. (2014) Onset

967 of oxidative weathering of continents recorded in the geochemistry of ancient glacial diamictites.

968 *Earth and Planetary Science Letters* **408**, 87-99.

969

970 Goldschmidt VM (1933) Grundlagen der quantitativen geochemie. *Fortschritte der Mineralogie*,

971 *Krystallographie und Petrographie* **17**:112

972

973 Goldschmidt, V. M. (1958) *Geochemistry*. Oxford University Press, Oxford.

974

975 Green, T.H. (1995) Significance of Nb/Ta as an indicator of geochemical processes in the crust-

976 mantle system. *Chemical Geology* **120**, 347-359.

977

978 Grimes CB, Ushikubo T, John BE, Valley JW (2011) Uniformly mantle-like $\delta^{18}\text{O}$ in zircons
979 from oceanic plagiogranites and gabbros, *Contr. Min. Pet.* **161**, 13-33.

980

981 Hambrey, M.J., Harland, W.B., eds., (1981) *Earth's Pre-Pleistocene Glacial Record*. Cambridge
982 University Press, Cambridge.

983

984 Hildes, D.H.D., Clarke, G.K.C., Flowers, G.E. and Marshall, S.J. (2004) Subglacial erosion and
985 englacial sediment transport modelled for North American ice sheets. *Quaternary Science*
986 *Reviews* **23**, 409-430.

987 Hoffman, P.F. (2011) Strange bedfellows; glacial diamictite and cap carbonate from the
988 Marinoan (635 Ma) glaciation in Namibia. *Sedimentology* **58**, 57-119.

989

990 Hoffmann, J.E., Münker, C., Næraa, T., Rosing, M.T., Herwartz, D., Garbe-Schönberg, D.,
991 Svahnberg, H. (2011) Mechanisms of Archean crust formation inferred from high-precision
992 HFSE systematics in TTGs. *Geochimica et Cosmochimica Acta* **75**, 4157-4178.

993

994 Hofmann, M., Linnemann, U., Hoffmann, K.-H., Germs, G., Gerdes, A., Marko, L., Eckelmann,
995 K., Gärtner, A. and Krause, R. (2015) The four Neoproterozoic glaciations of southern Namibia
996 and their detrital zircon record: The fingerprints of four crustal growth events during two
997 supercontinent cycles. *Precambrian Research* **259**, 176-188.

998

999 Holland, H.D. (2006) The oxygenation of the atmosphere and oceans. *Philosophical*
 1000 *Transactions of the Royal Society B: Biological Sciences* **361**, 903-915.
 1001
 1002 Houston, R.S., Karlstrom, K.E., Graff, P.J., Flurkey, A.J. (1992) New stratigraphic subdivisions
 1003 and redefinition of subdivisions of Late Archean and Early Proterozoic metasedimentary and
 1004 metavolcanic rocks of the Sierra Madre and Medicine Bow Mountains, southern Wyoming. *U.S.*
 1005 *Geological Survey Professional Paper* **1520**.
 1006
 1007 Jochum, K.P., Hofmann, A.W., Seufert, H.M. (1993) Tin in mantle-derived rocks: Constraints on
 1008 Earth evolution. *Geochimica et Cosmochimica Acta* **57**, 3585-3595.
 1009
 1010 Kamber, B.S., Greig, A., Collerson, K.D. (2005) A new estimate for the composition of
 1011 weathered young upper continental crust from alluvial sediments, Queensland, Australia.
 1012 *Geochimica et Cosmochimica Acta* **69**, 1041-1058.
 1013
 1014 Kaufman, A.J., Sial, A.N., Frimmel, H.E., Misi, A. (2009) Neoproterozoic to Cambrian
 1015 Palaeoclimatic Events in Southwestern Gondwana. In *Neoproterozoic-Cambrian tectonics,*
 1016 *global change and evolution: a focus on southwestern Gondwana* (eds. Gaucher, C., Sial, A.N.,
 1017 Halverson, G.P., Frimmel, H.E.). Elsevier, Amsterdam. pp. 369-388.
 1018
 1019 Keeley, J.A., Link, P.K., Fanning, C.M., Schmitz, M.D. (2013) Pre- to synglacial rift-related
 1020 volcanism in the Neoproterozoic (Cryogenian) Pocatello Formation, SE Idaho: New SHRIMP
 1021 and CA-ID-TIMS constraints. *Lithosphere* **5**, 128-150.

1022

1023 Keller, C.B., Schoene, B. (2012) Statistical geochemistry reveals disruption in secular
 1024 lithospheric evolution about 2.5 Gyr ago. *Nature* **485**, 490-493.

1025

1026 Klassen, R.A. (1999) The application of glacial dispersal models to the interpretation of till
 1027 geochemistry in Labrador, Canada. *Journal of Geochemical Exploration* **67**, 245-269.

1028

1029 Knauth, L.P., Lowe, D.R. (2003) High Archean climatic temperature inferred from oxygen
 1030 isotope geochemistry of cherts in the 3.5 Ga Swaziland Supergroup, South Africa. *Geological*
 1031 *Society of America Bulletin* **115**, 566-580.

1032

1033 Le Heron, D.P., Busfield, M.E., Kamona, F. (2013) An interglacial on snowball Earth? Dynamic
 1034 ice behaviour revealed in the Chuos Formation, Namibia. *Sedimentology* **60**, 411-427.

1035

1036 Li, S., Gaschnig, R.M., Rudnick, R.L. (2016) Insights into chemical weathering of the upper
 1037 continental crust from the geochemistry of ancient glacial diamictites: *Geochimica et*
 1038 *Cosmochimica Acta* **176**, 96-117.

1039

1040 Liu, C.-Q., Masuda, A., Okada, A., Yabuki, S., Zhang, J., Fan, Z.-L. (1993) A geochemical study
 1041 of loess and desert sand in northern China: Implications for continental crust weathering and
 1042 composition. *Chemical Geology* **106**, 359-374.

1043

1044 Liu, X., Gao, S., Diwu, C., Ling, W. (2008) Precambrian crustal growth of Yangtze Craton as
 1045 revealed by detrital zircon studies. *American Journal of Science* **308**, 421-468.
 1046
 1047 Marschall, H.R., Dohmen, R. and Ludwig, T. (2013) Diffusion-induced fractionation of niobium
 1048 and tantalum during continental crust formation. *Earth and Planetary Science Letters* **375**, 361-
 1049 371.
 1050
 1051 Matthey, D., Lowry, D., Macpherson, C. (1994) Oxygen isotope composition of mantle peridotite.
 1052 *Earth and Planetary Science Letters* **128**, 231-241.
 1053
 1054 McDonough, W.F. (1990) Constraints on the composition of the continental lithospheric mantle.
 1055 *Earth and Planetary Science Letters* **101**, 1-18.
 1056
 1057 McLennan, S.M. (1993) Weathering and global denudation. *The Journal of Geology* 101, 295-
 1058 303.
 1059
 1060 McLennan, S.M. (2001) Relationships between the trace element composition of sedimentary
 1061 rocks and upper continental crust. *Geochem. Geophys. Geosyst.* **2**, 1021.
 1062
 1063 McLennan, S.M., Fryer, B.J., Young, G.M. (1979) Rare earth elements in Huronian (Lower
 1064 Proterozoic) sedimentary rocks: Composition and evolution of the post-Kenoran upper crust.
 1065 *Geochimica et Cosmochimica Acta* **43**, 375-388.
 1066

1067 McLennan, S.M., Nance, W.B., Taylor, S.R. (1980) Rare earth element-thorium correlations in
 1068 sedimentary rocks, and the composition of the continental crust. *Geochimica et Cosmochimica*
 1069 *Acta* **44**, 1833-1839.
 1070
 1071 McLennan, S.M., Taylor, S.R. (1980) Th and U in sedimentary rocks: crustal evolution and
 1072 sedimentary recycling. *Nature* **285**, 621-624.
 1073
 1074 McLennan, S.M., Taylor, S.R., Kroner, A. (1983) Geochemical evolution of Archean shales
 1075 from South Africa. I. The Swaziland and Pongola Supergroups. *Precambrian Research* **22**, 93-
 1076 124.
 1077
 1078 Melezhik, V.A., Young, G.M., Eriksson, P.G., Altermann, W., Kump, L.R., Lepland, A., (2013)
 1079 Huronian-age glaciation. In *Global Events and the Fennoscandian Arctic Russia - Drilling Earth*
 1080 *Earth Project* (eds. Melezhik, V.A., Prave, A.R., Hanski, E.J., Fallick, A.E., Lepland, A., Kump,
 1081 L.R., Strauss, H.). Springer-Verlag, Berlin. pp. 1059-1109.
 1082
 1083 Moyen, J., Martin, H. (2012) Forty years of TTG research. *Lithos* **148**, 312-336.
 1084
 1085 Muehlenbachs, K. (1998) The oxygen isotopic composition of the oceans, sediments and the
 1086 seafloor. *Chemical Geology* **145**, 263-273.
 1087
 1088 Nance, W.B., Taylor, S.R. (1976) Rare earth element patterns and crustal evolution - I.
 1089 Australian post-Archean sedimentary rocks. *Geochimica et Cosmochimica Acta* **40**, 1539-1551.

1090

1091 Nance, W.B., Taylor, S.R. (1977) Rare earth element patterns and crustal evolution - II. Archean
 1092 sedimentary rocks from Kalgoorlie, Australia. *Geochimica et Cosmochimica Acta* **41**, 225-231.
 1093

1094 Nesbitt, H.W., Young, G.M. (1982) Early Proterozoic climates and plate motions inferred from
 1095 major element chemistry of lutites. *Nature* **299**, 715-717.
 1096

1097 Ohta, T., Arai, H. (2007) Statistical empirical index of chemical weathering in igneous rocks: A
 1098 new tool for evaluating the degree of weathering. *Chemical Geology* **240**, 280-297.
 1099

1100 Payne, J.L., Hand, M., Pearson, N.J., Barovich, K.M., McInerney, D.J. (2015) Crustal thickening
 1101 and clay: Controls on O isotope variation in global magmatism and siliciclastic sedimentary
 1102 rocks. *Earth and Planetary Science Letters* **412**, 70-76.
 1103

1104 Pearce, J.A. and Peate, D.W. (1995) Tectonic implications of the composition of volcanic arc
 1105 magmas. *Annual Review of Earth and Planetary Sciences* **23**, 251-285.
 1106

1107 Peucker-Ehrenbrink, B., Jahn, B.-M. (2001) Rhenium-osmium isotope systematics and platinum
 1108 group element concentrations: Loess and the upper continental crust. *Geochem. Geophys.*
 1109 *Geosyst.* **2**. 2001GC000172.
 1110

1111 Potter, P.E., Maynard, J.B., Depetris, P.J. (2005) *Mud and Mudstones: Introduction and*
 1112 *Overview*. Springer-Verlag, Hiedelberg, 297 p.

1113

1114 Pfander, J.A., Munker, C., Stracke, A., Mezger, K. (2007) Nb/Ta and Zr/Hf in ocean island
 1115 basalts -- Implications for crust-mantle differentiation and the fate of Niobium. *Earth and*
 1116 *Planetary Science Letters* **254**, 158-172.

1117

1118 Prave, A.R., Hoffmann, K.H., Hegenberger, W., Fallick, A.E. (2011) The Witvlei Group of east-
 1119 central Namibia. In *The Geological Record of Neoproterozoic Glaciations* (eds. Arnaud, E.,
 1120 Halverson, G.P., Shields-Zhou, G.). The Geological Society, London. pp. 211-216.

1121

1122 Rankin, D.W. (1993) The volcanogenic Mount Rogers Formation and the overlying glaciogenic
 1123 Konnarock Formation; two late Proterozoic units in southwestern Virginia. *U.S. Geological*
 1124 *Survey Bulletin* **2029**.

1125

1126 Rey, P.F. and Coltice, N. (2008) Neoarchean lithospheric strengthening and the coupling of
 1127 Earth's geochemical reservoirs. *Geology* **36**, 635-638.

1128

1129 Ronov, A.B., Yaroshevsky, A.A. (1967) Chemical structure of the Earth's crust. *Geokhimiya* **11**,
 1130 1285-1309.

1131

1132 Ronov, A.B., Yaroshevsky, A.A. (1976) A new model for the chemical structure of the Earth's
 1133 crust. *Geokhimiya* **12**, 1761-1795.

1134

1135 Rudnick, R.L., Barth, M., Horn, I., McDonough, W.F. (2000) Rutile-bearing refractory eclogites:
 1136 missing link between continents and depleted mantle. *Science* **287**, 278-281.
 1137
 1138
 1139 Rudnick, R.L., Gao, S. (2003) The composition of the continental crust, in: Rudnick, R.L. (Ed.),
 1140 The Crust. Elsevier-Pergamon, Oxford, pp. 1-64.
 1141
 1142 Ryan, J.G., 2002. Trace-element systematics of beryllium in terrestrial materials. *Rev. in*
 1143 *Mineral. Geochem* **50**, 121-146.
 1144
 1145 Sauzéat, L., Rudnick, R.L., Chauvel, C., Garçon, M., Tang, M. (2015) New perspectives on the
 1146 Li isotopic composition of the upper continental crust and its weathering signature. *Earth and*
 1147 *Planetary Science Letters* **428**, 181-192.
 1148
 1149 Shaw, D.M., Dostal, J., Keays, R.R. (1976) Additional estimates of continental surface
 1150 Precambrian shield composition in Canada. *Geochimica et Cosmochimica Acta* **40**, 73-83.
 1151
 1152 Shaw, D.M., Reilly, G.A., Muysson, J.R., Pattenden, G.E., Campbell, F.E. (1967) An estimate of
 1153 the chemical composition of the Canadian Precambrian Shield. *Canadian Journal of Earth*
 1154 *Sciences* **4**, 829-853.
 1155

1156 Spicuzza, M.J., Valley, J.W., McConnell, V.S. (1998) Oxygen isotope analysis of whole rock via
 1157 laser fluorination: an air-lock approach. *Abstracts with Programs - Geological Society of*
 1158 *America* **30**, 80.
 1159
 1160 Starck, D., Papa, C.d. (2006) The northwestern Argentina Tarija Basin: Stratigraphy,
 1161 depositional systems, and controlling factors in a glaciated basin. *Journal of South American*
 1162 *Earth Sciences* **22**, 169-184.
 1163
 1164 Tang, M., Chen, K., Rudnick, R.L. (2016) Archean upper crust transition from mafic to felsic
 1165 marks the onset of plate tectonics. *Science* 351, 372-375.
 1166
 1167 Tang, M., Rudnick, R. L., McDonough, W. F., Gaschnig, R. M., and Huang, Y. (2015) Europium
 1168 anomalies constrain the mass of recycled lower continental crust: *Geology* **43**, 703-706.
 1169
 1170 Taylor, S.R., McLennan, S.M., McCulloch, M.T. (1983) Geochemistry of loess, continental
 1171 crustal composition and crustal model ages. *Geochimica et Cosmochimica Acta* **47**, 1897-1905.
 1172
 1173 Taylor, S.R., McLennan, S.M. (1985) *The Continental Crust: It's Composition and Evolution*.
 1174 Blackwell Scientific, Oxford, 312 p.
 1175
 1176 Taylor, S.R., McLennan, S.M. (1995) The geochemical evolution of the continental crust.
 1177 *Reviews of Geophysics* **33**, 241-265.
 1178

1179 Togashi, S., Imai, N., Okuyama-Kusunose, Y., Tanaka, T., Okai, T., Koma, T., Murata, Y.
 1180 (2000) Young upper crustal chemical composition of the orogenic Japan Arc. *Geochemistry,*
 1181 *Geophysics, and Geosystems* **1**, 1049.
 1182
 1183 Valley, J.W., Kitchen, N., Kohn, M.J., Niendorf, C.R., Spicuzza, M.J. (1995) UWG-2, a garnet
 1184 standard for oxygen isotope ratios: Strategies for high precision and accuracy with laser heating.
 1185 *Geochimica et Cosmochimica Acta* **59**, 5223-5231.
 1186
 1187 Valley, J. W., Kinny, P. D., Schulze, D. J., and Spicuzza, M. J. (1998) Zircon Megacrysts from
 1188 Kimberlite: Oxygen isotope heterogeneity among mantle melts. *Contributions to Mineralogy*
 1189 *and Petrology* **133**, 1-11.
 1190
 1191 Valley, J.W., Lackey, J.S., Cavoisie, A.J., Clechenko, C.C., Spicuzza, M.J., Basei, M.A.S.,
 1192 Bindeman, I.N., Ferreira, V.P., Sial, A.N., King, E.M., Peck, W.H., Sinha, A.K., Wei, C.S.
 1193 (2005) 4.4 billion years of crustal maturation: oxygen isotope ratios of magmatic zircon.
 1194 *Contributions to Mineralogy and Petrology* **150**, 561-580.
 1195
 1196 Visser, J.N.J. (1982) Upper Carboniferous glacial sedimentation in the Karoo Basin near Prieska,
 1197 South Africa. *Palaeogeography, Palaeoclimatology, Palaeoecology* **38**, 63-92.
 1198
 1199 Vorster, C. (2013) Laser ablation ICP-MS age determination of detrital zircon populations in the
 1200 Phanerozoic Cape and lower Karoo Supergroup (South Africa) and correlatives in Argentina.
 1201 Ph.D. thesis, University of Johannesburg.

1202

1203 Wedepohl, K.H. (1995) The composition of the continental crust. *Geochimica et Cosmochimica*

1204 *Acta* **59**, 1217-1232.

1205

1206 Whalen, J.B., Percival, J.A., McNicoll, V.J., Longstaffe, F.J. (2004) Geochemical and isotopic

1207 (Nd–O) evidence bearing on the origin of late- to post-orogenic high-K granitoid rocks in the

1208 Western Superior Province: implications for late Archean tectonomagmatic processes.

1209 *Precambrian Research* **132**, 303-326.

1210

1211 Wronkiewicz, D.J., Condie, K.C. (1987) Geochemistry of Archean shales from the

1212 Witwatersrand Supergroup, South Africa: Source-area weathering and provenance. *Geochimica*

1213 *et Cosmochimica Acta* **51**, 2401-2416.

1214

1215 Wronkiewicz, D.J., Condie, K.C. (1989) Geochemistry and provenance of sediments from the

1216 Pongola Supergroup, South Africa: Evidence for a 3.0-Ga-old continental craton. *Geochimica et*

1217 *Cosmochimica Acta* **53**, 1537-1549.

1218

1219 Wronkiewicz, D.J., Condie, K.C. (1990) Geochemistry and mineralogy of sediments from the

1220 Ventersdorp and Transvaal Supergroups, South Africa: Cratonic evolution during the early

1221 Proterozoic. *Geochimica et Cosmochimica Acta* **54**, 343-354.

1222

1223 Yonkee, W.A., Dehler, C.D., Link, P.K., Balgord, E.A., Keeley, J.A., Hayes, D.S., Wells, M.L.,

1224 Fanning, C.M., Johnston, S.M. (2014) Tectono-stratigraphic framework of Neoproterozoic to

1225 Cambrian strata, west-central U.S.: Protracted rifting, glaciation, and evolution of the North
 1226 American Cordilleran margin. *Earth-Science Reviews* **136**, 59-95.

1227

1228 Young, G.M., Minter, W.E.L., Theron, J.N. (2004) Geochemistry and palaeogeography of upper
 1229 Ordovician glaciogenic sedimentary rocks in the Table Mountain Group, South Africa.
 1230 *Palaeogeography, Palaeoclimatology, Palaeoecology* **214**, 323-345.

1231

1232 Young, G.M., von Brunn, V., Gold, D.J.C., Minter, W.E.L. (1998) Earth's oldest reported
 1233 glaciation: physical and chemical evidence from the Archean Mozaan Group (~2.9 Ga) of South
 1234 Africa. *Journal of Geology* **106**, 523–538.

1235

1236 Zhang, W., Hu, Z., Liu, Y., Chen, H., Gao, S., Gaschnig, R.M. (2012a) Total Rock Dissolution
 1237 Using Ammonium Bifluoride (NH₄HF₂) in Screw-Top Teflon Vials: A New Development in
 1238 Open-Vessel Digestion. *Analytical Chemistry* **84**, 10686-10693.

1239

1240 Zhang, W., Hu, Z., Liu, Y., Chen, L., Chen, H., Li, M., Zhao, L., Hu, S., Gao, S. (2012b)
 1241 Reassessment of HF/HNO₃ decomposition capability in the high-pressure digestion of felsic
 1242 rocks for multi-element determination by ICP-MS. *Geostandards and Geoanalytical Research*
 1243 **36**, 271-289.

1244

1245 Zhou, C., Tucker, R., Xiao, S., Peng, Z., Yuan, X., Chen, Z. (2004) New constraints on the ages
 1246 of Neoproterozoic glaciations in south China. *Geology* **32**, 437-440.

Table 1. Glacial diamictites investigated in this study.

Stratigraphic unit	Country (state/province)	Drill core or Outcrop?	General reference	Approximate Age (Ga)
<i>Archean</i>				
Mozaan Grp	S Africa	O	Young et al. (1998)	2.9
Afrikander Frm*	S Africa	O	Guy et al. (2010)	2.9
Coronation Frm*	S Africa	DC	Guy et al. (2010)	2.9
Promise Frm*	S Africa	DC	Guy et al. (2010)	2.9
<i>Paleoproterozoic</i>				
Duitschland Frm	S Africa	DC	Melezhik et al. (2013)	2.4
Timeball Hill Frm**	S Africa	DC	Melezhik et al. (2013)	2.2
Makganyene Frm	S Africa	DC	Melezhik et al. (2013)	2.3
Ramsay Lake Frm	Canada (Ontario)	O	Melezhik et al. (2013)	2.4
Pecors Frm	Canada (Ontario)	O	Melezhik et al. (2013)	2.4
Bruce Frm	Canada (Ontario)	O	Melezhik et al. (2013)	2.4
Gowganda Frm	Canada (Ontario)	O	Melezhik et al. (2013)	2.4
Bottle Creek Frm	USA (Wyoming)	O	Houston et al (1992)	2.4
<i>Neoproterozoic</i>				
Pocatello Frm	USA (Idaho)	O	Keeley et al (2013)	0.7
Konnarock Frm	USA (Virginia)	O	Rankin (1993)	0.7
Gaskiers Frm	Canada (Newfoundland)	O	Carto and Eyles (2011)	0.58
Nantuo Frm	China (Hubei)	O	Zhou et al. (2004)	0.64
Gucheng Frm	China (Hubei)	O	Liu et al (2008)	0.7
Blaubeker Frm	Namibia	O	Prave et al. (2011)	0.7
Kaigas Frm	Namibia	O	Frimmel (2011)	0.75
Blasskranz	Namibia	O	Kaufman et al. (2010)	0.64
Chuos Frm	Namibia	O	Le Heron et al. (2013)	0.7
Numees Frm	Namibia	O	Frimmel (2011)	0.6
Ghaub Frm	Namibia	O	Hoffman (2011)	0.64
<i>Paleozoic</i>				
Machareti Group	Bolivia	O	Starck and Papa (2006)	0.3
Mandiyuti Group	Bolivia	O	Starck and Papa (2006)	0.3
Pakhuis Frm	S Africa	DC	Young et al. (2004)	0.44
Dwyka Group	S Africa and Namibia	O & DC	Visser (1982)	0.3

O = outcrop, DC = drill core

* - The Afrikander, Coronation, and Promise Formations are components of the Witwatersrand Supergroup and are grouped as such in figures. Note that the Afrikander diamictite is sometimes referred to as the Lagerspoort or the Blue Grit diamictite in the literature.

** - The Timeball Hill diamictite is sometimes referred to as the Rietfontein diamictite in the literature

Table 2

Sample	Stratigr:	SiO2	TiO2	Al2O3	Fe2O3	MnO	MgO	CaO	Na2O	K2O	P2O5	LOI**	CIA***	M%***	W%***	F%***
XRF*																
Mesoarchean																
South Africa																
13RMG005	Mozaan	58.84	0.39	8.95	23.41	0.11	2.89	0.38	0.78	1.67	0.07	2.51	71	19	75	7
13RMG006	Mozaan	58.13	0.37	8.75	24.44	0.13	3.02	0.52	0.63	1.42	0.05	2.54	72	24	70	6
13RMG007	Mozaan	58.86	0.38	8.94	23.59	0.16	3.24	0.13	0.08	0.96	0.06	3.62	88	2	98	0
13RMG008	Mozaan	59.31	0.41	9.07	23.01	0.15	2.90	0.14	0.72	1.78	0.06	2.46	74	6	91	3
13RMG009	Mozaan	53.99	0.41	9.52	26.42	0.13	3.30	0.76	0.66	1.51	0.06	3.25	72	28	66	6
13RMG010	Mozaan	57.64	0.36	8.51	23.21	0.13	2.89	1.25	0.56	1.69	0.06	3.70	70	23	70	7
13RMG011	Mozaan	55.92	0.33	7.82	28.54	0.11	2.99	0.16	0.51	0.80	0.05	2.78	81	10	88	2
13RMG001	Afrikan	63.97	0.24	8.49	8.37	0.13	9.31	5.94	0.50	0.41	0.04	2.61	80	37	58	5
13RMG002	Afrikan	64.09	0.24	8.24	8.38	0.13	9.02	6.09	0.38	0.49	0.04	2.90	82	27	68	4
13RMG003	Afrikan	63.86	0.26	8.32	8.62	0.14	9.37	5.99	0.42	0.34	0.04	2.65	83	33	63	4
13RMG004	Afrikan	62.05	0.24	8.53	8.52	0.15	9.64	5.87	0.90	1.07	0.04	2.99	67	46	43	12
BAB1-4031	Promise	61.01	0.66	15.48	11.59	0.11	3.90	0.54	0.54	2.43	0.06	3.69	78	16	77	6
BAB1-4042	Promise	85.64	0.09	7.18	1.23	0.02	0.40	1.25	0.93	1.48	0.02	1.76	60	8	18	74
BABI 4053	Promise	63.37	0.44	11.13	15.61	0.11	3.71	0.66	0.44	0.59	0.04	3.91	84	23	74	3
BABI 3137	Coronat	74.34	0.39	15.07	2.31	0.03	1.26	0.21	0.35	3.29	0.07	2.68	78	3	84	12
BABI-3171	Coronat	72.33	0.47	15.08	3.34	0.04	1.84	0.46	0.33	3.00	0.07	3.04	78	6	84	11
BABI 3201	Coronat	68.75	0.54	15.74	5.57	0.06	2.73	0.47	0.38	2.72	0.10	2.94	80	7	85	7
BAB1-3216	Coronat	72.15	0.52	15.67	3.94	0.06	1.95	0.70	0.30	2.57	0.06	2.07	81	6	86	8
BABI 3277	Coronat	71.70	0.55	13.94	5.15	0.05	2.01	0.14	0.23	2.65	0.04	3.53	80	3	94	3
Paleoproterozoic																
Wyoming																
12RMG032	Bottle C	72.19	0.46	13.12	3.90	0.05	2.19	0.61	4.39	1.82	0.11	1.15	57	22	34	44
12RMG033	Bottle C	70.62	0.47	13.42	4.50	0.05	2.70	0.49	3.92	2.00	0.12	1.71	59	20	43	37
12RMG034	Bottle C	71.66	0.48	13.18	4.08	0.05	2.43	0.52	3.97	1.89	0.12	1.62	59	21	40	39
12RMG035	Bottle C	70.90	0.46	13.49	4.31	0.03	2.45	0.46	4.10	1.76	0.12	1.92	59	20	43	37
Ontario																
12RMG036	Ramsay	82.09	0.26	7.91	2.84	0.06	0.98	1.29	1.86	1.60	0.05	1.07	53	23	23	54
12RMG037	Ramsay	77.60	0.38	10.81	3.01	0.05	2.23	0.15	0.16	4.06	0.07	1.48	69	2	94	4
12RMG040	Gowgar	69.30	0.47	13.83	5.84	0.03	2.17	0.34	5.08	1.40	0.11	1.43	57	18	50	32
12RMG041	Gowgar	70.49	0.44	13.77	4.77	0.06	2.21	0.26	4.13	2.21	0.08	1.59	59	12	53	34
12RMG042	Bruce F	71.27	0.41	14.12	2.87	0.02	2.19	0.35	4.23	2.90	0.10	1.54	57	11	40	48
12RMG043	Bruce F	62.28	0.61	17.37	5.16	0.04	3.81	0.41	3.60	3.84	0.12	2.76	62	15	54	31
12RMG044	Bruce F	70.47	0.40	12.96	5.02	0.03	3.54	0.32	3.51	1.46	0.09	2.20	63	19	53	28
12RMG050	Ramsay	61.19	0.72	15.82	9.93	0.10	4.19	0.53	2.31	1.86	0.18	3.17	71	22	64	13
12RMG051	Ramsay	62.61	0.63	12.87	9.77	0.15	3.96	2.51	1.43	1.38	0.18	4.50	68	44	43	13
12RMG052	Ramsay	59.20	0.89	15.02	12.15	0.14	4.74	0.71	1.41	1.55	0.18	4.02	76	28	63	8
12RMG053	Gowgar	65.47	0.56	14.86	6.78	0.06	3.03	0.45	4.52	2.07	0.16	2.05	59	20	51	29
12RMG054	Gowgar	65.32	0.59	15.07	6.49	0.05	2.95	0.48	4.80	2.17	0.16	1.92	58	21	49	31
12RMG055	Gowgar	70.18	0.44	13.59	4.77	0.05	1.98	0.37	4.96	2.06	0.09	1.52	56	17	42	42
12RMG056	Gowgar	66.70	0.54	14.16	6.93	0.05	2.86	0.27	4.64	1.66	0.11	2.08	59	16	60	24
12RMG057	Gowgar	65.18	0.67	15.24	5.62	0.11	2.52	1.16	5.62	2.27	0.13	1.48	53	34	26	40
12RMG058	Gowgar	64.71	0.72	15.32	5.70	0.14	2.60	1.15	5.67	2.39	0.11	1.50	53	34	27	39
12RMG059	Gowgar	71.80	0.40	13.53	3.16	0.07	1.22	0.72	4.98	3.14	0.08	0.91	52	14	23	63
12RMG063	Gowgar	65.20	0.56	14.67	7.37	0.06	2.58	0.71	4.75	2.20	0.14	1.76	57	27	39	34
12RMG064	Gowgar	66.72	0.54	14.49	6.59	0.06	2.67	0.68	5.34	0.91	0.13	1.89	58	36	36	28
12RMG065	Gowgar	74.13	0.35	10.96	3.06	0.04	2.06	1.13	2.52	3.83	0.14	1.79	52	19	26	55
12RMG066	Gowgar	66.71	0.59	12.60	6.87	0.11	3.21	3.16	3.90	1.24	0.10	1.51	49	59	16	25
12RMG067	Gowgar	66.89	0.57	12.48	5.85	0.11	3.63	2.14	2.31	1.56	0.10	4.37	58	50	26	24
12RMG048	Pecors I	59.15	0.75	16.80	9.59	0.11	3.57	1.74	1.42	2.84	0.18	3.85	69	34	49	17
12RMG049	Pecors I	56.55	0.90	21.05	8.25	0.07	2.89	0.18	0.99	5.01	0.13	3.98	75	0	98	2
South Africa																
GTF01.25	Makgar	50.93	0.36	7.04	18.17	0.65	5.33	6.50	0.06	0.54	0.09	10.33	92	1	99	0

Table 3 - Precision and reproducibility based on analysis of AGV-2 as an unknown

	Average (n=7) *	1 std	%RSD	GEOREM	Average/GEOREM	Std add value** (n=6)	Std add %RSD
Li	11.5	0.3	3.0	11	1.05		
Be	2.28	0.09	4.0	2.3	0.99		
Sc	12.6	1.6	12.9	13	0.97		
V	122	7	6.0	120	1.02		
Cr	15.5	0.5	2.9	17	0.91		
Co	15.2	0.3	1.8	16	0.95		
Ni	19.1	0.4	2.0	19	1.00		
Cu	52.8	1.4	2.6	53	1.00		
Ga	20.4	0.3	1.5	20	1.02	20.2	2.0
Ge	1.22	0.043	3.5	1.43	0.86	1.15	3.4
Rb	66.2	0.5	0.8	66.3	1.00		
Sr	649	5	0.7	661	0.98		
Y	20.2	0.2	1.1	20	1.01		
Zr	230	17	7.2	240	0.96		
Nb	13.8	0.2	1.4	14.6	0.95		
Mo	2.21	0.25	11.2	2.08	1.06	1.93	2.4
Cd***				0.0820		0.137	3.6
In***				0.0480		0.0445	2.0
Sn	2.14	0.45	21.1	2.08	1.03	1.90	3.2
Sb	0.469	0.021	4.5	0.5	0.94	0.574	8.1
Cs	8.83	0.07	0.8	7.57	1.17		
Ba	1133	14	1.2	1140	0.99		
La	40.2	0.3	0.9	37.9	1.06		
Ce	73.1	0.5	0.7	68.6	1.07		
Pr	8.83	0.07	0.8	7.57	1.17		
Nd	32.2	0.2	0.7	30.2	1.06		
Sm	5.93	0.07	1.2	5.49	1.08		

Eu	1.62	0.02	1.1	1.53	1.06		
Gd	4.58	0.05	1.1	4.52	1.01		
Tb	0.694	0.008	1.1	0.64	1.08		
Dy	3.60	0.05	1.3	3.47	1.04		
Ho	0.710	0.009	1.3	0.653	1.09		
Er	1.93	0.02	1.2	1.81	1.06		
Tm	0.272	0.003	1.0	0.26	1.05		
Yb	1.68	0.04	2.3	1.62	1.04		
Lu	0.257	0.003	1.3	0.247	1.04		
Hf	5.65	0.16	2.7	5.04	1.12		
Ta	0.907	0.040	4.4	0.89	1.02		
W	0.651	0.107	16.4	0.55	1.18	0.489	4.2
Tl	0.317	0.003	1.1	0.32	0.99	0.272	5.6
Pb	15.9	0.4	2.3	14.2	1.12		
Bi	0.0431	0.0103	23.9	0.052	0.83	0.0509	9.0
Th	6.62	0.17	2.6	6.16	1.07		
U	2.07	0.05	2.5	1.9	1.09		

* - average values for analysis by conventional external calibration with USGS standards

** - average values for standard addition analysis

*** - Cd and In were determined by standard addition only

Table 4 - Proposed model for UCC based on Neoproterozoic and Paleozoic glacial diamictites

	Filtered to remove carbonate-rich samples		All samples		% difference (filtered vs unfiltered)	UCC _{RG}	%RSD *
	(in wt %)	Filtered Median (Preferred values)	Median absolute deviation	Unfiltered Median	Median absolute deviation		
SiO ₂		70.4	2.3	70.2	2.5	0%	2%
TiO ₂		0.7	0.08	0.7	0.08	0%	13%
Al ₂ O ₃		14.6	0.9	14.4	1.3	1%	5%
Fe ₂ O ₃		5.5	0.64	5.6	1.3	-1%	10%
T							
MnO		0.1	0.03	0.1	0.04	-3%	13%
MgO		2.0	0.35	2.2	0.42	-9%	14%
CaO		1.4	0.63	1.7	0.85	-17%	6%
Na ₂ O		1.6	0.70	1.5	0.70	7%	15%
K ₂ O		3.6	0.33	3.5	0.43	2%	8%
P ₂ O ₅		0.2	0.03	0.2	0.03	2%	15%
(in ppm)							
Li		31.3	5.2	31.4	5.8	0%	21%
Be		2.34	0.41	2.19	0.59	7%	41%
Sc		12.1	1.8	12.0	2.0	1%	6%
V		73.8	7.2	75.0	12.1	-2%	11%
Cr		53.0	10.8	55.7	14.5	-5%	19%
Co		11.8	2.0	12.3	3.0	-4%	3%
Ni		26.3	5.4	27.0	7.3	-3%	24%
Cu		20.2	7.7	20.5	7.8	-1%	14%
Zn		73.0	11.4	72.0	15.0	1%	9%
Ga		18.2	2.1	17.6	2.5	3%	4%
Ge		1.66	0.14	1.61	0.21	3%	9%
Rb		110	20	106	21	4%	20%
Sr		91.5	30.6	99.1	30.4	-8%	14%
Y		33.2	4.5	31.0	5.0	7%	11%
Zr		220	31	205	39	7%	14%
Nb		12.7	1.8	12.4	2.1	2%	12%
Mo		0.30	0.09	0.35	0.13	-15%	28%
Cd		0.11	0.04	0.13	0.04	-21%	15%
In		0.0620	0.0136	0.0593	0.0109	4%	14%
Sn		2.41	0.34	2.38	0.42	1%	26%
Sb		0.418	0.174	0.382	0.156	9%	28%
Cs		4.30	0.85	4.30	0.94	0%	31%
Ba		731	113	696	121	5%	13%
La		36.5	7.2	34.9	8.7	4%	9%
Ce		72.9	16.5	70.0	18.5	4%	6%
Pr		8.54	1.87	8.25	1.95	3%	
Nd		33.3	6.9	31.5	6.7	6%	8%
Sm		6.84	1.21	6.36	1.34	7%	6%

Eu	1.37	0.23	1.31	0.26	4%	1.02	14%
Gd	6.05	1.18	5.60	1.07	8%	4.01	7%
Tb	0.983	0.162	0.920	0.174	7%	0.650	1%
Dy	5.57	0.82	5.28	0.96	5%	3.91	
Ho	1.13	0.16	1.08	0.19	4%	0.83	
Er	3.27	0.46	3.08	0.55	6%	2.30	
Tm	0.506	0.069	0.487	0.080	4%	0.300	
Yb	3.07	0.33	3.02	0.45	1%	2.04	18%
Lu	0.479	0.053	0.464	0.067	3%	0.306	17%
Hf	6.50	1.00	5.98	1.18	8%	5.26	14%
Ta	0.940	0.131	0.901	0.152	4%	0.882	13%
W	1.10	0.16	1.11	0.15	-1%	1.90	54%
Tl	0.588	0.187	0.544	0.179	8%	0.941	57%
Pb	15.8	5.1	15.1	4.9	5%	17.0	3%
Bi	0.24	0.08	0.22	0.09	8%	0.16	38%
Th	11.3	3.1	10.5	3.1	7%	10.1	10%
U	2.66	0.87	2.40	0.99	10%	2.63	21%

* - %RSD is based on uncertainties reported in Rudnick and Gao (2003)

Table 5. Average upper crustal abundances of select infrequently analyzed elements derived from correlation of element abundances in UCC from

		Correlation		Pinning element		uncertainty	
Best Est.*		uncertainty	derived**	uncertainty	median)		
Be	2.4	0.4	1.8	0.1	Sm	2.2	0.6
Ga	17.6	1.2	19.3	1.1	Al	17.6	2.5
In	0.066	0.009	0.044	0.003	Tm	0.059	0.011
Sn	2.54	0.30	1.74	0.27	Yb	2.38	0.42
Tl	0.69	0.13	0.52	0.12	Rb	0.54	0.18

All concentrations in ppm

* - Neoproterozoic/Paleozoic diamictite median for pinning element is used to calculate element of interest

** - Rudnick and Gao (2003) estimate for pinning element is used to calculate element of interest

Figure 1

FIGURES

Figure &

Figure 3

Figure 4

Figure 5

Figure 6

Figure +

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

Figure 13

Figure 14

Figure 15