**Platinum-group element abundances and Re-Os
isotopic systematics of the upper continental crust
through time: evidence from glacial diamictites**

Kang Chen ^{a, b, *}, Richard J. Walker ^b, Roberta L. Rudnick ^{b, 1}, Shan Gao ^{a, 2},
Richard M. Gaschnig ^{b, 3}, Igor S. Puchtel ^b, Ming Tang ^b, Zhao-Chu Hu ^a

^a State Key Laboratory of Geological Processes and Mineral Resources,
School of Earth Sciences, China University of Geosciences, Wuhan 430074,
China

^b Department of Geology, University of Maryland, College Park, Maryland
20742, United States

¹ Present address: Department of Earth Science, University of California,
Santa Barbara, CA 93106-9630, United States

² Deceased.

³ Present address: School of Earth and Atmospheric Sciences,
Georgia Institute of Technology, Atlanta 30332-0340, United States

* Corresponding author. E-mail address: kangchen@cug.edu.cn

28 Abstract

29 The fine-grained matrix of glacial diamictites, deposited periodically by
30 continental ice sheets over much of Earth history, provides insights into the
31 average composition and chemical evolution of the upper continental crust
32 (UCC) (Gaschnig et al., 2016, and references therein). The concentrations of
33 platinum-group elements (PGEs, including Os, Ir, Ru, Pt and Pd) and the
34 geochemically related Re, as well as $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios, are
35 reported here for globally-distributed glacial diamictites that were deposited
36 during the Mesoarchean, Paleoproterozoic, Neoproterozoic and Paleozoic
37 eras. The medians and averages of PGE concentrations of these diamictite
38 composites decrease from the Mesoarchean to the Neoproterozoic, mimicking
39 decreases in the concentrations of first-row transition elements (Sc, V, Cr, Co
40 and Ni). By contrast, Re concentrations are highly variable with no discernable
41 trend, owing to its high solubility. Assuming these diamictites are
42 representative of average UCC through time, the new data are fully consistent
43 with the previous inference that the Archean UCC contained a greater
44 proportion of mafic-ultramafic rocks relative to younger UCC. Linear
45 regressions of PGEs versus Cr and Ni concentrations in all the diamictite
46 composites from the four time periods are used to estimate the following
47 concentrations of the PGEs in the present-day UCC: 0.059 ± 0.016 ng/g Os,
48 0.036 ± 0.008 ng/g Ir, 0.079 ± 0.026 ng/g Ru, 0.80 ± 0.22 ng/g Pt and $0.80 \pm$
49 0.26 ng/g Pd (2σ of 10,000 bootstrapping regression results). These PGE
50 estimates are slightly higher than the estimates obtained from loess samples.
51 We suggest this probably results from loess preferentially sampling younger
52 UCC rocks that have lower PGE concentrations, or PGEs being fractionated
53 during loess formation. A Re concentration of 0.25 ± 0.12 ng/g (2σ) is obtained
54 from a regression of Re versus Mo. From this, time-integrated $^{187}\text{Re}/^{188}\text{Os}$ and

$^{187}\text{Os}/^{188}\text{Os}$ ratios for the UCC are calculated, assuming an average UCC residence duration of ~ 2.0 Ga, yielding ratios of 20 ± 12 and 0.80 ± 0.38 (2σ), respectively.

Keywords: platinum-group elements; Re-Os isotopes; glacial diamictite; secular variation; upper continental crust

1. Introduction

Quantification of the average composition of the continental crust with associated uncertainties, and recognition of secular compositional changes are critical for understanding the origin, differentiation and evolution of the continents. Among the three commonly-defined layers of the continental crust (upper, middle and lower), the upper continental crust (UCC) is the most accessible region and its chemical composition has been the focus of numerous studies (Rudnick and Gao, 2003, and references therein; Kamber et al., 2005; Hu and Gao, 2008; Gaschnig et al., 2016). The average abundances of the major elements and some trace elements (most of the transition elements, Rb, Sr, Y, Zr, Nb, Ba, rare earth elements, Hf, Ta, Pb, Th and U) in the UCC are generally well known, with most estimates falling within a 20% deviation of the averages of all estimates. Other element abundances are less well constrained, especially the halogens, S, Ge, As, Se, In, Sn, Au, Re and platinum-group elements (PGEs, including Os, Ir, Ru, Rh, Pt and Pd), where estimates of average UCC can vary by factors of two or more. This study seeks to provide new constraints on the average abundances of the PGEs and the geochemically related Re, as well as Re-Os isotopic systematics in the UCC through time.

Owing to their highly siderophile and chalcophile nature, the PGEs tend to partition strongly into metallic and sulfide phases, and as a consequence, ~98% of the Earth's PGEs are estimated to be in the core (McDonough and Sun, 1995; McDonough, 2003). Accordingly, PGE concentrations in the silicate Earth are typically low (low ng/g to pg/g range). In addition, PGEs are often concentrated in minute accessory phases, which may be heterogeneously distributed within sample powder aliquots (giving rise to the so-called "nugget effect", Ravizza and Pyle, 1997) so that multiple analyses of the same sample powders may yield concentrations that vary beyond the precision of a single measurement. Both of these factors present challenges to estimating PGE abundances in the UCC.

Rhenium is a trace element that is geochemically similar to the PGEs in that it is strongly siderophile and can also be chalcophile. ^{187}Re decays to ^{187}Os (half-life = 4.16×10^{10} y), providing a radiogenic isotopic system that is useful for dating melt depletion events in the mantle (e.g., Carlson and Irving, 1994), molybdenites (e.g., Stein et al., 1997) and black shales (e.g., Ravizza and Turekian, 1989). In addition, during mantle melting, Re typically behaves as a moderately incompatible trace element, whereas Os is strongly compatible. Therefore, average crust has a much higher Re/Os ratio than the mantle (Shirey and Walker, 1998). Over time the high parent/daughter ratio results in the development of highly radiogenic $^{187}\text{Os}/^{188}\text{Os}$ in crustal rocks that can be used as a tracer for crust-mantle interaction (e.g., Saal et al., 1998). Characterizing the average Re/Os and $^{187}\text{Os}/^{188}\text{Os}$ ratios of the UCC, is, therefore, critical for mass-balance calculations involving this geochemically important reservoir.

Only a few attempts have previously been made to estimate the PGE and Re

abundances, and $^{187}\text{Os}/^{188}\text{Os}$ ratio of the UCC (Table 1, Fig. 1). Some of these efforts have estimated the UCC PGE abundances by taking the average concentrations in different rock types and weighting them in proportion to the relative surface area over which they are exposed (Shaw et al., 1976; Gao et al., 1998). Another approach is to use data for fine-grained sediments and sedimentary rocks (e.g., loess and shale) to infer the PGE and Re abundances in the UCC exposed to weathering and erosion (Esser and Turekian, 1993; Wedepohl, 1995; Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012). In addition, Schmidt et al. (1997a) used impact melts to estimate the PGE and Re abundances in the upper crust of the Baltic Shield. As shown in Table 1 and Fig. 1, the abundance estimates for Os, Ir and Re fall within a relatively narrow range of ~30%, 50% and ~30% deviation from their average values, respectively. However, Ru, Pt, and Pd estimates obtained from fine-grained sediments and sedimentary rocks (Wedepohl, 1995; Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012), are generally lower, in some cases by more than one order of magnitude, compared to averages determined from large area surface sampling (Gao et al., 1998) and impact melt samples (Schmidt et al., 1997a). In addition, most of the previous estimates are not accompanied by uncertainties, except for the relatively large uncertainties reported by Schmidt et al. (1997a, b).

In this study, we use **precise analytical techniques** to determine PGE and Re concentrations, as well as $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios for globally-distributed glacial diamictites deposited during the Mesoarchean, Paleoproterozoic, Neoproterozoic and Paleozoic eras. The major and trace element concentrations of these individual samples and composite samples (see section 2) have been reported by Gaschnig et al. (2014, 2016). Our objectives are to: 1) investigate how PGE and Re abundances, as well as

Re-Os isotopic systematics changed in the UCC over time, 2) provide new estimates of PGE and Re abundances, as well as $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios in the present-day UCC, and 3) assess the uncertainty of the new estimates.

2. Samples

About 150 individual diamictite samples were collected globally from four geological eras: the Mesoarchean (~2.9 Ga), Paleoproterozoic (~2.2-2.5 Ga), Neoproterozoic (~0.58-0.75 Ga) and Paleozoic (0.30 and 0.33 Ga) (Tables A1 and A2). Samples are composed of a fine-grained matrix containing a wide range of clasts with different shapes and sizes. The individual diamictite samples come from 24 stratigraphic units or time periods and, thus, 24 diamictite composites were made, with each composite representing one stratigraphic unit or time period (Tables A1 and A2). Geological background information about the stratigraphic units and the GPS co-ordinates of the individual samples are reported in Gaschnig et al. (2014).

3. Analytical techniques

Individual diamictites were first broken into smaller fragments using a rock hammer with the sample placed between thick plastic sheets. The fragments were then crushed in an alumina ceramic jaw crusher, and clasts larger than ~5 mm in diameter were removed from the chips by hand. The remaining, clast-free chips were powdered in an alumina ceramic swing mill. The composite samples were made from individual samples using equal proportions of sample powder (by mass) for each unit, combining 3 to 16

individual powders for each unit or time period. A “composite tree” is provided in Table A2.

About three grams of powder were spiked with appropriate amounts of ^{190}Os , ^{191}Ir , ^{99}Ru , ^{194}Pt , ^{105}Pd and ^{185}Re enriched isotopes prior to digestion. Digestion was conducted in a mixture of 2.5 mL of triple-distilled, concentrated HCl and 5 mL triple-distilled, concentrated HNO_3 in sealed Pyrex borosilicate Carius tubes heated to 270 °C for 4 days (Puchtel et al., 2014). Upon opening the cooled tubes, Os was extracted from the acid phase into CCl_4 and then back-extracted into concentrated HBr (Cohen and Waters, 1996). This was followed by further purification by microdistillation (Birck et al., 1997). Iridium, Ru, Pt, Pd and Re were separated and purified using anion exchange chromatography (Rehkämper and Halliday, 1997). It has been suggested that some Ru measurements reported in the literature may have been compromised due to the possible presence of Ni-argide interferences (Ravizza and Pyle, 1997; Park et al., 2012). This is especially problematic for the NiS fire assay digestion method, which involves addition of a large amount of Ni that can accompany the PGE fraction through the column chemistry. This is not a problem for the present study because no Ni was added to the samples in the Carius tube digestion method, and Ni from the samples was efficiently separated from the PGE fractions by the column separations used.

Osmium isotopic compositions were measured via negative thermal ionization mass spectrometry (N-TIMS, Creaser et al., 1991) at the University of Maryland (UMD), using a secondary electron multiplier (SEM) detector on a *Thermo-Fisher Triton* instrument. Purified Os, dissolved in HBr, was loaded onto baked Pt filaments (99.99% purity; *Materion*), dried down, then covered by a $\text{Ba}(\text{OH})_2$ activator solution. The instrumental mass bias was assessed by

191 correcting to $^{192}\text{Os}/^{188}\text{Os} = 3.083$ (Shirey and Walker, 1998), using the
 192 exponential fractionation law. The precision of measured $^{187}\text{Os}/^{188}\text{Os}$ ratios for
 193 each analysis of a sample or reference material (usually 100 cycles) was
 194 typically better than 0.1% ($2\sigma_m$, where σ_m refers to standard deviation of the
 195 mean, i.e., the standard deviation divided by the square root of the number of
 196 observations). An in-house Johnson-Matthey Os isotope standard reference
 197 solution yielded an average $^{187}\text{Os}/^{188}\text{Os}$ of 0.11374 ± 0.00016 (2σ , $n = 5$, σ
 198 refers to standard deviation), comparable to the result of 0.11378 obtained
 199 from precise measurements using Faraday cups during the entire analytical
 200 campaign on this *Triton*. The precision and accuracy of sample measurements
 201 were determined by replicate analyses of the reference material TDB-1
 202 (CANMET Diabase), which is widely used for low abundance PGE studies
 203 (Peucker-Ehrenbrink et al., 2003; Meisel and Moser, 2004a; Ishikawa et al.,
 204 2014). Analyses of TDB-1 yielded Os concentrations of 0.146 ± 0.016 (2σ , $n =$
 205 5) ng/g and $^{187}\text{Os}/^{188}\text{Os}$ of 0.94 ± 0.16 (2σ , $n = 5$) (Table 2). These values are
 206 within precisions of previously reported values (Table 2). The results of two
 207 other reference materials (UB-N and OKUM) are also consistent with values
 208 reported in the literature (Table 2). All Os measurements were blank corrected.
 209 The average total analytical blank for Os was 0.34 ± 0.44 (2σ , $n = 4$) pg, with a
 210 $^{187}\text{Os}/^{188}\text{Os}$ ratio was 0.141 ± 0.052 , making blank contributions for all
 211 analyzed samples less than 1% of the total Os measured.

212
 213 Measurements of Re, Ru, Pd, Ir and Pt were made at UMD using a single
 214 collector, sector field, inductively coupled plasma mass spectrometer (*Thermo*
 215 *Finnigan Element II*) with an electron multiplier detector using an *Aridus II*
 216 micro flow nebulizer sample introduction system. The precision was better
 217 than 1% ($2\sigma_m$) for the ratio measurements of $^{185}\text{Re}/^{187}\text{Re}$, $^{99}\text{Ru}/^{101}\text{Ru}$,
 218 $^{105}\text{Pd}/^{106}\text{Pd}$, $^{191}\text{Ir}/^{193}\text{Ir}$ and $^{194}\text{Pt}/^{195}\text{Pt}$ (100 cycles). Standard solutions of Ir and

Ru (from *Johnson-Matthey*), Re, Pd and Pt (from the *National Institute of Standards and Technology*) were run before and after every four samples to correct for mass bias. The reference material TDB-1 yielded 0.074 ± 0.016 ng/g Ir, 0.212 ± 0.023 ng/g Ru, 5.0 ± 1.1 ng/g Pt, 22.0 ± 2.2 ng/g Pd and 1.05 ± 0.14 ng/g Re (2σ , $n = 5$), in good agreement with published results (Table 2). The results of two other reference materials (UB-N and OKUM) are also consistent with values reported in the literature (Table 2). The average total analytical blanks were 0.23 ± 0.13 pg Ir, 3.2 ± 2.7 pg Ru, 62 ± 23 pg Pt, 7.0 ± 2.0 pg Pd and 1.78 ± 0.28 pg Re (2σ , $n = 4$). The blank contributions for all analyzed samples were less than 1.2% for Ir, 7.0% for Ru (excluding the Gaskiers composite, which is 13%), 8.1% for Pt (excluding the Konnarock composite, which is 14%), 1.2% for Pd and 5.4% for Re (excluding the Pocatello composite, which is 8.8%), and were corrected for all samples.

The analytical methods for total organic carbon (TOC) content can be found in Gaschnig et al. (2014). Briefly, about 5 g aliquots of powder were sequentially acidified with 3 M HCl for digesting carbonate and repeatedly washed with H₂O to get pH neutral solutions. Decalcified residues were dried down in an 80°C oven and then homogenized. About 2-5 mg residues were weighed into tin cups for organic carbon determination with a Eurovector elemental analyzer. Contents measured were normalized to whole rock values according to the percent carbonate determined during acidification. The Cr and Ni concentrations were determined by solution ICP-MS using a high-pressure bomb dissolution method (original data, as well as completed methods, are reported in Gaschnig et al. (2016)). As some of the Mesoarchean diamictites are chromite-bearing, it was important to completely digest chromite to extract all Cr and PGEs. Because it is black, it was easy to see if chromite remained undigested. No chromite was observed following dissolution in our study, or in

Gaschnig et al. (2016). Moreover, the much higher measured Cr and PGE concentrations relative to younger diamictites suggest complete dissolution of any original chromite.

4. Results

4.1 Individual diamictite samples

To evaluate sample-to-sample heterogeneity within stratigraphic units, and to assess whether the compositions of the composites are representative of the units and/or time periods, three individual diamictite samples from each of the Mozaan, Gowganda, and Nantuo Formations and the two Bolivian formations were analyzed (Table A3 and Fig. A1). The results show that, for a single formation and/or time period, PGE concentrations in the three individual samples are within the 50% deviation of their average, excluding Ru in the Gowganda Formation (69%), Pd in the Nantuo Formation (54%) and Os in the Bolivian formations (70%). In addition, concentrations of the four composites are within the concentration ranges of the corresponding three individual samples, respectively. However, the Re concentrations are highly variable, and potential reasons for this are discussed below. In general, the individual diamictite samples from one stratigraphic unit are relatively homogeneous in terms of PGE concentrations, and the PGE concentrations of the corresponding composites are representative of that of the stratigraphic units from which they derive.

4.2 Diamictite composites

Platinum-group element and Re concentrations, as well as Re-Os isotopic

data for the 24 diamictite composites are provided in Table 3. As shown in Fig. 2, measured concentrations are double-normalized, with each analysis corrected for quartz, carbonate, or iron dilution by normalizing it to the average UCC Al_2O_3 content (15.4%: Rudnick and Gao, 2003), and then normalizing individual elements to estimates for primitive upper mantle (PUM, Becker et al., 2006). To obtain the common Os concentrations (Table 3), radiogenic ^{187}Os produced by ^{187}Re decay was subtracted, based on the depositional ages reported in the literature (Table A1) and the measured $^{187}\text{Re}/^{188}\text{Os}$. Due to the high solubility of Re, the measured $^{187}\text{Re}/^{188}\text{Os}$ may not represent the time-integrated $^{187}\text{Re}/^{188}\text{Os}$. Thus, the correction is not very precise, but has little impact on the conclusions. The radiogenic Os accounts for <30% of the measured Os concentrations, except for the Ramsay Lake (36%) and Gowganda (35%) composites.

The four Mesoarchean composites have similar Os, Ir, and Re abundances (Fig. 2a). However, the Afrikander composite is characterized by higher Ru, Pt, and Pd concentrations, compared to the Mozaan, Promise, and Coronation composites. The Afrikander composite also has higher first-row transition metal concentrations than the other three Archean composites (Fig. A2a), suggesting a greater fraction of chromite derived from mafic to ultramafic rocks. The high Cr (1132 ug/g) and Ru (1.89 ng/g) concentrations for the Afrikander composite are most likely caused by the presence of laurite-bearing chromites.

Among the seven Paleoproterozoic diamictite composites, the Duitschland composite has the highest PGE concentrations (Fig. 2b) and generally higher first-row transition metal contents (Fig. A2b), like the Afrikander composite, suggesting a greater contribution from ultramafic to mafic lithologies. The six remaining composites have roughly similar PUM-normalized PGE and Re

patterns, which are systematically lower than those of Mesoarchean
composites (Fig. 3).

The PGE and Re results for 10 Neoproterozoic composites (Fig. 2c) are
characterized by similar PUM-normalized PGE patterns and generally similar
first-row transition metal contents (Fig. A2c), but scattered Re. These
composites have systematically lower PGE and Re concentrations than the
older composites (Fig. 3).

Among the three Paleozoic composites (Fig. 2d), the Dwyka West composite,
with the highest PGE concentrations, is similar to the Mesoarchean
composites. Moreover, its first-row transition metal contents are between those
of the Mesoarchean and Paleoproterozoic composites (Fig. A2d), and the age
distribution of detrital zircons from this formation, at this locality, suggests that
it is dominated by a pre-2.0 Ga provenance (Gaschnig et al., 2015).
Accordingly, this composite is not included with the other Paleozoic
composites in this study, both of which have similar PGE concentrations to the
Neoproterozoic composites (Fig. 3).

The Re-Os isotopic systematics of the composites are provided in Table 2 and
Fig. A3. The $^{187}\text{Os}/^{188}\text{Os}$ ratios are scattered, ranging from 0.2184 to 4.468.
The $^{187}\text{Re}/^{188}\text{Os}$ ratios are highly variable, ranging between 1.28 and 813.

5. Discussion

5.1 Sample representativeness

The idea of using glacial diamictites to estimate the average composition of the

UCC originated with Goldschmidt (1933), who pointed out that the major-element contents obtained from glacial loams in Norway favorably corresponded to the averages of igneous -rock compositions. The ability of glaciers to sample wide areas of exposed crust are documented in studies of modern glacial tills from the Guys Bight Basin (Nesbitt and Young, 1996) and British Columbia (Canil and Lacourse, 2011), both of which show that such tills provide robust averages of the rocks that the glaciers traversed. The diamictites studied here represent lithified tills (e.g., Huber et al., 2001). Using the major and trace element concentrations of the diamictite composites investigated here, Gaschnig et al. (2016) demonstrated that they are good proxies not only for average elemental abundances of the present-day UCC, but also for tracking the compositional changes of the average UCC through time.

Below we discuss three important factors regarding the representativeness of the diamictites for average UCC PGE concentrations.

First, the ice sheets must sample sufficiently large areas of the continent so as to minimize the influence of local sources. Although the concentrations of some elements (e.g., Nb and Ta) in certain samples, like those comprising the Dwyka West and the Pocatello composites, are affected by local provenance (Gaschnig et al., 2016), other geochemical features are observed globally amongst samples of similar ages, implying that these signatures are representative of the exposed UCC (Gaschnig et al., 2014, 2016). For example, within a given time period the rare earth elements (REE) show similar concentrations and similar UCC-normalized patterns, with Al-normalized REE concentrations decreasing systematically from the Phanerozoic/Neoproterozoic, to the Paleoproterozoic, to the Archean (Fig. A4;

Figs. 2 and 10 of Gaschnig et al., 2016). In a complementary fashion, the insoluble first-row transition element concentrations increase systematically from the Phanerozoic/Neoproterozoic, to the Paleoproterozoic, to the Archean (Fig. A2; Fig. 3 of Gaschnig et al., 2016). Both features reflect a greater proportion of mafic/ultramafic rocks in the Archean crust, as has been previously documented (Taylor and McLennan, 1985; Condie, 1993; Keller and Schoene, 2012; Tang et al., 2016). The glacial diamictites thus appear to be sampling large continental areas.

A second concern is whether the samples derive from a wide enough geographic region to represent a global average. While the diamictites from the three younger periods were sampled from different continents that were geographically dispersed at the time of deposition (Gaschnig et al., 2014), the Mesoarchean diamictites are all from two adjacent Archean sedimentary basins in South Africa, the Afrikander, Promise and Coronation Formations from the Witwatersrand Basin, and the Mozaan Formation from the Pongola Basin. Unfortunately, the incomplete nature of the geologic record leaves no better approach for studying the average composition of the Archean UCC through diamictites, as these are the only such samples of which we are aware. Moreover, there are no loess deposits of appropriate ages, and other fine-grained sediments (e.g., shale, mudstone) are the products of chemical weathering, which preferentially removes more soluble elements. The similarity in REE patterns (Fig. A5) and insoluble transition metals (Fig. 3 of Tang et al., 2016) between Archean diamictites and Archean shales suggest that the diamictites, like the shales, provide a representative sample of the Archean UCC. Several previous studies have suggested that some diamictites could be produced by catastrophic impact events (e.g., Oberbeck et al., 1993; Koeberl and Montanari, 2000), which may elevate the PGE concentrations of

the diamictites, leading to higher PGE concentrations in the Mesoarchean diamictite composites. However, based on petrographic and geochemical evidence, Huber et al. (2001) concluded that neither geochemical nor petrographic evidence supports an impact origin of the diamictites from the Witwatersrand Supergroup in South Africa.

Third, the composition of the diamictites should not be modified by syn- or post-depositional processes, such as density sorting and chemical weathering. The PGEs are mainly hosted in platinum group minerals that have densities up to 10 times higher than silicates (in the case of Os-Ir nuggets). Thus, transportation by wind may create a severe segregation of PGEs within these sediments. Such density sorting has impacted loess compositions, which show anomalously high concentrations of zirconium (477 ± 135 ug/g, 1σ , Taylor et al., 1983; Barth et al., 2000), which has been attributed to the eolian concentration of zircon. By contrast, the zirconium concentrations in diamictites (188 ± 99 ug/g, 1σ) are similar to the average UCC Zr abundance (193 ± 28 ug/g, 1σ , Rudnick and Gao, 2003), and are much lower than in loess (Taylor et al., 1983; Barth et al., 2000). Moreover, there is no evidence for wind sorting in the diamictites, which are, by definition, extremely poorly sorted. Consequently, unlike loess, wind sorting is unlikely to affect the composition of the glacial diamictites.

By contrast, nearly all of the diamictites record a weathering signature in their chemical compositions (Gaschnig et al., 2014, 2016; Li et al., 2016). For example, the chemical index of alteration ($CIA = \text{molar } Al_2O_3 / (K_2O + Na_2O + CaO^*)$, where CaO^* is CaO associated with the silicate fraction of the bulk sample) for most of the diamictites is high (~ 55 to ~ 75), above those of igneous rocks (< 55) (Gaschnig et al., 2016). Using Li and Pb isotopes, Li et al.

(2016) showed that the intensity of the weathering signature in the diamictites increases as one goes back in time and concluded that the weathering signature is largely inherited from the provenance of the diamictites; there has been very limited post- or syn-depositional chemical weathering. This inference is supported by correlations between PGE and insoluble transition elements, described in section 5.2.1, below (see also Figs. 4, A6 and A7), which imply little post- or syn-depositional modification of the PGE concentrations. If so, then the diamictites should provide a faithful record of the average UCC composition, including any weathered regolith that existed at the surface.

5.2 Present-day UCC PGE abundances and Re-Os isotopes

There are two ways to estimate the chemical composition of the present-day average UCC using fine-grained sedimentary rocks. One way is to adopt the median or average elemental concentrations in fine-grained terrigenous sedimentary rocks to represent the average UCC composition. This method can only be applied to insoluble elements (e.g., Th, Fe, Al and REE) that are not fractionated or affected during sedimentary processing (Taylor and McLennan, 1985; Rudnick and Gao, 2003; Gaschnig et al., 2016). In order to obtain a good estimate (e.g., small uncertainty), a large number of samples from different areas are generally needed. The other approach is to use the correlation between an element of interest and another element whose concentration is well constrained in the UCC (e.g., La) (Taylor and McLennan, 1985; McLennan, 2001; Rudnick and Gao, 2003; Hu and Gao, 2008; Gaschnig et al., 2016). Below we explore the utility of both approaches in estimating the average PGE and Re concentrations of the present-day UCC.

5.2.1 PGE abundances: median approach

In order to use this method, the mobility of the PGEs during chemical weathering and any chemical fractionation associated with physical weathering, sediment transport and sorting, as well as diagenesis, must be evaluated. Such influences may differ from element to element. Of the limited work that has been done to constrain PGE solubility, Os appears to be moderately soluble during oxidative weathering (Peucker-Ehrenbrink and Blum, 1998; Peucker-Ehrenbrink and Hannigan, 2000; Jaffe et al., 2002). Based on a study of fresh and weathered black shales, Peucker-Ehrenbrink and Hannigan (2000) concluded that Ir, Pt and Pd were also moderately soluble, whereas Jaffe et al (2002) inferred from a weathering profile through black shale that the three elements were essentially insoluble. By contrast, it is well established that Re is highly soluble in the near-surface, oxidized environment (Colodner et al., 1993; Peucker-Ehrenbrink and Hannigan, 2000; Jaffe et al., 2002). Therefore, PGEs tend to stay in the solid phase during chemical weathering and their concentrations in diamictites most likely represent their abundances in the UCC, but the same is not true for Re.

Here we compare inter-element correlations between PGEs and insoluble transition metals (e.g., Cr, Ni; Taylor and McLennan, 1985) with those between PGEs and soluble transition metals (e.g., Re, Mo; Taylor and McLennan, 1985) to infer the relative solubility of the PGEs during the formation of the diamictites. If the PGEs show significant positive correlations with insoluble transition metals, but no correlations with soluble transition metals, they were likely insoluble. In Figs. 4, A6 and A7, strong positive correlations of Ir, Ru, Pt, Pd versus Cr and Ni ($R^2 \geq 0.70$) and fair correlations of Os versus Cr and Ni ($R^2 = 0.58$) are observed for the diamictite composites. No correlations, however, are observed between the PGEs and Re or Mo (not shown). Therefore, Ir, Ru,

Pt and Pd are presumed to be insoluble, and Os is likely moderately insoluble, during the formation of these diamictites.

We use the median PGE concentrations of the post-Paleoproterozoic diamictite composites, which is less affected by extremely high or low values than average values, in order to estimate their abundances in the present-day UCC. The same method has been used by Gaschnig et al. (2016) to infer the major and trace element abundances in the present-day UCC from the post-Paleoproterozoic diamictites. Many studies, based on both sedimentary and igneous rocks, have demonstrated that the UCC composition changed at the Archean-Proterozoic boundary, after which no obvious change in composition is observed (Taylor and McLennan, 1985; Condie, 1993; Taylor and McLennan, 2008; Keller and Schoene, 2012; Tang et al., 2016). This is why Taylor and McLennan (1985) used post-Archean shales to estimate the REE abundances in the modern UCC. Since our Paleoproterozoic diamictites (~2.2-2.5 Ga) contain a higher proportion of Archean UCC rocks than the younger diamictites, we still see compositional changes between Paleoproterozoic and Neoproterozoic diamictites (e.g., REE and first-row transition metals). Therefore, we excluded both Archean and Paleoproterozoic diamictites when applying the median method. The Dwyka West composite (Paleozoic) was not incorporated here because its provenance is Archean crust (Gaschnig et al., 2015). The median values and the median absolute deviation (MAD, defined as the median of the absolute deviations of the measurements from the population median) are listed in Table 4. The average values of the post-Paleoproterozoic diamictite composites, which are not shown here, are similar to the medians.

5.2.2 PGE abundances: regression approach

The least-squares regression approach has been widely used to estimate trace element abundances in the UCC (McLennan, 2001; Rudnick and Gao, 2003; Hu and Gao, 2008) by calculating the slope and intercept of the regression, and then the projected element concentration of interest. However, the uncertainty of the projected concentration cannot be obtained by this method when the uncertainty of the regression itself is considered. Thus, to calculate the uncertainty on the projected concentration of the element of interest, we used a bootstrapping resampling method (Efron and Tibshirani, 1993; see Appendix "how bootstrapping works"). For example, for the Cr-Ir correlation (Fig. 4a), we first randomly resampled the Cr-Ir dataset ($n = 22$), allowing a single point to be sampled more than once. The size of each resample is equal to that of the original Cr-Ir dataset ($n = 22$). For each resample, we calculated a slope and an intercept using the least-squares regression, and a projected Ir concentration at the average upper crustal Cr abundance of 82 ± 16 ug/g ($2\sigma_m$) (see next section for derivation of this value). The uncertainties on the Cr and Ir concentrations for each composite and the recommended UCC Cr abundance are all taken into account. We repeated this step 10,000 times to get 10,000 resamples, which gives us 10,000 regressions and 10,000 projected Ir concentrations, respectively. We took the final average Ir concentration of the 10,000 projected Ir concentrations and the standard deviation as the UCC Ir abundance and associated precision, respectively. Below we perform the regressions using the composites from all four time periods ($n = 24$, unless otherwise indicated), as the modern UCC contains rocks with ages ranging from ~ 4.0 Ga to ~ 0 Ma and all the UCC rocks contribute to the composition of the modern UCC. Regressions were also performed using a subset that excluded the four Mesoarchean samples (results not shown here). The resulting PGE abundance estimates were found

to be identical, within uncertainty, with those obtained from the entire data set, which supports the idea that samples with all ages should be incorporated into the regression.

As shown in Fig. 4, Ir is strongly positively correlated with Cr ($R^2 = 0.92$) and Ni ($R^2 = 0.94$). The averages of published UCC abundance estimates for Cr (82 ± 16 ug/g, $2\sigma_m$, $n = 6$) and Ni (42 ± 10 ug/g, $2\sigma_m$, $n = 6$) (Condie, 1993; Gao et al., 1998; McLennan, 2001; Rudnick and Gao, 2003; Hu and Gao, 2008; Gaschnig et al., 2016) are taken as their UCC abundances. The Ir concentrations obtained from the above two regressions are within uncertainty of one another, with values of 0.036 ± 0.008 and 0.044 ± 0.012 ng/g (2σ), respectively. Considering the lower relative standard deviation, we adopt 0.036 ± 0.008 ng/g as the Ir abundance estimate of the UCC. Our Ir abundance is higher by 60-80% than the two prior estimates based on data from loess (Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012), and the surface sampling-based estimate (Shaw et al., 1976) (Table 1, Fig. 1). It is lower by ~40% than the estimates of Esser and Turekian (1993) and Wedepohl (1995), based upon river sediments and greywackes, respectively, and is similar to the value reported by Schmidt et al. (1997a), based upon impact melt rocks (Table 1, Fig. 1).

Common Os shows a strong positive correlation with Ir ($R^2 = 0.89$, Fig. 5a), and fair correlations with Cr ($R^2 = 0.58$) and Ni ($R^2 = 0.58$) (Fig. A6a-b), respectively. Using the above Ir concentration estimate, the good correlation between Ir and Os leads to an Os abundance of 0.054 ± 0.016 ng/g (2σ). This Os value is within uncertainties of those obtained from the other two regressions, which give concentrations of 0.066 ± 0.024 and 0.077 ± 0.028 ng/g (2σ) using the above UCC Cr and Ni estimates, respectively. Since Os correlates best with Ir and the Os-Ir correlation yields a lower relative standard

deviation, the concentration of 0.054 ± 0.016 ng/g obtained from the Ir-Os regression is taken as the average common Os abundance for the UCC. Adding radiogenic Os back to common Os using the Re abundance of 0.25 ± 0.12 ng/g (section 5.2.3), an Os abundance of 0.059 ± 0.016 ng/g (2σ) is obtained (Table 4). Our Os estimate is within uncertainty of those obtained by Esser and Turekian (1993), but is about twice as high as those of Schmidt et al. (1997a) and Peucker-Ehrenbrink and Jahn (2001) (Table 1, Fig. 1).

Figures 5b and A6c-d illustrate the good correlations observed between Ru and Ir ($R^2 = 0.86$), Cr ($R^2 = 0.98$) and Ni ($R^2 = 0.88$), respectively. These regressions yield Ru concentrations of 0.079 ± 0.026 , 0.082 ± 0.034 and 0.112 ± 0.046 ng/g (2σ), respectively. In view of the lower relative standard deviation of the result obtained from the Ir-Ru regression, the concentration obtained from this correlation (0.079 ± 0.026 ng/g) is adopted as our Ru concentration estimate for the UCC. Previous Ru abundance estimates for the UCC varied by nearly two orders of magnitude (0.03-1.06 ng/g, Table 1, Fig. 1) (Wedepohl, 1995; Schmidt et al., 1997a; Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012). The highest Ru abundance estimate of 1.06 ng/g (Table A5) was obtained from impact melt samples by Schmidt et al. (1997a), using NiS fire assay combined with neutron activation. The reason for the unexpected high Ru estimate remains unclear. Peucker-Ehrenbrink and Jahn (2001) estimated an abundance of 0.210 ng/g. This Ru estimate is nearly an order of magnitude higher than the estimate of 0.0030 ng/g reported by Park et al. (2012). Park et al. (2012) suggested the higher Ru concentrations reported by Peucker-Ehrenbrink and Jahn (2001) may have resulted from an analytical artifact. Our Ru estimate of 0.079 ng/g falls between the estimates of Park et al. (2012) and Wedepohl (1995) (0.1 ng/g, Table 1, Fig. 1).

As shown in Figs. 6a and A7a-b, Pt is strongly positively correlated with Ru ($R^2 = 0.89$), Cr ($R^2 = 0.87$) and Ni ($R^2 = 0.70$). These correlations yield similar Pt concentrations of 0.80 ± 0.22 , 0.80 ± 0.26 and 0.95 ± 0.34 ng/g (2σ), respectively. In light of the lower relative standard deviation of the result obtained from the Ru-Pt regression correlation, 0.80 ± 0.22 ng/g is taken as the Pt UCC abundance estimate. Palladium shows an excellent correlation with Pt ($R^2 = 0.99$) (Fig. 6b), and good correlations with Cr ($R^2 = 0.87$) and Ni ($R^2 = 0.74$) (Fig. A7c-d), yielding similar Pd concentrations of 0.80 ± 0.26 , 0.80 ± 0.28 and 0.96 ± 0.36 ng/g (2σ), respectively. Here, we use 0.80 ± 0.26 ng/g as the Pd abundance for the UCC because of the lower relative standard deviation. **Because Ir, Ru, Pt and Pd become progressively more incompatible in many silicate systems from Ir to Pd (e.g., Pearson et al., 2004), correlations between Ru-Pt are better than Ir-Pt, and Pt-Pd is better than Ir-Pd. Although the Pt and Pd abundance estimates obtained from Ir-Pt and Ir-Pd correlations are within uncertainties of those from Ru-Pt and Pt-Pd correlations, the latter yield lower uncertainties. This is why here we use Ru-Pt and Pt-Pd correlations, instead of Ir-Pt and Ir-Pd correlations.** The Pt and Pd concentrations obtained by Gao et al. (1998) and Schmidt et al. (1997a), are generally three times higher than those derived from sedimentary averages (Table 1, Fig. 1) (Wedepohl, 1995; Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012). Our Pt and Pd estimates lie between these two types of estimates (Table 1, Fig. 1). In contrast to the differences in absolute concentrations, all of the previous Pd/Pt ratio estimates, as well as the one from this study, are similar, ranging from 0.88 to 1.02 (with a slightly higher ratio of 1.33 from Schmidt et al., 1997a). This suggests that these two elements are not fractionated during **weathering and sediment transport**, and have similar abundances in the UCC.

5.2.3 Re-Os isotopes

Rhenium is highly soluble in the near-surface, oxidized environment (Colodner et al., 1993; Peucker-Ehrenbrink and Hannigan, 2000; Jaffe et al., 2002), as is reflected in the weak correlation between Re and the soluble element Mo (Fig. 7b). This results in highly variable Re concentrations in different types of sedimentary samples, and a general lack of correlation between Re and other trace elements. Estimating the UCC Re abundance and $^{187}\text{Re}/^{188}\text{Os}$ ratio is, thus, challenging. Esser and Turekian (1993) used Fe-Ir-Os correlations in river sediments to estimate the UCC Os abundance. Then they used sample $^{187}\text{Os}/^{188}\text{Os}$ ratios and associated Nd model ages to calculate a time-integrated $^{187}\text{Re}/^{188}\text{Os}$ ratio for each sample, assuming that Os has the same crustal residence age as Nd. The average of the obtained $^{187}\text{Re}/^{188}\text{Os}$ ratios of all samples was taken as the present-day UCC $^{187}\text{Re}/^{188}\text{Os}$ ratio. Thus, the present-day UCC $^{187}\text{Os}/^{188}\text{Os}$ ratio and Re concentration can be recalculated using the above average $^{187}\text{Re}/^{188}\text{Os}$ ratio and the average Nd model age for the North American basement (~2.2 Ga, Esser and Turekian, 1993). A similar approach was applied to loess samples from China by Peucker-Ehrenbrink and Jahn (2001).

This method, however, cannot be applied to all of the diamictite composites, because they are characterized by much more variable $^{187}\text{Os}/^{188}\text{Os}$ ratios than river sediments and loess samples (Fig. A3). Therefore, contrary to the two previous studies (Esser and Turekian, 1993; Peucker-Ehrenbrink and Jahn, 2001) in which the UCC Os abundance and $^{187}\text{Re}/^{188}\text{Os}$ ratio were first estimated, and then the UCC $^{187}\text{Os}/^{188}\text{Os}$ ratio and Re abundance were calculated, here we first estimate the UCC Os and Re abundances using the regression approach. We then calculate the UCC $^{187}\text{Re}/^{188}\text{Os}$ ratio and finally obtain the UCC $^{187}\text{Os}/^{188}\text{Os}$ ratio.

The geochemical behavior of Re is generally similar to that of Mo. In oxidizing aqueous environments, Re and Mo are oxidized to ReO_4^- and MoO_4^{2-} , which are highly soluble, and thus enriched in seawater relative to UCC rocks (Ravizza et al., 1991; Anbar et al., 1992; Colodner et al., 1993). Under sulfidic conditions, both elements can be enriched in marine sediments (Crusius et al., 1996; Morford and Emerson, 1999). The difference is that Re, but not Mo, can also be enriched in sediments under suboxic conditions (Crusius et al., 1996; Morford and Emerson, 1999; Miller et al., 2011), suggesting that Re is scavenged more readily than Mo. This difference is reflected in the diamictites, which generally show a weathering signature inherited from their provenance (Li et al., 2016). A weak correlation between Re and total organic carbon (TOC) is observed in the diamictite composites (Fig. 7a), whereas there is no correlation between Mo and TOC (not shown). The organic carbon contents in these diamictite composites are relatively low ($<0.4\%$), and the observed correlations (or lack thereof) suggests limited Mo scavenging, but some Re scavenging. Excluding the five diamictites with the highest TOC in order to eliminate samples that experienced Re scavenging, we find a weak correlation between Re and Mo in the diamictite composites (Fig. 7b).

From this correlation, an UCC Re abundance of 0.25 ± 0.12 ng/g (2σ) can be obtained by the bootstrapping resample method, using the average of UCC Mo estimates (0.9 ± 0.3 ug/g, $2\sigma_m$, $n = 6$) from the literature (Sims et al., 1990; Wedepohl, 1995; Gao et al., 1998; Rudnick and Gao, 2003; Hu and Gao, 2008; Gaschnig et al., 2016). The relatively large standard deviation on this estimate is caused by the large uncertainty of the UCC Mo abundance and the weak correlation between Re and Mo. Additionally, the Re concentrations in igneous rocks are significantly lower (up to three to four orders of magnitude) than

those of organic-rich sediments (Dubin and Peucker-Ehrenbrink, 2015). The extreme heterogeneity of Re concentrations in UCC rocks and the disproportionate influence of heterogeneously distributed organic-rich sediments on the UCC average (Dubin and Peucker-Ehrenbrink, 2015) also provide challenges in estimating UCC Re abundance with a relatively small uncertainty. Our estimate lies between the estimates reported by the two previous studies (Table 1, Fig. 1, Esser and Turekian, 1993; Peucker-Ehrenbrink and Jahn, 2001).

Using the new Re and Os estimates from this work, the $^{187}\text{Re}/^{188}\text{Os}$ ratio of the UCC is calculated to be 20 ± 12 (2σ) (Table 1). Assuming an average age of the UCC of ~ 2.0 Ga (Allègre and Rousseau, 1984), the present-day $^{187}\text{Os}/^{188}\text{Os}$ ratio of the UCC is 0.80 ± 0.38 (2σ). Our lower estimates of $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios than the previous studies (Esser and Turekian, 1993; Peucker-Ehrenbrink and Jahn, 2001; Dubin and Peucker-Ehrenbrink, 2015) are mainly caused by our higher UCC Os abundance estimate (Table 1). The large uncertainty of our estimated $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios mainly result from the large uncertainty of the estimated Re concentration.

Another way to estimate the $^{187}\text{Os}/^{188}\text{Os}$ ratio of the present-day UCC is, similar to the “Median approach”, to use the median value of the Neoproterozoic and Paleozoic composites (again excluding the Dwyka West composite for the reasons stated above), yielding a result of 1.26 ± 0.32 ($n = 12$, MAD). This ratio is consistent, within uncertainty, with the average $^{187}\text{Os}/^{188}\text{Os}$ ratio of the Neoproterozoic and Paleozoic composites, which is 1.47 ± 0.61 ($n = 12$, 95% confidence interval). Both estimates are slightly higher, but within uncertainty of the ratio (0.80 ± 0.38 (2σ)) calculated from the

UCC Re and Os abundances. Considering that the sample $^{187}\text{Os}/^{188}\text{Os}$ ratio can be readily affected by the concentration of Re, which is a highly soluble element and shows scattered concentrations in diamictite composites (from 0.0068 to 3.68 ppb), and the $^{187}\text{Os}/^{188}\text{Os}$ ratios of the Neoproterozoic and Paleozoic composites are still variable (from 0.4229 to 3.381), the median or average $^{187}\text{Os}/^{188}\text{Os}$ ratio may not be representative of the average UCC. Here we take 0.80 ± 0.38 (2σ) and 20 ± 12 (2σ) as our best estimate of $^{187}\text{Os}/^{188}\text{Os}$ and $^{187}\text{Re}/^{188}\text{Os}$ ratios of the present-day UCC.

5.3 The role of old crust in the present-day UCC

Both loess and diamictite have been used as proxies for the composition of the UCC (Taylor et al., 1983; Gallet et al., 1998; Jahn et al., 2001; McLennan, 2001; Peucker-Ehrenbrink and Jahn, 2001; Rudnick and Gao, 2003; Hu and Gao, 2008; Park et al., 2012; Gaschnig et al., 2016). Our present-day UCC PGE abundance estimates from the medians or averages of post-Paleoproterozoic composites are similar to those obtained from loess samples, (Tables 1 and 4, Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012). These estimates are systematically lower than those obtained by the regression approach using all diamictite composites in this study (Table 4). We suggest the reason for the difference is that loess and our Neoproterozoic and Paleozoic diamictites preferentially sample younger UCC rocks, which have lower PGE, Cr and Ni concentrations than older crust, as reflected in shales (Taylor and McLennan, 1985; Condie, 1993), large scale surface averages (Eade and Fahrig, 1973; Condie, 1993), and data compilation (Keller and Schoene, 2012). This is supported by the fact that loess samples generally have Nd model ages of ~ 1.0 - 1.7 Ga (Taylor et al., 1983; Gallet et al., 1998; Peucker-Ehrenbrink and Jahn, 2001), suggesting a relatively younger

provenance than the average age of the UCC (~1.9-2.3 Ga, Allègre and Rousseau, 1984). Glaciers may also preferentially sample younger crust given that many of the diamictites are submarine deposits formed on the margins of continents, which can be younger than continental interiors (Peucker-Ehrenbrink et al., 2010). Additional studies of diamictites, such as detrital zircon ages (Gaschnig et al., 2015 and in progress), and whole rock Hf-Nd isotope analyses are needed to evaluate the age of crust sampled by the diamictites. Another possibility for the lower estimates obtained from loess samples is that PGEs are fractionated during wind transportation owing to the larger densities of platinum-group minerals (PGM) relative to silicates.

Therefore, we suggest that the PGE abundance estimates obtained from the median PGE concentrations of the Neoproterozoic and Paleozoic diamictites and from the Quaternary loess samples are biased toward lower values, resulting from the foregoing sampling bias. This bias can be ameliorated by using the regression approach that encompasses diamictites of all ages, ranging from the Mesoarchean to Paleozoic. If an unknown element correlates with another element in rocks or sediments with all ages (not only relatively young samples, e.g., loess) for which a robust average concentration has been established for the UCC, the concentration for the unknown element obtained from the regression should be a reasonable estimate for its abundance in the modern UCC. We consider the results derived from the regression as our best estimates of the UCC PGE and Re abundances.

5.4 Secular changes in PGE-Re abundances in the UCC

Previous estimates of PGE and Re abundances in the UCC focused only on the present-day UCC composition (Esser and Turekian, 1993; Wedepohl, 1995;

Schmidt et al., 1997a; Gao et al., 1998; Peucker-Ehrenbrink and Jahn, 2001; Park et al., 2012). No studies have been conducted to assess whether PGE and Re concentrations in the UCC have varied through time. From the observed higher concentrations of first-row transition metals in averages of surface outcrops and shales, the early continental crust is hypothesized to have had a greater proportion of mafic and ultramafic lithologies than younger continental crust (Eade and Fahrig, 1973; Taylor and McLennan, 1985; Condie, 1993; Keller and Schoene, 2012; Tang et al., 2016). Similar secular changes have also been recognized in individual diamictites (Gaschnig et al., 2014) and the diamictite composites studied here (Gaschnig et al., 2016). The similarities seen in glacial diamictites and shales suggest that the former are also useful proxies for studying the composition and evolution of the UCC (Gaschnig et al., 2016). It may be expected, then, that the older diamictite composites have higher PGE concentrations than younger ones.

Figures 2 and 3 show that PGE concentrations decrease in the diamictite composites progressing from the Mesoarchean to the Neoproterozoic, after which no change is apparent. Student's *t*-tests (Table A7) demonstrate that the older diamictite composites generally have statistically higher PGE concentrations than younger ones and that the Neoproterozoic composites are not statistically different from the Paleozoic composites. This trend is consistent with the higher concentrations of first-row transition metals and lower Al-normalized REE concentrations in the older diamictites (Figs. A2 and A4), and is also consistent with previous inference that there was a greater proportion of mafic to ultramafic lithologies in the Archean UCC compared to younger UCC. Indeed, Condie (1993) estimated that the Early Archean UCC had higher proportions of komatiite (~6%) and basalt (~15%) compared to the Proterozoic (~0% and 10%, respectively) and the Phanerozoic UCC (~0% and

9%, respectively). Using the average PGE concentrations of rock types compiled from GeoRoc (Table A5), and the relative lithological proportions in the UCC estimated by Condie (1993) and Tang et al. (2016) (Table A6), one can observe the important influence of komatiite on UCC PGE concentrations. For example, adding 1% more komatiite to the Early Archean UCC results in an Ir concentration that is ~10% higher in Condie's model. In contrast to PGE concentrations, Re concentrations are highly variable and show no trend, likely owing to its high solubility (see section 5.1.2.4).

Using the PGE data in Table A5 and the rock proportions in Table A6, we estimate the average PGE abundances for the Mesoarchean and Phanerozoic UCC (Table 4). Owing to the rarity of PGE concentration data in Proterozoic rocks, the average PGE abundances for Proterozoic UCC are not calculated here. Four points can be drawn from Table 4. First, the medians of the Mesoarchean diamictite PGE data lie between the PGE abundances of the Mesoarchean UCC calculated from Condie's model and Tang's et al model. The diamictite Ru median is close to the Ru abundance obtained from Tang's et al., model considering the large uncertainty. And the PGE abundances for the Phanerozoic UCC obtained from Condie's (1993) model calculation are quite similar to the medians of the Phanerozoic diamictite composites. Second, the PGE abundances in the Mesoarchean UCC obtained from diamictite composites and the model calculations are higher than those for the Phanerozoic UCC. Third, given the similarity between the results from model calculation and the composite medians for both the Mesoarchean and Phanerozoic periods, we suggest that the medians of the PGE concentrations for Paleoproterozoic and Neoproterozoic diamictite composites are representative of the PGE abundances for the Paleoproterozoic and Neoproterozoic UCC. Finally, the medians of the PGE concentrations of

Neoproterozoic diamictite composites are indistinguishable from the Paleozoic diamictite composites, consistent with their similar concentrations of first-row transition metals as well as other elements (Fig. A2). These observations are consistent with the observation that the UCC composition has not changed significantly since the Archean-Proterozoic boundary (e.g., Gaschnig et al., 2016; Tang et al., 2016).

Conclusions

Globally-distributed glacial diamictites, which were deposited during the Mesoproterozoic, Paleoproterozoic, Neoproterozoic and Paleozoic eras, are good proxies not only for estimating the average composition of the present-day UCC, but also for tracking the UCC composition through time (Gaschnig et al., 2014, 2016; Li et al., 2016). The concentrations of platinum-group elements (PGEs, including Os, Ir, Ru, Pt and Pd) and the geochemically related Re, as well as $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios reported here lead to the following conclusions:

1) The correlations between PGEs and Cr, Ni, and between Re and Mo, in all the diamictite composites can be used to estimate the PGE-Re abundances in the present-day UCC, with associated uncertainties (0.059 ± 0.016 ng/g Os, 0.036 ± 0.008 ng/g Ir, 0.079 ± 0.026 ng/g Ru, 0.80 ± 0.22 ng/g Pt, 0.80 ± 0.26 ng/g Pd and 0.25 ± 0.12 ng/g Re, 2σ of 10,000 bootstrapping regression results). From this, time-integrated $^{187}\text{Re}/^{188}\text{Os}$ and $^{187}\text{Os}/^{188}\text{Os}$ ratios for the UCC are calculated, yielding ratios of 20 ± 12 and 0.80 ± 0.38 (2σ), respectively.

2) These PGE abundance estimates are slightly higher than the estimates

obtained from loess samples. We suggest two possible reasons for the difference: 1) loess preferentially samples younger UCC rocks that have lower PGE concentrations; 2) PGEs are fractionated during wind transportation during the formation of loess.

3) The PGE concentrations decrease in the diamictite composites progressing from the Mesoarchean to the Neoproterozoic, after which no change is apparent. This trend is consistent with the higher concentrations of first-row transition metals and lower Al-normalized REE concentrations in the older diamictites, and is also consistent with previous inference that there was a greater proportion of mafic to ultramafic lithologies in the Archean UCC compared to younger UCC.

Acknowledgements

This research was supported by the U.S. National Science Foundation (EAR-1321954 and FESD EAR-1338810), the National Nature Science Foundation of China (41373026 and 41173016), Chinese Ministry of Education (B07039) and the State Key Laboratory of Geological Processes and Mineral Resources (MSFGPMR01 and GPMR201202). We thank Richard Ash, Gregory Archer, Emily Worsham and Katherine Bermingham for help with analytical work and William McDonough, Yongsheng Liu and Huan Cui for helpful discussions. We thank Bernhard Peucker-Ehrenbrink and Ian H. Campbell for comments on a previous draft of this manuscript. Finally, we greatly appreciate the editorial work of Munir Humayun and detailed comments from Greg Ravizza, Bernhard Peucker-Ehrenbrink, and an anonymous reviewer that helped to improve the presentation.

Appendices

Appendices (Tables A1, A2, A3, A4, A5, A6 and A7, Figs. A1, A2, A3, A4, A5, A6 and A7, and “How bootstrapping works”,) associated with this article can be found with the online version.

References

- Allègre C. J. and Rousseau D. (1984) The growth of the continent through geological time studied by Nd isotope analysis of shales. *Earth Planet. Sci. Lett.* **67**, 19-34.
- Anbar A. D., Creaser R. A., Papanastassiou D. A. and Wasserburg G. J. (1992) Rhenium in seawater: Confirmation of generally conservative behavior. *Geochim. Cosmochim. Acta* **56**, 4099-4103.
- Becker H., Horan M. F., Walker R. J., Gao S., Lorand J. P. and Rudnick R. L. (2006) Highly siderophile element composition of the Earth's primitive upper mantle: Constraints from new data on peridotite massifs and xenoliths. *Geochim. Cosmochim. Acta* **70**, 4528-4550.
- Birck J. L., RoyBarman M. and Capmas F. (1997) Re-Os isotopic measurements at the femtomole level in natural samples. *Geostan. Newsl.* **21**, 19-27.
- Canil D. and Lacourse T. (2011) An estimate for the bulk composition of juvenile upper continental crust derived from glacial till in the North American Cordillera. *Chem. Geol.* **284**, 229-239.
- Carlson R. W. and Irving A. J. (1994) Depletion and enrichment history of subcontinental lithospheric mantle: An Os, Sr, Nd and Pb isotopic study of ultramafic xenoliths from the northwestern Wyoming Craton. *Earth Planet. Sci. Lett.* **126**, 457-472.

876 Cohen A. S. and Waters F. G. (1996) Separation of osmium from geological
 877 materials by solvent extraction for analysis by thermal ionisation mass
 878 spectrometry. *Anal. Chim. Acta* **332**, 269-275.

879 Colodner D., Sachs J., Ravizza G., Turekian K., Edmond J. and Boyle E. (1993)
 880 The geochemical cycle of rhenium: a reconnaissance. *Earth Planet. Sci.*
 881 *Lett.* **117**, 205-221.

882 Condie K. C. (1993) Chemical composition and evolution of the upper
 883 continental crust: Contrasting results from surface samples and shales.
 884 *Chem. Geol.* **104**, 1-37.

885 Creaser R. A., Papanastassiou D. A. and Wasserburg G. J. (1991) Negative
 886 thermal ion mass spectrometry of osmium, rhenium and iridium. *Geochim.*
 887 *Cosmochim. Acta* **55**, 397-401.

888 Crusius J., Calvert S., Pedersen T. and Sage D. (1996) Rhenium and
 889 molybdenum enrichments in sediments as indicators of oxic, suboxic and
 890 sulfidic conditions of deposition. *Earth Planet. Sci. Lett.* **145**, 65-78.

891 Dubin A. and Peucker-Ehrenbrink B. (2015) The importance of organic-rich
 892 shales to the geochemical cycles of rhenium and osmium. *Chem. Geol.*
 893 **403**, 111-120.

894 Eade K. E. and Fahrig W. F. (1973) Regional, lithological, and temporal
 895 variation in the abundances of some trace elements in the Canadian
 896 Shield. *Geol. Surv. Can. Paper* **72**, 46.

897 Efron B. and Tibshirani R. J. (1993) *An Introduction to the Bootstrap*. Taylor &
 898 Francis.

899 Esser B. K. and Turekian K. K. (1993) The osmium isotopic composition of the
 900 continental crust. *Geochim. Cosmochim. Acta* **57**, 3093-3104.

901 Gallet S., Jahn B.-m., Van Vliet Lanoë B., Dia A. and Rossello E. (1998) Loess
 902 geochemistry and its implications for particle origin and composition of
 903 the upper continental crust. *Earth Planet. Sci. Lett.* **156**, 157-172.

904 Gao S., Luo T. C., Zhang B. R., Zhang H. F., Han Y. W., Zhao Z. D. and Hu Y. K.
 905 (1998) Chemical composition of the continental crust as revealed by
 906 studies in East China. *Geochim. Cosmochim. Acta* **62**, 1959-1975.

907 Gaschnig R. M., Rudnick R. L., McDonough W. F., Kaufman A. J., Hu Z. and
 908 Gao S. (2014) Onset of oxidative weathering of continents recorded in
 909 the geochemistry of ancient glacial diamictites. *Earth Planet. Sci. Lett.*
 910 **408**, 87-99.

911 Gaschnig R. M., Rudnick R. L., McDonough W. F., Kaufman A. J., Vervoort J.
 912 D. and Fisher C. (2015) Insights on crustal growth from detrital zircons in
 913 ancient glacial deposits, American Geophysical Union Fall Meeting, San
 914 Francisco, pp. V51C-3053.

915 Gaschnig R. M., Rudnick R. L., McDonough W. F., Kaufman A. J., Valley J. W.,
 916 Hu Z., Gao S. and Beck M. L. (2016) Compositional evolution of the
 917 upper continental crust through time, as constrained by ancient glacial
 918 diamictites. *Geochim. Cosmochim. Acta* **186**, 316-343.

919 Goldschmidt V. M. (1933) Grundlagen der quantitativen Geochemie.
 920 *Fortschritte der Mineralogie, Kristallographie und Petrographie* **17**,
 921 112-156.

922 Hu Z. and Gao S. (2008) Upper crustal abundances of trace elements: A
 923 revision and update. *Chem. Geol.* **253**, 205-221.

924 Huber H., Koeberl C., McDonald I. and Reimold W. U. (2001) Geochemistry
 925 and petrology of Witwatersrand and Dwyka diamictites from South Africa:
 926 search for an extraterrestrial component. *Geochim. Cosmochim. Acta* **65**,
 927 2007-2016.

928 Ishikawa A., Senda R., Suzuki K., Dale C. W. and Meisel T. (2014)
 929 Re-evaluating digestion methods for highly siderophile element and
 930 ¹⁸⁷Os isotope analysis: Evidence from geological reference materials.
 931 *Chem. Geol.* **384**, 27-46.

- 932 Jaffe L. A., Peucker-Ehrenbrink B. and Petsch S. T. (2002) Mobility of rhenium,
933 platinum group elements and organic carbon during black shale
934 weathering. *Earth Planet. Sci. Lett.* **198**, 339-353.
- 935 Jahn B.-m., Gallet S. and Han J. (2001) Geochemistry of the Xining, Xifeng
936 and Jixian sections, Loess Plateau of China: eolian dust provenance and
937 paleosol evolution during the last 140 ka. *Chem. Geol.* **178**, 71-94.
- 938 Kamber B. S., Greig A. and Collerson K. D. (2005) A new estimate for the
939 composition of weathered young upper continental crust from alluvial
940 sediments, Queensland, Australia. *Geochim. Cosmochim. Acta* **69**,
941 1041-1058.
- 942 Keller C. B. and Schoene B. (2012) Statistical geochemistry reveals disruption
943 in secular lithospheric evolution about 2.5 Gyr ago. *Nature* **485**, 490-493.
- 944 Koeberl C. and Montanari A. (2000) Impact stratigraphy : the Italian record.
- 945 Li S., Gaschnig R. M. and Rudnick R. L. (2016) Insights into chemical
946 weathering of the upper continental crust from the geochemistry of
947 ancient glacial diamictites. *Geochim. Cosmochim. Acta* **176**, 96-117.
- 948 Maier W. D., Peltonen P., McDonald I., Barnes S. J., Barnes S. J., Hatton C.
949 and Viljoen F. (2012) The concentration of platinum-group elements and
950 gold in southern African and Karelian kimberlite-hosted mantle xenoliths:
951 Implications for the noble metal content of the Earth's mantle. *Chem.*
952 *Geol.* **302–303**, 119-135.
- 953 McDonough W. F. and Sun S. S. (1995) The composition of the Earth. *Chem.*
954 *Geol.* **120**, 223-253.
- 955 McDonough W. F. (2003) 2.15 - Compositional Model for the Earth's Core, in:
956 Holland, H.D., Turekian, K.K. (Eds.), *Treatise on Geochemistry*.
957 Pergamon, Oxford, pp. 547-568.
- 958 McLennan S. M. (2001) Relationships between the trace element composition
959 of sedimentary rocks and upper continental crust. *Geochem. Geophys.*

960 Geosyst. **2**, 2000GC000109.

961 Meisel T. and Moser J. (2004a) Reference materials for geochemical PGE
 962 analysis: new analytical data for Ru, Rh, Pd, Os, Ir, Pt and Re by isotope
 963 dilution ICP-MS in 11 geological reference materials. *Chem. Geol.* **208**,
 964 319-338.

965 Meisel T. and Moser J. (2004b) Platinum-Group Element and Rhenium
 966 concentrations in low abundance reference materials. *Geostand.*
 967 *Geoanal. Res.* **28**, 233-250.

968 Miller C. A., Peucker-Ehrenbrink B., Walker B. D. and Marcantonio F. (2011)
 969 Re-assessing the surface cycling of molybdenum and rhenium. *Geochim.*
 970 *Cosmochim. Acta* **75**, 7146-7179.

971 Morford J. L. and Emerson S. (1999) The geochemistry of redox sensitive
 972 trace metals in sediments. *Geochim. Cosmochim. Acta* **63**, 1735-1750.

973 Nesbitt H. W. and Young G. M. (1996) Petrogenesis of sediments in the
 974 absence of chemical weathering: effects of abrasion and sorting on bulk
 975 composition and mineralogy. *Sedimentology* **43**, 341-358.

976 Oberbeck V. R., Marshall J. R. and Aggarwal H. (1993) Impacts, Tillites, and
 977 the Breakup of Gondwanaland. *J. Geol.* **101**, 1-19.

978 Park J.-W., Hu Z., Gao S., Campbell I. H. and Gong H. (2012) Platinum group
 979 element abundances in the upper continental crust revisited-New
 980 constraints from analyses of Chinese loess. *Geochim. Cosmochim. Acta*
 981 **93**, 63-76.

982 Pearson D. G., Irvine G. J., Ionov D. A., Boyd F. R. and Dreibus G. E. (2004)
 983 Re-Os isotope systematics and platinum group element fractionation
 984 during mantle melt extraction: a study of massif and xenolith peridotite
 985 suites. *Chem. Geol.* **208**, 29-59.

986 Peucker-Ehrenbrink B. and Blum J. D. (1998) Re-Os isotope systematics and
 987 weathering of Precambrian crustal rocks: implications for the marine

988 osmium isotope record. *Geochim. Cosmochim. Acta* **62**, 3193-3203.

989 Peucker-Ehrenbrink B. and Hannigan R. E. (2000) Effects of black shale
 990 weathering on the mobility of rhenium and platinum group elements.
 991 *Geology* **28**, 475-478.

992 Peucker-Ehrenbrink B. and Jahn B.-m. (2001) Rhenium-osmium isotope
 993 systematics and platinum group element concentrations: Loess and the
 994 upper continental crust. *Geochem. Geophys. Geosyst.* **2**,
 995 2001GC000172.

996 Peucker-Ehrenbrink B., Bach W., Hart S. R., Blusztajn J. S. and Abbruzzese T.
 997 (2003) Rhenium-osmium isotope systematics and platinum group
 998 element concentrations in oceanic crust from DSDP/ODP Sites 504 and
 999 417/418. *Geochem. Geophys. Geosyst.* **4**, 8911.

1000 Peucker-Ehrenbrink B., Miller M. W., Arsouze T. and Jeandel C. (2010)
 1001 Continental bedrock and riverine fluxes of strontium and neodymium
 1002 isotopes to the oceans. *Geochem. Geophys. Geosyst.* **11**, Q03016.

1003 Puchtel I. S., Walker R. J., Touboul M., Nisbet E. G. and Byerly G. R. (2014)
 1004 Insights into early Earth from the Pt–Re–Os isotope and highly
 1005 siderophile element abundance systematics of Barberton komatiites.
 1006 *Geochim. Cosmochim. Acta* **125**, 394-413.

1007 Ravizza G. and Turekian K. K. (1989) Application of the ¹⁸⁷Re-¹⁸⁷Os system
 1008 to black shale geochronometry. *Geochim. Cosmochim. Acta* **53**,
 1009 3257-3262.

1010 Ravizza G., Turekian K. K. and Hay B. J. (1991) The geochemistry of rhenium
 1011 and osmium in recent sediments from the Black Sea. *Geochim.*
 1012 *Cosmochim. Acta* **55**, 3741-3752.

1013 Ravizza G. and Pyle D. (1997) PGE and Os isotopic analyses of single sample
 1014 aliquots with NiS fire assay preconcentration. *Chem. Geol.* **141**, 251-268.

1015 Rehkämper M. and Halliday A. N. (1997) Development and application of new

ion exchange techniques for the separation of the platinum group
and other siderophile elements from geological samples. *Talanta* **44**,
663-672.

Rudnick R. L. and Gao S. (2003) 3.01 - Composition of the Continental Crust,
in: Holland, H.D., Turekian, K.K. (Eds.), *Treatise on Geochemistry*, 1st ed.
Pergamon, Oxford, pp. 1-64.

Saal A. E., Rudnick R. L., Ravizza G. E. and Hart S. R. (1998) Re-Os isotope
evidence for the composition, formation and age of the lower continental
crust. *Nature* **393**, 58-61.

Savard D., Barnes S.-J. and Meisel T. (2010) Comparison between
Nickel-Sulfur Fire Assay Te Co-precipitation and Isotope Dilution with
High-Pressure Asher Acid Digestion for the Determination of
Platinum-Group Elements, Rhenium and Gold. *Geostand. Geoanal. Res.*
34, 281-291.

Schmidt G., Palme H. and Kratz K.-L. (1997a) Highly siderophile elements (Re,
Os, Ir, Ru, Rh, Pd, Au) in impact melts from three European impact
craters (Sääksjärvi, Mien, and Dellen): Clues to the nature of the
impacting bodies. *Geochim. Cosmochim. Acta* **61**, 2977-2987.

Schmidt G., Palme H. and Kratz K.-L. (1997b) Fractionation of highly
siderophile elements in the Earth's upper continental crust. *Yearbook of
Institute of Nuclear Chemistry, Johannes Gutenberg-Universität, Mainz,
Germany*.

Shaw D. M., Dostal J. and Keays R. R. (1976) Additional estimates of
continental surface Precambrian shield composition in Canada. *Geochim.
Cosmochim. Acta* **40**, 73-83.

Shirey S. B. and Walker R. J. (1998) The Re-Os isotope system in
cosmochemistry and high-temperature geochemistry. *Annu. Rev. Earth
Planet. Sci.* **26**, 423-500.

1044 Sims K. W. W., Newsom H. E. and Gladney E. S. (1990) Chemical
 1045 fractionation during formation of the Earth's core and continental crust:
 1046 clues from As, Sb, W, and Mo, in: Newsom, H.E., Jones, J.H., Newsom,
 1047 J.H. (Eds.), *Origin of the Earth*. Oxford Univ. Press, pp. 291-317.

1048 Stein H. J., Markey R. J., Morgan J. W., Du A. and Sun Y. (1997) Highly
 1049 precise and accurate Re-Os ages for molybdenite from the East Qinling
 1050 molybdenum belt, Shaanxi Province, China. *Econ. Geol.* **92**, 827-835.

1051 Tang M., Chen K. and Rudnick R. L. (2016) Archean upper crust transition
 1052 from mafic to felsic marks the onset of plate tectonics. *Science* **351**,
 1053 372-375.

1054 Taylor S. R., McLennan S. M. and McCulloch M. T. (1983) Geochemistry of
 1055 loess, continental crustal composition and crustal model ages. *Geochim.*
 1056 *Cosmochim. Acta* **47**, 1897-1905.

1057 Taylor S. R. and McLennan S. M. (1985) *The continental crust: its composition*
 1058 *and evolution*. Blackwell Scientific, Oxford.

1059 Taylor S. R. and McLennan S. (2008) *Planetary Crusts: Their Composition,*
 1060 *Origin and Evolution*. Cambridge University Press, New York, 378.

1061 Wedepohl K. H. (1995) The composition of the continental crust. *Geochim.*
 1062 *Cosmochim. Acta* **59**, 1217-1232.

1063

1064 **Figure captions**

1065 Fig. 1. Primitive upper mantle (PUM) normalized PGE and Re abundance
 1066 estimates for the UCC from previous studies and this work. Primitive upper
 1067 mantle values are from Becker et al. (2006) (here and elsewhere).

1068

1069 Fig. 2. Doubled normalized PGE and Re concentrations of all 24 diamictite
 1070 composites grouped by depositional ages. Measured concentrations are

double-normalized, with each analysis corrected for quartz dilution by normalizing it to the average UCC_{R&G} Al₂O₃ content (15.4%: Rudnick and Gao, 2003), and then normalizing individual elements to PUM. Osmium concentrations have been corrected for radiogenic ingrowth of ¹⁸⁷Os from ¹⁸⁷Re decay.

Fig. 3. Doubled normalized median PGE and Re concentrations of diamictite composites from each time interval. Osmium concentrations for diamictite composites have been corrected for radiogenic ingrowth of ¹⁸⁷Os from ¹⁸⁷Re decay. Errors quoted at MAD (median absolute deviation).

Fig. 4. Correlations between Ir and Cr (a), Ni (b) for diamictite composites from this study. Black, blue, red and green circles denote Mesoarchean, Paleoproterozoic, Neoproterozoic and Paleozoic diamictite composites, respectively. Symbols with letters near them (D: Duitschland; A: Afrikander) are not incorporated in the regression calculations. The “n” is the sample numbers used in the regression calculation. Equations are obtained using a bootstrapping resample method (see section 5.2.2).

Fig. 5. Correlations between Ir and Os (a), Ru (b) for diamictite composites from this study. The Os concentrations plotted for diamictite composites are common Os (Table 2). Symbols with letters near them (M: Makganyene; A: Afrikander) are not incorporated in the regression calculation. The “n” is the sample numbers used in the regression calculation. See Fig. 4 for symbols.

Fig. 6. Correlations between Pt and Ru (a), and between Pd and Pt (b). Symbols with letters near them (D: Duitschland) are not incorporated in the regression calculations. The “n” is the sample numbers used in the regression

1099 calculation. See Fig. 4 for symbols.

1100

1101 Fig. 7. Correlations between Re and TOC (a) and between Re and Mo (b) in
1102 diamictite composites. Five composites (Timeball Hill, Makganyene, Bolivian,
1103 Dwyka West and Dwyka East) with relatively high TOC contents and Re
1104 concentrations are not used in the Re-Mo regression calculation and are
1105 shown in the inset. See Fig. 4 for symbols.

Table 1

Table 1 Estimates of PGE abundances (ng/g) and Re-Os isotopes for the upper continental crust

References	Shaw et al., 1976	Esser and Turekian, 1993	Wedepohl, 1995 ^a	Schmidt et al., 1997a, b	Gao et al., 1998	Peucker-Ehrenbrink and Jahn, 2001	Park et al., 2012	This study ^b
Os		0.05		0.03 ± 0.02		0.031		0.059 ± 0.016
Ir	0.02	0.05	0.05	0.03 ± 0.02		0.022	0.022	0.036 ± 0.008
Ru			0.1	1.06 ± 0.4		0.21	0.030	0.079 ± 0.026
Pt			0.4	1.5 ± 0.5	1.5	0.51	0.599	0.80 ± 0.22
Pd			0.4	2 ± 0.5	1.46	0.52	0.526	0.80 ± 0.26
Re		0.4				0.198		0.25 ± 0.12
¹⁸⁷ Os/ ¹⁸⁸ Os		1.92				1.40		0.80 ± 0.38
¹⁸⁷ Re/ ¹⁸⁸ Os		48				35		20 ± 12

^a The raw data, analyzed by G. Hartmann using NiS fire assay followed by Te coprecipitation, have not been published.

^b Obtained from regression approach, see section 5.2.2.

Table 2

Table 2 PGE-Re concentrations and Os isotopes in reference materials

	Os	Ir	Ru	Pt	Pd	Re	¹⁸⁷ Os/ ¹⁸⁸ Os
References	ng/g	ng/g	ng/g	ng/g	ng/g	ng/g	
TDB-1, Diabase							
This study, n = 5	0.146 ± 0.016	0.074 ± 0.016	0.212 ± 0.023	5.0 ± 1.1	22.0 ± 2.2	1.05 ± 0.14	0.94 ± 0.16
Peucker-Ehrenbrink et al., 2003, n = 8	0.122 ± 0.013	0.078 ± 0.012		4.40 ± 0.57	24.8 ± 1.8		0.832 ± 0.099
Meisel and Moser, 2004a, b, n = 6	0.117 ± 0.022	0.074 ± 0.020	0.199 ± 0.017	5.00 ± 0.39	24.4 ± 4.1	0.796 ± 0.049	0.92 ± 0.19
Ishikawa et al., 2014, n = 32	0.150 ± 0.070	0.077 ± 0.042	0.236 ± 0.092	4.8 ± 1.3	23.1 ± 3.8	0.99 ± 0.16	0.88 ± 0.33
UB-N, Serpentinite							
This study, n = 1	3.640	3.610	7.040	7.41	6.0	0.200	0.127
Puchtel et al., 2014, n = 5	3.79 ± 0.27	3.55 ± 0.32	6.88 ± 0.37	7.71 ± 0.07	6.10 ± 0.10	0.211 ± 0.010	0.12706 ± 16
Meisel and Moser, 2004b, n = 14	3.85 ± 0.13	3.38 ± 0.22	6.30 ± 0.30	7.42 ± 0.30	6.11 ± 0.18	0.210 ± 0.004	
OKUM, Komatiite							
This study, n = 1	0.749	0.843	4.190	10.70	11.10	0.475	0.269
Savard et al., 2010	0.790	0.943 ± 0.040	4.15 ± 0.08	11.44 ± 0.22	12.2 ± 0.73	0.566 ± 0.040	
Maier et al., 2012	0.780	0.990	4.53	11.44	11.9		

Errors quoted at 2σ.

Table 3

Table 3 PGE concentrations and Re-Os isotopes of diamictite composites.

Sam. Name	Os ^a	2σ _m ^b	Os ^c	2σ _m	Ir	2σ _m	Ru	2σ _m	Pt	2σ _m	Pd	2σ _m	Re	2σ _m	¹⁸⁷ Os/ ¹⁸⁸ Os	2σ _m	¹⁸⁷ Re/ ¹⁸⁸ Os	Cr	Ni	Mo	TOC ^d
Mozaan	0.225	0.001	0.207	0.001	0.137	0.001	0.384	0.002	1.45	0.02	2.03	0.09	0.580	0.006	0.9317	0.0008	13.7	344	141	0.88	0.023
Mozaan*	0.208	0.001	0.190	0.001	0.115	0.001	0.394	0.002	1.85	0.03	2.09	0.10	0.575	0.006	0.9988	0.0007	14.8				
Afrikander	0.186	0.001	0.177	0.001	0.166	0.001	1.89	0.02	7.79	0.41	7.89	1.11	0.278	0.003	0.4548	0.0004	7.51	1132	227	0.47	0.025
Promise	0.315	0.001	0.301	0.001	0.203	0.002	0.635	0.005	1.55	0.02	1.78	0.06	0.448	0.004	0.4753	0.0004	7.15	386	152	0.77	0.053
Coronation	0.155	0.001	0.137	0.001	0.123	0.001	0.342	0.002	2.22	0.04	2.35	0.12	0.589	0.007	1.177	0.001	20.7	294	114	3.0	0.046
Duitschland	0.265	0.002	0.263	0.002	0.194	0.002	0.308	0.002	10.9	0.8	12.2	2.7	0.0791	0.0007	0.2184	0.0003	1.45	164	61.3	0.55	0.130
Timeball Hill	0.0643	0.0001	0.0486	0.0002	0.0399	0.0003	0.0922	0.0005	0.964	0.009	0.993	0.026	0.659	0.006	2.851	0.002	66.9	123	38.7	1.3	0.183
Makganyene	0.236	0.001	0.191	0.001	0.0412	0.0003	0.0881	0.0005	0.781	0.007	0.894	0.025	1.81	0.03	1.921	0.001	45.6	82.4	37.3	1.0	0.174
Ramsay Lake	0.0737	0.0001	0.0537	0.0003	0.0565	0.0003	0.0736	0.0006	2.24	0.04	2.77	0.18	0.786	0.009	4.468	0.002	80.4	74.7	39.7	2.3	0.035
Bruce	0.0699	0.0001	0.0526	0.0002	0.0448	0.0003	0.0931	0.0005	1.20	0.01	1.11	0.03	0.680	0.007	3.348	0.002	66.5	111	32.4	1.5	0.039
Gowganda	0.0583	0.0001	0.0466	0.0001	0.0462	0.0003	0.100	0.001	1.33	0.02	1.49	0.05	0.477	0.004	2.723	0.002	52.8	121	55.9	1.8	0.033
Bottle Creek	0.0338	0.0001	0.0331	0.0001	0.0318	0.0002	0.0676	0.0005	0.719	0.007	0.499	0.008	0.0268	0.0003	2.409	0.002	4.95	96.2	37.8	0.58	0.045
Gaskiers	0.0121	0.0001	0.0116	0.0001	0.0066	0.0001	0.0080	0.0002	0.276	0.002	0.241	0.002	0.0798	0.0007	0.7083	0.0008	34.1	64.2	25.9	1.2	0.035
Pocatello	0.0146	0.0001	0.0146	0.0001	0.0141	0.0001	0.0278	0.0002	0.329	0.004	0.224	0.003	0.0068	0.0001	1.597	0.002	2.66	44.7	21.9	0.48	0.034
Pocatello*	0.0153	0.0001	0.0152	0.0001	0.0187	0.0002	0.0305	0.0004	0.364	0.005	0.220	0.002	0.0068	0.0001	1.494	0.001	2.51				
Konnarock	0.0403	0.0004	0.0402	0.0004	0.0084	0.0001	0.0154	0.0001	0.145	0.002	0.169	0.002	0.0110	0.0001	0.4229	0.0020	1.37	25.6	11.4	0.56	0.050
Nantuo	0.0358	0.0001	0.0352	0.0001	0.0264	0.0002	0.0236	0.0002	0.637	0.015	0.402	0.013	0.0877	0.0004	0.7528	0.0007	12.8	55.7	28.8	0.32	0.080
Gucheng	0.0205	0.0001	0.0199	0.0001	0.0229	0.0002	0.0268	0.0002	0.566	0.009	0.453	0.007	0.0851	0.0004	0.9602	0.0011	22.2	66.0	34.9	0.33	0.150
Blaubeker	0.0549	0.0001	0.0548	0.0001	0.0263	0.0003	0.0417	0.0004	0.586	0.011	0.468	0.008	0.0124	0.0001	1.498	0.001	1.28	39.0	14.1	0.49	0.029
Kaigas	0.132	0.002	0.129	0.002	0.0216	0.0002	0.0336	0.0003	0.553	0.010	0.305	0.004	0.316	0.002	1.500	0.004	13.6	58.9	29.5	1.0	0.063
Chuos	0.0611	0.0002	0.0603	0.0002	0.0336	0.0003	0.0452	0.0003	0.526	0.005	0.576	0.010	0.112	0.001	0.9860	0.0012	9.8	31.0	29.5	0.61	0.111
Numees	0.0162	0.0001	0.0157	0.0001	0.0094	0.0001	0.0192	0.0002	0.257	0.003	0.364	0.006	0.0742	0.0005	3.359	0.005	31.3	29.0	16.0	0.43	0.093
Ghaub	0.0537	0.0002	0.0533	0.0002	0.0245	0.0002	0.0384	0.0003	0.341	0.003	0.323	0.005	0.0516	0.0003	1.143	0.002	5.24	44.9	17.4	0.46	0.090
Bolivian	0.0290	0.0001	0.0171	0.0002	0.0179	0.0002	0.0285	0.0003	0.328	0.004	0.206	0.004	3.43	0.06	3.381	0.003	813	50.0	18.4	0.24	0.250
Dwyka West	0.132	0.001	0.120	0.001	0.0939	0.0007	0.399	0.004	2.74	0.10	2.46	0.10	3.68	0.04	1.825	0.002	165	164	79.9	2.9	0.369
Dwyka East	0.0442	0.0002	0.0421	0.0002	0.0336	0.0003	0.0867	0.0011	0.963	0.021	0.931	0.018	0.665	0.003	1.381	0.002	84.2	95.4	32.5	0.73	0.190

The concentration unit for PGEs and Re is ng/g and for Cr, Ni and Mo is µg/g. ^a Measured Os concentrations.

^b The σ_m (standard deviation of the mean) was obtained by error propogation from the σ_m of measured isotopic ratios (100 cycles).

^c Os concentrations with radiogenic Os removed (common Os). ^d Total organic carbon (wt%) * Replicate The Cr, Ni and Mo data are from Gaschnig et al. (2016).

Table 4

Table 4 Average PGE abundance estimates (ng/g) in the UCC of various time periods.

	Mesoarchean UCC		Paleoproterozoic UCC	Neoproterozoic UCC	Phanerozoic UCC		Present-day UCC	
	Mesoarchean composites, n = 4 ^a	Model calculation ^b	Paleoproterozoic composites, n = 7 ^a	Neoproterozoic composites, n = 10 ^a	Phanerozoic composites, n =2 ^a	Model calculation ^b	Median approach ^c	Regression approach ^d
Os	0.192 ± 0.035 ^e		0.053 ± 0.006 ^e	0.038 ± 0.020 ^e	0.030 ± 0.013 ^e		0.038 ± 0.017	0.059 ± 0.016
Ir	0.152 ± 0.022	0.08-0.16	0.045 ± 0.005	0.022 ± 0.006	0.026 ± 0.008	0.019	0.022 ± 0.006	0.036 ± 0.008
Ru	0.51± 0.15	0.21-0.46	0.092 ± 0.007	0.027 ± 0.010	0.059 ± 0.029	0.030	0.028 ± 0.010	0.079 ± 0.026
Pt	1.89 ± 0.39	1.8-2.9	1.20 ± 0.42	0.43 ± 0.14	0.65 ± 0.32	0.72	0.43 ± 0.14	0.80 ± 0.22
Pd	2.19 ± 0.29	1.8-2.7	1.11 ± 0.38	0.34 ± 0.11	0.57 ± 0.36	0.76	0.34 ± 0.11	0.82 ± 0.26

^a For each time period, the median values (consistent with averages) from diamictites are taken as the average PGE abundances for the UCC in that period. The errors quoted at MAD (median absolute deviation).

^b Calculated using the average PGE concentrations of rocks in Table A5 and the relative proportions of rock types in Condie's (1993) restoration model and the model from Tang et al. (2016) in Table A6. No PGE data for felsic rocks are published and their PGE concentrations are assumed to be zero. The PGE abundances in the Proterozoic UCC are not provided here as few data have been reported for Proterozoic rocks.

^c Obtained from the "Median approach", see section 5.2.1. Errors quoted at MAD.

^d Obtained from the "Regression approach", see section 5.2.2.

^e The Os concentrations used for medians are common Os.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7