Fathering in Rodents: Neurobiological Substrates and Consequences for Offspring

Karen L. Bales^{1,2} and Wendy Saltzman³

- 91. Department of Psychology, University of California, Davis
- 102. California National Primate Research Center
- 113. Department of Biology, University of California, Riverside

Please address correspondence to: Karen L. Bales, Department of Psychology, One Shields Ave.,
University of California, Davis, CA, 95616. Ph: 530-754-5890, Email: klbales@ucdavis.edu

Abstract

Paternal care, though rare among mammals, is routinely displayed by several species of rodents. Here we review the neuroanatomical and hormonal bases of paternal behavior, as well as the behavioral and neuroendocrine consequences of paternal behavior for offspring. Fathering behavior is subserved by many of the same neural substrates which are also involved in maternal behavior (for example, the medial preoptic area of the hypothalamus). While gonadal hormones such as testosterone, estrogen, and progesterone, as well as hypothalamic neuropeptides such as oxytocin and vasopressin, and the pituitary hormone prolactin, are implicated in the activation of paternal behavior, there are significant gaps in our knowledge of their actions, as well as pronounced differences between species. Removal of the father in biparental species has long-lasting effects on behavior, as well as on these same neuroendocrine systems, in offspring. Finally, individual differences in paternal behavior can have similarly long-lasting, if more subtle, effects on offspring behavior. Future studies should examine similar outcome measures in multiple species, including both biparental species and closely related uniparental species. Careful phylogenetic analyses of the neuroendocrine systems presumably important to male parenting, as well as their patterns of gene expression, will also be important in establishing the next generation of hypotheses regarding the regulation of male parenting behavior.

Key Words: paternal care, testosterone, prolactin, vasopressin, oxytocin, alloparenting

Introduction

Male parenting behavior is rare among mammals and displayed mostly by socially monogamous species (Kleiman, 1977; Lukas and Clutton-Brock, 2013). The fact that fathering behavior is displayed by humans (Cabrera and Tamis-LeMonda, 2012), and that human paternal behavior is highly variable, has led to a keen interest in the hormonal and neural substrates of this behavior (Bales et al., 2011b; Saltzman and Ziegler, 2014), as well as its effects on offspring (Braun and Champagne, 2014). Rodent species have been particularly informative in this area, due to the ease of experimentation with them, but also to the relatively high number of biparental species in this order, including prairie voles (*Microtus ochrogaster*), mandarin voles (*Microtus mandarinus*), Mongolian gerbils (*Meriones unguiculatus*), California mice (*Peromyscus californicus*), Djungarian hamsters (*Phodopus campbelli*), and Octodon degus (*Octodon degus*). In this paper, we review the current understanding of the neuroanatomical and hormonal bases of paternal care in rodents. We also summarize what is known about the effects of fathering behavior on offspring, which has mostly been studied by removing fathers from the family group and, more rarely, by examining effects of individual variation in paternal behavior on offspring in intact groups.

Neuroanatomical basis of paternal behavior

The neural circuitry underlying maternal behavior has been studied extensively in the rat and provides a useful starting point for investigating the neuroanatomical basis of male parental care. Maternal behavior in Norway rats (*Rattus norvegicus*) is thought to be regulated largely by two opposing neural systems, both of which are activated in response to output from the main and accessory olfactory systems to the medial nucleus of the amygdala (MeA) (reviewed by (Numan, 2014; Numan and Insel, 2003). In the absence of specific hormonal and neurochemical inputs (estrogen, progesterone, oxytocin), MeA activity leads to activation of the anterior hypothalamus and ventromedial nucleus of the hypothalamus. These regions in turn project to the periaqueductal gray, which promotes aversion to pup stimuli as well as

defensiveness and avoidance, thereby inhibiting the expression of maternal behavior. In contrast, in the hormonal milieu associated with late pregnancy and parturition, stimulation of the MeA leads to activation of the bed nucleus of the stria terminalis (BST) and the medial preoptic area of the hypothalamus (MPOA), stimulating attraction to stimuli (primarily odors) from pups and promoting maternal behavior.

As in females, the brain region most consistently implicated in male parental behavior is the MPOA. Adult male Norway rats do not engage in spontaneous parental behavior but can be induced to behave paternally through continuous exposure to pups, as is also the case for virgin female rats (i.e., sensitization; (Rosenblatt, 1967). Several studies indicate that the MPOA is essential for this process in adult males. For example, Rosenblatt et al. (1996) found that radiofrequency lesions of the MPOA prevented sensitization in adult males, at least for the 13 days of pup exposure over which males were tested. Sturgis and Bridges (Sturgis and Bridges, 1997) used the neurotoxin NMA to lesion the MPOA in castrated adult males treated with estrogen and progesterone. In contrast to radiofrequency lesions, NMA selectively targets cell bodies, sparing fibers of passage. MPOA lesions in this study inhibited the expression of paternal behavior (i.e., retrieving and crouching over pups) in previously sensitized rats. Collectively, these findings indicate that the MPOA is important for the expression of several specific behavioral components of paternal care during both the initiation and maintenance of pup-induced paternal behavior in the adult male rat. In addition, the MPOA appears to be a critical site for estrogenic facilitation of rat paternal behavior (Rosenblatt and Ceus, 1998). Most recently, a series of studies in the uniparental laboratory house mouse found that galaninin-expressing neurons in the MPOA are essential for expression of paternal behavior (Wu et al., 2014b).

Because male rats do not show spontaneous infant care, this species is not a particularly appropriate model for studies of paternal behavior. In fact, Rosenblatt et al. (1996) refer to sensitization in male rats

as maternal behavior, because in this species only females display parental behavior under natural conditions. In this context, it is notable that both biparental California mice and uniparental white-footed mice (*Peromyscus maniculatus*) show increases in Fos-ir in the MPOA in response to pups (Lambert et al., 2013). More recent studies, therefore, have focused on neural and endocrine influences on paternal behavior in biparental species, in which both parents provide infant care under natural conditions (Bales et al., 2011b). In rodents, most of these studies have focused on the biparental prairie vole and California mouse. In these species, as in female rodents and male rats, the MPOA is crucial for the expression of paternal behavior. Lee and Brown (Lee and Brown, 2002, 2007) characterized pup-directed behavior in male California mice that underwent electrolytic lesions of the MPOA three days after the birth of their first litter. Over the subsequent 10 days of testing, lesioned males showed a slower onset of paternal behavior, as well as less time engaging in paternal behavior (retrieving, sniffing, licking, or crouching over pups), less time in proximity to pups, and longer latencies to retrieve pups, compared to sham-lesioned males. Consistent with these findings, several studies of Fos-immunoreactivity (Fos-ir), an index of neuronal activation, have found increased Fos expression in the MPOA following exposure of males to pups. California mouse fathers, but not virgins or males housed with primigravid females (females in their first pregnancy), had elevated Fos-ir in the MPOA following exposure to a foster pup, compared to fathers similarly exposed to a control object (De Jong et al., 2009); but see (De Jong et al., 2010). In prairie voles, virgin males had increased MPOA Fos-ir following exposure to an unrelated pup compared to exposure to a novel object (Kirkpatrick et al., 1994b). (Please note that behavior towards an unrelated pup, also known as alloparenting, is a common behavior in prairie voles, especially in males).

Several studies of biparental rodents provide evidence that the amygdala, in addition to the MPOA, is involved in paternal behavior. In California mice, lesions of the basolateral nucleus of the amygdala (BLA) had effects on parental behavior that were very similar to those of MPOA lesions (Lee and Brown, 2007). Specifically, new, first-time fathers with BLA lesions spent less time engaging in paternal

behavior, licking pups, and in proximity to pups, as well as longer latencies to retrieve pups, compared to sham-lesioned males. In contrast, Kirkpatrick et al. (Kirkpatrick et al., 1994a) found that in young adult male prairie voles housed with ovariectomized females, electrolytic lesions of the BLA had no effect on responses to pups, whereas electrolytic lesions of the corticomedial amygdala or the MeA reduced males' contact time with pups. Immunohistochemical studies have further implicated the amygdala and the BST, considered part of the "extended amygdala" (Davis et al., 2010), in paternal behavior. Virgin male prairie voles exposed to a pup had higher Fos-ir in the medial amygdala (MeA) and medial BST, compared to males exposed to a control object (Kirkpatrick et al., 1994b), while in California mice exposed to a foster pup, fathers had significantly higher Fos-ir in both the medioventral and medial posteromedial amygdala, compared to virgin males (De Jong et al., 2009).

Not surprisingly, the olfactory bulbs also appear to play a critical role in rodent paternal behavior. This region has received surprisingly little attention; however, bilaterally bulbectomized, virgin adult male prairie voles were significantly more likely to attack pups than were sham-lesioned males (Kirkpatrick et al., 1994c), suggesting that olfactory cues normally inhibit pup-directed aggression. In the same species, exposure to a pup elevated Fos-ir in the accessory olfactory bulbs, compared with exposure to a control object (Kirkpatrick et al., 1994b).

Immunohistochemical studies have identified several additional brain regions that might be associated with paternal care in biparental rodents. In California mouse fathers, exposure to a pup increased Fos-ir in the caudal dorsal raphe nucleus and the lateral habenula (De Jong et al., 2009; De Jong et al., 2010), whereas in virgin male prairie voles, pup exposure elevated Fos-ir in the lateral septum, paraventricular nucleus of the thalamus, and nucleus reuniens of the thalamus (Kirkpatrick et al., 1994b). Virgin male California mice also exhibited elevated Fos-ir in the lateral septum when compared to either pup-exposed virgins or fathers (Lambert et al., 2011). The roles of these regions in paternal care, if any, are unknown.

138

139 In summary, findings from immunohistochemical and lesion studies suggest that paternal care in
140 biparental rodents is associated with some of the same brain regions implicated in maternal care,
141 including the olfactory bulbs, MeA, BST, and MPOA. The precise roles of these and other brain regions
142 in both the initiation and maintenance of paternal behavior are not yet known, however, and the
143 generalizability of these findings are not clear, as they come from only two cricetid rodents.

144

145 Intriguingly, recent studies have found that fatherhood influences neural plasticity in several rodent
146 species. Lieberwirth and colleagues (Lieberwirth et al., 2013) used the cell-division marker
147 bromodeoxyuridine (BrdU) to investigate neurogenesis in male prairie voles, and found that fatherhood
148 reduced the survival of new cells in the amygdala, dentate gyrus, and hypothalamus, but not the main
149 olfactory bulbs. In the California mouse, Glasper and colleagues (Glasper et al., 2011) found that
150 fatherhood inhibited hippocampal neurogenesis in the California mouse. Lambert et al. (2011) found that
151 biparental California mice had higher levels of nestin-ir in CA2 and CA3, while uniparental white-footed
152 mice had higher glial fibrillary acid protein in dentate gyrus and hippocampal fissure (Lambert et al.,
153 2011). They interpreted this as indicating higher neuroplasticity in the biparental species and higher glial
154 plasticity in the uniparental species. Finally, a fascinating series of studies in the uniparental house mouse
155 (*Mus*) indicated that interactions of adult male mice with their own pups stimulated neurogenesis in the
156 father's subventricular zone and dentate gyrus under the influence of prolactin signaling (Mak et al.,
157 2013; Mak and Weiss, 2010). Some of the new cells matured into olfactory interneurons in the olfactory
158 bulb, where they responded preferentially to offspring odors and appeared to subserve later recognition of
159 mature offspring. These results highlight potential differences in the role of pup cues in neurogenesis in
160 males of different species, and may highlight differences between males of biparental and uniparental
161 species; however, additional studies will be necessary to tease apart these differences.

162

Hormonal influences on paternal behavior

Neuroendocrine regulation of parental behavior has been studied extensively in female rodents, mostly rats. In new mothers, maternal responsiveness toward pups is activated under the hormonal conditions of late pregnancy, parturition, and lactation. Declining progesterone levels, as well as high levels of estrogen, oxytocin, and prolactin, are especially important in this process (Numan and Insel, 2003). These same hormones and neuropeptides are implicated in the modulation of paternal behavior in male rodents.

As described above, adult male rats do not display paternal behavior spontaneously; however, they can be induced to behave paternally by continuous exposure to pups (Rosenblatt, 1967). The onset of this pup-induced paternal behavior can be influenced by exposure of males to specific hormones both in the early postnatal period and in adulthood. Most importantly, gonadal steroids act early in life to organize paternal behavior: castration during the early postnatal period increases males' paternal responsiveness in adulthood, whereas early androgen treatment reverses this effect (McCullough et al., 1974; Rosenberg and Herrenkohl, 1976). In adult males, display of pup-induced parental behavior is not dependent on activational effects of hormones (Rosenblatt, 1967); nonetheless, both sensitization and infanticide can be influenced by gonadal steroids in adulthood. In males castrated as adults, treatment with high doses of estrogen, following priming with estrogen and progesterone, decreases the latency to the onset of paternal behavior during sensitization, whereas testosterone treatment increases the likelihood of infanticide (Lubin et al., 1972; Rosenberg, 1974; Rosenblatt et al., 1996). The onset of pup-induced paternal behavior can also be facilitated by treatment with prolactin (Sakaguchi et al., 1996).

As mentioned above, findings from rats and other uniparental rodents may not be particularly informative with respect to influences on spontaneously occurring paternal care in biparental species. Accordingly,

recent studies of hormonal and neuropeptide consequences of fatherhood and influences on paternal care have focused on species in which fathers routinely provide care to their offspring under natural conditions (reviewed by (Bales et al., 2011b; Saltzman and Ziegler, 2014).

Developmental effects of hormones

Studies of developmental influences on paternal behavior in rodents have focused on the prairie vole and yield a complex picture of early organizational effects of hormones. First, treatment of pregnant dams with either testosterone propionate (TP), corticosterone, the androgen receptor antagonist flutamide (FTD), or the aromatase inhibitor 1,4,6-androstatriene-3,17-dione (ATD) did not alter behavioral responses of male offspring to unrelated pups during either the juvenile period or adulthood (Lonstein and De Vries, 2000; Lonstein et al., 2002; Roberts et al., 1996). Similarly, treatment of pups with TP, FTD or ATD during the first week after birth had minimal effects on males' responses to pups in adulthood (Lonstein and De Vries, 2000). On the other hand, paternal behavior during the juvenile period or adulthood was reduced markedly by either castration immediately after birth (Lonstein et al., 2002) or treatment with ATD or FTD during the second postnatal week (Kramer et al., 2009). Together, these findings suggest that early-life exposure to both androgens and estrogens is critical for the later expression of paternal behavior in male prairie voles, and that the sensitive period for these actions of gonadal steroids is during the second week after birth, rather than earlier.

Paternal behavior in prairie voles can also be modulated by early exposure to neuropeptides. Male pups that received a single injection of an oxytocin (OT) receptor antagonist ([d(CH₂)⁵, Tyr(Me)², Orn⁸]-Vasotocin) on the first postnatal day exhibited reduced parental behavior and increased aggression toward unrelated pups during the juvenile period, but no effects were seen in adulthood (Bales et al., 2004b). In the same study, however, a single injection of neonates with exogenous OT had no discernible effects on

males' later paternal or aggressive responses to pups (Bales et al., 2004b). This dosage of OT given neonatally did alter other outcome measures, including pair-bond formation (Bales and Carter, 2003), and the neonatal rodent blood-brain barrier is not fully formed on day 1 (Vorbodt, 1993), suggesting that neonatal treatment with OT can potentially alter behavior via central actions. However, it is also possible that these effects (and the intranasal ones described below), could be due to peripheral binding of OT, and/or binding of OT to V1a receptors (Gimpl and Fahrenholz, 2001). In fact, daily intranasal treatment of juvenile males with OT did not alter their alloparental responsiveness to pups in early adulthood, compared to controls (Bales et al., 2013), whereas it did alter pair-bonding behavior. The failure of exogenous OT to enhance alloparental responsiveness in these two studies is likely due to the high baseline levels of alloparental behavior in young male prairie voles, which might obscure any potential stimulatory effects of OT treatment.

Activational effects of hormones

In many biparental rodents, fathers undergo systematic changes in central and/or peripheral concentrations of several hormones and neuropeptides in association with reproductive condition. In some cases, changes in central expression of hormone or neuropeptide receptors have also been described. While some of these changes in hormone and neuropeptide signaling appear to facilitate the expression of paternal behavior, most are of unknown significance (Bales et al., 2011b; Saltzman and Ziegler, 2014). Below we review profiles of several hormones (prolactin, gonadal steroids, glucocorticoids) and neuropeptides (vasopressin [AVP], OT) associated with fatherhood, as well as the evidence that these profiles may play a causal role in the activation of paternal care.

Prolactin

Effects of fatherhood

In several biparental rodents, including Mongolian gerbils (Brown et al., 1995), California mice (Gubernick and Nelson, 1989), and Djungarian hamsters (Reburn and Wynne-Edwards, 1999), circulating prolactin levels are significantly higher in fathers living with their mate and pups than in virgin males, newly mated males, and/or expectant fathers (see also (Schradin, 2008) on striped mice, *Rhabdomys pumilio*). Male Djungarian hamsters also have increased prolactin receptor mRNA transcript levels in the choroid plexus of the hypothalamus during their mate's early postpartum period, indicating that prolactin signaling in the brain is elevated during periods when males are interacting with pups (Ma et al., 2005) .

Effects on paternal behavior

In contrast to the numerous studies demonstrating positive correlations between circulating prolactin levels and paternal behavior, no evidence is available, to our knowledge, to support a causal role of prolactin in the onset or maintenance of paternal care in biparental rodents (Wynne-Edwards and Timonin, 2007). In the only study that experimentally tested the effect of prolactin on paternal behavior in biparental rodents, Brooks et al. (Brooks et al., 2005) treated first-time Djungarian hamster fathers with the dopamine agonists bromocriptine and cabergoline in order to suppress secretion of prolactin. Although this treatment markedly reduced plasma prolactin concentrations, no differences in paternal behavior (pup-retrieval) in fathers, or growth and survival of pups, were found between agonist-treated and control males.

Gonadal steroids

Effects of fatherhood

Testosterone generally shows a negative relationship with paternal care. In some biparental rodents, testosterone levels of expectant fathers rise across the mate's pregnancy (e.g., Djungarian hamster: (Reburn and Wynne-Edwards, 1999) and decline following the birth of their pups (e.g., Mongolian gerbil:

((Brown et al., 1995); Djungarian hamster: (Reburn and Wynne-Edwards, 1999); California mouse: (Trainor et al., 2003); (Gubernick and Nelson, 1989).

Few studies have characterized changes in estrogen signaling in rodent fathers, and in the small number of species that have been studied, no consistent picture has emerged. In male Djungarian hamsters, serum estradiol levels do not appear to change with paternal status (Schum and Wynne-Edwards, 2005), and estrogen receptor alpha (ER α)-ir in several brain regions (MPOA, BST, and MeA) does not differ between fathers and non-fathers (Timonin ME et al., 2008). Similarly, a recent study comparing California mouse fathers and virgin males found no differences in mRNA expression for ER α in the MPOA, BST, or MeA (Perea-Rodriguez et. al., in press). In contrast, biparental mandarin vole (*M. mandarinus*) fathers have lower levels of ER α -ir in the MPOA and BST, and higher levels in the ventromedial hypothalamus, compared to non-fathers (Song et al., 2010). Virgin male mandarin voles that display high levels of paternal care had more ER α -ir in the MPOA, BST, the arcuate nucleus, and the MeA (Li et al., 2015).

Findings on progesterone are sparse and inconsistent. Male Djungarian hamsters show a significant increase in circulating progesterone levels at the end of the mate's pregnancy (Schum and Wynne-Edwards, 2005), whereas in California mice, circulating progesterone levels (Trainor et al., 2003) and expression of mRNA for progesterone receptors in the BST (Perea-Rodriguez et al., in press) are significantly lower in fathers than in virgin males (Trainor et al., 2003).

Effects on paternal behavior

Although circulating or excreted testosterone levels correlate negatively with measures of paternal behavior in several rodents (reviewed by (Numan and Insel, 2003; Wynne-Edwards and Timonin, 2007),

experimental studies indicate that effects of testosterone on paternal care differ markedly across species. In the Djungarian hamster (Hume and Wynne-Edwards, 2005) and in one study of prairie voles (Lonstein and De Vries, 1999), for example, castration of adult males had minimal effects on paternal behavior, whereas castrated male Mongolian gerbils engaged in significantly more paternal behavior than either castrated, testosterone-treated males or sham-castrated males (Clark and Galef, 1999). The opposite pattern has been found in another study of prairie voles (Wang and De Vries, 1993) and in California mice (Trainor and Marler, 2001), in which castration reduced and testosterone treatment restored paternal behavior. In the California mouse, this effect is mediated by aromatization of testosterone to estrogen within the brain: paternal behavior of castrated, reproductively experienced males was restored by treatment with testosterone or estrogen, but not by treatment with the non-aromatizable androgen dihydrotestosterone (Trainor and Marler, 2002). Moreover, fathers had higher aromatase activity within the MPOA than non-fathers (Trainor et al., 2003). Therefore, even though California mouse fathers have lower circulating concentrations of testosterone than non-fathers, they might have higher local concentrations of estrogen within the MPOA and possibly other brain regions.

The MeA appears to be an important site for estrogenic modulation of male parental behavior. Cushing et al. (Cushing et al., 2008) used a viral vector to increase ER α expression in the MeA of adult male prairie voles. Subsequently, these males showed reduced rates of paternal behavior toward an unrelated pup, as compared to controls. Using the same methods, Lei and colleagues (Lei et al., 2010) found no effect of increased ER α expression in the BST on behavioral responses to a pup in adult male prairie voles. These findings suggest that estrogen has site-specific effects within the neural circuitry underlying paternal behavior.

In the uniparental house mouse (*Mus* spp.), progesterone signaling has been shown to promote infanticide and inhibit paternal behavior in adult males (Lightman et al., 2001; Schneider et al., 2003). Progesterone

receptor knockout mice, as well as mice treated with the progesterone receptor antagonist RU486, showed markedly reduced aggression toward pups and enhanced paternal behavior, whereas progesterone treatment of wild-type males significantly increased aggression toward pups. To our knowledge, however, effects of progesterone on paternal care have not been tested in biparental mammals.

Glucocorticoids

Effects of fatherhood

Studies of basal glucocorticoid (corticosterone and cortisol) concentrations in biparental rodents typically find no differences between fathers and non-breeding males (e.g., California mouse: (Chauke et al., 2011; Harris and Saltzman, 2013), prairie vole: (Campbell et al., 2009)). In the Djungarian hamster, however, males had significantly higher serum cortisol levels prior to pair formation than during their mate's pregnancy, and higher levels during the mate's late pregnancy than during mid-pregnancy and the early lactation period (Reburn and Wynne-Edwards, 1999).

Effects on paternal behavior

Very few studies have examined effects of glucocorticoids on paternal care. In California mouse fathers, acute treatment with high doses of corticosterone had neither short-term effects on paternal behavior nor longer-term effects on pup survival or development (Harris et al., 2011). To determine effects of chronic stress, including chronic elevations of circulating corticosterone levels, on paternal care, Harris et al. (Harris et al., 2013) subjected California mouse fathers to a chronic variable stress paradigm for 7 days. Stressed fathers showed both significant elevations in plasma corticosterone levels and subtle reductions in their interactions with their mate and pups compared to control fathers, but no differences were detected in pup survival or development. In male prairie voles, a swim stressor, which reliably doubles corticosterone concentrations in this species (Taymans et al., 1997), led to increased huddling over a

strange pup. These effects were not seen in females (Bales et al., 2006), although long-term effects of the stressor on fathers or offspring were not studied. These findings suggest that neither acute nor chronic glucocorticoid elevations have pronounced effects on parental care in males, but clearly, additional data are needed from other species.

Vasopressin

Effects of fatherhood

AVP in fathers has been studied most thoroughly in the prairie vole, with a strong emphasis on AVP signaling within the brain. In this species, new fathers have reduced densities of AVP-immunoreactive fibers in the lateral septum and lateral habenular nucleus compared to both sexually naïve males and males housed with late-pregnant mates. This finding might reflect increased synthesis and release of AVP from the BST and MeA, the likely source of the vasopressinergic fibers, following both copulation and parturition (Bamshad et al., 1994; Bamshad et al., 1993). Prairie vole fathers (as well as mothers) also exhibit elevated AVP mRNA levels in the hypothalamic paraventricular nucleus (PVN) and supraoptic nucleus (SON) in the postpartum period compared to sexually naïve controls (Wang et al., 2000). In the California mouse, in contrast, AVP mRNA levels in the PVN do not differ between fathers and either virgin males or males housed with tubally ligated females (De Jong et al., 2013), but levels of mRNA for AVP V1a receptors in the BST are significantly lower in fathers than in virgin males (Perea-Rodriguez et al., in press). In virgin male prairie voles, exposure to a pup increased activation (as determined by Fos-ir expression) of AVP-immunoreactive cells in the PVN, but not plasma AVP (Kenkel et al., 2012). Effects of fatherhood on AVP signaling therefore appear to differ across species and might be clarified by additional studies.

Effects on paternal behavior

Both within and among species, paternal behavior correlates with patterns of AVP -ir and AVP binding, especially in the lateral septum and other parts of the extended amygdala (Bester-Meredith et al., 1999; Insel and Shapiro, 1992; Parker et al., 2001). Moreover, central infusion of AVP or AVP receptor antagonists promotes or inhibits paternal behavior, respectively, in the biparental prairie vole (Wang et al., 1994) and in the facultatively biparental meadow vole (*M. pennsylvanicus*: (Parker and Lee, 2001). Nonetheless, castration of male prairie voles virtually eliminates AVP-ir in the lateral septum and lateral habenula but does not affect paternal behavior, suggesting that AVP signaling in these areas is not essential for the expression of paternal care (Lonstein and De Vries, 1999).

Oxytocin

Effects of fatherhood

Several studies have investigated changes in intracerebral OT signaling or peripheral OT concentrations in rodent fathers. In the facultatively biparental meadow vole (*Microtus pennsylvanicus*), sexually and paternally experienced, paternally behaving males were found to have significantly higher OT receptor binding in several brain regions (accessory olfactory nucleus, lateral septum, BST, and lateral amygdala) than sexually and paternally inexperienced, non-paternally behaving males; however, it could not be determined whether these neural differences were caused by differences in sexual activity, cohabitation with a female, and/or paternal experience (Parker et al., 2001). Similarly, fathers have increased numbers of OT-immunoreactive fibers in the PVN and SON in mandarin voles, compared to virgin males; however, similar effects were seen in non-fathers that were either exposed briefly to pups or housed with a female prior to parturition (Song et al., 2010). Virgin mandarin vole males that engaged in more paternal behavior had elevated OT-immunoreactive fibers in the PVN and SON as well (Li et al., 2015).

In male prairie voles, fathers, compared to virgins, were recently found to have more OT-immunoreactive neurons in the PVN, as well as higher densities of OT-immunoreactive fibers in the nucleus ambiguus and nucleus of the solitary tract, but fewer OT-immunoreactive neurons in the BST (Kenkel et al., 2014). Another study on prairie voles, however, found no differences between fathers and virgin males in OT gene expression in either the PVN or the SON (Wang et al., 2000), or in OT receptor binding in the lateral septum, BST, or the ventromedial nucleus of the hypothalamus (Wang et al., 2000). Finally, California mouse fathers have reduced expression of mRNA for the OT receptor in the BST, but not in the MPOA or medial amygdala, compared to virgin males (Perea-Rodriguez et. al., in press). Thus, effects of fatherhood on the brain's OT system seem to be inconsistent across species, and where effects have been detected, they may correspond to cohabitation with a female rather than fatherhood *per se*.

In addition to central OT signaling, circulating OT concentrations may be altered by fatherhood and interactions with pups. Male California mice exhibit elevated circulating OT concentrations during the first half of their mate's pregnancy, but OT levels decline prior to parturition and remain low throughout the postpartum period (Gubernick et al., 1995). Moreover, in virgin male prairie voles (which had not been exposed to a female), pups elicited a rise in plasma OT levels as well as an increase in OT-ir staining in the PVN (Kenkel et al., 2012), suggesting that pups present a potent stimulus to the OT system. It is unclear, however, whether circulating OT concentrations reliably reflect levels in the brain (Churchland and Winkielman, 2012).

Effects on paternal behavior

In virgin male prairie voles, paternal behavior was inhibited by combined treatment with an AVP receptor antagonist and an OT receptor antagonist, but not either antagonist alone (Bales et al., 2004a). No other

published studies, to our knowledge, have manipulated OT signaling in adult male rodents to identify possible effects on paternal behavior.

Effects of Male Parenting on Offspring

Paternal deprivation

Paternal deprivation, or paradigms where the father in a biparental species is removed during offspring development, has often been used to explore the role of the father in offspring development and behavior. These studies have revealed extensive, often sex-specific effects on offspring, including effects on survival, especially under adverse conditions (Cantoni and Brown, 1997; Gubernick et al., 1993; Wright and Brown, 2002), developmental markers (Elwood and Broom, 1978; Piovanotti and Vieira, 2004; Wang and Novak, 1992), and behavior (Ahern et al., 2010; Ahern and Young, 2009; Bredy et al., 2004).

Behavioral effects of father absence are particularly intriguing and span cognitive, emotional, and reproductive behavior (Table 3). In California mice removal of the father interacted with experimenter handling to predict offspring performance in cognitive tasks such as the Barnes maze, a test of spatial learning (Bredy et al., 2004). Non-handled, father-absent offspring received the least licking/grooming and performed worst in the Barnes maze, although this effect was primarily in male offspring, and father absence had little effect on top of the handling effect. In prairie voles, female offspring that had been raised without fathers displayed less alloparenting (Ahern and Young, 2009), did not form normal pair-bonds (Ahern et al., 2010; Ahern and Young, 2009), and exhibited less licking and grooming of their own offspring when later raising them without a male pair-mate (Ahern et al., 2010), compared to female offspring that had been raised by both parents. Male offspring raised without a father also displayed deficits in pair-bonding (Ahern and Young, 2009).

A growing body of evidence shows systematic changes in behavior caused by paternal deprivation in mandarin voles. Paternally deprived pups of both sexes displayed higher levels of anxiety in an open field and less time in social behavior in a social interaction test, compared to pups raised by both parents (Jia et al., 2009). Social recognition in a habituation-dishabituation paradigm was also impaired in both sexes of paternally deprived offspring (Cao et al., 2014). Paternally deprived male voles displayed less play behavior (boxing and wrestling) on PND 35, and higher contact with littermates (Wang et al., 2012). Finally, paternal deprivation significantly inhibited partner preference formation in females and increased aggression towards their pair-mate and towards an unfamiliar male (Yu et al., 2012). In males, paternal deprivation reduced contact with both the partner and the stranger, as well as increasing attacks towards the strange female (Yu et al., 2012). The overall picture of paternally deprived mandarin voles is broadly consistent with that of paternally deprived prairie voles, with offspring being more anxious and less social on multiple measures (although in a socially monogamous species, aggression towards an outside animal can be interpreted as mate-guarding, an essential aspect of the pair-bond).

Recent studies in voles (mandarin and prairie) and *Octodon degus* have also examined the neuroendocrine consequences of paternal deprivation for offspring (Table 3). Female prairie vole pups in the study cited above (Ahern and Young, 2009), when raised in paternally deprived groups, showed increased numbers of clusters of OT mRNA in the PVN compared to females raised in biparental groups. In addition, both sexes had higher density of receptor binding of corticotropin-releasing factor type 2 receptors (CRFR2) in the dorsal raphe; these two measures (OT mRNA and CRFR2 in the dorsal raphe) correlated in females, but not in males.

Similarly, in mandarin voles, paternal deprivation resulted in sex-specific changes in neuroendocrine systems. In females, paternal deprivation resulted in decreases in glucocorticoid receptors (GR) and brain-derived neurotrophic factor (BDNF) in the dentate gyrus (Wu et al., 2014a), increased serum levels of adrenocorticotrophin (ACTH) (Wu et al., 2014a), reduced levels of dopamine type 1 (D1) and type 2

(D2) receptors in the nucleus accumbens (Nacc) (Yu et al., 2012), reduced serum OT, and reduced ER α mRNA in the MeA, with no change in OTR mRNA in the MeA but a reduction in OTR mRNA in the Nacc (Cao et al., 2014). Corticosterone in paternally deprived females was higher than in pair-reared females in one study (Wu et al., 2014a) and lower in another (Yu et al., 2012).

In contrast, male mandarin voles did not display changes in GR or BDNF in dentate gyrus, or changes in ACTH (Wu et al., 2014a), and had enhanced expression of D1 and D2 in the NAcc (Yu et al., 2012), increased AVP-ir in the PVN, reduced AVP-ir in the anterior hypothalamus and reduced OT-ir in the PVN (Wang et al., 2012). Further, paternally deprived males showed no change in serum OT or in ER α in the MeA, and reductions of OTR in both the MeA and NAcc (Cao et al., 2014). Corticosterone in paternally deprived males did not change in one study (Wu et al., 2014a) and increased in two others (Wang et al., 2012; Yu et al., 2012).

Both sexes of paternally deprived mandarin voles displayed decreased levels of GR and BDNF in CA1 and CA 2/3 of the hippocampus (Wu et al., 2014a). Timelines of corticosterone and ACTH levels throughout early development (Wang et al., 2014) suggest that paternal deprivation alters development of these systems, including disruption of the stress-hyporesponsive period. As a whole, father absence in this socially monogamous species results in an anxious, less social phenotype in both sexes, with sex differences in the underlying neurobiology generally suggesting lower responsiveness of the OT system, altered AVP systems particularly in males, sex-specific changes in the reward system, and conflicting effects on corticosterone.

Octodon degus are an interesting comparison species in that they are from a different rodent lineage than the vole species and California mice, are not socially monogamous, but do often display care by males (Ebensperger et al., 2010). While research on paternal deprivation in this species has mostly focused on the cerebral cortex, some studies have included limbic areas as well. In male degus, numbers of CRF-

positive neurons decreased globally over development; however, paternal deprivation led to even greater reductions in numbers of CRF-positive pyramidal cells in CA1 at PND21 and 90, reduced cells in the dentate gyrus at day 21 but not 90, increased cells in the BLA at day 21 (Seidel et al., 2011), decreased cells in the medial BST at day 21, and no change in the PVN (Gos et al., 2014). Tyrosine hydroxylase fibers, indicating catecholaminergic innervations, were elevated at PND21 in many areas in paternally deprived male degus, including the NAcc, CeA, and CA1. These elevations remained in adulthood in the hippocampus, but the effect was reversed in the NAcc (Braun et al., 2013). Paternal deprivation in degus also results in widespread developmental changes in interneuron expression patterns (Braun et al., 2011), reduced dendritic spine numbers in the somatosensory neurons (Pinkernelle et al., 2009), as well as in the neurons of the anterior cingulate cortex (Otscharoff et al., 2006).

In summary, the absence of the father in biparental species has profound effects on the offspring. The behavioral effects seem to be relatively consistent in that, when they occur, they result in less social and more anxious animals. These effects can be sex- and species-specific. Consistent in the underlying neurobiology are changes in stress-responsive systems including catecholaminergic systems. However, different outcomes of paternal deprivation have been studied in different species; it is thus not always easy to make direct comparisons between species.

Effects of individual differences in paternal behavior on offspring

Less common than studies of paternal deprivation are those in which the role of the father is explored in intact groups. These effects are often more subtle than those described above for paternal deprivation, but suggest that male care affects pup development in a species- and sex-specific manner.

In California mice, retrievals by fathers predict aggression in male pups (Frazier et al., 2007), which appears to be subserved by AVP-immunoreactivity in the ventral BST (Frazier et al., 2007). Retrievals also lead to increased testosterone levels in prepubertal pups 45 minutes post-retrieval (Becker et al.,

2010). Castrated males lick and groom young at lower rates than intact males. Male pups raised by castrated males also lick and groom at lower rates, however they also retrieve pups more frequently, which their fathers did not (Gleason and Marler, 2013). Male California mouse pups cross-fostered to white-footed mouse (*Peromyscus leucopus*) parents were retrieved less frequently as pups, and then retrieved their own pups less frequently as adults (Bester-Meredith and Marler, 2003).

In prairie voles, fathers displayed apparent compensation for mothers that spent low amounts of time in contact with offspring (Perkeybile et al., 2013), with offspring in which mothers displayed high contact receiving less fathering, and offspring that received low contact from mothers receiving more fathering. Low-contact offspring developed at a higher rate, but were less social with a strange juvenile, displayed more autogrooming, and showed more retrievals but less contact with a strange pup compared to high-contact offspring. They also showed greater potentiation of an acoustic startle response. However, it is unclear whether these changes can be attributed specifically to changes in paternal care. When parental care was analyzed by principal components analysis and a “paternal care factor” identified, correlations between this factor and offspring behavioral outcomes did not survive statistical correction.

In another series of studies, prairie vole pups underwent a handling manipulation that had long-lasting effects on social and anxiety-like behavior (Bales et al., 2011a; Bales et al., 2007). Specifically, adults that experienced reduced handling as pups had F1 offspring that displayed deficits in social behavior (alloparenting and pair-bonding). The F2 generation continued to display these deficits if either their mother or father had come from a reduced-handling litter (Stone and Bales, 2010), thus suggesting that in this case, maternal and paternal experience were interchangeable in their effects on the offspring.

Alloparenting experience in cooperatively breeding species can potentially have long-lasting effects on the alloparent’s behavior (Greenberg et al., 2012), neuroendocrine systems such as BDNF (Greenberg et al., 2012), and reproductive success (Tardif et al., 1984); as well as lasting effects on the offspring of

experienced alloparents. Stone and colleagues (Stone et al., 2010) paired male and female prairie voles with varying levels of alloparenting experience, and looked at offspring outcomes. Offspring of parents that had two litters of alloparental experience (either through two litters of experience for one parent, or one litter of experience for each parent) developed faster and displayed more male alloparenting themselves. This effect did not depend on which parent had the alloparenting experience. Another aspect of paternal experience that might affect offspring development is previous experience as a father (as opposed to alloparenting experience). Offspring of experienced Mongolian gerbil fathers locomote more outside the nest than offspring of inexperienced fathers (Piovanotti and Vieira, 2004).

Discussion

We have reviewed what is known regarding the neurobiological substrates of male parenting behavior in rodents. This has provided a clear view of the multitude of species differences in the biological regulation and the consequences of male parenting behavior. Male parenting is typically displayed in socially monogamous or cooperatively breeding species, and as such it is clear that it has evolved many different times. Recent phylogenetic analyses of social monogamy, biparental care, and infanticide have debated the order in which these traits evolved (Dixson, 2013; Lukas and Clutton-Brock, 2013; Lukas and Huchard, 2014; Opie et al., 2013a, b, 2014); however, mostly they agree that biparental care is a consequence rather than a cause of social monogamy. If biparental care evolved independently so many times, why would we expect mechanisms to be similar across species? It is possible that convergent evolution has repeatedly co-opted the same neural and hormonal systems. One recent study, while on a completely different topic, is enlightening in this respect (Parker et al., 2013). Parker and colleagues identified convergence in genes for echolocation in 200 loci in bats and the rather distantly related bottlenose dolphin. A similar study on biparental care in rodents, or mammals in general, would be highly informative as to the underlying evolution and mechanisms of male care.

The work that we review here thus suggests that while fathering is clearly regulated by neuroendocrine systems, our current state of knowledge is insufficient. We need more studies that use similar outcome measures and similar methodologies, carried out across multiple species. For instance, the effects of paternal deprivation have been studied in depth in both mandarin voles and degus, but the focus has been on very different brain areas and outcomes. One great advantage of many biparental rodent species is that they come from taxa that display differing levels of male parenting at either the species level (i.e. *Peromyscus californicus* vs. *Peromyscus maniculatus*), or even within populations of the same species (i.e. strongly biparental and monogamous Illinois prairie voles vs. mostly uniparental Kansas prairie voles (Roberts et al., 1998)). Continued use of closely related species that are either obligately biparental, facultatively biparental, or uniparental will be highly informative.

Other areas in the literature that need to be fleshed out include the distinctions between fathering in obligate and facultative species, and between the neurobiology of parental care in parents and alloparents. Literature on fathering in “facultatively paternal” species such as rats and mice has greatly informed our knowledge of neural substrates (Rosenblatt, 1967; Rosenblatt and Ceus, 1998), especially with regard to induced hormonal changes and to communication between the mother and father in these species (Liang et al., 2014; Liu et al., 2013). Similarly, work with non-fathers in “obligately paternal” species such as prairie voles can inform our understanding of what may *not* be necessary for the display of care. However, we do need to carefully distinguish these different categories in the literature, particularly in respect to their natural occurrence under field conditions rather than artificial conditions of captivity.

Future studies might also focus on what is “unique” to paternal care vs. other types of care. For instance, studies of paternal deprivation have provided ample evidence for long-term consequences on offspring development. However, we were unable to find any studies in which a father was removed but the total number of caregivers remained the same (for example, by leaving an adult sibling in the cage with the mother to help her provide care). As such, the quantity of caregiving, in contrast to (potentially) the

quality, varied in all of the studies cited here, and we are unable to make statements about whether consequences for offspring are the results specifically of father absence or simply less care. This provides an important avenue for future studies.

Finally, relationships between parents, for example the effects of the mother's behavior on the father and vice versa, have rarely been studied in biparental rodents and provide an important area in which to consider the family as a system. Exceptions include a recent, extensive study of parental behavior in California mice (Rosenfeld et al., 2013), as well as the prairie vole study in which principal components analysis was used to examine the latent variables underlying the expression of parental behavior (Perkeybile et al., 2013). However, neither of these studies examined the effects of one parent on the other in a manner that measured bidirectional statistical dependence over time (Butler and Randall, 2013). Dyadic statistical techniques used to study relationships in humans could be useful in characterizing the interdependent nature of the parents' behaviors. This understanding could be crucial to illuminating variation in the behavior of each parent and consequences for offspring development.

Acknowledgements

During the writing of this article, KLB was supported by HD071998, HD053555, OD P51OD01107 to the California National Primate Research Center, and the Good Nature Institute. WS's research on paternal care has been supported by NIH grants R21MH087806 and R21HD075021, and by NSF grant IOS-1256572. We thank the many students, colleagues, technical staff, veterinarians, and animal caretakers who have contributed to our own studies on paternal care in rodents.

604

605 References

606

607 Ahern, T.H., Hammock, E.A., Young, L.J., 2010. Parental division of labor, coordination, and the effects
 608 of family structure on parenting in monogamous prairie voles (*Microtus ochrogaster*). Dev. Psychobiol.
 609 53, 118-131.

610 Ahern, T.H., Young, L.J., 2009. The impact of early life family structure on adult social attachment,
 611 alloparental behavior, and the neuropeptide systems regulating affiliative behaviors in the monogamous
 612 prairie vole (*Microtus ochrogaster*). Front. Behav. Neurosci. 3, 17.

613 Bales, K.L., Boone, E., Epperson, P., Hoffman, G., Carter, C.S., 2011a. Are behavioral effects of early
 614 experience mediated by oxytocin? Front. Child Neurodevel. Psychiat. 2, 24.

615 Bales, K.L., Carter, C.S., 2003. Developmental exposure to oxytocin facilitates partner preferences in
 616 male prairie voles (*Microtus ochrogaster*). Behav. Neurosci. 117, 854-859.

617 Bales, K.L., Kim, A.J., Lewis-Reese, A.D., Carter, C.S., 2004a. Both oxytocin and vasopressin may
 618 influence alloparental behavior in male prairie voles. Horm. Behav. 45, 354-361.

619 Bales, K.L., Kramer, K.M., Lewis-Reese, A.D., Carter, C.S., 2006. Effects of stress on parental care are
 620 sexually dimorphic in prairie voles. Physiol. Behav. 87, 424-429.

621 Bales, K.L., Lewis-Reese, A.D., Pfeifer, L.A., Kramer, K.M., Carter, C.S., 2007. Early experience affects
 622 the traits of monogamy in a sexually dimorphic manner. Devel. Psychobiol. 49, 335-342.

- 623 Bales, K.L., Maninger, N., Hinde, K., 2011b. New directions in the neurobiology and physiology of
 624 paternal care, in: Gillath, O., Adams, G., Kunkel, A.D. (Eds.), *New Directions in Relationship Research:
 625 Integrating Across Disciplines and Theoretical Approaches*. American Psychological Association.
- 626 Bales, K.L., Perkeybile, A.M., Conley, O.G., Lee, M.H., Guoynes, C.D., Downing, G.M., Yun, C.R.,
 627 Solomon, M., Jacob, S., Mendoza, S.P., 2013. Chronic intranasal oxytocin causes long-term impairment
 628 in partner preference formation in male prairie voles. *Biol. Psychiat.* 74, 180-188.
- 629 Bales, K.L., Pfeifer, L.A., Carter, C.S., 2004b. Sex differences and effects of manipulations of oxytocin
 630 on alloparenting and anxiety in prairie voles. *Dev. Psychobiol.* 44, 123-131.
- 631 Bamshad, M., Novak, M., de Vries, G., 1994. Cohabitation alters vasopressin innervation and paternal
 632 behavior in prairie voles (*Microtus ochrogaster*). *Physiol. Behav.* 56, 751-758.
- 633 Bamshad, M., Novak, M.A., De Vries, G.J., 1993. Sex and species differences in the vasopressin
 634 innervation of sexually naive and parental prairie voles, *Microtus ochrogaster*, and meadow voles,
 635 *Microtus pennsylvanicus*. *J. Neuroendocrinol.* 5, 247-255.
- 636 Becker, E.A., Moore, B.M., Auger, C., Marler, C.A., 2010. Paternal behavior increases testosterone levels
 637 in offspring of the California mouse. *Horm. Behav.* 58, 385-389.
- 638 Bester-Meredith, J.K., Marler, C.A., 2003. Vasopressin and the transmission of paternal behavior across
 639 generations in mated, cross-fostered *Peromyscus* mice. *Behav. Neurosci.* 117, 455-463.

- 640 Bester-Meredith, J.K., Young, L.J., Marler, C.A., 1999. Species differences in paternal behavior and
 641 aggression in *Peromyscus* and their associations with vasopressin immunoreactivity and receptors. *Horm.*
 642 *Behav.* 36, 25-38.
- 643 Braun, K., Champagne, F.A., 2014. Paternal influences on offspring development: behavioural and
 644 epigenetic pathways. *J. Neuroendocrinol.* 26, 697-706.
- 645 Braun, K., Seidel, K., Holetschka, R., Groeger, N., Poeggel, G., 2013. Paternal deprivation alters the
 646 development of catecholaminergic innervation in the prefrontal cortex and related limbic brain regions.
 647 *Brain Struct. Funct.* 218, 859-872.
- 648 Braun, K., Seidel, K., Weigel, S., Roski, C., Poeggel, G., 2011. Paternal deprivation alters region- and
 649 age-specific interneuron expression patterns in the biparental rodent, *Octodon degus*. *Cere. Cort.* 21,
 650 1532-1546.
- 651 Bredy, T.W., Lee, A.W., Meaney, M.J., Brown, R.E., 2004. Effect of neonatal handling and paternal care
 652 on offspring cognitive development in the monogamous California mouse (*Peromyscus californicus*).
 653 *Horm. Behav.* 46, 30-38.
- 654 Brooks, P.L., Vella, E.T., Wynne-Edwards, K.E., 2005. Dopamine agonist treatment before and after the
 655 birth reduces prolactin concentration but does not impair paternal responsiveness in Djungarian hamster,
 656 *Phodopus campbelli*. *Horm. Behav.* 47, 358-366.
- 657 Brown, R.E., Murdoch, T., Murphy, P.R., Moger, W.H., 1995. Hormonal responses of male gerbils to
 658 stimuli from their mate and pups. *Horm. Behav.* 29, 474-491.

- 659 Butler, E.A., Randall, A.K., 2013. Emotional coregulation in close relationships. *Emotion Rev.* 5, 202-
660 210.
- 661 Cabrera, N., Tamis-LeMonda, C.S., 2012. *Handbook of Father Involvement: Multidisciplinary*
662 *Perspectives*. Taylor and Francis, Inc.
- 663 Campbell, J.C., Laugero, K.D., Van Westerhuyzen, J.A., Hostetler, C.M., Cohen, J.D., Bales, K.L., 2009.
664 Costs of pair-bonding and parenting in male prairie voles (*Microtus ochrogaster*). *Physiol. Behav.* 98,
665 367-373.
- 666 Cantoni, D., Brown, R.E., 1997. Paternal investment and reproductive success in the California mouse,
667 *Peromyscus californicus*. *Anim. Behav.* 54, 377-386.
- 668 Cao, Y., Wu, R., Tai, F., Zhang, X., Yu, P., An, X., Qiao, X., Hao, P., 2014. Neonatal paternal
669 deprivation impairs social recognition and alters levels of oxytocin and estrogen receptor alpha mRNA
670 expression in the MeA and NAcc, and serum oxytocin in mandarin voles. *Horm. Behav.* 65, 57-65.
- 671 Chauke, M., Malisch, J.L., Robinson, C., de Jong, T.R., Saltzman, W., 2011. Effects of reproductive
672 status on behavioral and endocrine responses to acute stress in a biparental rodent, the California mouse
673 (*Peromyscus californicus*). *Horm. Behav.* 60, 128-138.
- 674 Churchland, P.S., Winkielman, P., 2012. Modulating social behavior with oxytocin: how does it work?
675 What does it mean? *Horm. Behav.* 61, 392-399.
- 676 Clark, M.M., Galef, B.G.J., 1999. A testosterone-mediated trade-off between parental and sexual effort in
677 male Mongolian gerbils (*Meriones unguiculatus*). *J. Comp. Psychol.* 113, 388-395.

- 678 Cushing, B.S., Perry, A., Musatov, S., Ogawa, S., Papademetriou, E., 2008. Estrogen receptors in the
679 medial amygdala inhibit the expression of male prosocial behavior. *J. Neurosci.* 28, 10399-10403.
- 680 Davis, M., Walker, D.L., Miles, L., Grillon, C., 2010. Phasic vs. sustained fear in rats and humans: Role
681 of the extended amygdala in fear and anxiety. *Neuropsychopharmacology* 35, 105-135.
- 682 De Jong , T., Harris, B., Perea-Rodriguez, J., Saltzman, W., 2013. Physiological and neuroendocrine
683 responses to chronic variable stress in male California mice (*Peromyscus californicus*): influence of social
684 environment and paternal state. . *Psychoneuroendocrinology* 38, 2023-2033.
- 685 De Jong, T.R., Chaulke, M., Harris, B.N., Saltzman, W., 2009. From here to paternity: neural correlates
686 of the onset of paternal behavior in California mice (*Peromyscus californicus*). *Horm. Behav.* 56, 220-
687 231.
- 688 De Jong, T.R., Measor, K.R., Chaulke, M., Harris, B.N., Saltzman, W., 2010. Brief pup exposure induces
689 Fos expression in the lateral habenula and serotonergic caudal dorsal raphe nucleus of paternally
690 experienced male California mice (*Peromyscus californicus*). *Neuroscience* 169, 1094-1104.
- 691 Dixon, A.F., 2013. Male infanticide and primate monogamy. *Proc. Nat. Acad. Sci.* 110, E4937.
- 692 Ebensperger, L.A., Ramirez-Otarola, N., Leon, C., Ortiz, M., Croxatto, H.B., 2010. Early fitness
693 consequences and hormonal correlates of parental behaviour in the social rodent, *Octodon degus*. *Physiol.*
694 *Behav.* 101, 509-517.
- 695 Elwood, R.W., Broom, D.M., 1978. The influence of litter size and parental behaviour in the development
696 of Mongolian gerbil pups. *Anim. Behav.* 26, 438-454.

- 697 Frazier, C.R., Trainor, B.C., Cravens, C.J., Whitney, T.K., Marler, C.A., 2007. Paternal behavior
 698 influences development of aggression and vasopressin expression in male California mouse offspring.
 699 Horm. Behav. 50, 699-707.
- 700 Gimpl, G., Fahrenholz, F., 2001. The Oxytocin Receptor System: Structure, function, and regulation.
 701 Physiol. Rev. 81, 629-683.
- 702 Glasper, E.R., Kozorovitskiy, Y., Pavlic, A., Gould, E., 2011. Paternal experience suppresses adult
 703 neurogenesis without altering hippocampal function in *Peromyscus californicus*. J. Comp. Neurol. 519,
 704 2271-2281.
- 705 Gleason, E.D., Marler, C.A., 2013. Non-genomic transmission of paternal behaviour between fathers and
 706 sons in the monogamous and biparental California mouse. Proc. Roy. Soc. Lond. B 280, 20130824.
- 707 Gos, T., Schulkin, J., Gos, A., Bock, J., Poeggel, G., Braun, K., 2014. Paternal deprivation affects the
 708 functional maturation of corticotropin-releasing hormone (CRH)- and calbindin-D28k-expressing neurons
 709 in the bed nucleus of the stria terminalis (BNST) of the biparental *Octodon degus*. Brain Struct. Funct.
 710 219, 1983-1990.
- 711 Greenberg, G.D., Van Westerhuyzen, J.A., Bales, K.L., Trainor, B.C., 2012. Is it all in the family? The
 712 effects of early social structure on neural-behavioral systems of prairie voles (*Microtus ochrogaster*).
 713 Neuroscience 216, 46-56.
- 714 Gubernick, D.J., Nelson, R.J., 1989. Prolactin and paternal behavior in the biparental California mouse,
 715 *Peromyscus californicus*. Horm. Behav. 23, 203-210.

- 716 Gubernick, D.J., Winslow, J.T., Jensen, P., Jeanotte, L., Bowen, J., 1995. Oxytocin Changes in Males
717 Over the Reproductive-Cycle in the Monogamous, Biparental California Mouse, *Peromyscus-*
718 *Californicus*. *Horm. Behav.* 29, 59-73.
- 719 Gubernick, D.J., Wright, S., Brown, R., 1993. The significance of father's presence for offspring survival
720 in the monogamous California mouse, *Peromyscus californicus*. *Anim. Behav.* 46, 539-546.
- 721 Harris, B., Saltzman, W., 2013. Effect of reproductive status on hypothalamic-pituitary-adrenal (HPA)
722 activity and reactivity in male California mice (*Peromyscus californicus*). *Physiol. Behav.* 112, 70-76.
- 723 Harris, B.N., De Jong, T.R., Yang, V., Saltzman, W., 2013. Chronic variable stress in fathers alters
724 paternal and social behavior but not pup development in the biparental California mouse (*Peromyscus*
725 *californicus*). *Horm. Behav.* 64, 799-811.
- 726 Harris, B.N., Perea-Rodriguez, J.P., Saltzman, W., 2011. Acute effects of corticosterone injection on
727 paternal behavior in California mouse (*Peromyscus californicus*) fathers. *Horm. Behav.* 60, 666-675.
- 728 Hume, J.M., Wynne-Edwards, K.E., 2005. Castration reduces male testosterone, estradiol, and territorial
729 aggression, but not paternal behavior in dwarf hamsters (*Phodopus campbelli*). *Horm. Behav.* 48, 303-
730 310.
- 731 Insel, T.R., Shapiro, L.E., 1992. Oxytocin receptor distribution reflects social organization in
732 monogamous and polygamous voles. *Proc. Nat. Acad. Sci.* 89, 5981-5985.

- 733 Jia, R., Tai, F., Shucheng, A., Zhang, X., Broders, H., 2009. Effects of neonatal paternal deprivation or
 734 early deprivation on anxiety and social behaviors of the adults in mandarin voles. *Behav. Proc.* 82, 271-
 735 278.
- 736 Kenkel, W.M., Paredes, J., Yee, J.R., Pournajafi-Nazarloo, H., Bales, K.L., Carter, C.S., 2012.
 737 Neuroendocrine and behavioural responses to exposure to an infant in male prairie voles. *J.*
 738 *Neuroendocrinol.* 24, 874-886.
- 739 Kirkpatrick, B., Carter, C.S., Newman, S.W., Insel, T.R., 1994a. Axon-sparing lesions of the medial
 740 nucleus of the amygdala decrease affiliative behaviors in the prairie vole (*Microtus ochrogaster*) -
 741 behavioral and anatomical specificity. *Behavioral Neuroscience* 108, 501-513.
- 742 Kirkpatrick, B., Kim, J.W., Insel, T.R., 1994b. Limbic system fos expression associated with paternal
 743 behavior. *Brain Res.* 658, 112-118.
- 744 Kirkpatrick, B., Williams, J.R., Slotnick, B.M., Carter, C.S., 1994c. Olfactory bulbectomy decreases
 745 social behavior in male prairie voles (*Microtus ochrogaster*). *Physiol. Behav.* 55, 885-889.
- 746 Kleiman, D.G., 1977. Monogamy in mammals. *Quart. Rev. Biol.* 52, 39-69.
- 747 Kramer, K.M., Perry, A.N., Golbin, D., Cushing, B.S., 2009. Sex steroids are necessary in the second
 748 postnatal week for the expression of male alloparental behavior in prairie voles (*Microtus ochrogaster*).
 749 *Behav. Neurosci.* 123, 958-963.
- 750 Lambert, K.G., Franssen, C.L., Bardi, M., Hampton, J.E., Hainley, L., Karsner, S., Tu, E.B., Hyer, M.M.,
 751 Crockett, A., Baranova, A., Ferguson, T., Ferguson, T., Kinsley, C.H., 2011. Characteristic

- 752 neurobiological patterns differentiate paternal responsiveness in two *Peromyscus* species. *Brain Behav.*
 753 *Evol.* 77, 159-175.
- 754 Lambert, K.G., Franssen, C.L., Hampton, J.E., Rzucidlo, A.M., Hyer, M.M., True, M., Kaufman, C.,
 755 Bardi, M., 2013. Modeling paternal attentiveness: distressed pups evoke differential neurobiological and
 756 behavioral responses in paternal and nonpaternal mice. *Neuroscience* 234, 1-12.
- 757 Lee, A.W., Brown, R.E., 2002. Medial preoptic lesions disrupt parental behavior in both male and female
 758 California mice (*Peromyscus californicus*). *Behav. Neurosci.* 116, 968-975.
- 759 Lee, A.W., Brown, R.E., 2007. Comparison of medial preoptic, amygdala, and nucleus accumbens lesions
 760 on parental behavior in California mice (*Peromyscus californicus*). *Physiol. Behav.* 92, 617-628.
- 761 Lei, K., Cushing, B.S., Musatov, S., Ogawa, S., Kramer, K.M., 2010. Estrogen receptor alpha in the bed
 762 nucleus of the stria terminalis regulates social affiliation in male prairie voles (*Microtus ochrogaster*).
 763 *PLoS One* 5, e8931.
- 764 Li, Y., Lian, Z., Wang, B., Tai, F., Wu, R., Hao, P., Qiao, X., 2015. Natural variation in paternal behavior
 765 is associated with central estrogen receptor alpha and oxytocin levels. *J. Comp. Physiol. A* 201, 285-293.
- 766 Liang, M., Zhong, J., Liu, H.-X., Lopatina, O., Nakada, R., Yamauchi, A.M., Higashida, H., 2014.
 767 Pairmate-dependent pup retrieval as parental behavior in male mice. *Front. Neurosci.* 8, 186.
- 768 Lieberwirth, C., Wang, Y., Jia, X., Liu, Y., Wang, Z., 2013. Fatherhood reduces the survival of adult-
 769 generated cells and affects various types of behavior in the prairie vole (*Microtus ochrogaster*). *Eur. J.*
 770 *Neurosci.* 38, 3345-3355.

- 771 Lightman, S.L., Windle, R.J., Wood, S.A., Kershaw, Y.M., Shanks, N., Ingram, C.D., 2001. Peripartum
772 plasticity within the hypothalamo-pituitary-adrenal axis. *Prog. Brain Res.* 133, 111-129.
- 773 Liu, H.-X., Lopatina, O., Higashida, C., Fujimoto, H., Akther, S., Inzhutova, A., Liang, M., Zhong, J.,
774 Tsuji, T., Yoshihara, T., Sumi, K., Ishiyama, M., Ma, W.J., Ozaki, M., Yagitani, S., Yokoyama, S.,
775 Mukaida, N., Sakurai, T., Hori, O., Yoshioka, K., Hirao, A., Kato, Y., Ishihara, K., Kato, I., Okamoto, H.,
776 Cherepanov, S.M., Salmina, A.B., Hirai, H., Asano, M., Brown, D.A., Nagano, I., Higashida, H., 2013.
777 Displays of paternal pup retrieval following communicative interaction with maternal mates. *Nature*
778 *Comm.* 4, 1346.
- 779 Lonstein, J.S., De Vries, G.J., 1999. Sex differences in the parental behaviour of adult virgin prairie voles:
780 Independence from gonadal hormones and vasopressin. *J. Neuroendocrinol.* 11, 441-449.
- 781 Lonstein, J.S., De Vries, G.J., 2000. Influence of gonadal hormones on the development of parental
782 behavior in adult virgin prairie voles (*Microtus ochrogaster*). *Behav. Brain Res.* 114, 79-87.
- 783 Lonstein, J.S., Rood, B.D., De Vries, G.J., 2002. Parental responsiveness is feminized after neonatal
784 castration in virgin male prairie voles, but is not masculinized by perinatal testosterone in virgin females.
785 *Horm. Behav.* 41, 80-87.
- 786 Lubin, M., Leon, M., Moltz, H., Numan, M., 1972. Hormones and maternal behavior in the male rat.
787 *Horm. Behav.* 3, 369-374.
- 788 Lukas, D., Clutton-Brock, T.H., 2013. The evolution of social monogamy in mammals. *Science* 341, 526-
789 530.

- 790 Lukas, D., Huchard, E., 2014. Sexual conflict. The evolution of infanticide by males in mammalian
791 societies. *Science* 346, 841-844.
- 792 Ma, E., Lau, J., Grattan, D., Lovejoy, D., Wynne-Edwards, K., 2005. Male and female prolactin receptor
793 mRNA expression in the brain of a biparental and a uniparental hamster, *Phodopus*, before and after the
794 birth of a litter. *J. Neuroendocrinol.* 17, 81-90.
- 795 Mak, G.K., Antle, M.C., Dyck, R.H., Weiss, S., 2013. Bi-parental care contributes to sexually dimorphic
796 neural cell genesis in the adult mammalian brain. *PLoS One* 8, e62701.
- 797 Mak, G.K., Weiss, S., 2010. Paternal recognition of adult offspring mediated by newly generated CNS
798 neurons. *Nature Neurosci.* 13, 753-758.
- 799 McCullough, J., Quadagno, D.M., Goldman, B.D., 1974. Neonatal gonadal hormones: Effect on maternal
800 and sexual behavior in the male rat. *Physiol. Behav.* 12, 183-188.
- 801 Numan, M., 2014. *Neurobiology of Social Behavior: Toward an Understanding of the Prosocial and*
802 *Antisocial Brain*. Academic Press, CA.
- 803 Numan, M., Insel, T.R., 2003. *The Neurobiology of Parental Behavior*. Springer-Verlag, New York.
- 804 Opie, C., Atkinson, Q.D., Dunbar, R.I.M., Shultz, S., 2013a. Male infanticide leads to social monogamy
805 in primates. *Proc. Nat. Acad. Sci.* 110, 13328-13332.
- 806 Opie, C., Atkinson, Q.D., Dunbar, R.I.M., Shultz, S., 2013b. Reply to Dixson: infanticide triggers primate
807 monogamy. *Proc. Nat. Acad. Sci.* 110, e4938.

- 808 Opie, C., Atkinson, Q.D., Dunbar, R.I.M., Shultz, S., 2014. Reply to Lukas and Clutton-Brock:
809 infanticide still drives primate monogamy. *Proc. Nat. Acad. Sci.* 111, e1675.
- 810 Outscharoff, W.J., Helmeke, C., Braun, K., 2006. Lack of paternal care affects synaptic development in
811 the anterior cingulate cortex. *Brain Res.* 1116, 58-63.
- 812 Parker, J., Tsagkogeorga, G., Cotton, J.A., Liu, Y., Provero, P., Stupka, E., Rossiter, S.J., 2013. Genome-
813 wide signatures of convergent evolution in echolocating mammals. *Nature* 503, 228-231.
- 814 Parker, K.J., Kinney, L.F., Phillips, K.M., Lee, T.M., 2001. Paternal behavior is associated with central
815 neurohormone receptor binding patterns in meadow voles (*Microtus pennsylvanicus*). *Behav. Neurosci.*
816 115, 1341-1348.
- 817 Parker, K.J., Lee, T.M., 2001. Central vasopressin administration regulates the onset of facultative
818 paternal behavior in *Microtus pennsylvanicus* (meadow voles). *Horm. Behav.* 39, 285-294.
- 819 Perea-Rodriguez, J.P., Takahashi, E.Y., Amador, T.M., Hao, R.C., Saltzman, W., Trainor, B.C., 2015.
820 Effects of reproductive experience on central expression of progesterone, oestrogen receptor (alpha),
821 oxytocin, and vasopressin (V1a) receptor mRNA in the male California mouse (*Peromyscus californicus*).
822 *J. Neuroendocrinol.* 27, 245-252.
- 823 Perkeybile, A.M., Griffin, L.L., Bales, K.L., 2013. Natural variation in early parental care correlates with
824 social behaviors in adolescent prairie voles (*Microtus ochrogaster*). *Front. Behav. Neurosci.* 7, 21.

- 825 Pinkernelle, J., Abraham, A., Seidel, K., Braun, K., 2009. Paternal deprivation induces dendritic and
826 synaptic changes and hemispheric asymmetry of pyramidal neurons in the somatosensory cortex. *Dev.*
827 *Neurobiol.* 69, 663-673.
- 828 Piovanotti, M.R.A., Vieira, M.L., 2004. Presence of the father and parental experience have differentiated
829 effects on pup development in Mongolian gerbils (*Meriones unguiculatus*). *Behav. Proc.* 66, 107-117.
- 830 Reburn, C.J., Wynne-Edwards, K.E., 1999. Hormonal changes in males of a naturally biparental and a
831 uniparental mammal. *Horm. Behav.* 35, 163-176.
- 832 Roberts, R.L., Williams, J.R., Wang, A.K., Carter, C.S., 1998. Cooperative breeding and monogamy in
833 prairie voles: influence of the sire and geographical variation. *Anim. Behav.* 55, 1131-1140.
- 834 Roberts, R.L., Zullo, A.S., Carter, C.S., 1996. Perinatal steroid treatments alter alloparental and affiliative
835 behavior in prairie voles. *Horm. Behav.* 30, 576-582.
- 836 Rosenberg, K.M., 1974. Effects of pre- and post-pubertal castration and testosterone on pup-killing
837 behavior in the male rat. *Physiol. Behav.* 13, 159-161.
- 838 Rosenberg, P.A., Herrenkohl, L.R., 1976. Maternal behavior in male rats: Critical times for the
839 suppressive action of androgens. *Physiol. Behav.* 16, 293-297.
- 840 Rosenblatt, J.S., 1967. Nonhormonal basis of maternal behavior in the rat. *Science* 156, 1512-1514.
- 841 Rosenblatt, J.S., Ceus, K., 1998. Estrogen implants in the medial preoptic area stimulate maternal
842 behavior in male rats. *Horm. Behav.* 33, 23-30.

- 843 Rosenblatt, J.S., Hazelwood, S., Poole, J., 1996. Maternal behavior in male rats: Effects of medial
844 preoptic area lesions and presence of maternal aggression. *Horm. Behav.* 30, 201-215.
- 845 Rosenfeld, C.S., Johnson, S.A., Ellersieck, M.R., Roberts, R.M., 2013. Interactions between parents and
846 parents and pups in the monogamous California mouse (*Peromyscus californicus*). *PLoS One* 8.
- 847 Sakaguchi, K., Tanaka, M., Ohkubo, T., Doh-ura, K., Fujikawa, T., Sudo, S., Nakashima, K., 1996.
848 Induction of brain prolactin receptor long-form mRNA expression and maternal behavior in pup-
849 contacted male rats: promotion by prolactin administration and suppression by female contact.
850 *Neuroendocrinology* 63, 559-568.
- 851 Saltzman, W., Ziegler, T.E., 2014. Functional significance of hormonal changes in mammalian fathers. *J.*
852 *Neuroendocrinol.* 26, 685-696.
- 853 Schneider, J.S., Stone, M.K., Wynne-Edwards, K.E., Horton, T.H., Lydon, J., O'Malley, B., Levine, J.E.,
854 2003. Progesterone receptors mediate male aggression toward infants. *Proc. Nat. Acad. Sci.* 100, 2951-
855 2956.
- 856 Schradin, C., 2008. Differences in prolactin levels between three alternative male reproductive tactics in
857 striped mice (*Rhabdomys pumilio*). *Proc. Biol. Sci.* 275, 1047-1052.
- 858 Schum, J.E., Wynne-Edwards, K.E., 2005. Estradiol and progesterone in paternal and non-paternal
859 hamster (*Phodopus*) becoming fathers: conflict with hypothesized roles. *Horm. Behav.* 47, 410-418.

- 860 Seidel, K., Poeggel, G., Holetschka, R., Helmeke, C., Braun, K., 2011. Paternal deprivation affects the
861 development of corticotrophin-releasing-factor-expressing neurones in prefrontal cortex, amygdala, and
862 hippocampus of the biparental *Octodon degus*. J. Neuroendocrinol. 23, 1166-1176.
- 863 Song, Z., Tai, F., Yu, C., Wu, R., Zhang, X., Broders, H., He, F., Guo, R., 2010. Sexual or paternal
864 experiences alter alloparental behavior and the central expression of ERalpha and OT in male mandarin
865 voles (*Microtus mandarinus*). Behav. Brain Res. 214, 290-300.
- 866 Stone, A.I., Bales, K.L., 2010. Intergenerational transmission of the behavioral consequences of early
867 experience in prairie voles. Behav. Proc. 84, 732-738.
- 868 Stone, A.I., Mathieu, D., Griffin, L., Bales, K.L., 2010. Alloparenting experience affects future parental
869 behavior and reproductive success in prairie voles (*Microtus ochrogaster*). Behav. Proc. 83, 8-15.
- 870 Sturgis, J.D., Bridges, R.S., 1997. N-methyl-DL-aspartic acid lesions of the medial preoptic area disrupt
871 ongoing parental behavior in male rats. Physiol. Behav. 62, 305-310.
- 872 Tardif, S., Richter, C.B., Carson, R.L., 1984. Effects of sibling-rearing experience on future reproductive
873 success in two species of Callitrichidae. Am. J. Primatol. 6, 377-380.
- 874 Taymans, S.E., DeVries, A.C., DeVries, M.B., Nelson, R.J., Friedman, T.C., Detera-Wadleigh, S., Carter,
875 C.S., Chrousos, G.P., 1997. The hypothalamic-pituitary-adrenal axis of prairie voles (*Microtus*
876 *ochrogaster*): evidence for target tissue glucocorticoid resistance. Gen. Comp. Endocrinol. 106, 48-61.

- 877 Timonin ME, Cushing BS, KE, W.-E., 2008. In three brain regions central to maternal behaviour, neither
878 male nor female *Phodopus dwarf* hamsters show changes in oestrogen receptor alpha distribution with
879 mating or parenthood. *J. Neuroendocrinol.* 20, 1301-1309.
- 880 Trainor, B.C., Bird, M., Alday, M.A., Schlinger, B.A., Marler, C.A., 2003. Variation in aromatase activity
881 in the medial preoptic area and plasma progesterone is associated with the onset of paternal behavior.
882 *Neuroendocrinology* 78, 36-44.
- 883 Trainor, B.C., Marler, C.A., 2001. Testosterone, paternal behavior, and aggression in the monogamous
884 California mouse (*Peromyscus californicus*). *Horm. Behav.* 40, 32-42.
- 885 Trainor, B.C., Marler, C.A., 2002. Testosterone promotes paternal behaviour in a monogamous mammal
886 via conversion to oestrogen. *Proc. Roy. Soc. of Lond. B* 269, 823-829.
- 887 Vorbrodt, A.W., 1993. Morphological evidence of the functional polarization of brain microvascular
888 epithelium, in: Pardridge, W.M. (Ed.), *The Blood-Brain Barrier*. Raven Press, New York, pp. 137-164.
- 889 Wang, J., Tai, F., Yan, X., Yu, P., 2012. Paternal deprivation alters play-fighting, serum corticosterone
890 and the expression of hypothalamic vasopressin and oxytocin in juvenile male mandarin voles. *J. Comp.*
891 *Physiol. A* 198, 787-796.
- 892 Wang, L., Zhang, W., Wu, R., Kong, L., Feng, W., Cao, Y., Tai, F., Zhang, X., 2014. Neuroendocrine
893 responses to social isolation and paternal deprivation at different postnatal ages in mandarin voles. *Dev.*
894 *Psychobiol.* 56, 1214-1228.

- 895 Wang, Z., De Vries, G.J., 1993. Testosterone effects on paternal behavior and vasopressin
896 immunoreactive projections in prairie voles (*Microtus ochrogaster*). Brain Res. 631, 156-160.
- 897 Wang, Z.X., Ferris, C.F., Devries, G.J., 1994. Role of septal vasopressin innervation in paternal behavior
898 in prairie voles (*Microtus ochrogaster*). Proc. Nat. Acad. Sci. 91, 400-404.
- 899 Wang, Z.X., Liu, Y., Young, L.J., Insel, T.R., 2000. Hypothalamic vasopressin gene expression increases
900 in both males and females postpartum in a biparental rodent. J. Neuroendocrinol. 12, 111-120.
- 901 Wang, Z.X., Novak, M.A., 1992. Influence of the social environment on parental behavior and pup
902 development of meadow voles (*Microtus pennsylvanicus*) and prairie voles (*Microtus ochrogaster*). J.
903 Comp. Psychol. 106, 163-171.
- 904 Wright, S.L., Brown, R.E., 2002. The importance of paternal care on pup survival and pup growth in
905 *Peromyscus californicus* when required to work for food. Behav. Proc. 60, 41-52.
- 906 Wu, R., Song, Z., Wang, S., Shui, L., Tai, F., Qiao, X., He, F., 2014a. Early paternal deprivation alters
907 levels of hippocampal brain-derived neurotrophic factor and glucocorticoid receptor and serum
908 corticosterone and adrenocorticotropin in a sex-specific way in socially monogamous mandarin voles.
909 Neuroendocrinology 100, 119-128.
- 910 Wu, Z., Autry, A.E., Bergan, J.F., Watabe-Uchida, M., Dulac, C.G., 2014b. Galanin neurons in the
911 medial preoptic area govern parental behaviour. Nature 509, 325-330.

- 912 Wynne-Edwards, K., Timonin, M., 2007. Paternal care in rodents: weakening support of hormonal
913 regulation of the transition to behavioral fatherhood in rodent animal models of biparental care. *Horm.*
914 *Behav.* 52, 114-121.
- 915 Yu, P., Shucheng, A., Tai, F., Zhang, X., He, F., Wang, J., An, X., Wu, R., 2012. The effects of neonatal
916 paternal deprivation on pair bonding, NAcc dopamine receptor mRNA expression and serum
917 corticosterone in mandarin voles. *Horm. Behav.* 61, 669-677.
- 918

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Table 1. Summary of findings on neural influences on and correlates of paternal care in biparental rodents. IHC - immunohistochemistry; ER α - estrogen receptor alpha

Species	Brain Region	Method	Experimental Group	Results	Reference
California mouse	Medial preoptic area	Electrolytic lesions	First-time fathers	Increased latencies to retrieve, lick, and hover over pups; decreased time spent sniffing, licking, and near pups	Lee and Brown, 2002
			Fathers	Decreased time licking and hovering over pups; increased latencies to retrieve pups	
California mouse	Medial preoptic area	Fos IHC	Adult males	Increased Fos-ir in new fathers exposed to a pup; no effect in virgin males or first-time expectant fathers	De Jong et al., 2009
California mouse	Basolateral amygdala	Electrolytic lesions	Fathers	Increased latency to approach pups; decreased time licking pups and near pups	Lee and Brown, 2007
California mouse	Medioventral amygdala	Fos IHC	Adult males	Increased Fos-ir following exposure to a pup in fathers compared to virgin males	De Jong et al., 2009
California mouse	Medial posteromedial amygdala	Fos IHC	Adult males	Increased Fos-ir following exposure to a pup in fathers compared to virgin males	De Jong et al., 2009
California mouse	Nucleus accumbens	Electrolytic lesions	Fathers	Decreased latency to retrieve pups	Lee and Brown, 2007
California mouse	Caudal dorse raphe nucleus	Fos IHC	Fathers	Increased Fos-ir following exposure to a pup	De Jong et al., 2009; De Jong et al., 2010
California mouse	Lateral habenula	Fos IHC	Fathers	Increased Fos-ir following exposure to a pup	De Jong et al., 2009; De Jong et al., 2010

California mouse	Medial preoptic area	Fos IHC	Fathers	Increased Fos-ir following exposure to a pup	Lambert et al., 2013
	Posterior cingulate cortex			Decreased Fos-ir following exposure to a pup	Lambert et al., 2011, 2013
Prairie vole	Basolateral amygdala	Excitotoxic lesions	Adult virgin males	No effect on responses to pups ²²	Kirkpatrick et al., 1994a
	Medial amygdala	Electrolytic lesions		Decreased contact time with pup	Kirkpatrick et al., 1994a
	Corticomedial amygdala	Electrolytic lesions		Decreased contact time with pup, increased time away from pup	
Prairie vole	Medial preoptic area, Medial amygdala, Lateral septum, Paraventricular nucleus of the thalamus, Nucleus reuniens, Accessory olfactory bulbs, Medial bed nucleus of the stria terminalis	Fos IHC	Young adult virgin males	Increased Fos-ir following exposure to a pup	Kirkpatrick et al., 1994b
Prairie vole	Medial amygdala	Increased expression of ER α via viral vector	Adult males	Decreased alloparental behavior	Cushing et al., 2008
Prairie vole	Bed nucleus of stria terminalis	Increased expression of	Adult males	No effect	Lei et al., 2010

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

		ERα via viral vector			
Prairie vole	Olfactory bulbs	Surgical removal (bilateral)	Adult virgin males	Increased likelihood of attacking pups	Kirkpatrick et al., 1994c

Table 2. Summary of findings on hormonal and neuropeptide influences on paternal care in biparental rodents. All manipulations were performed on adult animals unless otherwise indicated. T – testosterone, E – estrogen, DHT – dihydrotestosterone, ER α – estrogen receptor alpha, PR – progesterone receptor, BST – bed nucleus of the stria terminalis, ICV – intracerebroventricular, AVP – vasopressin, OT – oxytocin

Species	Hormone/ Neuropeptide	Organizational or Activational	Method	Experimental Group	Results	Reference
Djungarian hamster	Prolactin	Activational	Treatment with dopamine agonists bromocriptine or cabergoline to decrease prolactin secretion	First-time fathers	No effect on paternal behavior	Brooks et al., 2005
Djungarian hamster	Gonadal steroids	Activational	Castration	Fathers	No effect on paternal behavior	Hume and Wynne-Edwards, 2005
Djungarian hamster	Estrogen	Activational	Treatment with letrozole (aromatase inhibitor)	First-time fathers	No effect on paternal behavior	Hume and Wynne-Edwards, 2006
Mongolian gerbil	Testosterone	Activational	Castration + T replacement	Males paired with pregnant females	Castration increased and T replacement decreased paternal behavior	Clark and Galef, 1999
California mouse	Gonadal steroids	Activational	Castration + E, T, or DHT replacement	Fathers	Paternal behavior was decreased by castration and reinstated by treatment with E or T, but not with DHT (non-aromatizable) or T + fadrozole (aromatase inhibitor)	Trainor and Marler, 2001, 2002
Prairie vole	Estrogen	Activational	Castration + estradiol replacement	Adult virgin males	No effect on paternal behavior	Lonstein and DeVries, 1999
Prairie vole	Estrogen	Activational	Increase expression of ER α in medial amygdala via viral vector	Adult males	Decreased paternal behavior	Cushing et al., 2008

Prairie vole	Estrogen	Activational	Increase expression of ER α in BST via viral vector	Adult virgin males	No effect on paternal behavior	Lei et al., 2010
Prairie vole	Testosterone	Activational	Castration + T replacement	Adult virgin males	Paternal behavior and pup-directed aggression were decreased by castration and reinstated by T replacement	Wang and De Vries, 1993
California mouse	Corticosterone	Activational	Acute corticosterone treatment	First-time fathers	No effect on paternal behavior	Harris et al., 2011
California mouse	Corticosterone	Activational	Variable chronic stress, leading to chronic increase in corticosterone	First-time fathers	Transient decreases in paternal behavior	Harris et al., 2013
Prairie vole	Corticosterone	Activational	Acute swim stressor, leading to likely increases in corticosterone	Adult virgin males	Increased paternal behavior	Bales et al., 2006
Prairie vole	Vasopressin	Activational	Infusion of AVP or AVP receptor V1a antagonist into lateral septum	Adult virgin males	AVP increased and V1a receptor antagonist decreased paternal behavior	Wang et al., 1994
Meadow vole (facultative paternal care)	Vasopressin	Activational	ICV infusion of AVP or AVP V1a receptor antagonist	Adult virgin males	AVP decreased pup-directed aggression and increased paternal behavior; V1a antagonist decreased paternal behavior	Parker and Lee, 2001
Prairie vole	Vasopressin, oxytocin	Activational	ICV infusion of AVP, OT or both; or of AVP, antagonist, OT antagonist, or both	Adult virgin males	Combined treatment with AP antagonist and OT antagonist increased latency to approach pup, increased proportion of males attacking pup, decreased kyphosis, and decreased proportion of males	Bales et al., 2004a

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

					behaving paternally	
Prairie vole	Oxytocin	Organizational	OT or OT antagonist injection on PND1	Weanlings and adult virgin males	OT antagonist decreased paternal behavior and increased pup-directed aggression in weanlings	Bales et al., 2004b
Prairie vole	Oxytocin	Organizational (developmental)	Intranasal OT treatment during juvenile period	Adult virgin males	No effect on paternal behavior	Bales et al., 2013

Table 3. Summary of findings on effects of paternal deprivation on brain in biparental species, organized by species. PVN = paraventricular nucleus of the hypothalamus, OT = oxytocin, GR = glucocorticoid receptors, BDNF = brain-derived neurotrophic factor, ACTH = adrenocorticotrophin, D1R = dopamine type 1 receptors, D2R = dopamine type 2 receptors, ER α = estrogen receptor alpha, OTR = oxytocin receptor, CRFR2 = corticotrophin-release factor type 2 receptors

Species	Brain Region/ Behavior	Experimental Group	Results (compared to animals raised by both parents)	Reference
Prairie vole	PVN	Paternaly deprived females	Increased numbers of clusters of OT mRNA	Ahern et al., 2010
	Dorsal raphe	Paternaly deprived, both sexes	Higher density of CRFR2 receptors	
Mandarin vole	Dentate gyrus CA1, CA 2/3	Paternaly deprived females	Decreases in GR and BDNF	Wu et al., 2014
Mandarin vole	Nucleus accumbens	Paternaly deprived females	Reduced D1R and D2R Reduced OTR mRNA	Yu et al., 2012 Cao et al., 2014
Mandarin vole	Medial amygdala	Paternaly deprived females	Reduced ER α No change in OTR mRNA	Cao et al., 2014
	Serum		Reduced serum OT	
Mandarin vole	Dentate gyrus CA1, CA 2/3	Paternaly deprived males	No change in GR and BDNF in Dentate Reductions in GR and BDNF in CA1, CA 2/3	Wu et al., 2014
Mandarin vole	Nucleus accumbens	Paternaly deprived males	Higher D1R and D2R Reduced OTR mRNA	Yu et al., 2012 Cao et al., 2014
Mandarin vole	Medial amygdala	Paternaly deprived males	No change in ER α Lower OTR mRNA	Cao et al., 2014
Mandarin	Hypothalamus	Paternaly deprived	Reduced AVP-ir in anterior hypothalamus	Wang et al., 2012

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

vole		males	Reduced OT-ir in PVN	
Mandarin vole	Serum	Paternally deprived males	No change in serum OT	Cao et al., 2014
Octodon degus	Dentate gyrus CA1	Paternally deprived males	Fewer CRF-positive pyramidal cells at PND21 Fewer CRF-positive pyramidal cells at PND21 and 90	Seidel et al., 2011
Octodon degus	Medial BST	Paternally deprived males	Fewer CRF-positive cells at PND21	Gos et al., 2014
	PVN			
		Paternally deprived males	No change in CRF-positive cells	
	Basolateral amygdala	Paternally deprived males	More CRF-positive cells at PND21	
Octodon degus	CA1, nucleus accumbens, central amygdala	Paternally deprived males	Elevated tyrosine hydroxylase	Braun et al.,2013