Maternal stress and plural breeding with communal care affect development of the endocrine stress response in a wild rodent

Carolyn M. Bauer ^{a,*}, Loren D. Hayes ^b, Luis A. Ebensperger ^c, Juan Ramírez-Estrada ^c, Cecilia León ^c, Garrett T. Davis ^b, and L. Michael Romero ^a

^a *Department of Biology, Tufts University, Medford, MA, USA*

^b *Department of Biological and Environmental Sciences, University of Tennessee at Chattanooga, TN, USA*

^c *Departamento de Ecología, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile*

* *Corresponding author:* Carolyn M. Bauer

Mailing address: Biological Sciences, Dept. 2715, North Dakota State University, PO Box 6050, Fargo, ND, 58108-6050, USA

Phone: +1 701 231 8999

E-mail address: carolyn.marie.bauer@gmail.com

[‡] *Current address:* Biological Sciences, Dept. 2715, North Dakota State University, PO Box 6050, Fargo, ND, 58108-6050, USA

24

25 **Abstract**

26

27 Maternal stress can significantly affect offspring fitness. In laboratory rodents,
28 chronically stressed mothers provide poor maternal care, resulting in pups with
29 hyperactive stress responses. These hyperactive stress responses are characterized by
30 high glucocorticoid levels in response to stressors plus poor negative feedback, which can
31 ultimately lead to decreased fitness. In degus (*Octodon degus*) and other plural breeding
32 rodents that exhibit communal care, however, maternal care from multiple females may
33 buffer the negative impact on pups born to less parental mothers. We used wild, free-
34 living degus to test this hypothesis. After parturition, we manipulated maternal stress by
35 implanting cortisol pellets in 0%, 50–75%, or 100% of adult females within each social
36 group. We then sampled pups for baseline and stress-induced cortisol, negative feedback
37 efficacy, and adrenal sensitivity. From groups where all mothers were implanted with
38 cortisol, pups had lower baseline cortisol levels and male pups additionally had weaker
39 negative feedback compared to 0% or 50–75% implanted groups. Contrary to
40 expectations, stress-induced cortisol did not differ between treatment groups. These data
41 suggest that maternal stress impacts some aspects of the pup stress response, potentially
42 through decreased maternal care, but that presence of unstressed mothers may mitigate
43 some of these effects. Therefore, one benefit of plural breeding with communal care may
44 be to buffer post-natal stress.

45

46 **Keywords**

Communal care; Cortisol; Hypothalamic-Pituitary-Adrenal axis; Maternal effects;
Maternal stress; Negative feedback; *Octodon degus*; Post-natal stress

Introduction

Maternal effects are important contributors to phenotypic variation among individuals (Mousseau and Fox, 1998). Independent of genetic contributions, mothers can significantly influence the phenotypes of their offspring through habitat selection and resource allocation during development. After birth, mothers and social group members can continue to affect offspring phenotypes by altering the quality and quantity of parental care. These parental care behaviors can significantly influence offspring fitness by affecting future reproductive output (Lindstrom, 1999), growth rate (Metcalf and Monaghan, 2001), immune function (Saino et al., 1997), and stress responsiveness (Francis et al., 1999; Liu et al., 1997). Low rates of parental care have generally been associated with negative or maladaptive offspring development (Cottrell and Seckl, 2009; Meaney et al., 2007) although there has been increasing recognition of maternal effects as potential adaptive responses to environmental heterogeneity, as what may be considered a “maladaptive phenotype” in a benign environment can be beneficial in a stressful or challenging environment (Dantzer et al., 2013; Mousseau and Fox, 1998; Sheriff and Love, 2013).

For mammals, parental care encompasses many different behaviors including thermoregulatory huddling, nursing, predator protection, and licking and grooming. In rodents, licking and grooming, which are almost exclusively performed by the mother,

are important for development of the hypothalamic-pituitary-adrenal (HPA) axis (Francis et al., 1999; Liu et al., 1997). Chronically stressed rodent mothers have lower rates of licking and grooming (Brummelte and Galea, 2010), and pups receiving relatively low amounts of licking and grooming increase glucocorticoid (GC) secretion (Liu et al., 1997). Pups exposed to higher, sustained levels of GCs are more likely to develop hyper-reactive stress responses (Francis et al., 1999; Liu et al., 1997), which are characterized by increased GC output in response to stressors, plus poor negative feedback. Together, these factors increase long-term GC exposure, which has been linked to lower adult survival and reproductive success (Cottrell and Seckl, 2009; Matthews and Phillips, 2010; Monaghan et al., 2012).

Because the stress response is a key component in maintaining physiological homeostasis (McEwen and Wingfield, 2003; Romero et al., 2009), maternal-induced alterations to the offspring HPA-axis can have widespread effects on numerous physiological and behavioral processes. When an animal encounters a stressor, corticotropin-releasing factor (CRF) is secreted from the hypothalamus, which stimulates release of adrenocorticotrophic hormone (ACTH) from the anterior pituitary, which then increases secretion of GCs from the adrenal glands. When circulating GC levels are elevated, a variety of physiological effects occur that essentially shift energy away from non-necessary processes to those crucial for survival. In the short-term, these effects include increased energy mobilization, inhibition of the reproductive system, and priming of the fight-or-flight response (Sapolsky et al., 2000). However, these “adaptive” physiological effects can shift to being pathological if GC levels remain elevated for long periods of time, resulting in diabetes, immunosuppression, reproductive shut-down, and

cardiovascular disease (McEwen, 1998; Sapolsky et al., 2000). Animals with hyper-reactive stress responses may be more susceptible to developing these pathologies (Romero and Wikelski, 2010), as amplified GC responses to stressors and poor negative feedback cause greater, integrated GC exposure.

While the relationships between post-parturition maternal stress, maternal care, and the offspring stress response have been well studied in laboratory rodents (Hennessy, 2003; Liu et al., 1997; Own and Patel, 2013), this has remained relatively unexplored in wild, free-living animals. Additionally, there are few studies examining whether care from other members within a social group can significantly contribute to the development of the offspring stress response (Birnie et al., 2013; Branchi et al., 2013; Harris et al., 2013). We assessed whether post-parturition maternal stress affected the development of the offspring stress response in wild, free-living degus (*Octodon degus*). Degus are social, caviomorph rodents endemic to central Chile. Cortisol is the predominant glucocorticoid hormone in degus (Kenagy et al., 1999), and while degus are precocial rodents, their post-natal basal glucocorticoid profiles are more similar to altricial rodents such as rats and mice than to other precocial caviomorph rodents such as domestic guinea pigs (*Cavia porcellus*) (Gruss et al., 2006). Degus are plural breeders, which means that several females reproduce and raise their offspring within a group (Hayes, 2000). Additionally, degus exhibit communal care, where mothers within the group will lick, groom, and even nurse other group member's offspring (Hayes et al., 2009). There are obvious fitness consequences to plural breeding with communal care, but a recent study using a long-term dataset suggests that degus benefit from plural breeding with communal care only during years of low food availability (Ebensperger et al., 2014). The mechanisms behind

116 this relationship are unknown, however, as several laboratory and field studies have
117 failed to find any direct benefits degus could be deriving from this unique reproductive
118 strategy (Ebensperger et al., 2007; Ebensperger et al., 2011; Ebensperger et al., 2015;
119 Hayes et al., 2009). One possible benefit degus may receive from plural breeding with
120 communal care is a “buffering effect” against post-natal stress; if a chronically stressed
121 mother is not providing sufficient care for her offspring, other mothers within the group
122 may be able to partially compensate.

123 Wild, free-living adult female degus were radio-collared to determine social
124 group membership (Hayes et al., 2009). Directly after parturition, each social group was
125 assigned to one of three treatments: CORT, Mixed, or Control. In CORT groups, all
126 females were implanted with 21-day release cortisol pellets, while 50–75% and 0% of
127 females were implanted in Mixed and Control groups, respectively. When pups emerged
128 from burrows at 4 weeks of age, we sampled them to determine baseline and stress-
129 induced cortisol levels. Additionally, we assessed their negative feedback efficacy via a
130 dexamethasone suppression test and their adrenal sensitivity via an ACTH challenge.

131 The purpose of this study was to address two hierarchical questions and their
132 associated hypotheses: 1) how does post-parturition maternal stress affect the
133 development of the offspring stress response in wild, free-living degus and 2) how does
134 plural breeding with communal care influence these effects of maternal stress on
135 offspring stress response development? If maternal stress affects the development of the
136 offspring stress response (Hypothesis 1), then we predicted that pups from CORT groups
137 would have hyper-reactive stress responses (high stress-induced GCs and weak negative
138 feedback) compared to pups from Control groups (Fig. 1 a–b). If plural breeding with

communal care helps buffer post-natal stress (Hypothesis 2), then we predicted that only pups from CORT groups would have hyper-reactive stress responses compared to pups from Control and Mixed groups (Fig. 1c).

Methods

Study animals

This study was conducted from August 3rd–October 29th, 2011 and August 9th–November 9th, 2012 near Santiago, Chile at Estación Experimental Rinconada de Maipú (33°23'S, 70°31'W, altitude 495 m), a field site owned and maintained by the Universidad de Chile. Although degus are precocial, pups spend the first 4 weeks of life in underground burrows (Bauer et al., 2014). When pups emerge aboveground, they forage near the maternal burrow but will continue nursing for another 1–3 weeks (Becker et al., 2007). Laboratory studies have confirmed that mothers will nurse both their pups and other group member pups (Ebensperger et al., 2002; Ebensperger et al., 2006), as well as provide other forms of maternal care such as licking, grooming, and huddling. While male degus may also provide some of the later forms of care (Braun et al., 2013), their contribution to parental care does not impact offspring quality or survival (Ebensperger et al. 2010).

Social group determination

To capture degus, we used Tomahawk live traps (Tomahawk Live Trap Company, Hazelhurst, WI, USA) baited with plain, rolled oats. Newly captured degus were sexed, ear-tagged for identification purposes, and weighed to the nearest 0.1 g. For female degus, pregnancy status was determined via abdominal palpation. All female degus captured in this study were pregnant and showed signs of lactation after parturition, typical of this field site (Ebensperger et al., 2011).

Degus were then fitted with either 8 g (BR radiocollars, AVM Instrument Co., Colfax, CA, USA) or 7–9 g radiotransmitters (RI-2D, Holohil Systems Limited, Carp, ON, Canada, and SOM-2190A, Wildlife Materials Incorporated, Murphysboro, IL, USA). All radiocollars had unique pulse frequencies that could be tracked with hand-held three-element Yagi antennas (AVM Instrument Co., or Advanced Telemetry Systems, Isanti, MN, USA) fitted with either LA 12-Q (for 150.000-151.999 MHz frequency transmitters; AVM Instrument Co.) or FM-100 receivers (for 164.000-164.999 MHz frequency transmitters; Advanced Telemetry Systems). We used the same radiotelemetry techniques, trapping schemes, and data analysis as described in Hayes *et al.* (2009) to determine social group membership. Briefly, individuals that spent 50% or more of their nights together in the same burrow system were assigned to the same social group. We removed radiocollars immediately prior to parturition as changes in group membership typically occur between mating and early/mid pregnancy.

Cortisol implants

Social groups were divided into three different treatments: Control, Mixed, and
CORT. Control groups ($n = 13$) received no cortisol implants, whereas 50–75% of
females within each Mixed group ($n = 5$) and 100% of females within each CORT group
($n = 9$) were implanted with cortisol pellets. Because social groups from a separate study
were opportunistically used as Control groups, we had more Control groups than Mixed
or CORT groups. Because surgery is a major stressor, we did not implant Control females
and certain Mixed females with placebo implants because the aim of our experiment was
to manipulate stress levels, not solely cortisol levels. Consequently, surgical implantation
of a placebo is the proper control for cortisol increases, but lack of surgery is the proper
control for stress per se. Female degus were implanted 1–2 days after parturition.
Females were anesthetized via inhaled Isoflurane and pellets were implanted on the back
of the neck, under the skin. In 2011, we used 60-day release, 35 mg hydrocortisone
pellets (Innovative Research of America, Sarasota, FL, USA). In 2012, we switched to
using smaller pellets that gave the same daily dose (21-day release, 10 mg hydrocortisone
pellets, Innovative Research of America). We switched to these pellets because we found
in 2011 that pups emerged aboveground at 3–4 weeks of age, and therefore maternal care
after 3 weeks of age was unlikely to have significant effects on HPA-axis development in
pups. We verified that cortisol implants elevated circulating cortisol levels in female
degus 2–4 days after implantation by determining baseline cortisol levels in a sub-sample
of un-implanted and implanted female degus (166 ± 42 vs. 357 ± 93 ng mL⁻¹,
respectively, mean $\pm$ sem; $t_{(13)} = 2.2$, $P = 0.02$). Inducing and/or mimicking chronic stress
in a free-living animal is difficult because chronic stress is more than just GC exposure
and also includes activation of the fight-or-flight response (Cyr and Romero, 2006) and

exposure to CRF and other hormones higher up in the HPA-axis. However, by manipulating cortisol levels, we were able to mimic at least one arm of the chronic stress response.

Blood sampling

Degu pups were trapped and sampled between 0700 and 1300, but time of day does not correlate with baseline cortisol levels in this species (Bauer et al., 2013). One to two observers with binoculars continuously monitored nearby burrow systems so exact time of capture could be determined. Pups were then bled from the saphenous vein into heparinized, microhematocrit capillary tubes for a baseline blood sample (~30 μ L) within 3 min of capture (Romero and Reed, 2005). Pups were then ear-tagged, sexed, and weighed to the nearest 0.1 g. Stress-induced blood samples (~15 μ L) were taken 30 min after capture, immediately after which pups were injected intra-peritoneally with a 1 mg kg^{-1} of body weight dose of dexamethasone (Bauer et al., 2013). Dexamethasone (DEX) is a synthetic glucocorticoid that inhibits endogenous glucocorticoid release and is commonly used to test negative feedback efficacy (Boonstra and Singleton, 1993; Carroll et al., 1981; Romero and Wikelski, 2010; Sapolsky and Altmann, 1991). A blood sample (~30 μ L) was taken 90 min after DEX injection, after which pups were given an intra-peritoneal injection of ACTH (adrenocorticotrophic hormone from porcine pituitary, Sigma Aldrich, St. Louis, Mo., USA) at a dose of 50 IU kg^{-1} of body weight. This dose should induce maximal secretion of cortisol, as it is much higher than that used in most small mammal studies (Boonstra et al., 1998; Boonstra et al., 2008; Boonstra et al., 2011).

A final blood sample (~15 µL) was taken 15 min after ACTH injection. The total amount of blood taken (~90 µL) was below 1% of each individual's body weight, as per the guidelines of the American Society of Mammalogists for the use of wild mammals in research (Sikes et al., 2011). Between blood samples, pups were kept in shaded traps and supplied with oats. Pups were then released at their home burrow system.

Radioimmunoassays

Blood samples were kept in a cooler with ice packs for no more than 9 h and were then centrifuged for 2–3 min at ~230 G. Plasma was drawn off and stored in sealed Eppendorf tubes at –20 °C until further analysis. Samples were measured for cortisol with commercial I¹²⁵ radioimmunoassay kits (Corti-Cote Solid Phase Component System, MP Biomedicals LLC, Irvine, CA, USA) that we previously validated for measuring cortisol in degus (Bauer et al., 2014). Total sample volume was brought up to 50 µL with distilled water and samples were then assayed in duplicate. Assay sensitivity was 0.7 ng mL⁻¹ and intra- and inter-assay variation were 3.8% and 9.6%, respectively.

Statistical analyses

Negative feedback efficacy was calculated as percent decrease from stress-induced cortisol levels after DEX injection, taking into account baseline cortisol levels: (stress-induced cortisol – DEX cortisol) / (stress-induced cortisol – baseline cortisol) * 100. If DEX cortisol levels were higher than stress-induced cortisol levels, then a

negative feedback level of zero was assigned since it is not clear what the biological relevance is of having a response worse than no response. Because it was difficult to get multiple blood samples from pups on cold mornings, we do not have all four samples for each individual pup. Therefore, sample size is not the same for all stress response parameters. Because baseline cortisol, stress-induced cortisol, negative feedback efficacy, and adrenal sensitivity are independently regulated (Reul et al., 1987; Romero, 2006), we analyzed each variable separately. Data were analyzed with mixed model ANOVAs (PROC MIXED) in SAS (Version 9.3; SAS Institute, Cary, North Carolina, USA). We controlled for the effect of social group on pup body mass, baseline cortisol, stress-induced cortisol, negative feedback efficacy, and adrenal sensitivity by including social group as a random variable in the mixed models. We included year, sex, number of adult females within the social group, social group treatment, and sex*treatment as fixed effects. We included year*treatment as a fixed effect because we only had one “Mixed” group in 2011. Effect sizes for significant main fixed factor effects were reported using Cohen’s f^2 , which is considered an appropriate effect size measure for mixed models (Selya et al., 2012). For significant main effects, we tested for pair-wise differences by using Least Squares Means post-hoc tests. Cohen’s d was reported as the effect size for pair-wise differences. For significant interactive effects, we examined simple effects separately and controlled for multiple comparisons using Bonferroni’s alpha adjustment procedure (Pedhazur and Schmelkin, 1991).

Results

Body mass

There was no significant effect of treatment ($F_{2,61} = 0.03$, $P = 0.97$), year ($F_{1,61} = 1.02$, $P = 0.32$), sex ($F_{1,61} = 0.38$, $P = 0.54$), number of adult females ($F_{2,61} = 0.26$, $P = 0.77$), year*treatment ($F_{1,61} = 2.68$, $P = 0.11$), or sex*treatment ($F_{2,61} = 0.64$, $P = 0.53$) on pup body mass.

Baseline cortisol

CORT pups had significantly lower baseline cortisol compared to Control and Mixed pups (Fig. 2a; $F_{2,59} = 4.63$, $P = 0.01$, Cohen's $f^2 = 0.05$, LSM post-hoc: $P < 0.01$, Cohen's $d = -0.60$ and $P = 0.02$, Cohen's $d = -0.76$, respectively). Year had a significant effect on baseline cortisol ($F_{1,59} = 5.95$, $P = 0.02$, Cohen's $f^2 = 0.08$); pups from 2011 had lower baseline cortisol levels compared to pups from 2012 (111 ± 17 vs. 230 ± 37 ng mL⁻¹, respectively, mean $\pm$ sem). However, there was no significant year*treatment effect ($F_{2,59} = 2.52$, $P = 0.09$). There was also no significant effect of sex ($F_{1,59} = 0.80$, $P = 0.38$), number of adult females ($F_{2,59} = 1.84$, $P = 0.17$), or sex*treatment ($F_{2,59} = 3.12$, $P = 0.06$) on pup baseline cortisol levels.

Stress-induced cortisol

There was no significant effect of treatment ($F_{2,61} = 0.88, P = 0.42$), year ($F_{1,61} = 0.71, P = 0.40$), sex ($F_{1,61} = 0.27, P = 0.61$), number of adult females ($F_{2,61} = 0.55, P = 0.58$), year*treatment ($F_{2,61} = 0.81, P = 0.45$), or sex*treatment ($F_{2,61} = 0.69, P = 0.51$) on pup stress-induced cortisol levels (Fig. 2b).

Negative feedback

While negative feedback efficacy was not significantly affected by treatment ($F_{2,51} = 2.20, P = 0.12$), year ($F_{1,51} = 0.12, P = 0.73$), number of adult females ($F_{2,51} = 0.23, P = 0.80$), or year*treatment ($F_{2,51} = 0.28, P = 0.76$), there was a significant effect of sex ($F_{1,51} = 7.70, P < 0.01$, Cohen's $f^2 = 0.14$) and sex*treatment ($F_{2,51} = 5.02, P = 0.01$). When we examined each sex separately, we found a significant treatment effect in males (Fig. 2c; $F_{2,19} = 4.66, P = 0.02$, Cohen's $f^2 = 0.15$) but not females ($F_{2,21} = 0.76, P = 0.48$). Within males, Control and Mixed pups had significantly stronger negative feedback efficacy than CORT pups (LSM post-hoc: $P < 0.05$, Cohen's $d = 0.98$ and $P < 0.01$, Cohen's $d = 1.40$, respectively). When we examined each treatment separately, we found that males had stronger negative feedback only within the Mixed treatment (Fig. 2c; Control: $F_{1,17} = 0.24, P = 0.63$; Mixed: $F_{1,19} = 12.00, P < 0.01$, Cohen's $f^2 = 0.50$; CORT: $F_{1,18} = 1.20, P = 0.29$).

Adrenal sensitivity

Pup cortisol response to ACTH injection was not significantly affected by treatment (Fig. 2d; $F_{2,50} = 0.23$, $P = 0.80$), year ($F_{1,50} = 0.26$, $P = 0.61$), sex ($F_{1,50} = 0.85$, $P = 0.36$), number of adult females ($F_{2,50} = 1.41$, $P = 0.25$), year*treatment ($F_{2,50} = 0.66$, $P = 0.52$), or sex*treatment ($F_{2,50} = 0.87$, $P = 0.42$).

Discussion

The purpose of this study was to determine if post-parturition maternal stress affected the development of the offspring stress response in wild, free-living degus, and whether plural breeding with communal care could help buffer this effect (Fig. 1). Our results provide the first evidence that post-parturition maternal stress can significantly influence the development of the offspring stress response in a wild free-living mammal, and that plural-breeding with communal care may help buffer some of these effects.

Baseline glucocorticoid levels are commonly used as indicators of an animal's health or condition (Bonier et al., 2009). Degu pups from social groups where all mothers were implanted with cortisol pellets (CORT groups) had lower baseline cortisol levels compared to pups from groups where only some or none of the mothers were implanted (Mixed and Control groups). We did not originally predict that baseline cortisol levels would differ between our treatment groups since rat pups (*Rattus norvegicus*) from low licking/grooming arched-back nursing (LG/ABN) mothers had similar baseline GC levels compared to pups from high LG/ABN mothers (Liu et al., 1997). However, laboratory rat studies that directly manipulated GC levels in mothers or pups found different trends. Juvenile rats whose mothers were exposed to GCs after parturition had decreased

baseline GC concentrations (Catalani et al., 1993; McCormick et al., 2001), as did rats whose mothers were implanted with corticosterone pellets 3–18 days after birth (Turner and Taylor, 1976). Our findings may fit better with the latter studies since we directly manipulated adult female degu GC levels in our study whereas Liu *et al.* (1997) used rat dams with naturally varying LG/ABN rates. Our results also fit well with the single, non-rat mammalian study that also examined the effects of maternal stress and care on offspring baseline GC levels. Weaver *et al.* (2000) found that male piglets (*Sus domesticus*) separated from their mothers for 10 min/day over the first 2 weeks of life had lower baseline GC concentrations as adults. Therefore, while we had no a priori predictions concerning pup baseline GC levels, our results generally support that pups from burrows where all mothers were implanted with cortisol pellets had hypo-reactive stress responses, as low baseline GC levels decrease an individual's integrated GC exposure.

Negative feedback is an integral part of the HPA-axis, and hyper-reactive stress responses are often typified by weak negative feedback. Male degu pups from CORT groups had weaker negative feedback compared to male pups from Mixed and Control burrows. These findings are in line with our original predictions and are supported by studies demonstrating that rat pups from low LG/ABN mothers also have weak GC negative feedback (Liu et al., 1997; Weaver et al., 2004). However, rat pups given GC-laced drinking water directly after parturition displayed stronger negative feedback (Catalani et al., 2002). Additionally, in studies where neonates were repeatedly separated from their mothers for short periods of time during post-natal development, maternally deprived rats (Ogawa et al., 1994), squirrel monkeys (*Saimiri sciureus*) (Lyons et al.,

2000), and male piglets (Weaver et al., 2000) had stronger or similar negative feedback efficacy compared to controls. These maternal deprivation and LG/ABN studies may suggest that the development of the post-natal HPA-axis is affected by multiple maternal behaviors and that different maternal care behaviors can have opposing effects.

Altered levels of maternal behaviors may also affect male and female pups differently. Decreased negative feedback efficacy in only male offspring could be attributed to differential care between male and female offspring. Studies in laboratory rats have shown that licking and grooming may not be evenly distributed among all pups within a litter (Cavigelli et al., 2010) and mothers perform more ano-genital licking on male pups compared to female pups (Moore and Morelli, 1979). If degu mothers show the same sex bias in ano-genital licking and grooming, then it is possible that male degu pups are more sensitive to the effects of post-parturition maternal stress. Additionally, Gruss *et al.* (2006) found that when repeatedly separating degu pups from their mothers, males had higher stress-induced cortisol concentrations compared to females during the separations. This may suggest that male degu pups are generally more sensitive to stressors during the first few weeks after parturition.

Although increased GC responses to stress are another indicator of a hyper-reactive HPA-axis (Francis et al., 1999), the laboratory rodent literature does not indicate a consistent response. Rat pups that receive low rates of LG/ABN have higher, stress-induced GC levels (Liu et al., 1997; Weaver et al., 2004), but maternal deprivation studies report higher (Banerjee et al., 2012), lower (Ogawa et al., 1994), or unchanged (Schmidt et al., 2014; Weaver et al., 2000) stress-induced GC levels in offspring subjected to repeated, acute maternal deprivation. Studies that directly manipulated

maternal stress levels also found mixed results, with some showing increased (Levine and Thoman, 1969; McCormick et al., 2001) or decreased (Catalani et al., 2002; Macri et al., 2007) stress-induced GC concentrations in offspring with stressed mothers. Since maternal stress can affect several different maternal behaviors (Brummelte and Galea, 2010), and because different maternal behaviors may have opposing effects on the development of the offspring stress response, it is perhaps not surprising that the data on stress-induced cortisol did not meet our original predictions. Additionally, we found that our treatment groups did not significantly differ in their cortisol response to ACTH injection. We had no a priori predictions concerning this as there are no studies, to our knowledge, that have examined effects of maternal stress or maternal care on adrenal sensitivity in a mammalian species. However, a recent study on song sparrows (*Melospiza melodia*) found that nestlings fed GC-laced food had increased adrenal sensitivity as adults (Schmidt et al., 2014). Future studies are needed to determine if maternal care and stress may affect the development of offspring adrenal sensitivity in other vertebrate species.

Although we did not directly test the impact of our manipulations on offspring fitness, evidence suggests that decreased fitness could result from the changes documented here. Weak negative feedback may significantly affect survival, as studies in Galapagos marine iguanas (*Amblyrhynchus cristatus*) found that individuals with poor negative feedback were less likely to survive long-term food shortages caused by El Niño episodes (Romero and Wikelski, 2010) and studies in snowshoe hares (*Lepus americanus*) found that individuals had weaker negative feedback during population declines (Boonstra et al., 1998). Baseline GC levels may also affect survival rates (Goutte

et al., 2010; Rivers et al., 2012), although there is no consistent pattern across species (Bonier et al., 2009). Furthermore, there is no correlation between fecal glucocorticoid metabolite levels and adult survival rates in degus (Ebensperger et al., 2013). Further studies are needed to determine which aspects of the degu stress response might be correlated with survival and to determine if correlations between survival and the stress response change with different environmental conditions. If maternal stress is an indicator of current environmental conditions, then maternally-derived stress may adaptively program offspring to either increase offspring or maternal fitness (Dantzer et al., 2013; Sheriff and Love, 2013).

Interestingly, plural-breeding with communal care did seem to help buffer effects of post-natal maternal stress (Fig. 1). Our results are consistent with the hypothesis that un-implanted mothers in Mixed groups partially compensate for the lack of care provided by cortisol-implanted mothers. Another prediction of this hypothesis could be that larger groups are more effective at “buffering.” Therefore, while we would have predicted a higher proportion of pups with normal stress responses from larger Mixed groups, the social group size within our Mixed groups (range: 2–4 adult females) did not vary enough to sufficiently test this prediction. Care from other social group members has also been shown to affect the development of the stress response in young marmosets (*Callithrix geoffroyi*) (Birnie et al., 2013), but not in California mice (*Peromyscus californicus*) (Harris et al., 2013). It is tempting to conclude that buffering of maternal stress might be a benefit of plural breeding with communal care in degus. For this to be true, however, changes to the pup HPA-axis induced by maternal stress must be maladaptive so that reversing the changes through buffering would increase fitness. Future studies will need

to show this link in order to support the conclusion that stress buffering is a potential benefit of the degu's unique reproductive strategy.

Because degus live in inaccessible underground burrows, we were never able to verify if our cortisol implants changed the rate or quality of maternal care behaviors. Development of the pup stress response may have been affected by several other mechanisms such as adult inter-female aggression, chemical cues, etc. Additionally, post-parturition maternal stress could affect offspring development via transmission of cortisol during lactation. Laboratory rat studies have shown that maternal GCs can pass into the mother's milk, albeit in very small quantities, and then later absorbed from the pup's stomach into their bloodstream (Zarrow et al., 1970). If cortisol-implanted mothers were significantly affecting pup cortisol levels via lactation, then we would have expected pups from Mixed groups to have stress profiles intermediate to pups from Control and CORT groups. However, we found the opposite, with pups from Mixed groups having stress profiles similar to pups from Control groups. Future studies should test whether maternal stress alters the rate or quality of maternal care in degus, and whether these changes in maternal care are directly responsible for changes in pup HPA-axis development, in addition to exploring other ways maternal stress may impact pup development.

Conclusions

This study demonstrates that maternal stress affects the development of the offspring stress response in wild, free-living degus, and that plural breeding with

communal care may help buffer post-natal stress. This is, to our knowledge, the first time post-parturition maternal stress has been shown to affect the development of the offspring HPA-axis in a wild, free-living animal. Future studies should determine if maternal care is the mechanism through which maternal stress affects HPA-axis development in degus. While we doubt plural breeding with communal care evolved solely to help buffer pups from the negative effects of maternal stress, this could be a lasting benefit and potentially a reason this reproductive strategy has been maintained since original evolutionary pressures vanished.

Acknowledgements

The authors thank A. Bernard, J. Burger, L. Correa, P. Correa Galaz, J. Ho, C. Le, N. Skaff, and J. Trevino for help collecting data. We thank the Universidad de Chile for use of their field site, J.M. Reed for statistical advice, and two anonymous reviewers for their comments on an earlier version of this manuscript. C.M.B. was funded by Sigma Xi, the Tufts University Graduate School, the American Society of Mammalogists, the Animal Behavior Society, NSF DDIG grant (IOS-1210492), the Tufts University International Research Program, and the Tufts Institute of the Environment. L.M.R. was funded by NSF grant IOS-1048529. L.A.E. was funded by FONDECYT grant #1090302 and by the Center for the Advanced Studies in Ecology and Biodiversity (CASEB), Program 1. L.D.H. was funded by an NSF IRES grant (OISE-0853719). This study was approved by the Tufts University Animal Care and Use Committee and the Chilean Servicio Agrícola y Ganadero (permit no. 1-902011 #4731).

References

- Banerjee, S.B., Arterbery, A.S., Fergus, D.J., Adkins-Regan, E., 2012. Deprivation of maternal care has long-lasting consequences for the hypothalamic-pituitary-adrenal axis of zebra finches. *Proc. R. Soc. B-Biol. Sci.* 279, 759-766.
- Bauer, C.M., Hayes, L.D., Ebensperger, L.A., Romero, L.M., 2014. Seasonal variation in the degu (*Octodon degus*) endocrine stress response. *Gen. Comp. Endocrinol.* 197, 26-32.
- Bauer, C.M., Skaff, N.K., Bernard, A.B., Trevino, J.M., Ho, J.M., Romero, L.M., Ebensperger, L.A., Hayes, L.D., 2013. Habitat type influences endocrine stress response in the degu (*Octodon degus*). *Gen. Comp. Endocrinol.* 186, 136.
- Becker, M.I., De Ioannes, A.E., Leon, C., Ebensperger, L.A., 2007. Females of the communally breeding rodent, *Octodon degus*, transfer antibodies to their offspring during pregnancy and lactation. *J. Reprod. Immunol.* 74, 68-77.
- Birnie, A.K., Taylor, J.H., Cavanaugh, J., French, J.A., 2013. Quality of maternal and paternal care predicts later stress reactivity in the cooperatively-breeding marmoset (*Callithrix geoffroyi*). *Psychoneuroendocrinology* 38, 3003-3014.
- Bonier, F., Martin, P.R., Moore, I.T., Wingfield, J.C., 2009. Do baseline glucocorticoids predict fitness?. *Trends in Ecology & Evolution* 24, 634-642.
- Boonstra, R., Hik, D., Singleton, G., Tinnikov, A., 1998. The impact of predator-induced stress on the snowshoe hare cycle. *Ecol. Monogr.* 68, 371-394.

503 Boonstra, R., Singleton, G.R., 1993. Population Declines in the Snowshoe Hare and the
504 Role of Stress. *Gen. Comp. Endocrinol.* 91, 126-143.

505 Boonstra, R., Bradley, A.J., Delehanty, B., 2011. Preparing for hibernation in ground
506 squirrels: adrenal androgen production in summer linked to environmental severity in
507 winter. *Funct. Ecol.* 25, 1348-1359.

508 Boonstra, R., Lane, J.E., Boutin, S., Bradley, A., Desantis, L., Newman, A.E.M., Soma,
509 K.K., 2008. Plasma DHEA levels in wild, territorial red squirrels: Seasonal variation
510 and effect of ACTH. *Gen. Comp. Endocrinol.* 158, 61-67.

511 Branchi, I., Santarelli, S., D'Andrea, I., Alleva, E., 2013. Not all stressors are equal: Early
512 social enrichment favors resilience to social but not physical stress in male mice.
513 *Horm. Behav.* 63, 503-509.

514 Braun, K., Seidel, K., Holetschka, R., Groeger, N., Poeggel, G., 2013. Paternal
515 deprivation alters the development of catecholaminergic innervation in the prefrontal
516 cortex and related limbic brain regions. *Brain Struct. Funct.* 218, 859-872.

517 Brummelte, S., Galea, L.A.M., 2010. Chronic corticosterone during pregnancy and
518 postpartum affects maternal care, cell proliferation and depressive-like behavior in
519 the dam RID A-6775-2010. *Horm. Behav.* 58, 769-779.

520 Carroll, B.J., Feinberg, M., Greden, J.F., Tarika, J., Albala, A.A., Haskett, R.F., James,
521 N.M., Kronfol, Z., Lohr, N., Steiner, M., Devigne, J.P., Young, E., 1981. A Specific
522 Laboratory Test for the Diagnosis of Melancholia - Standardization, Validation, and
523 Clinical Utility. *Arch. Gen. Psychiatry* 38, 15-22.

524 Catalani, A., Casolini, P., Cigliana, G., Scaccianoce, S., Consoli, C., Cinque, C., Zuena,
525 A., Angelucci, L., 2002. Maternal corticosterone influences behavior, stress response

526 and corticosteroid receptors in the female rat. *Pharmacol. Biochem. Behav.* 73, 105-
527 114.

528 Catalani, A., Marinelli, M., Scaccianoce, S., Nicolai, R., Muscolo, L., Porcu, A., Koranyi,
529 L., Piazza, P., Angelucci, L., 1993. Progeny of Mothers Drinking Corticosterone
530 during Lactation has Lower Stress-Induced Corticosterone Secretion and Better
531 Cognitive Performance. *Brain Res.* 624, 209-215.

532 Cavigelli, S.A., Ragan, C.M., Barrett, C.E., Michael, K.C., 2010. Within-litter variance in
533 rat maternal behaviour. *Behav. Processes* 84, 696-704.

534 Cottrell, E.C., Seckl, J.R., 2009. Prenatal stress, glucocorticoids and the programming of
535 adult disease. *Front. Behav. Neurosci.* 3, 19.

536 Cyr, N.E., Romero, L.M., 2006. Chronic stress alters the avian cardiovascular and
537 glucorticoid stress response. *Integrative and Comparative Biology* 46, E30-E30.

538 Dantzer, B., Newman, A.E.M., Boonstra, R., Palme, R., Boutin, S., Humphries, M.M.,
539 McAdam, A.G., 2013. Density Triggers Maternal Hormones That Increase Adaptive
540 Offspring Growth in a Wild Mammal. *Science* 340, 1215-1217.

541 Ebensperger, L.A., Villegas, A., Abades, S., Hayes, L.D., 2014. Mean ecological
542 conditions modulate the effects of group living and communal rearing on offspring
543 production and survival. *Behavioral Ecology*.

544 Ebensperger, L., Veloso, C., Wallem, P., 2002. Do female degus communally nest and
545 nurse their pups?. *J. Ethol.* 20, 143-146.

546 Ebensperger, L.A., Chesh, A.S., Castro, R.A., Ortiz Tolhuysen, L., Quirici, V., Burger,
547 J.R., Sobrero, R., Hayes, L.D., 2011. Burrow limitations and group living in the
548 communally rearing rodent, *Octodon degus*. *J. Mammal.* 92, 21-30.

549 Ebensperger, L.A., Hurtado, M.J., Leon, C., 2007. An experimental examination of the
 550 consequences of communal versus solitary breeding on maternal condition and the
 551 early postnatal growth and survival of degu, *Octodon degus*, pups. *Anim. Behav.* 73,
 552 185-194.

553 Ebensperger, L.A., Hurtado, M.J., Valdivia, I., 2006. Lactating females do not
 554 discriminate between their own young and unrelated pups in the communally
 555 breeding rodent, *Octodon degus*. *Ethology* 112, 921-929.

556 Ebensperger, L.A., León, C., Ramírez-Estrada, J., Abades, S., Hayes, L.D., Nova, E.,
 557 Salazar, F., Bhattacharjee, J., Becker, M.I., 2015. Immunocompetence of breeding
 558 females is sensitive to cortisol levels but not to communal rearing in the degu
 559 (*Octodon degus*). *Physiol. Behav.* 140, 61-70.

560 Ebensperger, L.A., Ramirez-Estrada, J., Leon, C., Castro, R.A., Ortiz Tolhuysen, L.,
 561 Sobrero, R., Quirici, V., Burger, J.R., Soto-Gamboa, M., Hayes, L.D., 2011. Sociality,
 562 glucocorticoids and direct fitness in the communally rearing rodent, *Octodon degus*
 563 RID G-2290-2011. *Horm. Behav.* 60, 346-352.

564 Ebensperger, L.A., Tapia, D., Ramírez-Estrada, J., León, C., Soto-Gamboa, M., Hayes,
 565 L.D., 2013. Fecal cortisol levels predict breeding but not survival of females in the
 566 short-lived rodent, *Octodon degus*. *Gen. Comp. Endocrinol.* 186, 164-171.

567 Francis, D., Diorio, J., Liu, D., Meaney, M., 1999. Nongenomic transmission across
 568 generations of maternal behavior and stress responses in the rat. *Science* 286, 1155-
 569 1158.

570 Goutte, A., Angelier, F., Welcker, J., Moe, B., Clement-Chastel, C., Gabrielsen, G.W.,
 571 Bech, C., Chastel, O., 2010. Long-term survival effect of corticosterone manipulation
 572 in Black-legged kittiwakes. *Gen. Comp. Endocrinol.* 167, 246-251.
 573 Gruss, M., Westphal, S., Luley, C., Braun, K., 2006. Endocrine and behavioural plasticity
 574 in response to juvenile stress in the semi-precocial rodent *Octodon degus*.
 575 *Psychoneuroendocrinology* 31, 361-372.
 576 Harris, B.N., de Jong, T.R., Yang, V., Saltzman, W., 2013. Chronic variable stress in
 577 fathers alters paternal and social behavior but not pup development in the biparental
 578 California mouse (*Peromyscus californicus*). *Horm. Behav.* 64, 799-811.
 579 Hayes, L.D., 2000. To nest communally or not to nest communally: a review of rodent
 580 communal nesting and nursing. *Anim. Behav.* 59, 677-688.
 581 Hayes, L.D., Chesh, A.S., Castro, R.A., Tolhuysen, L.O., Burger, J.R., Bhattacharjee, J.,
 582 Ebensperger, L.A., 2009. Fitness consequences of group living in the degu *Octodon*
 583 *degus*, a plural breeder rodent with communal care. *Anim. Behav.* 78, 131-139.
 584 Hennessy, M., 2003. Enduring maternal influences in a precocial rodent. *Dev. Psychobiol.*
 585 42, 225-236.
 586 Kenagy, G.J., Place, N.J., Veloso, C., 1999. Relation of glucocorticosteroids and
 587 testosterone to the annual cycle of free-living degus in semiarid Central Chile. *Gen.*
 588 *Comp. Endocrinol.* 115, 236-243.
 589 Levine, S., Thoman, E., 1969. Physiological and Behavioral Consequences of Postnatal
 590 Maternal Stress in Rats. *Physiol. Behav.* 4, 139-&.
 591 Lindstrom, J., 1999. Early development and fitness in birds and mammals. *Trends Ecol.*
 592 *Evol.* 14, 343-348.

593 Liu, D., Diorio, J., Tannenbaum, B., Caldji, C., Francis, D., Freedman, A., Sharma, S.,
 594 Pearson, D., Plotsky, P.M., Meaney, M.J., 1997. Maternal care, hippocampal
 595 glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress.
 596 Science 277, 1659-1662.

597 Lyons, D., Yang, C., Mobley, B., Nickerson, J., Schatzberg, A., 2000. Early
 598 environmental regulation of glucocorticoid feedback sensitivity in young adult
 599 monkeys. J. Neuroendocrinol. 12, 723-728.

600 Macri, S., Pasquali, P., Bonsignore, L.T., Pieretti, S., Cirulli, F., Chiarotti, F., Laviola, G.,
 601 2007. Moderate Neonatal Stress Decreases Within-Group Variation in Behavioral,
 602 Immune and HPA Responses in Adult Mice. PLoS One 2, e1015.

603 Matthews, S.G., Phillips, D.I.W., 2010. Minireview: Transgenerational Inheritance of the
 604 Stress Response: A New Frontier in Stress Research. Endocrinology 151, 7-13.

605 McCormick, C., Rioux, T., Fisher, R., Lang, K., MacLaury, K., Teillon, S., 2001. Effects
 606 of neonatal corticosterone treatment on maze performance and HPA axis in juvenile
 607 rats. Physiol. Behav. 74, 371-379.

608 McEwen, B.S., Wingfield, J.C., 2003. The concept of allostasis in biology and
 609 biomedicine. Horm. Behav. 43, 2-15.

610 McEwen, B., 1998. Protective and damaging effects of stress mediators. N. Engl. J. Med.
 611 338, 171-179.

612 Meaney, M.J., Szyf, M., Seckl, J.R., 2007. Epigenetic mechanisms of perinatal
 613 programming of hypothalamic-pituitary-adrenal function and health. Trends Mol.
 614 Med. 13, 269-277.

615 Metcalfe, N.B., Monaghan, P., 2001. Compensation for a bad start: grow now, pay later?.

616 Trends in Ecology & Evolution 16, 254-260.

617 Monaghan, P., Heidinger, B.J., D'Alba, L., Evans, N.P., Spencer, K.A., 2012. For better

618 or worse: reduced adult lifespan following early-life stress is transmitted to breeding

619 partners. Proc. R. Soc. B-Biol. Sci. 279, 709-714.

620 Moore, C., Morelli, G., 1979. Mother Rats Interact Differently with Male and Female

621 Offspring. J. Comp. Physiol. Psychol. 93, 677-684.

622 Mousseau, T.A., Fox, C.W., 1998. The adaptive significance of maternal effects. Trends

623 Ecol. Evol. 13, 403-407.

624 Ogawa, T., Mikuni, M., Kuroda, Y., Muneoka, K., Mori, K., Takahasi, K., 1994. Periodic

625 Maternal-Deprivation Alters Stress-Response in Adult Offspring - Potentiates the

626 Negative Feedback-Regulation of Restraint Stress-Induced Adrenocortical-Response

627 and Reduces the Frequencies of Open Field-Induced Behaviors. Pharmacol. Biochem.

628 Behav. 49, 961-967.

629 Own, L.S., Patel, P.D., 2013. Maternal behavior and offspring resiliency to maternal

630 separation in c57bl/6 mice. Horm. Behav. 63, 411-417.

631 Pedhazur, E.J., Schmelkin, L.P., 1991. Measurement, Design, and Analysis: An

632 Integrated Approach. Lawrence Erlbaum Associates, Hillsdale, NJ.

633 Reul, J., van den Bosch, F., Dekloet, E., 1987. Relative Occupation of Type-i and Type-Ii

634 Corticosteroid Receptors in Rat-Brain Following Stress and Dexamethasone

635 Treatment - Functional Implications. J. Endocrinol. 115, 459-467.

636 Rivers, J.W., Liebl, A.L., Owen, J.C., Martin, L.B., Betts, M.G., 2012. Baseline
637 corticosterone is positively related to juvenile survival in a migrant passerine bird.
638 *Funct. Ecol.* 26, 1127-1134.

639 Romero, L.M., Reed, J.M., 2005. Collecting baseline corticosterone samples in the field:
640 is under 3 min good enough?. *Comparative Biochemistry and Physiology A-
641 Molecular & Integrative Physiology* 140, 73-79.

642 Romero, L.M., 2006. Seasonal changes in hypothalamic-pituitary-adrenal axis sensitivity
643 in free-living house sparrows (*Passer domesticus*). *Gen. Comp. Endocrinol.* 149, 66-
644 71.

645 Romero, L.M., Dickens, M.J., Cyr, N.E., 2009. The reactive scope model - A new model
646 integrating homeostasis, allostasis, and stress. *Horm. Behav.* 55, 375-389.

647 Romero, L.M., Wikelski, M., 2010. Stress physiology as a predictor of survival in
648 Galapagos marine iguanas. *Proceedings of the Royal Society B-Biological Sciences*
649 277, 3157-3162.

650 Saino, N., Bolzern, A.M., Moller, A.P., 1997. Immunocompetence, ornamentation, and
651 viability of male barn swallows (*Hirundo rustica*). *Proc. Natl. Acad. Sci. U. S. A.* 94,
652 549-552.

653 Sapolsky, R.M., Altmann, J., 1991. Incidence of Hypercortisolism and Dexamethasone
654 Resistance Increases with Age among Wild Baboons. *Biol. Psychiatry* 30, 1008-1016.

655 Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How do glucocorticoids influence
656 stress responses? Integrating permissive, suppressive, stimulatory, and preparative
657 actions. *Endocr. Rev.* 21, 55-89.

Schmidt, K.L., MacDougall-Shackleton, E.A., Soma, K.K., MacDougall-Shackleton, S.A., 2014. Developmental programming of the HPA and HPG axes by early-life stress in male and female song sparrows. *Gen. Comp. Endocrinol.* 196, 72-80.

Selye, A.S., Rose, J.S., Dierker, L.C., Hedeker, D., Mermelstein, R.J., 2012. A practical guide to calculating Cohen's $f(2)$, a measure of local effect size, from PROC MIXED. *Front. Psychol.* 3, 111.

Sheriff, M.J., Love, O.P., 2013. Determining the adaptive potential of maternal stress. *Ecol. Lett.* 16, 271-280.

Sikes, R.S., Gannon, W.L., Amer Soc Mammalogists, 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *J. Mammal.* 92, 235-253.

Turner, B., Taylor, A., 1976. Persistent Alteration of Pituitary-Adrenal-Function in Rat by Prepuberal Corticosterone Treatment. *Endocrinology* 98, 1-9.

Weaver, I., Cervoni, N., Champagne, F., D'Alessio, A., Sharma, S., Seckl, J., Dymov, S., Szyf, M., Meaney, M., 2004. Epigenetic programming by maternal behavior. *Nat. Neurosci.* 7, 847-854.

Weaver, S., Aherne, F., Meaney, M., Schaefer, A., Dixon, W., 2000. Neonatal handling permanently alters hypothalamic-pituitary-adrenal axis function, behaviour, and body weight in boars. *J. Endocrinol.* 164, 349-359.

Zarrow, M., Philpott, J., Denenber, V., 1970. Passage of C-14-4-Corticosterone from Rat Mother to Foetus and Neonate. *Nature* 226, 1058-&.

Figure Captions

Fig. 1. Relationship between maternal stress, maternal care, the offspring stress response, and offspring fitness. Hypothesis 1 posits that maternal stress can affect the development of the offspring stress response. If stressed mothers provide low rates of maternal care to offspring, this may cause offspring to develop hyper-reactive stress responses, which may then lead to decreased fitness. Under this hypothesis, we predicted that compared to pups from **a)** Control groups (all mothers unimplanted), pups from **b)** CORT groups (all mothers implanted with CORT) would develop hyper-reactive stress responses. Hypothesis 2 posits that because all mothers within the group provide care for all pups, then the potential consequences of maternal stress are buffered. Under this hypothesis, we predicted within the **c)** Mixed groups (~50% of mothers implanted with CORT), *all* pups should develop normal stress responses and avoid negative fitness consequences. If plural breeding with communal does not help buffer the stress response, then we predicted that ~50% of pups from Mixed groups would have normal stress responses, as those with CORT-implanted mothers would have hyper-reactive stress responses.

Fig. 2. Mean $\pm$ sem **a)** baseline and **b)** stress-induced cortisol levels of degu pups from Control ($n = 35$ and 35), Mixed ($n = 26$ and 27), and CORT ($n = 28$ and 29) treatment groups, respectively. Mean $\pm$ sem **c)** negative feedback efficacy of female and male degu pups from Control ($n = 12$ and 15), Mixed ($n = 14$ and 12), and CORT ($n = 16$ and 12)

704 treatment groups, respectively. Note that a larger decrease indicates stronger negative
705 feedback. Individuals with negative feedback values of less than -100% were able to
706 bring cortisol concentrations below baseline levels after DEX injection. Mean $\pm$ sem **d)**
707 cortisol response to adrenocorticotrophic hormone (ACTH) injection of degu pups from
708 Control ($n = 32$), Mixed ($n = 22$), and CORT ($n = 25$) treatment groups. Different letters
709 indicate significant differences within treatment (but only within males for negative
710 feedback efficacy). Asterisks indicate significant sex differences within treatments.

Figure 1

Figure 2

a) Baseline

b) Stress-induced

c) Negative feedback

d) Adrenal sensitivity