The g -good-neighbor conditional diagnosability of n -dimensional hypercubes under the MM* model *

Shiying Wang¹ Weiping Han

Henan Engineering Laboratory for Big Data Statistical Analysis and Optimal Control, School of Mathematics and Information Science, Henan Normal University, Xinxiang, Henan 453007 PR China

Abstract

Diagnosability of a multiprocessor system is one important study topic. In 2012, Peng et al. proposed a new measure for fault diagnosis of the system, which is called g -good-neighbor conditional diagnosability that restrains every fault-free node containing at least g fault-free neighbors. As a famous topology structure of interconnection networks, the n -dimensional hypercube has many good properties. In this paper, we give the g -good-neighbor conditional diagnosability of the n -dimensional hypercube under the MM* model.

Key words: interconnection network; graph; diagnosability; g -good-neighbor conditional diagnosability; MM* model; hypercube

1. Introduction

As a famous topology structure of interconnection networks, the n -dimensional hypercube Q_n has many good properties, such as connectivity, symmetry, short communication distance, and high fault tolerance of network. Q_n is the first network structure which is taken into practice on IBM machine diagram. With the development of interconnection network topologies, there are many deformation networks which are based on the hypercube, such as the Crossed cube, the Twisted cube and the Möbius cube. As the size

* This work is supported by the National Science Foundation of China (61370001, U1304601)

¹ E-mail address: wangshiying@htu.edu.cn, shiying@sxu.edu.cn (S. Wang)

of multiprocessor systems increases continuously, the possibility of the systems existing more fault processors increases greatly. In order to maintain the reliability of the systems, whenever a processor is found faulty, it should be replaced by a fault-free processor. Several diagnosis models were proposed to identify the faulty processors. One major approach is the Preparata, Metze, and Chien's (PMC) diagnosis model introduced by Preparata et al. [7]. The diagnosis of the system is achieved through two linked processors testing each other. Another important model, namely the comparison diagnosis model (MM model), was proposed by Maeng and Malek [5]. In the MM model, to diagnose a system, a node sends the same task to two of its neighbors, and then compares their responses. A system is t -diagnosable if all faulty processors can be identified without replacement, provided that the number of faults presented does not exceed t . Diagnosability of a system is an important parameter for system diagnosis, which is the maximum value of t such that the system is t -diagnosable. For a t -diagnosable system, Dahbura and Masson [4] proposed an algorithm with time complex $O(n^{2.5})$, which can effectively identify the set of faulty processors.

An n -dimensional hypercube Q_n ($n \geq 1$) is the graph whose vertex set is the set of all n -tuples of 0s and 1s, where two n -tuples are adjacent if they differ in precisely one coordinate. Q_n has $\binom{2^n}{n}$ vertex subsets of size n , among which there are only 2^n vertex subsets which contain all the neighbors of some vertex. Since the ratio $2^n / \binom{2^n}{n}$ is very small for large n , the probability of a fault set with size n containing all the neighbors of any vertex is very low. So Lai et al. [2] introduced the restricted diagnosability of multiprocessor systems called conditional diagnosability. They consider the situation that any fault set cannot contain all the neighbors of any vertex in a system. Inspired by this concept, Peng et al. [6] proposed a new measure for fault diagnosis of the system, namely, g -good-neighbor conditional diagnosability, which requires that every fault-free node contains at least g fault-free neighbors. In [6], they studied the g -good-neighbor conditional diagnosability of the n -dimensional hypercube Q_n under PMC model. Later, Yuan et al. [9] studied that the g -good-neighbor conditional diagnosability of the k -ary n -cube ($k \geq 4$) under the PMC model and MM* model. In this paper, we give that the g -good-neighbor conditional diagnosability of Q_n under the MM* model is $(n - g + 1)2^g - 1$, where $n \geq 5$ and $0 \leq g \leq n - 3$.

2. Terminology and notations

2.1. Terminology

A multiprocessor system is modeled as an undirected simple graph $G = (V, E)$, whose vertices represent processors and edges represent communication links. Given a nonempty vertex subset V' of V , the induced subgraph by V' in G , denoted by $G[V']$, is a graph, whose vertex set is V' and the edge set is the set of all the edges of G with both endpoints in V' . The degree $d_G(v)$ of a vertex v is the number of edges incident with v . We denote by $\delta(G)$ the minimum degrees of vertices of G . For any vertex v , we define the neighborhood $N_G(v)$ of v in G to be the set of vertices adjacent to v . u is called a neighbor or a neighbor vertex of v for $u \in N_G(v)$. Let $S \subseteq V(G)$. We use $N_G(S)$ to denote the set $\cup_{v \in S} N_G(v) \setminus S$, $C_G(S)$ to denote the set $N_G(S) \cup S$. For neighborhoods and degrees, we will usually omit the subscript for the graph when no confusion arises. A graph G is said to be k -regular if for any vertex v , $d_G(v) = k$. The connectivity $\kappa(G)$ of a graph G is the minimum number of vertices whose removal results in a disconnected graph or only one vertex left. We say that the g -good-neighbor property of H , $P_g(H)$ holds for H if and only if every vertex in H has at least g neighbors in H . Let $G = (V, E)$. A fault set $F \subseteq V$ is called a g -good-neighbor conditional faulty set if $|N(v) \cap (V \setminus F)| \geq g$ for every vertex v in $V \setminus F$. A g -good-neighbor conditional cut of a graph G is a g -good-neighbor conditional faulty set F such that $G - F$ is disconnected. The cardinality of the minimum g -good-neighbor conditional cut is said to be the R_g -connectivity of G , denoted by $\kappa^{(g)}(G)$. Let F_1 and F_2 be two distinct subsets of V , and let the symmetric difference $F_1 \triangle F_2 = (F_1 \setminus F_2) \cup (F_2 \setminus F_1)$. For graph-theoretical terminology and notation not defined here we follow [1].

2.2. The MM^* model

In the MM model [5,9], to diagnose a system, a vertex sends the same task to two of its neighbors, and then compares their responses. To be consistent with the MM model, we have the following assumptions.

1. All faults are permanent.

2. A faulty processor produces incorrect outputs for each of its given tasks.
3. The output of a comparison performed by a faulty processor is unreliable.
4. Two faulty processors given the same input and task do not produce the same output.

The comparison scheme of a system G is modeled as a multigraph, denoted by $M(V(G), L)$, where L is the labeled-edge set. A labeled edge $(u, v)_w \in L$ represents a comparison in which two vertices u and v are compared by a vertex w , which implies $uw, vw \in E(G)$. The collection of all comparison results in $M(V(G), L)$ is called the syndrome, denoted by σ^* , of the diagnosis. If the comparison $(u, v)_w$ disagrees, then $\sigma^*((u, v)_w) = 1$; otherwise, $\sigma^*((u, v)_w) = 0$. Hence, a syndrome is a function from L to $\{0, 1\}$. The MM* model is a special case of the MM model. In the MM* model, all comparisons of G are in the comparison scheme of G , i.e., if $uw, vw \in E(G)$, then $(u, v)_w \in L$.

3. Diagnosability of the n -dimensional hypercube under the comparison model

The n -dimensional hypercube Q_n is proposed by Saad and Schultz [8], which is a famous topology structure of interconnection network for multiprocessor systems. Q_n is an undirected n -regular graph containing 2^n vertices, which has the form $u_1u_2 \cdots u_n$, where $u_i \in \{0, 1\}$ for $1 \leq i \leq n$. Two vertices $u = u_1u_2 \cdots u_n$ and $v = v_1v_2 \cdots v_n$ are adjacent if and only if there exists exactly a dimension j such that $u_j \neq v_j$ for every $1 \leq j \leq n$, i.e., $\sum_{i=1}^n |u_i - v_i| = 1$. In this paper, we give the g -good-neighbor conditional diagnosability of Q_n under the MM* model.

Before discussing the g -good-neighbor conditional diagnosability of the n -dimensional hypercube under the MM* model, we first give some useful known theorems.

Theorem 3.1 ([4,9]). *A system G is g -good-neighbor conditional t -diagnosable under the MM* model if and only if for each distinct pair of g -good-neighbor conditional faulty subsets F_1 and F_2 of V with $|F_1| \leq t$ and $|F_2| \leq t$ satisfies one of the following conditions.*

- (1). *There are two vertices $u, w \in V(G) \setminus (F_1 \cup F_2)$ and there is a vertex $v \in F_1 \Delta F_2$ such that $uw \in E$ and $vw \in E$.*
- (2). *There are two vertices $u, v \in F_1 \setminus F_2$ and there is a vertex $w \in V(G) \setminus (F_1 \cup F_2)$*

such that $uw \in E$ and $vw \in E$.

(3). There are two vertices $u, v \in F_2 \setminus F_1$ and there is a vertex $w \in V(G) \setminus (F_1 \cup F_2)$ such that $uw \in E$ and $vw \in E$.

Theorem 3.2 ([6,8]). Let $n \geq 3$ and $0 \leq g \leq n - 2$. Suppose that F is a minimum cardinality cut of Q_n such that $|N_{Q_n}(x) \setminus F| \geq g$ for all $x \in V(Q_n) \setminus F$. Then $|F| = (n - g)2^g$.

Theorem 3.3 ([3]). Let H be a subgraph of Q_n satisfying $P_g(H)$. Then $|V(H)| \geq 2^g$, for $0 < g \leq n$.

Let $X \in \{0, 1\}$, $0^n = \underbrace{00 \cdots 0}_n$, and let $X^n = \underbrace{XX \cdots X}_n$.

Lemma 3.4. Let $A = \{0^{n-g}X^g\}$, $0 \leq g \leq n-3$, and let $F_1 = N_{Q_n}(A)$, $F_2 = A \cup N_{Q_n}(A)$. Then $|F_1| = (n - g)2^g$, $\delta(Q_n - F_1) \geq g$ and $\delta(Q_n - F_2) \geq n - 2$.

Proof. By the definition of Q_n , we have $|A| = 2^g$. Let $0 \leq p \leq n - g - 1$. Then $\{0^p 10^{n-g-1-p}X^g\} \subseteq N_{Q_n}(A)$. Note $|\{0^p 10^{n-g-1-p}X^g\}| = 2^g$. When $p \neq x$, it is easy to see that $0^p 10^{n-g-1-p}X^g \neq 0^x 10^{n-g-1-x}X^g$. Therefore, we have $|F_1| = |N_{Q_n}(A)| = (n - g)2^g$. Note that the form of $N(F_2) = N(A \cup N_{Q_n}(A))$ is $0^p 10^q 10^{n-g-p-q-2}X^g$, where $0 \leq p, q \leq n - g - 2$. Thus, each of $N(F_2)$ is adjacent to two vertices of F_2 . Each of $V(Q_n) \setminus (F_2 \cup N(F_2))$ is not adjacent to any vertex of F_2 . Since Q_n is n -regular, $\delta(Q_n - F_2) \geq n - 2$. By the definition of A , $Q_n - F_1$ has two parts $Q_n[A]$ and $Q_n - F_2$ (for convenience). Note $\delta(Q_n[A]) = g$. Combining this with $0 \leq g \leq n - 3$ and $\delta(Q_n - F_2) \geq n - 2$, we have $\delta(Q_n - F_1) \geq g$. $\square$

Lemma 3.5. Let $n \geq 5$ and $0 \leq g \leq n-3$. Then g -good-neighbor conditional diagnosability of the n -dimensional hypercube Q_n under the MM^* model, $t_g(Q_n) \leq (n - g + 1)2^g - 1$.

Proof. Let A , F_1 and F_2 be defined as above. By Lemma 3.4, $|F_1| = (n - g)2^g$, $|F_2| = (n - g)2^g + 2^g = (n - g + 1)2^g$, $\delta(Q_n - F_1) \geq g$ and $\delta(Q_n - F_2) \geq n - 2$. So both F_1 and F_2 are g -good-neighbor conditional faulty sets. By the definitions of F_1 and F_2 , $F_1 \triangle F_2 = A$. Note $F_1 \setminus F_2 = \emptyset$, $F_2 \setminus F_1 = A$ and $(V(Q_n) \setminus (F_1 \cup F_2)) \cap A = \emptyset$. Therefore, both F_1 and F_2 are not satisfied with any one condition in Theorem 3.1, and Q_n is not g -good-neighbor conditional $(n - g + 1)2^g$ -diagnosable. Hence, $t_g(Q_n) \leq (n - g + 1)2^g - 1$. $\square$

Lemma 3.6. *Let $n \geq 5$ and $0 \leq g \leq n - 3$. Then g -good-neighbor conditional diagnosability of the n -dimensional hypercube under the MM^* model, $t_g(Q_n) \geq (n - g + 1)2^g - 1$.*

Proof. By the definition of g -good-neighbor conditional diagnosability, it is sufficient to show that Q_n is g -good-neighbor conditional $((n - g + 1)2^g - 1)$ -diagnosable. By Theorem 3.1, suppose, on the contrary, that there exist two distinct g -good-neighbor conditional faulty sets F_1 and F_2 with $|F_1| \leq (n - g + 1)2^g - 1$ and $|F_2| \leq (n - g + 1)2^g - 1$, but the vertex set pair (F_1, F_2) is not satisfied with any one condition in Theorem 3.1. Without loss of generality, assume that $F_2 \setminus F_1 \neq \emptyset$. We shall discuss the following two cases.

Case 1. $V(Q_n) = F_1 \cup F_2$.

Let $f(x) = (n - x + 1)2^x$ for $0 \leq x \leq n - 3$. Then $f(x)$ is an increasing function and $n + 1 \leq f(x) \leq 2^{n-1}$. Since $0 \leq g \leq n - 3$, $2^n = |V(Q_n)| = |F_1 \cup F_2| = |F_1| + |F_2| - |F_1 \cap F_2| \leq |F_1| + |F_2| = 2[(n - g + 1)2^g - 1] \leq 2[2^{n-1} - 1] = 2^n - 2 < 2^n$, which is a contradiction.

Case 2. $V(Q_n) \neq F_1 \cup F_2$.

Claim 1. $Q_n - F_1 - F_2$ has no isolated vertex. We shall consider the following two subcases.

Subcase 2.1. $g = 1$.

Suppose, on the contrary, that $Q_n - F_1 - F_2$ has at least one isolated vertex w . Since F_1 is one 1-good neighbor conditional faulty set, there is a vertex $u \in F_2 \setminus F_1$ such that u is adjacent to w . Meanwhile, since the vertex set pair (F_1, F_2) is not satisfied with any one condition in Theorem 3.1. By the condition (3) of Theorem 3.1, there is at most one vertex $u \in F_2 \setminus F_1$ such that u is adjacent to w . Thus, there is just a vertex $u \in F_2 \setminus F_1$ such that u is adjacent to w . Similarly, we can deduce that there is just a vertex $v \in F_1 \setminus F_2$ such that v is adjacent to w when $F_1 \setminus F_2 \neq \emptyset$. Let $W \subseteq V(Q_n) \setminus (F_1 \cup F_2)$ be the set of isolated vertices in $Q_n[V(Q_n) \setminus (F_1 \cup F_2)]$, and let H be the induced subgraph by the vertex set $V(Q_n) \setminus (F_1 \cup F_2 \cup W)$. Then for any $w \in W$, there are $(n - 2)$ neighbors in $F_1 \cap F_2$. Since $|F_2| \leq (n - g + 1)2^g - 1$ and $g = 1$, we have $|F_2| \leq 2n - 1$. Thus, $\sum_{w \in W} |N_{Q_n[(F_1 \cap F_2) \cup W]}(w)| = |W|(n - 2) \leq \sum_{v \in F_1 \cap F_2} d_{Q_n}(v) \leq |F_1 \cap F_2|n \leq (|F_2| - 1)n \leq 2n(n - 1)$. It follows that $|W| \leq \frac{2n(n-1)}{n-2} = 2n + \frac{2n}{n-2} \leq 2n + 4$ when $n \geq 4$. Note $|F_1 \cup F_2| \leq |F_1| + |F_2| - |F_1 \cap F_2| \leq 2(2n - 1) - (n - 2) = 3n$. Assume $H = \emptyset$. Then $2^n = |V(Q_n)| = |F_1 \cup F_2| + |W| \leq 3n + 2n + 4 = 5n + 4$. Let $f(x) = 2^x - 5x - 4$. Then $f(x)$ is

an increasing function and $f(5) = 3$, which contradicts $n \geq 5$. So $H \neq \emptyset$. Since the vertex set pair (F_1, F_2) is not satisfied with the condition (1) of Theorem 3.1, and any vertex of $V(H)$ is not isolated in H , we induce that there is no edge between H and $F_1 \triangle F_2$. Thus, $F_1 \cap F_2$ is a vertex cut of Q_n and $\delta(Q_n - (F_1 \cap F_2)) \geq 1$, i.e., $F_1 \cap F_2$ is one 1-good-neighbor conditional cut of Q_n . By Theorem 3.2, $|F_1 \cap F_2| \geq (n-g)2^g = 2(n-1) = 2n-2$. Because $|F_1| \leq (n-g+1)2^g - 1 = 2n-1$ and $|F_2| \leq (n-g+1)2^g - 1 = 2n-1$, we have neither $F_1 \setminus F_2$ nor $F_2 \setminus F_1$ is empty and $|F_1 \setminus F_2| = |F_2 \setminus F_1| = 1$. Let $F_1 \setminus F_2 = \{v_1\}$ and $F_2 \setminus F_1 = \{v_2\}$. Then for any vertex $w \in W$, w are adjacent to v_1 and v_2 . Since there are at most two common neighbors for any pair of vertices in Q_n , it follows that there are at most two isolated vertices in $Q_n - F_1 - F_2$.

Suppose that there is exactly one isolated vertex v in $Q_n - F_1 - F_2$. Then v_1 and v_2 are adjacent to v and so $N_{Q_n}(v) \setminus \{v_1, v_2\} \subseteq F_1 \cap F_2$. Since Q_n contains no triangle, it follows that $N_{Q_n}(v_1) \setminus \{v\} \subseteq F_1 \cap F_2$, $N_{Q_n}(v_2) \setminus \{v\} \subseteq F_1 \cap F_2$, $[N_{Q_n}(v) \setminus \{v_1, v_2\}] \cap [N_{Q_n}(v_1) \setminus \{v\}] = \emptyset$ and $[N_{Q_n}(v) \setminus \{v_1, v_2\}] \cap [N_{Q_n}(v_2) \setminus \{v\}] = \emptyset$. Since there are at most two common neighbors for any pair of vertices in Q_n , it follows that $|[N_{Q_n}(v_1) \setminus \{v\}] \cap [N_{Q_n}(v_2) \setminus \{v\}]| \leq 1$. Thus, $|F_1 \cap F_2| \geq |N_{Q_n}(v) \setminus \{v_1, v_2\}| + |N_{Q_n}(v_1) \setminus \{v\}| + |N_{Q_n}(v_2) \setminus \{v\}| = (n-2) + (n-1) + (n-1) - 1 = 3n-5$. It follows that $|F_2| = |F_2 \setminus F_1| + |F_1 \cap F_2| \geq 1 + 3n-5 = 3n-4 > 2n-1$ ($n \geq 5$), which contradicts $|F_2| \leq (n-g+1)2^g - 1 = 2n-1$.

Suppose that there is another isolated vertex v' in $Q_n - F_1 - F_2$ and $v' \neq v$. Then v_1 and v_2 are adjacent to v' . Since Q_n contains no triangle and there are at most two common neighbors for any pair of vertices in Q_n , it follows that the four vertex sets $N_{Q_n}(v) \setminus \{v_1, v_2\}$, $N_{Q_n}(v') \setminus \{v_1, v_2\}$, $N_{Q_n}(v_1) \setminus \{v, v'\}$ and $N_{Q_n}(v_2) \setminus \{v, v'\}$ do not pairwise intersect. Therefore, $|F_1 \cap F_2| \geq |N_{Q_n}(v) \setminus \{v_1, v_2\}| + |N_{Q_n}(v') \setminus \{v_1, v_2\}| + |N_{Q_n}(v_1) \setminus \{v, v'\}| + |N_{Q_n}(v_2) \setminus \{v, v'\}| = (n-2) + (n-2) + (n-2) + (n-2) = 4n-8$. It follows that $|F_2| = |F_2 \setminus F_1| + |F_1 \cap F_2| \geq 1 + 4n-8 = 4n-7 > 2n-1$ ($n \geq 5$), which contradicts $|F_2| \leq (n-g+1)2^g - 1 = 2n-1$.

Suppose $F_1 \setminus F_2 = \emptyset$. Then $F_1 \subseteq F_2$. Since F_2 is a 1-good neighbor conditional faulty set, $Q_n - F_2 = Q_n - F_1 - F_2$ has no isolated vertex.

Subcase 2.2. $2 \leq g \leq n-3$.

Since F_1 is a g -good-neighbor conditional faulty set, $|N_{Q_n-F_1}(x)| \geq g$ for any $x \in V(Q_n) \setminus F_1$. As the vertex set pair (F_1, F_2) is not satisfied with any one condition in

Theorem 3.1. By the condition (3) of Theorem 3.1, for any pair of vertices $u, v \in F_2 \setminus F_1$, there is no vertex $w \in V(Q_n) \setminus (F_1 \cup F_2)$ such that $uw \in E(Q_n)$ and $vw \in E(Q_n)$. Therefore, any vertex w in $V(Q_n) \setminus (F_1 \cup F_2)$ has at most one neighbor in $F_2 \setminus F_1$. Thus, for any vertex $w \in V(Q_n) \setminus (F_1 \cup F_2)$, $|N_{Q_n - F_1 - F_2}(w)| \geq g - 1 \geq 1$, i.e., every vertex of $Q_n - F_1 - F_2$ is not an isolated vertex. The proof of Claim is complete.

Let $u \in V(Q_n) \setminus (F_1 \cup F_2)$. By Claim 1, u has at least one neighbor in $Q_n - F_1 - F_2$. Since the vertex set pair (F_1, F_2) is not satisfied with any one condition in Theorem 3.1, by the condition (1) of Theorem 3.1, for any pair of adjacent vertices $u, w \in V(Q_n) \setminus (F_1 \cup F_2)$, there is no vertex $v \in F_1 \triangle F_2$ such that $uw \in E(Q_n)$ and $vw \in E(Q_n)$. It follows that u has no neighbor in $F_1 \triangle F_2$. Since u is arbitrary, there is no edge between $V(Q_n) \setminus (F_1 \cup F_2)$ and $F_1 \triangle F_2$.

Since $F_2 \setminus F_1 \neq \emptyset$ and F_1 is a g -good-neighbor conditional faulty set, $\delta_{Q_n}([F_2 \setminus F_1]) \geq g$. By Theorem 3.3, $|F_2 \setminus F_1| \geq 2^g$ when $0 < g \leq n$. Since both F_1 and F_2 are g -good-neighbor conditional faulty sets and there is no edge between $V(Q_n) \setminus (F_1 \cup F_2)$ and $F_1 \triangle F_2$, $F_1 \cap F_2$ is a g -good-neighbor conditional cut of Q_n . By Theorem 3.2, we have $|F_1 \cap F_2| \geq (n - g)2^g$. Therefore, $|F_2| = |F_2 \setminus F_1| + |F_1 \cap F_2| \geq 2^g + (n - g)2^g = (n - g + 1)2^g$, which contradicts $|F_2| \leq (n - g + 1)2^g - 1$. Suppose that $g = 0$. Since $F_2 \setminus F_1 \neq \emptyset$, $|F_2 \setminus F_1| \geq 1$. Therefore, $|F_2| = |F_2 \setminus F_1| + |F_1 \cap F_2| \geq 1 + (n - 0)2^0 = n + 1$, which contradicts $|F_2| \leq (n - 0 + 1)2^0 - 1 = n$. Therefore, Q_n is g -good-neighbor conditional $((n - g + 1)2^g - 1)$ -diagnosable and hence $t_g(Q_n) \geq (n - g + 1)2^g - 1$. The proof is complete. $\square$

Combining Lemmas 3.5 and 3.6, we have the following theorem.

Theorem 3.7. *Let $n \geq 5$ and $0 \leq g \leq n - 3$. Then g -good-neighbor conditional diagnosability of the n -dimensional hypercube under the MM^* model is $(n - g + 1)2^g - 1$.*

4. Conclusions

In this paper, we investigate the problem of g -good-neighbor conditional diagnosability of the n -dimensional hypercube under the MM^* model. It is proved that the g -good-neighbor conditional diagnosability of the n -dimensional hypercube under the MM^* model is $(n - g + 1)2^g - 1$, where $n \geq 5$ and $0 \leq g \leq n - 3$. The above results show that

the g -good-neighbor conditional diagnosability is several times larger than the classical diagnosability of Q_n depending on the condition g . The work will help engineers to develop more different measures of g -good-neighbor conditional diagnosability based on application environment, network topology, network reliability, and statistics related to fault patterns.

References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory with Applications, The Macmillan Press Ltd, New York, 1976.
- [2] Pao-Lien Lai, Jimmy J.M. Tan, Chien-Ping Chang, Lih-Hsing Hsu, Conditional diagnosability measures for large multiprocessor systems, IEEE Transactions on Computers 54 (2005) 165-175.
- [3] S. Latifi, M. Hedge, M. Naraghi-pour, Conditional connectivity measures for large multiprocessor systems, IEEE Transactions on Computers 43 (1994) 218-222.
- [4] A.T. Dahbura, G.M. Masson, An $O(n^{2.5})$ fault identification algorithm for diagnosable systems, IEEE Transactions on Computers c-33 (6) (1984) 486-492.
- [5] J. Maeng, M. Malek, A comparison connection assignment for self-diagnosis of multiprocessor systems, in: Proceeding of 11th International Symposium on Fault-Tolerant Computing, 1981, pp. 173-175.
- [6] Shao-Lun Peng, Cheng-Kuan Lin, Jimmy J.M. Tan, Lih-Hsing Hsu, The g -good-neighbor conditional diagnosability of hypercube under PMC model, Applied Mathematics and Computation 218 (21) (2012) 10406-10412.
- [7] F.P. Preparata, G. Metze, R.T. Chien, On the connection assignment problem of diagnosable systems, IEEE Transactions on Computers EC-16 (1967) 848-854.
- [8] Youcef Saad, Martin H. Schultz, Topological properties of hypercubes, IEEE Transactions on Computers 37 (1988) 867-872.
- [9] Jun Yuan, Aixia Liu, Xue Ma, Xiuli Liu, Xiao Qin, Jifu Zhang, The g -good-neighbor conditional diagnosability of k -ary n -cubes under the PMC model and MM^* model, IEEE Transactions on Parallel and Distributed Systems 26 (2015) 1165-1177.