Molecular cages from dinuclear ruthenium(II) polypyridyl complexes

Shen Chao^a, Aaron D. W. Kennedy^a, William A. Donald^a, Allan M. Torres^b,
William S. Price^b, Jonathon E. Beves^{a,c*}.

^a School of Chemistry, UNSW Australia, Sydney 2052, Australia

^b School of Science and Health, Western Sydney University, Penrith 2751, Australia

^c Key State Laboratory for Coordination Chemistry, Nanjing University, Nanjing, China

* Corresponding author: j.beves@unsw.edu.au

Abstract

Four dinuclear ruthenium(II) terpyridine complexes containing terminal pyridyl groups were synthesised for assembly into larger heterometallic supramolecular structures. These 4 nm long metalloligands are capable of binding additional metal ions. Combination of tetrakis(acetonitrile)palladium(II) tetrafluoroborate with one of the dinuclear complexes lead to rapid self-assembly of a $[\text{Pd}_2\text{L}_4]^{20+}$ cage, which was characterised using high resolution electrospray mass spectrometry and NMR spectroscopy, including diffusion measurements.

1 Introduction

Metal-ion directed self-assembly methods^[1] have allowed construction of diverse topologies including macrocycles,^[2] interlocked architectures^[1], polygons^[2b] and polyhedra.^[2b, 3] Of particular value are those three-dimensional geometries with an internal cavity such as tetrahedra,^[4] barrels^[5] and spheres.^[6] These internal cavities are capable of remarkable properties,^[7] including stabilising highly reactive species, and catalysing diastereoselective reactions. Most studies have focused on homometallic assemblies with organic ligands bridging the metal centres where the role of the metal ion is limited to that of a structural unit, rather than exploiting the rich chemistry of the metal complexes. Labile metal ions such as palladium(II)^[2, 5c, 8] are typically used for self-assembly to enable a degree of reversibility and therefore error-correction to be built into the resulting structure(s). This lability requirement precludes the use of inert metal ions such as ruthenium(II) or iridium(III) which are enjoying a renaissance in new applications including for photoredox catalysis^[9] and light emitting devices.^[10] Ruthenium(II) polypyridyl complexes have been extensively studied due to their tuneable photophysical properties, stability and relative ease of functionalisation.^[11] Dinuclear,^[11-12] polynuclear^[12b, 13] and polymeric^[14] species containing these subunits have been widely reported, yet examples of discrete 3D assemblies remain relatively rare.^[12a, 15] The introduction of inert metal centres as components of bridging ligands could lead to structures where the catalytic, photoactive or redox properties can be combined with the ability to encapsulate small molecules to result in novel reactions in confined spaces.^[16] The metalloligand,^[17] or expanded ligand^[18] approach has proven highly successful in preparing linker units which retain the metal ion binding properties, while incorporating additional functionality in the ligand. Heteroleptic ruthenium(II) terpyridine complexes have been incorporated into several supramolecular motifs^{[19],[15d, 20]} due to their stability, predictable coordination geometry and ability to form achiral metal complexes, including examples

decorated with pendant pyridyl rings which can act as building blocks for the construction of larger polynuclear assemblies.^[15e] In this study we report the preparation and characterisation of a series of dinuclear ruthenium(II) complexes featuring pendant pyridyl groups and demonstrate their self-assembly to form molecular cages upon reaction with labile palladium(II) ions.

2 Experimental

2.1 General

The numbering scheme adopted for the mononuclear and dinuclear complexes is shown in Scheme 1. ¹H and ¹³C{¹H} NMR spectra were recorded on a Bruker AV-400 spectrometer, Bruker Avance DRX 500, and Bruker Avance 600M spectrometer; The chemical shifts for the ¹H and ¹³C{¹H} NMR spectra are referenced to residual solvent resonance (CD₂HClN ¹H δ 1.94; ¹³C δ 1.32). Coupling constants (*J*) are reported in hertz (Hz). Standard abbreviations indicating multiplicity were used as follows: s = singlet, d = doublet, t = triplet, dd = double doublet, m = multiplet, ddd = doublet of doublets of doublets. ¹H and ¹³C{¹H} NMR assignments were made using 2D-NMR methods (COSY, NOESY, HSQC, HMBC) and are unambiguous unless stated otherwise. Complexes **5-8** were prepared according to previously reported methods.^[21]

Low resolution ESI-MS measurements were performed using a Thermo LCQ Fleet Ion Trap mass spectrometer. High-resolution ESI-MS were performed on a hybrid linear quadrupole ion trap mass spectrometer coupled to a high performance orbitrap mass spectrometer (Thermo LTQ Orbitrap XL) equipped with an external electrospray ionization (ESI) source. To generate intact ions of the molecular cages, ESI solutions containing ca. 1 mg/mL of the cages were used. ESI was initiated and maintained by applying a voltage of 2.5 kV to the ESI emitter relative to the capillary entrance to the mass spectrometer. ESI solutions were infused into the ion source at 5 μL min⁻¹ and the temperature of the capillary entrance to the mass spectrometer was reduced to below 100°C. These conditions were selected to minimize the dissociation of highly charged non-covalently bound assemblies during the ion formation and transfer processes prior to mass analysis using the orbitrap mass spectrometer.

NMR diffusion experiments were performed on a Bruker Avance 400 spectrometer using a 5 mm BBO probe with single axis z-gradient capable of achieving a maximum gradient strength of 54.7 G/cm. In all experiments the pulsed gradient stimulated-echo pulse sequence (PGSTE)^[22] was used and probe temperature maintained at 298 K. The diffusion delays Δ were set to 60 ms and the gradient pulse δ fixed to 2 ms. Samples were dispensed in both regular 5 mm NMR tubes and 2 mm capillary NMR tubes (Wilmad WG-1364-2) which were then inserted and centered in a regular 5mm NMR tubes. These capillary tubes have internal diameter (ID) of 1.6 mm, outer diameter (OD) of 2.0 mm and length of 100 mm. The 2mm capillary tubes were used to eliminate convection effects that were observed in CD₃CN samples in 5 mm tubes.

2.2 Synthesis of dinuclear complexes

2.2.1 Synthesis of complex **1**(PF₆)₄

Complex **5** (50 mg, 0.047 mmol), Cs₂CO₃ (310 mg, 0.95 mmol), 1,3-dibromobenzene (2.8 mg,

0.012 mmol) were combined in an oven-dried Schlenk flask with DMF (5 mL). The solution was degassed via three freeze-pump thaw cycles, and Pd(PPh₃)₄ (11 mg, 0.0094 mmol) was added. The reaction was heated at 80 °C under Ar for 48 h, resulting in a blood-red solution. Excess saturated aqueous NH₄PF₆ solution was added to give a red solid. The precipitate was collected on Celite and washed with H₂O (2 x 5 mL), ethanol (5 mL) and ether (5 mL). The red solid was dissolved in CH₃CN and purified by column chromatography (SiO₂; CH₃CN/H₂O/sat. aq. KNO₃/TEA (14:2:1:0.02)). Addition of excess aqueous NH₄PF₆ and removal of CH₃CN gave a red precipitate which was collected and washed with H₂O (2 x 5 mL), ethanol (5 mL) and ether (5 mL). The precipitate was redissolved in CH₃CN and the solvent removed to give the product as a red solid (Yield: 9.7 mg, 0.0054 mmol, 45%). ¹H NMR (400 MHz, CD₃CN) δ 9.41 (d, *J* = 2.3 Hz, 2H, H^{C2}), 9.13 (s, 4H, H^{B3'}), 9.07 (s, 4H, H^{B3}), 8.86 (dd, *J* = 4.8, 1.4 Hz, 2H, H^{C6}), 8.72 (dt, *J* = 8.0, 1.1 Hz, 4H, H^{A6/A6'}), 8.67 (dt, *J* = 8.0, 1.1 Hz, 4H, H^{A6/A6'}), 8.56 (ddd, *J* = 7.9, 2.5, 1.6 Hz, 2H, H^{C4}), 8.41 (d, *J* = 8.4 Hz, 4H, H^{D1}), 8.31 (t, *J* = 1.7 Hz, 1H, H^{E4}), 8.22 (d, *J* = 8.4 Hz, 4H, H^{D2}), 8.03 – 7.93 (m, 10H, H^{A5+A5'+E2}), 7.83 – 7.70 (m, 3H, H^{E1+C5}), 7.49 (ddd, *J* = 5.6, 1.5, 0.7 Hz, 4H, H^{A3/A3'}), 7.45 (ddd, *J* = 5.6, 1.5, 0.7 Hz, 4H, H^{A3/A3'}), 7.18–7.25 (m, H^{A4+A4'}). ¹³C NMR (151 MHz, CD₃CN) δ 159.2 (C^{A2/A2'}), 159.1 (C^{A2/A2'}), 156.7 (C^{B2/B2'}), 156.5 (C^{B2/B2'}), 153.51 (C^{A6/A6'}), 153.46 (C^{A6/A6'}), 152.2 (C^{C6}), 149.7 (C^{C2}), 148.8 (C^{D6}), 146.3 (C^{B4}), 143.6 (C^{B4'}), 141.6 (C^{E3}), 139.2 (C^{A4/A4'}), 139.1 (C^{A4/A4'}), 137.0 (C^{C1}), 136.2 (C^{C4}), 131.0 (C^{D5}), 129.4 (C^{D1/D2}), 129.4 (C^{D1/D2}), 128.6 (C^{A5/A5'}), 128.5 (C^{A5/A5'}), 126.9 (C^{E2}), 125.64 (C^{A3/A3'}), 125.63 (C^{A3/A3'}), 125.3 (C^{E4}), 122.8 (C^{B3}), 122.5 (C^{B3'}). LR ESI-MS (CH₃CN) *m/z* 903.25 [**M** – 2(PF₆)]²⁺ (calc. 903.13), 553.58 [**M** – 3(PF₆)]³⁺ (calc. 553.76), 379.25 [**M** – 4PF₆]⁴⁺ (calc. 379.08).

2.2.2 Synthesis of 2(PF₆)₄

This complex was prepared following the same conditions as **1** starting from complex **6** (Yield: 9.5 mg, 0.0053 mmol, 44%). ¹H NMR (400 MHz, CD₃CN) δ 9.15 (s, 4H, H^{B3'}), 9.10 (s, 4H, H^{B3}), 9.04 – 8.95 (m, 4H, H^{C2}), 8.74 (dt, *J* = 8.1, 1.1 Hz, 4H, H^{A6/A6'}), 8.71 (dt, *J* = 8.1, 1.1 Hz, 4H, H^{A6/A6'}), 8.44 (d, *J* = 8.5 Hz, 4H, H^{D1}), 8.34 (t, *J* = 1.7 Hz, 1H, H^{E4}), 8.25 (d, *J* = 8.4 Hz, 4H, H^{D2}), 8.21 – 8.14 (m, 4H, H^{C3}), 8.05–7.96 (m, 10H, H^{A5+A5'+E2}), 7.81 (t, *J* = 7.7 Hz, 1H, H^{E1}), 7.53 (ddd, *J* = 5.6, 1.5, 0.7 Hz, 4H, H^{A3/A3'}), 7.46 (ddd, *J* = 5.6, 1.5, 0.7 Hz, 4H, H^{A3/A3'}), 7.24 (m, 8H, H^{A4+A4'}). ¹³C NMR (151 MHz, CD₃CN) δ 159.2 (C^{A2/A2'}), 159.0 (C^{A2/A2'}), 156.9 (C^{B2/B2'}), 156.4 (C^{B2/B2'}), 153.5 (C^{A6/A6'}), 153.4 (C^{A6/A6'}), 152.1 (C^{C2}), 149.4 (C^{D6}), 146.3 (C^{B4}), 145.1 (C^{C4}), 143.2 (C^{B4'}), 142.2 (C^{E3}), 139.1 (C^{A4+A4'}), 138.6 (C^{C1}), 131.3 (C^{E1}), 128.6 (C^{D1/D2}), 128.5 (C^{D1/D2}), 128.1 (C^{A5+A5'}), 127.7 (C^{E2}), 127.3 (C^{E4}), 125.7 (C^{A3/A3'}), 125.7 (C^{A3/A3'}), 123.0 (C^{C3}), 122.9 (C^{B3}), 122.8 (C^{B3'}). LR ESI-MS (CH₃CN) *m/z* 903.26 [**M** – 2(PF₆)]²⁺ (calc. 903.13), 553.84 [**M** – 3(PF₆)]³⁺ (calc. 553.76), 379.20 [**M** – 4PF₆]⁴⁺ (calc. 379.08).

2.2.3 Synthesis of 3(PF₆)₄

This complex was prepared following the same procedure as **1** starting from complex **7** (Yield: 8.2 mg, 0.0046 mmol, 38%). ¹H NMR (400 MHz, CD₃CN) δ 9.40 (dd, *J* = 2.5, 0.9 Hz, 2H, H^{C2}), 9.16 (s, 4H, H^{B3'}), 9.05 (s, 4H, H^{B3}), 8.86 (dd, *J* = 4.8, 1.6 Hz, 2H, H^{C6}), 8.72 (dt, *J* = 8.1, 1.3, 4H, H^{A6/A6'}), 8.67 – 8.64 (m, 6H, H^{D1+A6/A6'}), 8.55 (ddd, *J* = 8.0, 2.5, 1.6 Hz, 2H, H^{C4}), 8.42 (t, *J* = 1.7 Hz, 1H, H^{E1}), 8.27 (dt, *J* = 7.8, 2H, H^{D5}), 8.14 (d, *J* = 7.8, 2H, H^{D3}), 8.05 (dd, *J* = 7.8, 1.8 Hz, 2H, H^{E3}), 7.97 – 7.90 (m, 10H, H^{A5/A5'+D4}), 7.83 (t, *J* = 7.7 Hz, 1H, H^{E4}), 7.74 (ddd, *J* = 8.0, 4.8, 0.9 Hz, 2H, H^{C5}), 7.47 (ddd, *J* = 5.6, 1.5 Hz, 0.8 Hz, 4H, H^{A3/A3'}), 7.42 (ddd, *J* = 5.6, 1.5 Hz, 0.8 Hz, 4H, H^{A3/A3'}), 7.17 (ddd, *J* = 7.6, 5.6, 1.3 Hz, 8H, H^{A4/A4'}). ¹³C NMR (151 MHz, CD₃CN) δ 159.2 (C^{A2/A2'}), 159.1 (C^{A2/A2'}), 156.7 (C^{B2/B2'}), 156.4 (C^{B2/B2'}), 153.5 (C^{A6/A6'}), 153.4 (C^{A6/A6'}), 152.2 (C^{C6}), 149.7 (C^{C2}), 149.26 (C^{D6}), 146.33 (C^{B4}), 143.20 (C^{E2}), 142.2 (C^{B4'}), 139.1 (C^{A4/A4'}), 139.1 (C^{A4/A4'}), 138.7 (C^{D2}), 136.2 (C^{C4}), 133.7 (C^{C3}), 131.3 (C^{D4}), 131.0 (C^{E4}), 130.3 (C^{D3}), 128.6 (C^{A5/A5'}), 128.5 (C^{A5/A5'}), 128.1 (C^{E3+D5}), 127.68 (C^{D1}), 127.3 (C^{E1}), 125.6

(C^{A3+A3'}), 125.3 (C^{C5}), 123.0 (C^{B3}), 122.8 (C^{B3'}). LR ESI-MS (CH₃CN) *m/z* 903.13 [**M** – 2(PF₆)]²⁺ (calc. 903.13), 553.92 [**M** – 3(PF₆)]³⁺ (calc. 553.76), 379.25 [**M** – 4PF₆]⁴⁺ (calc. 379.08).

2.2.4 Synthesis of **4**(PF₆)₄

This complex was prepared following the same procedure as **1** starting from complex **8** (Yield: 8.7 mg, 0.0048 mmol, 40%). ¹H NMR (400 MHz, CD₃CN) δ 9.15 (s, 4H, H^{B3'}), 9.06 (s, 4H, H^{B3}), 9.00 – 8.93 (m, 4H, H^{C2}), 8.71 (dt, *J* = 8.3, 0.9 Hz, 4H, H^{A6/A6'}), 8.67 (dt, 7.9, 0.8 Hz, 4H, H^{A6/A6'}), 8.63 (dd, *J* = 1.7, 1.7 Hz, 2H, H^{D1}), 8.41 (t, *J* = 1.6 Hz, 1H, H^{E1}), 8.27 (d, *J* = 8.6 Hz, 2H, H^{D5}), 8.18 – 8.10 (m, 6H, H^{D3+C3}), 8.05 (dd, *J* = 7.7, 1.8 Hz, 2H, H^{E3}), 7.98 – 7.89 (m, 10H, H^{A5+A5'+D4}), 7.86 (t, *J* = 7.8 Hz, 1H, H^{E4}), 7.47 (ddd, *J* = 5.7, 1.6, 0.8 Hz, 4H, H^{A3/A3'}), 7.42 (ddd, *J* = 5.6, 1.6, 0.7 Hz, 4H, H^{A3/A3'}), 7.17 (m, 8H, H^{A4+A4'}). ¹³C NMR (151 MHz, CD₃CN) δ 159.1 (C^{A2/A2'}), 159.0 (C^{A2/A2'}), 156.9 (C^{B2/B2'}), 156.4 (C^{B2/B2'}), 153.54 (C^{A6/A6'}), 153.45 (C^{A6/A6'}), 152.1 (C^{C2}), 148.9 (C^{D6}), 146.3 (C^{B4}), 145.1 (C^{C4}), 143.6 (C^{E2}), 141.6 (C^{B4'}), 139.2 (C^{A4+A4'}), 137.0 (C^{D2}), 131.0 (C^{D3}), 129.42 (C^{A5/A5'}), 129.37 (C^{A5/A5'}), 128.7 (C^{E3/D5}), 128.5 (C^{E3/D5}), 128.1 (C^{D1}), 126.9 (C^{E1}), 125.7 (C^{A3/A3'}), 125.6 (C^{A3/A3'}), 122.9 (C^{B3'}), 122.8 (C^{C3}), 122.6 (C^{B3}). LR ESI-MS (CH₃CN) *m/z* 903.33 [**M** – 2(PF₆)]²⁺ (calc. 903.13), 553.76 [**M** – 3(PF₆)]³⁺ (calc. 553.76), 379.25 [**M** – 4PF₆]⁴⁺ (calc. 379.08).

2.3 Synthesis of Pd₂(**4**)₄(PF₆)_x(BF₄)_{20-x}

Self-assembled complex [Pd₂(**4**)₄]²⁰⁺ was synthesised by combination of [Pd(CH₃CN)₄](BF₄)₂ (0.81 mg, 1.8 μmol, 0.6 eq.) and complex **4** (6.4 mg, 3.0 μmol, 1 eq.) in CD₃CN leading to formation of a deep red solution which was analysed without future purification. ¹H NMR (500 MHz, CD₃CN) δ 9.95 (d, *J* = 6.1 Hz, 16H, H^{C1}), 9.27 (s, 16H, H^{B3'}), 9.09 (s, 16H, H^{B3}), 9.05 (t, *J* = 1.6 Hz, 8H, H^{D1}), 8.84 (t, *J* = 1.9 Hz, 4H, H^{E1}), 8.79 – 8.74 (m, 16H, H^{C2}), 8.64 (d, *J* = 8.0 Hz, 16H, H^{A6}), 8.60 (d, *J* = 8.1 Hz, 16H, H^{A6'}), 8.38 (dt, *J* = 7.7, 1.3 Hz, 8H, H^{D5}), 8.15 (dt, *J* = 8.2, 1.2 Hz, 8H, H^{D3}), 8.03 (dd, *J* = 7.9, 1.8 Hz, 8H, H^{E3}), 7.95 (t, *J* = 7.8 Hz, 8H, H^{D4}), 7.82 (t, *J* = 7.9 Hz, 4H, H^{E4}), 7.72 (td, *J* = 7.8, 1.5 Hz, 16H, H^{A5}), 7.47 (dt, *J* = 5.6, 1.1 Hz, 16H, H^{A3}), 7.36 (td, *J* = 8.0, 1.6 Hz, 16H, H^{A5'}), 7.25 (dt, *J* = 5.6, 1.1 Hz, 16H, H^{A3'}), 6.89 (ddd, *J* = 7.2, 5.5, 1.3 Hz, 16H, H^{A4}), 6.44 (ddd, *J* = 7.1, 5.7, 1.0 Hz, 16H, H^{A4'}). ¹³C NMR (151 MHz, CD₃CN) δ 158.8 (C^{A2+A2'}), 156.8 (C^{B2/B2'}), 156.0 (C^{B2/B2'}), 153.6 (C^{A3'}), 153.3 (C^{A3}), 152.9 (C^{C1}), 150.8 (C^{B4}), 149.2 (C^{B4'}), 143.5 (C^{C3}), 142.9 (C^{D2}), 141.7 (C^{E2}), 138.9 (C^{A5}), 138.5 (C^{A5'}), 137.8 (C^{D6}), 131.4 (C^{D4}), 131.1 (C^{E4}), 130.4 (C^{D3}), 128.5 (C^{A4}), 128.0 (C^{D1+E3}), 127.8 (C^{A4'}), 127.6 (C^{C2}), 127.2 (C^{D5}), 126.9 (C^{E1}), 125.9 (C^{A6}), 125.5 (C^{A6'}), 123.5 (C^{B3}), 122.5 (C^{B3'}). ESI-MS characterization given in Figure S3-8.

3 Results and Discussion

3.1 Dinuclear ruthenium(II) complexes

Following procedures we have previously optimised,^[21] the dinuclear complexes **1–4** were synthesised by Suzuki coupling of the appropriate boronic acid functionalised ruthenium(II) complex (**5–8**) and 1,3-dibromobenzene with Pd(PPh₃)₄ as the palladium source and Cs₂CO₃ as the base (Scheme 1). The complexes were purified by column chromatography and isolated in moderate yields (38–45%). Protonation of the pendant pyridine rings^[23] leads to highly polar products which are difficult to recover from silica, which contributes to the low isolated yields.

Scheme 1 Synthesis of ruthenium(II) dinuclear complexes. (i) (a): $\text{Pd}(\text{PPh}_3)_4$, Cs_2CO_3 , DMF, 80 °C, 2d (b) aq. NH_4PF_6 .

As a representative example, Figure 1 shows a comparison of the ^1H NMR spectra (acetonitrile- d_3) of the mononuclear complex **5** and dinuclear complex **1**. The signal corresponding to the terminal $\text{B}(\text{OH})_2$ at 6.28 ppm is not present in the coupled product and the signals corresponding to the terpyridine groups are more clearly separated in the dinuclear complexes (notably the $\text{H}^{\text{B}3}$ and $\text{H}^{\text{B}3'}$ signals). Similar changes in ^1H NMR spectra were observed for the other isomers, which are shown in Figure 2. All ^1H and ^{13}C NMR signals were assigned unambiguously using 2D homo- and heteronuclear techniques, except where specified; see Figure S9 for assigned $^{13}\text{C}\{^1\text{H}\}$ NMR spectra.

Figure 1 ^1H NMR spectra (400 MHz, CD_3CN , 298 K) of (a) mononuclear complex **5** and (b) dinuclear complex **1**. Red: terpyridine protons. Black: phenyl protons.

Comparing the ^1H NMR spectra (Figure 2) of complexes **1–4** allows some general trends to be observed. Firstly, signals corresponding to the $\text{H}^{\text{A}4}$, $\text{H}^{\text{A}5}$ and $\text{H}^{\text{A}6}$ protons (see Scheme 1 for atom labelling) are distinguishable in the non-equivalent terpyridine ligands, but have very similar chemical environments. The signals corresponding to proton $\text{H}^{\text{B}3}$ are more sensitive to substitution at the 4'-position, as expected. The signals corresponding to protons on the pendant pyridyl rings (3-pyridyl or 4-pyridyl) are unaffected by the substitution at the phenyl linker at the centre of the complex, as would also be expected. Similarly, the signals corresponding to the central linker group (1,3:1'3' or 1,4:1',3'-triphenyl substitution) are essentially unaffected by the substitution of the pyridyl rings. All dinuclear complexes were analysed by electrospray

ionization mass spectrometry (ESI-MS) and in each case signals corresponding to sequential loss of PF_6^- anions (i.e. $[\text{M} - n(\text{PF}_6^-)]^{n+}$ where $2 \leq n \leq 4$) were observed with the simulated isotopes patterns agreeing well with experiment (Tables S1-4).

Figure 2 ^1H NMR (400 MHz, CD_3CN , 298 K) spectra of dinuclear complexes **1-4** (a-d), Red: terpyridine protons. Blue: terminal pyridine protons. Black: phenyl protons.

3.2 Larger assemblies with palladium(II)

Having synthesised the dinuclear complexes we turned our attention to the self-assembly of larger structures. Reaction of $\text{Pd}(\text{NO}_3)_2$ or $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2$ (0.6 eq.) with any of the dinuclear complexes **1-4** in $\text{DMSO}-d_6$ ^[6b, 24] for 12h at 80 °C did not result in significant changes in the ^1H NMR spectra. Significant precipitation occurred, presumably a metallopolymer formed as the kinetic product of the reaction. Reaction of $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2$ (0.6 eq.) with complex **1-3** in acetonitrile- d_3 , as previously described,^[5a, 25] resulted in only minor ^1H NMR peak shifts and significant precipitation, presumably of polymeric material. However, reaction of $[\text{Pd}(\text{CH}_3\text{CN})_4](\text{BF}_4)_2$ (0.6 eq.) with complex **4** in acetonitrile- d_3 led to a rapid change in the ^1H NMR spectra (Figure 3), to give a single clean product. Firstly, both terminal pyridine signals are shifted downfield, as expected^[5c] upon coordination to a palladium(II) ion. All terpyridine signals shift upfield to some extent, consistent with being encapsulated in the shielding environment of a larger structure.^[26] Similar upfield shifts were observed for a related cage complex^[15e] also based on a ruthenium bisterpyridyl core.

Figure 3 ^1H NMR (400 MHz, CD_3CN , 298 K) spectra of cage formation in CD_3CN . (a) Dinuclear complex **4**; (b) Self-assembled cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$.

The resulting structure is of high symmetry, with no change in the number of ^1H NMR signals with respect to the parent dinuclear complex. The lack of significant peak broadening suggests that the ruthenium(II) terpyridine units can rotate freely on the NMR time scale, unlike that seen in related complexes which show restricted rotation.^[15e] Due to the low solvent viscosity, the structure is not slowly tumbling which was supported by measurements of T_1 relaxation times in both the dinuclear complexes and the self-assembled structure (See SI Tables 5-6). Slow tumbling in more viscous solvents commonly leads to broadening of the signals for related supramolecular structures.^[6b, 24]

Significant shifting of the ^1H NMR signals corresponding to the phenyl groups, which are distant from the pendant pyridyl groups, is evidence for the formation of a large self-assembled molecule. Protonation, or coordination of one palladium(II) ion to the terminal pyridine rings would lead to downfield shifting of the resonances on that ring,^[23] however this would not affect distant protons such as those on the A' ring. Similarly, formation of a species without a discrete interior cavity would not show the significant shielding seen. This shielding effect can result due to close proximity to aromatic surfaces, as commonly observed within the solvophobic cavities of molecular cages.^{[15e], [27]}

The relative spatial orientation of the dinuclear complex was examined by selective NOE techniques.^[28] Figure 4 shows the NOE interactions for H^{E1} and H^{E3} in both the dinuclear complex **4** and the cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$. In the free complex **4**, proton H^{E1} shows NOE interactions with both H^{D1} and H^{D3} , reflecting the free rotation about the bond between the D and E rings. Upon formation of the larger structure, the NOE interaction between H^{E1} and H^{D3} is no longer observed, suggesting the complex can no longer freely rotate due to inclusion into a more rigid structure. Similarly, H^{E3} does not show an NOE with H^{D1} but retains the NOE with H^{D3} . This supports the proposed orientation of the phenylene backbone shown in Figure 4. This orientation allows formation of the M_2L_4 model proposed for the self-assembled structure (*vide infra*).

Figure 4 Selective ^1H NOE (500 MHz, CD_3CN , 298 K) spectra of a) Irradiation of H^{E1} in complex **4**; b) ^1H spectrum of complex **4**; c) Irradiation of H^{E3} in complex **4**; d) Irradiation of H^{E1} in cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$; e) ^1H spectrum of cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$; f) Irradiation of H^{E3} in cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$

High resolution electrospray ionisation mass spectrometry (HR ESI-MS) analysis of a concentrated acetonitrile solution of the complex confirmed the formation of a large, highly charged structure (Figure 5). A series of signals were observed corresponding to $[\text{Pd}_2(\mathbf{4})_4]^{n+}$ species where $n = 5-10$, formed by loss of PF_6^- and BF_4^- counterions. The mass spectrum is complicated by the presence of different counter-ions (PF_6^- and BF_4^-), leading to the statistical mixture of $[\text{Pd}_2(\mathbf{4})_4 - x(\text{PF}_6^-) - y(\text{BF}_4^-)]^{n+}$ peaks where $x+y = n$. The spectra are in agreement with simulated isotopic distributions (See SI Figure 3-8) and are consistent with the formation of the $[\text{Pd}_2(\mathbf{4})_4]^{20+}$ cage. Note that peaks corresponding to the parent complex, notably $[\mathbf{4} - 2(\text{PF}_6^-)]^{2+}$ and $[\mathbf{4} - (\text{BF}_4^-) - (\text{PF}_6^-)]^{2+}$ species, are also seen; however, this is attributed to fragmentation in the mass spectrometry conditions.

Figure 5 High resolution ESI-MS (CH_3CN) of $[\text{Pd}_2(\mathbf{4})_2]^{20+}$. M^{n+} peaks correspond to $x + y = n$. Selected expansions are shown for the $y = 0$ case. Black = experimental; Red = Simulated

Diffusion measurements of dinuclear complexes **1–4** and cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$ were performed to determine their hydrodynamic radii.^[22] The use of standard 5 mm NMR tubes resulted in anomalous behaviour, which we attribute to convection due to the low viscosity of acetonitrile. This is most apparent looking at the fit of the data (Figure S17) and identifying the systematic deviation from the fit. This phenomena applies to all diffusion (or DOSY) NMR data recorded in low viscosity solvents, where much care must be taken in interpreting results. This convection was overcome by the use of 2 mm capillaries, which minimise convection and allowed collection of data with a better fit to the expected diffusion function. It is also worth noting that the absolute values obtained differed by approximately 15% compared with data collected in standard NMR tubes.

The calculated diffusion constants are summarised in Table 1. Each of the dinuclear complexes has diffusion coefficients ranging from 5.24 to $5.92 \times 10^{10} \text{ cm}^2 \cdot \text{s}^{-1}$, corresponding to calculated hydrodynamic radii ranging from 1.22 to 1.08 nm respectively. The cage complex $[\text{Pd}_2(\mathbf{4})_4]^{20+}$ has a significantly smaller diffusion coefficient of $3.28 \times 10^{10} \text{ cm}^2 \cdot \text{s}^{-1}$, corresponding to a hydrodynamic radius of 2.0 nm , consistent with the formation of a larger structure.^[29]

Table 1 Diffusion coefficients for selected compounds calculated by NMR.^a

Compound	Diffusion coefficient (D) / $10^{10} \text{ cm}^2 \text{ s}^{-1}$	Hydrodynamic radius / nm
1 ⁴⁺	5.25 ± 0.28	1.22 ± 0.06
2 ⁴⁺	5.47 ± 0.23	1.17 ± 0.05
3 ⁴⁺	5.24 ± 0.19	1.22 ± 0.04
4 ⁴⁺	5.92 ± 0.24	1.08 ± 0.05
$[\text{Pd}_2(\mathbf{4})_4]^{20+}$	3.28 ± 0.22	2.0 ± 0.1

^a CD₃CN, 400 MHz, 298 K. Hydrodynamic radii calculated from the Stokes-Einstein equation. Errors were calculated from standard deviations, see SI for details.

In the absence of crystallographic data, a MMFF (Spartan '14) model of the proposed $[\text{M}_2\text{L}_4]^{20+}$ structure is shown in Figure 6. The dimensions of this rudimentary model are in excellent agreement with the measured diffusion data, with longest diagonal axis of the molecule being 4.1 nm. The orientation of the triphenylene bridge reduces the divergence between the *N*-donor pyridine rings, allowing coordination to form the burger-like architecture shown in Figure 6a, the smallest possible discrete species for palladium(II) and this ligand. Coordination to palladium(II) locks the orientation of the triphenylene bridge, reducing the number of observed NOE interactions for protons H^{E3} and H^{E1}, and the C₄ rotation axis accounts for the symmetry of the ¹H NMR spectrum. The distance between the ruthenium(II) units of the same dinuclear unit (~13 Å) is sufficient to allow unrestricted rotation of the Ru(tpy)₂ units on the NMR timescale, leading to sharp peaks in the ¹H NMR spectrum.

Figure 6 Molecular mechanics model (Spartan 14) of proposed cage $[\text{Pd}_2(\mathbf{4})_4]^{20+}$ showing a) the enclosed cavity and b) the 4-fold symmetry along the Pd-Pd axis. grey: carbon; white: hydrogen; blue: nitrogen; green: ruthenium; magenta: palladium.

4 Conclusion

We have synthesised four dinuclear ruthenium(II)-terpyridine complexes with terminal pyridine units for constructing heterometallic supramolecular structures. Combination of complex **4** with palladium(II) led to the rapid formation of a discrete $[\text{Pd}_2(\mathbf{4})_4]^{20+}$ architecture, established by detailed NOE and diffusion NMR measurements and ESI-MS. This class of Ru₈Pd₂ molecular cage has potential for binding guest molecules within its cavity which is sufficiently large to accommodate a wide range of organic guest molecules.

5 Acknowledgements

We acknowledge Hasti Iranmanesh for her useful discussions and attempts to crystalize these complexes for X-ray studies. This work was supported by the National Science Foundation of China (NSFC) Research Fund for International Young Scientists Project (No. 21450110060), the Australian Research Council [DP160100870] and UNSW Australia.

6 Addendum

During the review of this work, Severin and co-workers reported 3 nm $[\text{Pd}_2\text{L}_4]^{4+}$ cages using dinuclear metalloligands with iron(II) centres.^[30] The synthetically simple, sterically bulky metalloligands which feature hydrophobic regions were shown to be extremely effective in driving the selective cage formation.

7 References

- [1] J. E. Beves, B. A. Blight, C. J. Campbell, D. A. Leigh, R. T. McBurney, *Angew. Chem., Int. Ed.* **2011**, *50*, 9260-9327.
- [2] a) R. Chakrabarty, P. S. Mukherjee, P. J. Stang, *Chem. Rev.* **2011**, *111*, 6810-6918; b) T. R. Cook, P. J. Stang, *Chem. Rev.* **2015**, *115*, 7001-7045.
- [3] a) M. D. Ward, *Chem. Commun.* **2009**, 4487-4499; b) T. K. Ronson, S. Zarra, S. P. Black, J. R. Nitschke, *Chem. Commun.* **2013**, *49*, 2476-2490; c) M. M. J. Smulders, I. A. Riddell, C. Browne, J. R. Nitschke, *Chem. Soc. Rev.* **2013**, *42*, 1728-1754.
- [4] For selected examples of tetrahedral cages, see a) D. L. Caulder, R. E. Powers, T. N. Parac, K. N. Raymond, *Angew. Chem., Int. Ed.* **1998**, *37*, 1840-1843; b) J. S. Fleming, K. L. V. Mann, C.-A. Carraz, E. Psillakis, J. C. Jeffery, J. A. McCleverty, M. D. Ward, *Angew. Chem., Int. Ed.* **1998**, *37*, 1279-1281; c) D. L. Caulder, C. Brückner, R. E. Powers, S. König, T. N. Parac, J. A. Leary, K. N. Raymond, *J. Am. Chem. Soc.* **2001**, *123*, 8923-8938; d) R. L. Paul, Z. R. Bell, J. C. Jeffery, J. A. McCleverty, M. D. Ward, *Proc. Natl Acad. Sci. U. S. A.* **2002**, *99*, 4883-4888; e) J. K. Clegg, L. F. Lindoy, B. Moubaraki, K. S. Murray, J. C. McMurtrie, *Dalton Trans.* **2004**, 2417-2423; f) P. Mal, B. Breiner, K. Rissanen, J. R. Nitschke, *Science* **2009**, *324*, 1697-1699; g) C. R. K. Glasson, J. K. Clegg, J. C. McMurtrie, G. V. Meehan, L. F. Lindoy, C. A. Motti, B. Moubaraki, K. S. Murray, J. D. Cashion, *Chem. Sci.* **2011**, *2*, 540-543; h) R. A. Bilbeisi, T. K. Ronson, J. R. Nitschke, *Angew. Chem., Int. Ed.* **2013**, *52*, 9027-9030.
- [5] For selected examples of barrel structures, see a) G. H. Clever, S. Tashiro, M. Shionoya, *Angew. Chem., Int. Ed.* **2009**, *48*, 7010-7012; b) A. M. Johnson, R. J. Hooley, *Inorg. Chem.* **2011**, *50*, 4671-4673; c) J. E. M. Lewis, E. L. Gavey, S. A. Cameron, J. D. Crowley, *Chem. Sci.* **2012**, *3*, 778-784; d) M. Han, R. Michel, B. He, Y.-S. Chen, D. Stalke, M. John, G. H. Clever, *Angew. Chem., Int. Ed.* **2013**, *52*, 1319-1323; e) M. Han, D. M. Engelhard, G. H. Clever, *Chem. Soc. Rev.* **2014**, *43*, 1848-1860; f) A. Schmidt, A. Casini, F. E. Kühn, *Coord. Chem. Rev.* **2014**, *275*, 19-36.
- [6] For selected examples of spherical cages, see a) M. Tominaga, K. Suzuki, T. Murase, M. Fujita, *J. Am. Chem. Soc.* **2005**, *127*, 11950-11951; b) Q.-F. Sun, J. Iwasa, D. Ogawa, Y. Ishido, S. Sato, T. Ozeki, Y. Sei, K. Yamaguchi, M. Fujita, *Science* **2010**, *328*, 1144-1147; c) K. Harris, D. Fujita, M. Fujita, *Chem. Commun.* **2013**, *49*, 6703-6712; d) Q.-Q. Wang, S. Gonell, S. H. A. M. Leenders, M. Dürr, I. Ivanović-Burmazović, J. N. H. Reek, *Nat. Chem.* **2016**, *8*, 225-230.
- [7] a) Y. Inokuma, M. Kawano, M. Fujita, *Nat. Chem.* **2011**, *3*, 349-358; b) B. Breiner, J.

- K. Clegg, J. R. Nitschke, *Chem. Sci.* **2011**, 2, 51-56.
- [8] For reviews of the use of Pd(II) ions for self-assembly, see Reference 2, 5e and a) N. B. Debata, D. Tripathy, D. K. Chand, *Coord. Chem. Rev.* **2012**, 256, 1831-1945; b) T. R. Cook, Y.-R. Zheng, P. J. Stang, *Chem. Rev.* **2013**, 113, 734-777.
- [9] For select examples of transition metal mediated photoredox catalysis, see a) D. A. Nicewicz, D. W. C. MacMillan, *Science* **2008**, 322, 77-80; b) T. P. Yoon, M. A. Ischay, J. Du, *Nat. Chem.* **2010**, 2, 527-532; c) J. M. R. Narayanam, C. R. J. Stephenson, *Chem. Soc. Rev.* **2011**, 40, 102-113; d) C. K. Prier, D. A. Rankic, D. W. C. MacMillan, *Chem. Rev.* **2013**, 113, 5322-5363.
- [10] S. Ardo, G. J. Meyer, *Chem. Soc. Rev.* **2009**, 38, 115-164.
- [11] S. Campagna, F. Puntoriero, F. Nastasi, G. Bergamini, V. Balzani, in *Photochemistry and Photophysics of Coordination Compounds I*, Vol. 280 (Eds.: V. Balzani, S. Campagna), Springer-Verlag Berlin, Berlin, **2007**, pp. 117-214.
- [12] For selected examples of dinuclear Ru(II) complexes, see a) G. R. Newkome, T. J. Cho, C. N. Moorefield, R. Cush, P. S. Russo, L. A. Godínez, M. J. Saunders, P. Mohapatra, *Chem.-Eur. J.* **2002**, 8, 2946-2954; b) C. Chiorboli, M. T. Indelli, F. Scandola, *Top. Curr. Chem.* **2005**, 257, 63-102.
- [13] For selected examples of polynuclear Ru(II) complexes, see a) S.-S. Sun, A. J. Lees, *Inorg. Chem.* **2001**, 40, 3154-3160; b) J.-L. Wang, X. Li, C. D. Shreiner, X. Lu, C. N. Moorefield, S. R. Tummalapalli, D. A. Medvetz, M. J. Panzner, F. R. Fronczek, C. Wesdemiotis, G. R. Newkome, *New J. Chem.* **2012**, 36, 484-491.
- [14] For selected examples of polymeric Ru(II) polypyridyl complexes, including metal-organic frameworks, see a) V. Balzani, A. Juris, M. Venturi, S. Campagna, S. Serroni, *Chem. Rev.* **1996**, 96, 759-834; b) C. A. Kent, B. P. Mehl, L. Ma, J. M. Papanikolas, T. J. Meyer, W. Lin, *J. Am. Chem. Soc.* **2010**, 132, 12767-12769; c) I. Eryazici, O. K. Farha, O. C. Compton, C. Stern, J. T. Hupp, S. T. Nguyen, *Dalton Trans.* **2011**, 40, 9189-9193; d) A. Kobayashi, T. Ohba, E. Saitoh, Y. Suzuki, S.-i. Noro, H.-C. Chang, M. Kato, *Inorg. Chem.* **2014**, 53, 2910-2921; e) S. Zhang, L. Li, S. Zhao, Z. Sun, J. Luo, *Inorg. Chem.* **2015**, 54, 8375-8379.
- [15] For discrete 3D Ru(II) polypyridyl containing structures, see a) S. Roche, C. Haslam, S. L. Heath, J. A. Thomas, *Chem. Commun.* **1998**, 1681-1682; b) K. Li, L.-Y. Zhang, C. Yan, S.-C. Wei, M. Pan, L. Zhang, C.-Y. Su, *J. Am. Chem. Soc.* **2014**, 136, 4456-4459; c) A. J. Methereell, M. D. Ward, *Chem. Commun.* **2014**, 50, 6330-6332; d) T.-Z. Xie, S.-Y. Liao, K. Guo, X. Lu, X. Dong, M. Huang, C. N. Moorefield, S. Z. D. Cheng, X. Liu, C. Wesdemiotis, G. R. Newkome, *J. Am. Chem. Soc.* **2014**, 136, 8165-8168; e) J. Yang, M. Bhadbhade, W. A. Donald, H. Iranmanesh, E. G. Moore, H. Yan, J. E. Beves, *Chem. Commun.* **2015**, 51, 4465-4468; f) A. J. Methereell, M. D. Ward, *RSC Advances* **2016**, 6, 10750-10762; g) K. Wu, K. Li, Y.-J. Hou, M. Pan, L.-Y. Zhang, L. Chen, C.-Y. Su, *Nat. Commun.* **2016**, 7, 10487.
- [16] S. H. A. M. Leenders, R. Gramage-Doria, B. de Bruin, J. N. H. Reek, *Chem. Soc. Rev.* **2015**, 44, 433-448.
- [17] a) S. R. Halper, L. Do, J. R. Stork, S. M. Cohen, *J. Am. Chem. Soc.* **2006**, 128, 15255-15268; b) E. C. Constable, E. L. Dunphy, C. E. Housecroft, M. Neuburger, S. Schaffner, F. Schaper, S. R. Batten, *Dalton Trans.* **2007**, 4323-4332.
- [18] E. C. Constable, *Coord. Chem. Rev.* **2008**, 252, 842-855.
- [19] E. C. Constable, *Chem. Soc. Rev.* **2007**, 36, 246-253.
- [20] For examples of [Ru(tpy)₂]²⁺ containing macrocycles and related structures, see a) J. D. Crowley, A. J. Goshe, B. Bosnich, *Chem. Commun.* **2003**, 2824-2825; b) T. Schröder, R. Brodbeck, M. C. Letzel, A. Mix, B. Schnatwinkel, M. Tonigold, D. Volkmer, J. Mattay, *Tetrahedron Lett.* **2008**, 49, 5939-5942; c) M. Wang, C. Wang, X.-Q. Hao, X.

- Li, T. J. Vaughn, Y.-Y. Zhang, Y. Yu, Z.-Y. Li, M.-P. Song, H.-B. Yang, X. Li, *J. Am. Chem. Soc.* **2014**, *136*, 10499-10507; d) S.-Y. Wang, J.-H. Fu, Y.-P. Liang, Y.-J. He, Y.-S. Chen, Y.-T. Chan, *J. Am. Chem. Soc.* **2016**, *138*, 3651-3654; e) T.-Z. Xie, K. Guo, J.-Y. Li, B. Zhang, K. Zheng, C. N. Moorefield, M. J. Saunders, K. J. Endres, S. Sallam, C. Wesdemiotis, G. R. Newkome, *J. Inorg. Organomet. Polym. Mater.* **2016**, *26*, 907-913; f) Y. Li, Z. Jiang, J. Yuan, D. Liu, T. Wu, C. N. Moorefield, G. R. Newkome, P. Wang, *Chem. Commun.* **2015**, *51*, 5766-5769; g) Y.-T. Chan, X. Li, C. N. Moorefield, C. Wesdemiotis, G. R. Newkome, *Chem.-Eur. J.* **2011**, *17*, 7750-7754.
- [21] C. Shen, P. Wang, J. E. Beves, *Polyhedron* **2016**, *103*, 241-247
- [22] a) J. E. Tanner, *J. Chem. Phys.* **1970**, *52*, 2523-2526; b) A. M. Torres, R. Dela Cruz, W. S. Price, *J. Magn. Reson.* **2008**, *193*, 311-316.
- [23] J. E. Beves, E. L. Dunphy, E. C. Constable, C. E. Housecroft, C. J. Kepert, M. Neuburger, D. J. Price, S. Schaffner, *Dalton Trans.* **2008**, 386-396.
- [24] K. Suzuki, M. Tominaga, M. Kawano, M. Fujita, *Chem. Commun.* **2009**, 1638-1640.
- [25] D. Preston, J. E. Barnsley, K. C. Gordon, J. D. Crowley, *J. Am. Chem. Soc.* **2016**, *138*, 10578-10585.
- [26] W. Meng, B. Breiner, K. Rissanen, J. D. Thoburn, J. K. Clegg, J. R. Nitschke, *Angew. Chem., Int. Ed.* **2011**, *50*, 3479-3483.
- [27] a) S. M. Biroš, J. J. Rebek, *Chem. Soc. Rev.* **2007**, *36*, 93-104; b) J. Rebek, *Acc. Chem. Res.* **2009**, *42*, 1660-1668.
- [28] K. Stott, J. Keeler, Q. N. Van, A. J. Shaka, *J. Magn. Reson.* **1997**, *125*, 302-324.
- [29] Molecular mechanisms molecular modelling was performed using Spartan '14 to estimate sizes.
- [30] S. M. Jansze, M. D. Wise, A. V. Vologzhanina, R. Scopelliti, K. Severin, *Chem. Sci.* **2017**, DOI: 10.1039/C6SC04732G.