1
2 Walk or ride? Phoretic behavior of amblyceran and ischnoceran lice
3
4

5 Andrew W. Bartlow*, Scott M. Villa, Michael W. Thompson, Sarah E. Bush
6
7

8 *Department of Biology, University of Utah, 257 S. 1400 E., Salt Lake City, Utah, USA 84112*
9
10

11 *Corresponding author. Tel.: 1-801-585-9742.

12 *E-mail address:* andrew.bartlow@utah.edu
13
14
15
16
17
18
19

ABSTRACT

Phoresy is a behavior where one organism hitches a ride on another more mobile organism. This is a common dispersal mechanism among relatively immobile species that specialize on patchy resources. Parasites specialize on patchily distributed resources: their hosts. Although host individuals are isolated in space and time, parasites must transmit between hosts or they will die with their hosts. Lice are permanent obligate ectoparasites that complete their entire life cycle on their host. They typically transmit when hosts come into direct contact; however, lice are also capable of transmitting phoretically. Yet, phoresy is rare among some groups of lice. Fundamental morphological differences have traditionally been used to explain the phoretic differences among different suborders of lice; however, these hypotheses do not fully explain observed patterns. We propose that a more fundamental natural history trait may better explain variation in phoresy. Species able to disperse under their own power should be less likely to engage in phoresy than more immobile species. Here we experimentally tested the relationship between independent louse mobility and phoresy using a system with four species of lice (Phthiraptera: Ischnocera and Amblycera) that all parasitize a single host species, the Rock Pigeon (*Columba livia*). We quantified the relative ability of all four species of lice to move independently off the host, and we quantified their ability to attach to, and remain attached to, hippoboscids flies (*Pseudolynchia canariensis*). Our results show that the most mobile louse species is the least phoretic, and the most phoretic species is quite immobile off the host. Our findings were consistent with the hypothesis that phoretic dispersal should be rare among species of lice that are capable of independent dispersal; however other factors such as interspecific competition may also play a role.

Keywords: Amblycera, Dispersal, Hippoboscids fly, Ischnocera, Lice, Phoresy

1. Introduction

Organisms often specialize on resources that are patchily distributed in space and time (MacArthur and Pianka, 1966). Although patches can be resource-rich, dispersing among these spatially isolated and ephemeral patches can be difficult. This is particularly true of free-living and parasitic organisms that are relatively immobile such as wingless insects, mites and worms. Some organisms have solved this dispersal problem by being phoretic. Phoresy is a behavior where a relatively immobile organism disperses by hitching a ride on another more mobile organism (Farish and Axtell, 1971; Houck and OConnor, 1991).

Phoresy has evolved in several phyla and is relatively common among nematodes, mites, lice, beetles and pseudoscorpions, some of which are obligate parasites or mutualists of vertebrate hosts (Treat, 1956; Keirans, 1975a; Roubik and Wheeler, 1982; Houck and OConnor, 1991; Zeh and Zeh, 1992; Athias-Binche and Morand, 1993). Hosts are patchily distributed because each host individual is, in essence, an island of exploitable resources (Kuris et al., 1980). Moreover, hosts are temporally patchy because all hosts eventually die. Thus, dispersal among host individuals (also referred to as transmission) is critical for the persistence of parasite and mutualist lineages.

Lice (Phthiraptera) are permanent, obligate ectoparasites of birds and mammals. Lice most commonly transmit between hosts when individuals come into direct, physical contact, such as contact between mates and contact between parents and offspring (Rothschild and Clay, 1952; Johnson and Clayton, 2003b; Clayton et al., 2016). However, lice also engage in phoretic transmission. In most cases, lice hitch rides on hippoboscids flies, which are blood-feeding parasites of birds and mammals (Keirans, 1975a; Durden, 1990). Rarely, lice also hitch rides on other insects such as fleas, dragonflies, bees and butterflies (Worth and Patterson, 1960; Keirans, 1975b; Durden, 1990; Kirk-Spriggs and Mey, 2014).

There are three major suborders of lice: Anoplura, Amblycera and Ischnocera (Price et al., 2003). Although there are records of phoretic lice from all three suborders (Table 1), phoresy appears to be exceptionally rare among amblyceran lice (Table 2). To date, there is only one documented case of phoresy among over 1300 species of amblyceran lice (Hopkins, 1946), whereas at least 33 of the more than 3,000 spp. of the suborder Ischnocera are known to engage in phoresy (Table 2). Some species of ischnoceran lice even engage in phoresy quite regularly. Studies by Markov (1938), Edwards (1952), Corbet (1956), Bennett (1961), and Baum (1968) report that 20 – 43.5% of hippoboscids in their field studies carried ischnoceran lice. Moreover, flies frequently carry more than one louse; for example, Peters (1935) found a fly with 31 lice attached.

Keirans (1975a) noted that lice attached to flies with their mandibles, and he hypothesized that the differences in the frequency of phoresy among lice is determined by the ability or inability of lice to grab onto flies with their mouthparts. Lice of the suborder Ischnocera have dorso-ventrally aligned, mandibulate mouthparts that are used to bite or scrape the host's integument (Johnson and Clayton, 2003b). Species of the suborder Ischnocera, which use their mandibles to cling tightly to the hair or feathers of their host, also use their mandibles to grasp setae on the body of hippoboscids (Keirans, 1975a). In contrast, lice of the suborder Amblycera have chewing mouthparts that are essentially modified to suck blood and lice of the suborder Anoplura have piercing, sucking mouthparts for sucking blood (Johnson and Clayton, 2003b).

Despite these extreme morphological differences, anopluran lice are as phoretic as ischnocerans (Table 2). Instead of using mouthparts, anopluran lice cling to flies with their tarsal claws (Mitzmain, 1912; Allingham, 1987; Durden, 1990). Recent phoretic records indicate that some ischnocerans also attach to flies using their legs and claws (Harbison et al., 2009). The single published record of phoresy of an amblyceran louse noted that the louse was attached by its

mouthparts to one of the fly's legs (Hopkins, 1946). In addition, we recently examined hippoboscids
 flies (*Pseudolynchia canariensis*) leaving captive Rock Pigeons (*Columba livia*) and found
 amblyceran lice (*Hohorstiella lata*) attached to several flies (Fig. 1). These lice appeared to be
 attached using their tarsal claws; these lice may also be attached with their mouthparts but scanning
 electron microscope images of this point of attachment were inconclusive (Fig. 1).

Regardless of the precise way in which lice attach to flies, it is clear that lice from all three
 suborders can be phoretic. Given that all lice face similar transmission barriers, why is phoresy so
 rare among some groups of lice? We suggest that a fundamental life-history difference is
 responsible for these phoretic differences.

Dispersal is risky. However, phoresy may be an especially risky dispersal strategy given that
 lice typically disperse by moving from one host to another during periods of direct, physical contact
 between two host individuals (Clayton et al., 2016). Off the vertebrate host, lice can only live a few
 days (Johnson and Clayton, 2003b). Lice that are not transported to a compatible host during this
 short time-frame will die. Lice that fall off or that are groomed from the fly will also die without
 reaching a new host. Furthermore, hippoboscids are not as host-specific as lice. Flies may
 transport lice to a novel host species where the lice cannot survive or reproduce. The benefits of
 phoresy may outweigh the costs of phoretic dispersal for highly immobile species, but species that
 can move independently between host individuals may not need to engage in such risky behavior.
 For example, agile amblyceran lice will crawl away from dead or distressed hosts in search of a new
 host (Keirans, 1975a; Johnson and Clayton, 2003a; Clayton et al., 2016). It is not clear what cues
 most lice use to locate new host individuals, but the few studies that have been done indicate that
 chemosensory and tactile mechanisms are likely involved (Wigglesworth, 1941; Clayton et al.,
 2016). Regardless of the types of cues used, the probability of crawling to a new host may be high,

especially among gregarious or colonial host species that roost or nest in close proximity. In contrast, ischnoceran lice are so specialized for life on hair or feathers that they typically remain on the host, even if it is dead (Keirans, 1975a). For these lice, the benefits of phoresy may outweigh the costs.

Here, we investigated the relationship between independent mobility and phoresy among four species of lice that are ecological "replicates" in many respects (Johnson and Clayton, 2003b; Johnson et al., 2005). We compared two ischnoceran species: *Columbicola columbae* and *Campanulotes compar*; and two amblyceran species: *H. lata* and *Menacanthus stramineus* (Fig. 2). All four of these species are permanent obligate parasites that are found on Rock Pigeons (*C. livia*; Brown, 1971; Dranzoa et al., 1999; Price et al., 2003; Musa et al., 2011). Rock Pigeons are also parasitized by a hippoboscids fly (*P. canariensis*). These flies feed on blood and crawl throughout the bird's plumage (Harbison, C.W., 2008. Ecology and Evolution of Transmission in Feather-Feeding Lice (Phthiraptera: Ischnocera). Ph.D. Thesis, University of Utah, USA). Flies are found most commonly on the abdomen of the bird (Harbison et al., 2008), a region of the body where all four lice species occur. Thus, all four species have had opportunities to engage in phoresy over macroevolutionary time.

Another aspect of this host-parasite system that makes it especially suitable for testing the relative role of phoretic dispersal versus independent dispersal is that Rock Pigeons nest in flocks. Rock Pigeon nests are often very close together; neighboring nests are frequently 0.1 – 1.0 m apart (Johnston and Janiga, 1995). Thus, all four species of lice have had opportunities to disperse to new host individuals by crawling to neighboring hosts over macroevolutionary time.

Despite these ecological similarities, these four species of lice have very different phoretic records. *Columbicola columbae* has been observed on hippoboscids flies on several occasions (Table

1), and has been shown to be phoretic in experiments (Harbison et al., 2009). The other ischnoceran louse, *Ca. compar*, has not been observed on flies and did not engage in phoresy in experimental settings (Harbison et al., 2009; Harbison and Clayton, 2011). Of the two amblyceran species in this study, neither species has been observed phoretically dispersing. However, one closely related species (*Hohorstiella gigantea*, a parasite of Stock Doves, *Columba oenas*) was found on a hippoboscid fly and is the only known case of phoresy from the suborder Amblycera (Hopkins, 1946, Table 1).

Here, we investigate the relationship between the ability of these four species of lice to move independently off the surface of the host relative to their phoretic ability. Specifically, we quantify how far each species can crawl off the surface of the host in a specified amount of time. We also conduct three assays to determine the relative abilities of these four species of lice to i) attach to hippoboscid flies; ii) remain attached to hippoboscid flies while the flies groom, and iii) remain attached to hippoboscid flies during flight.

2. Materials and methods

2.1. Relative mobility off the host

We compared the "off-host mobility" of four species of lice (*Co. columbae*, *Ca. compar*, *H. lata* and *M. stramineus*) by measuring the distance each species traveled on a filter paper in 2 min. We placed each louse, one at a time, in the center of a filter paper that was 15 cm in diameter. Next, we placed the cover of a glass petri dish over the filter paper. To stimulate louse movement, a light (Leica Illuminator, Model 31-35-28) was briefly shone on each louse. Lice are negatively phototactic and they immediately move away from light. As each louse moved, we traced its path on the cover glass with a marker. Each path was photographed and digitally measured using ImageJ

software (Rasband W.S., 2015. ImageJ. U.S. National Institutes of Health, Bethesda, Maryland, USA, imagej.nih.gov/ij/). We quantified off-host mobility for 20 lice of each of the four species. Flies and all four species of lice were obtained from culture stocks raised on captive Rock Pigeons kept in an animal facility at the University of Utah, USA, Institutional Animal Care and Use Committee # 11-07018). Our small culture of *M. stramineus* was supplemented with lice from B. Mullens, University of California, Riverside, USA.

2.2. Attachment of lice to flies

Three assays were conducted to compare the ability of *Co. columbae*, *Ca. compar*, *H. lata*, and *M. stramineus* to i) attach to a fly; ii) remain attached to a grooming fly, and iii) remain attached to a fly during flight. Flies used in assays all came from a captive culture population that was founded with flies from Salt Lake City, UT, USA.

First, the relative ability of lice to attach to flies was assessed by placing lice, one at a time, in a small (5 mL) glass vial with a single hippoboscid fly. These small vials ensured that each louse came into contact with the fly, and thus had an opportunity to attach. During each “trial” the louse and fly were monitored for 2 min, or until the louse attached, whichever came first. Lice were considered “attached” if they held onto the fly for at least 2 s. We conducted attachment trials for at least 55 lice of each species. Trials were ordered randomly with respect to louse species, and flies were replaced every 20 trials with a new fly, which was a fly that had not previously been used in any assay. Lice that attached during the first assay were immediately placed in the grooming assay (see below).

In the grooming assay, we compared the ability of lice to withstand grooming by flies. Flies with attached lice were placed into a small (10x10x18 cm) clear plastic observation chamber. Upon

181 entering the chamber, the flies would perch on a wall of the chamber and use their legs to groom
 182 their own bodies. This behavior is easily observable with the naked eye. The length of time that
 183 each louse stayed attached to a grooming fly was recorded. The observation period lasted 5 min, or
 184 until the louse detached, whichever came first.

185 Third, we compared the ability of three species of lice to remain attached to flies that flew
 186 approximately 3 m. *Menacanthus stramineus* was not used in this assay because very few *M.*
 187 *stramineus* stayed attached to flies during the previous assays (see Section 3). Flight trials were
 188 conducted in a small room (3.5 x 3.0 x 1.5 m) with a large window at the end (as in Harbison et al.,
 189 2009). A single fly and a louse were first constrained within a small vial, so that the louse had an
 190 opportunity to attach to the fly. Once the louse was attached for 2 s, the fly was released; if a louse
 191 did not attach within 2 min, it was discarded. Flies are positively phototactic and flew directly
 192 towards the window upon release. Once a fly landed on the window, the cover of a glass petri dish
 193 was placed over the fly and the presence or absence of the louse was recorded. A new louse was
 194 used for each trial. We conducted flight trials for 55 lice of each species. Trials were ordered
 195 randomly with respect to louse species and flies were replaced every 12 trials with a new fly.

196

197 **3. Results**

198 *3.1. Relative mobility off the host*

199 We compared the relative ability of four species of lice to move off the host by quantifying
 200 the distance that each species moved on a filter paper. The four species of lice significantly differed
 201 in the distance they traveled in 2 min (Fig. 3A; Kruskal-Wallis, $\chi^2 = 58.3$, degrees of freedom = 3, P
 202 < 0.0001). Both amblyceran species traveled significantly farther than either species of ischnoceran

louse. *Menacanthus stramineus* traveled three-fold farther than *H. lata* and over 22-fold farther than either ischnoceran species.

3.2. Attachment of lice to flies

We compared the relative ability of four species of lice to attach to hippoboscid flies that were placed in close contact. The four species of lice significantly differed in attachment to flies ($\chi^2 = 119.4$, degrees of freedom (df) = 3, $P < 0.0001$; final sample sizes indicated in Fig. 3B). Nearly all ischnoceran lice attached to flies (100% of *Co. columbae* and 96.9% of *Ca. compar*). The majority of *H. lata* also attached to flies (89.2%), but they did so significantly less than *Co. columbae* (Fisher's Exact test, $P = 0.013$). *Menacanthus stramineus* attached to flies in only 30.9% of trials, which was significantly less than any of the other species (Fisher's Exact test, $P < 0.0001$, in all three cases).

Next, we compared the ability of the four species of lice to withstand fly grooming. Flies did not groom in 5% of trials and these trials were excluded from analyses (final sample sizes are indicated in Fig. 3C). The amount of time that the four species of lice remained attached to grooming flies was significantly different (Kruskal-Wallis, $\chi^2 = 43.3$, df = 3, $P < 0.0001$). A majority of *Co. columbae* (61%) stayed attached for the entire 5 min trial. In contrast, only 4.9% of *Ca. compar*, 18.2 % of *H. lata*, and 0% of *M. stramineus* remained attached to the fly for the full 5 min trial.

Lastly, we compared the ability of three species of lice to remain attached to flying flies. Rarely (3%), flies did not fly towards the window; these trials were excluded from analyses. In addition, four (7.5%) *H. lata* did not attach to the fly within the allotted time (2 min), and these

trials were also excluded from analyses (Fig. 3D). The percentage of lice that remained attached to flies during flight differed significantly among the three species ($\chi^2 = 13.39$, $df = 2$, $P = 0.001$).

4. Discussion

We tested the ability of four species of lice to move, independently, on substrates off the host. We found that the two species of ischnoceran lice, *Co. columbae* and *Ca. compar*, are quite immobile off the host. The mean crawling speed of either ischnoceran spp. did not exceed 8 mm/min off the host. By comparison, *Co. columbae* and *Ca. compar* are much more mobile when they are on feathers. Using the same experimental setup, Bush (Bush, S.E., 2004. Evolutionary Ecology of Host Specificity in Columbiform Feather Lice. Ph.D. Thesis, Department of Biology, University of Utah, USA) placed lice on feathers from their preferred microhabitats on the host and recorded the crawling speed of each species. In this study, *Co. columbae* traveled (mean $\pm$ S.E.) 121 ± 8.7 mm/min and *Ca. compar* crawled 161 ± 13.6 mm/min. In short, both species of lice can travel 15-fold faster on feathers than on surfaces off the host.

In contrast, the two amblyceran species in the study were quite mobile off the host. *Hohorstiella lata* crawled at a speed of 55 mm/min, and *M. stramineus*, the most mobile louse species in the study, crawled 174 mm/min, with the fastest individual louse traveling at 401 mm/min. Moreover, both *M. stramineus* and *H. lata* readily crawled off filter paper and on to the glass petri dish, which indicates they could even cross slick surfaces they might encounter in the wild. In contrast, neither species of ischnoceran louse can move on slick surfaces such as glass (personal observation).

At these speeds, amblyceran lice may be able to crawl between nests in the wild. Rock Pigeons frequently nest in close proximity, often less than 1 m apart (Johnston and Janiga, 1995). At

the observed speeds, *M. stramineus* could travel between nests 1 m apart in 6 min and *H. lata* could travel between the same nests in 18 min. Given how fast these lice crawl, it seems that crawling between neighboring nests could be an effective dispersal strategy for these lice. In contrast, it would take *C. columbae* 2.1 h and *C. compar* 3.5 h to cover the same distance, assuming the lice do not encounter any surfaces too slick to cross. Both *Co. columbae* and *Ca. compar* can survive a few days off the body of the host (Johnson and Clayton, 2003b), so it is conceivable that these lice could survive such a journey, however improbable it may be. Whether or not any of the four species of lice actually crawls between hosts in the wild is unknown. Our data indicate that crawling for several hours may prevent ischnoceran lice from successfully dispersing between nests, whereas amblycerans should be able to use this dispersal strategy. A study in the field is needed to test this hypothesis.

We also tested the ability of these lice to move phoretically on hippoboscids flies. The four species of lice in our study differed in their abilities to attach to a fly and remain attached to the fly while it was grooming or flying. Both ischnoceran species and one amblyceran species (*H. lata*) readily attached to flies when they were placed in close contact. All *Co. columbae* attached, and nearly all (>89%) *Ca. compar* and *H. lata* attached to the flies. In contrast, the second amblyceran species (*M. stramineus*) attached to flies only 29% of the time, and the few *M. stramineus* that initially attached to flies quickly detached from grooming flies. On grooming flies, *Co. columbae* stayed attached much longer than any other species. Both *Ca. compar* and *H. lata* remained attached to grooming flies but for less than half as long as *Co. columbae*.

It is difficult to determine how important it is that lice are able to withstand grooming by the fly, because there is no information about the amount of time that hippoboscids flies spend grooming in the wild. The screwworm (*Cochliomyia hominivorax*: Calliphoridae), a distantly related parasitic

fly, spends over 19% of its time engaged in grooming behavior (Thomas, 1991). If the pigeon fly *P. canariensis* also spends a considerable amount of time grooming then this could explain, at a proximal level, why *Ca. compar* does not engage in phoretic dispersal even though it can attach to, and remain attached to, flying flies as well as *Co. columbae* (Harbison et al., 2008). The amblyceran louse *H. lata* detached from grooming flies as frequently as *Ca. compar* and, as with *Ca. compar*, there are no records of *H. lata* being phoretic under natural conditions.

Interestingly, the louse species *M. stramineus* is found on at least eight host species from two different host orders (Brown, 1971; Dranzoa et al., 1999; Price et al., 2003; Musa et al., 2011). By comparison, other lice in this study are restricted to either a single host genus (*Co. columbae*) or a single host species (*Ca. compar* and *H. lata*). *Menacanthus stramineus* is, by far, the least host-specific of any of the species of lice used in this study. One could argue that this species should be the most likely to engage in phoresis because it could potentially survive and reproduce on other host species that the fly is likely to visit (Maa, 1969). Despite the ability of *M. stramineus* to parasitize many host species, however, this louse was the least likely to engage in phoresis. In a comparative study, Harbison (Harbison, C.W., 2008. Ecology and Evolution of Transmission in Feather-Feeding Lice (Phthiraptera: Ischnocera). Ph.D. Thesis, University of Utah, USA) showed that phoretic species are more likely to be found on multiple host species than are closely related species that are not phoretic. Our data examines this relationship from another angle, and shows that species found on multiple host species are not necessarily more phoretic.

We hypothesized that parasites, which are able to move independently off the surface of the host, should be less likely to engage in phoresy than relatively immobile parasites. We found that the species that was best able to crawl off the surface of the host, *M. stramineus*, performed poorly in all assays designed to quantify phoretic ability. In contrast, one of the least mobile species, *Co.*

294 *columbae*, was able to attach to, and remain attached to, flies more often and for a longer period of
 295 time than any other species. *Hohorstiella lata* exhibited intermediate performance in the off-host
 296 mobility assay as well as the three assays quantifying phoretic ability. Perplexingly, however, *Ca.*
 297 *compar* is as immobile off the host as *Co. columbae*, yet *Ca. compar* exhibited intermediate
 298 performance in the assays quantifying phoretic ability. Moreover, there are no records of *Ca.*
 299 *compar* being phoretic, whereas *Co. columbae* has repeatedly been found on flies (Table 1).

300 Other ecological factors may be driving the phoretic differences between *Co. columbae* and
 301 *Ca. compar*. Bush and Malenke (2008) found that *Co. columbae* is competitively inferior to *Ca.*
 302 *compar*. Harbison et al. (2008) found that phoresy can be an important mechanism for escaping
 303 competition with other species of lice on the same host. Our results are also consistent with a
 304 competition-colonization trade-off model between these two species. If by dispersing phoretically,
 305 *Co. columbae* are better able to escape from competition with *Ca. compar*, then this benefit may
 306 outweigh the potential risks to phoretic dispersal.

307 Competition is thought to drive phoretic behavior in other systems. For example, the
 308 phoretic behavior of the pseudoscorpion *Paratemnoides nidificator* seems to be driven, in part, by
 309 intraspecific competition (Tizo-Pedroso and Del-Claro, 2007). This species is most likely to attempt
 310 phoresy when there are many pseudoscorpions living together in a single colony (Tizo-Pedroso and
 311 Del-Claro, 2007). More studies are needed among "ecological replicates" to determine whether
 312 mobility and/or competition frequently trigger the evolution of phoretic behavior in other animal
 313 systems.

314

315

316

317 **Acknowledgements**

318 We thank D. H. Clayton, A. Hansen, S. McNew, D. Gustafsson for various forms of
319 assistance. We thank B. Mullens for providing some of the *M. stramineus* used in this study. We
320 also thank two anonymous reviewers for providing helpful comments. This work was supported by
321 the USA National Science Foundation Division of Environmental Biology (DEB-1342600) and the
322 USA National Science Foundation Graduate Research Fellowship Program (GRFP-1256065) to
323 AWB.

References

- Allingham, P.G., 1987. Phoresy involving a nymph of *Haematopinus eurysternus* (Nitzsch) and *Haematobia irritans exigua* De Meijere. J. Aust. Ent. Soc. 26, 237 - 238.
- Ansari, M.A.R., 1947. Associations between the Mallophaga and the Hippoboscidae infesting birds. J. Bombay Nat. Hist. Soc. 46, 509 - 516.
- Ash, J., 1952. Records of Hippoboscidae (Dipt.) from Berkshire and Co., Durham, in 1950, with notes on their bionomics. Entomol. Mon. Mag. 88, 25 - 30.
- Athias-Binche, F., Morand, S., 1993. From phoresy to parasitism: the example of mites and nematodes. Res. Rev. Parasit. 53, 73 - 79.
- Baker, J.R., Blackie, E.J., 1963. Phoresy of *Brueelia varia* (Burm.) (Mallophaga, Philopteridae) on *Ornithomyia avicularia* L. (Dipt. Hippoboscidae). Entomol. Mon. Mag. 99, 95.
- Baum, V.H., 1968. Biologie und ökologie der amselfederlause. Angew. Parasitol. 9, 129 – 176.
- Bay, D.E., 1977. Cattle-biting louse, *Bovicola bovis* (Mallophaga: Trichodectidae), phoretic on the horn fly, *Haematobia irritans* (Diptera: Muscidae). J. Med. Entomol. 13, 628.
- Bedford, G.A.H., 1929. Anoplura (Siphunculata) and Mallophaga from South African hosts. Rep. Vet. Res. S. Afr. 15, 501 - 549.
- Bennett, G.F., 1961. On three species of Hippoboscidae (Diptera) on birds in Ontario. Can. J. Zool. 39, 379 - 406.
- Blagoveshtchenski, D.I., 1956. Stroenie i. systematicheskoe znacenie polovejsistemy puchoedov (Mallophaga). Parazitologičeskij Zbornik 16, 5 - 88.
- Braack, L.E.O., Emerson, K.C., 1986. A louse phoretic on a haematophagous muscid fly. J. Entomol. Soc. S. Afr. 49, 161 - 162.

- 346 Brown, N.S., 1971. A survey of the arthropod parasites of pigeons (*Columba livia*) in Boston. J.
347 Parasitol. 57, 1379 - 1380.
- 348 Bush, S.E., Malenke, J.R., 2008. Host defense mediates interspecific competition in parasites. J.
349 Anim. Ecol. 77, 558 - 564.
- 350 Buttiker, W., 1949. Über die Übertragungsweise von Ektoparasiten bei Vögeln. Die Vogel der
351 Heimat 19, 74 - 77.
- 352 Calandruccio, S., 1890. Animali parassiti dell'uomo in Sicilia. Atti d. Accad. Gioenia di Sc. Nat. in
353 Catania 2, 95 - 135.
- 354 Callot, J., 1946. Nouveau cas de phoresie. Ann. Parasit. Hum. Comp. 21, 379 - 380.
- 355 Clay, T., 1949. Some problems in the evolution of a group of ectoparasites. Evolution 3, 279 - 299.
- 356 Clay, T., Meinertzhagen, R., 1943. The relationship between Mallophaga and hippoboscids flies.
357 Parasitology 35, 11 - 16.
- 358 Clayton, D.H., Bush, S.E., Johnson, K.P., 2016. Coevolution of Life on Hosts: Integrating Ecology
359 and History. University of Chicago Press, Chicago and London.
- 360 Corbet, G.B., 1956. The phoresy of Mallophaga on a population of *Ornithomya fringillina* Curtis
361 (Dipt., Hippoboscidae). Entomol. Mon. Mag. 92, 207 - 211.
- 362 Couch, A.B., 1962. Phoretic mallophagans from hippoboscids of Mourning Doves, *Zenaidura*
363 *macroura*. J. Parasitol. 48, 497.
- 364 Dranzoa, C., Ocaido, M., Katete, P., 1999. The ecto-, gastro-intestinal and haemo-parasites of live
365 pigeons (*Columba livia*) in Kampala, Uganda. Avian Path. 28, 119 - 124.
- 366 Dubinin, V. B., 1947. Investigation of the adaptation of ectoparasites. II. Ecological adaptations of
367 feather-mites and Mallophaga. Parazitologicheskii Sbornik Moscow 9, 191 - 222.

- 368 Durden, L.A., 1990. Phoretic relationships between sucking lice (Anoplura) and flies (Diptera)
 369 associated with humans and livestock. Entomologist 109, 191 - 192.
- 370 Edwards, R., 1952. Flatflies taken in the laboratory during 1951. Fair Isle Bird Obs. Bull. 6, 37 - 38.
- 371 Eichler, W., 1939. Deutsche lausfliegen, ihre lebensweise und ihre hygienische bedeutung. A. Hyg.
 372 Zool. 31, 210 - 226.
- 373 Eichler, W., 1946. Problems der Mallaphagenfarschung. Die Vogel der Heimat 16, 205 - 217.
- 374 Ewing, H.E., 1927. The hippoboscids fly, *Ornithomyia avicularia* Linnaeus, as carrier of
 375 Mallophaga. Ann. Entomol. Soc. Am. 20, 245 - 250.
- 376 Farish, D.J., Axtell, R.C., 1971. Phoresy redefined and examined in *Macrocheles muscaedomesticae*
 377 (Acarina: Macrochelidae). Acarologia 13, 16-29.
- 378 Forsius, R., 1912. Uber den transport von Mallophagen durch hippobosciden. Medd. Soc. Fauna
 379 Flora Fenn. 38, 58 - 60.
- 380 Guimaraes, L.R., 1944. Mais um caso de associacao entre Mallophaga e Hippoboscidae. Pap. Avul.
 381 Depto. Zool. São Paulo 4, 79 - 84.
- 382 Harbison, C.W., Bush, S.E., Malenke, J.R., Clayton, D.H., 2008. Comparative transmission
 383 dynamics of competing parasite species. Ecology 89, 3186 - 3194.
- 384 Harbison, C.W., Clayton, D.H., 2011. Community interactions govern host-switching with
 385 implications for host-parasite coevolutionary history. Proc. Natl. Acad. Sci. USA. 108,
 386 9525-9529.
- 387 Harbison, C.W., Jacobsen, M.V., Clayton, D.H., 2009. Hitchhiker's guide to parasite transmission:
 388 phoretic behavior of feather lice. Int. J. Parasitol. 39, 569 - 575.
- 389 Harrison, L., 1913. Notes and exhibits. P. Linn. Soc. N. S. W. 38, 108 - 109.

- 390 Hathaway, C.R., 1943. Associacao entre Mallophaga e Hippoboscidae. Mem. Inst. Oswaldo Cruz
391 38, 413 - 417.
- 392 Hopkins, G.H., 1946. Stray notes on Mallophaga. Ann. Mag. Nat. Hist. 13, 170 - 183.
- 393 Houck, M.A., OConnor, B.M., 1991. Ecological and evolutionary significance of phoresy in the
394 Astigmata. Ann. Rev. Entomol. 611 – 636.
- 395 Iannacone, J.A., 1992. Registro de un caso de phoresis: *Columbicola columbae* (L) (Phthiraptera:
396 Insecta) por *Psuedolynchia canariensis* (Diptera: Insecta) en la zona de Lima, Peru. Bol.
397 Lima 84, 17 - 18.
- 398 Johnson, K.P., Clayton, D.H., 2003a. Coevolutionary history of ecological replicates: comparing
399 phylogenies of wing and body lice to Columbiform hosts. In: Page, R.D.M., (Ed.) Tangled
400 Trees: Phylogeny, Cospeciation and Coevolution. University of Chicago Press, Chicago and
401 London, pp. 262 - 286.
- 402 Johnson, K.P., Clayton, D.H., 2003b. The biology, ecology, and evolution of chewing lice. In: Price,
403 R.D., Hellenthal, R.A., Palma, R.L., Johnson, K.P., Clayton, D.H. (Eds.), The Chewing
404 Lice: World Checklist and Biological Overview. Illinois Natural History Survey Special
405 Publication 24, Champaign, pp. 451 - 475.
- 406 Johnston, R.F., Janiga M., 1995. Feral Pigeons. Oxford University Press, New York, USA.
- 407 Keirans, J.E., 1975a. A review of the phoretic relationship between Mallophaga (Phthiraptera:
408 Insecta) and Hippoboscidae (Diptera: Insecta). J. Med. Ent. 12, 71 - 76.
- 409 Keirans, J.E., 1975b. Records of phoretic attachment of Mallophaga (Insecta: Phthiraptera) on
410 insects other than Hippoboscidae. J. Med. Entomol. 12, 476.

- 411 Kirk-Spriggs, A.H., Mey, E., 2014. Phoresy of a sucking louse, *Linognathus* sp. (Phthiraptera:
 412 Anoplura: Linognathidae), by *Musca* (Byomya) *conducens* Walker (Diptera: Muscidae) in
 413 South Africa. African Invertebr. 55, 119 - 123.
- 414 Kuris, A.M., Blaustein, A.R., Alio, J.J., 1980. Hosts as islands. Am. Nat. 116, 570 - 586.
- 415 MacArthur, R.H., Pianka, E.R., 1966. On optimal use of a patchy environment. Am. Nat. 100, 603
 416 - 609.
- 417 Macchioni, F., Magi, M., Mancianti, F., Perruci, S., 2005. Phoretic association of mites and
 418 Mallophaga with the pigeon fly *Pseudolynchia canariensis*. Parasite 12, 277 - 279.
- 419 Main, A.J., Anderson, K.S., 1970. The genera *Ornithoica*, *Ornithomyia*, and *Ornithoctona* in
 420 Massachusetts (Diptera: Hippoboscidae). Bird-Banding 41, 300 - 306.
- 421 Main, A.J., Anderson, K.S., 1971. Avian ectoparasites at Manomet Bird Observatory. Part II. Fleas
 422 Flies. Manomet Bird Observ. Res. Report 3, 1 - 5.
- 423 Markov, G.S., 1938. The presence of phoresy in Mallophaga. Zool. Zh. 17, 634 - 636.
- 424 Martin, M., 1934. Life history and habits of the pigeon louse (*Columbicola columbae* (Linnaeus)).
 425 Can. Entomol. 66, 6 - 16.
- 426 Matthyse, J.G., 1946. Cattle lice, their biology and control. Bull. Cornell Univ. Agric. Expt. Stn.
 427 832, 1 - 67.
- 428 McAtee, W.L., 1922. Bird lice (Mallophaga) attaching themselves to bird flies (Diptera,
 429 Hippoboscidae). Entomol. News 33, 90.
- 430 Mitzmain, M.B., 1912. Collected notes on the insect transmission of surra in Carabaos. Philipp.
 431 Agric. Rev. 5, 670 - 681.
- 432 Mjoberg, E., 1910. Studien uber Mallophagen und Anopluren. Arkiv Zool. 6, 1 - 297.

- 433 Musa, S., Afroz, S.D., Khanum, H., 2011. Occurrence of ecto- and endo parasites in pigeon
 434 (*Columba livia* Linn.). Univ. J. Zool. Rajshahi Univ. 30, 73 - 75.
- 435 Nutall, G.H.F., 1917. The biology of *Pediculus humanus*. Parasitology 10, 80 - 185.
- 436 Overgaard Nielsen, B., 1990. Phoresy of *Cervicola meyeri* (Taschenb.) on *Aedes communis* (Deg.)
 437 (Mallophaga: Ischnocera & Diptera: Culicidae). Entomol. Medd. 58, 43 - 45.
- 438 Pajot, F.X., Germain, M., 1971. Note sur un cas nouveau de phorésie chez les insectes transport de
 439 *Linognathus breviceps* (Piaget) [Anoplura Linognathidae] par des *Eretmapodites* du groupe
 440 *chrysogaster* [Diptera Culicidae]. Bull. Soc. Entomol. Fr. 76, 5 – 6.
- 441 Peus, F., 1933. Tansport von Mallophaga durch stechmucken. Z. Parasitenk. 5, 740-741.
- 442 Peters, H.S., 1935. Mallophaga carried by hippoboscids. Annls. Carnegie Mus. 24, 57 – 58.
- 443 Price, R.D., Hellenthal, R.A., Palma, R.L., Johnson, K.P., Clayton, D.H., 2003. The Chewing Lice:
 444 World Checklist and Biological Overview. Illinois Natural History Survey Special
 445 Publication 24, Champaign, USA.
- 446 Proctor, H.C., Jones, D.N., 2004. Geographical structuring of feather mite assemblages from the
 447 Australian Brush-turkey (Aves: Megapodiidae). J. Parasitol. 90, 60 - 66.
- 448 Rothschild, M., Clay, T., 1952. Fleas, Flukes & Cuckoos: A Study of Bird Ectoparasites. Collins,
 449 London, UK.
- 450 Roubik, D.W., Wheeler, Q.D., 1982. Flightless beetles and stingless bees: Phoresy of scotocryptine
 451 beetles (Leiodidae) on their meliponine hosts (Apidae). J. Kansas Entomol. Soc. 55, 125 – 135.
- 452 Spencer, G.J., 1928. External parasites on certain birds of British Columbia. Can. Entomol. 60, 257
 453 - 260.
- 454 Thomas, D.B., 1991. Time-activity budget of adult screwworm behavior (Diptera: Calliphoridae). J.
 455 Med. Entomol. 28, 372 - 377.

- 456 Thompson, G.B., 1933. Association of hippoboscids with lice. *Nature* 132, 605 - 606.
- 457 Thompson, G.B., 1934. The parasites of British birds and mammals. I. Notes and records. *Entomol.*
 458 *Mon. Mag.* 70, 133 - 136.
- 459 Thompson, G.B., 1937. Further notes on the association of Hippoboscidae and Mallophaga. *Ann.*
 460 *Mag. Nat. Hist.* 20, 441 - 442.
- 461 Thompson, G.B., 1947. Association of Hippoboscidae and Mallophaga: Further notes and records.
 462 *Entomol. Mon. Mag.* 83, 212 - 214.
- 463 Treat, A.E., 1956. A pseudoscorpion on moths. *Lepid. News* 10, 87 - 89.
- 464 Warburton, C., 1928. *Ornithomyia avicularia* (Diptera: Hippoboscidae) as the carrier of
 465 Mallophaga, with some remarks on phoresis in insects. *Parasitology* 20, 175 - 178.
- 466 Walter, V.G., 1989. Phoresie und hyperparasitismus bei *Ornithomyia* (Diptera, Hippoboscidae) in
 467 der Bundesrepublik Deutschland. *Angew. Parasitol.* 30, 43 - 46.
- 468 Ward, R.A., 1953. Additional record of phoresy of Mallophaga on Hippoboscidae. *Bull. Brooklyn*
 469 *Entomol. Soc.* 48, 128.
- 470 Wilson, N., 1964. Records of Hippoboscidae (Diptera) from Indiana. *J. Med. Entomol.* 1, 128 - 130.
- 471 Wigglesworth, V.B., 1941. The sensory physiology of the human louse *Pediculus humanus corporis*
 472 de Geer (Anoplura). *Parasitology* 33, 67 - 109.
- 473 Worth, B., Patterson, H.E., 1960. Phoresy of sucking lice (Siphunculata: Linognathidae) by a
 474 mosquito (Diptera: culicidae). *J. Entomol. Soc. S. Afr.* 23, 228 - 230.
- 475 Zeh, D.W., Zeh, J.A., 1992. On the function of harlequin beetle-riding in the pseudoscorpion,
 476 *Cordylochernes scorpioides*. *J. Arachnol.* 20, 47 - 51.

Legends to Figures

Fig. 1. Photo of a hippoboscid fly (*Pseudolynchia canariensis*) captured from a captive Rock Pigeon (*Columba livia*) with two amblyceran lice (*Hohorstiella lata*) attached to the abdomen of the fly. Inset: false-colored scanning electron microscope image.

Fig. 2. Four species of lice (to scale) from Rock Pigeons (*Columba livia*): (A) *Columbicola columbae* (Ischnocera), (B) *Campanulotes compar* (Ischnocera), (C) *Hohorstiella lata* (Amblycera), and (D) *Menacanthus stramineus* (Amblycera).

Fig. 3. Comparisons of off-host mobility and fly attachment for four species of lice (*Columbicola columbae*, *Campanulotes compar*, *Hohorstiella lata*, *Menacanthus stramineus*) from Rock Pigeons (*Columba livia*). Bars with different letters differ significantly. (A) Distance lice traveled off the host in 2 min. The four species of lice differed significantly in off-host mobility ($P < 0.0001$, see Section 3; Mann-Whitney U post hoc tests, $P < 0.05$). (B) Percentage of lice that attached to pigeon flies (based on post hoc Fisher's Exact tests of raw data, $P < 0.05$). (C) The amount of time that lice stayed attached to grooming flies differed significantly among species ($P < 0.0001$, see section 3; Mann-Whitney U post hoc tests, $P < 0.05$). (D) Percentage of lice remaining attached to a fly during flight (post hoc Fisher's Exact tests of raw data, $P < 0.05$).

497 **Table 1.** Species of lice that have been found attached to flies, based on a comprehensive survey of
 498 literature from 1890 to 2014 (updated from Harbison, C.W., 2008. Ecology and Evolution of
 499 Transmission in Feather-Feeding Lice (Phthiraptera: Ischnocera). Ph.D. Thesis, University of Utah,
 500 USA). Cases in which the phoretic lice were not identified to genus are not included.

Suborder	Louse Species	Host	References
Amblycera	<i>Hohorstiella giganteus</i> ^a	bird	Hopkins (1946)
Anoplura	<i>Haematopinus eurysternus</i>	mammal	Allingham (1987)
	<i>Haematopinus tuberculatus</i>	mammal	Mitzmain (1912)
	<i>Linognathus africanus</i>	mammal	Braack and Emerson (1986)
	<i>Linognathus breviceps</i>	mammal	Pajot and Germain (1971)
	<i>Linognathus vituli</i>	mammal	Bedford (1929)
	<i>Linognathus</i> sp.	mammal	Worth and Patterson (1960)
	<i>Linognathus</i> sp.	mammal	Kirk-Spriggs and Mey (2014)
	<i>Pediculus humanus</i>	mammal	Calandruccio (1890), Nuttall (1917)
Ischnocera	<i>Ardeicola botauri</i>	bird	Peters (1935)
	<i>Bovicola meyeri</i>	mammal	Keirans (1975b)
	<i>Brueelia amsel</i>	bird	Baum (1968), Walter (1989)
	<i>Brueelia clayae</i>	bird	Main and Anderson (1970, 1971)
	<i>Brueelia deficiens</i>	bird	Spencer (1928)
	<i>Brueelia glandarii</i>	bird	Eichler (1939), Callot (1946), Buttiker (1949)
	<i>Brueelia hectica</i>	bird	Harrison (1913)
	<i>Brueelia interposita</i>	bird	Ewing (1927), Thompson (1937)
	<i>Brueelia marginata</i>	bird	Ash (1952), Harrison (1913), Warburton (1928), Thompson (1933), Blagoveshtchenski (1956)
	<i>Brueelia merulensis</i>	bird	Ash (1952), Clay (1949), Clay and Meinerzthagen (1943) Eichler (1939), Walter (1989)
	<i>Brueelia rotundata</i>	bird	McAtee (1922)
	<i>Brueelia turdinulae</i>	bird	Walter (1989)
	<i>Brueelia unicosia</i>	bird	Forsius (1912), Dubinin (1947), Blagoveshtchenski (1956)
	<i>Brueelia varia</i>	bird	Baker and Blackie (1963)
	<i>Brueelia zeropunctata</i>	bird	Wilson (1964)
	<i>Cervicola meyeri</i>	mammal	Overgaard Nielsen (1990)
	<i>Columbicola columbae</i>	bird	Martin (1934), Ward (1953), Ansari (1947), Hathaway (1953), Ianacone (1992), Macchioni et al. (2005)

<i>Columbicola macrourae</i>	bird	Couch (1962)
<i>Damalinia bovis</i>	mammal	Matthysse (1946), Bay (1977)
<i>Damalinia meyeri</i>	mammal	Keirans (1975b)
<i>Damalinia tibialis</i>	mammal	Peus (1933)
<i>Gyropus ovalis</i>	mammal	Keirans (1975b)
<i>Lagopoecus lyrurus</i>	bird	Forsius (1912)
<i>Lipeurus crassus</i>	bird	Proctor and Jones (2004)
<i>Nirmus sp.</i>	bird	Keirans (1975b)
<i>Philopterus coarctatus</i>	bird	Eichler (1946)
<i>Physconelloides zenaidurae</i>	bird	Couch (1962)
<i>Strigiphilus crenulatus</i>	bird	Blagoveshtchenski (1956)
<i>Sturnidoecus pastoris</i>	bird	Eichler (1939)
<i>Sturnidoecus simplex</i>	bird	Harrison (1913)
<i>Sturnidoecus sturni</i>	bird	Mjoberg (1910), Harrison (1913), Thompson (1934, 1947), Markov (1938), Eichler (1939), Clay and Meinerzthagen (1943), Corbet (1956)
<i>Trogoninirmus odontopleuron</i>	bird	Guimares (1944)
<i>Trichodectes melis</i>	mammal	Keirans (1975b)

^a Species formerly known as *Menacanthus giganteus*

501
502
503
504
505
506
507

Table 2. Relative frequency of phoresy among the three major suborders of lice. Data based on published phoretic records (Table 1). The frequency of phoresy differs significantly among suborders ($\chi^2 = 14.06$, degrees of freedom = 2, $P = 0.001$). Different letters indicate significant differences (post hoc Fisher's exact tests, $P < 0.001$).

Suborder	Total species	Phoretic species	% Phoretic species
Ischnocera	3060	33	1.08 ^a
Anoplura	532	8	1.50 ^a
Amblycera	1334	1	0.07 ^b

Figure 1

Figure 2

Figure 3

Graphical Abstract