1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Thermodynamic properties of xanthone: Heat capacities, phase-transition properties, and thermodynamic-consistency analyses using computational results^{*}

Robert D. Chirico,^{a,†} Andrei F. Kazakov,^a

^a *Applied Chemicals and Materials Division*

National Institute of Standards and Technology

Boulder, CO 80305-3337, U.S.A.

^{*} Contribution of the National Institute of Standards and Technology, not subject to copyright in the United States.

[†] Corresponding author. E-mail address: robert.chirico@nist.gov

Abstract

Heat capacities and phase-transition properties for xanthone (IUPAC name *9H*-xanthen-9-one and Chemical Abstracts registry number [90-47-1]) are reported for the temperature range $5 < T/K < 524$. Statistical calculations were performed and thermodynamic properties for the ideal gas were derived based on molecular geometry optimization and vibrational frequencies calculated at the B3LYP/6-31+G(d,p) level of theory. These results are combined with sublimation pressures from the literature to allow critical evaluation of inconsistent enthalpies of sublimation for xanthone, also reported in the literature. Literature values for the enthalpy of combustion of xanthone are reassessed, a revision is recommended for one result, and a new value for the enthalpy of formation of the ideal gas is derived. Comparisons with thermophysical properties reported in the literature are made for all other reported and derived properties, where possible.

Keywords: xanthone; computational chemistry; enthalpy of combustion, enthalpy of formation; enthalpy of fusion; enthalpy of sublimation; heat capacity; ideal-gas properties; phase-transition; triple point temperature

1. Introduction

This article is part of a series in which we have reported studies of experimental and computed thermodynamic properties for aromatic ring systems, including an extensive series of azaaromatics (2-methylquinoline [1], 8-methylquinoline [1], 2,6-dimethylquinoline [1], phenazine [2], acridine [2], phenanthridine [3], 1,10-phenanthroline [3], and 7,8-benzoquinoline [3]), some hydroaromatics (1-phenyl and 2-phenylnaphthalene [4]), and oxygen-containing compounds (9-fluorenone [5], and 1-naphthol [6]). For this series, computations were performed at the B3LYP/6-31+G(d,p) model chemistry with excellent accord demonstrated between computed entropies for the ideal gas and those based on the experimental property measurements. In the present work, this approach is applied to xanthone (IUPAC name *9H*-xanthen-9-one and Chemical Abstracts registry number [90-47-1]).

Results of our previous work [1-6] have demonstrated that for rigid fused-ring systems thermodynamic properties for the ideal-gas state can be computed with relative standard uncertainties near 0.1 percent. In the present work, we use this finding together with new condensed-phase thermodynamic properties determined with adiabatic calorimetry to assess and resolve inconsistencies between enthalpies of sublimation reported in the literature for xanthone [7-9]. These results allow evaluation of the enthalpy of formation of xanthone in the ideal gas state with high confidence, based on enthalpies of combustion reported in the literature [7,8,10]. As part of this assessment, an apparent error in one report [10] was detected and corrected.

Here, we report results of measurements of heat capacities and phase-transition properties for xanthone for the temperature range $5 < T/\text{K} < 524$. The crystal-liquid-gas

triple-point temperature and enthalpy of fusion were determined, together with properties associated with a small lambda-like transition in the solid state. Measured and derived thermodynamic property values are compared with those reported in the literature, where possible.

2. Experimental

2.1 Materials

The research group of Professor E. J. “Pete” Eisenbraun (retired) of Oklahoma State University purified the sample of xanthone used in this research. A commercial sample was recrystallized twice from ethanol (95 percent) and twice from a 1:1 by volume mixture of (hexane + ethyl acetate). The resulting crystals were filtered and dried by evacuation at room temperature. The mole fraction purity x of the sample ($x = 0.9999$) was determined by fractional melting as part of the adiabatic calorimetric studies. Sample handling during purification and transfer was done under a dry atmosphere of argon, nitrogen, or helium.

2.2. Physical constants and standards

Property values for xanthone are reported in terms of molar mass $M = 196.19 \text{ g}\cdot\text{mol}^{-1}$ [11] based on the formula $\text{C}_{13}\text{H}_8\text{O}_2$ and the gas constant $R = 8.3144621 \text{ J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$ adopted in 2010 by CODATA [12]. The platinum resistance thermometer used here was calibrated by comparison with a standard thermometer calibrated at the National Bureau of Standards (NBS), now the National Institute of Standards and Technology (NIST). Temperatures were measured in terms of IPTS-68 [13] and were converted to ITS-90 with published differences [14]. The thermometer was calibrated below $T/\text{K} = 13.81$

with the method of McCrackin and Chang [15]. Mass, time, electrical resistance, and potential difference were measured in terms of standards traceable to calibrations at NIST.

2.3. *Adiabatic calorimetry*

Heat-capacities, enthalpy increments, and the triple-point temperature for xanthone were measured with an adiabatic calorimetric system described previously [16]. Sample-container characteristics and sealing conditions are given in table 1. Energies were measured with a relative standard uncertainty of 0.01 percent. Temperatures were measured with a repeatability of 0.0001 K, which represents the lower limit in the standard uncertainty for measured temperature increments. The standard uncertainty for a single temperature measurement was 0.005 K, due to uncertainties associated with the thermometer calibration, and represents the lower limit for the standard uncertainty in reported absolute temperatures, including phase-transition temperatures. The energy increments to the filled platinum calorimeter were corrected for enthalpy changes in the empty calorimeter and for the helium exchange gas. Corrections for vaporization of the sample into the free space of the sealed vessel were estimated to be insignificant, so none were applied. The maximum correction to the measured energy for the helium exchange gas was 0.4 percent near $T/K = 5$. The approximate size of the correction for the enthalpy changes in the empty calorimeter is given in table 1.

3. Experimental Results

3.1. Heat capacities, phase-transition properties, and derived thermodynamic functions for the condensed phases

Measurements were made for the temperature range ($5 < T/\text{K} < 524$), and included determination of heat capacities for the solid and liquid phases, the triple-point temperature T_{tp} , the molar enthalpy of fusion $\Delta_{\text{cr}}^{\text{l}}H_{\text{m}}$, and the temperature of a lambda-like transition T_{tr} in the solid state. The density of liquid xanthone under saturation pressure at temperature $T/\text{K} = 473$ $\{\rho_{\text{sat}} = 1090 \pm 20 \text{ kg}\cdot\text{m}^{-3}\}$ was determined from the sample mass and volume measured as part of the filling procedure for the calorimetric vessel.

Crystallization of xanthone in the calorimeter was initiated by slow cooling (approximately $7 \text{ mK}\cdot\text{s}^{-1}$) the liquid sample to approximately 10 K below T_{tp} . Complete crystallization was obtained by maintaining the sample under adiabatic conditions in the partially melted state (10 percent to 20 percent liquid) until ordering of the crystals was complete, as evidenced by the absence of spontaneous warming of the sample. In these experiments, no spontaneous warming was observed, indicating rapid and complete crystallization of the sample. The sample was then cooled at an effective rate of $7 \text{ mK}\cdot\text{s}^{-1}$ to crystallize the remaining liquid and was thermally cycled from approximately 100 K to within 3 K of the triple-point temperature ($T_{\text{tp}} = 448.79_5 \text{ K}$), where it was held for a minimum of 16 h to provide further tempering. All of the solid-phase measurements were performed upon crystals pre-treated in this way. Excellent repeatability (within 0.01 percent) was observed for three separate $\Delta_{\text{cr}}^{\text{l}}H_{\text{m}}$ determinations, which further indicates complete conversion to the crystalline state.

The triple-point temperature T_{tp} and mole fraction purity x for xanthone were determined by measurement of the equilibrium melting temperatures $T(F)$ as a function of fraction F of the sample in the liquid state [17]. Equilibrium melting temperatures were determined by measuring temperatures for ~ 1.5 h after an energy input and extrapolating to infinite time by assuming an exponential decay toward the equilibrium value. The observed temperatures at the end of the equilibration period were within 2 mK of the calculated equilibrium temperatures for F values listed in table 2 and used in the determination of T_{tp} . Results showed no evidence for the presence of solid-soluble impurities. The calculated triple-point temperature and sample purity are included in table 2.

The experimental heat capacities under vapor saturation pressure $C_{\text{sat,m}}$ determined by adiabatic calorimetry for xanthone are listed in table 3 and shown in figure 1. The values listed in table 3 were corrected for pre-melting with the mole fraction impurity value given in table 2. Relative expanded uncertainties (0.95 level of confidence) for the heat-capacity measurements $U_r(C_{\text{sat,m}})$ ranged from 0.03 at $T/\text{K} = 5$, to 0.005 at $T/\text{K} = 10$, and improved gradually to 0.001 for temperatures $T/\text{K} \geq 30$. Extrapolation of the heat-capacity results to $T/\text{K} \rightarrow 0$ was made with a plot of $C_{\text{sat,m}}/T$ against T^2 for temperatures below $T/\text{K} = 10$.

Equilibrium was reached in less than 1 h for all measurements in the liquid phase and for measurements in the solid phase below $T/\text{K} = 420$. As the triple-point temperature ($T_{\text{tp}} = 448.79_5$ K) was approached, equilibration periods increased gradually from 1 h near $T/\text{K} = 420$ to 6 h near $T/\text{K} = 445$.

As seen in figure 1, there is a small lambda-like transition near the temperature $T/K = 304.5$. Equilibration times remained less than 1 hour throughout this region, and enthalpy measurements across the transition were highly reproducible, as shown in table 4. No hysteresis was observed. Measurements were made with 3-K temperature increments (Series 8) and the transition temperature T_{tr} was derived through extrapolation of the heat capacities for each phase, followed by adjustment of the transition temperature to yield zero excess enthalpy for measurements made across the transition region with large (~ 20 K) temperature increments (Series 9 and 11). The transition temperature so determined was $T_{tr}/K = 304.5$ with an estimated expanded uncertainty of 0.5 K (0.95 level of confidence). No attempt was made to determine the detailed shape of the heat capacity curve in the transition region with temperature increments less than ~ 3 K, but as the variation of the heat capacity in the upper and lower phases was nearly linear with temperature, the assignment of the T_{tr} value could be made with high certainty.

Experimental molar enthalpy measurements are summarized in table 4, which includes results used in the determination of the enthalpy of fusion, determination of T_{tr} , and measurements in single-phase regions, which serve as consistency checks for the integrated heat-capacity values. Corrections for pre-melting [18] caused by impurities were made in these evaluations with the impurity x_{pre} listed in table 1. Results with the same series number N in tables 3 and 4 were taken without interruption of adiabatic conditions.

Entropies and enthalpies under vapor saturation pressure for the condensed phases relative to that of the crystals at $T \rightarrow 0$ K for xanthone are listed in table 5. These were

derived by integration of the smoothed heat capacities corrected for pre-melting, together with the entropy and enthalpy of fusion. All pre-melting corrections were made using standard methods [17].

4. Discussion

4.1 Comparisons with literature melting temperatures and enthalpies of fusion

The normal melting temperature T_m and enthalpy of fusion $\Delta_{cr}^1H_m$ for xanthone were reported by Sabbah and El Watik [7] $\{T_m/K = (449.63 \pm 0.3), \Delta_{cr}^1H_m/(kJ \cdot mol^{-1}) = (26.12 \pm 0.24)\}$ and Monte *et al.* $\{T_m/K = (448.3 \pm 0.1), \Delta_{cr}^1H_m/(kJ \cdot mol^{-1}) = (26.6 \pm 0.2)\}$ using differential calorimetric methods. The melting temperatures are within 0.8 K of the triple-point temperature T_{tp} determined here (table 2). For the enthalpies of fusion, the value reported by Monte *et al.*[9] is 3 percent low and that reported by Sabbah and El Watik [7] is 5 percent low. Such differences are common in comparisons of results from differential calorimetric methods with those of high-precision adiabatic calorimetry. It is also common that uncertainties reported with the differential methods are underestimated. There are numerous reports of T_m for xanthone in the literature, and these are, generally, within a few Kelvins of the value of T_{tp} reported here. Such differences are expected due to the relatively large uncertainties associated with the methods used for the literature values. In addition, the quality of the sample used in the present research far exceeds that used in most literature reports of normal melting temperature.

4.2 Assessment of literature enthalpies of sublimation through analysis of thermodynamic consistency

As noted in the introduction, we have demonstrated previously [1-6] that thermodynamic properties in the ideal-gas state for rigid fused-ring systems, such as xanthone, can be computed with relative uncertainties near 0.1 percent. Recently, this was demonstrated recently for the 3-ring ketone, 9-fluorenone [5]. Vibrational frequencies for xanthone were calculated at the B3LYP/6-31+G(d,p) model chemistry with the Gaussian 09 software [19]. We have demonstrated for many compounds [1-6] that the single scale factor $F = 0.975$ for the vibrational frequencies provides excellent agreement (within 0.1 percent) between computed ideal-gas entropies and those derived from calorimetric and thermophysical-property measurements. We also provided further justification for this scale factor through comparisons of computed and experimental gas-phase vibrational spectra [2] for quinoline and isoquinoline reported in the literature for all frequencies $\bar{\nu}$ less than 2000 cm^{-1} . Higher frequencies, which are all greater than 3000 cm^{-1} for most molecules, were not considered important in the comparison because they make no significant contribution to thermodynamic properties in the temperature range of interest in this research. The scale factor $F = 0.975$ and symmetry number 2 were used for xanthone, and the calculated vibrational frequencies and rotational constants are given in table 6. There is general agreement between the calculated values and those measured experimentally for condensed phases by Zwarich and Binbrek [20], Sinha *et al.* [21], and Connors and Fratini [22]. There are no gas-phase spectra available for xanthone to allow direct comparison with the vibrational frequencies computed here.

The enthalpy of sublimation for xanthone at given temperature T can be calculated with the relationship

$$\Delta_{\text{cr}}^{\text{g}}H_{\text{m}} = T \{ \Delta_0^T S_{\text{m}}^{\text{o}}(\text{ideal}) - \Delta_0^T S_{\text{m}}\{\text{cr(I)}\} - S_{\text{comp}} \} \quad (1)$$

where $\Delta_0^T S_{\text{m}}^{\text{o}}(\text{ideal})$ is the standard entropy of the ideal gas computed with parameters given in table 6 ($p^{\circ} = 101.325$ kPa) and the methods described in the text, $\Delta_0^T S_{\text{m}}(\text{cr})$ is the standard entropy of crystalline xanthone interpolated from table 5, and S_{comp} is the entropy of compression $\ln(p/p^{\circ})$, calculated with sublimation pressures reported by Monte *et al.*[9].

Sabbah and El Watik [7] reported a value of $\Delta_{\text{cr}}^{\text{g}}H_{\text{m}}$ at $T/\text{K} = 353$ based on a direct calorimetric method, and Monte *et al.*[9] reported this property at $T/\text{K} = 352.22$, based on the slope of sublimation pressures measured with Knudsen effusion. The sublimation enthalpies do not vary rapidly with temperature, so these values may be compared directly with the value at $T/\text{K} = 352.2$ derived in this research with equation (1). The experimental and derived values are listed in table 7. The value of $\Delta_{\text{cr}}^{\text{g}}H_{\text{m}}$ reported by Monte *et al.*[9] is $1.8 \text{ kJ}\cdot\text{mol}^{-1}$ higher than that derived here; however, the uncertainty claimed for their value ($0.6 \text{ kJ}\cdot\text{mol}^{-1}$ or ~ 0.6 percent) is probably too small, based on expected uncertainties of at least 1 percent for the method used. Consequently, we conclude that the agreement is good. In contrast, the value reported by Sabbah and El Watik [7] is $5 \text{ kJ}\cdot\text{mol}^{-1}$ lower than that derived here. This value is either in error or the uncertainty reported by Sabbah and El Watik [7] is more than one order of magnitude too small.

A value for $\Delta_{\text{cr}}^{\text{g}}H_{\text{m}}$ at $T/\text{K} = 298.15$ was reported by Freitas *et al.*[8] based on measurement of enthalpy differences between the gas at temperature $T/\text{K} = 417$ and

crystals at $T/K = 298.15$ with Calvet microcalorimetry, combined with a computed enthalpy for the gas phase for the interval $T/K = 298.15$ to $T/K = 417$. The value reported by Freitas *et al.*[8] is in good agreement with that of this research, as seen in Table 7. The origin of the uncertainty given by Freitas *et al.*($2.3 \text{ kJ}\cdot\text{mol}^{-1}$) is not clear, because the reported uncertainty for their measured enthalpy difference is $1.3 \text{ kJ}\cdot\text{mol}^{-1}$. Additional uncertainty may have been added due to unknown reliability of the computational methods used. The computational methods used in the present research have been validated [1-6] and shown to produce gas-phase properties with small uncertainties, so we revised the value of $\Delta_{\text{cr}}^{\text{g}}H_{\text{m}}$ at $T/K = 298.15$ reported by Freitas *et al.*[8] by substituting the computational results of the present research and decreasing the uncertainty. The revised value (104.2 ± 1.3) $\text{kJ}\cdot\text{mol}^{-1}$ is in excellent agreement with that of the present research (103.26 ± 0.4) $\text{kJ}\cdot\text{mol}^{-1}$, justifying the revised smaller uncertainty.

4.3 Assessment of literature enthalpies of combustion

The enthalpy of combustion for xanthone has been studied with oxygen-bomb calorimetry by Freitas *et al.*[8], Sabbah and El Watik [7], and Kim *et al.*[10]. The values reported by Freitas *et al.* and Sabbah and El Watik are in excellent agreement, while the value reported by Kim [10] is $5 \text{ kJ}\cdot\text{mol}^{-1}$ higher. The value reported by Kim *et al.*[10] includes a significant “impurity correction” that we believe was added in error. The sample used by Kim *et al.*[10] was carefully purified by crystallization from ethanol and dried, as was the sample in the present research. Kim *et al.* also used a gas chromatograph/mass spectrometer to look for organic impurities, and none were found. In addition, no mention of an impurity correction is given in the text of the article. We

recalculated the enthalpy of formation of xanthone with the experimental results reported by Kim *et al.*[10], but without the “impurity correction,” and the revised value is given at the bottom of table 8. Agreement of the revised value with those of Freitas *et al.*[8] and Sabbah and El Watik [7] is excellent; within 0.6 kJ·mol⁻¹, in contrast to a difference of 5 kJ·mol⁻¹ with the “correction” included.

4.4 Enthalpy of formation of xanthone in the ideal-gas state

The enthalpy of formation of xanthone in the ideal-gas state is now calculated with the weighted average of the enthalpies of formation for the crystals reported by Freitas *et al.*[8], Sabbah and El Watik [7], and Kim *et al.*[10] (revised) { $\Delta_f H_m^o(\text{cr}) = -(197.0 \pm 4.0)$ }, together with the enthalpy of sublimation derived in this research { $\Delta_{\text{cr}}^g H_m = (103.6 \pm 0.4)$ kJ·mol⁻¹ (table 7)} to yield $\Delta_f H_m^o(\text{ideal gas}) = -(93.4 \pm 4.0)$.

Conclusion

Computational chemistry is a powerful tool for evaluation of thermodynamic properties in the ideal-gas state, particularly for rigid molecules, such as xanthone. In this research we have demonstrated that computed gas-phase properties with validated uncertainties can provide key information that cannot be obtained from other sources for thermodynamic-consistency analysis and literature assessments.

Acknowledgements

We acknowledge the contributions of I. Alex Hossenlopp in vapor-transfer of the sample prior to heat-capacity studies and Stephen E. Knipmeyer for maintenance of all apparatus. The authors gratefully acknowledge the Office of Fossil Energy of the U.S. Department of Energy (DOE) for financial support of the experimental studies. This

1
2
3
4 research was funded within the Processing and Downstream Operations section of the
5
6 Advanced Oil Recovery (AOR) program. The experiments were completed through
7
8 BDM-Oklahoma under its contract with DOE for Management and Operations of the
9
10 National Oil and Related Programs (NORP), Contract Number DE-AC22-94C91008.
11
12
13 Manuscript preparation and all computations were done at NIST.
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- [1] R.D. Chirico, R.D. Johnson III, W.V. Steele, J. Chem. Thermodyn. 39 (2007) 698-711.
- [2] R.D. Chirico, A.F. Kazakov, W.V. Steele, J. Chem. Thermodyn. 42 (2010) 571-581.
- [3] R.D. Chirico, A.F. Kazakov, W.V. Steele, J. Chem. Thermodyn. 42 (2010) 581-590.
- [4] R.D. Chirico, A.F. Kazakov, W.V. Steele, J. Chem. Thermodyn. 73 (2014) 241-254.
- [5] R.D. Chirico, A.F. Kazakov, W.V. Steele, J. Chem. Thermodyn. 54 (2012) 278-287.
- [6] R.D. Chirico, W.V. Steele, A.F. Kazakov, submitted to J. Chem. Thermodyn. (2014).
- [7] R. Sabbah, L. El Watik, Bull. Soc. Chim. Fr. 4 (1988) 626-630.
- [8] V.L.S. Freitas, J.R.B. Gomes, M.D.M.C. Ribeiro da Silva, J. Therm. Anal. Calorim. 97 (2009) 827-833.
- [9] M.J.S. Monte, C.A.D. Sousa, J.M.S. Fonseca, L.M.N.B.F. Santos, J. Chem. Eng. Data 55 (2010) 5264-5270.
- [10] K.Y. Kim, R.E. Winans, W.N. Hubbard, C.E. Johnson, J. Phys. Chem. 82 (1978) 402-405.
- [11] M.E. Wieser, N. Holden, T.B. Coplen, J.K. Böhlke, M. Berglund, W.A. Brand, P. De Bièvre, M. Gröning, R.D. Loss, J. Meija, T. Hirata, T. Prohaska, R. Schoenberg, G. O'Connor, T. Walczyk, S. Yoneda, X-K. Zhu, Pure Appl. Chem. 85 (2013) 1047-1078.
- [12] 2010 CODATA Recommended Values: <http://physics.nist.gov/cgi-bin/cuu/Value?r> (Accessed March 8, 2012).
- [13] Metrologia 5 (1969) 35-44.

-
- [14] R.N. Goldberg, R.D. Weir, *Pure Appl. Chem.* 64 (1992) 1545-1562.
- [15] F.L. McCrackin, S.S. Chang, *Rev. Sci. Instrum.* 46 (1975) 550-553.
- [16] W.V. Steele, D.G. Archer, R.D. Chirico, W.B. Collier, I.A. Hossenlopp, A. Nguyen, N.K. Smith, B.E. Gammon, *J. Chem. Thermodyn.* 20 (1988) 1233–1264
- [17] J.P. McCullough, G. Waddington, *Anal. Chim. Acta* 17 (1957) 80-96.
- [18] E.F. Westrum Jr., G.T. Furukawa, J.P. McCullough, in: J.P. McCullough, D.W. Scott (Eds.), *Experimental Thermodynamics*, vol. 1, Butterworths, London, 1968 (Chapter 5).
- [19] M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H. P. Hratchian, A. F. Izmaylov, J. Bloino, G. Zheng, J. L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J. A. Montgomery, Jr., J. E. Peralta, F. Ogliaro, M. Bearpark, J. J. Heyd, E. Brothers, K. N. Kudin, V. N. Staroverov, T. Keith, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J. C. Burant, S. S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J. M. Millam, M. Klene, J. E. Knox, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, R. L. Martin, K. Morokuma, V. G. Zakrzewski, G. A. Voth, P. Salvador, J. J. Dannenberg, S. Dapprich, A. D. Daniels, O. Farkas, J. B. Foresman, J. V. Ortiz, J. Cioslowski, and D. J. Fox, *Gaussian 09*, Revision C.01, Gaussian, Inc., Wallingford CT, 2010.
- [20] R. Zwarich, O.S. Binbrek, *Spectrochimica Acta A*, 41 (1985) 537-544.
- [21] H.K. Sinha, L. Chantranupong, R.P. Steer, *J. Mol. Spectrosc.* 169 (1995) 302-314.
- [22] R.E. Connors, C.M. Fratini, *J. Mol. Struct.* 553 (2000) 235-241.

2
3
4
5 **TABLE 1**
6 **Calorimeter and sample characteristics for adiabatic calorimetric studies on xanthone ^a**
7

18

19

20 m / g	47.893
21 $V_{\text{i}}(298.15 \text{ K}) / \text{cm}^3$	61.78
22 $T_{\text{cal}} / \text{K}$	298.2
23 $p_{\text{cal}} / \text{kPa}$	4.53
24 $r(T_{\text{max}})$	3.1
25 r_{min}	1.6
26 x_{pre}	0.0001

27
28
29
30
31
32

33

34

35 ^a m is the sample mass; V_{i} is the internal volume of the calorimeter vessel; T_{cal} is the

36 temperature of the calorimeter when sealed; p_{cal} is the pressure of the helium and sample

37 when sealed; $r(T_{\text{max}})$ is the ratio of the heat capacity of the full calorimeter to that of the

38 empty at the highest temperature $T_{\text{max}} \approx 524 \text{ K}$ of these measurements; r_{min} is the

39 minimum value of r observed in this study; and x_{pre} is the mole-fraction impurity used for

40 pre-melting corrections.

41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

TABLE 2
Melting-study summary for xanthone ^a

<i>F</i>	<i>T</i> /K
0.2046	448.755 ₀
0.4047	448.774 ₅
0.5547	448.780 ₁
0.7048	448.783 ₃
0.8548	448.785 ₀
<i>T</i> _{tp} = (448.79 ₅ ± 0.02) K ^b	
<i>x</i> = 0.9999	

^a *F* is the fraction melted at observed temperature *T*, *T*_{tp} is the triple-point temperature, *x* is the derived mole-fraction purity of the sample.

^b The uncertainty is the expanded uncertainty with 0.95 level of confidence.

TABLE 3

Molar heat capacities $C_{\text{sat,m}}$ at vapor-saturation pressure measured with adiabatic calorimetry for xanthone ($R = 8.3144621 \text{ J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$) ^a

N^b	$\langle T \rangle / \text{K}$	$C_{\text{sat,m}} / R^c$	$\Delta T / \text{K}$	N^b	$\langle T \rangle / \text{K}$	$C_{\text{sat,m}} / R^c$	$\Delta T / \text{K}$
crystal(II)							
10	4.948	0.050	1.102	2	107.097	10.257	9.739
10	5.884	0.088	0.892	2	116.862	10.994	9.773
10	6.774	0.136	1.018	2	126.790	11.749	10.021
10	7.756	0.202	1.063	2	136.843	12.522	9.896
10	8.886	0.300	1.120	2	146.729	13.295	9.866
10	10.012	0.420	1.069	2	156.734	14.087	10.070
10	11.155	0.554	1.168	2	166.814	14.902	10.015
10	12.420	0.709	1.327	2	176.933	15.728	10.106
10	13.824	0.911	1.480	2	187.026	16.571	10.078
10	15.408	1.133	1.752	2	197.157	17.433	10.184
10	17.209	1.398	1.857	2	207.332	18.309	10.167
10	19.141	1.687	2.005	2	217.492	19.195	10.158
10	21.272	1.999	2.251	2	227.644	20.098	10.154
10	23.691	2.359	2.582	2	237.794	20.998	10.155
10	26.445	2.758	2.922	2	247.000	21.829	10.185
10	29.516	3.185	3.219	2	257.215	22.770	10.193
10	32.987	3.649	3.686	2	267.433	23.722	10.200
10	36.761	4.125	3.865	8	271.494	24.108	10.097
10	40.808	4.592	4.231	2	277.658	24.697	10.212
10	45.210	5.087	4.575	8	281.579	25.098	10.111
10	49.987	5.579	4.980	9	285.783	25.491	8.203
10	55.295	6.089	5.638	2	287.891	25.693	10.220
10	61.275	6.630	6.320	8	287.931	25.716	2.998
2	67.424	7.162	6.111	8	290.919	26.012	2.978
2	74.047	7.704	7.121	8	293.888	26.303	2.960
2	81.422	8.298	7.617	8	296.837	26.623	2.939
2	89.275	8.903	8.081	8	299.768	26.967	2.921
2	97.771	9.553	8.904	8	302.679	27.291	2.902
crystal(I)							
8	308.544	26.522	2.944	3	416.688	35.189	10.198
9	314.516	27.030	8.309	12	419.613	35.413	16.583
2	319.049	27.448	10.319	2	422.488	35.639	10.229
2	329.404	28.250	10.373	6	426.255	35.934	10.285
2	339.755	29.101	10.317	3	426.893	35.967	10.173
2	350.053	29.937	10.266	4	428.370	36.089	10.048
2	360.249	30.758	10.111	2	432.603	36.416	9.990
2	370.496	31.596	10.397	12	434.538	36.592	13.201
2	380.935	32.427	10.464	6	436.392	36.756	10.132

2	391.393	33.242	10.426	7	436.865	36.790	10.113
2	401.806	34.052	10.392	3	437.001	36.762	9.921
12	403.537	34.188	15.537	4	438.344	36.883	9.892
6	411.031	34.759	20.544	2	441.529	37.115	7.847
2	412.187	34.846	10.363	3	444.475	37.376	4.804
				liquid			
4	456.708	42.811	9.225	13	488.563	44.410	12.455
12	456.822	42.825	9.776	14	497.530	44.856	12.387
7	456.898	42.826	9.404	13	500.955	45.038	12.380
13	465.069	43.247	11.668	13	512.381	45.606	10.385
4	465.905	43.275	9.169	13	521.082	46.026	6.575
13	476.597	43.820	11.583				

^a The relative expanded uncertainty for the heat capacities $U_r(C_{\text{sat,m}}) = 0.03$ at $T/\text{K} = 5$, 0.005 at $T/\text{K} = 10$, and 0.001 for temperatures $T/\text{K} \geq 30$ with 0.95 level of confidence. The standard uncertainty in the mean temperatures $u(\langle T \rangle)/\text{K} = 0.01$.

^b Adiabatic series number.

^c Average heat capacity for a temperature increment of ΔT with a mean temperature $\langle T \rangle$.

TABLE 4

Measurements of molar energy increment $\Delta_{\text{tot}}U_{\text{m}}$ and derived results for xanthone

N^a	h^b	T_i	T_f	T_{trs}	$\Delta_{\text{tot}}H_{\text{m}}^c$	$\Delta_{\text{trs}}H_{\text{m}}^d$
		K	K	K	$\text{kJ}\cdot\text{mol}^{-1}$	$\text{kJ}\cdot\text{mol}^{-1}$
single-phase measurements in phase cr(II)						
8	1	205.300	266.467		10.605	0.003
11	1	61.168	119.955		4.388	-0.001
11	1	119.967	203.716		10.122	0.000
11	1	203.667	290.036		15.703	0.003
cr(II)-to-cr(I) phase transition						
9	1	289.885	310.371	304.50	4.532	0.000
11	1	290.047	310.534		4.533	0.000
					average	0.000
single-phase measurements in phase cr(I)						
7	1	330.388	396.576		17.071	0.002
7	1	396.537	431.805		10.263	0.000
11	1	310.986	392.846		20.477	0.002
11	1	392.858	440.913		14.070	0.000
cr(I)-to-liquid phase transition						
4	2	443.285	452.097	448.79 ₅	30.362	27.478
7	6	441.931	452.195		30.813	27.475
12	2	441.196	451.925		30.941	27.471
					average	27.475
single-phase measurements in the liquid phase						
14	1	451.180	491.439	---	14.577	0.000

^a Adiabatic series number.^b Number of heating increments.^c $\Delta_{\text{tot}}H_{\text{m}}$ is the molar energy input from the initial temperature T_i to the final temperature T_f . The relative expanded uncertainty for $\Delta_{\text{tot}}H_{\text{m}}$ is 0.001 with 0.95 level of confidence.^d $\Delta_{\text{trs}}H_{\text{m}}$ is the net molar enthalpy of transition at the transition temperature T_{trs} or the excess enthalpy for single-phase measurements relative to the heat-capacity curve described in the text and defined in table 5.

TABLE 5

Molar thermodynamic functions at vapor-saturation pressure for xanthone

 $(R = 8.3144621 \text{ J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1})^a$

T/K	$C_{\text{sat,m}}/R$	$\Delta_0^T S_{\text{m}}/R$	$\Delta_0^T H_{\text{m}}/RT$	T/K	$C_{\text{sat,m}}/R$	$\Delta_0^T S_{\text{m}}/R$	$\Delta_0^T H_{\text{m}}/RT$
crystal(II)							
5.00	0.054	0.018	0.014	140.00	12.768	13.045	6.882
10.00	0.418	0.143	0.107	160.00	14.350	14.852	7.716
20.00	1.813	0.843	0.598	180.00	15.983	16.636	8.543
30.00	3.251	1.854	1.247	200.00	17.677	18.407	9.372
40.00	4.502	2.967	1.909	220.00	19.418	20.173	10.205
50.00	5.581	4.090	2.539	240.00	21.196	21.939	11.047
60.00	6.517	5.192	3.125	260.00	23.028	23.708	11.898
70.00	7.375	6.262	3.672	280.00	24.930	25.483	12.760
80.00	8.185	7.300	4.186	298.15	26.772	27.105	13.556
90.00	8.960	8.309	4.673	300.00	26.980	27.271	13.638
100.00	9.722	9.292	5.140	304.50 ^b	27.521	27.677	13.839
120.00	11.232	11.198	6.029				
crystal(I)							
304.50 ^b	26.150	27.677	13.839	400.00	33.912	35.841	17.722
320.00	27.516	29.009	14.469	420.00	35.441	37.532	18.529
340.00	29.121	30.726	15.284	440.00	37.021	39.218	19.334
360.00	30.740	32.436	16.097	448.79 ₅ ^b	37.687	39.957	19.687
380.00	32.354	34.142	16.911				
liquid							
448.79 ₅ ^b	42.422	47.320	27.050	500.00	44.990	52.039	28.756
460.00	42.977	48.373	27.431	520.00	45.975	53.823	29.399
480.00	43.988	50.223	28.100				

^a Relative expanded uncertainties for all properties are 0.03 at $T = 5 \text{ K}$, 0.005 at $T = 10 \text{ K}$, and 0.001 for temperatures $T \geq 30 \text{ K}$ with 0.95 level of confidence.

^b Values at this temperature were calculated with heat capacities extrapolated linearly.

TABLE 6

Calculated fundamental vibrational wavenumbers $\bar{\nu}$ used in the computation of the ideal-gas entropies for xanthone ^a

$\bar{\nu}/\text{cm}^{-1}$	$\bar{\nu}/\text{cm}^{-1}$	$\bar{\nu}/\text{cm}^{-1}$	$\bar{\nu}/\text{cm}^{-1}$	$\bar{\nu}/\text{cm}^{-1}$
61.7	535.5	873.3	1208.4	1609.6
116.0	585.0	920.7	1210.5	1620.7
145.0	624.6	955.5	1234.2	1680.4
225.6	647.1	957.3	1269.4	3111
238.6	665.0	978.7	1317.8	3111
281.4	693.4	979.6	1341.1	3124
311.2	708.6	1023.0	1346.9	3124
375.1	751.0	1027.1	1457.3	3135
409.4	758.8	1096.7	1457.7	3135
434.9	792.4	1105.4	1467.9	3139
448.4	826.4	1143.3	1478.8	3139
513.8	858.4	1146.9	1570.4	
515.2	862.4	1170.6	1591.9	

^a The scale factor 0.975 was used in the calculation of the listed wavenumbers. The rotational constants are $A = 0.04858 \text{ cm}^{-1}$, $B = 0.01543 \text{ cm}^{-1}$, and $C = 0.01171 \text{ cm}^{-1}$, and the symmetry number for xanthone is 2.

TABLE 7

Comparison of literature enthalpies of sublimation and values derived in this research.

T/K	$\Delta_0^T S_m(\text{cr})/R^a$	S_{comp}/R^b	$\Delta_0^T S_m^o(\text{ideal})/R^c$	$\Delta_{\text{cr}}^g S_m/R^d$	$\Delta_{\text{cr}}^g H_m/(\text{kJ}\cdot\text{mol}^{-1})$
352.2	(31.77 ± 0.03)	(-12.53 ± 0.10)	(54.11 ± 0.11)	(34.87 ± 0.13)	$(102.1 \pm 0.4)^e$
				Monte <i>et al.</i> [9]	(103.9 ± 0.6)
				Sabbah and El Watik [7]	(97.1 ± 0.3)
298.15	(27.11 ± 0.03)	(-18.97 ± 0.10)	(49.93 ± 0.10)	(41.80 ± 0.14)	$(103.6 \pm 0.4)^e$
				Freitas <i>et al.</i> (original) [8]	$(102.7 \pm 2.3)^f$
				Freitas <i>et al.</i> (revised here)	$(104.2 \pm 1.3)^g$

^a Standard entropy of crystalline xanthone interpolated from table 5.^b Entropy of compression calculated with the sublimation pressures p reported by Monte *et al.*[9]. S_{comp} is $\ln(p/p^\circ)$, where p° is 101.325 kPa.^c Standard entropy of the ideal gas computed with parameters given in table 6 ($p^\circ = 101.325$ kPa) and the methods described in the text.^d Entropy of sublimation calculated with the values given in this table and equation 1.^e Enthalpy of sublimation $\Delta_{\text{cr}}^g H_m$ calculated with $T \cdot \Delta_{\text{cr}}^g S_m$.^f Enthalpy of sublimation based on the calorimetric results and computations reported by Freitas [8]^g Enthalpy of sublimation based on the calorimetric results by Freitas [8] and the computed gas-phase enthalpies of the present research.

TABLE 8

Comparison of enthalpies of formation based on measured enthalpies of combustion for xanthone reported in the literature ^a

Literature Source (Year)	$\Delta_f H_m^o(\text{cr})/(\text{kJ}\cdot\text{mol}^{-1})$ ^a
Freitas <i>et al.</i> (2009)	$-(196.7 \pm 4.0)$
Sabbah and El Watik (1988)	$-(196.8 \pm 3.6)$
Kim <i>et al.</i> (1974)	$-(191.5 \pm 2.7)$
Kim <i>et al.</i> (revised here)	$-(197.3 \pm 2.7)$ ^b

^a Uncertainties are those reported in the original sources. These are equivalent to the expanded uncertainty with 0.95 level of confidence.

^b This value was calculated with the experimental results reported by Kim *et al.*[10], but without the “impurity correction” that appears to have been applied in error.

Figure

FIGURE 1. Plot of molar heat capacities against temperature at saturation pressure $C_{\text{sat,m}}$ for xanthone measured in this research. The vertical lines indicate the cr(II)-to-cr(I) transition temperature T_{tr} and triple-point temperature T_{tp} . $\circ$, Experimental values determined in this research.