Disentangling fine- and broad- scale effects of habitat on predator-prey interactions

D.H. Chacin *, C.D. Stallings

College of Marine Science, University of South Florida, 140 7th Ave S, St. Petersburg,

Florida 33701, USA

* Corresponding author

E-mail address: dchacin@mail.usf.edu

24 ABSTRACT

25 Predator-prey interactions can be influenced by habitat at different spatial scales. In
26 seagrass systems, blade density can provide refugia for prey at fine scales, which are further
27 embedded within broad-scale features such as variation in biotic (e.g., predator
28 assemblages) and abiotic attributes (e.g., turbidity, salinity). Fine-scale effects of seagrass
29 habitats on predator-prey interactions involving invertebrates have been well studied while
30 less is known about their effects on fish as prey. A field experiment was conducted in
31 Tampa Bay, Florida, USA to examine and separate the effects of habitat across fine and
32 broad scales on the relative predation rates of tethered Pinfish (*Lagodon rhomboides*).
33 Artificial seagrass units (ASUs) were used at three levels of blade density and deployed in
34 different locations within the seascape. Predation rates on Pinfish decreased with increasing
35 seagrass blade density. The effects of blade density were consistent across locations, but
36 overall mortality was higher in the lower Bay, where the water was less turbid, higher in
37 salinity, and characterized by a different suite of predators compared to the mid Bay. Using
38 controlled-laboratory experiments, it was found that Pinfish reduced their activity levels in
39 more turbid water as well as in response to the presence of a common predator in both clear
40 and more turbid waters. Thus, predation rates were influenced by the combined effects of
41 refugia (fine scale), variation in prey behavior (broad scale), and detection by predators
42 (both scales). This study demonstrates the strong influence habitat can have at different
43 spatial scales in mediating predator-prey interactions of mobile species in estuarine
44 environments.

45 **Keywords:** Water clarity · Predation risk · Mortality · Submerged aquatic vegetation ·
46 Landscape

1. Introduction

The strength of ecological processes, such as predation, can be contextual upon location, thus requiring a landscape perspective for their study. In the marine environment, seagrass beds are appropriate for using a broad-scale approach because they cover extensive areas of coastal habitats (Robbins and Bell, 1994; Bell et al., 2006), support an abundant and diverse community of fishes and invertebrates (Hemminga and Duarte, 2000; Gillanders 2006, Stallings et al., 2015), and can have high levels of heterogeneity (Bell et al., 1995; Bowden et al., 2001; Balestri et al., 2003). Broad-scale studies in seagrass beds have mainly focused on assessing faunal responses to geomorphic and geospatial habitat features such as patchiness, fragmentation, configuration, and proximity to other habitats (Bell and Hicks, 1991; Bell et al., 2001; Healey and Hovel, 2004; Bell et al., 2006; Heck and Orth, 2006; Larkum et al., 2006; Carroll and Peterson, 2013).

Other ecological (e.g., predator communities) and environmental features (e.g., water turbidity) can also vary across the seascape in seagrass beds, with potential effects on predator-prey dynamics. These variations in the seascape often result in discrepancies observed when experiments are repeated at different locations in the field. For example, both survival and recolonization of mussels were contextual upon the spatial scale and location in the rocky intertidal where identically-designed experiments were conducted (Menge et al., 1994). Similarly, Menge (1976) observed several aspects of community structure (such as abundance of predators and species richness) to vary across six areas of intertidal communities in Maine and Massachusetts spanning a wave exposure gradient.

Embedded within these broad-scale features, fine-scale variation in the complexity of seagrass habitats (e.g., blade density) can further influence predator-prey dynamics.

Structurally complex habitats can provide physical and visual barriers that can reduce predation risk (Stoner and Lewis, 1985; Wahle and Steneck, 1991; Beukers and Jones, 1998; Pirtle et al., 2012) and this has been a rich area of study in seagrass systems (Irlandi 1994, Murphey and Fonseca, 1995; Irlandi, 1997; Hovel and Fonseca, 2005). At fine spatial scales the survival of organisms varies with the presence, density, or biomass of seagrass blades (Heck and Crowder, 1991; Orth, 1992; Irlandi, 1994; Rooker et al., 1998; Hovel and Fonseca, 2005). Most of these studies, however, have been conducted in controlled-laboratory settings or have largely focused on invertebrates in the field (Heck and Thoman, 1981; Orth et al., 1984; Main, 1987; Gotceitas, 1990; Heck and Crowder, 1991; Hovel and Lipcius, 2002; Heck et al., 2003).

Fishes are an important component of seagrass bed communities, both as predators and prey (Heck and Orth, 2006) and thus may also be influenced by complex seagrass habitats. In the laboratory, Gotceitas (1997) observed that the survival of Atlantic Cod (*Gadus morhua*) increased with eelgrass density only after a threshold of seagrass density was reached. Harris et al. (2004) found that refuge from predation for Tautog (*Tautoga onitis*), Cunnners (*Tautogolabrus adspersus*), and Silversides (*Brevoortia tyrannus*) can be modeled as an asymptotic function of plant density. Others have found no differences in mortality between vegetated and unvegetated habitat types (Laurel and Brown, 2006; Horinouchi, 2007a). Additionally, macrophyte complexity in littoral zones of lakes has been identified as an important factor in reducing predation and altering behavior in predator-prey interactions among fishes (Savino and Stein, 1982; Gotceitas, 1990). While most of these studies suggest that seagrass complexity may influence the survival of fishes, the results are equivocal. Furthermore, few studies have focused on how blade density may directly affect

mortality of fishes in the field (Horinouchi, 2007b; Schultz and Kruschel, 2010), thus requiring further research *in situ* to better understand generalities.

Tampa Bay is a large, shallow embayment with approximately 12,000 hectares of seagrass coverage (Greening et al., 2011). Because of its large size, it has a great potential for variation in biotic and abiotic factors over multiple spatial scales. These features present a suitable opportunity to evaluate the integration of broad- and fine- scale effects of seagrass habitats on predator-prey interactions. This study examined the broad-scale effects of location and fine-scale effects of seagrass complexity (via blade density) on predation rates of a common marine fish. Specifically, a field experiment was used to examine 1) the broad-scale effects of distinct locations of the Bay and 2) the fine-scale effects of seagrass blade density (high, medium, and none) on the relative mortality rates on Pinfish (*Lagodon rhomboides*; Family Sparidae). A laboratory experiment was also used to examine how the behavior of Pinfish varied with exposure to different biotic and abiotic features. By manipulating fine-scale complexity of seagrass habitats deployed in locations embedded in broad-scale seascapes, it was sought to provide insight on how ecological processes can be affected by habitat over multiple spatial scales.

2. Materials and methods

2.1. Study organism

The Pinfish inhabits coastal waters from the Yucatan Peninsula of Mexico eastward through the Gulf of Mexico and up to Massachusetts, USA (Hoese and Moore, 1977; Darcy, 1985; Nelson, 2002; Harter and Heck, 2006). Pinfish are an ecologically important

species for several reasons. First, they are commonly the most abundant vertebrate found in seagrass beds (Nelson et al., 2013; Stallings et al., 2015). Second, juveniles feed on a suite of invertebrates influencing entire assemblages of benthic macrofauna (Young et al., 1976; Young and Young, 1977; Nelson, 1978; Stoner, 1980; Nelson, 2002). Third, Pinfish are an important link between primary and secondary production due to their consumption of plant matter (Stoner, 1982; Weinstein et al., 1982; Nelson, 2002). Because all life stages are preyed upon by other fishes (Kjelson and Johnson, 1976; Seaman and Collins, 1983; Nelson, 2002; Stallings, 2010; Stallings et al., 2010; Hall-Scharf and Stallings, 2014; Hall-Scharf et al., in press) they also serve as an important linkage to offshore food webs (Nelson et al., 2013).

In Florida, adult Pinfish spawn in offshore waters from late fall to early winter and have peak larval settlement in February and March (Tabb and Manning, 1961; Chacin thesis, 2014). Post-larval Pinfish settle in a variety of habitats including shallow and vegetated flats where juveniles remain most of their first year. These vegetated flats (mostly composed of seagrass beds) are nearshore habitats often located in estuarine environments, where biotic and abiotic factors can be highly variable.

2.2. Study region

This study was conducted in Tampa Bay, Florida, USA (Fig. 1), during the months of July through September 2013. Tampa Bay is a large, open-water estuary with a surface area of ~1000 km² and mean water depth of ~4.0 m (Chen et al., 2007). Two locations of the Bay were selected for this study, the middle-west and south-west, which we hereafter refer to as “mid Bay” and “lower Bay,” respectively. These two locations were chosen because

they have the highest densities of Pinfish observed in the Bay (Chacin thesis, 2014). *In situ* tethering experiments were deployed at each location.

2.3. Field experiment – study design

An orthogonally designed field experiment was used to test whether Pinfish experienced different predation rates under different habitat complexities (via three blade density levels) cross-factored with different locations of the Bay. Artificial seagrass units (ASUs) were used to control for the density of seagrass blades, which were constructed on plastic mesh (0.5 m x 0.5 m squares) with 1,500 shoots m² (high), 600 shoots m² (medium), and 0 shoots m² (none); these values were meant to simulate *Thalassia testudinum* densities found in Tampa Bay (Meyer and Bell, 2010). Each shoot was composed of two 30 cm long blades made with 1 cm wide green polypropylene ribbon tied to the plastic mesh. Pinfish (size range standard length (SL): 50 – 80 mm) were tethered using a 0.25 m microfilament braided line (18.1 kg test) connected to a 2 m long microfilament braided line attached to the center of the ASUs (Fig. 2). A small plastic float was attached on the other end of the 2 m braided line to keep it vertical in the water column. Although the sizes of the ASUs were small, they allowed for replication and experimental control to determine mechanism, and were large enough to allow the Pinfish space to swim both horizontally and vertically in a 0.39 m³ cylinder. Pinfish used for the study were collected using seine and cast nets, and immediately tethered after capture. All fishes other than the study species were released instantly upon capture.

The deployment of the ASUs required three steps. First, the plot was fixed to the substrate with metal stakes (one in each corner) on the edge of the natural and continuous

seagrass bed. By retaining a constant edge position, rather than confounding with interior habitat locations, it was assumed that if edge effects occurred, they were consistent across trials. Second, the tether line was secured to a Pinfish by inserting it through the mouth and out the opercular opening to form a loose loop forward of its snout. The entire process was fairly rapid (~ 10-15 sec) and the tethered Pinfish was quickly returned to the water without the need for recovery from anesthesia, which could have artificially inflated mortality. This same approach was previously used to tether Pinfish in Biscayne Bay, Florida, USA (Hammerschlag et al., 2010) and is advantageous over more invasive methods (e.g., running tether through soma) as it minimizes tissue damage and the release of body fluids, which could increase detection by predators. Last, the tether was secured to the centerline using a loose loop, allowing Pinfish to move in a vertical cylinder with a radius of 0.25 m (Fig. 2) but preventing it from accessing adjacent habitats. If bending of the line occurred despite the preventions used, it was assumed it occurred equally across all treatments.

During each tethering event, fifteen plots were deployed with five replicates for each blade density (high, medium, none) randomly arranged at the seagrass edges to avoid potential confounding effects of predation differences between edge and interior locations (Peterson et al., 2001). Deployments were always positioned at the shoreward edge to avoid potential predation differences due to variation in fish abundances known to occur at shoreward versus seaward edges (Smith et al., 2008) and were conducted at both locations throughout the experimental months.

Although tethering studies cannot produce estimates of *absolute* mortality rates, given the potential artifacts of the approach, they are powerful tools for comparing relative mortality among experimental treatments. Pilot deployments were video recorded and

analyzed for potential alterations of Pinfish behavior caused by tethering and artificial seagrass units. Pinfish swimming behavior was not altered; they readily swam both horizontally and vertically through the water column. It was concluded that although the distance Pinfish could swim was restricted, the tether did not hamper their hiding and swimming behaviors. No losses from Pinfish breaking free from the tether during the pilot deployments (five per seagrass blade density level) were observed, nor were there any entanglements during either the pilot trials or when tethers were retrieved. Therefore, it was assumed that any losses in tethered fish were due to predation events.

A series of pilot deployments were conducted to determine optimal experimental time. It was found that 10-hr deployments were sufficiently long given the observed mortality rates while also short enough to avoid potential confounding effects of increased predation during crepuscular periods. Deployments began two hours after sunrise and ended two hours before sunset (0800-1800). Deployments were replicated a minimum of 15 times for each ASU density * location combination, with one tethered fish per individual deployment. The absence of the Pinfish or the presence of an injured fish at the conclusion of the deployment was counted as a predation event.

2.4. Field experiment – data analysis

Differences in mortality rates were compared separately across 1) seagrass density (three levels) and 2) location (two levels) using non-parametric analysis of variance (npANOVA). The non-parametric ANOVAs were permutation-based and distribution-free, due to data not meeting assumptions of normality and to avoid concerns about zero inflations; this method is equivalent to a univariate PERMANOVA. Pair-wise comparisons

were included for the tests involving seagrass densities. For all analyses, a dissimilarity matrix was produced for the response variable (mortality) based on Euclidean distance using Fathom toolbox for Matlab (Jones, 2014). A two-way npANOVA was conducted to test for the interactions between seagrass density with location on mortality. A one-way ANOVA was used (in addition to the two-way) to perform the pairwise comparisons among seagrass densities and also because one of the factors is kept constant when a two-way ANOVA is performed, yielding inaccurate *F*-statistics. The level of significance for all tests was based on an alpha value of 0.05.

2.5. Exploration of other abiotic and biotic variables

Because differences in Pinfish mortality were observed between study locations (see Results), potential correlated biotic and abiotic variables that could have contributed to the results were explored. Seagrass habitats were chosen at both study locations that were dominated by *Thalassia testudinum*, with lower and similar compositions of *Syringodium filiforme* and *Halodule wrightii*. Canopy heights of the beds were 25-35 cm. Both locations contained continuous seagrass cover, with a bed area ~ 600 m² and had an average depth of 1.9 m. Locations also had similar adjacent land habitat (thin sand/vegetation mix set in an urban area). Other variables, however, that could have influenced the results of the experiments, such as turbidity, temperature, salinity, and predatory assemblage were explored.

The experiments were restricted to the summer season in Tampa Bay, when a turbidity gradient is the strongest with higher values in the upper and mid Bay and lower values in the lower Bay. It was assumed that since juvenile Pinfish remain in the same

relative area prior to egress, they experienced consistently different turbidities depending on which general location of the Bay they occupied. Turbidity levels during the study were quantified with data provided by the Environmental Protection Commission of Hillsborough County's (EPCHC) Tampa Bay water quality monitoring program. Water samples were collected at mid depth of a station if it was over 3 m in depth, otherwise surface water samples were collected. Turbidity of water samples was measured with Hach® Model 2100N Turbidimeter. Turbidity values were then averaged over the summer months (corresponding to the months when experiments were conducted) and were compared using paired *t*-tests to determine if turbidity varied between study locations.

The remaining variables explored were obtained from Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute (FWRI). The state agency's Fishery Independent Monitoring Program (FIM) conducts sampling trips each month throughout Tampa Bay (and other locations in Florida) and thus can provide the most extensive database for determining spatial dynamics in the assemblages of Pinfish predators as well as environmental variables at their sampling locations (e.g., temperature, salinity). The FIM data were collected with a 6.1 m otter trawl, towed over an area of 4,120 m². All data used for the analyses were collected at the same locations and time period of the tethering experiment. Since predation is one of the primary drivers of juvenile fish mortality, it was explored whether there were differences in the predatory assemblages that could have influenced the patterns observed between the study locations. Although there was no direct way to determine which species consumed the tethered Pinfish, all fishes with reported evidence of piscivory that were ≥ 180 mm (SL) as a conservative size due to gape limitation were included. These characteristics narrowed the analysis to include predators

that most likely consume Pinfish. Since the predator data were zero-inflated and did not meet assumptions of normality, a permutation-based two-tailed t -test was conducted to determine whether species richness and density of predators differed between study locations. Additionally, predator assemblage composition was compared between the study locations using a permutation-based MANOVA followed by a Canonical Analysis of Principal Coordinates (CAP) to create a bi-plot and correlation vector plot to visualize which species, if any, were driving the observed differences. Salinity and temperature values were averaged over the summer months (similarly to turbidity) and compared with paired t -tests.

2.6. Laboratory experiment – study organisms

Previous studies on fishes have suggested that turbidity levels below 10 NTUs may not influence behaviors and interactions (Meager and Batty, 2007; Meager et al., 2006; De Robertis, 2003). It is possible that turbidity effects on behavior could be species dependent. Although turbidity was < 5 NTU during the study, it was statistically different between regions of the Bay (see Results), and the potential effects it may have had on anti-predatory behaviors of the visually-oriented Pinfish were intriguing. Therefore, in addition to the field experiment, two laboratory experiments were conducted to examine the behavior of Pinfish 1) under two turbidity levels (low and high) and 2) in response to visual cues from a common predator under different levels of turbidity.

The predatory species used was Spotted Seatrout (*Cynoscion nebulosus*). Spotted Seatrout is an inshore fish found in coastal waters from the Yucatan Peninsula of Mexico to the Chesapeake Bay, USA. They inhabit the same vegetated areas as Pinfish (e.g., seagrass

beds) and a substantial portion of their diets include Pinfish (FWRI-FIM, unpublished data; Hall-Scharf and Stallings, 2014; Hall-Scharf et al., in press).

Pinfish and Spotted Seatrout were collected using seine nets. They were placed in 40-l holding tanks containing aerated seawater and were transported to the aquarium laboratory at the University of South Florida, College of Marine Science. Once in the lab, Pinfish were held in 621-l tanks and Spotted Seatrout in 1,200-l tanks until the time of the experimental trials. Each tank was supplied with flow-through water from Tampa Bay, with aerators to maintain constant oxygenation ($DO \geq 6.5$ mg/l) and heaters to ensure a constant temperature (26.7–28.3 °C).

2.7. Laboratory experiment – study design

Turbidity treatments were prepared by mixing artificial seawater (ASW; from Instant Ocean ® Sea Salt product) with Kaolin powder clay in trial tanks. Two turbidity levels were produced: low (0 grams Kaolin per liter) and high (0.13 grams Kaolin per liter). The turbidity measurements were not represented in Nephelometric Turbidity Units (NTU) since Kaolin is formed by particles of similar size and reflectivity and therefore would differ from the field where the presence of varying suspended particles will influence the NTUs. It is recognized that this is a limitation in the study and caution must be exerted when comparing to field settings. Nevertheless, valuable insights could be gained from further examination of mechanisms in such relative conditions. Thus, the low turbidity level was meant to simulate areas where water turbidity is the lowest in Tampa Bay (e.g., lower Bay) while the high level was meant to simulate areas such the upper and mid regions of the Bay. After the turbidity preparation, individual Pinfish (size range SL: 56 – 90 mm)

were moved into the trial tanks for a 1-hr acclimation period. Lightning conditions were kept constant before, during, and among experimental trials.

Behavioral observations were conducted in rectangular, 38-l tanks filled with 19 l ASW to a water depth of 20 cm. A 15 cm x 9 cm plot of artificial seagrass was submerged in the center of the tank (Fig. 3A). The plot was used to provide a small shelter for the Pinfish to associate, whether hiding inside or adjacent to it, as are common anti-predatory behaviors. While the sizes of the tank and seagrass plots were small, Pinfish behavior did not appear to be affected. The fish were observed swimming while foraging around the tank or using the plot as shelter by hiding in it. Sixteen food pellets were evenly scattered across the bottom of the tank to encourage foraging. An air diffuser was placed at the corner of the tank to maintain constant oxygenation. A grid with dimensions 5 cm x 5 cm was placed under the trial tanks to provide a framework for measuring activity level, recorded as the number of times the experimental Pinfish crossed a line of the grid per minute (Fig. 3B). Trials were recorded with a downward-looking video camera to reduce observer effects on fish behavior.

In the first laboratory experiment, 70 10-minute trials were conducted to measure Pinfish behavior between the low ($n = 35$) and high ($n = 35$) turbidity levels in the absence of any visual cues. In the second laboratory experiment, turbidity (low vs. high) and visual cues from a predator were crossed. As controls for the predator treatment, a non-predatory conspecific (a large Pinfish) and an empty tank filled with ASW-only (i.e., no fish) were used. The order of the trials was randomized and trials were conducted during the day between 1200-1600. Preparation of visual cues consisted of placing the Spotted Seatrout (size range SL: 230 – 310 mm), the large Pinfish (SL: 140 – 150 mm), or ASW-only into a

separate tank (with the same dimensions of the trial tank) adjacent to the trial tank prior the start of the experiment (Fig. 3B). The visual cue tank was filled with ASW with no Kaolin powder present. A black barrier was placed between the trial and visual cue tanks (Fig. 3B). After a 10-minute observational period, the barrier was removed exposing the experimental Pinfish to the visual cue tank. The movements of the large Pinfish and Spotted Seatrout used as visual cues were restricted with a mesh barrier placed to guarantee they remained near the experimental tank; this was done to ensure that the experimental Pinfish had full view of the visual cue in the adjacent tank. Observations of activity levels were recorded for 10 additional minutes after the barrier was removed. Thirty-six 20-minute trials were completed (6 replicates for each turbidity * visual cue combination).

2.8. Laboratory experiment – data analysis

For the controlled laboratory experiments, the number of lines crossed per minute measured between turbidity levels (experiment 1) was compared using a two-sample *t*-test. Additionally, line crosses per minute before and after the visual stimuli (experiment 2) were compared using a paired *t*-test. Data were log transformed to meet the assumption of homoscedasticity and the level of significance for all tests was based on an alpha value of 0.05.

3. Results

In total, 136 tethering deployments were conducted across the mid and lower regions of the Bay (Table 1). There was a significant effect of seagrass density on relative mortality rates of Pinfish ($F_{2, 26} = 4.41, p = 0.021$). The posteriori, permutation-based, pair-wise

comparisons indicated that the relative mortality rates in seagrass plots with zero and medium densities of blades were not statistically different from each other ($t = 0.96, p = 0.340$). Relative mortality rates in seagrass plots with high blade density were lower than those with both zero ($t = 2.99, p = 0.009$) and medium blade densities ($t = 2.17, p = 0.042$, Fig. 4).

Relative mortality was higher for deployments made in the lower Bay ($F_{1, 27} = 18.75, p < 0.001$). Although not significant at the alpha 0.05 level, there was marginal support of an interaction between seagrass density and location ($F_{2, 23} = 0.69, p = 0.055$, Table 2, Fig. 4), suggesting that habitat structure may have been less critical under higher turbidity conditions.

During the summer months, Tampa Bay experienced a turbidity gradient with higher levels in the upper and mid regions of the Bay (3.74 ± 0.19 Nephelometric Turbidity Units; NTU $\pm$ SE) compared to the lower Bay (2.17 ± 0.15 NTU, 2000-2009, $t_{29} = 8.18, p = 0.03$). Temperature was similar between locations (mid Bay = 29.6 ± 0.20 °C and lower Bay = 30.2 ± 0.16 °C, $t_{36} = 2.76, p = 0.109$) and salinity was higher in the lower Bay (32.5 ± 2.17 ppt) than the mid Bay (26.1 ± 0.70 ppt, $t_{36} = 14.2, p = 0.001$).

There were no differences in either the abundance ($t_{40} = 0.77, p = 0.258$, Fig. 5A) or species richness of predators ($t_{40} = 0.86, p = 0.226$ Fig. 5B) between mid and lower locations of the Bay from FWRI-FIM data collected during the field experiment. The assemblage composition, however, varied between the two locations ($F_{1, 40} = 4.00, p = 0.001$, Fig. 6) with Common Snook (*Centropomus undecimalis*) and Red Drum (*Sciaenops ocellatus*) driving the patterns in the mid Bay while Gulf Flounder (*Paralichthys albigutta*) and Inshore Lizardfish (*Synodus foetens*) did so in the lower Bay.

In the behavioral study, Pinfish were typically observed swimming and foraging around the tank, which it was interpreted as an indication of normal behavior in the absence of anti-predatory responses; this occurred mainly in the low turbidity treatments. Pinfish tended to associate more closely with the ASU (either inside or directly adjacent) in the high turbidity trials, and reduced their activity levels compared to low turbidity trials ($t_{68} = 2.67$, $p = 0.010$, Fig. 7A). Pinfish also reduced their activity after the predator visual cue was presented in both turbidity levels (low $t_5 = 2.64$, $p = 0.046$ and high $t_5 = 2.81$, $p = 0.038$, Fig. 7B), but not in the conspecific control (low $t_5 = 1.58$, $p = 0.175$ and high $t_5 = 1.27$, $p = 0.260$), or ASW-only trials lacking fish (low $t_5 = 0.26$, $p = 0.808$ and high $t_5 = 0.38$, $p = 0.721$).

4. Discussion

Using an experimental approach in the field, it was observed that predation rates differed strongly among treatment levels at two habitat scales. The effects of habitat at the fine scale (blade density) on mortality were consistent between locations in Tampa Bay but overall mortality differed at broader scales (between locations). The locations where deployments were conducted were similar in many habitat-related aspects but also had broad-scale differences in turbidity, salinity, and predator assemblage that could have influenced the observed results. Additionally, through controlled-laboratory experiments it was observed that Pinfish activity levels decreased as water turbidity increased as well as when they detected the presence of a common predator. Therefore, predation rates were likely influenced by the combined effects of refugia (fine scale), variation in prey behavior

(broad scale), and detection and feeding of predators with different predatory tactics (both scales).

At the fine scale of seagrass habitats that were used in this study, relative predation rates of Pinfish were lowest on ASU plots with a high density of seagrass blades, consistent with previous studies using invertebrate macrofauna as prey and fishes as predators (Nelson, 1978; Heck and Thoman, 1981; Crowder and Cooper, 1982; Stoner, 1982; Orth et al., 1984; Heck and Crowder, 1991; Hovel and Lipcius, 2002; Heck et al., 2003). Thus, increased density of seagrass blades appeared to have provided Pinfish with protection from their predators likely due to the dense structure inhibiting efficient foraging as it has been observed in other studies (e.g., Crowder and Cooper, 1982). This study is among the few (e.g., Horinouchi, 2007a; Schultz and Kruschel, 2010) that have examined the effects of seagrass blade density on relative mortality rates of mobile fish prey *in situ*. Interestingly, predation rates were equally high in low (no blades) and medium complexity habitats, suggesting that the presence alone of seagrass blades was not enough to provide protection. These results corroborate laboratory studies that involved fish as prey. In their examination of foraging efficiency of 3-year old Atlantic Cod (*Gadus morhua*) on age-0 cod, Gotceitas et al. (1997) found that there was no difference in mortality among the lower densities of seagrass patches, but did find a significant reduction in mortality in high density patches. Harris et al. (2004) found that the mortality of three prey fishes in *Zostera marina* declined at a low-blade density but did not continue to decrease with increased densities of the seagrass. Since the field experiment in this study was only composed of three levels of blade density, it is difficult to conclude whether a complexity threshold existed. Additional levels of blade density would be required to determine the functional response, as has been

reported in other studies (Gotceitas, 1990; Nelson and Bonsdorff, 1990; Harris et al., 2004).

In contrast, others have found mortality to either decrease in a linear manner or have no relationship with increased seagrass complexity (e.g., Harris et al., 2004; Horinouchi 2007b). Notably, studies that have shown predation to decrease with an increase in plant density have observed this relationship primarily with predators that use chase-and-attack foraging tactics in the laboratory (Horinouchi, 2007a; Horinouchi, 2009). Conversely, the experiment in this study was conducted in the field, thereby exposing experimental Pinfish to the natural suite of predators that would normally prey upon them.

Pinfish mortality was higher in the lower Bay compared to the mid Bay. These results were not surprising since Tampa Bay experiences environmental and biological differences from the innermost to outermost locations of the Bay. The turbidity gradient observed at broader scales in the Tampa Bay seascape may have been one responsible factor. It is known that turbidity in Tampa Bay is influenced by a combination of organic matter, phytoplankton, and suspended solids resulting mainly from wind-driven bottom resuspension and riverine input (Johansson, 1991; Janicki and Wade, 1996; Chen et al., 2007). This is especially true in the mid and innermost regions of the Bay during the summer, which is highly influenced by river discharges. Turbidity resulting from dissolved organic matter, suspended solids, and plankton can absorb and scatter light and hence decrease the visual capacities of fishes (Gregory and Northcote, 1993; Gregory, 1993; Abrahams and Kattenfeld, 1997; Vogel and Beauchamp, 1999; Meager et al., 2006). Turbidity has been shown to influence predator-prey interactions by altering foraging efficiency, anti-predator responses, habitat preference, prey searching activity, and consumption (Boehlert and Morgan, 1985; Miner and Stein, 1993; Miner and Stein, 1996;

Gregory and Levings, 1998; De Robertis et al., 2003, Meager et al., 2006; Meager and Utne-Palm, 2007). Although differences in the suspended particles used to manipulate turbidity in the simulated conditions in the laboratory prevented a direct quantitative comparison to turbidity levels in the field, the relative water turbidities had interesting effects on Pinfish behaviors. In the lab, Pinfish behavior did not seem affected by the small dimensions of the tank. They were observed swimming and foraging when no turbidity was present in the water. Pinfish reduced their activity levels (swimming), stopped foraging, and became more closely associated with the ASUs when the water was more turbid. It is possible that Pinfish employed these anti-predatory behaviors in both laboratory and field settings to compensate for the reduction of vision used to detect the threat of predators in more turbid waters. In addition, increased turbidity has been shown to interfere with the foraging of predators by making it more difficult for them to locate prey (Meager and Batty, 2007). By reducing their activity levels and remaining closer to the seagrass refugia, Pinfish might have been less visible to predators, hence decreasing the risk of capture and consumption and lowering the predation rates observed in the mid Bay compared to the lower Bay. Pinfish have been shown to change their selection of habitat and behavior based on the presence of predators (Jordan et al., 1996), which can result in the use of suboptimal foraging habitats and slower growth (Harter and Heck, 2006). This can serve as further support that alteration of Pinfish behavior is possible and could have influenced the observed predation differences between locations. Compared to other studies where turbidity acts as an ecological constraint on predator recognition and hampers anti-predatory responses (Ferrari et al., 2010; Becker and Gabor, 2012), Pinfish in the laboratory experiments in this study were able to detect predators even at higher turbidity levels,

suggesting they may have a visual advantage over some predators as turbidity increases.

This speculation may be further supported by the observation that Pinfish can exist at incredibly high densities, including a wide range of individual sizes, across variable levels of turbidity (Chacin thesis, 2014; Stallings et al., 2015) and are visual feeders (Luczkovich, 1988).

Furthermore, marginal support for the presence of an interaction between seagrass complexity and location was found. This could indicate that differences in mortality between sites were stronger when habitat structure was absent compared to when it was more complex. Assuming that the mortality patterns were due to the difference in turbidity between locations, then habitat structure may have been less important for fish predation risk in high turbidity versus low turbidity conditions. To fully clarify this relationship, however, further experiments involving several levels of turbidity and habitat structure must be conducted.

The locations where experiments were conducted also differed in distance to the mouth of the Bay, which is likely responsible for the variation observed in salinity which in turn might have influenced the structure of the predator assemblages. The predator assemblage composition could have been another factor that influenced mortality rates between locations. While the abundance and species richness of predators was similar between the lower and mid Bay, assemblage structure differed between locations. Thus, species identity may have been an important factor contributing to differences in mortality (e.g., Stallings, 2009) potentially due to differences in foraging mode (Huey and Pianka, 1981). Gulf flounder (*Paralichthys albigutta*) and Inshore Lizardfish (*Synodus foetens*) drove the patterns in the lower Bay. These fishes are voracious, lie-and-wait predators and it is

possible that less turbid conditions benefited their type of foraging and behavior. For example, in water where visibility is higher, predators that are mainly stationary on the bottom (and thus non-conspicuous) may have a foraging advantage over conspicuous species. Additionally, fishes in lower turbidity waters have longer reactive distances and thus larger search volumes and higher prey encounter rates (Brett and Groot, 1963; Vinyard and O'Brien, 1976; Gerking, 1994; Benfield and Minello, 1996). Hence, it is likely that predators at the less turbid location, the lower Bay, were able to locate Pinfish more efficiently compared to the mid Bay where the water was more turbid. In the mid Bay, Red Drum (*Sciaenops ocellatus*) and Common Snook (*Centropomus undecimalis*) were key piscivores driving the assemblage patterns. These species are more active and ambush their prey; this feeding strategy could have been beneficial in water where visibility was lower. By being more active, predators can increase their encounter rates with prey. It is probable that the increased scattering of light due to higher turbidity in the mid Bay may have reduced their reactive distance and encounter rates yielding the lower predation rates of Pinfish were observed.

Other studies have reported similar results in that ecological responses (e.g., survival, recruitment) from identically designed, replicated, and controlled field experiments have yielded different outcomes when conducted at multiple sites along environmental gradients (Menge et al., 1976; Menge and Sutherland, 1987). The existence of such experiments (also termed “comparative experimental approach”) allows scientists to understand the importance and generality of factors varying along environmental gradients in the seascape (Dayton, 1975; Menge, 1976; Lubchenco and Menge, 1978; Menge et al., 1994). This study suggests that moderately turbid seagrass environments can have

important refuge value for an ecologically important species. Although eutrophication can be a major cause of seagrass loss (Burkholder et al., 2007), moderate nutrient input (usually correlated with higher turbidity) can stimulate epiphytic algal growth, which can support increased secondary productivity. Embracing the nursery function (e.g., Beck et al., 2001; Dahlgren et al., 2006), it is possible that more turbid seagrass habitats (to some extent) could contribute a considerable number of recruits to the adult population, via reduced mortality rates (as observed in this study). It is not being implied that low turbidity seagrass habitats are less essential, but it is an attempt to emphasize the importance of accounting for more turbid habitats that may be considered less “aesthetically” appealing by human observers.

In this study it was determined that both fine-scale effects of seagrass blade density and broad-scale effects of locations varying in turbidity and predator assemblage can influence relative predation rates on Pinfish. This study may serve as a baseline for continuing investigations on how these and other effects of habitat may mediate predator-prey interactions across spatial scales. Extensive declines in the presence and abundance of seagrass habitats have occurred worldwide in recent years, often with associated changes in blade densities, turbidity levels, and fish assemblages (Blaber and Blaber, 1980; Cyrus and Blaber, 1987; Breitburg, 1988; Shepherd et al., 1989; Hauxwell et al., 2003; Orth et al., 2006; Horinouchi, 2007b; Horinouchi et al., 2009; Waycott et al., 2009). In response to the continuing and accelerating changes to coastal marine ecosystems, often as a consequence of anthropogenic activities (Jackson et al., 2001), different species will experience positive, negative, and null effects. Thus, there is a need to better understand generalities to allow for

528 a more predictive framework that can be used to inform conservation and management
529 efforts.

530

531

532

533

534

535 **Acknowledgements**

536 The following people are thanked for their assistance with fish collections and experimental
537 deployments: Matt Farnum, Tiffany Boisvert, Gabriella Chisari, Peter Simard, Natasha
538 Mendez-Ferrer, Alisha Gray, Ileana Freytes-Ortiz, Matt Garret, Lindsey Berger, Orian
539 Tzadik, Kara Wall, Brock Houston, and the crew from FWRI's Fishery Independent
540 Monitoring Program. Ted Switzer is thanked for his assistance with the predator database,
541 David Jones for help with the statistical analysis, and both Susan Bell and Cameron
542 Ainsworth for advice as graduate committee members for D.H.C. Edits and comments on
543 the manuscript by Pamela Hallock-Muller and the USF Fish Ecology Lab (Orian Tzadik,
544 Ileana Freytes-Ortiz, Joseph Curtis, Kara Wall, Timothy Pusack, Benjamin Kurth, Oscar
545 Ayala) are appreciated. Funding to D.H.C. was provided by the NSF Graduate Research
546 Fellowship, the Peter Betzer Fellowship from the USF College of Marine Science, and the
547 McKnight Doctoral Fellowship from USF.

548

549 **References**

550 Abrahams, M.V., Kattenfeld, M.G., 1997. The role of turbidity as a constraint on predator-
551 prey interactions in aquatic environments. *Behav. Ecol Sociobiol.* 40, 169-174.

- Baltestri, E., Cinelli, F., Lardicci, C., 2003. Spatial variation in *Posidonia oceanica* structural, morphological and dynamic features in a northwestern Mediterranean coastal area: a multi-scale analysis. *Mar. Ecol. Prog. Ser.* 250, 51-60.
- Beck, M.W., Heck, K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M., Halpern, B., Hays, C.G., Hoshino, K., Minello, T.J., Orth, R.J., Sheridan, P.F., Weinstein, M.P., 2001. The identification, conservation, management of estuarine and marine nurseries for fish and invertebrates. *BioScience* 51, 633-641.
- Becker, L.J., Gabor, C.R., 2012. Effects of turbidity and visual vs. chemical cues on anti-predator response in the endangered Fountain Darter. *Ethology* 118, 994-1000.
- Bell, S.S., Hall, M.O., Robbins, B.D., 1995. Toward a landscape approach in seagrass beds: Using macroalgal accumulation to address questions of scale. *Oecologia* 104, 163-168.
- Bell, S.S., Brooks, R.A., Robbins, B.D., Fonseca, M.S., Hall, M.O., 2001. Faunal response to fragmentation in seagrass habitats: Implications for seagrass conservation. *Biol. Conserv.* 100, 115-123.
- Bell, S.S., Fonseca, M.S., Stafford, N.B., 2006. Seagrass ecology: new contributions from a landscape perspective. In: Larkum, A.W.D., Orth, R.J., Duarte, C.M. (Eds) *Seagrasses: Biology, ecology and conservation*. Springer, New York, pp 625-645.
- Bell, S.S., Hicks, G.R.F., 1991. Marine landscapes and faunal recruitment: A field-test with seagrasses and copepods. *Mar. Ecol. Prog. Ser.* 73, 61-68.
- Benfield, M.C., Minello, T.J., 1996. Relative effects of turbidity and light intensity on reactive distance and feeding of an estuarine fish. *Env. Biol. Fish.* 46, 211-216.
- Beukers, J.S., Jones, G.P., 1998. Habitat complexity modifies the impact of piscivores on a coral reef fish population. *Oecologia* 114, 50-59.
- Boehlert, G.W., Morgan, J.B., 1985. Turbidity enhances feeding abilities of larval Pacific herring, *Clupea harengus pallasii*. *Hydrobiologia* 123, 161-170.
- Bowden, D.A., Rowden, A.A., Attrill, M.J., 2001. Effect of patch size and in-patch location on the infaunal macroinvertebrate assemblages of *Zostera marina* seagrass beds. *J. Exp. Mar. Biol. Ecol.* 259, 133-154.
- Blaber, S.J.M., Blaber, T.G., 1980. Factors affecting the distribution of juvenile estuarine and inshore fish. *J. Fish Biol.* 17, 143-162.
- Breitburg, D.L., 1988. Effects of turbidity on prey consumption by striped bass larvae. *Trans. Amer. Fish Soc.* 117, 72-77.
- Bret, J.R., Groot, C., 1963. Some aspects of olfactory and visual responses in Pacific salmon. *J Fish Res. Board Can.* 20, 287-301.
- Burkholder, J.M., Tomasko, D.A., Touchette, B.W., 2007. Seagrasses and eutrophication. *J. Exp. Mar. Bio. Ecol.* 350, 46-72.
- Carroll, J.M., Peterson, B.J., 2013. Ecological trade-offs in seascape ecology: Bay Scallop survival and growth across a seagrass seascape. *Landsc. Ecol.* 28, 1401-1413.
- Chacin, D.H., 2014. A multi-scale approach to study predator-prey interactions and habitat use of Pinfish, *Lagodon rhomboides*. Master Thesis. University of South Florida, College of Marine Science, Florida, USA.
- Chen, Z.Q., Hu, C.M., Muller-Karger, F., 2007. Monitoring turbidity in Tampa Bay using MODIS/Aqua 250-m imagery. *Remote Sens. Environ.* 109, 207-220.
- Crowder, L.B., Cooper, W.E., 1982. Habitat structural complexity and the interaction between Bluegills and their prey. *Ecology* 63, 1802-1813.

- Cyrus, D.P., Blaber, S.J.M., 1987. The influence of turbidity on juveniles marine fishes in estuaries. Part 2. Laboratory studies, comparisons and conclusions. *J. Exp. Mar. Biol. Ecol.* 109, 71-91.
- Dahlgren, C.P., Kellison, G.T., Adams, A.J., Gillanders, B.M., Kendall, M.S., Layman, C.A., Ley, J.A., Nagelkerken, I., Serafy, J.E., 2006. Marine nurseries and effective juvenile habitats: concepts and applications. *Mar. Ecol. Prog. Ser.* 312, 291-295.
- Darcy, G.H., 1985. Synopsis of biological data on the Pinfish, *Lagodon rhomboides* (Pisces:Sparidae). U.S. Dep Commer, NOAA Tech Rep NMFS 23.
- Dayton, P.K., 1975. Experimental evaluation of ecological dominance in a rocky intertidal algal community. *Ecolog. Monog.* 45, 137-159.
- De Robertis, A., Ryer, H.C., Veloza, A., Brodeur, R.D., 2003. Differential effects of turbidity on prey consumption of piscivorous and planktivorous fish. *Can. J. Fish Aquat. Sci.* 60, 1517-1526.
- Ferrari, M.C.O., Wisenden, B.D., Chivers, D.P., 2010. Chemical ecology of predator-prey interactions in aquatic ecosystems: A review and prospectus. *Can. J. Zool.* 88, 698-724.
- Gillanders, B.M., 2006. Seagrasses, fish, and fisheries. In: Larkum, A.W.D., Orth, R.J., Duarte, C.M. (Eds) *Seagrasses: biology, ecology and conservation*. Springer, New York, pp 503-536.
- Gerkin, S.D., 1994. Particulate Feeding. In: *Feeding Ecology of Fish*. Academic Press, Inc, London pp 011-138.
- Gotceitas, V., 1990. Foraging and predator avoidance: A test of a patch choice model with juvenile Bluegill Sunfish. *Oecologia* 83, 346-351.
- Gotceitas, V., Fraser, S., Brown, J., 1997. Use of eelgrass (*Zostera marina*) beds by juvenile Atlantic Cod (*Gadus morhua*). *Can. J. Fish Aquat. Sci.* 54, 1306-1319.
- Greening, H.S., Cross, L.M., Sherwood, T., 2011. A multiscale approach to seagrass recovery in Tampa Bay, Florida. *Ecolog. Rest.* 29, 82-93.
- Gregory, R.S., 1993. Effect of turbidity on the predator avoidance behavior of juvenile chinook salmon (*Oncorhynchus tshawytscha*). *Can. J. Fish Aquat. Sci.* 50, 241-246.
- Gregory, R.S., Northcote, T.G., 1993. Surface, planktonic, and benthic foraging by juvenile chinook salmon (*Oncorhynchus tshawytscha*) in turbid laboratory conditions. *Can. J. Fish Aquat. Sci.* 50, 233-240.
- Gregory, R.S., Levings, C.D., 1998. Turbidity reduces predation on migrating juvenile Pacific salmon. *Trans. Amer. Fish. Soc.* 127, 275-285.
- Hall-Scharf, B.J., Stallings, C.D., 2014. Experimental test of preference by a generalist piscivore on morphologically- and behaviorally- different prey. *J. Exp. Mar. Bio. Ecol.* 460, 193-196.
- Hall-Scharf, B.J., Switzer, T.S., Stallings, C.D. in press. Ontogenetic and long-term diet shifts of a predatory fish in an urban estuary undergoing dramatic changes in water quality and habitat dynamics. *Trans. Amer. Fish. Soc.*
- Hammerschlag, N., Morgan, A.B., Serafy, J.E., 2010. Relative predation risk for fishes along a subtropical mangrove-seagrass ecotone. *Mar. Ecol. Prog. Ser.* 401, 259-267.
- Harris, L.A., Buckley, B., Nixon, S.W., Allen, B.T., 2004. Experimental studies of predation by Bluefish *Pomatomus saltatrix* in varying densities of seagrass and macroalgae. *Mar. Ecol. Prog. Ser.* 281, 233-239.

- Harter, S.L., Heck, K.L., 2006. Growth rates of juvenile Pinfish (*Lagodon rhomboides*): effects of habitat and predation risk. *Estuar. Coast.* 29, 318-327.
- Hauxwell, J., Cebrian, J., Valiela, I., 2003. Eelgrass *Zostera marina* loss in temperate estuaries: relationship to land-derived nitrogen loads and effect of light limitation imposed by algae. *Mar. Ecol. Prog. Ser.* 247, 59-73.
- Healey, D., Hovel, K.A., 2004. Seagrass bed patchiness: Effects on epifaunal communities in San Diego Bay, USA. *J. Exp. Mar. Biol. Ecol.* 313, 155-174.
- Heck, K.L., Crowder, L.B., 1991. Habitat structure and predator-prey interactions in vegetated aquatic systems. In: Bell, S.S., McCoy, E.D., Mushinsky, H.R. (Eds) *Habitat complexity: The physical arrangement of objects in space*. Chapman and Hall, New York, pp 280-299.
- Heck, K.L., Hays, G., Orth, R.J., 2003. Critical evaluation of the nursery role hypothesis for seagrass meadows. *Mar. Ecol. Prog. Ser.* 253, 123-136.
- Heck, K.L., Orth, R.J., 2006. Predation in seagrass beds. In: Larkum AWD, Orth RJ, Duarte CM (Eds) *Seagrasses: biology, ecology and conservation*. Springer, New York, pp 537-550.
- Heck, K.L., Thoman, T.A., 1981. Experiments on predator-prey interactions in vegetated aquatic habitats. *J. Exp. Mar. Biol. Ecol.* 53, 125-134.
- Hemminga, M.A., Duarte, CM., 2000. *Seagrass ecology*. Cambridge University Press. The Edinburgh Building, Cambridge, UK.
- Hoese, H.D., Moore, R.H., 1977. *Fishes of the Gulf of Mexico: Texas, Louisiana, and adjacent waters*. Texas AandM University Press, College Station, pp 327.
- Horinouchi, M., 2007a. Review of the effects of within-patch scale structural complexity on seagrass fishes. *J. Exp. Mar. Biol. Ecol.* 350, 111-129.
- Horinouchi, M., 2007b. Distribution patterns of benthic juvenile gobies in and around seagrass habitats: Effectiveness of seagrass shelter against predators. *Estuar. Coast. Shelf S.* 72, 657-664.
- Horinouchi, M., Mizuno, N., Jo, Y., Fujita, M., Sano, M., Suzuki, Y., 2009. Seagrass habitat complexity does not always decrease foraging efficiencies of piscivorous fishes. *Mar. Ecol. Prog. Ser.* 377, 43-49.
- Hovel, K.A., Fonseca, M.S., 2005. Influence of seagrass landscape structure on the juvenile Blue Crab habitat-survival function. *Mar. Ecol. Prog. Ser.* 300, 179-191.
- Hovel, K.A., Lipcius, R.N., 2002. Effects of seagrass habitat fragmentation on juvenile Blue Crab survival and abundance. *J. Exp. Mar. Biol. Ecol.* 271, 75-98.
- Huey, R.B., Pianka, E.R., 1981. Ecological consequences of foraging mode. *Ecology* 62, 991-999.
- Irlandi, E.A., 1994. Large-scale and small-scale effects of habitat structure on rates of predation: How percent coverage of seagrass affects rates of predation and siphon nipping on an infaunal bivalve. *Oecologia* 98, 176-183.
- Irlandi, E.A., 1997. Seagrass patch size and survivorship of an infaunal bivalve. *Oikos* 78, 511-518.
- Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R.R., 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293, 629-637.

- Janicki, A.J., Wade, D.L., 1996. Estimating critical nitrogen loads for the Tampa Bay estuary: An empirically based approach to setting management targets. Technical Publication #06-96 of the Tampa Bay National Estuary Program. St. Petersburg, Florida.
- Johansson, J.O.R., 1991. Long-term trends of nitrogen loading, water quality and biological indicators in Hillsborough Bay, Florida. In: Treat SF, Clark PA (Eds) Proc, Tampa Bay Area Sci Inf Symp 2, Tampa, Florida, pp157-176.
- Jones, D.L., 2014. Fathom Toolbox for Matlab: software for multivariate ecological and oceanographic data analysis. College of Marine Science, University of South Florida, St. Petersburg, FL, USA. Available from: www.marine.usf.edu/user/djones/
- Jordan, F., Bartolini, M., Christine, N., Patterson, P.E., Soulen, H.L., 1996. Risk of predation affects habitat selection by the Pinfish *Lagodon rhomboides* (Linnaeus). J Exp. Mar. Bio. Ecol. 208, 45-56.
- Kjelson, M.A., Johnson, G.N., 1976. Further observations of the feeding ecology of post-larval Pinfish, *Lagodon rhomboides*, and Spot, *Leiostomus xanthurus*. Fish Bull. 74, 423-432.
- Larkum, A.W.D., Drew, E.A., Ralph, P.J., 2006. Photosynthesis and metabolism in seagrass at the cellular level. In: Larkum AWD, Orth RJ, Duarte CM (Eds) Seagrasses: biology, ecology and conservation. Springer, New York, pp 323-345.
- Laurel, B.J., Brown, J.A., 2006. Influence of cruising and ambush predators on 3-dimensional habitat use in age 0 juvenile Atlantic Cod *Gadus morhua*. J. Exp. Mar. Biol. Ecol. 329, 34-46.
- Lubchenco, J., Menge, B.A., 1978. Community development and persistence in a low rocky intertidal zone. Ecolog. Monog.
- Luczkovich, J.J., 1988. The role of prey detection in the selection of prey by pinfish *Lagodon rhomboides* (Linnaeus). J. Exp. Mar. Biol. Ecol. 123, 15-30.
- Main, K.L., 1987. Predator avoidance in seagrass meadows: prey behavior, microhabitat selection, and cryptic coloration. Ecology 68, 170-180.
- Meyer, A.C., Bell, S.S., 2010. Depositional dynamics in seagrass systems of Tampa Bay, FL: Influence of hydrodynamic regime and vegetation density on ecosystem function. PhD dissertation, University of South Florida, Tampa, FL
- Meager, J.J., Domenici, P., Shingles, A., Utne-Palm, A.C., 2006. Escape responses in juvenile Atlantic cod *Gadus morhua* L.: the effects of turbidity and predator speed. J. Exp. Biol. 209, 4174-4184.
- Meager, J.J., Batty, R., 2007. Effects of turbidity on the spontaneous and prey-searching activity of juvenile Atlantic Cod (*Gadus morhua*). Phil. Trans. R. Soc. B. 362, 2123-2130.
- Meager, J.J., Utne-Palm, A.C., 2007. Effect of turbidity on habitat preference of juvenile Atlantic Cod, (*Gadus morhua*). Envir. Biol. Fish. DOI 10.1007/s10641-007-9183-z
- Menge, B.A., 1976. Organization of the New England rocky intertidal community: role of predation, competition and environmental heterogeneity. Ecolog. Monog. 46, 355-393.
- Menge, B.A., Sutherland, J.P., 1987. Community regulation: variation in disturbance, competition, and predation in relation to environmental stress and recruitment. Am. Nat. 130, 730-757.
- Menge, B.A., Berlow, E.L., Blanchette., 1994. The keystone species concept: variation in

- interactions strength in a rocky intertidal habitat. Ecol. Monog. 64, 249-286.
- Miner, J.G., Stein, R.A., 1993. Interactive influence of turbidity and light on larval bluegill (*Lepomis macrochis*) foraging. Can. J. Fish. Aquat. Sci. 50, 781-788.
- Miner, J.G., Stein, R.A., 1996. Detection of predators and habitat choice by small bluegills: Effects of turbidity and alternative prey. Trans. Amer. Fish. Soc. 125, 97-103.
- Murphey, P.L., Fonseca, M.S., 1995. Role of high and low-energy seagrass beds as nursery areas for *Penaeus duorarum* in North Carolina. Mar. Ecol. Prog. Ser. 121, 91-98.
- Nelson, G.A., 1978. Organization of a subtidal seagrass amphipod guild: the roles of predation, competition, and physical stress. PhD dissertation, Duke University, Durham, NC.
- Nelson, W.G., Bonsdorff, 1990. Fish predation and habitat complexity: are complexity thresholds real? J. Exp. Mar. Bio. Ecol. 141, 183-194
- Nelson, G.A., 2002. Age, growth, mortality, and distribution of Pinfish (*Lagodon rhomboides*) in Tampa Bay and adjacent Gulf of Mexico waters. Fish Bull. 100, 582-592.
- Nelson, J.A., Stallings, C.D., Landing, W.M., Chanton, J., 2013. Biomass transfer subsidizes nitrogen to offshore food webs. Ecosystems 16, 1130-1138.
- Orth, R.J., Carruthers, T.J.B., Dennison, W.C., Duarte, C.M., Fourqurean, J.W., Heck, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Olyarnik, S., Short, F.T., Waycott, M., Williams, S.L., 2006. A global crisis for seagrass ecosystems. BioScience 56, 987-996.
- Orth, R.J., Heck, K.L., Vanmontfrans, J., 1984. Faunal communities in seagrass beds: A review of the influence of plant structure and prey characteristics on predator prey relationships. Estuaries 7, 339-350.
- Orth, R.J., 1992. A perspective on plant-animal interactions in seagrasses: Physical and biological determinants influencing plant and animal abundance. In: John, D.M., Hawkins, S.J. and Price, J.H. (Eds) Plant-animal interactions in the marine benthos. Systematics Association Special 46, 147-164.
- Peterson, B.J., Thompson, K.R., Cowan, J.H., Heck, K.L., 2001. Comparison of predation pressure in temperate and subtropical seagrass habitats based on chronographic tethering. Mar. Ecol. Prog. Ser. 224, 77-85.
- Pirtle, J.L., Eckert, E.L., Stoner, A.W., 2012. Habitat structure influences the survival and predator-prey interactions of early juvenile Red King Crab *Paralithodes camtschaticus*. Mar. Ecol. Prog. Ser. 465, 169-184.
- Robbins, B.D., Bell, S.S., 1994. Seagrass landscapes: A terrestrial approach to the marine subtidal environment. Trends Ecol. Evol. 9, 301-304.
- Rooker, J.R., Holt, G.J., Holt, S.A., 1998. Vulnerability of newly settled red drum (*Sciaenops ocellatus*) to predatory fish: is early-life survival enhanced by seagrass meadows. Mar. Biol. 131:145-151.
- Savino, J.F., Stein, R.A., 1982. Predator-prey interaction between Largemouth Bass and Bluegills as influenced by simulated, submersed vegetation. Trans. Am. Fish. Soc. 111, 255-266.
- Seaman, W.Jr., Collins, M., 1983. Species profiles: Life histories and environmental requirements of coastal fishes and invertebrates (South Florida)-- Snook. U.S. Fish Wildl. Serv. FWS/OBS-82/11.16. U.S. Army Corps of Engineers, pp 16.

- Schultz, S.T., Kruschel, C., 2010. Frequency and succes of ambush and chase predation in fish assemblages associated with seagrass and bare sediment in an Adriatic lagoon. *Hydrobiologia* 649, 25-37.
- Shepherd, S.A., McComb, A.J., Bulthuis, D.A., Neverauskas, V., Steffensen, D.A., West, R., 1989. Decline of seagrasses. In: Larkum, AWD, McComb AJ, Shepard SA (eds) *Biology of seagrasses: A treatise on seagrass with special reference to the Australian region*. Elsevier, North Holland, pp 346-387.
- Smith, T.M., Hindell, J.S., Jenkins, G.P., Connolly, R.M., 2008. Edge effects on fish associated with seagrass and sand patches. *Mar. Ecol. Prog. Ser.* 359, 203-213.
- Stallings, C.D., 2009. Predator identity and recruitment of coral-reef fishes: indirect effects of fishing. *Mar. Ecol. Prog. Ser.* 383, 251-259.
- Stallings, C.D., 2010. Experimental test of preference by a predatory fish for prey at different densities. *J. Exp. Mar. Biol. Ecol.* 389, 1-5.
- Stallings, C.D., Coleman F.C., Koenig, C.C., Markiewicz, D.A., 2010. Energy allocation in juveniles of a warm-temperate reef fish. *Environ. Biol. Fish.* 88, 389-398.
- Stallings, C.D., Mickle, A., Nelson, J.A., McManus, M.G., Koenig, C.C., 2015. Faunal communities and habitat characteristics of the Big Bend seagrass meadows, 2009-2010. *Ecology* 96, 304 (Ecological Archives E096-030).
- Stoner, A.W., 1980. Feeding ecology of *Lagodon rhomboides* (Pisces:Sparidae) variation and functional responses. *Fish Bull.* 78, 337-352.
- Stoner, A.W., 1982. The influence of benthic macrophytes on the foraging behavior of Pinfish, *Lagodon rhomboides* (Linnaeus). *J. Exp. Mar. Biol. Ecol.* 58, 271-284.
- Stoner, A.W., Lewis, F.G., 1985. The influence of quantitative and qualitative aspects of habitat complexity in tropical seagrass meadows. *J. Exp. Mar. Biol. Ecol.* 94, 19-40.
- Tabb, D.C., Manning, R.B., 1961. A checklist of the flora and fauna of northern Florida Bay and adjacent brackish waters of the Florida mainland collected during Jul 1957-Sep 1960. *Bull. Mar. Sci. Gulf. Carib.* 4, 552-649.
- Vogel, J.L., Beauchamp, D.A., 1999. Effects of light, prey size, and turbidity on reaction distances of lake trout (*Salvelinus namaycush*) to salmonid prey. *Can. J. Fish Aquat. Sci.* 56, 1293-1297.
- Vinyard, G.L., O'Brien, W.J., 1976. Effects of light and turbidity on the reactive distance of bluegill (*Lepomis macrochirus*). *J. Fish Res. Board Can.* 33, 2845-2849.
- Wahle, R.A., Steneck, R.S., 1991. Recruitment habitats and nursery grounds of the American Lobster *Homarus americanus*: a demographic bottleneck. *Mar. Ecol. Prog. Ser.* 69, 231-243.
- Waycott, M., Duarte, C.M., Carruthers, T.J.B., Orth, R.J., Dennison, W.C., Olyarnik, S., Calladine, A., Fourqurean, J.W., Heck, K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Short, F.T., Williams, S.L., 2009. Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc. Nat. Acad. Sci. USA* 106, 12377-12381.
- Weinstein, M.P., Heck, K.L., Giebel, P.E., Gates, J.E., 1982. The role of herbivory in Pinfish (*Lagodon rhomboides*): A preliminary investigation. *Bull. Mar. Sci.* 32, 791-795.
- Young, D.K., Buzas, M.A., Young, M.W., 1976. Species densities of macrobenthos associated with seagrass field experimental-study of predation. *J. Mar. Res.* 34, 577-592.

Young, D.K., Young, M.W., 1977. Community structure of the macrobenthos associated with seagrass of the Indian River estuary, Florida. In: Coull BC (Ed) Ecology of marine benthos. Belle W. Baruch Library in Marine Science 6, Unive. SC Press, Columbia, SC, pp 359-381.

Table 1. Number of Pinfish preyed upon over tethering deployments conducted at each study location and by seagrass density treatment.

Study location	Seagrass density		
	Low	Medium	High
Mid Bay	3/22	2/20	0/22
Lower Bay	20/23	14/25	5/24

Table 2. Non-parametric one-way and two-way analysis of variance tests for main effects and interactions between seagrass density with location effects on relative mortality rates of Pinfish. Values in bold indicate significant differences.

Source	df	SS	MS	<i>F</i>	p
Location	1	1.448	1.448	18.746	0.0002
Residual	27	2.085	0.077		
Total	28	3.533			
Seagrass density	2	0.895	0.447	11.284	0.0006
Location	1	1.448	1.448	36.513	0.0002
Seagrass density X Location	2	0.278	0.139	3.511	0.0546
Residual	23	0.912	0.040		
Total	28	3.533			

Figure Captions

847

848 Fig. 1. Map of Tampa Bay, Florida, USA, showing the general turbidity gradient and the
849 study locations (stars).

850

851 Fig. 2. Diagram of the tethering set up on an Artificial Seagrass Unit. The diagram shows a
852 tethered Pinfish on a high density ASU. Pinfish illustration was kindly provided with
853 permission by Diane R. Peebles.

854

855 Fig. 3. Laboratory experimental set up from both a side view (A) and a top view (B). Note
856 the grid shown in (B), used to quantify behavior of Pinfish. The dark line between the
857 experimental tank containing the Pinfish and the predator/control tank represents the visual
858 barrier, which was removed after the 10-minute pre-stimulus period. Pinfish illustration was
859 kindly provided with permission by Diane R. Peebles.

860

861 Fig. 4. Mean $\pm$ standard error of relative predation rates of Pinfish across the three blade
862 densities of seagrass and two study locations.

863

864 Fig. 5. Mean $\pm$ standard error density (A) and richness (B) of Pinfish predators per sample
865 trawl tow (per hectare) in the two sections of the Bay where tethering experiments were
866 conducted.

867

Fig. 6. CAP plot of Pinfish predator assemblage structure (A). Circles represent data points in the mid Bay and upside down triangles the lower Bay. The Y-axis data are jittered to ease visual assessment of the patterns. Species vectors pointing to the right were the drivers in the lower Bay while those pointing to the left were drivers in the mid Bay (B).

Fig. 7. Activity of Pinfish in two turbidity levels (A) and in response to a cross-factored turbidity * predator cue experiment (B).

Figure 1

Figure 3

B

Experimental tank

Predator/control tank

Figure 4

Figure 5

Figure 6

Figure 7

Table 1

Table 1. Number of Pinfish preyed upon over tethering deployments conducted at each study location and by seagrass density treatment.

Study location	Seagrass density		
	Low	Medium	High
Mid Bay	3/22	2/20	0/22
Lower Bay	20/23	14/25	5/24

Table 2

Table 2. Non-parametric one-way and two-way analysis of variance tests for main effects and interactions between seagrass density with location effects on relative mortality rates of Pinfish. Values in bold indicate significant differences.

Source	df	SS	MS	<i>F</i>	p
Location	1	1.448	1.448	18.746	0.0002
Residual	27	2.085	0.077		
Total	28	3.533			
Seagrass density	2	0.895	0.447	11.284	0.0006
Location	1	1.448	1.448	36.513	0.0002
Seagrass density X Location	2	0.278	0.139	3.511	0.0546
Residual	23	0.912	0.040		
Total	28	3.533			