The ectoparasitic mite *Tropilaelaps mercedesae* reduces western honey bee, *Apis mellifera*, longevity and emergence weight, and promotes Deformed wing virus infections

Kitiphong Khongphinitbunjong^{1,2,*}, Peter Neumann^{2,3,4}, Panuwan Chantawannakul^{2,*},
and Geoffrey R. Williams^{2,3,4}

¹School of Science, Mae Fah Luang University, Chiang Rai 57100, Thailand

²Bee Protection Laboratory, Department of Biology, Faculty of Science, Chiang Mai University, Chiang Mai 50200, Thailand

³Institute of Bee Health, Vetsuisse Faculty, University of Bern, 3003 Bern, Switzerland

⁴Agroscope, Swiss Bee Research Centre, 3003 Bern, Switzerland

* Corresponding authors

E-mail address: khongphinit@gmail.com (KK)

E-mail address: panuwan@gmail.com (PC)

Abstract

Historically an ectoparasite of the native Giant honey bee *Apis dorsata*, the mite *Tropilaelaps mercedesae* has switched hosts to the introduced western honey bee *Apis mellifera* throughout much of Asia. Few data regarding lethal and sub-lethal effects of *T. mercedesae* on *A. mellifera* exist, despite its similarity to the devastating mite *Varroa destructor*. Here we artificially infested worker brood of *A. mellifera* with *T. mercedesae* to investigate lethal (longevity) and sub-lethal (emergence weight, Deformed wing virus (DWV) levels and clinical symptoms of DWV) effects of the mite on its new host. The data show that *T. mercedesae* infestation significantly reduced host longevity and emergence weight, and promoted both DWV levels and associated clinical symptoms. Our results suggest that *T. mercedesae* is a potentially important parasite to the economically important *A. mellifera* honey bee.

Keywords: *Apis mellifera*; Deformed wing virus; *Tropilaelaps mercedesae*; vector; longevity

1. Introduction

Global trade of managed western honey bees, *Apis mellifera*, from its native Eurasian and African ranges has provided bountiful opportunities for novel host-parasite relationships (Mutinelli, 2011; Neumann et al., 2016). This is especially true for Southeast Asia, which hosts at least eight indigenous honey bee species (Crane, 1999) in addition to the introduced *A. mellifera* (Moritz et al., 2005). *Tropilaelaps mercedesae*, an indigenous and ubiquitous parasitic mite of the giant honey bee *Apis dorsata* (Laigo and Morse, 1968), is an example of one species that has exploited sympatric *A. mellifera* populations across much of Asia since introduction of the latter (Anderson and Morgan 2007). Similar to the more well studied mites *Varroa* spp. that historically parasitized the eastern honey bee *Apis cerana* and now also exploit *A. mellifera* (*V. destructor*: Anderson and Trueman 2000; *V. jacobsoni*, Roberts et al., 2015), *T. mercedesae* exploits adult hosts, but is primarily a brood parasite (Kapil and Aggarwal, 1987; Sammataro et al. 2000; Oldroyd and Wongsiri 2006; Rosenkranz et al., 2010). Fertile female mites oviposit within cells that contain developing honey bees; progeny subsequently hatch and pierce the cuticle of the host to feed on hemolymph (Burgett et al., 1983; Woyke, 1987c). Comparative studies of *T. mercedesae* and *V. destructor* parasitizing *A. mellifera* in Asia have demonstrated that the former produces a higher number of offspring and requires a shorter phoretic period on adult hosts, thereby promoting relatively higher population levels within colonies that could potentially have devastating consequences (Woyke, 1987a,b,c; Sammataro et al., 2000; Garrido and Rosenkranz, 2003). Indeed, anecdotal reports from beekeepers suggest that *T. mercedesae* can have a severe impact on local *A. mellifera* colonies (Buawangpong et al., 2015).

Lethal and sub-lethal effects of parasitism by *V. destructor* on *A. mellifera* are well-documented (Rosenkranz et al., 2010). Globally, it is considered to be the single most important biotic stressor to colonies of European *A. mellifera* honey bee subspecies (Dietemann et al., 2012; Neumann et al., 2012); feeding on haemolymph results in anaemia, the vectoring of viruses, host immune suppression, reduced host longevity (De Jong et al., 1982; De Jong and De Jong, 1983; Yue and Genersch, 2005; Chen and Siede, 2007; Yang and Cox-Foster, 2007) and ultimately colony mortality (Neumann and Carreck, 2010). Conversely, relatively few information exists about the effects of *T. mercedesae* parasitism on *A. mellifera*. Recent studies suggest that similar to *V. destructor*, *T. mercedesae* can vector honey bee parasites such as Deformed wing virus (DWV) (Dainat et al., 2009; Forsgren et al., 2009) and can influence host immune responses (Khongphinitbunjong et al., 2015b), but no data currently exist regarding common lethal and sub-lethal measures of mite parasitism. In light of anecdotal reports by beekeepers in Asia that *T. mercedesae* might be more devastating to colonies of *A. mellifera* than *V. destructor*, data on the effects of this mite are urgently required considering the potential for this parasite to expand its range due to movement of bees and their products. Here, we investigated possible lethal (longevity) and sub-lethal (emergence weight, DWV infection levels and clinical symptoms) effects of *T. mercedesae* on *A. mellifera*.

2. Materials and methods

2.1 Experimental design

This experiment was conducted at Chiang Mai University, Chiang Mai, Thailand during October and November 2013 using local *A. mellifera* western honey

bees of mixed European origin and *T. mercedesae* mites; the latter was confirmed by molecular analyses according to Khongphinitbunjong et al. (2012).

Three queenright *A. mellifera* colonies were used in this study; all had non-patchy-brood frames, low brood infestation levels of *T. mercedesae* (0.29 ± 0.17 %) and *V. destructor* (0 %), respectively (estimated following Pettis et al., 2013), and no visible clinical symptoms of other diseases (e.g. chalkbrood). Acaricide treatments were not applied to the experimental colonies within three months of the study to facilitate sufficient *T. mercedesae* numbers. Adult female *T. mercedesae* mites were obtained by uncapping newly sealed worker brood of highly infested colonies (Kirrane et al., 2011; Khongphinitbunjong et al., 2013). Age cohorts of bees for the experiment were obtained by confining queens in each colony on an empty brood comb for 24 hours using a standard cage frame (Williams et al., 2013). Within six hours of capping, *T. mercedesae* were introduced into age cohort brood cells using an established artificial infestation method that inserted a single mature female mite via a small incision (~1 mm) created at the base of the cell capping (Garrido and Rosenkranz, 2003; Dietemann et al., 2013; Khongphinitbunjong et al., 2013). Cohort brood cells in each colony were randomly assigned to one of three groups (n = 40 per group per colony) that consisted of: 1.) cells undisturbed (C/C = controls); 2.) cells with an incision but without an introduced mite (I/C = controls for effect of experimental manipulation); and 3.) cells with an incision and with an introduced mite (I/T = mite treatment). The position of each experimental cell was mapped on a transparent sheet to enable future re-examination. After the experimental treatments were performed, test frames were maintained at 34°C and 70% relative humidity following standard protocols (Williams et al., 2012) in a growth chamber. Each pupa was removed from its cell

~10-11 days after inoculation during the grey thorax stage to ensure that all the mother mites and progeny could be collected (Anderson and Roberts 2013; Khongphinitbunjong et al., 2015b). Pupae were transferred to modified 1.5 ml ventilated microcentrifuge tubes and maintained under previously described conditions to allow proper development to maturity. Within 24 hours of removal from the cells, each individual was weighed and examined for deformed wings according to Williams et al. (2009). Each bee was subsequently marked with a worker Opalith platelet (Neumann et al., 2003), and transferred to plastic hoarding cages (Retschnig et al. 2014a,b). For each treatment and colony, two cages were used (n = 20 bees per cage). Each cage was provided sucrose solution *ad libitum* (1 water: 1 sugar), and again maintained at previously described growth chamber conditions. Cages were observed daily until all bees had died; dead bees were individually collected and kept at -20°C before RNA extraction.

2.2 Deformed wing virus analyses

Clinical symptoms of DWV are easily diagnosed in the field, and include deformed wings and shortened abdomens (Bailey and Ball, 1991). A total of 70 bees that died in cages at Day 2 ± 1 not exhibiting (C/C=15, I/C=15, I/T=13) or exhibiting (C/C=2, I/C=5, I/T= 20) clinical symptoms were analyzed for DWV levels according to Khongphinitbunjong et al. (2015a, b). These dead bees provided a ‘snap-shot’ using a suitable sample size of individuals collected within a compressed time period in order to prevent extended parasite replication from influencing results (Huang and Solter, 2013; Williams et al., 2014). Total RNA was extracted from individual bees using TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer’s protocol. The concentration (ng/μl) and purity (A260/280) of total

RNA was determined using a spectrophotometer (BioDrop™). Reverse transcription reactions for cDNA synthesis were performed using an iScript™ cDNA Synthesis Kit (Biorad). The presence of DWV virus was determined by qPCR using IQ5 Real-Time PCR thermal cycler (Bio-Rad). Amplification was performed in 20 µl reaction volumes using SsoFast EvaGreen (BioRad), consisting of 10 µl SSo SYBER mix, 1 µl of 10 µM of each primer [DWV(AY292384.1) and *β-actin* (Simone et al., 2009)], 7 µl of nuclease free water and 1 µl cDNA. Reactions were run at 95°C for 30 s, 40 cycles of 95°C for 1 s, 59°C for 5 s followed by a melt-curve dissociation analysis. All reactions included three replicates. The qPCR data were expressed as the threshold cycle (Ct) value and were determined by normalizing the Ct value of the reference gene (*β-actin*) to the target genes (ΔCt). To compare DWV levels across treatments, the qPCR data were interpreted using the $2^{-\Delta\Delta Ct}$ method (Livak and Schnittgen, 2001, Chaimanee et al., 2012, Boncristiani et al., 2013). The treatment group with the lowest viral level was used as the calibrator and the levels of viruses in all other groups were expressed as n-fold differences relative to the calibrator (Chen et al., 2005).

2.3 Statistical analyses

Statistical analyses were performed using JMP (version 11.0) and SPSS (version 17.0). Figures visualizing data were created using Microsoft Excel (Office version 2010), except for longevity which was created using SPSS (version 17.0). Some data were not normally distributed; emergence weight and longevity data were therefore square-root transformed, whereas proportional deformed wing data were arcsine square-root transformed. Emergence mass, wing deformity, and DWV levels were analyzed using a mixed model that incorporated colony as a random factor.

When differences were found, means were compared using Tukey-HSD (emergence weight, proportions of bees with deformed wings and DWV levels for all individuals) and Student's *t* test (DWV levels considering clinical symptoms) with 95% confidence. Differences in bee longevity among treatments were compared using a Kaplan-Meier survival analysis; Log Rank multiple comparison tests were used to determine difference between specific treatments.

3. Results

Overall, 84% of the introduced female mites reproduced; the average number of progeny was 1.86 ± 0.12 . No *V. destructor* mites were found in any cells of the experimental brood frames. Bees infested with *T. mercedesae* (I/T = 96.2 ± 0.94 mg) had a significantly reduced weight ($F = 22.83$, d.f. = 2, $P < 0.0001$; Tukey's HSD test, both P s < 0.0001) (Figure 1), whereas no significant difference was observed between the two control groups (C/C = 103.9 ± 0.79 mg, I/C = 101.2 ± 0.71 mg) (Tukey's HSD test, $P = 0.059$) (Figure 1).

The proportion of bees with deformed wings significantly differed among treatment groups ($F = 24.61$, d.f. = 2, $P = 0.0056$) (Figure 2); bees from the *T. mercedesae* group showed significantly more wing deformities compared to both controls (Tukey's HSD test; both P s < 0.008), whereas no difference was observed between the two control groups (Tukey's HSD test, $P = 0.997$) (Figure 2).

The mean longevity of bees infested with *T. mercedesae* was significantly shorter compared to the controls (Log rank test, $\chi^2 = 32.915$, $P < 0.0001$) (Figure 3). No difference was observed between the two controls (Log rank test, $P > 0.05$) (Figure 3)

T. mercedesae-infested bees had significantly higher DWV levels ($F = 36.26$, d.f. = 2, $P = 0.0027$; Tukey's HSD test, both P s < 0.0222) (Figure 4a). For those bees not exhibiting clinical symptoms, DWV levels were also significantly higher when bees were infested by *T. mercedesae* compared to both controls ($F = 2.51$, d.f. = 5, $P = 0.0388$; Student's *t*-test, both P s < 0.009) (Figure 4b); however, DWV levels did not differ among treatments when bees exhibited DWV clinical symptoms (Student's *t* test; all P s > 0.556) (Figure 4b). Furthermore, presence or absence of wing deformities did not affect DWV levels in bees exposed to *T. mercedesae* (Student's *t* test; $P = 0.414$) (Figure 4b).

4. Discussion

Our results demonstrate for the first time experimentally that *T. mercedesae* can reduce *A. mellifera* longevity and emergence weight, as well as promote DWV infection levels and associated clinical symptoms. Since *T. mercedesae* occurs nearly ubiquitously throughout Asia, our results support the need for further research on this parasite.

Low levels of *V. destructor* from *T. mercedesae*-infested *A. mellifera* colonies observed in this study confirms earlier reports from China (Dainat et al. 2009) and Thailand (Pettis et al., 2012; Buawangpong et al., 2015). The underlying reasons for these infestation levels might be due to inter-specific competition (e.g. Williams et al., 2014; Buawangpong et al., 2015); however, further investigation is required.

The significant reduction in worker longevity due to *T. mercedesae* parasitism observed in this study is in line with previous studies that demonstrate that honey bee lifespan can be reduced by ectoparasitic mites (De Jong and De Jong, 1983; Yang and Cox-Foster, 2007; Annoscia, et al. 2012). The overall reduced longevity of hoarding

cage workers in this study likely reflects early uncapping of all experimental workers (~1-2 days prior to emergence in order to obtain mite infestation data).

The data also suggest that *T. mercedesae* promotes wing deformation and increased DWV levels in *A. mellifera*, despite comparable DWV levels between deformed control bees (C/C and I/C groups) and deformed *T. mercedesae*-parasitized bees. This is because proportionally fewer deformed control individuals (2 C/C and 5 I/C) were observed compared to deformed mite-infested individuals (20 I/T). This relatively small proportion also supports the importance of the primary mode of transmission for DWV virulence (Möckel et al. 2011). Furthermore, future studies should investigate the role of particular DWV mutants and virus replication using only bee heads to further understand factors responsible for inducing clinical symptoms of DWV infection (Yue and Genersch, 2005; Yue et al., 2007; Gisder et al., 2009). Similar to *V. destructor* (Yue and Genersch, 2005; Gisder et al., 2009; Nazzi et al., 2012), it is likely that higher DWV infection loads and a higher prevalence of wing deformities in *T. mercedesae* infested bees is because mite feeding vectors the virus (Dainat et al. 2009; Forsgren et al. 2009) or suppresses bee host immune responses (e.g. altering anti-viral immune defenses such as the NF- κ B gene family) when the virus is already present (Khongphinitbunjong et al., 2015b). Low DWV levels in the controls provide strong evidence that DWV levels were also low in our experimental colonies and that bees from our other experimental treatments initially had low DWV levels prior to the experiment. Indeed, latent DWV levels are common place in honey bees (Kojima et al., 2011; Ai et al., 2012; Martin et al., 2012), particularly in Southeast Asia (Sanpa and Chantawannakul 2009; Forsgren et al., 2015).

Emergence weight is an important indicator of overall bee health (Pettis et al.; 2004, 2012). Our results showed that the weight of newly emerged *A. mellifera* workers infested with *T. mercedesae* was much lower than those of uninfested ones. Similar results were also reported by De Jong et al. (1982), wherein bee weight at emergence also decreased due to *V. destructor* parasitism. Moreover, it has also been reported that feeding by *V. destructor* and *T. mercedesae* could induce immune responses (Gregory et al., 2005; Yang and Cox-foster, 2005, Nazzi et al. 2012, Khongphinitbunjong et al. 2015b).

In summary, we have demonstrated that parasitism by the ectoparasitic mite *T. mercedesae* can have both important lethal and sublethal effects on *A. mellifera* at the individual worker level. This is worrisome considering the importance of *A. mellifera* to agricultural pollination and the massive economic penalties already caused *V. destructor* (Rosenkranz et al., 2010; Dietemann et al., 2012). It is interesting to note that both genera of mites are not considered to be harmful towards the original corresponding hosts, *Apis cerana* and *A. dorsata* respectively. Future comparative investigations between *T. mercedesae* and *V. destructor* in both their respective native and novel host honey bees have a significant potential to enhance our knowledge of these two emerging ectoparasitic mites. This is true for both the fundamental understanding of host-pathogen interactions as well as for practical beekeeping in Asia and elsewhere. Our data provide support that *T. mercedesae* has the potential to become an important global pest of *A. mellifera* apiculture, thereby creating demand for more studies on its biology and control.

Acknowledgments

We wish to thank the Vinetum Foundation (PN), Agroscope (GW), the Ricola Foundation – *Nature and Culture* (PN), the Swiss National Science Foundation (PN), Chiang Mai University (PC, KK, GW), and the Thailand research fund (PC and KK) for their financial support.

References

- Ai, H., Yan, X., Han, R., 2012. Occurrence and prevalence of seven bee viruses in *Apis mellifera* and *Apis cerana* apiaries in China. J. Invertebr. Pathol 109, 160-164.
- Anderson, D. L., Morgan, M. J., 2007. Genetic and morphological variation of bee parasitic *Tropilaelaps* mites (Acari: Laelapidae): New and re-defined species. Exp. Appl. Acarol. 43, 1-24. doi:10.1007/s10493-007-9103-0.
- Anderson, D. L., Trueman, J. W. H., 2000. *Varroa jacobsoni* (Acari: Varroidae) is more than one species. Exp. Appl. Acarol. 24, 165-189. doi:10.1023/A:1006456720416.
- Anderson, D.L., M K Roberts, J.M.K., 2013. Standard methods for *Tropilaelaps* mites research. J. Apic. Res. 52 doi:10.3896/IBRA.1.52.4.21.
- Annoscia, D., Del Piccolo, F., Nazzi, F., 2012. How does the mite *Varroa destructor* kill the honeybee *Apis mellifera*? Alteration of cuticular hydrocarbons and water loss in infested honeybees. J. Insect Physiol. 58: 1548-1555.
- Bailey L., Ball B.V. Honey Bee Pathology. Academic Press Inc.; San Diego, CA, USA: 1991
- Boncristiani, H.F., Evans, J.D., Chen, Y.P., Pettis, J., Murphy, C., Lopez, D.L., Simone-Finstrom, M., Strand, M., Tarpy, D.R., Rueppell, O., 2013, In vitro infection of pupae with Israeli Acute Paralysis Virus suggests disturbance of transcriptional homeostasis in honey bees (*Apis mellifera*). PLoS ONE, 8, e73429.

- Buawangpong, N., De Guzman, L.I., Khongphinitbunjong, K., Frake, A.M., Burgett, M., Chantawannakul, P., 2015. *Tropilaelaps mercedesae* and *Varroa destructor*: prevalence and reproduction in concurrently infested *Apis mellifera* colonies. *Apidologie* doi:10.1007/s13592-015-0368-8.
- Burgett, M., Akratanakul, P., Morse, R.A., 1983. *Tropilaelaps clareae*: a parasite of honey bees in south-east Asia. *Bee World* 64, 25-28.
- Chaimanee, V., Chantawannakul, P., Chen, Y.P., Evans, J.D., Pettis, J., 2012. Differential expression of immune genes of adult honey bee (*Apis mellifera*) after inoculated by *Nosema ceranae*. *J. Insect Physiol.* 58, 1090-1095.
- Chen, Y. P., Higgins, J. A., Feldlaufer, M.F., 2005. Quantitative real time reverse transcription-PCR analysis of deformed wing virus infection in honeybee (*Apis mellifera* L.). *Appl. Environ. Microbiol.* 71, 436-441.
- Chen, Y.P., Siede R., 2007. Honey Bee Viruses. *Advances in Virus Research.* 70, 33-80.
- Crane, E., 1999. *The World History of Beekeeping and Honey Hunting*. Gerald Duckworth & Co. Ltd, London.
- Dainat, B., Ken, T., Berthoud, H., Neumann, P., 2009. The ectoparasitic mite *Tropilaelaps mercedesae* (Acari, Laelapidae) as a vector of honeybee viruses. *Insectes Sociaux*, 56, 40-43.
- De Jong, D., De Jong, P., Gonçalves, L., 1982. Weight loss and other damage to developing worker honeybees from infestation with *Varroa jacobsoni*. *J. Apic. Res.* 21, 165-167.
- De Jong, D., De Jong, P., 1983. Longevity of Africanized Honey bees (Hymenoptera: Apidae) infested by *Varroa jacobsoni* (Parasitiformes: Varroidae). *J. Econ. Entomol.* 76, 766-768.

- Dietemann, V., Pflugfelder, J. Anderson, D., Charrière, J.D., Chejanovsky, N.,
Dainat, B. de Miranda, J., Delaplane, K., Diller, F., Fuch, S., Gallman, P.,
Gauthier, L., Imdorf, A., Koeniger, N., Kralj, J., Meikle, W., Pettis, J.,
Rosenkranz, P., Sammartaro, D. Smith, D., Yañez, O., Neumann, P., 2012.
Varroa destructor: research avenues towards sustainable control. J. Apic. Res.
51, 125-132.
- Dietemann, V., Nazzi, F.; Martin S.J., Anderson, D.L., Barbara Locke, B., Keith, S.,
Delaplane, K.S., Wauquiez, Q., Tannahill, C., Frey, E.; Ziegelmann, B.,
Rosenkranz, P., D Ellis, J.D., 2013. Standard methods for varroa research.
Journal of Apicultural Research, 52 doi:10.3896/ibra.1.52.1.09.
- Forsgren, E., de Miranda, J.R., Isaksson, M., Wei, S., Fries, I., 2009. Deformed wing
virus associated with *Tropilaelaps mercedesae* infesting European honey bees
(*Apis mellifera*). Exp. Appl. Acarol. 47, 87-97.
- Forsgren, E., Wei, S., Guiling, D., Zhiguang, L., Tran, T.V., Tang, P.T., Truong, T.A.,
Dinh, T.Q., Fries, I., 2015. Preliminary observations on possible pathogen spill-
over from *Apis mellifera* to *Apis cerana*. Apidologie 46, 265-275.
- Garrido, C., Rosenkranz, P., 2003. The reproductive program of female *Varroa*
destructor mites is triggered by its host, *Apis mellifera*. Exp. Appl. Acarol. 31,
269-273.
- Gisder, S., Aumeier, P., Genersch, E., 2009. Deformed wing virus: replication and
viral load in mites (*Varroa destructor*). J. Gen. Virol. 90, 463.
- Gregory, P.G., Evans, J.D., Rinderer, T., de Guzman, L.I., 2005. Conditional
immune-gene suppression of honeybees parasitized by Varroa mites. J.
Insect Sci. 5, 1-5.

- Huang, W.F., Solter, L.F., 2013. Comparative development and tissue tropism in *Nosema apis* and *Nosema ceranae*. J. Invertebr. Pathol. 113, 35-41.
- Kapil, R. P., Aggarwal, K., 1987. Some observations on the concurrent parasitization of *Apis florea* by *Tropilaelaps clareae* and *Euvarroa sinhai*. Exp. Appl. Acarol. 3, 267-269.
- Khongphinitbunjong, K., de Guzman, L.I., Burgett, D.M., Rinderer, T.E., Chantawannakul, P., 2012. Behavioural responses underpinning resistance and susceptibility of honeybees to *Tropilaelaps mercedesae*. Apidologie. doi:10.1007/s13592-012-0129-x.
- Khongphinitbunjong, K., de Guzman, L.I., Buawangpong, N., Rinderer, T.E., Frake, A.M., Chantawannakul, P., 2013. Observations on the removal of brood inoculated with *Tropilaelaps mercedesae* (Mesostigmata: Laelapidae) and the mite's reproductive success in *Apis mellifera* colonies. Exp. Appl. Acarol. 62, 47-55.
- Khongphinitbunjong, K., de Guzman, L.I., Matthew, R.T., Rinderer, T.E., Chen, Y., Chantawannakul, P., 2015a. Differential viral levels and immune gene expression in three stocks of *Apis mellifera* induced by different numbers of *Varroa destructor*. J. Insect Physiol. 72, 28-34.
- Khongphinitbunjong, K., de Guzman, L.I., Matthew, R.T., Rinderer, T.E., Chantawannakul, P., 2015b. Interactions of *Tropilaelaps mercedesae*, honey bee viruses and immune response in *Apis mellifera*. J. Apic. Res. 54, 40-47.
- Kirrane, M.J., de Guzman, L.I., Rinderer, T.E., Frake, A.M., Wagnitz, J., Whelan, P.M., 2011. Asynchronous development of honey bee host and *Varroa destructor* (Mesostigmata: Varroidae) influences reproductive potential of mites. J. Econ. Entomol. 104, 1146-1152.

- Kojima, Y., Toki, T., Morimoto, T., Yoshiyama, M., Kimura, K., Kadowaki, T., 2011. Infestation of Japanese native honey bees by tracheal mite and virus from non-native European honey bees in Japan. *Microbiol. Ecol.* 62, 895-906.
- Laigo, F.M., Morse, R.A., 1968. The mite *Tropilaelaps clareae* in *Apis dorsata* colonies in the Philippines. *Bee World* 49, 116-118.
- Livak, K.J., Schnittgen, T.D., 2001. Analysis of relative gene expression data using Real-Time Quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. *Methods* 25, 402-408.
- Martin S.J., Highfield A.C., Brettell L., Villalobos E.M., Budge G.E., Powell M., Nikaido S., Schroeder D.C., 2012 Global honey bee viral landscape altered by a parasitic mite. *Science* 336, 1304-1306.
- Möckel N., Gisder S., Genersch E. 2011. Horizontal transmission of deformed wing virus: pathological consequences in adult bees (*Apis mellifera*) depend on the transmission route. *J. Gen. Virol.* 92, 370-377.
- Moritz, R.F.A., Härtel, S., Neumann, P., 2005. Global invasions of the western honeybee (*Apis mellifera*) and the consequences for biodiversity. *Ecoscience* 12, 289-301.
- Mutinelli, F., 2011. The spread of pathogens through trade in honey bees and their products (including queen bees and semen): overview and recent developments. *Rev. sci. tech. Off. int. Epiz.* 30, 257-271.
- Nazzi, F., Brown, S.P., Annoscia, D., Del Piccolo, F., Di Prisco, G., Varricchio, P., Vedova, G.D., Cattonaro, F., Caprio, E., Pennacchio, F., 2012. Synergistic parasite-pathogen interactions mediated by host immunity can drive the collapse of honeybee colonies. *PLoS Pathog.* 8, e1002735.

- Neumann, P., Carreck, N., 2010. Honey bee colony losses. *J. Apic. Res.* 49, 1-6.
- Neumann, P., Radloff, S.E., Pirk, C.W.W., Hepburn, H.R., 2003. The behaviour of drifted Cape honeybee workers (*Apis mellifera capensis* Esch.): predisposition for social parasitism? *Apidologie* 34, 585-590.
- Neumann, P., Yañez, O., Fries, I., de Miranda, J.R., 2012. Varroa invasion and virus adaptation. *Trends in Parasitology* 28, 353-354.
- Neumann, P., Pettis, J.S., Schäfer, M.O. (2016) *Quo vadis Aethina tumida?* Biology and control of small hive beetles. *Apidologie* doi:10.1007/s13592-016-0426-x.
- Oldroyd, B.P., Wongsiri, S., 2006. Asian honey bees. Harvard University Press, Cambridge
- Pettis, J.S., Collins, A.M., Wilbanks, R., Feldlaufer, M.F., 2004. Effects of coumaphos on queen rearing in the honey bee, *Apis mellifera*. *Apidologie* 35, 605-610.
- Pettis, J.S., vanEngelsdorp, D., Johnson, J., Dively, G., 2012. Pesticide exposure in honey bees results in increased levels of the gut pathogen *Nosema*. *Naturwissenschaften* 99, 153-158.
- Pettis, J.S., Rose, R., Lichtenberg, E.M., Chantawannakul, P., Buawangpong, N. Somana, W., Sukumalanand, P., vanEngelsdorp, D., 2013. A rapid survey technique for *Tropilaelaps* mite (Mesostigmata: Laelapidae) detection. *J. Econ. Entomol.* 106, 1535-1554.
- Retschnig, G., Neumann, P., Williams, G.R., 2014a. Thiacloprid–*Nosema ceranae* interactions in honey bees: Host survivorship but not parasite reproduction is dependent on pesticide dose. *J. Invertebr. Pathol.* 118, 18-19.

- Retschnig, G., Williams, G.R., Mehmman, M.M., Yañez, O., de Miranda, J., Neumann, P., 2014b. Sex-specific differences in pathogen susceptibility in honey bees (*Apis mellifera*). PLoS One 9, e85261
- Rosenkranz, P., Aumeier, P., Ziegelmann, B., 2010. Biology and control of *Varroa destructor*. J. Invertebr. Pathol. 103, 96-119.
- Roberts, J.M.K., Anderson, D.L., Tay, W.T., 2015. Multiple host shifts by the emerging honeybee parasite, *Varroa jacobsoni*. Mol. Ecol. 24, 2379-2391.
- Sammataro, D., Gerson, U., Needham, G., 2000. Parasitic mites of honey bees: life history, implications, and impact. Ann. Rev. Entomol. 45, 519-548.
- Sanpa, S., Chantawannakul, P., 2009. Survey of six bee viruses using RT-PCR in Northern Thailand. J. Invertebr. Pathol. 100, 116-119.
- Simone-Finstrom, M., Evans, J. D., Marla, S., 2009. Resin collection and social immunity in honey bees. Evolution 63, 3016-3022. doi:10.1111/j.1558-5646.2009.00772.x.
- Williams, G.R., Rogers, R.E.L., Kalkstein, A.L., Taylor, B.A., Shutler, D., Ostiguy, N., 2009 Deformed wing virus in western honey bees (*Apis mellifera*) from Atlantic Canada and the first description of an overtly-infected emerging queen. J. Invertebr. Pathol. 101, 77-79.
- Williams, G.R., Dietemann, V., Ellis, J.D., Neumann, P., 2012. An update on the COLOSS network and the 'BEEBOOK: standard methodologies for *Apis mellifera* research'. J. Apic. Res. 51, 151-153.

- Williams, G.R., Alaux, C., Costa, C., Csáki, T., Doublet, V., Eisenhardt, D., Fries, I., Kuhn, R., McMahon, D.P., Medrzycki, P., Murray, T.E., Natsopoulou, M.E., Neumann, P., Oliver, R., Paxton, R.J., Pernal, S.F., Shutler, D., Tanner, G., van der Steen, J.J.M., Brodschneider, R., 2013. Standard methods for maintaining adult *Apis mellifera* in cages under in vitro laboratory conditions. In V. Dietemann., J.D. Ellis., P. Neumann (Eds.) The COLOSS BEEBOOK, Volume I: standard methods for *Apis mellifera* research. J. Apic. Res. 52
doi:10.3896/IBRA.1.52.1.04.
- Williams, G.R., Shutler, D., Burgher-MacLellan, K.L., Rogers, R.E.L., 2014. Intra-population and -community dynamics of the parasites *Nosema apis* and *Nosema ceranae*, and consequences for honey bee (*Apis mellifera*) hosts. PLoS ONE 9, e99465.
- Woyke, J., 1987a. Length of stay of the parasitic mite *Tropilaelaps clareae* outside sealed honeybee brood cells as a basis for its effective control. J. Apic. Res. 26, 104-109.
- Woyke, J., 1987b. Length of successive stages in the development of the mite *Tropilaelaps clareae* in relation to honeybee brood age. J. Apic. Res. 26, 110-114.
- Woyke, J., 1987c. Comparative population dynamics of *Tropilaelaps clareae* and *Varroa jacobsoni* mites on honey bees. J. Apic. Res. 26, 196-202.
- Yang, X., Cox-Foster, D.L., 2005. Impact of an ectoparasite on the immunity and pathology of an invertebrate: evidence for host immunosuppression and viral amplification. Proc. Natl. Acad. Sci. U.S.A. 102, 7470-7475.

- Yang, X., Cox-Foster, D., 2007. Effects of parasitization by *Varroa destructor* on survivorship and physiological traits of *Apis mellifera* in correlation with viral incidence and microbial challenge. *Parasitology* 134, 405-412.
- Yue, C., Genersch, E., 2005. RT-PCR analysis of Deformed wing virus in honeybees (*Apis mellifera*) and mites (*Varroa destructor*). *J. Gen. Virol.* 86, 3419-3424.
- Yue, C., Schröder, M., Gisder, S., Genersch E., 2007. Vertical transmission routes for deformed wing virus of honeybees (*Apis mellifera*). *J. Gen. Virol.* 88, 2329-2336.

Figure captions

Figure 1 Weight of newly emerged *Apis mellifera* workers. Bars with different letters are significantly different (Tukey-HSD test, $P < 0.001$). C/C = undisturbed brood, brood without *Tropilaelaps mercedesae* mite inoculation, I/C = brood with their cell cappings opened and closed without mite infestation, I/T = brood infested with the mite, n = number of individual bees.

Figure 2 Proportion of newly emerged *Apis mellifera* workers with deformed wings. Bars with different letters are significant different (Tukey-HSD test, $P < 0.01$). C/C = undisturbed brood, brood without *Tropilaelaps mercedesae* mite inoculation, I/C = brood with their cell cappings opened and closed, without mite infestation, I/T = brood infested with the mite, n = number of individual colonies.

Figure 3 Longevity of *Apis mellifera* workers. Number of days honey bees survived in cages among treatments. Bars with different letters are significantly different from each other (Log rank test, $P < 0.0001$). C/C = undisturbed brood, brood without *Tropilaelaps mercedesae* mite inoculation, I/C = brood with their cell cappings opened and closed without mite infestation, I/T = brood infested with the mite.

Figure 4 Log fold change of DWV levels in adult bees for: (a) all treatment groups regardless of clinical symptoms, and (b) all treatment groups considering clinical symptoms. For each group, bars with different letters are significantly different from each other ((a): Tukey-HSD test, $P < 0.01$; (b): Student's t test, $P < 0.05$). C/C = undisturbed brood, brood without *Tropilaelaps mercedesae* mite inoculation, I/C = brood with their cell cappings opened and closed without mite infestation, I/T = brood infested with *T. mercedesae*, n = number of individual bees, * calibrated to the lowest DWV level.

Figure 1

Figure 2

Figure 3

Figure 4

The mite *Tropilaelaps mercedesae* reduces western honey bee, *Apis mellifera*, host longevity and emergence weight, and promotes both Deformed wing virus levels and associated clinical symptoms. This suggests that this mite is a potentially important parasite to *A. mellifera*.