Phylogenetic structure and habitat associations of *Beauveria* species isolated from soils in Slovakia

Juraj Medo^{1*}, Jaroslav Michalko^{2,4}, Janka Medová³, Ľudovít Cagán¹

¹ Slovak University of Agriculture in Nitra, Tr. A. Hlinku 2, 949 76 Nitra, Slovakia

² Institute of Plant Genetics and Biotechnology, Slovak Academy of Sciences, Akademická 2, 950 07 Nitra, Slovakia

³ Constantine the Philosopher University in Nitra, Tr. A. Hlinku 1, 949 74 Nitra, Slovakia

⁴ Institute of Forest Ecology, Arboretum Mlynany, Slovak Academy of Sciences, Vieska nad Zitavou 178, 951 52 Slepčany, Slovakia

*corresponding author e-mail juraj.medo@uniag.sk, tel +421 37 641 5810

Abstract

The phylogenetic structure of 109 soil-borne entomopathogenic *Beauveria* isolates acquired using the *Galleria mellonella* bait method from different habitat types in Slovakia was determined by sequence analysis of their ITS and Bloc loci. Three *Beauveria* species were identified; *Beauveria bassiana*, *B. pseudobassiana* and *B. brongniartii*, represented by 51.4 %, 43.1 % and 5.5 % of acquired isolates, respectively, which were resolved into 15, 1 and 1 distinguishable haplotypes. Correlation analysis with the habitat type and individual habitat characteristics showed strong preferences of the most prevalent haplotypes for agricultural (*B. bassiana* A1) and forest habitats (*B. pseudobassiana*) which has possible implications for conservative biocontrol strategies.

Keywords: phylogenetic structure, *Beauveria* spp., entomopathogenic fungi, habitat preference

1 Introduction

Anamorphic entomopathogenic fungi of the genus *Beauveria* belong to the most commonly isolated facultative pathogens of insect pests (Chandler et al., 1997; Bidochka et al., 1998; Vega et al., 2012). A high level of genetic diversity is commonly detected within natural *Beauveria* populations. General phylogenetic markers such as intergenic spacer (ITS) or elongation factor 1- α are suitable for distinguishing between *Beauveria* species but are not

able to characterize this diversity in deep. Recently developed highly variable markers, specifically the Bloc marker (a nuclear intergenic marker developed for phylogenetic investigation of *Beauveria*) as well as microsatellites enable further resolution of *Beauveria* isolates at the level of individual genotypes (Coates et al., 2002; Meyling and Eilenberg, 2006; Castrillo et al., 2007; Hollingsworth et al., 2011). Phylogenetic studies using these markers have shown that diversity within the *Beauveria* genus could be both on inter- and intra-specific level. For example, it has been found that *B. bassiana sensu lato* and *B. brongniartii* represent a complex of morphologically cryptic phylogenetic species (Rehner, 2005; Rehner et al., 2006), which led to a revision of *Beauveria* taxonomy. Three morphologically similar phylogenetic lineages within the *B. bassiana* complex were reclassified as *B. pseudobassiana* (previously designated as Clade C), *B. varroae* and *B. kipukae*. Additionally, two sister lineages within the *B. brongniartii* complex were reclassified as *B. asiatica* and *B. australis*, which increased the number of currently known *Beauveria* species to twelve (Rehner et al., 2011). To distinguish between the remaining thus far unresolved lineages within the *B. bassiana* complex a temporary annotation system has been proposed, which designates lineages based on the continent from which they were isolated followed by a digit indicating the order of their discovery, e.g. Eu_1 (Meyling et al., 2009). However, many detected lineages have a worldwide distribution; therefore, an *ad hoc* naming system based on Rehner and Buckley (2005) without indication of continent should be preferred (e.g. A1), and a reference isolate for each lineage should be deposited in a culture collection (Stephen A. Rehner, personal communication).

Soil is the most commonly examined environment when assessing *Beauveria* diversity. Baiting techniques using susceptible larvae of *Galleria mellonella* (Lepidoptera: Pyralidae), commonly referred to as the “*Galleria* bait method” (GBM), or *Tenebrio molitor* (Coleoptera: Tenebrioidae) are most widely used for isolation of soil-borne *Beauveria* fungi, although the use of selective media or a combination of both methods may improve the isolation efficacy (Meyling et al., 2009; Medo and Cagán, 2011). Three species of *Beauveria* are commonly detected using these methods: *B. bassiana*, *B. pseudobassiana*, and *B. brongniartii*. Using the GBM *B. bassiana* A1 was found by Pérez-González et al. (2014) to be the most common genotype in Mexican agricultural soils. The GBM was also used by Sevim et al. (2010) to find that *B. pseudobassiana* and *B. bassiana* A1 and A11 were the most abundant lineages in Turkey. Meyling et al. (2009), using baiting techniques as well as selective media for isolation of fungi from soil, detected lineages A1 (labeled as Eu_1), A3 (Eu_3) and A11 (Eu_6) in their study in Denmark. Lineages A6 (named as Wd-2) and A3

(Eu_3) were also identified by Garrido-Jurado et al. (2011), who analyzed the EF1- α region of Spanish *B. bassiana* isolates from culture collection. Additionally, lineages Eu_8 and Eu_9 were reported by these authors. However, Rehner and Buckley (2005) reported much higher diversity of *Beauveria* isolates from culture collections collected throughout Europe.

Despite improvements made in isolation, taxonomy, and genotyping of *Beauveria* fungi, major factors determining their genotypic structure at the population level remain poorly understood. Previous studies have linked a particular genetic group of the fungus with its geographical origin (Reay et al., 2010; Sevim et al., 2010; Meyling et al., 2012b; Cai et al., 2013), although some genetic groups have a world-wide distribution. A large number of studies have tried to find an association between a particular genetic group and its insect host (Maurer et al., 1997; Rehner et al., 2006; Reay et al., 2008; Meyling et al., 2012a; Wang et al., 2013). However, only limited evidence has been found to support this association and the general conclusion is that genetic groups in *Beauveria* spp. do not have any particular host preferences (Meyling et al., 2009). Certain studies have proposed that habitat-related environmental factors constitute the prime selective forces shaping the genetic structure of facultative entomopathogenic fungi (Vanninen, 1996; Bidochka et al., 1998, 2001, 2002; Meyling and Eilenberg, 2007; Quesada-Moraga et al., 2007; Meyling et al., 2009; Medo and Cagán, 2011). Despite the existing consensus, few studies have investigated this hypothesis (Bidochka et al., 2002; Meyling et al., 2009). In our study we addressed this question by phylogenetic and correlation analysis of a subset of *Beauveria* soil isolates from different habitat types across Slovakia isolated using the GBM.

2 Material and methods

2.1 Soil sampling and isolation of fungi

Soil samples were collected from various locations within Slovakia during the year 2008 (Figure 1A, Supplementary Table S1) and processed as described in Medo and Cagán (2011). The sampling sites were selected with the aim of maximizing the spectrum of environmental conditions covered (altitude, soil characteristics, vegetation type). The minimum distance between the adjacent sampling sites was 1 km. For each soil sample the habitat type (forest, field, meadow or hedgerow), altitude of the sampling site, type of the vegetation canopy, and basic soil characteristics (pH, texture) were recorded (Supplementary Table S1). *Beauveria*

fungi were isolated from soil samples using the *Galleria mellonella* baiting technique as described previously (Medo and Cagán, 2011) and cultured on Sabouraud dextrose agar (SDA) (Merck KGaA).

2.2 Genetic analysis

DNA was extracted from two-week old single spore fungal cultures using the ZR fungal/bacterial DNA extraction kit (Zymo Research Corp. USA). All PCR reaction mixtures contained 200 mM dNTPs, 1x DreamTaq buffer, 0.5 unit DreamTaq DNA polymerase (Life technologies, USA), 0.5 mM of the corresponding primer, and 0.5 µl of non-diluted DNA. Cycling conditions were as follows: 95 °C for 3 min followed by 35 cycles of 95 °C for 30 s, annealing at corresponding temperature for each primer pair for 45 s, 72 °C for 90 s, and a final elongation at 72°C for 10 min. Primers used for PCR and sequencing of ITS region were ITS1 and ITS4 (White et al., 1990). Primers 5.1F and B822Ldg were used for *B. pseudobassiana* and 5.1F and 3.1R for other *Beauveria* isolates to amplify the nuclear intergenic region Bloc. Amplification products were sequenced with primers B822Ldg and B22Udg (Rehner et al., 2011).

Sequences were analyzed in an ABI310 genetic analyzer using BigDye 3.1 sequencing chemistry (Life technologies, USA). Acquired sequences were assembled and processed using the Seaview software (Gouy et al., 2010). Alignment was made using MUSCLE (Edgar, 2004). The acquired unique haplotype sequences as well as the reference sequences (for Genbank accessions see Supplementary Table S2) were used for phylogenetic analysis. A phylogenetic tree was constructed using Phy ML (Guindon et al., 2010) with a GTR substitution model, a BioNJ starting tree with the best of NNI and SPR tree searching, and followed by 1000 bootstrap replications.

2.3 Statistical analyses

The presence/absence of certain genetic groups between two habitats was evaluated using cross-tabulation followed by the Fisher exact test and the odds ratio was then calculated. The occurrence of genetic groups at different soil pH's and altitudes was evaluated using the analysis of variance along with the Tukey HSD test. These statistics were only computed for *Beauveria* groups with a sufficient number of representatives, i.e. *B. pseudobassiana* and *B. bassiana* clade A1. All statistical analyses were performed using the Statgraphics XV package (StatPoint Technologies Inc. USA). The estimates of haplotypic diversity were calculated

according to Stoddart and Taylor (1988) and expressed as the index of haplotypic diversity (G).

3 Results and discussion

Maximum likelihood analysis of the combined ITS and Bloc data resolved 109 analyzed *Beauveria* isolates into 8 phylogenetic clades (Figure 1B) represented by three *Beauveria* species: *B. bassiana*, *B. pseudobassiana* and *B. brongniartii* containing 56 (54.1 %), 47 (43.1 %) and 6 (5.5 %) isolates, respectively. Isolates of *B. pseudobassiana* and *B. brongniartii* were each represented by a single phylogenetic clade and haplotype while isolates of *B. bassiana* were resolved into six distinct phylogenetic clades (A1, A2, A3, A6, A11 and AX containing respectively 42, 2, 9, 1, 1 and 1 isolates) comprising 15 fungal haplotypes (Figure 1B).

The recognized species composition of *Beauveria* fungi in Slovakian soils is similar to that reported by other authors in and outside of Europe (Meyling et al., 2009; Sevim et al., 2010; Garrido-Jurado et al., 2011; Pérez-González et al., 2014; Mayerhofer et al., 2015). The genotypic composition of *Beauveria* spp., on the other side, varies among studies from different geographic areas. Although the comparison of results among studies is not straightforward due to differences in employed protocols for fungal isolation and different genetic markers for isolate genotyping, the composition of main *Beauveria* lineages appears to be relatively stable across Europe with *B. bassiana* A1 and *B. pseudobassiana* being the predominant lineages. Whether this indicates that the less frequent lineages are worse competitors in the soil environment or that their occurrence is restricted to specific environmental conditions remains to be determined.

Within individual habitats, we found that the haplotypic diversity (G) of *Beauveria* isolates was highest in field soils ($G = 5.39$) and lowest in forests soils ($G = 1.53$). Only three (A1g, A3b and AX) out of 15 identified *B. bassiana* haplotypes were isolated from forest soils (Figure 2). The abundance of the two most frequent *Beauveria* clades, *B. bassiana* A1 and *B. pseudobassiana* differed significantly among habitats. The odds of finding *B. bassiana* A1 within field soils was 45.78 times higher than odds to find it within forest soils ($p < 0.001$). A significant difference was also found between meadow and forest ($p < 0.001$, odds ratio 476.19), hedgerow and forest ($p = 0.0043$, odds ratio 19.19), and meadow and field habitats ($p = 0.013$, odds ratio 10.20). In regards to *B. pseudobassiana*, differences in its occurrence were significant between the forest and field ($p < 0.001$, odds ratio 12.67), forest and meadow

($p < 0.001$, odds ratio 48.00), and forest and hedgerow habitats ($p = 0.0152$, odds ratio 8.00). *B. brongniartii* was isolated mostly from forest soils (66.6 %); however, it was only represented by 6 isolates. There is a well-known affinity of certain genus/species of entomopathogenic fungi for specific habitats (e.g. *Metharizium* is common in agricultural soils, while *Isaria* in forest soils). The differential preference of *B. pseudobassiana* towards natural habitats and *B. bassiana* towards agricultural habitats is in accordance with previous studies (Sevim et al., 2010; Pérez-González et al., 2014). In contrast, Quesada-Moraga et al., 2007 found Spanish isolates of *B. bassiana* morphospecies equivalently common in both natural and cultivated soils, although no genetic discrimination of isolates was made in their study.

To increase our understanding of environmental factors that could underlay *Beauveria* spp. habitat preferences we evaluated the influence of individual habitat characteristics on the haplotypic structure of *Beauveria* isolates (Figure 2A). The diversity of *Beauveria* spp. haplotypes was strongly affected by both the sampling altitude and the soil pH. The highest diversity was found at the lowest altitudes (< 205 m; $G = 5.36$) and in more basic soils ($\text{pH} > 7.4$, $G = 4.80$), whereas diversity was lowest at the highest altitudes (> 400 m, $G = 3.01$) and in more acidic soils ($\text{pH} < 6.6$, $G = 2.93$) (Figure 2B). Statistically significant differences were found in the occurrence of *B. bassiana* clade A1 and *B. pseudobassiana* at different altitudes (301 resp. 387 m, $p = 0.0342$) as well as at different soil pH (7.14 resp. 6.62, $p = 0.005$). Interestingly, *B. bassiana* haplotypes A1g and A1h were found to have inverse preferences in regards to altitude. Soils with loamy texture had much higher *Beauveria* haplotypic diversity ($G = 6.29$) than sandy, clay or organic soils ($G = 2.63$, 2.38 and 1.57, respectively). The haplotypic diversity of non-forest vegetation types ($G = 4.62 \pm 1.33$) was about three times higher than that observed in forest vegetation ($G = 1.54 \pm 0.23$) (Figure 2B). Similarly, Quesada-Moraga et al. (2007) isolated *Beauveria* spp. more commonly in slightly basic soils, in soils with lower organic content and at lower altitudes (below 700 m).

While all recorded habitat characteristics seem to influence to some extent the haplotypic composition of soil-borne *Beauveria* spp., their individual importance needs to be tested experimentally as several habitat characteristics tend to be inter-correlated (e.g. the analyzed forest soils had generally lower pH and occurred at higher altitudes than the field samples and *vice versa*). In addition, Bidochka et al. (2002) reported association of certain *Beauveria* and *Metarhizium* genetic groups with different level of UV irradiation or temperature regions which were not assessed in our study. It is therefore likely that complex relationships among habitat-associated factors influence genotypic composition of *Beauveria* fungi in nature. Nevertheless, the strong preferences of certain *Beauveria* phylogenetic groups

for specific habitats could provide useful information regarding their more targeted usage as biocontrol agents within a particular habitat, especially if their long-term persistence in an environment is desired.

6 Acknowledgements

The authors would like to thank Dr. Stephen A. Rehner, USDA-ARS, Beltsville MD, for advices and tips on *Beauveria* sequencing and phylogeny. We are also thankful to Allwin McDonald for language corrections. This work was financially supported by the national project VEGA 1/0732/14.

REFERENCES

- Bidochka, M. J., Kamp, A. M., Lavender, T. M., Dekoning, J., De Croos, J. N. A. 2001. Habitat association in two genetic groups of the insect-pathogenic fungus *Metarhizium anisopliae*: Uncovering cryptic species? *Appl. Environ. Microbiol.*, 67, 3, 1335-1342
- Bidochka, M. J., Kasperski, J. E., Wild, G. A. M. 1998. Occurrence of the entomopathogenic fungi *Metarhizium anisopliae* and *Beauveria bassiana* in soils from temperate and near-northern habitats. *Can. J. Bot.*, 76, 7, 1198-1204
- Bidochka, M. J., Menzies, F. V., Kamp, A. M. 2002. Genetic groups of the insect-pathogenic fungus *Beauveria bassiana* are associated with habitat and thermal growth preferences. *Arch. Microbiol.*, 178, 6, 531-537
- Cai, Y., Pu, S., Nie, Y., Rehner, S., Huang, B. 2013. Discrimination of Chinese *Beauveria* strains by DGGE genotyping and taxonomic identification by sequence analysis of the Bloc nuclear intergenic region. *Appl. Entomol. Zool.*, 48, 3, 255-263
- Castrillo, L.A., Ugine, T.A., Filotas, M.J., Sanderson, J.P., Vandenberg, J.D., Wraight, S.P. 2007. Molecular characterization and comparative virulence of *Beauveria bassiana* isolates (Ascomycota: Hypocreales) associated with the greenhouse shore fly, *Scatella tenuicosta* (Diptera: Ephydriidae). *Biological Control* 45, 154– 162
- Chandler, D., Hay, D., Reid, A.P. 1997. Sampling and occurrence of entomopathogenic fungi and nematodes in UK soils. *Appl. Soil Ecol.* 5, 133–141.
- Coates, B.S., Hellmich, R.L., Lewis L.C. 2002. *Beauveria bassiana* haplotype determination based on nuclear rDNA internal transcribed spacer PCR–RFLP. *Mycological Research* 106, 1, 40-50.
- Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research*, 32, 5, 1792-1797.
- Enkerli, J., Widmer, F. 2010. Molecular ecology of fungal entomopathogens: molecular genetic tools and their applications in population and fate studies. *Biocontrol*, 55, 1, 17-37
- Garrido-Jurado, I., Marquez, M., Ortiz-Urquiza, A., Santiago-Alvarez, C., Iturriaga, E., Quesada-Moraga, E., Monte, E., Hermosa, R. 2011. Genetic analyses place most Spanish isolates of *Beauveria bassiana* in a molecular group with world-wide distribution. *BMC Microbiology*, 11, 1, 84

- Gouy, M., Guindon, S., Gascuel, O. 2010. SeaView Version 4: A multiplatform graphical user interface for sequence alignment and phylogenetic tree building. *Mol. Biol. Evol.*, 27, 2, 221-224
- Guindon, S., Dufayard, J.-F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O. 2010. New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Syst. Biol.*, 59, 3, 307-321
- Hollingsworth, R. G., Lysy, A. M., & Matsumoto, T. K. 2011. Preliminary study of genetic variation in Hawaiian isolates of *Beauveria bassiana* [Hypocreales, Cordycipitaceae]. *Journal of invertebrate pathology* 106, 3, 422-425.
- Maurer, P., Couteaudier, Y., Girard, P. A., Bridge, P. D., Riba, G. 1997. Genetic diversity of *Beauveria bassiana* and relatedness to host insect range. *Mycol Res*, 101, 2, 159-164
- Mayerhofer, J., Enkerli, J., Zelger, R., Strasser, H. 2015. Biological control of the European cockchafer: persistence of *Beauveria brongniartii* after long-term applications in the Euroregion Tyrol. *BioControl*, 60, 5, 617-629.
- Medo, J., Cagán, Ľ. 2011. Factors affecting the occurrence of entomopathogenic fungi in soils of Slovakia as revealed using two methods. *Biol. Control*, 59, 2, 200-208
- Meyling, N.V., Eilenberg, J., 2006. Isolation and characterisation of *Beauveria bassiana* isolates from phylloplanes of hedgerow vegetation. *Mycological Research* 110, 188–195.
- Meyling, N. V., Eilenberg, J. 2007. Ecology of the entomopathogenic fungi *Beauveria bassiana* and *Metarhizium anisopliae* in temperate agroecosystems: potential for conservation biological control. *Biol. Control*, 43, 2, 145-155
- Meyling, N. V., Lübeck, M., Buckley, E. P., Eilenberg, J., Rehner, S. A. 2009. Community composition, host range and genetic structure of the fungal entomopathogen *Beauveria* in adjoining agricultural and seminatural habitats. *Mol. Ecol.*, 18, 6, 1282-1293
- Meyling, N. V., Pilz, C., Keller, S., Widmer, F., Enkerli, J. 2012a. Diversity of *Beauveria* spp. isolates from pollen beetles *Meligethes aeneus* in Switzerland. *J. Invertebr. Pathol.*, 109, 1, 76-82
- Meyling, N. V., Schmidt, N. M., Eilenberg, J. 2012b. Occurrence and diversity of fungal entomopathogens in soils of low and high Arctic Greenland. *Polar Biol.*, 35, 9, 1439-1445
- Pérez-González, V. H., Guzmán-Franco, A. W., Alatorre-Rosas, R., Hernández-López, J., Hernández-López, A., Carrillo-Benítez, M. G., Baverstock, J. 2014. Specific diversity of the entomopathogenic fungi *Beauveria* and *Metarhizium* in Mexican agricultural soils. *J. Invertebr. Pathol.*, 119, 0, 54-61

- Quesada-Moraga, E., Navas-Cortés, J. A., Maranhao, E. A. A., Ortiz-Urquiza, A., Santiago-Álvarez, C. 2007. Factors affecting the occurrence and distribution of entomopathogenic fungi in natural and cultivated soils. *Mycol. Res.*, 111, 8, 947-966
- Reay, S. D., Brownbridge, M., Cummings, N. J., Nelson, T. L., Souffre, B., Lignon, C., Glare, T. R. 2008. Isolation and characterization of *Beauveria* spp. associated with exotic bark beetles in New Zealand *Pinus radiata* plantation forests. *Biol. Control*, 46, 3, 484-494
- Reay, S. D., Brownbridge, M., Gicquel, B., Cummings, N. J., Nelson, T. L. 2010. Isolation and characterization of endophytic *Beauveria* spp. (Ascomycota: Hypocreales) from *Pinus radiata* in New Zealand forests. *Biol. Control*, 54, 1, 52-60
- Rehner, S. A. 2005. Phylogenetics of the insect pathogenic genus *Beauveria* (pp. 3-27). Oxford: Oxford University Press.
- Rehner, S., Buckley, E. 2005. A *Beauveria* phylogeny inferred from nuclear ITS and EF1-alpha sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia*, 97, 84 – 98
- Rehner, S. A., Minnis, D., Sung, G.-H., Luangsa-ard, J. J., DeVotto, L., Humber, R. A. 2011. Phylogeny and systematics of the anamorphic, entomopathogenic genus *Beauveria*. *Mycologia*, 10-302
- Rehner, S. A., Posada, F., Buckley, E. P., Infante, F., Castillo, A., Vega, F. E. 2006. Phylogenetic origins of African and Neotropical *Beauveria bassiana* s.l. pathogens of the coffee berry borer, *Hypothenemus hampei*. *J. Invertebr. Pathol.*, 93, 1, 11-21
- Sevim, A., Demir, I., Höfte, M., Humber, R. A., Demirbag, Z. 2010. Isolation and characterization of entomopathogenic fungi from hazelnut-growing region of Turkey. *Biocontrol*, 55, 2, 279-297.
- Stoddart, J. A., Taylor, J. F. 1988. Genotypic diversity: estimation and prediction in samples. *Genetics* 118, 4, 705-11.
- Vanninen, I. 1996. Distribution and occurrence of four entomopathogenic fungi in Finland: Effect of geographical location, habitat type and soil type. *Mycol. Res.*, 100, 93-101
- Vega, F. E., Meyling, N. V., Luangsa-ard, J. J., Blackwell, M. 2012. Chapter 6 - Fungal Entomopathogens. In F. E. Vega H. K. Kaya (Eds.), *Insect Pathology (Second Edition)* (pp. 171-220). San Diego: Academic Press.
- Wang, J.-j., Yang, L., Qiu, X., Liu, Y.-g., Zhou, W., Wan, Y.-J. 2013. Diversity analysis of *Beauveria bassiana* isolated from infected silkworm in southwest China based on

molecular data and morphological features of colony. *World J. Microbiol. Biotechnol.* 29, 7, 1263-1269

White, T., Bruns, T., Lee, S., Taylor, J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. Innis, D. Gelfand, J. Shinsky T. White (Eds.), *PCR Protocols: A Guide to Methods and Applications* (pp. 315-322): Academic Press.

A

Figure2