Stochastic Unrelatedness, Couplings, and Contextuality

Ehtibar N. Dzhafarov*
Purdue University

R. Duncan Luce once mentioned in a conversation that he did not consider Kolmogorov's probability theory well-constructed because it treats stochastic independence as a "numerical accident," while it should be treated as a fundamental relation, more basic than the assignment of numerical probabilities. I argue here that stochastic independence is indeed a "numerical accident," a special form of stochastic dependence between random variables (most broadly defined). The idea that it is fundamental may owe its attractiveness to the confusion of stochastic independence with stochastic unrelatedness, the situation when two or more random variables have no joint distribution, "have nothing to do with each other." Kolmogorov's probability theory cannot be consistently constructed without allowing for stochastic unrelatedness, in fact making it a default situation: any two random variables recorded under mutually incompatible conditions are stochastically unrelated. However, stochastically unrelated random variables can always be probabilistically coupled, i.e., imposed a joint distribution upon, and this generally can be done in an infinity of ways, independent coupling being merely one of them. The notions of stochastic unrelatedness and all possible couplings play a central role in the foundation of probability theory and, especially, in the theory of probabilistic contextuality.

KEYWORDS: contextuality, coupling, joint distribution, probability, random variable, stochastic relation, stochastic unrelatedness.

I. INTRODUCTION

Almost 15 years ago R. Duncan Luce mentioned in a conversation that the Kolmogorovian probability theory (KPT) was unsatisfactory because it treated stochastic independence as a "numerical accident" rather than a fundamental relation. If I roll a die today in Irvine, California, Duncan said, and on another day you roll a die in Lafayette, Indiana, the fact that the two outcomes are independent cannot be established by checking the multiplication rule. On the contrary, the applicability of the multiplication rule in this case is justified by determining that the two dice are stochastically independent, "have nothing to do with each other."

This simple example (some may think too simple to be of great interest) leads us to the very foundations of probability theory. Let us try to understand it clearly by comparing it to two other examples. The first one is about a situation when I repeatedly roll a single die, having defined two random variables:

$$A = \begin{cases} 1 & \text{if the outcome is even} \\ 0 & \text{otherwise} \end{cases},$$
$$B = \begin{cases} 1 & \text{if the outcome exceeds 3} \\ 0 & \text{otherwise} \end{cases}.$$

$\Pr[A = 1 \text{ and } B = 1]$, $\Pr[A = 1]$, and $\Pr[B = 1]$ (I will use $\Pr$ as a symbol for probability throughout this paper): if the joint probability turns out to be the product of the two marginal ones (statistical issues aside), the two events are determined to be independent. I cannot simply make this determination a priori, as it depends on what die I am rolling: if it is a fair die, A and B are not independent, but if the distribution of the outcomes is

$$\begin{array}{l} \text{value :} \quad 1 \quad 2 \quad 3 \quad 4 \quad 5 \quad 6 \\ \text{pr.mass :} \quad 0 \quad \frac{1}{4} \quad \frac{1}{4} \quad \frac{1}{4} \quad \frac{1}{4} \quad 0 \end{array},$$

then A and B are independent. The difference between this example and that of Duncan Luce's is not in the number of the dice being rolled: the second example would not change too much if I roll two dice together, having marked them "Left" and "Right," and define the random variables as

$$A = \begin{cases} 1 & \text{if the Left outcome is even} \\ 0 & \text{otherwise} \end{cases},$$
$$B = \begin{cases} 1 & \text{if the Right outcome exceeds 3} \\ 0 & \text{otherwise} \end{cases}.$$

The realizations of A and B again come together, this time the empirical meaning of the "togetherness" being "in the same trial," or "simultaneously." Again, one can also say that the two random variables co-occur because they are functions of one and the same "background" random variable Z , only this time it is the pair of values rather than a single one. And again, I can estimate from the observations the probabilities $\Pr[A = 1 \text{ and } B = 1]$, $\Pr[A = 1]$, and $\Pr[B = 1]$ and check their adherence to the multiplication rule. Whether the two random variables are stochastically independent is determined by the outcome of this test: the dice may very well be rigged not to be independent.

In Duncan Luce's example the situation is very different: the outcomes of rolling the two dice in two different places

* E-mail: ehtibar@purdue.edu

at two different times have no empirically defined pairing.
If I define my random variables as

$$A = \begin{cases} 1 & \text{if on Tuesday in Irvine the outcome is even} \\ 0 & \text{otherwise} \end{cases},$$

$$B = \begin{cases} 1 & \text{if on Friday in Lafayette the outcome exceeds 3} \\ 0 & \text{otherwise} \end{cases}$$

then I can estimate empirically the probabilities $\Pr[A = 1]$, and $\Pr[B = 1]$ and find out, e.g., that they are (statistical issues aside) 0.7 and 0.5, respectively. But I cannot estimate empirically $\Pr[A = 1 \text{ and } B = 1]$: the two random variables are not recorded in pairs. The experiment involves no empirical procedure by which one could find which value of B should be paired with which value of A . The two random variables therefore do not have an observable (estimable from frequencies) joint distribution, they cannot be presented as functions of one and the same “background” random variable. What one can do, however, is to *declare* the two random variables stochastically independent, based on one’s understanding that they “have nothing to do with each other.” If one does so, the validity of $\Pr[A = 1 \text{ and } B = 1]$ being equal to the product of two individual probabilities is true *by construction*, requiring no empirical testing and allowing for no empirical falsification.

This was Duncan Luce’s point: while the KPT defines stochastic independence through the multiplication rule, at least in some cases the determination of independence precedes and justifies the applicability of the multiplication rule. In Duncan Luce’s opinion, this warranted treating stochastic independence as a fundamental, “qualitative” relation preceding assignment of numerical probabilities. This opinion is in accordance with the general precepts of the representational theory of measurement. Thus, the authors of the first volume of Foundations of Measurement (Krantz et al., 1971) sympathetically refer to Zoltan Dornator 1969 dissertation in which he axiomatized probability theory treating stochastic independence as a primitive relation. As far as I know, however, it has not translated into a viable alternative to the KPT.

I accept Duncan Luce’s example as posing a genuine foundational problem, but I disagree that this problem is about defining independence by means other than the multiplication rule. The position I defend below in this paper is as follows.

1. Random variables that “have nothing to do with each other” are defined on different domains (sample spaces). Rather than being independent (which is a form of a joint distribution), they are *stochastically unrelated*, i.e., they possess no joint distribution.
2. It is not that we do not know the “true” distribution, or that in “truth” they are independent but we do not know how to justify this. A joint distribution simply is not defined (until imposed by us in one of multiple ways, discussed below).
3. The KPT is consistent with the idea of multiple sample spaces and in fact requires it for internal consistency:

the idea of a single sample space for all random variables imaginable is mathematically untenable.

4. Any given set of pairwise stochastically unrelated random variables can always be *coupled*, i.e., imposed a joint distribution on. This is equivalent to inventing a pairing scheme for their realizations, and this can be done in multiple ways, coupling them as independent random variables being just one of them.

II. ON RANDOM VARIABLES, UNRELATEDNESS, AND INDEPENDENCE

II.1. Informal introduction

Stochastic unrelatedness is easy to distinguish from stochastic independence: the latter assumes the existence of a joint distribution, which means that an empirical procedure exists by which each realization of one random variables can be paired (coupled) with that of another. The most familiar forms of empirical coupling are co-occurrence in the same trial and co-relation to the same person. In the table below,

$$\begin{array}{lcl} c : & 1 & 2 & 3 & 4 & 5 & \dots \\ X : & x_1 & x_2 & x_3 & x_4 & x_5 & \dots, \\ Y : & y_1 & y_2 & y_3 & y_4 & y_5 & \dots \end{array} \quad (1)$$

the indexing entity c can be the number of a trial (as in repeatedly rolling two marked dice together) or an ID of a person (as in relating heights and weights, or weights before and after dieting). The random variables X and Y here have a joint distribution: one can, e.g., estimate the probability with which X falls within an event E_X and (“simultaneously”) Y falls within an event E_Y ; and if

$$\Pr[X \in E_X \& Y \in E_Y] = \Pr[X \in E_X] \Pr[Y \in E_Y], \quad (2)$$

for any two such events E_X, E_Y , then X and Y are considered independent.

Suppose, however, that the information about c in (1) does not exist, and all one has is some set of values for X and some set of values for Y . Clearly, now the “togetherness” of $X \in E_X$ and $Y \in E_Y$ is undefined. Although $\Pr[X \in E_X]$ and $\Pr[Y \in E_Y]$ have the same meaning as before, $\Pr[X \in E_X \& Y \in E_Y]$ is undefined, and (2) cannot be tested. This is what stochastic unrelatedness is: lack of a joint distribution. A pair of stochastically unrelated random variables are neither independent nor interdependent, these terms do not apply.

Think, e.g., of a list of weights in some group of people before dieting (X) and a list of weights in some other group of people after dieting (Y): which value of X should be paired with which value of Y to try to estimate $\Pr[X \in E_X \& Y \in E_Y]$? Any pairing one can impose here will be as good as any other pairing, and none of them is determined by the empirical procedures involved (weighing people in the two groups). This simple example has counterparts in all experiments where random variables are recorded under two or more mutually exclusive conditions.

174 A question arises: couldn't one nevertheless treat
 175 stochastically unrelated X and Y as if they were inde-
 176 pendent? The answer is affirmative, but so is the answer
 177 to the question whether X and Y can be treated as if they
 178 were not independent. Treating the outcomes of rolling dice
 179 in Irvine on Tuesday (X) and in Lafayette on Friday (Y)
 180 as if they were jointly distributed means constructing an-
 181 other pair of random variables, $(\tilde{X}, \tilde{Y})$, this time a jointly
 182 distributed one, such that $\tilde{X}$ and $\tilde{Y}$ taken separately are
 183 distributed as X and Y , respectively. Such constructions
 184 form the subject of a special branch of probability theory
 185 called the *theory of coupling(s)* (Thorisson, 2000).

186 Let, e.g., both dice be fair. One can always construct
 187 $(\tilde{X}, \tilde{Y})$ by assigning probability mass $\frac{1}{36}$ to each of the 36
 188 pairs. This pair $(\tilde{X}, \tilde{Y})$ is the *independent coupling* of X
 189 and Y . Its choice corresponds to pairing every realization
 190 of X with every realization of Y (or with randomly chosen
 191 realization of Y). There is, however, no reason to single
 192 out the independent coupling. One can also make $\tilde{X}$ and $\tilde{Y}$
 193 perfectly correlated or perfectly anticorrelated by assigning
 194 the probability masses as, respectively,

$$\text{pr.mass} [\tilde{X} = x \& \tilde{Y} = y] = \begin{cases} 0 & \text{if } x \neq y \\ \frac{1}{6} & \text{if } x = y \end{cases}$$

195 OR

$$\text{pr.mass} [\tilde{X} = x \& \tilde{Y} = y] = \begin{cases} 0 & \text{if } x + y \neq 7 \\ \frac{1}{6} & \text{if } x + y = 7 \end{cases},$$

196 where $x, y \in \{1, \dots, 6\}$.

197 These couplings correspond to pairing each realization
 198 of X with only one specific realization of Y . If the dice
 199 producing X and Y have outcomes with different distribu-
 200 tions, a perfectly correlated or anticorrelated coupling will
 201 not be possible, while the independent coupling will, as it is
 202 universally applicable. But the independent coupling still
 203 will not be the only possible one (unless one of the dice
 204 is rigged to roll a single outcome, in which case the only
 205 possible coupling can be viewed as independent, perfectly
 206 correlated, or perfectly anticorrelated).

207 One may be tempted to think that the “true” pair-
 208 ing should involve ordering the observations of X and Y
 209 chronologically and pairing the outcomes with the same
 210 trial number. A brief reflection should show, however,
 211 that this is an arbitrary choice: what theoretical princi-
 212 ples would compel one to pair the first realization of X
 213 with the first realization of Y (occurring, in Duncan Luce’s
 214 example, at another time in another place), rather than
 215 with the tenth one, the last one, or one having the same
 216 quantile rank? Recall also that the chronological sequences
 217 need not be defined to begin with: instead of rolling a sin-
 218 gle die repeatedly one could roll a large number of identical
 219 dice and count the events.

220 Summarizing, a joint distribution for empirically ob-
 221 served X and Y exists only if there is an empirical proce-
 222 dure for coupling their realizations, such as relating them
 223 to one and the same value of c in (1). Otherwise X and

224 Y are stochastically unrelated. When they are, one can
 225 impose on them a joint distribution by creating a coupling
 226 $(\tilde{X}, \tilde{Y})$ for X and Y “on paper.” The individual distribu-
 227 tions of stochastically unrelated X and Y do impose some
 228 constraints on possible joint distributions of $(\tilde{X}, \tilde{Y})$, but,
 229 except in degenerate cases, do not determine it uniquely.
 230 The independent coupling is not the only possible coupling
 231 of stochastically unrelated random variables.

232 II.2. Formalizing the “naive” account of random 233 variables

234 Random variables are defined by their distributions (say
 235 probability masses associated with every possible roll of a
 236 die) and, in order to distinguish different variables having
 237 the same distribution, by their unique names (e.g., “the
 238 outcome of the die rolled in Irvine on Tuesday”). On a more
 239 general level, the distribution of a random variable called
 240 X is a probability space (S_X, Σ_X, μ_X) , with the standard
 241 meaning of the terms: S_X is the set of possible values for X ,
 242 Σ_X is a sigma-algebra of subsets of S_X , and μ a probability
 243 measure.¹ For each element E_X of Σ_X (an event) we define
 244 the probability of X “falling in E_X ” or “satisfying E_X ” as

$$\Pr[X \in E_X] = \mu_X(E_X). \quad (3)$$

245 Given another random variable, called Y and distributed
 246 as (S_Y, Σ_Y, μ_Y) , we say it is jointly distributed with X
 247 if there is a random variable $Z = (X, Y)$ whose name
 248 is “ordered pair of X and Y ” and whose distribution is
 249 $(S_X \times S_Y, \Sigma_X \otimes \Sigma_Y, \nu)$, subject to

$$\begin{aligned} \nu(E_X \times S_Y) &= \mu_X(E_X), \\ \nu(S_X \times E_Y) &= \mu_Y(E_Y), \end{aligned} \quad (4)$$

250 for any events $E_X \in \Sigma_X$ and $E_Y \in \Sigma_Y$. The meaning of
 251 $\Sigma_X \otimes \Sigma_Y$ is the smallest sigma-algebra on $S_X \times S_Y$ that
 252 contains pairwise products of events in Σ_X and Σ_Y .

253 If such a $Z = (X, Y)$ exists (is defined among the ran-
 254 dom variables one considers), then the joint distribution of
 255 X and Y is unique. The existence of this random variable,
 256 however, is not established by a mathematical derivation
 257 from the properties of X and Y , it is determined by the ex-
 258 istence of an empirical procedure in which the realizations
 259 of X and Y are observed “together.” If $Z = (X, Y)$ does
 260 not exist, one can always construct a coupling $\tilde{Z} = (\tilde{X}, \tilde{Y})$
 261 whose distribution is $(S_X \times S_Y, \Sigma_X \otimes \Sigma_Y, \nu)$, subject to
 262 (4). The only difference (but a critical one) is that the
 263 name of this $\tilde{Z}$ is not “ordered pair of X and Y ” but “or-
 264 dered pair of $\tilde{X}$ [whose distribution is the same as that of

¹ I could have said “distribution is determined by (S_X, Σ_X, μ_X) ,” but it is simpler to say “distribution is (S_X, Σ_X, μ_X) ,” as we do not have an independent general definition of a distribution.

265 $X]$ and $\tilde{Y}$ [whose distribution is the same as that of Y].”
 266 Such a $\tilde{Z}$ can be freely introduced and does not change the
 267 X and Y being coupled; in fact, $(\tilde{X}, \tilde{Y})$ is stochastically
 268 unrelated to X and to Y .

269 All of this can be easily generalized to an arbitrary set
 270 of random variables (see, e.g., Dzhaferov and Kujala, in
 271 press).

272 II.3. Random variables and joint distributions in 273 KPT

274 The formal account just given is not, however, that of
 275 the traditional KPT. The latter begins with the notion of a
 276 *sample space*, (S, Σ, μ) ,² and defines a random variables X
 277 as a measurable mapping of this space into a measurable
 278 space (S_X, Σ_X) , i.e., a function $X : S \rightarrow S'$ such that
 279 $X^{-1}(E_X) \in \Sigma$ for any $E_X \in \Sigma_X$.³ The mapping induces
 280 on the codomain space (S_X, Σ_X) a probability measure μ_X ,
 281 by the rule

$$\Pr[X \in E_X] = \mu_X(E_X) = \mu(X^{-1}(E_X)), \quad (5)$$

282 for any $E_X \in \Sigma_X$. The resulting triple (S_X, Σ_X, μ_X)
 283 is called the *distribution of X* . If another measurable
 284 mapping Y is defined on the same sample space, map-
 285 ping it into a codomain space (S_Y, Σ_Y) and resulting in
 286 the distribution (S_Y, Σ_Y, μ_Y) , then their joint distribution
 287 $(S_X \times S_Y, \Sigma_X \otimes \Sigma_Y, \nu)$ is derived from the relation

$$\Pr[X \in E_X \& Y \in E_Y] = \mu(X^{-1}(E_X) \cap Y^{-1}(E_Y)), \quad (6)$$

288 for any $E_X \in \Sigma_X$ and $E_Y \in \Sigma_Y$. Note that unlike in the
 289 “naive” approach above, the joint distribution of X and Y
 290 here is mathematically derived from their individual defi-
 291 nitions as measurable functions on the same sample space.
 292 Clearly, any two random variables defined on the same
 293 sample space are jointly distributed. This may create a
 294 temptation to assume the existence of a common sample
 295 space and a joint distribution (even if unknown to us) for
 296 *any two* random variables. In turn, this would mean the
 297 existence of a common sample space for *all possible* ran-
 298 dom variables, so that random variables in any set under
 299 consideration possess a joint distribution, and this distribu-
 300 tion is unique. Kolmogorov’s (1933/1956) book may seem

301 to reinforce this view, as it does not explicitly speak of
 302 multiple sample spaces. I disagree with this interpretation,
 303 even if not uncommon (see, e.g., the overview of interpreta-
 304 tions in Khrennikov, 2009c). Kolmogorov’s monograph ties
 305 the notion of a sample space⁴ to “a complex of conditions
 306 which allows of any number of repetitions” (Kolmogorov,
 307 1933/1956, §2 of Chapter 1): this can be interpreted as a
 308 position very close to if not the same as the one argued for
 309 below. Whatever the correct interpretation, however, the
 310 notion of a single sample space for all random variables is
 311 untenable as it contradicts the common mathematical prac-
 312 tices in dealing with random variables. In Dzhaferov and
 313 Kujala (2014a) we presented the following two arguments
 314 demonstrating this.

315 First of all, for any choice of a universal sample space
 316 (S, Σ, μ) all random variables X defined on it will have the
 317 cardinality of their set of possible values, defined as $S_X =$
 318 $X(S)$, less than or equal to the cardinality of S . There
 319 is, however, no justification, empirical or mathematical, for
 320 limiting the cardinality of the set S_X of all possible values
 321 for a random variable X .⁵

322 Second, even if we confine our attention to very simple
 323 random variables with one and the same distribution, there
 324 is no justification, empirical or mathematical, for limiting
 325 this set in any way. One can always add a new random
 326 variable to any given set thereof. Thus, given any set $\mathcal{N}$
 327 of unit-normally distributed random variables, one can in-
 328 troduce a unit-normally distributed $Y_{\mathcal{N}}$ such that its cor-
 329 relation with any $X \in \mathcal{N}$ is zero. If there were a universal
 330 sample space (S, Σ, μ) , then there would be a definite set
 331 $\mathcal{N}^*$ of all possible unit-normally distributed random vari-
 332 ables. But this would mean that our $Y_{\mathcal{N}^*}$ would have to
 333 belong to this set, which is impossible, as $Y_{\mathcal{N}^*}$ cannot have
 334 zero correlation with itself.

335 To further appreciate the untenability of a universal sam-
 336 ple space, observe that the identity mapping from this space
 337 into itself is a random variable, R . The idea of a univer-
 338 sal sample space therefore is equivalent to the existence
 339 of a random variable R of which all imaginable random
 340 variables are functions. This does not add a new formal
 341 argument against the idea, but it seems especially demon-
 342 strative: what this mysterious “super-variable” R could be?

343 II.4. A reinterpreted (or revised?) KPT

344 All these considerations lead us to a different picture of
 345 the KPT, in which there are different, stochastically un-
 346 related random variables $R, R', R'', \dots$, corresponding to
 347 different, mutually exclusive conditions under which they

² Note the terminological variance: “sample space” is more often than not used in the literature to designate just the set S rather than the entire domain probability space (S, Σ, μ) . I find it more in line with the general meaning of the terms “set” and “space” in mathematics to refer to S as a sample set (or set of possible outcomes). This set is promoted into a space by endowing it with a structure, which in this case is provided by the sigma algebra Σ and the measure μ .

³ Kolmogorov (1933/1956) only considered the case when S_X is a subset of reals and Σ_X is the Borel sigma-algebra restricted to this subset. In this paper I use the term “random variable” in the broad sense, with no restrictions imposed on (S_X, Σ_X) . Some authors prefer to use the term “random element” or “random entity” to designate random variables in the broad sense.

⁴ Kolmogorov’s terminology is not the same as the modern terminology (or variant thereof) I use in this paper; in particular, he does not speak of a sample space but of a “basic set with an algebra of subsets.”

⁵ Kolmogorov (1933/1956) did not have to deal with this issue, as the random variables in this book are confined to real-valued ones.

are observed;⁶ and for each of these random variables one can define various functions of it,

$$\begin{aligned} X &= f(R), Y = g(R), \dots \\ X' &= f'(R'), Y' = g'(R'), \dots \\ X'' &= f''(R''), Y'' = g''(R''), \dots \end{aligned} \quad (7)$$

so that any two random variables that are functions of one and the same member of the set $R, R', R'', \dots$ possess a joint distribution, while any functions of two different members of this set do not. To preserve Kolmogorov's definition of a random variable, the $R, R', R'', \dots$ can be thought of as identity functions on their separate sample spaces.

This picture is a step in the right direction, but it is still flawed if we think of $R, R', R'', \dots$ as some fixed set comprising all pairwise stochastically unrelated random variables. The reason for this is that if $R, R', R'', \dots$ were a fixed set, with the corresponding sample spaces (which, since they are identity functions, are simultaneously their distributions)

$$(S_R, \Sigma_R, \mu_R), (S_{R'}, \Sigma_{R'}, \mu_{R'}), (S_{R''}, \Sigma_{R''}, \mu_{R''}), \dots \quad (8)$$

then one could form a single random variables R^* of which $R, R', R'', \dots$ (hence also all other random variables imaginable) were functions. This “super-variable” R^* would have the set and sigma-algebra that are products of, respectively, sets and sigma-algebras in (8), and it would have a probability measure ν from which $\mu_R, \mu_{R'}, \mu_{R''}, \dots$ are computed as marginals, e.g.,

$$\mu_R(E_R) = \nu(E_R \times S_{R'} \times S_{R''} \times \dots),$$

for any $E_R \in \Sigma_R$. We have seen already that the idea of such a “super-variable” is untenable.

A logically consistent way out of this difficulty is to consider $R, R', R'', \dots$ as a *class with uncertain and/or flexible membership*.⁷ Indeed, it should be clear from the previous discussion that random variables can be freely introduced, so, e.g., there is no fixed set of random variables with any given distribution. Some random variables we observe have an empirically defined coupling scheme, and then they are jointly distributed. Other sets of random variables we observe are observed under different conditions each and do not have an empirical coupling. Then they can be modeled as stochastically unrelated random variables. However, we then can create “copies” of these random variables and couple them “on paper” in a multitude of ways. This seems to be a consistent view of random variables. KPT is by no means dismissed in this view, because any distribution (S_X, Σ_X, μ_X) for a random variable X is a probability space subject to Kolmogorov's axioms:

1. μ_X is a function $\Sigma_X \rightarrow [0, \infty)$;

$$2. \mu_X(S_X) = 1;$$

3. $\mu_X\left(\bigcup_{i=1}^{\infty} E_X^{(i)}\right) = \sum_{i=1}^{\infty} \mu_X\left(E_X^{(i)}\right)$ for any sequence of pairwise disjoint $E_X^{(1)}, E_X^{(2)}, \dots$ in Σ_X .

Moreover, insofar as one focuses on a given set of jointly distributed random variables, all of them can be presented as measurable functions on a single sample space (or functions of a single random variable).

II.5. Radical contextualism

Is there a unique way of determining which random variables are and which are not stochastically interrelated? A general answer to this question is negative: the definition of a jointly distributed set of random variables involves an empirical procedure of coupling their realizations, “observing them together.” The meaning of such an empirical procedure may be different for different situation and different observers. From the mathematical point of view, however, the question is about a language that makes the fundamental distinctions between stochastically related and stochastically unrelated random variables. Such a language is proposed in Dzhamalov and Kujala (2014b), Dzhamalov, Kujala, and Larsson (2015), and Kujala, Dzhamalov, and Larsson (2015). For an overview, see Dzhamalov and Kujala (2015a) and Dzhamalov, Kujala, and Cervantes (in press).

It is postulated that every random variable's identity is determined by two types of variables, referred to as *objects* (also properties, entities, contents, etc.) and *contexts* (also conditions, environment, etc.). Intuitively, the random variables are treated as “measurements,” the objects answer the question “what is measured?” whereas the contexts answer the question “how is it measured?”

Let Q be a set of objects and C a set of contexts considered in a given study. The mentioning of “a given study” is essential: in a different study one could choose a different set of objects to measure and a different set of contexts in which to measure them. The measurement R of an object $q \in Q$ in a context $c \in C$ is denoted by R_q^c .

The meaning of a context is that it provides an empirical coupling for the measurements within this context: the random variables R_q^c with different q measured within the same context c are “measured together,” i.e., they possess a joint distribution. Denoting by Q_c the subset of objects in Q that are measured in context c ,

$$R^c = \{R_q^c\}_{q \in Q_c} \quad (9)$$

is a random variable (which implies that it has a distribution, and this distribution is a joint distribution of its components). On the other hand, any two random variables R_q^c and $R_{q'}^{c'}$ with $c \neq c'$ are stochastically unrelated, whether q and q' are distinct objects or not. It follows that any random variables R^c and $R^{c'}$ defined as in (9) with $c \neq c'$ are stochastically unrelated.

The idea of such contextual notation and (mutatis mutandis) understanding of stochastic unrelatedness have precursors and analogues in the quantum-physical literature:

⁶ The notation $R, R', R'', \dots$ is informal and should not be interpreted as indicating a countable set.

⁷ A different approach is presented in Dzhamalov and Kujala (2015b), where we formally define, by means of a quasi-constructive procedure, the set of all random variables considered “existing” in a given study.

see Khrennikov (2005, 2008, 2009a-c), Simon, Brukner, and Zeilinger (2001), Larsson (2002), Svozil (2012), and Winter (2014). Khrennikov (2009c) points out that the contextual understanding of random variables is intrinsic features of von Mises’s “ensemble approach” to probabilities: the identity of an “ensemble” of observations corresponds to context in which these observations are made.⁸

Three aspects of our theory, however, set it aside from this literature:

1. Contextual labeling is universal, and no two random variables recorded in different contexts have a joint distribution.
2. Pairwise stochastically unrelated random variables $\{R^c\}_{c \in C}$ (each of which is a set of jointly distributed random variables) can be coupled at will, with no coupling being privileged.
3. The random variables $\{R^c\}_{c \in C}$ can be characterized by whether it is possible or impossible to couple them in a particular way (e.g., by a maximally connected coupling, as discussed in Section II.7).

Below I will give an example of how these principles work in solving the problem of selective influences in psychology, as well as its generalized version, the problem of contextuality, primarily studied in quantum mechanics. First, however, I have to address some obvious objections to the radical contextualism.

II.6. Possible objections

The first objection is that it is impossible to take into account all conditions in the world, and without knowing them one would not know if one deals with stochastically related or unrelated random variables. The response to this objection lies in the qualification “in a given study” I made when I introduced object sets Q and context sets C . The identification of random variables by what they measure and by how they measure it depends on what other variables in the world one records and relates to realizations of the random variables in question.

To give an example, let there be a very large group of husband-and-wife couples; to each of the husbands Alice poses one of two different Yes/No questions, a_1 or a_2 ; to each of the wives Bob poses one of two different Yes/No questions, b_1 or b_2 (that may be the same as or different from a_1, a_2). Alice decides (this is not a matter of truth or falsity but one of convention) to consider the responses to a_1 and a_2 in the group of husbands as realizations of random variables R_{a_1} and R_{a_2} , respectively; and Bob defines R_{b_1} and R_{b_2} for the group of wives analogously. This labeling indicates that Alice treats a_1 and a_2 as objects being

measured (by responses to these questions), and so does Bob for b_1 and b_2 .⁹

Let us ask now: what are the contexts in which Alice records her R_{a_1} and R_{a_2} ? By the rules of the survey each person answers a single question, so asking a_1 excludes asking a_2 and vice versa. In other words, the conditions under which one records answers to a_1 and a_2 are incompatible. Formally, this means that a_1 is measured in the context a_1 while a_2 is measured in the context a_2 : Alice has therefore stochastically unrelated random variables $R_{a_1}^{a_1}$ and $R_{a_2}^{a_2}$. Analogously, Bob has stochastically unrelated $R_{b_1}^{b_1}$ and $R_{b_2}^{b_2}$. Their stochastic unrelatedness is quite obvious: why should any response by a person to a_1 (or b_1) be paired with any response by another person to a_2 (respectively, b_2)?

By the same argument, either of Alice’s measurements is stochastically unrelated to either of Bob’s: the four measurements

$$R_{a_1}^{a_1}, R_{a_2}^{a_2}, R_{b_1}^{b_1}, R_{b_2}^{b_2} \quad (10)$$

are made in four different contexts. It is clear, however, that Alice and Bob could try to form joint distributions of their measurements using some empirical coupling procedure, e.g., the pairing of the measurements by the marital relation: that is, pairing a husband’s response to a_i with his wife’s response to b_j , for each of the four combinations of $i = 1, 2$ and $j = 1, 2$. To do this means to form new contexts, (a_1, b_1) , (a_1, b_2) , (a_2, b_1) , and (a_2, b_2) , and to re-label Alice’s random variables as

$$R_{a_1}^{(a_1, b_1)}, R_{a_1}^{(a_1, b_2)}, R_{a_2}^{(a_2, b_1)}, R_{a_2}^{(a_2, b_2)} \quad (11)$$

while Bob’s random variables become

$$R_{b_1}^{(a_1, b_1)}, R_{b_1}^{(a_2, b_1)}, R_{b_2}^{(a_1, b_2)}, R_{b_2}^{(a_2, b_2)}. \quad (12)$$

The previous four pairwise stochastically unrelated variables (10) are replaced now with the four pairwise stochastically unrelated variables

$$R^{(a_i, b_j)} = \left(R_{a_i}^{(a_i, b_j)}, R_{b_j}^{(a_i, b_j)} \right), \quad i, j \in \{1, 2\}. \quad (13)$$

There is no justification for saying that either of these representations, (10) or (13), is more “correct” than another: Alice who does not know whose husband her respondent is and Alice who knows this deal with different sets of random variables.

Another, related objection is that radical contextualism should lead to considering every realization of a random variable as being stochastically unrelated to every other realization. In the previous example, if Alice records the identities of the people she is posing the question a_1 to, then in place of a single $R_{a_1}^{a_1}$ she creates random variables $R_{a_1}^{John}$, $R_{a_1}^{Peter}$, etc., each with a single realization. In a typical behavioral experiment, John, Peter, etc. can be

⁸ Khrennikov thinks that in this respect von Mises’s approach is radically different from Kolmogorov’s, an opinion one can disagree with if the KPT is not confined to a single probability space.

⁹ This example is formally equivalent to the EPR/Bohm experiment in quantum physics (see Section II.7).

replaced with “trial 1,” “trial 2,” etc. As the contexts (upper indexes) differ, the variables are stochastically unrelated. Isn’t this a problem? In particular, does not stochastic unrelatedness of the realizations of a random variable clash with the standard statistical practice of viewing them as independent identically distributed variables?

The answer to these questions is essentially the same as to the previous objection. Alice does not have to record the identity of the people she queries, and if she does not, then $R_{a_1}^{a_1}$ is the random variable she forms. If she does, then she creates new contexts, and it is indeed true then that $R_{a_1}^{John}$, $R_{a_1}^{Peter}$, etc., are pairwise stochastically unrelated. They can, however, be coupled, and as always when coupling is not based on an empirical procedure, this can be done in a multitude of ways. One possible coupling is the independent coupling, the other is the identity coupling, and one could create an infinity of other couplings.

This may be difficult to understand. Suppose we know that John responded Yes, Peter responded No, Paul responded No, etc. — then how could one speak of the identity coupling? Or, if we know that the response in trial $n + 1$ repeats the response in trial n with probability 0.7 — how can one speak then of the independent coupling? To answer these questions one should recall that by coupling pairwise stochastically unrelated $R^{(1)}, R^{(2)}, R^{(3)}, \dots$ one does not mysteriously transform them into jointly distributed random variable. Instead one creates a new sequence $(\tilde{R}^{(1)}, \tilde{R}^{(2)}, \tilde{R}^{(3)}, \dots)$, in which each $\tilde{R}^{(i)}$ has the same distribution as $R^{(i)}$, and all these components have a joint distribution. The table below demonstrates the logical structure of the identity coupling:

$$\begin{array}{cccc} & R^{(1)} & R^{(2)} & R^{(3)} & \dots \\ \tilde{R}^{(1)} & \boxed{r_1} & r_2 & r_3 & \dots \\ \tilde{R}^{(2)} & r_1 & \boxed{r_2} & r_3 & \dots \\ \tilde{R}^{(3)} & r_1 & r_2 & \boxed{r_3} & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{array} \quad (14)$$

The boxed values are the ones factually observed, the rest of the values in each column are those attained by the corresponding components of the identity coupling. As we see, this has nothing to do with the observed values being or not being equal to each other.

The next table demonstrates the logical structure of the independent coupling:

$$\begin{array}{cccc} & R^{(1)} & R^{(2)} & R^{(3)} & \dots \\ \tilde{R}^{(1)} & \boxed{r_1} & r'_2 & r'_3 & \dots \\ \tilde{R}^{(2)} & r'_1 & \boxed{r_2} & r'_3 & \dots \\ \tilde{R}^{(3)} & r''_1 & r''_2 & \boxed{r_3} & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{array} \quad (15)$$

The boxed values, again, are those factually observed, and the primed values in the i th column are sampled from a coupling $(\tilde{R}^{(1)}, \tilde{R}^{(2)}, \tilde{R}^{(3)}, \dots)$ with stochastically independent

components and $\tilde{R}^{(i)} = r_i$. Again, this has nothing to do with the observed values forming or not forming a sequence with certain statistical properties.

Focusing on the statistical properties of the observed (“boxed”) values means, formally, that the observations in different trials (or responses from different persons) are treated as objects rather than contexts, all these objects being measured in a single context and therefore jointly distributed:

$$R_1^{(1,2,3,\dots)}, R_2^{(1,2,3,\dots)}, R_3^{(1,2,3,\dots)}, \dots \quad (16)$$

A third objection one can raise against the radically contextual reinterpretation (or revision) of the KPT is that the notions of an “object” and a “context” are not mathematically defined: they are primitives of the language proposed. How can one know what objects and what contexts to invoke in a specific situation? The response to this is that it is indeed not a mathematical issue. Mathematical analysis begins once one has specified a set Q of objects and a set C of contexts, and there is no single correct way of doing it.

Consider, e.g., the situation when two questions are asked in one of two orders, $a \rightarrow b$ or $b \rightarrow a$. One can take a to be the same object measured in two different contexts, and similarly for b , forming thereby four random variables (responses to the questions)

$$R_a^{a \rightarrow b}, R_b^{a \rightarrow b}, R_a^{b \rightarrow a}, R_b^{b \rightarrow a}. \quad (17)$$

By our rules, they are grouped into two stochastically unrelated random variables

$$R^{a \rightarrow b} = (R_a^{a \rightarrow b}, R_b^{a \rightarrow b}) \text{ and } R^{b \rightarrow a} = (R_a^{b \rightarrow a}, R_b^{b \rightarrow a}). \quad (18)$$

This view of the situation leads to an interesting contextual analysis (Dzhafarov, Zhang, & Kujala, 2015).

It is, however, possible to deny that “the same question a ” means “the same object a ” in the two contexts: one can maintain instead that a asked first is simply a different object from a asked second; and similarly for b . In this view we have four objects, a_1, a_2, b_1, b_2 (where index indicates whether the question is asked first or second), measured in two contexts, $a_1 \rightarrow b_2$ and $b_1 \rightarrow a_2$. One ends up with two stochastically unrelated random variables

$$\begin{aligned} R^{a_1 \rightarrow b_2} &= (R_{a_1}^{a_1 \rightarrow b_2}, R_{b_2}^{a_1 \rightarrow b_2}) \\ &\text{and} \\ R^{b_1 \rightarrow a_2} &= (R_{a_2}^{b_1 \rightarrow a_2}, R_{b_1}^{b_1 \rightarrow a_2}). \end{aligned} \quad (19)$$

This representation allows for no nontrivial contextual analysis (see below), as the stochastically unrelated random variables have no objects in common. It is, nevertheless, as legitimate a representation as the previous one. A psychologist will most probably choose (18) over (19) (Wang & Bussemeyer, 2013; Wang et al., 2014), but it is not mathematics that dictates this choice.

II.7. An example of contextual analysis

The problem of selective influences was introduced to psychology by Sternberg (1969) and developed through a series of publications (Schweickert & Townsend, 1989; Townsend, 1984, 1990; Townsend & Schweickert, 1989; Roberts & Sternberg, 1993; Townsend & Nozawa, 1995; Schweickert, Giorgini, & Dzhamfarov, 2000; Dzhamfarov 2003; Dzhamfarov, Schweickert, & Sung, 2005; Kujala & Dzhamfarov, 2008; Dzhamfarov & Kujala, 2010). Later, a link has been established between this problem and the quantum-mechanical analysis of entanglement (Dzhamfarov & Kujala, 2012a-b, 2013, 2014c) and, more generally, probabilistic contextuality (Dzhamfarov & Kujala, 2014a-b, 2015a-b; Dzhamfarov, Kujala, & Larsson, 2015; Kujala, Dzhamfarov, & Larsson, 2015).

I will formulate the problem using the contextual language introduced above. Let there be a system acted upon by two inputs, α and β , and reacting by two simultaneous distinct responses, R_α and R_β (or distinct aspects of the same response, such as response time and response accuracy). The indexation here reflects the belief (or hypothesis) that R_α is “primarily” influenced by α and R_β by β . One can also say that R_α measures α and R_β measures β . The question is whether R_α is also influenced by β and/or R_β is also influenced by α . Let us simplify the problem by assuming that $\alpha \in \{1, 2\}$ and $\beta \in \{1, 2\}$, and they vary in a completely crossed factorial design, $\{1, 2\} \times \{1, 2\}$. Each of the treatments $(\alpha, \beta) = (i, j)$ should be considered a context, wherefrom the responses of the system must be labeled

$$R^{(\alpha=i, \beta=j)} = (R_{\alpha=i}^{(\alpha=i, \beta=j)}, R_{\beta=j}^{(\alpha=i, \beta=j)}), \quad i, j \in \{1, 2\}. \quad (20)$$

To remind the interpretation, $R_{\alpha=i}^{(\alpha=i, \beta=j)}$ measures the object $\alpha = i$ in the context $(\alpha = i, \beta = j)$; $R_{\beta=j}^{(\alpha=i, \beta=j)}$ measures the object $\beta = j$ in the same context; being in the same context, these two measurements form a random variable $R^{(\alpha=i, \beta=j)}$ (whose components possess a joint distribution); however, the four random variables $R^{(\alpha=i, \beta=j)}$ are pairwise stochastically unrelated. To lighten the notation, let us put

$$R_{\alpha=i}^{(\alpha=i, \beta=j)} = A_i^{ij}, \quad R_{\beta=j}^{(\alpha=i, \beta=j)} = B_j^{ij}. \quad (21)$$

According to the definition of selective influences given in Dzhamfarov (2003) and elaborated in Dzhamfarov and Kujala (2010), one says that A_i^{ij} is not influenced by β and B_j^{ij} is not influenced by α (for all i, j) if one can find a coupling

$$(\tilde{A}_1^{11}, \tilde{B}_1^{11}, \tilde{A}_1^{12}, \tilde{B}_2^{12}, \tilde{A}_2^{21}, \tilde{B}_1^{21}, \tilde{A}_2^{22}, \tilde{B}_2^{22}) \quad (22)$$

in which the equalities

$$\begin{aligned} \tilde{A}_1^{11} &= \tilde{A}_1^{12}, & \tilde{A}_2^{21} &= \tilde{A}_2^{22}, \\ \tilde{B}_1^{11} &= \tilde{B}_1^{21}, & \tilde{B}_2^{12} &= \tilde{B}_2^{22} \end{aligned} \quad (23)$$

hold with probability 1. Put differently, the random variables $\tilde{A}_1^{11}$ and $\tilde{A}_1^{12}$ in the joint distribution of (22) always

attain one and the same value, even though the value of β changes; and analogously for the remaining three equalities.

Note that, by the definition of a coupling,

$$\tilde{A}_i^{ij} \stackrel{d}{=} A_i^{ij} \quad \text{and} \quad \tilde{B}_j^{ij} \stackrel{d}{=} B_j^{ij}, \quad i, j \in \{1, 2\}, \quad (24)$$

where $\stackrel{d}{=}$ means “has the same distribution as.”

If all A and B responses of the system have a finite number of possible values, this situation generalizes Bohm’s version of the Einstein-Podolsky-Rosen (EPR) paradigm (Bohm & Aharonov, 1957; Bell, 1964). Of course, the distributions of the A, B need not be generally in compliance with the quantum rules for entangled particles, but the existence or nonexistence of a coupling with the stipulated properties should be determinable for any observed A and B . Let us assume for simplicity that both A and B responses of the system are binary, and let us denote their values $+1$ and -1 . In this special case the necessary and sufficient conditions for the selectiveness of influences are given by

$$\begin{aligned} A_i^{i1} &\stackrel{d}{=} A_i^{i2} \quad \text{for } i = 1, 2 \\ B_j^{1j} &\stackrel{d}{=} B_j^{2j} \quad \text{for } j = 1, 2 \end{aligned} \quad (25)$$

and

$$\max_{k, l \in \{1, 2\}} \left| \sum_{i, j \in \{1, 2\}} E[A_i^{ij} B_j^{ij}] - 2E[A_k^{kl} B_l^{kl}] \right| \leq 2, \quad (26)$$

where E stands for expected value. This (in an algebraically different form) was first proved by Fine (1982), although (25) in his work is implied by the notation rather than stated explicitly. The distributional equalities (25) describe the condition known as *marginal selectivity*: the distribution of A_i^{ij} does not change with the value j of β , and the distribution of B_j^{ij} does not change with the value i of α . The numerical inequality (26) is known as the *CHSH inequality* (after the authors of Clauser et al., 1969). In quantum mechanics, violations of this inequality when the marginal selectivity (25) holds is described by saying that the system is *contextual* (see, e.g., Kurzynski, Ramanathan, & Kaszlikowski, 2012).

If the marginal selectivity (25) is violated, the CHSH inequality (26) cannot be derived, and it makes no difference whether it is satisfied or not. Moreover, if marginal selectivity is violated, it seems unnecessary to look at anything else: clearly then A is not selectively influenced by α alone, and/or B is not selectively influenced by β alone. As it turns out, however, one may still be interested in the question: is the influence of β upon A and/or of α upon B entirely described by the violations of marginal selectivity? Indeed, since the CHSH inequality (26) may very well be violated when the marginal selectivity (25) holds, and since we then conclude that selectiveness of influences is violated too, we have to admit that the “wrong” influences (from β to A and/or from α to B) can be indirect, without manifesting themselves in changed marginal distributions. This leads

us to a generalized notion of contextuality (Dzhafarov, Kujala, & Larsson, 2015; Kujala, Dzhafarov, & Larsson, 2015; Dzhafarov & Kujala, 2015a-b; Dzhafarov, Zhang, Kujala, 2015; Dzhafarov, Kujala, & Cervantes, in press).

When applied to our example with two binary inputs α, β and two binary random outputs A, B , the definition is as follows. A system

$$(A_1^{11}, B_1^{11}), (A_1^{12}, B_2^{12}), (A_2^{21}, B_1^{21}), (A_2^{22}, B_2^{22})$$

is noncontextual if it has a *maximally connected coupling*. The latter is defined as a coupling (22) in which each of the equalities (23) holds with the maximal possible probability that is allowed by the individual distributions of the random variables. To explain, if $A_1^{11} \stackrel{d}{=} A_1^{12}$, then the maximal possible value for $\Pr[\tilde{A}_1^{11} = \tilde{A}_1^{12}]$ is 1. Applying this to all other equalities in (23), we get the previous definition. If, however, A_1^{11} and A_1^{12} have different distributions, then the maximal possible value for $\Pr[\tilde{A}_1^{11} = \tilde{A}_1^{12}]$ is

$$\begin{aligned} & \min \{ \Pr[A_1^{11} = 1], \Pr[A_1^{12} = 1] \} \\ & + \min \{ \Pr[A_1^{11} = -1], \Pr[A_1^{12} = -1] \} \\ & = 1 - |\Pr[A_1^{11} = 1] - \Pr[A_1^{12} = 1]|. \end{aligned} \quad (27)$$

If some coupling (22) has this and the analogously computed maximal values for other equalities in (23), then the system is noncontextual: the “wrong” influences in it are all confined to directly changing the distributions of the “wrong” random variables. If no such coupling exists, however, the system is contextual: the influence of β upon A and/or α upon B is greater than just distributional changes. As shown in Dzhafarov, Kujala, and Larsson (2015), Kujala, Dzhafarov, and Larsson (2015), and Kujala and Dzhafarov (in press), the necessary and sufficient condition for noncontextuality in accordance with this definition is

$$\begin{aligned} & \max_{k,l \in \{1,2\}} \left| \sum_{i,j \in \{1,2\}} \mathbb{E}[A_i^{ij} B_j^{ij}] - 2\mathbb{E}[A_k^{kl} B_l^{kl}] \right| \\ & \leq 2 + \sum_{i=1}^2 |\mathbb{E}[A_i^{i1}] - \mathbb{E}[A_i^{i2}]| + \sum_{j=1}^2 |\mathbb{E}[B_j^{1j}] - \mathbb{E}[B_j^{2j}]| \end{aligned} \quad (28)$$

For application of this and other criteria of contextuality to available experimental data in physics and psychology see, respectively, Kujala, Dzhafarov, and Larsson (2015) and Dzhafarov, Zhang, and Kujala (2015).

III. CONCLUSION

I have argued in this paper that the KPT (Kolmogorovian probability theory) must allow for stochastically unrelated random variables, and these must not be confused with stochastically independent ones. I have argued for radical contextualism: any two random variables recorded under different conditions (in different contexts) are stochastically unrelated. There is no fixed set of pairwise stochasti-

cally unrelated random variables: they can be freely introduced and freely coupled. To couple a given set of stochastically unrelated random variables means to create their jointly distributed “copies” (stochastically unrelated to the “originals”). The couplings for a given set of random variables are typically infinite in number, with no coupling being “more correct” than another. This applies also to couplings with stochastically independent components. The idea I and Janne Kujala have been promoting in recent publications is that stochastically unrelated random variables can be usefully characterized by their possible couplings, in particular, by determining whether these variables allow for couplings subject to certain constraints. I have illustrated this idea on the issue of selective influences, generalized into the issue of probabilistic contextuality.

Acknowledgments.

This research has been supported by NSF grant SES-1155956 and AFOSR grant FA9550-14-1-0318, and A. von Humboldt Foundation. I greatly benefited from discussions with Matt Jones of the University of Colorado, my doctoral students Ru Zhang and Victor H. Cervantes, and, of course, my long-term collaborator (and co-author of the theory presented in this paper) Janne Kujala. Finally, I should acknowledge my indebtedness to R. Duncan Luce: my conversations and debates with him have served as a major source of intellectual inspiration for me for many years.

REFERENCES

- Bell, J. (1964). On the Einstein-Podolsky-Rosen paradox. *Physics* 1, 195-200.
- Bohm, D., & Aharonov, Y. (1957). Discussion of experimental proof for the paradox of Einstein, Rosen and Podolski. *Physical Review* 108, 1070-1076.
- Clauser, J.F., Horne, M.A., Shimony, A., & Holt, R.A. (1969). Proposed experiment to test local hidden-variable theories. *Physical Review Letters* 23, 880-884.
- Dzhafarov, E.N. (2003). Selective influence through conditional independence. *Psychometrika*, 68, 7-26.,
- Dzhafarov, E.N., & Kujala, J.V. (2010). The Joint Distribution Criterion and the Distance Tests for selective probabilistic causality. *Frontiers in Psychology* 1:151, doi: 10.3389/fpsyg.2010.00151.
- Dzhafarov, E.N. & Kujala, J.V. (2012a). Selectivity in probabilistic causality: Where psychology runs into quantum physics. *Journal of Mathematical Psychology* 56, 54-63.
- Dzhafarov, E.N., & Kujala, J.V. (2012b). Quantum entanglement and the issue of selective influences in psychology: An overview. *Lecture Notes in Computer Science* 7620, 184-195.
- Dzhafarov, E.N., & Kujala, J.V. (2013). Order-distance and other metric-like functions on jointly distributed

- random variables. *Proceedings of the American Mathematical Society* 141, 3291-3301.
- Dzhafarov, E.N., & Kujala, J.V. (2014a). A qualified Kolmogorovian account of probabilistic contextuality. *Lecture Notes in Computer Science* 8369, 201-212.
- Dzhafarov, E.N., & Kujala, J.V. (2014b). Contextuality is about identity of random variables. *Physica Scripta* T163, 014009.
- Dzhafarov, E.N., & Kujala, J.V. (2014c). On selective influences, marginal selectivity, and Bell/CHSH inequalities. *Topics in Cognitive Science* 6, 121-128.
- Dzhafarov, E.N., & Kujala, J.V. (2015a). Conversations on contextuality. In E.N. Dzhafarov, J.S. Jordan, R. Zhang, V.H. Cervantes (Eds). *Contextuality from Quantum Physics to Psychology*, pp. 1-22. New Jersey: World Scientific Press.
- Dzhafarov, E.N., & Kujala, J.V. (2015b). Context-content systems of random variables: The contextuality-by-default theory. arXiv:1511.03516.
- Dzhafarov, E.N., & Kujala, J.V. (in press). Probability, random variables, and selectivity. In W. Batchelder, H. Colonius, E.N. Dzhafarov, J. Myung (Eds). *New Handbook of Mathematical Psychology*. Cambridge: Cambridge University Press.
- Dzhafarov, E.N., Kujala, J.V., & Cervantes, V.H. (in press). Contextuality-by-Default: A brief overview of ideas, concepts, and terminology. *Lecture Notes in Computer Science* 9535.
- Dzhafarov, E.N., Kujala, J.V., & Larsson, J.-A. (2015). Contextuality in three types of quantum-mechanical systems. *Foundations of Physics* 7, 762-782.
- Dzhafarov, E.N., Zhang, R., & Kujala, J.V. (2015). Is there contextuality in behavioral and social systems? *Philosophical Transactions of the Royal Society A* 374, 20150099.
- Fine, A. (1982). Hidden variables, joint probability, and the Bell inequalities. *Physical Review Letters* 48 291-295.
- Khrennikov, A. (2005). The principle of complementarity: A contextual probabilistic viewpoint to complementarity, the interference of probabilities, and the incompatibility of variables in quantum mechanics. *Foundations of Physics* 35, 1655-1693.
- Khrennikov, A. (2008). Bell-Boole inequality: Nonlocality or probabilistic incompatibility of random variables? *Entropy* 10, 19-32.
- Khrennikov, A. (2009a). Bell's inequality: Physics meets Probability. *Information Science* 179, 492-504.
- Khrennikov, A. (2009b). *Contextual Approach to Quantum Formalism*. Berlin: Springer.
- Khrennikov, A. (2009c) *Interpretations of Probability*. Berlin: De Gruyter.
- Kolmogoroff, A. N. (1933). *Grundbegriffe der Wahrscheinlichkeitsrech.* Berlin: Springer Verlag. English translation: Kolmogorov, A.N. (1956). *Foundations of the Probability Theory*. New York: Chelsea Publishing Company.
- Krantz, D.H., Luce, R.D, Suppes, P., & Tversky, A. (1971). *Foundations of Measurement*, vol. I: *Additive and Polynomial Representations*. New York, NY: Academic Press.
- Kujala, J.V., & Dzhafarov, E.N. (in press). Proof of a conjecture on contextuality in cyclic systems with binary variables. *Foundations of Physics*.
- Kujala, J.V., & Dzhafarov, E.N. (2015). Probabilistic contextuality in EPR/Bohm-type systems with signaling allowed. In E.N. Dzhafarov, J.S. Jordan, R. Zhang, V.H. Cervantes (Eds). *Contextuality from Quantum Physics to Psychology*, pp. 287-308. New Jersey: World Scientific.
- Kujala, J.V., Dzhafarov, E.N., & Larsson, J.-A. (2015). Necessary and sufficient conditions for maximal non-contextuality in a broad class of quantum mechanical systems. *Physical Review Letters* 115, 150401.
- P. Kurzynski, R. Ramanathan, & D. Kaszlikowski (2012). Entropic test of quantum contextuality. *Physical Review Letters* 109, 020404.
- Larsson, J.-Å. (2002). A Kochen-Specker inequality. *Europhysics Letters*, 58, 799-805.
- Luce, R.D. (1996). The ongoing dialog between empirical science and measurement theory. *Journal of Mathematical Psychology* 40, 78-98.
- Luce, R.D. (2013). Analogs in Luce's global psychophysical theory of Stevens's psychophysical regression effect. *American Journal of Psychology* 126, 47-52
- Simon, C., Brukner, Č., & Zeilinger, A. (2001). Hidden-variable theorems for real experiments. *Physical Review Letters* 86, 4427-4430.
- Svozil, K. (2012). How much contextuality? *Natural Computing* 11, 261-265.
- Thorisson, H. (2000). *Coupling, Stationarity, and Regeneration*. New York: Springer.
- von Mises, R. (1957). *Probability, Statistics and Truth*. London: Macmillan.
- Wang, Z., Busemeyer, J.R. (2013). A quantum question order model supported by empirical tests of an a priori and precise prediction. *Topics in Cognitive Science* 5, 689-710.
- Wang, Z., Solloway, T., Shiffrin, R.M., Busemeyer, J.R. (2014). Context effects produced by question orders reveal quantum nature of human judgments. *Proceedings of the National Academy of Sciences* 111, 9431-9436.
- Winter, A. (2014). What does an experimental test of quantum contextuality prove or disprove? *Journal of Physics A: Mathematical and Theoretical* 47, 42403.