Undifferentiated ES cells

Differentiated ES cells

A Structural Investigation into *Oct4* Regulation by Orphan Nuclear Receptors, Germ Cell Nuclear Factor (GCNF) and Liver Receptor Homolog-1 (LRH-1)

Emily R. Weikum^{a¶}, Micheal L. Tuntland^{a¶}, Michael N. Murphy^a, and Eric A. Ortlund^{a*}

Abstract

Oct4 is a transcription factor required for maintaining pluripotency and self-renewal in stem cells. Prior to differentiation, *Oct4* must be silenced to allow for the development of the three germ layers in the developing embryo. This fine-tuning is controlled by the nuclear receptors, liver receptor homolog-1 and germ cell nuclear factor. Liver receptor homolog-1 is responsible for driving the expression of *Oct4* where germ cell nuclear factor represses its expression upon differentiation. Both receptors bind to a DR0 motif located within the *Oct4* promoter. Here, we present the first structure of mouse germ cell nuclear factor DNA binding domain in complex with the *Oct4* DR0. The overall structure revealed two molecules bound in a head-to-tail fashion on opposite sides of the DNA. Additionally, we solved the structure of the human liver receptor

homolog-1 DNA binding domain bound to the same element. We explore the structural elements that govern *Oct4* recognition by these two nuclear receptors.

Keywords: Nuclear receptors, X-ray crystallography, Germ Cell Nuclear Factor, Liver Receptor Homolog-1, Oct4 Regulation

Introduction

The pluripotency of embryonic stem (ES) cells is maintained by a specific group of factors including leukemia inhibitor factor (LIF), and transcription factors Oct4, Sox2, and Nanog [1]. All of these proteins are critical for maintaining precise gene expression profiles in ES cells. Oct4, a member of the POU-domain family is widely recognized as the gatekeeper, preventing ES cell differentiation by maintaining pluripotent gene expression and inhibiting lineage-determining factors [2]. During this time, *Oct4* expression must be tightly regulated to ensure proper expression levels. Upon exposure to differentiation cues, such as the presence of retinoic acid (RA), Oct4 and the other pluripotency factors are repressed in a temporal and spatial manner to ensure proper development of the three germ layers. Members of the nuclear receptor superfamily of ligand-regulated transcription factors are responsible for ensuring this specific pattern of *Oct4* expression [3, 4].

Nuclear receptors (NR) play key roles in diverse biological processes, including maintaining homeostasis, metabolism, development, and many others [4,5]. These receptors all share the same core domain structure: a N-terminal transactivation domain (NTD), a DNA binding domain (DBD) with two highly conserved zinc fingers, a flexible hinge region, and a ligand binding domain (LBD) that contains an activator function-2 helix that is critical for binding coregulators (**Fig. 2a**). These receptors bind to palindromic DNA sequences as a monomer, homodimer, or heterodimer to regulate transcription from their target genes [6]. A

1
2
3
4 number of these receptors including, liver receptor homolog-1 (LRH-1, *NR5A2*), germ cell
5
6 nuclear factor (GCNF, *NR6A1*), steroidogenic factor (SF-1, *NR5A1*), and retinoic acid receptor
7
8 (RAR, *NR1B1*) are known to regulate of *Oct4* expression by binding to response elements within
9
10 the promoter regions [3, 7, 8, 9].
11
12
13

14 LRH-1 regulates *Oct4* gene expression by binding DNA as a monomer to a 9 nucleotide
15
16 recognition element comprised of a YCA followed by a NR half-site (AGGCCCR) sequence (Y=
17
18 pyrimidine, R= purine) [10]. LRH-1 contains a canonical NR DBD composed of two helical zinc
19
20 fingers, which recognize the NR-halfsite. LRH-1, and all NR5A family members, contain a
21
22 unique *fushi tarazu* factor 1 (Ftz-F1) domain located C-terminal to the DBD that has been shown
23
24 to play a functionally important role as a protein interaction module required for the recruitment
25
26 of other transcription factors [11-14]. This domain was structurally characterized in the LRH-1
27
28 DBD *CYP7A1* complex by X-ray crystallography then by NMR for SF-1 [10, 15], and assumes
29
30 an α -helix fold that packs against body of the zinc-figure domain. Mutations that untether the
31
32 Ftz-F1 helix from the body of the protein dramatically reduce both DNA binding and
33
34 transactivation. LRH-1 makes structurally conserved contacts with its DNA half-site through
35
36 both the canonical DNA-reading helix, located within the zinc finger domain, and the C-terminal
37
38 extension (CTE), located between the zinc-finger domain and Ftz-F1 domain. The CTE-DNA
39
40 interactions are influenced by the Ftz-F1 helix which serves to orient the CTE in the DNA minor
41
42 groove [10].
43
44
45
46
47
48
49

50 In early embryonic development, LRH-1 is highly expressed within the inner cell mass
51
52 and primitive endoderm of the blastocyst, where other NRs are secluded to other cell types [3].
53
54 To maintain pluripotency, LRH-1 binds directly to DR0 sequences within the proximal promoter
55
56 (PP) and proximal enhancer (PE) to drive expression of *Oct4* [16]. This regulation via LRH-1 is
57
58
59
60
61
62
63
64
65

critical, as loss of LRH-1 results in embryonic lethality at day 6.5. LRH-1^{-/-} embryos also exhibit low *Oct4* expression and die before liver development is established. Recently, it has even been shown that LRH-1 can replace Oct4 in the reprogramming of somatic cells into pluripotent stem cells [17]. However, upon signals for differentiation, LRH-1 levels are dramatically reduced while GCNF is recruited to repress *Oct4*.

GCNF is an orphan nuclear receptor that was first identified from mouse heart tissue and shows high expression levels in developing germ cells, oocytes, and spermatogenic cells [18]. GCNF comprises its own unique NR superfamily subclass with a DBD that resembles retinoid X receptor (RXR) but a LBD more closely related to COUP transcription factor 2 (COUP-TF). Additionally, GCNF does not have the typical AF-2 helix within its LBD but instead contains a predicted β -sheet that interacts with the transcriptional corepressors nuclear receptor corepressor-1 (NCoR) and nuclear receptor corepressor-2 (NCoR2/SMRT) [7,19]. GCNF represses *Oct4* expression by binding directly to a DR0 element within the PP region, the same site as LRH-1 [18-21]. This transrepression of *Oct4* is required for cell differentiation as loss of GCNF results in embryonic lethality at day E10.5 from cardiovascular complications and other severe developmental abnormalities [19,22]. DNA binding by GCNF is functionally critical as a deletion of the DBD phenocopies the GCNF^{-/-} mice [23]. GCNF binds the *Oct4* DR0 element in the PP as a homodimer and recruits transcriptional corepressors and DNA methyltransferases in order to ultimately silence *Oct4* expression [24]. This promoter methylation is maintained well beyond GCNF expression in order to maintain *Oct4* silencing [25,26]. This process is depicted in

Figure 1.

Here, we present the crystal structures of the mGCNF DBD-TA and hLRH-1 DBD-FtzF1 bound to the *mOct4* DR0 proximal promoter element. Throughout the rest of this manuscript

mGCNF DBD-TA and hLRH-1 DBD-FtzF1 will be referred to as GCNF DBD and LRH-1 DBD, respectively. This work represents the first crystal structure of GCNF and permitted the visualization of the sequence specific contacts that facilitate recognition of this element for both of these orphan NRs.

Results

GCNF and LRH-1 directly bind the mOct4 DR0

To characterize *in vitro* binding affinity and kinetics of GCNF and LRH-1 we monitored the ability of recombinant GCNF DBD, LRH-1 DBD, and full-length LRH-1 to bind a FAM labeled 16 bp *Oct4* DR0 fragment via fluorescence polarization (**Fig 2**). Intact, GCNF has been shown to bind the *Oct4* DR0 as a homodimer via EMSA [18, 19]. Here, GCNF bound the *Oct4* DR0 with a two-site binding mechanism with the K_d of the high affinity site at 170 nM and 2.2 μ M for the low affinity site (**Fig 2c**). This apparent two-site binding, rather than cooperative binding, may be due to the lack of the LBD which presumably facilitates dimerization [24]. The LRH-1 DBD displayed a one site binding mechanism with an apparent K_d of 60 nM (**Fig. 2b**). The full-length LRH-1 construct also fit a one site binding curve with the 16 bp *Oct4* element with an apparent affinity of 30 nM. The data is summarized in a **Figure 2d**.

Structural analysis of GCNF and LRH-1 – Oct4 Complexes

To determine how these individual NRs recognize the same *Oct4* DR0, we solved crystal structures of each protein-DNA complex. The GCNF-*Oct4* 16 bp DR0 DNA complex crystallized in the P2₁2₁2₁ space group and data were collected to 2.1 Å with 96.2% completeness. The LRH-1-*Oct4* DR0 12 bp DNA complex crystallized in the space group P4₃ and data were collected to 2.2 Å with 99.8% completeness (**Table 1**).

1
2
3
4 The GCNF structure shows two molecules bound to opposite sides of the DNA in a head-
5
6 to-tail fashion (**Fig. 3a**). This orientation on DNA is very similar to the structure of the RXR α -
7
8 DR1 complex (PDB ID: **1BY4**) [27]. The GCNF DBD adopts the canonical NR DBD fold [6],
9
10 and strong electron density allowed for modeling of the T and A box residues within the C-
11
12 terminal extension (CTE) for molecule 2. One molecule, colored purple, positions the DNA
13
14 reading helix into the major groove of the DNA at the first AGGTCA repeat (bases 106-111).
15
16 This molecule makes three base-specific contacts (**Fig. 3b**). The Lys96 side chain makes
17
18 hydrogen bonds to the O6 and N7 positions on guanine 107. Arg101 make additional hydrogen
19
20 bonds to the on guanine 97 at the N7 position on the other side of the DNA as well as to the
21
22 backbone phosphate of a thymine 98. The second molecule, colored deep purple, makes similar
23
24 contacts to the second AGGCTA sequence (bases 112-117) (**Fig. 3b**). Here, Lys96 again makes
25
26 hydrogen bonds to the O6 position of guanine 113. Arg101 makes hydrogen bonds to the N7
27
28 position on adenine 91 instead of a guanine and also makes additional hydrogen bonds the
29
30 backbone of thymine 90. These interactions are supported by excellent electron density and
31
32 additional side chains participating in backbone and water-mediated interactions are highlighted
33
34 in **Figure 3c**.
35
36
37
38
39
40
41
42

43 The GCNF CTE, including the T/A box residues, was observed for molecule 2 (residues
44
45 141-156). There is strong electron density for these residues, which is shown in **Figure 5d**. The
46
47 GCNF CTE dips into the minor groove at the TCAA sequence (bases 109-112) to make
48
49 additional sequence specific contacts. Arg86 makes hydrogen bonds to the O2 positions on both
50
51 thymine 109 and cytosine 110. Arg78 makes hydrogen bonds to the polypeptide backbone
52
53 oxygen at Gly152, which may help lock the CTE into a more stable conformation for interaction
54
55 with the DNA minor groove. In addition to DNA contacts, the CTE also makes a series of
56
57
58
59
60
61
62
63
64
65

1
2
3
4 contacts with GNCF molecule 1 via a set of highly conserved residues (**Fig 4**). The largely
5
6 negative CTE of molecule 2 rests within the minor groove but also makes contacts with the
7
8 positive surface of molecule 1(**Fig 4a**). Namely, Asp148 makes weak electrostatic contacts with
9
10 Arg121. This intermolecular interface is also supported by a series of hydrophobic contacts
11
12 driven largely by the conserved Met150, which interacts with Arg121 and Arg124 (**Fig 4b,c**).
13
14

15
16
17 The LRH-1 DBD crystallized as a monomer on a 12 bp *Oct4* fragment, with its DNA
18
19 recognition helix resting in the major groove formed by the DR0 motif, flanked by two
20
21 conserved zinc fingers (**Fig. 5a**). This structure is similar to the previously solved LRH-1 –
22
23 *hCYP7A1* complex (PDB ID: **2A66**) [10]. The DNA reading helix contains four residues that
24
25 make distinct base-specific contacts with the *Oct4* DR0, including hydrogen bonds between
26
27 Glu104 and cytosine 93, Lys107 and the O6 atom on guanine 113, Lys111 and the N7 atom on
28
29 guanine 114, and Arg112 and the N7 atom on adenine 91 (**Fig. 5b**). In addition to the core DBD
30
31 there was also density to model the CTE and Ftz-F1 domains. The CTE of hLRH-1 wedges into
32
33 the minor groove of the DNA duplex (**Fig. 6c**). Base-specific contacts formed by the CTE
34
35 include hydrogen bonds between Arg162 and thymine 95 and Arg165 and cytosine 110. Other
36
37 stabilizing interactions with the phosphate backbone and the water network are shown in **Figure**
38
39 **5c**.
40
41
42
43
44
45
46
47
48

49 *LRH-1 and GCNF differentially recognize the DR0 of the Oct4 PP*

50
51

52
53 To understand how these distinct proteins recognize the same *Oct4* DR0 DNA, we
54
55 compared both the primary amino acid sequence and structure (**Fig. 6**). Alignment of mGCNF
56
57 and hLRH-1 DBDs show 44.7% identity with 61.8% similarity. Though human LRH-1 DBD
58
59 was used, sequence alignments show that human and mouse LRH-1 DBDs are 94.4 % identical,
60
61
62
63
64
65

1
2
3
4 with the critical DNA binding residues being 100% conserved. Mouse GCNF (495 aa) and
5
6 human GCNF (480 aa) are 95.4% identical, with both the DBD and CTE 100% identical.
7
8
9 Furthermore, the core DR0 motif within the *Oct4* PP is conserved between mice and humans,
10
11 suggesting these interactions would be maintained across species.
12
13
14

15 In order to visualize differences in the structure, we superimposed both to identify how
16
17 each recognizes the same DNA element (**Fig. 6**). Unlike, GCNF, LRH-1 binds to DNA as a
18
19 monomer. While this is uncommon among NRs, other monomeric receptors include SF-1,
20
21 ERR2, REV-ERB [15,28,29]. Perhaps not surprising, the LRH-1 DBD binds the *Oct4* DR0
22
23 element with greater affinity than the GCNF DBD, which is lacking the LBDs that are known to
24
25 participate in receptor dimerization [24]. Superposition of the structures revealed LRH-1 makes
26
27 four additional base-specific contacts within the major and minor groove. The root-mean-square
28
29 deviation (r.m.s.d.) between the DBD core domains of LRH-1 (83-154) and GCNF (74-147) is
30
31 0.46 Å (63 Cα aligned). In the DNA reading helix, LRH-1 displays two additional base-specific
32
33 contacts with Glu104 and Lys107 making hydrogen bonds to cytosine 93 and guanine 114,
34
35 respectively (**Fig. 5b**). The GCNF and LRH-1 CTE share 56.3% identity with 62.5% similarity
36
37 and a similar backbone position within the DNA minor groove (r.m.s.d 0.51 Å; 13 Cα aligned).
38
39 However, LRH-1 makes two additional H-bonding contacts vs GCNF via Arg160 and 162 (**Fig.**
40
41 **5c,e**) where GCNF contains glycine and proline residues at those positions (**Fig. 5d,e**). Mutation
42
43 of the GCNF CTE glycine and proline to arginine residues (Gly149Arg and Pro151Arg), to
44
45 mimic the LRH-1 CTE, increased the GCNF DBD affinity from 170 nM to 20 nM (**Fig. 5f**).
46
47 Conversely, simultaneous Arg160Gly and Arg162Pro mutations within the LRH-1 CTE reduced
48
49 binding affinity from 60 nM to 750 nM (**Fig. 5f**). Taken together, difference within the CTE
50
51 drive differential affinity for the isolated DBDs for the *Oct4* promoter. The difference in affinity
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 between these isolated DBDs should not be used to predict in-cell kinetics; however, it is clear
5
6 that these proteins compete for *Oct4* proximal promoter binding when co-expressed *in vivo*.
7
8

9 **Discussion**

10
11 The orphan nuclear receptors, LRH-1 and GCNF play a critical role in regulating genes
12
13 central to embryonic development [3,7]. Notably, LRH-1 and GCNF reciprocally regulate the
14
15 pluripotency factor, Oct4, by binding to a DR0 response element within the promoter [3,7].
16
17

18
19 Though many have studied this mechanism, the structural basis for this regulation has yet to be
20
21 explored. Here, we not only present the first structure of GCNF DBD but also LRH-1 DBD
22
23 bound to the *Oct4* DR0 sequence and examine the sequence-specific contacts that guide *Oct4*
24
25 regulation.
26
27

28
29 Understanding the DNA-binding properties of transcription factors is required to
30
31 understand their biological function. Initial gel mobility shift assays show GCNF binds direct
32
33 repeats of the AGGTCA sequence with no spacer or to extended half-sites of TCAAGGTCA
34
35 sequence as a homodimer, though GCNF bound to the DR0 with higher affinity than the half-
36
37 sites [18,19]. Deletion studies show that removal of the ligand-binding domain had no effect of
38
39 DNA binding and *in vivo* removal of this domain did not affect transrepression [30, 31].
40
41 Additionally, N-terminal domain deletions still bound DNA to the same levels of WT GCNF
42
43 [20]. In contrast, removal of the DBD *in vivo* phenocopies the GCNF complete knockout mice
44
45 [23]. Additionally, the DBD and the CTE are strictly required to bind DNA making them
46
47 necessary to repress Oct4 [20,30]. For this reason, we purified the minimal residues required for
48
49 DNA recognition (**Fig. 2A**).
50
51
52
53
54

55
56 We show that purified GCNF DBD binds to the *Oct4* DR0 via a two-site binding
57
58 mechanism with a high and low affinity binding event (**Fig. 2c**). In contrast, intact *in vitro*
59
60
61
62
63
64
65

1 translated full-length protein binds DNA as a homodimer [15, 20, 30]. The overall structure
2
3 shows two GCNF molecules bound to opposite sides of the DNA and the sister DBDs only
4
5 weakly interact (**Fig. 3,4**). This structure is similar to retinoic X receptor bound to a DR1
6
7 sequence [27]. The GCNF DBD sequence has been shown to be most similar to the RXR DBD,
8
9 which requires an intact LBD for receptor dimerization; therefore, it is likely that strong
10
11 homodimerization of GCNF also requires the LBD [19,24]. Dimerization motifs have been
12
13 proposed in the DBD, CTE, and LBD. Structural analysis reveals that the CTE makes a number
14
15 of hydrophobic contacts with the other GCNF molecule (**Fig. 4**). Additionally, the side chains
16
17 that participate in homodimerization are very well conserved (**Fig. 4c**). Therefore, structural
18
19 studies of the GCNF LBD or full-length will be required complete the understanding of
20
21 GCNF:DNA recognition [24, 30].
22
23
24
25
26
27
28
29
30

31 LRH-1 is required to maintain *Oct4* expression [3]. Monomeric LRH-1 interacts with
32
33 extended half-site sequences in both the proximal promoter and proximal enhancer to activate
34
35 *Oct4* expression prior to differentiation [16]. The only reported structures of NR5A DBDs are
36
37 the x-ray structure of the LRH-1-*hCYP7A1* complex (PDB ID: **2A66**) and the NMR solution
38
39 structure of SF-1 in complex with a 9 bp fragment of *inhibin-α* (PDB ID: **2FF0**; 16 conformers)
40
41 [10,15]. While SF-1 and LRH-1 bind the same response elements, previous data show that SF-1
42
43 is not expressed at detectable levels in ES cells [3]. We show that purified LRH-1 DBD and full-
44
45 length LRH-1 bind the *Oct4* DR0 with high affinity as a monomer positioned over the extended
46
47 LRH-1 recognition sequence (**Fig. 2b,d, 5**).
48
49
50
51
52

53 Comparing the GCNF and LRH-1 structures revealed that their respective response
54
55 elements directly overlap in the DR0 of the PP. (**Fig. 6**). LRH-1 is highly expressed in ES cells
56
57 when *Oct4* expression is up regulated [3]. However, upon signals to differentiate, both LRH-1
58
59
60
61
62
63
64
65

1
2
3
4 and *Oct4* expression is rapidly decreased [3, 28]. During this time GCNF expression is high (**Fig.**
5
6
7 **1**) [7]. LRH-1 has an apparent higher affinity *in vitro*, potentially due to additional side chain –
8
9 DNA base interactions (**Fig. 5,6**). Mutational analysis of GCNF CTE to mimic the LRH-1 CTE
10
11 shows an increased affinity for the *Oct4* DR0 and the high LRH-1 affinity can be drastically
12
13 reduced when the Arg residues of the CTE are mutated to the GCNF glycine and proline residues
14
15 at the same positions (**Fig. 5f**). Since binding was only being tested with isolated domains, this
16
17 result is not surprising. Kinetic studies with intact GCNF may reveal similar affinities for the
18
19 element. Yet, it is possible that GCNF will not have to compete for binding to the DR0 during
20
21 embryonic development due to the inverse expression patterns of LRH-1 and GCNF (**Fig 6**)
22
23
24
25
26 [3,18].
27
28
29

30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
Mutational analyses of C-terminal extensions from numerous nuclear receptors such as
SF-1, RXR, LRH-1, and GCNF have shown this region to be critical for DNA recognition and
transcriptional regulation [10, 24, 32, 33]. The GCNF structure shows electron density for the
CTE within molecule 2 (**Fig 5d**). Removal of these residues from GCNF prevents DNA binding,
highlighting their importance [19]. Furthermore, DNA sequence analysis revealed that the TCA
sequence preceding the second direct repeat, that forms the minor groove CTE interaction, is
most important for GCNF to bind DNA [19, 20]. Here, GCNF uses an Arg residue to make two
base-specific contacts within the minor groove (**Fig 5d**). PISA analysis of the GCNF structure
reveals that the average gain on complex formation for molecule 2 and DNA is -9.1 kcal/mol
(complex score 1.0) but with the CTE removed it is -5.2 kcal/mol (complex score 0.190) and has
a much weaker complex score (1.0 being highly favorable complex) [34]. Like GCNF, the LRH-
1 CTE also dips into the minor groove to make additional base-specific contacts (**Fig 5c**).

Structure alignments of GCNF, LRH-1, RXR, and SF-1 bound to DNA elements show high conservation among the overall DBD structures. The r.m.s.d. of LRH-1 and GCNF on the *Oct4* DR0 is 0.57 Å (76 Cα aligned), while RXRα and GCNF is 0.66 Å (66 Cα aligned) and SF-1 and LRH-1 is 1.12 Å (88 Cα aligned). Though GCNF is cited as most similar to RXR, the CTE regions are very different. When overlaid, these structures show that the RXRα CTE does not rest in the DNA minor groove but trails off away from the DNA. Furthermore, sequence alignments of GCNF₁₂₆₋₁₄₁ and RXRα₂₀₁₋₂₁₆ only have 27% identity and 40% similarity. These differences also set GCNF apart in its own subclass of the nuclear receptor superfamily. LRH-1₁₅₂₋₁₆₇ and SF-1₇₉₋₉₄ on the other hand, are 100% identical. Aligned structures show that both of these proteins' CTEs rest in the minor groove of DNA leading into NR5A conserved Ftz-F1 helices. The NMR solutions of SF-1 show that, although the CTE adopts multiple conformations, all make contacts in the minor groove, as in both LRH-1 structures (PDB ID: [2A66](#) and [5L0M](#)). Interestingly, replacing the GCNF CTE with that from LRH-1/SF-1, induces a monomeric DNA binding preference likely by removing conserved GCNF intermolecular contacts (i.e. Met150) and increasing half-site affinity [24]. In agreement, the WT GCNF binds to the *Oct4* DR0 with an apparent two-site binding curve but the GCNF Gly142Arg/Pro151Arg mutant binds the same element using a one-site binding mechanism (**Fig. 5f**). As these mutations now mimic the LRH-1 and SF-1 CTE, this result is not surprising. Structural studies using intact LRH-1 and GCNF would shed deep insight *Oct4* recognition; however, it is clear that the CTEs of these receptors play a critical role not only in sequence specificity but oligomerization on DNA [3].

Materials and Methods

Protein expression and purification

The DNA binding domain (DBD) of mouse GCNF (residues 69-180, UniProt Q64249)

1
2
3
4 with a C104S mutation was cloned into a pMCGS7 vector with a 6X-Histidine tag. GCNF was
5
6 expressed in BL21 (DE3) pLysS *E. coli* cells. The protein was grown at 37 °C for 2 hrs, then
7
8 reduced to 20 °C and grown until an OD₆₀₀ of 0.6, and then induced with 0.78 mM Isopropyl β-
9
10 D-1-thiogalactopyranoside (IPTG). Cultures were grown overnight at 20 °C. Cells were lysed in
11
12 20 mM Tris-HCl (pH 7.4), 1M NaCl, 25 mM imidazole, and 5% glycerol via sonication. Protein
13
14 was purified using affinity chromatography (His-Trap FF, GE Healthcare) followed by further
15
16 purification via gel filtration chromatography. Protein was then concentrated to 2-3 mg/ml in 20
17
18 mM Tris-HCL (pH 7.4), 150 mM NaCl, and 5% glycerol, flash frozen in liquid N₂, and stored at
19
20 -80 °C.
21
22
23
24

25
26 The DBD-FtzF1 domains of human LRH-1 (LRH-1 DBD) were expressed and purified
27
28 similar to previous [10]. Briefly, cultures were grown in Terrific Broth (TB) at 37 °C to OD₆₀₀ of
29
30 0.6, induced with 0.3 mM IPTG at 18 °C and grown overnight. Fusion protein was purified via
31
32 affinity chromatography (His-Trap FF, GE Healthcare) and the MBP tag was removed following
33
34 TEV protease cleavage with an additional pass over His-Trap FF resin. LRH-1 DBD was further
35
36 purified using gel filtration (Superdex 75; GE Healthcare) equilibrated with 20 mM Tris-HCL
37
38 (pH 7.4), 150 mM NaCl, and 5% glycerol. Eluted protein was concentrated to 11.5 mg/mL and
39
40 either flash frozen in liquid N₂ and stored at -80 °C or used directly for crystallization
41
42 experiments.
43
44
45
46
47

48 hLRH-1 has multiple full-length functional isoforms, the canonical sequence (isoform 2;
49
50 1-541) was used to number the residues of the crystal structure [10]. Experimentally, isoform 1
51
52 (1-495) lacking residues 22-67 in the N-terminal domain, was used to generate a construct
53
54 (amino acids 2-495) preceded by a TEV protease site, was cloned into the pE-SUMO-Amp
55
56 vector (LifeSensors) and recombinantly expressed in *E. coli* BL21 DE3 cells. Cells were grown
57
58
59
60
61
62
63
64
65

1
2
3
4 in LB at 37 °C to OD₆₀₀ of 0.6, induced with 0.5 mM IPTG at 20 °C and grown overnight. Cells
5
6 were lysed in 20 mM Tris-HCl (pH 8.5), 0.5 M NaCl, 25 mM imidazole, 2 mM CHAPS, 0.2%
7
8 Triton-100X and pierce protease inhibitor tablets (Thermo Fisher Scientific) via sonication.
9
10 Fusion protein was purified by affinity chromatography (His-Trap FF and HiTrap Heparin, GE
11
12 Healthcare). The His-SUMO fusion tag was removed by incubation with TEV overnight and
13
14 affinity chromatography used to isolate pure target protein.
15
16
17

18 19 *Generation of GCNF DBD Gly149Arg/Pro151Arg and LRH-1 DBD Arg160Gly/Arg162Pro*

20

21 Mutagenesis was performed following the MEGAWHOP protocol. Briefly, megaprimers
22
23 were generated containing the desired mutations, purified and used for whole plasmid PCR [35].
24
25
26 Mutant proteins were prepared as wild-type.
27

28 29 *Sequence Alignments and Analyses*

30

31 The following sequences were obtained from UniProt: mGCNF (UniProt Q64249-1),
32
33 hGCNF (Uniprot Q15406-1), mLRH-1 (UniProt P45448-1), hLRH-1 (UniProt O00482-1), hSF-1
34
35 (UniProt Q13285-1) and hRXR α (UniProt P19793-1) and aligned using Clustal Omega [36].
36
37
38 Jalview [37] was used for visualization and manipulation of the alignments. The Sequence
39
40 Manipulation Suite (SMS) was used for percent identity and similarity calculations [38].
41
42

43 44 *Nucleic acid binding assays*

45

46 Synthesized FAM-labeled nucleic acid duplexes (Integrated DNA Technologies) of
47
48 mouse *Oct4* DR0 (5' – [FAM] AGAGGTCAAGGCTAGA – 3') were annealed in a 1 L water
49
50 bath heated to 90 °C then cooled slowly to room temperature. Fluorescence polarization assays
51
52 were performed by adding increasing concentrations of GCNF, LRH-1 DBD, or FL LRH-1 (1
53
54 nM- 50 μ M for the DBDs; 1 nM- 20 μ M for FL hLRH-1) to 10 nM of the FAM-labeled DNA.
55
56
57
58 Reactions were performed in 20 mM Tris-HCL (pH 7.4), 150 mM NaCl, and 5% glycerol.
59
60
61
62
63
64
65

Polarization was monitored on a Biotek Synergy plate-reader at an excitation/emission wavelength of 485/528 nm. The program GraphPad Prism 6 was used to analyze binding data and generate graphs. Binding data were analyzed with an F-test to compare a two-site binding event to a one-site binding event with Hill slope. This test generated an F-statistic and p-value supporting a two-site binding model. These values are represented in Figure 2. Additionally, dissociation values (K_d) and coefficient of determination (r^2) are included.

Crystallization, data collection, and structure determination

Crystals of the GCNF-*mOct4* (16bp - 5' – AGAGGTCAAGGCTAGA – 3') complex were grown by hanging drop vapor diffusion in 0.1M Tris pH 8.5, 20% PEG 3350, 3% glycerol with a 2:1 protein:DNA molar ratio. Crystals were cryo-protected with 0.1M Tris pH 8.5, 30% PEG 3350, and 15% glycerol and flash cooled in liquid N₂. The hLRH-1 DBD-*mOct4* (12bp duplex - 5' – GGTC AAGGCTAG – 3') complex was formed by mixing 1:1.2 molar ratios of protein to DNA and incubating at 25 °C. Crystallization conditions were screened using a Phoenix nanolitre drop dispensing robot (Art Robbins) with a 1:1 protein (5-6 mg/mL) to well solution ratio. Single well-formed crystals appeared overnight at 18 °C in 0.2 M calcium acetate and 20% (w/v) PEG 3350. Larger crystals were grown by hanging drop vapor diffusion in the same solution. Crystals were cryo-protected with an additional 20% (v/v) glycerol and flash cooled in liquid N₂.

Data were collected at the 22-ID beamline (Advanced Photon Source, Argonne, IL) and processed using HKL-2000 [39]. The structures were phased using a low-resolution model GCNF-*Oct4*, previously generated in the lab or the hLRH-1 DBD-*CYP7a1* complex (PDB ID: **2A66**) [10]. Structure refinement and validation was performed using PHENIX refine software and model building was performed in COOT [40,41]. PDB Redo was used iteratively to optimize

refinement parameters and geometry [42]. PyMOL v1.8.2 was used to visualize structures and generate figures (Schrödinger, LLC). Both structures showed good overall geometry with one Ramachandran outlier in the LRH-1 DBD – DNA complex and all other residues (93) in favored or allowed regions and zero Ramachandran outliers in the GCNF DBD – DNA complex and all in favored or allowed regions.

Accession Numbers

Coordinates and structure factors have been deposited in the Protein Data Bank with the accession numbers, 5KRB for GCNF:*Oct4* complex and 5L0M for LRH-1:*Oct4* complex.

Acknowledgements

X-ray data were collected at Southeast Regional Collaborative Access Team (SER-CAT) 22-ID beamline at the Advanced Photon Source, Argonne National Laboratory. Supporting institutions may be found at www.ser-cat.org/members/html. Use of the Advanced Photon source was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. W-31-109-Eng-38.

Author Contributions

Conceived and designed experiments: ERW MLT EAO. Performed the experiments ERW MLT MNM EAO. Analyzed the data ERW MLT EAO. Contributed to writing the manuscript ERW MLT EAO.

References

1. Paranjpe SS, Veenstra GJ. Establishing pluripotency in early development. *Biochim Biophys Acta*. 2015;1849:626-36.
2. Wu G, Scholer HR. Role of Oct4 in the early embryo development. *Cell Regen (Lond)*. 2014;3:7.

3. Gu P, Goodwin B, Chung AC, Xu X, Wheeler DA, Price RR, et al. Orphan nuclear receptor LRH-1 is required to maintain Oct4 expression at the epiblast stage of embryonic development. *Molecular and cellular biology*. 2005;25:3492-505.
4. Sylvester I, Scholer HR. Regulation of the Oct-4 gene by nuclear receptors. *Nucleic Acids Res*. 1994;22:901-11.
5. Mullen EM, Gu P, Cooney AJ. Nuclear Receptors in Regulation of Mouse ES Cell Pluripotency and Differentiation. *PPAR Res*. 2007;2007:61563.
6. Olefsky JM. Nuclear receptor minireview series. *The Journal of biological chemistry*. 2001;276:36863-4.
7. Fuhrmann G, Chung AC, Jackson KJ, Hummelke G, Baniahmad A, Sutter J, et al. Mouse germline restriction of Oct4 expression by germ cell nuclear factor. *Dev Cell*. 2001;1:377-87.
8. Palmieri SL, Peter W, Hess H, Scholer HR. Oct-4 transcription factor is differentially expressed in the mouse embryo during establishment of the first two extraembryonic cell lineages involved in implantation. *Dev Biol*. 1994;166:259-67.
9. Barnea E, Bergman Y. Synergy of SF1 and RAR in activation of Oct-3/4 promoter. *The Journal of biological chemistry*. 2000;275:6608-19.
10. Solomon IH, Hager JM, Safi R, McDonnell DP, Redinbo MR, Ortlund EA. Crystal structure of the human LRH-1 DBD-DNA complex reveals Ftz-F1 domain positioning is required for receptor activity. *Journal of molecular biology*. 2005;354:1091-102.
11. Yussa M, Löhr U, Su K, Pick L. The nuclear receptor Ftz-F1 and homeodomain protein Ftz interact through evolutionarily conserved protein domains. *Mechanisms of Development*. 2001;107:39-53.

- 1
2
3
4 12. Pare, J. F., Malenfant, D., Courtemanche, C., Jacob-Wagner, M., Roy, S., Allard, D. &
5
6 Belanger, L. (2004). The fetoprotein transcription factor (FTF) gene is essential to
7
8 embryogenesis and cholesterol homeostasis and is regulated by a DR4 element. *J Biol Chem*
9
10 279, 21206-16.
11
12
13 13. Takemaru, K., Li, F. Q., Ueda, H. & Hirose, S. (1997). Multiprotein bridging factor 1
14
15 (MBF1) is an evolutionarily conserved transcriptional coactivator that connects a regulatory
16
17 factor and TATA element-binding protein. *Proc Natl Acad Sci U S A* 94, 7251-6.
18
19
20 14. Cai, Y. N., Zhou, Q., Kong, Y. Y., Li, M., Viollet, B., Xie, Y. H. & Wang, Y. (2003). LRH-
21
22 1/hB1F and HNF1 synergistically up-regulate hepatitis B virus gene transcription and DNA
23
24 replication. *Cell Res* 13, 451-8.
25
26
27 15. Little TH, Zhang Y, Matulis CK, Weck J, Zhang Z, Ramachandran A, et al. Sequence-
28
29 specific deoxyribonucleic acid (DNA) recognition by steroidogenic factor 1: a helix at the
30
31 carboxy terminus of the DNA binding domain is necessary for complex stability. *Mol*
32
33 *Endocrinol.* 2006;20:831-43.
34
35
36 16. Sung B, Do HJ, Park SW, Huh SH, Oh JH, Chung HJ, et al. Regulation of OCT4 gene
37
38 expression by liver receptor homolog-1 in human embryonic carcinoma cells. *Biochem Biophys*
39
40 *Res Commun.* 2012;427:315-20.
41
42
43 17. Heng JC, Feng B, Han J, Jiang J, Kraus P, Ng JH, et al. The nuclear receptor Nr5a2 can
44
45 replace Oct4 in the reprogramming of murine somatic cells to pluripotent cells. *Cell Stem Cell.*
46
47 2010;6:167-74.
48
49
50 18. Chen F, Cooney AJ, Wang Y, Law SW, O'Malley BW. Cloning of a novel orphan receptor
51
52 (GCNF) expressed during germ cell development. *Mol Endocrinol.* 1994;8:1434-44.
53
54
55
56
57
58
59
60
61
62
63
64
65

19. Yan ZH, Medvedev A, Hirose T, Gotoh H, Jetten AM. Characterization of the response element and DNA binding properties of the nuclear orphan receptor germ cell nuclear factor/retinoid receptor-related testis-associated receptor. *The Journal of biological chemistry*. 1997;272:10565-72.
20. Cooney AJ, Hummelke GC, Herman T, Chen F, Jackson KJ. Germ cell nuclear factor is a response element-specific repressor of transcription. *Biochem Biophys Res Commun*. 1998;245:94-100.
21. Wang H, Wang X, Xu X, Kyba M, Cooney AJ. Germ Cell Nuclear Factor (GCNF) Represses Oct4 Expression and Globally Modulates Gene Expression in Human Embryonic Stem (hES) Cells. *The Journal of biological chemistry*. 2016;291:8644-52.
22. Chung AC, Katz D, Pereira FA, Jackson KJ, DeMayo FJ, Cooney AJ, et al. Loss of orphan receptor germ cell nuclear factor function results in ectopic development of the tail bud and a novel posterior truncation. *Molecular and cellular biology*. 2001;21:663-77.
23. Lan ZJ, Chung AC, Xu X, DeMayo FJ, Cooney AJ. The embryonic function of germ cell nuclear factor is dependent on the DNA binding domain. *The Journal of biological chemistry*. 2002;277:50660-7.
24. Greschik H, Wurtz JM, Hublitz P, Kohler F, Moras D, Schule R. Characterization of the DNA-binding and dimerization properties of the nuclear orphan receptor germ cell nuclear factor. *Molecular and cellular biology*. 1999;19:690-703.
25. Gu P, Xu X, Le Menuet D, Chung AC, Cooney AJ. Differential recruitment of methyl CpG-binding domain factors and DNA methyltransferases by the orphan receptor germ cell nuclear factor initiates the repression and silencing of Oct4. *Stem Cells*. 2011;29:1041-51

- 1
2
3
4 26. Sato N, Kondo M, Arai K. The orphan nuclear receptor GCNF recruits DNA
5
6 methyltransferase for Oct-3/4 silencing. *Biochem Biophys Res Commun*. 2006;344:845-51.
7
8
9 27. Zhao Q, Chasse SA, Devarakonda S, Sierk ML, Ahvazi B, Rastinejad F. Structural basis of
10
11 RXR-DNA interactions. *Journal of molecular biology*. 2000;296:509-20.
12
13
14 28. Gearhart MD, Holmbeck SM, Evans RM, Dyson HJ, Wright PE. Monomeric complex of
15
16 human orphan estrogen related receptor-2 with DNA: a pseudo-dimer interface mediates
17
18 extended half-site recognition. *Journal of molecular biology*. 2003;327:819-32.
19
20
21 29. Zhao Q, Khorasanizadeh S, Miyoshi Y, Lazar MA, Rastinejad F. Structural elements of an
22
23 orphan nuclear receptor-DNA complex. *Molecular cell*. 1998;1:849-61.
24
25
26 30. Borgmeyer U. Dimeric binding of the mouse germ cell nuclear factor. *Eur J Biochem*.
27
28 1997;244:120-7.
29
30
31 31. Okumura LM, Lesch BJ, Page DC. The ligand binding domain of GCNF is not required for
32
33 repression of pluripotency genes in mouse fetal ovarian germ cells. *PLoS One*. 2013;8:e66062.
34
35
36 32. Wilson TE, Fahrner TJ, Milbrandt J. The orphan receptors NGFI-B and steroidogenic factor
37
38 1 establish monomer binding as a third paradigm of nuclear receptor-DNA interaction. *Mol Cell*
39
40 *Biol*. 1993;13:5794-804.
41
42
43 33. Melvin VS, Roemer SC, Churchill ME, Edwards DP. The C-terminal extension (CTE) of the
44
45 nuclear hormone receptor DNA binding domain determines interactions and functional response
46
47 to the HMGB-1/-2 co-regulatory proteins. *The Journal of biological chemistry*. 2002;277:25115-
48
49 24.
50
51
52 34. Krissinel E, Henrick K. Inference of macromolecular assemblies from crystalline state. *J Mol*
53
54 *Biol*. 2007;372:774-97.
55
56
57
58
59
60
61
62
63
64
65

35. Miyazaki K. MEGAWHOP cloning: a method of creating random mutagenesis libraries via megaprimer PCR of whole plasmids. *Methods Enzymol.* 2011;498:399-406.
36. Stothard P. The sequence manipulation suite: JavaScript programs for analyzing and formatting protein and DNA sequences. *Biotechniques.* 2000;28:1102, 4.
37. Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics.* 2009;25:1189-91.
38. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol.* 2011;7:539.
39. Otwinowski Z, Minor W. Processing of X-ray Diffraction Data Collected in Oscillation Mode. *Methods in Enzymology.* 1997; 276: 307–362.
40. Adams PD, Afonine PV, Bunkoczi G, Chen VB, Davis IW, Echols N, et al. PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr.* 2010;66:213-21.
41. Emsley P, Lohkamp B, Scott WG, Cowtan K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr.* 2010;66:486-501.
42. Joosten RP, Salzemann J, Bloch V, Stockinger H, Berglund AC, Blanchet C, et al. PDB_REDO: automated re-refinement of X-ray structure models in the PDB. *J Appl Crystallogr.* 2009;42:376-84.

Figure Legends

Figure 1: Schematic representation of differential regulation of Oct4 by LRH-1 and GCNF.

In undifferentiated ES cells, LRH-1 expression is high and drives *Oct4* expression by binding to DR0 sequences in the proximal enhancer and promoter. This binding recruits coactivators and

1
2
3
4 the transcriptional machinery to drive gene expression. Upon signals to differentiate, LRH-1
5
6 expression is rapidly reduced while GCNF expression is high. GCNF then binds to the DR0
7
8 within the proximal promoter to repress Oct4 expression. Binding by GCNF recruits
9
10 corepressors, such as NCoR, to block Oct4 expression. GCNF also recruits DNA
11
12 methyltransferases (DMNT) and methyl-binding proteins (MBP) to methylate the Oct4 gene in
13
14 order to efficiently shut off its expression.
15
16
17

18
19 **Figure 2: GCNF and LRH-1 bind directly to the Oct4 DR0.** (a) Diagram of the GCNF and
20
21 LRH-1 modular structure. In this study, GCNF DBD-TA (residues 69-180), LRH-1 DBD Ftz-F1
22
23 (residues 79-187), and full-length LRH-1 (residues 2-495) were used. (b) GCNF DBD-TA bound
24
25 to the Oct4 DR0 in a two-site binding mechanism. (c) LRH-1 DBD Ftz-F1 (orange) and full-
26
27 length LRH-1 (green) bind the Oct4 DR0 in a one-site binding mechanism. Binding data are
28
29 represented as mean $\pm$ s.e.m from three replicates and from three independent fluorescence
30
31 polarization experiments. (d) Summary of binding data.
32
33
34

35
36 **Figure 3: Structural Analysis of GCNF - mOct4 Complex.** (a) Overall structure of GCNF
37
38 DBD (purple) bound to the Oct4 DR0 (gray). GCNF DBD and DNA shown as cartoon in purple
39
40 and white, respectively and zinc atoms as spheres. Two molecules of GCNF bound to opposite
41
42 sides of the DNA in a head-to-tail fashion. The DR0 sequence is shown below with arrows
43
44 denoting the direction of GCNF over the sequence. (b) Each molecule of GCNF makes base-
45
46 specific contacts with the DR0 (bases in white) mediated by hydrogen bonds (red) made between
47
48 Arg101 and Lys96. Mesh shows 2Fo-Fc electron density map contoured to 2σ around the DNA
49
50 bases. (c) Schematic view of protein-DNA interactions. Larger, bold side chains denoted base-
51
52 contacting side chains. Water molecules are indicated as red spheres.
53
54
55
56
57

58 **Figure 4: Interactions between GCNF molecules.** (a) Electrostatic surface of overall GCNF
59
60
61
62
63
64
65

1
2
3
4 structure. (b) Zoom in of contacts (black) made between the two molecules. Side chains from
5
6 molecule 1 are colored purple and dark purple from molecule 2. (c) Sequence alignments from
7
8 numerous species reveal that the side chains that participate in homodimer formation are highly
9
10 conserved.
11
12
13

14 **Figure 5: Structural Analysis of LRH-1 - mOct4 Complex.** (a) Overall structure of LRH-1
15
16 DBD (orange) bound to the 12 bp Oct4 DR0 (gray) with zinc atoms as spheres. The Oct4 DR0
17
18 sequence is shown below with arrows denoting the footprint and orientation of the LRH-1
19
20 binding site. (b) LRH-1 makes base-specific contacts with the DR0 (bases in white) mediated by
21
22 hydrogen bonds (red) made between Glu104, Lys107, Lys 111, Arg112. Mesh shows 2Fo-Fc
23
24 electron density map contoured to 2σ around the DNA bases. (c) Schematic view of protein-
25
26 DNA interactions. Larger, bold side chains denoted base-contacting side chains. Water
27
28 molecules are indicated as red spheres.
29
30
31
32

33 **Figure 6: GCNF and LRH-1 comparison.** (a) Overlay of GCNF and LRH-1 structures.
34
35 Molecule 2 of GCNF (Dark purple) sites directly on top of the LRH-1 (orange) recognition site.
36
37 (b) Close up view of the base-specific contacts mediated by these two receptors. LRH-1 makes
38
39 an additional contact with Lys 107 making hydrogen bonds to guanine 114. (c) LRH-1 CTE has
40
41 good electron density, mesh shows 2Fo-Fc electron density map contoured to 1σ around the
42
43 residues. Arg162 and 165 make hydrogen bonds (red) to thymine 95 and cytosine 110. (d) GCNF
44
45 CTE has good electron density, mesh shows 2Fo-Fc electron density map contoured to 2σ
46
47 around the residues. Arg86 makes hydrogen bonds to thymine 109 and cytosine 110. Arg78 also
48
49 folds into the CTE to make hydrogen bonds to Gly152. (e) Sequence alignment of GCNF and
50
51 LRH-1 CTEs show LRH-1 to contain two additional Arg residues that are used for DNA
52
53 binding. (f) Mutational analysis of CTE residues on *Oct4* binding: GCNF DBD
54
55
56
57
58
59
60
61
62
63
64
65

Gly149Arg/Pro151Arg (open purple circles) bound to the *Oct4* DR0 with an affinity 20 nM, where WT GCNF DBD (faded closed purple circles) binds with an affinity of 170 nM. LRH-1 DBD Arg160Gly/Arg162Pro (open orange circles) bound with an affinity of 750 nM, where WT LRH-1 DBD (faded closed orange circles) bound with an affinity of 60 nM.

Table 1: Data collection and refinement statistics

	GCNF DBD - <i>mOct4</i> (16bp)	LRH-1 DBD - <i>mOct4</i> (12bp)
Data Collection		
Space Group	P2 ₁ 2 ₁ 2 ₁	P4 ₃
Cell Dimension	a=53.6, b=69.5, c=84.5	a=40.9, b=40.9, c=105.1
Resolution (Å)	2.10 (2.18-2.10)	50-2.2 (2.28-2.20)
R _{pim}	6.9 (37.1)	3.7 (30.0)
I/σ	9.7 (2.0)	24.3 (2.1)
Completeness	96.2 (80.2)	99.8 (98.8)
Redundancy	5.2 (3.5)	7.2 (5.7)
Refinement		
Resolution	2.10	2.20
No. Reflections	18277	8789
R _{work} /R _{free}	21.6/26.8	15.4/19.7
No. Atoms		
Protein	1244	779
DNA	651	485
Water	43	50
B-factors		
Protein	56.0	59.1
DNA	51.8	52.6
Water	54.6	53.4
R.m.s. deviations		
Bond lengths (Å)	0.003	0.016
Bond angles (°)	0.57	1.65
PDB code	5KRB	5L0M

*Data collected from a single crystal; values in parentheses are for the highest-resolution shell

Figure 1

Figure 1

Undifferentiated ES cells

Expression Levels

LRH-1 ↑

Oct4 ↑

GCNF ↓

Differentiated ES cells

LRH-1 ↓

Oct4 ↓

GCNF ↑

Fully differentiated cells

Oct4 ↓

GCNF ↓

Figure 2

Figure 2

Figure 3

Figure 3

Figure 4

Figure 5

Figure 5

Figure 6