First analysis of the rotationally-resolved ν_2 and $2\nu_2 - \nu_2$ bands of sulfur dioxide, $^{33}\text{S}^{16}\text{O}_2$

T.A. Blake ^{a*}, J.-M. Flaud ^b, W.J. Lafferty ^c

Abstract:

A Fourier transform spectrum of sulfur dioxide $^{33}\text{S}^{16}\text{O}_2$ has been recorded in the 18.3 μm spectral region at a resolution of 0.002 cm^{-1} using a Bruker IFS 125HR spectrometer leading to the observation of the ν_2 and $2\nu_2 - \nu_2$ vibrational bands of the $^{33}\text{S}^{16}\text{O}_2$ molecule. The corresponding upper state ro-vibrational levels were fit using Watson-type Hamiltonians. In this way it was possible to reproduce the upper state ro-vibrational levels to within the experimental uncertainty; i.e., $\sim 0.20 \times 10^{-3} \text{ cm}^{-1}$. Very accurate rotational and centrifugal distortion constants were derived from the fit together with the following band centers: $\nu_0(\nu_2) = 515.659089(50) \text{ cm}^{-1}$, $\nu_0(2\nu_2) = 1030.697723(20) \text{ cm}^{-1}$.

Keywords: sulfur dioxide; sulfur-33 isotope; high resolution Fourier transform infrared spectroscopy; ro-vibrational constants.

1. Introduction

Sulfur dioxide plays an important role in the atmospheres of various planets [1-4]. Providing laboratory spectroscopic information on the less abundant isotopic species is then of interest for detecting them in those atmospheres. The fundamental bands of several isotopic species of SO₂ have been widely studied both in the microwave (see Refs. [5-13] and references therein) and infrared regions (see Refs. [14-21] and references therein). The ³³S¹⁶O₂ species has been the subject of spectroscopic studies in the microwave region [5, 9, 11, 13], but to our knowledge no high resolution study has been performed in the infrared region. In this paper we present the first detailed extensive analysis of the spectrum of this molecule in the 18.3 μm spectral region where the ν_2 and the $2\nu_2 - \nu_2$ bands absorb. The work was performed using a high-resolution Fourier transform spectrum (0.002 cm⁻¹ resolution) recorded at room temperature. The transition assignments have been extended to very high quantum numbers both for the ν_2 and the $2\nu_2 - \nu_2$ bands. The upper state rotational constants were derived through the fit of the upper state levels using a Watson-type Hamiltonian written in the I^r (x = b, y = c, z = a representation) [22]. It is also worth noting that the ground state constants were slightly improved through a simultaneous fit of the existing microwave data and the ground state combination differences derived in this work.

2. Experimental.

The ³³S¹⁶O₂ ro-vibrational spectra were recorded using a Bruker IFS 125HR spectrometer¹. The instrument was configured to operate between 400 and 900 cm⁻¹ using a KBr beamsplitter and a resistively heated silicon carbide infrared source. The spectrometer was evacuated below 0.015 Torr for these measurements. The sulfur dioxide sample was held in an adjustable path length White cell, the fore optics of which are seated in a sample compartment of the spectrometer. Wedged cesium iodide windows are used to separate the interior of the White cell from the interior of the spectrometer. The White cell optical path length for these

---

¹ Certain commercial equipment instruments or materials are identified in this paper to adequately specify the experimental procedure. Such identification does not imply recommendation or endorsement by the National Institute of Standards and Technology nor does it imply that the materials or equipment identified are necessarily the best available for the purpose.

measurements was $1604\text{ cm} \pm 10\text{ cm}$. Infrared light can be directed into the sample compartment without breaking vacuum. The White cell is hard-plumbed to a vacuum manifold that is used to evacuate the cell and introduce metered amounts of sample into the cell. Sample pressures were measured with a capacitance manometer (MKS Instruments model 690A01TRA; accuracy 0.05% of reading). The $^{33}\text{SO}_2$ sample pressure was 0.14483 hPa in the White cell. The temperature of the cell was not regulated, and was at room temperature, $25\text{ }^\circ\text{C} \pm 1\text{ }^\circ\text{C}$. For the 400 to 900 cm^{-1} range a liquid helium-cooled Si:Ga photodetector (Infrared Laboratories, Inc.) was used as the detector; it is fronted with a Winston cone and a 727 to 441 cm^{-1} passband filter. No other optical filters were used, the interferometer's aperture was set to 1.7 mm, and the scanner velocity was set to 40 kHz. Bruker's OPUS software (v7.0) was used to record the spectra.

A total of three-hundred twenty-eight coadded interferograms were processed by a Cooley-Tukey FFT with boxcar apodization, Mertz phase correction, and 1 cm^{-1} phase resolution. A zero filling factor of two was applied to the interferogram when it was initially recorded and an additional zero filling factor of eight was applied to bring the total zero filling factor to sixteen. Absorbance spectra were calculated as $\text{Abs} = -\ln(\text{sample}/\text{background})$, where background spectra were recorded at sixteen times the instrument resolution and sixteen times the zero filling factor used for the associated sample spectrum. The White cell was being actively evacuated while background spectra were being recorded.

Wavenumber calibration was performed by recording the spectrum of a sample of OCS in the White cell using the same instrument parameters as those used for recording the $^{33}\text{S}^{16}\text{O}_2$ spectra. For the 400 to 900 cm^{-1} wavenumber region, 0.1433 hPa of OCS was placed in the White cell with the path length set to 1604 cm and the absorption spectrum recorded. The average of the wavenumber differences between the recorded spectrum and those given in the online NIST wavenumber tables (see <https://www.nist.gov/pml/data/atlas-and-wavenumber-tables>) for fifty-nine OCS transitions is $0.001002(107)\text{ cm}^{-1}$. This number was added to the wavenumber scale of the $^{33}\text{SO}_2$ spectrum for this wavenumber range.

The $^{33}\text{S}^{16}\text{O}_2$ sample was prepared by weighing out 0.026 g of sulfur-33 elemental powder (Trace Sciences International, Corp.; 99.8% ^{33}S enrichment) into a 100 mL vacuum Schlenk tube

with a side arm (Kimble/Kontes model 218720-0100). The tube was then connected to the vacuum manifold, the air in the tube was evacuated, and the tube was then backfilled with 211 hPa of $^{16}\text{O}_2$ (Matheson). The tube was then wrapped with heat tape and baked at 488 °C for approximately 16 hours. The sample was allowed to cool to room temperature and then was put through two freeze/pump/thaw cycles at liquid nitrogen temperature. As each spectroscopic measurement was finished, the sample was cryotrapped back into the Schlenk tube, and two freeze/pump/thaw cycles were performed prior to putting the sample into a gas cell for the next measurement.

3. Assignments and Results

The line assignment process proved to be somewhat difficult given the density of lines in the spectrum. The analysis was started by analyzing the ν_2 band which is the strongest infrared band in the 18.3 μm spectral region. The first estimation of the ground state rotational levels was made using the rotational constants quoted in Ref. [23]. As far as the 2^1 rotational levels are concerned, a first calculation was made using rotational constants estimated from the rotational constants of the $^{32}\text{S}^{16}\text{O}_2$ [16] and $^{34}\text{S}^{16}\text{O}_2$ [17] species. In this way it was possible to assign a few lines with rather low rotational quantum numbers. Then, as soon as a few lines were assigned, the corresponding upper state energy levels were fit with a computer program [24] using a Watson-type Hamiltonian (Table 1) and the refined upper state Hamiltonian constants were used to perform more reliable predictions allowing one to assign new lines. The process was repeated until it was no longer possible to locate and assign new lines with confidence.

However, it is worth noting that for high J and K_a quantum numbers the observed ground state combination differences were marginally different from those derived from the original calculation performed with the constants of Ref. [23]. Accordingly, we fit simultaneously the ground state microwave data together with the combination differences derived in this work and the corresponding ground state constants are given in Table 2. The 2^1 upper state rotational levels were fit together with the existing 2^1 microwave data [5, 8] and the corresponding constants are given in Table 2. The results of the analysis are given in Table 3 showing that the fit was very satisfactory since the standard deviation is fully consistent with the experimental uncertainty.

For the $^{32}\text{SO}_2$ [16] and $^{34}\text{SO}_2$ [17] isotopic species a $\Delta K = \pm 2$ Fermi-type resonance coupling the 1^1 and 2^2 interacting energy levels was included in the fit of the experimental data. For $^{33}\text{SO}_2$, however, adding the Fermi resonance term did not alter in any statistically significant way the fit of the data, so that term was not included in the final fit.

Concerning the $2\nu_2$ - ν_2 band, a first estimation of the 2^2 upper state Hamiltonian constants was made by extrapolating the 2^1 and ground state constants. As with the ν_2 band, we were able to assign a few lines with low quantum numbers which were used to refine the upper state constants. In this way new lines could be assigned and the process was repeated until no more assignments could be performed. The corresponding upper state constants are given in Table 2 and the results of the fit in Table 3. As with the 2^1 state, the fit of the 2^2 state is very satisfactory as shown by its standard deviation, which is on the order of the experimental uncertainty.

The method used to compute line intensities is described in detail in Ref. [25]. However, it should be noted that in the present case only relative intensities were computed using the $^{32}\text{S}^{16}\text{O}_2$ transition moment [25, 26] since no attempt was made to derive experimental absolute intensities. Figures 1 and 2 show detailed portions of ν_2 and $2\nu_2$ - ν_2 bands, respectively. The very good agreement between observation and simulation² demonstrates the quality of the analysis.

Using an empirically refined potential energy surface and *ab initio* dipole moment surface, Huang, Schwenke and Lee have calculated, and recently published, empirical spectroscopic line lists (transition frequencies and intensities) for five sulfur dioxide isotopic species: $^{32/33/34/36}\text{S}^{16}\text{O}_2$ and $^{32}\text{S}^{18}\text{O}_2$ [27]. For $^{33}\text{SO}_2$ they compared their results with the published experimental ground state microwave spectra [5, 8, 9, 11] and found good agreement for line positions with the majority of differences between observed and calculated less than 0.004 cm^{-1} and relative intensity deviations in the range of -10% to 0%. The experimental data presented here will provide an opportunity for future comparisons with these empirical line lists.

4. Conclusion

Using a high-resolution Fourier transform spectrometer we have recorded a rotationally-resolved, room temperature spectrum of sulfur dioxide $^{33}\text{S}^{16}\text{O}_2$ and have performed an extensive

---

² The fit energy levels for the 2^1 and 2^2 states are available as Supplementary Material for this article. These tables are also available upon request from one of the authors: JMF.

analysis of the 18.3 μm spectral region. Microwave transitions and ground state combination differences formed from the infrared spectrum were combined and fit to provide a set of improved ground state spectroscopic constants. The upper state ro-vibrational energy levels of the 2^1 and 2^2 vibrational states have been fit to within the experimental accuracy ($\sim 0.20 \times 10^{-3} \text{ cm}^{-1}$) using Watson-type Hamiltonians. In this way, accurate band centers, rotational and centrifugal distortion constants have been determined. As for the intensities, more work is required to measure absolute and reliable individual line intensities. The high resolution infrared spectra of other $^{33}\text{SO}_2$ bands have been recorded and their analysis are the subject of another publication [28].

Acknowledgments

One of authors (JMF) thanks the Sensor Science Division for support during a stay at NIST. The infrared spectra were recorded at the Pacific Northwest National Laboratory (PNNL). PNNL is operated for the United States Department of Energy by the Battelle Memorial Institute under contract DE-AC05-76RLO 1830. Funding for recording the spectra was provided by PNNL's Laboratory Directed Research and Development program. TAB would like to thank Dr. James Moran for this support.

References

- [1] P.J. Wallace, J. Volcanol. Geotherm. Res. 108 (2001) 85-106.
- [2] R.M. Nelson, A.L. Lane, D.L. Matson, F.P. Fanale, D.B. Nash, T.V. Johnson, Science 210 (1980) 784-786.
- [3] J.R. Spencer, E. Lellouch, J.J. Richter, M.A. Lopez-Ververde, K.L. Jessup, T.K. Greathouse, J.-M. Flaud, Icarus 176 (2005) 283-304.
- [4] L.E. Snyder, J.M. Hollis, B.L. Ulich, F.J. Lovas, D.R. Johnson, D. Buhl, Ap. J. Lett. 198 (1975) L81-L84.
- [5] F.J. Lovas, J. Phys. Chem. Ref. Data 7 (1978) 1445-1750.
- [6] R. Van Riet, G. Steenbeckeliers, Ann. Soc. Sc. Brux. 97 III (1983) 117-153.
- [7] R. Van Riet, G. Steenbeckeliers, Ann. Soc. Sc. Brux. 97 IV (1984) 243-258.
- [8] F.J. Lovas, J. Phys. Chem. Ref. Data 14 (1985) 395-488.

- [9] E. Klisch, P. Schilke, S. P. Belov, G. Winnewisser, J. Mol. Spectrosc. 186 (1997) 314-318.
- [10] S.P. Belov, M.Y. Tretyakov, I.N. Kozin, E. Klisch, G. Winnewisser, W.J. Lafferty, J.-M. Flaud, J. Mol. Spectrosc. 191 (1998) 17-27.
- [11] H.S.P. Müller, J. Farhoomand, E.A. Cohen, B. Brupbacher-Gatehouse, M. Schäfer, A. Bauder, G. Winnewisser, J. Mol. Spectrosc. 201 (2000) 1-8.
- [12] H.S.P. Müller, S. Brunken, J. Mol. Spectrosc. 232 (2005) 213-222.
- [13] T. Helgaker, J. Gauss, G. Cazzoli, C. Puzzarini, J. Chem. Phys. 139 (2013) 244308-1-9.
- [14] J. Lindenmayer, H. Jones, H.D. Rudolph, J. Mol. Spectrosc. 101 (1983) 221-228.
- [15] J. Lindenmayer, H. Jones, J. Mol. Spectrosc. 126 (1987) 58-62.
- [16] J.-M. Flaud, A. Perrin, L.M. Salah, W.J. Lafferty, G. Guelachivili, J. Mol. Spectrosc. 160 (1993) 272-278.
- [17] W.J. Lafferty, J.-M. Flaud, R.L. Sams, El Hadji Abib Ngom, J. Mol. Spectrosc. 252 (2008) 72-76.
- [18] O.N. Ulenikov, E.S. Bekhtereva, S. Alanko, V.M. Horneman, O.V. Gromova, C. Le Roy, J. Mol. Spectrosc. 257 (2009) 137-156.
- [19] J.-M. Flaud, W.J. Lafferty, R.L. Sams, J. Quant. Spectrosc. Rad. Transfer. 110 (2009) 669-674.
- [20] O.N. Ulenikov, G.A. Onopenko, O.V. Gromova, E.S. Bekhtereva, V.-M. Horneman, J. Quant. Spectrosc. Rad. Transfer 130 (2013) 220-232.
- [21] O.N. Ulenikov, E.S. Bekhtereva, Yu.V. Krivchikova, Yu.B. Morzhikova, T. Buttersack, C. Sydow, S. Bauerecker, J. Quant. Spectrosc. Rad. Transfer 166 (2015) 13-22.
- [22] J.K.G. Watson, in *Vibrational Spectra and Structure*, edited by J.R. Durig (Elsevier, Amsterdam, The Netherlands, 1977), Vol 6, Ch. 1, pp. 1-89.
- [23] H.S.P Müller, F. Schloder, G. Winnewisser, J. Mol. Struct. 742 (2005) 215-227.
- [24] J.-M. Flaud, C. Camy-Peyret, Mol. Phys. 26 (1973) 811-823.
- [25] P.M. Chu, S.J. Wetzel, W.J. Lafferty, A. Perrin, J.-M. Flaud, Ph. Arcas, G. Guelachvili, J. Mol. Spectrosc. 189 (1998) 55-63.
- [26] L.S. Rothman, I.E. Gordon, Y. Babikov, A. Barbe, D. Chris Benner, P.F. Bernath, M. Birk, L. Bizzocchi, V. Boudon, L.R. Brown, A. Campargue, K. Chance, E.A. Cohen, L. Coudert,

V.M. Devi, B.J. Drouin, A. Fayt, J.-M. Flaud, R.R. Gamache, J.J. Harrison, J.-M. Hartmann, C. Hill, J.T. Hodges, D. Jacquemart, A. Jolly, J. Lamouroux, R.J. LeRoy, G. Li, D.A. Long, O.M. Lyulin, C.J. Mackie, S.T. Massie, S. Mikhailenko, H.S.P. Müller, O.V. Naumenko, A.V. Nikitin, J. Orphal, V. Perevalov, A. Perrin, E.R. Polovtseva, C. Richard, M.A.H. Smith, E. Starikova, K. Sung, S. Tashkun, J. Tennyson, G.C. Toon, V.I.G. Tyuterev, and G. Wagner, *J. Quant. Spectrosc. Rad. Transfer* 130 (2013) 4-50.

[27] X. Huang, D.W. Schwenke, T.J. Lee, *J. Mol. Spectrosc.* 311 (2015) 19-24.

[28] J.-M. Flaud, T.A. Blake, W.J. Lafferty, *Mol. Phys.* (2017). Accepted.

Table 1. Watson-type Hamiltonian used to calculate the 2^1 and 2^2 ro-vibrational levels of sulfur dioxide $^{33}\text{S}^{16}\text{O}_2$.

Table 2. Vibrational band centers, rotational and centrifugal distortion constant (in cm^{-1}) for the ground state and the 2^1 and 2^2 vibrational states of sulfur dioxide $^{33}\text{S}^{16}\text{O}_2$.

Footnotes:

^aUncertainties are given in parentheses in units of the last significant digits as stated, in terms of one standard deviation σ in the least squares adjustment (coverage factor $k=1$, B.N. Taylor and C.E. Kuyatt, NIST Technical Note 1297, U.S. Government Printing Office, 1994, pp. 1-20. The publication may be downloaded from <http://physics.nist.gov/Pubs/guidelines/contents.html>). The number of digits in the uncertainties kept for the parameters (2 for the band centers, 3 for the rotational constants and 4 for the centrifugal distortion constants) are required to reproduce the results of the fit within the experimental uncertainty.

^bFixed at ground state values from Ref. [11, 23].

^cExtrapolated from the corresponding constants of 0^0 and 2^1

Table 3. Range of quantum numbers observed for the experimental rotational energy levels of the 2^1 and 2^2 vibrational states of sulfur dioxide $^{33}\text{S}^{16}\text{O}_2$.

Table 1.

Watson-type Hamiltonian:

$$\begin{aligned}
 H_{\text{W}} = E_{\text{v}} &+ \left[A^{\text{v}} - \frac{1}{2} (B^{\text{v}} + C^{\text{v}}) \right] J_z^2 + \frac{1}{2} (B^{\text{v}} + C^{\text{v}}) J^2 + \frac{1}{2} (B^{\text{v}} - C^{\text{v}}) J_{\text{xy}}^2 \\
 &- \Delta_{\text{K}}^{\text{v}} J_z^4 - \Delta_{\text{JK}}^{\text{v}} J_z^2 J^2 - \Delta_{\text{J}}^{\text{v}} (J^2)^3 - \delta_{\text{K}}^{\text{v}} \{ J_z^2, J_{\text{xy}}^2 \} - 2\delta_{\text{J}}^{\text{v}} J_{\text{xy}}^2 J^2 \\
 &+ H_{\text{K}}^{\text{v}} J_z^6 + H_{\text{KJ}}^{\text{v}} J_z^4 J^2 + H_{\text{JK}}^{\text{v}} J_z^2 (J^2)^2 + H_{\text{J}}^{\text{v}} (J^2)^3 \\
 &+ h_{\text{K}}^{\text{v}} \{ J_z^4, J_{\text{xy}}^2 \} + h_{\text{KJ}}^{\text{v}} \{ J_z^2, J_{\text{xy}}^2 \} J^2 + 2h_{\text{J}}^{\text{v}} J_{\text{xy}}^2 (J^2)^2 + \dots
 \end{aligned}$$

with $\{A, B\} = AB + BA$, $J_{\pm} = J_x \mp iJ_y$ and $J_{xy}^2 = J_x^2 - J_y^2$.

Table 2.

	0⁰	2¹	2²
E_v/hc		515.659089(45)	1030.697723(20)
A	1.996595277(660) ^a	2.03519127(120)	2.0755724155(510)
B	0.344181308(130)	0.344255311(140)	0.3443181510(720)
C	0.292873783(120)	0.292341697(120)	0.2918044548(760)
$\Delta_K \times 10^4$	0.83801438(5200)	0.92860380(9100)	1.03008145(3000)
$\Delta_{JK} \times 10^5$	-0.38180636(9300)	-0.3984898(1000)	-0.41622140(7000)
$\Delta_J \times 10^6$	0.21976967(4000)	0.22026261(5500)	0.220776116(4500)
$\delta_K \times 10^6$	0.837348(2400)	1.025294(1200)	1.2247786(9700)
$\delta_J \times 10^7$	0.5708293(1400)	0.5744184(3300)	0.577717(1000)
$H_K \times 10^7$	0.1190711(1000)	0.1454599(2400)	0.17619965(5000)
$H_{KJ} \times 10^9$	-0.671247(4300)	-0.768561(6500)	-0.866080(2000)
$H_{JK} \times 10^{12}$	2.214621415 ^b	4.605(1500)	6.995 ^c
$H_J \times 10^{13}$	3.827390459 ^b	3.86274(8800)	4.06420(9400)
$h_K \times 10^9$	0.74134(1400)	0.89564(6500)	1.06534(1100)
$h_{KJ} \times 10^{13}$	-1.427642403 ^b	-1.427642403 ^b	-1.427642403 ^b
$h_J \times 10^{13}$	1.795770464 ^b	1.85734(6300)	1.91891 ^c
$L_K \times 10^{12}$	-2.635612983 ^b	-3.21188(2000)	-3.78815 ^c
$L_{KKJ} \times 10^{13}$	1.444367432 ^b	1.57021(8300)	1.69605 ^c
$L_{KJ} \times 10^{16}$	8.466570753 ^b	8.466570753 ^b	8.466570753 ^b
$L_{JK} \times 10^{17}$	-8.243466351 ^b	-8.243466351 ^b	-8.243466351 ^b
$L_J \times 10^{18}$	-1.331348331 ^b	-1.331348331 ^b	-1.331348331 ^b
$l_{JK} \times 10^{17}$	6.855534121 ^b	6.855534121 ^b	6.855534121 ^b
$l_J \times 10^{19}$	-8.610696124 ^b	-8.610696124 ^b	-8.610696124 ^b
$P_K \times 10^{16}$	6.531343786 ^b	6.531343786 ^b	6.531343786 ^b
$P_{KKKJ} \times 10^{17}$	-3.393376689 ^b	-3.393376689 ^b	-3.393376689 ^b
$P_{KKJ} \times 10^{17}$	3.800968760 ^b	3.800968760 ^b	3.800968760 ^b
$R_{K12} \times 10^{19}$	-1.208536717 ^b	-1.208536717 ^b	-1.208536717 ^b

Vibrational State	2 ¹	2 ²
Number of assigned lines	4650	2763
Number of levels	1146	813
J _{max}	73	63
K _{max}	24	21
Std. Deviation (10 ⁻³ cm ⁻¹)	0.15	0.17