**Synthesis, Structures, and Catalytic Reactivity of Bis(N-Heterocyclic Carbene) Supported
Diphenyldiazomethane and 1-Azidoadamantane Complexes of Nickel**

Nicole D. Harrold,* Amanda R. Corcos,[‡] and Gregory L. Hillhouse[⊥]

*Gordon Center for Integrative Science, Department of Chemistry, The University of Chicago,
929 East 57th Street, Chicago, Illinois, 60637, USA*

* Email: nharrold@uchicago.edu

ABSTRACT.

Diphenyldiazomethane reacts with $\text{Ni}(\text{NHC})_2$ (**1**, NHC = N-heterocyclic carbene) to afford complexes of the type $(\text{NHC})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ (**3**) in which the diazoalkane ligand is bound to the metal in an end-on κ^1 fashion. Additionally, **1-IMes** (IMes = 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene) reacts with 1-azidoadamantane (N_3Ad) to produce $(\text{IMes})_2\text{Ni}(\eta^2\text{-N}_3\text{Ad})$ (**4**) in which the azide ligand is bound to the metal in a side-bound η^2 -fashion. While thermolysis of **3** and **4** does not induce N_2 loss to form terminal carbenes and imidos, respectively, $(\text{IMes})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ (**3-IMes**) reacts with olefins at elevated temperatures to give cyclopropane products in excellent yields. Either **3-IMes** or **1-IMes** can also perform this reaction catalytically with a 1:1 ratio of diphenyldiazomethane to olefin. An olefin activation mechanism is proposed to explain the cyclopropanation reaction.

GRAPHICAL ABSTRACT.

Complexes of type $(\text{NHC})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ (**3**), in which diphenyldiazomethane is bound to an N-heterocyclic carbene-supported nickel center in an unusual κ^1 end-on fashion, react with olefins at elevated temperatures to give cyclopropane products in excellent yields. **3-IMes** or **1-IMes** can also perform this reaction catalytically with an efficient 1:1 ratio of diphenyldiazomethane to olefin.

KEYWORDS.

1. Nickel
2. *N*-heterocyclic carbene (NHC)
3. Catalysis
4. Cyclopropanation
5. Diazoalkane
6. Organoazide

1. INTRODUCTION.

Diazoalkanes are commonly used precursors for the synthesis of metallocarbenes,^[1-3] which in turn are versatile chemical moieties capable of olefin cyclopropanation, C–H activation, and other reactions.^[4-12] While N₂ elimination from diazoalkanes is generally facile upon reaction with metal complexes, the isolation and characterization of metal-bound diazoalkane adducts is possible and has yielded a plethora of complexes spanning multiple coordination modes.^[6, 7, 13, 14] Five coordination modes are possible for a diazoalkane moiety bound to a single transition metal center (Chart 1): κ^1 -N end-on coordination involving the terminal N (type **A**),^[15-22] η^2 -N,N side-on (type **B**),^[23-28] η^2 -C,N side-on (type **C**), κ^2 -C,N metallacyclic (type **D**), and κ^1 -C coordination (type **E**). Coordination of a diazoalkane to more than one metal atom is also possible.^[29]

Several limiting electronic configurations for the κ^1 -N binding configuration (**A**) are shown in Chart 2. In configuration **A**₁, the diazoalkane ligand serves as a neutral 2-electron donor with a dative interaction between the terminal nitrogen and the metal center. The diazoalkane ligands in configurations **A**₂ (singly bent) and **A**₃ (doubly bent) are bound to the metal center through two covalent bonds ($\sigma + \pi$). Configuration **A**₄ features two covalent bonds and one dative bond.

Chart 1. Diazoalkane binding modes to a single metal center.

Chart 2. Electronic configurations for a κ^1 -N end-on diazoalkane bound to a single metal center.

In a similar manner, organoazides are often used as imido precursors,^[30-32] which can undergo group transfer and C-H amination after reacting with a metal center to induce loss of N_2 .^[33-37] There are several possible coordination modes for organoazides to bind to a metal center. In the few instances where these complexes are stable enough for isolation, the ligand is usually bound as a diazenylimido (RN_3^{2-} as **F**, **G**, or **H** in Chart 3)^[38-42] or as a neutral RN_3 through the α or γ N-atoms (**I** and **J**).^[43-45]

Chart 3. Common organoazide binding modes to a metal center.

Herein we report the synthesis, structures, and reactivity of several new bis(N-heterocyclic carbene) (NHC) supported diphenyldiazomethane complexes of Ni in which the diazo ligand is bound to the metal in a κ^1 end-on fashion (**A**). These complexes mediate cyclopropane formation in high yields from stoichiometric and catalytic 1:1 mixtures of diazoalkanes and olefins, whereas typical cyclopropanation catalysis requires an excess of olefin for good selectivity and high yields. It is proposed that catalysis for this system occurs without the intermediacy of a metallacarbene. The synthesis and reactivity of a new bis(NHC) supported organoazide complex of Ni, in which the azide ligand is bound to the metal in an η^2 -fashion, is also reported and its reactivity discussed.

2. MATERIALS AND METHODS.

2.1. General Considerations. Unless stated otherwise, all operations were performed in an MBraun Lab Master dry box under an atmosphere of purified nitrogen. Anhydrous diethyl ether (Et₂O) and tetrahydrofuran (THF) were purchased from Fischer, stirred over sodium metal, and filtered through activated alumina.^[46] Pentane was purchased from Sigma Aldrich and dried by passage through activated alumina and Q-5 columns. C₆D₆ was purchased from Cambridge Isotope Laboratories, degassed by freeze-pump-thaw cycles, and dried over CaH₂ or activated 4 Å molecular sieves. Celite, alumina, and 4 Å molecular sieves were activated by evacuation overnight at 180 °C. N₂CPh₂,^[47] N₃Ad,^[48-50] IMes,^[51-53] SIMes,^[51] IPr,^[51, 52, 54] and Ni(IMes)₂^[55] were prepared according to the literature. All other chemicals were used as received. Please note: caution should be used when conducting photolytic and thermolytic experiments on diazoalkane and organoazide species, for such species have the potential to be explosive. Due to the small scale on which these reactions were performed and the low temperatures utilized, all reactions performed here are deemed safe.

2.2. Instruments. Elemental analyses were performed by Midwest Microlab (Indianapolis, IN). ¹H NMR spectra were recorded on a Bruker 500 MHz NMR spectrometer and reported with reference to solvent resonance (for example, residual C₆D₅H in C₆D₆ at 7.16 ppm). Infrared data (Fluorolube or Nujol mulls, CaF₂ or KBr plates, respectively) were measured using a Nicolet Nexus 670 FT-IR.

2.3. Synthesis.

2.3.1. Preparation of (NHC)₂Ni(κ¹-N₂CPh₂) (3). A typical procedure is given for the synthesis of **3-IMes**. A cold (-35 °C) solution of 145 mg (750 μmol) N₂CPh₂ in 5 mL diethyl ether was added to a cold stirring solution of 500 mg (750 μmol) Ni(IMes)₂ (**1-IMes**) in 10 mL diethyl ether. The reaction was stirred at room temperature for 1 hour after which the reaction mixture was reduced to dryness. The resulting solids were then washed with cold hexanes (3 x 5 mL). The resulting orange red solids were dried under vacuum for 2 hours. The characterization data for **3-IMes**, **3-SIMes**, and **3-IPr** are given below. All complexes are stable in the freezer (-35 °C) for up to 6 months but degrade if left at room temperature or exposed to light for more than one week.

2.3.1.1. $(IMes)_2Ni(\kappa^1-N_2CPh_2)$ (**3-IMes**). Yield: 620 mg (96%). 1H NMR (298 K, 500.13 MHz, C_6D_6): δ 7.68 (d, 4H, o- C_6H_6), 7.19 (t, 4H, m- C_6H_6), 6.86 (t, 2H, p- C_6H_6), 6.69 (s, 8H, C_6H_2), 6.02 (s, 4H, NC_2H_2N), 2.28 (s, 12H, $C_6H_2(CH_3)_3$), 1.87 (s, 24H, $C_6H_2(CH_3)_3$); $^{13}C\{^1H\}$ NMR (298 K, 125.76 MHz, C_6D_6): δ 199.8, 138.8, 138.4, 137.7, 136.4, 129.8, 128.3, 125.8, 123.6, 121.7, 21.9, 18.9. IR (Nujol, KBr): 3153 (m), 2915 (m), 2041 (s), 1703 (s), 1608 (s), 1560 (s), 1487 (s), 1447 (s), 1405 (s) cm^{-1} . Anal. Calcd. for $C_{55}H_{58}NiN_6$: C, 76.75; H, 6.78; N, 9.75. Found: C, 74.71; H, 6.30; N, 9.06. Please note these values are not within the standard range due to (unavoidable) decomposition of the complex during shipment to the EA facilities. All other characterizations and catalytic reactions presented here were performed with freshly-made (non-decomposed) materials.

2.3.1.2. $(SIMes)_2Ni(\kappa^1-N_2CPh_2)$ (**3-SIMes**). Yield: 595 mg (92%). 1H NMR (298 K, 500.13 MHz, C_6D_6): δ 7.71 (d, 4H, o- C_6H_6), 7.33 (t, 4H, m- C_6H_6), 6.99 (t, 2H, p- C_6H_6), 6.89 (d, 8H, C_6H_3), 6.76 (t, 4H, C_6H_3), 3.15 (s, 8H, NC_2H_4N), 6.69 (s, 8H, C_6H_2), 2.27 (s, 12H, $C_6H_2(CH_3)_3$), 1.98 (s, 24H, $C_6H_2(CH_3)_3$); $^{13}C\{^1H\}$ NMR (298 K, 125.76 MHz, C_6D_6): δ 138.9, 138.6, 137.2, 136.7, 130.3, 125.9, 121.8, 51.8, 21.9, 18.8. IR (Nujol, KBr): 2810 (m), 2043 (s), 1713 (s), 1606 (s), 1563 (s), 1487 (s), 1448 (s), 1410 (s) cm^{-1} .

2.3.1.3. $(IPr)_2Ni(\kappa^1-N_2CPh_2)$ (**3-IPr**). Yield: 631 mg (82%). 1H NMR (298 K, 500.13 MHz, C_6D_6): δ 7.65 (d, 4H, o- C_6H_6), 7.58 (t, 4H, m- C_6H_6), 7.22 (t, 2H, p- C_6H_6), 6.04 (s, 4H, NC_2H_2N), 2.23 (m, 8H, $CH(CH_3)_2$), 1.31 (d, 24H, $CH(CH_3)_2$), 1.07 (d, 24H, $CH(CH_3)_2$); $^{13}C\{^1H\}$ NMR (298 K, 125.76 MHz, C_6D_6): δ 204.8, 142.8, 137.4, 137.3, 134.3, 127.8, 126.4, 126.3, 125.8, 123.6, 121.7, 21.9, 18.9, 18.7. IR (Nujol, KBr): 2805 (m), 2042 (s), 1713 (s), 1605 (s), 1559 (s), 1484 (s), 1448 (s), 1410 (s) cm^{-1} .

2.3.2. *Stoichiometric Procedure for Formation of Cyclopropane*. A typical procedure is given for the formation of 1,2-diphenyl-2-butylcyclopropane. In a 25 mL glass bomb equipped with a stir bar was loaded **3-IMes** (100 mg, 0.116 mmol) and 10 mL toluene. To the solution was added 1-hexene (9.8 mg, 0.116 mmol) dissolved in 1 mL of toluene at 22 °C. The solution remained red in color upon addition. Upon heating (2 hours, 60 °C) the solution turned dark purple in color. After completion the solution was exposed to air and filtered through celite. The filtrate was then passed through a short SiO_2 column (1:1 hexanes: diethyl ether). The yellow band was collected and dried under reduced pressure for 2 hours to afford pure cyclopropane. The characterization data for cyclopropane products are given below.

2.3.2.1. *1,2-diphenyl-2-butylcyclopropane*. Yield: 26.1 mg (90%). ^1H NMR (298 K, 500.13 MHz, CDCl_3): δ 7.35 – 7.19 (m, 10H, Ph), 1.61 (m, 1H, C_4H_9), 1.43 (m, 2H, C_4H_9), 1.32 (m, 1H, C_4H_9), 1.25 (4 line, 1H, CH_2 , $\text{J}_{\text{HH}} = 8.8, 4.6$ Hz), 1.19 (4 line, 1H, CH_2 , $\text{J}_{\text{HH}} = 4.6, 6.3$ Hz), 0.85 (t, 3H, C_4H_9), 0.74 (m, 1H, C_4H_9); $^{13}\text{C}\{^1\text{H}\}$ NMR (298 K, 125.76 MHz, CDCl_3): δ 130.8, 128.6, 126.2, 125.7, 35.4, 31.8, 29.9, 26.6, 22.7, 21.1, 11.7; GC/MS (m/z): 250 (M^+), 193 (M-Bu), 165 ($\text{Ph}_2\text{C}^+-\text{H}$).

2.3.2.2. *1,1-diphenyl-2,2,3-trimethylcyclopropane*. Yield: 24.7 mg (90%). ^1H NMR (298 K, 500.13 MHz, CDCl_3): δ 7.27 – 6.95 (m, 10H, Ph), 1.34 (s, 3H, CH_3), 1.02 (s, 3H, CH_3), 1.00 (s, 3H, CH_3); $^{13}\text{C}\{^1\text{H}\}$ NMR (298 K, 125.76 MHz, CDCl_3): δ 146.1, 129.2, 128.2, 128.1, 127.4, 126.2, 126.1, 46.0, 33.7, 26.4, 22.4, 15.2; GC/MS (m/z): 236 (M^+), 221 (M^+-Me), 143 ($\text{M}^+-\text{Me-Ph}$).

2.3.2.3. *7,7-diphenylbicyclo[4.1.0]heptane*. Yield: 24.8 mg (86%). ^1H NMR (298 K, 500.13 MHz, CDCl_3): δ 7.35 – 7.01 (m, 10H, Ph), 2.30 – 0.62 (m, 10H, C_4H_8); $^{13}\text{C}\{^1\text{H}\}$ NMR (298 K, 125.76 MHz, CDCl_3): δ 149.7, 140.1, 131.6, 128.5, 127.8, 36.2, 25.2, 22.3, 21.4; GC/MS (m/z): 250 (M^+), 205 ($\text{M}^+-\text{C}_3\text{H}_6$), 165 ($\text{Ph}_2\text{C}^+-\text{H}$).

2.3.2.4. *1,1-diphenyl-2-butyl-3-methylcyclopropane*. Yield: 23.2 mg (80%). ^1H NMR (298 K, 500.13 MHz, CDCl_3): δ 7.28 – 7.01 (m, 10H, Ph), 1.64 – 1.39 (m, 7H), 1.08 – 1.04 (m, 3H), 0.96 – 0.91 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (298 K, 125.76 MHz, CDCl_3): δ 147.4, 147.3, 129.7, 129.5, 128.2, 128.1, 126.4, 126.1, 56.0, 35.1, 34.1, 24.6; GC/MS (m/z): 250 (M^+), 207 (M-Pr).

2.3.3. *Catalytic Procedure for 1,2-diphenyl-2-butylcyclopropane*. In a 25 mL glass bomb equipped with a stir bar was loaded $\text{Ni}(\text{IMes})_2$ (**1-IMes**) (205 mg, 0.238 mmol, 20 mol %) and 10 mL toluene. To the solution was added 1-hexene (100 mg, 1.19 mmol) and N_2CPh_2 (231 mg, 1.19 mmol) dissolved in 1 mL of toluene at 22 °C. The solution remained red in color upon addition. Upon heating (12 hours, 60 °C) the solution turned dark purple in color. After completion the solution was exposed to air and filtered through celite. The filtrate was then passed through a short SiO_2 column (1:1 hexanes: diethyl ether). The yellow band was collected and dried under reduced pressure for 2 hours to afford pure cyclopropane (250 mg, 84%).

2.3.4. *Preparation of $(\text{IMes})_2\text{Ni}(\eta^2\text{-N}_3\text{Ad})$ (**4**)*. A cold, 5 mL THF solution of N_3Ad (26.6 mg, 0.150 mmol) was added to a cold, 10 mL solution of **1-IMes** (100 mg, 0.150 mmol). The solution was stirred at room temperature for 1 hour. The resulting solution was reduced to dryness under reduced pressure. The crude light brown solid was then washed with cold hexanes (2

x 5 mL) resulting in a light tan solid, from which crystals were grown after concentration in cold toluene and layering with hexanes (68 mg, 55%). This complex is stable in the freezer (-35 °C) for up to 3 months but degrades if left at room temperature or exposed to light for more than one week. ¹H NMR (298 K, 500.13 MHz, C₆D₆): δ 6.69 (s, 8H, C₆H₂), 6.09 (s, 4H, NC₂H₂N), 2.29 (s, 12H, C₆H₂(CH₃)₃), 2.23 (s, 4H, Ad), 2.19 (s, 6H, C₆H₂(CH₃)₃), 2.03 (s, 12H, C₆H₂(CH₃)₃), 1.91 (d, 5H, Ad), 1.76 (d, 5H, Ad), 1.41 (s, 2H, Ad). ¹³C{¹H} NMR (298 K, 125.76 MHz, C₆D₆): δ 204.3, 137.7, 137.2, 137.1, 135.1, 132.7, 128.5, 126.4, 118.0, 51.0, 38.7, 36.5, 2.4, 21.9, 17.2, 17.1; IR (Nujol, KBr): 1349 (m), 1283 (w), 1181 (s), 1149 (s), 1116 (s), 1099 (s), 1078 (m), 1044 (w), 1022 (m), 938 (m), 898 (w), 845 (s), 813 (s), 783 (s), 724 (w), 686 (s), 657 (s), 582 (m), 496 (s) cm⁻¹. Anal. Calcd for C₅₂H₆₃NiN₇: C, 73.93; H, 7.52; N, 11.61. Found: C, 71.40; H, 7.37; N, 10.05. Please note these values are not within the standard range due to (unavoidable) decomposition of the complex during shipment to the EA facilities. All other characterizations and catalytic reactions presented here were performed with freshly-made (non-decomposed) materials.

2.3.5. Synthesis of (IMes)₂Ni(η^2 -C₆H₁₂) (5). A solution of **1-IMes** (10.0 mg) was put in an NMR tube and dissolved in C₆D₆. To the NMR tube was added 1-hexene (28 mg, 0.33 mmol) via syringe. Immediately the color changed from purple to red. The NMR was taken immediately. The NMR tube was heated at 60 °C and no visible or spectroscopic change occurred even after prolonged (12 h) in the oil bath. The ¹H NMR (shown below) of complex **5** was obtained in a C₆D₆ solution with excess (~ 40 equivalents) 1-hexene. Due to the large excess of 1-hexene the ¹³C shifts of **5** could not be obtained. Attempts to isolate complex **5** on a preparatory scale were unsuccessful as 1-hexene dissociates from the metal upon solvent removal. ¹H NMR (298 K, 500.13 MHz, C₆D₆): δ 6.72 (s, 8H, C₆H₂), 6.72 (m, 1H, 1-hex), 6.69 (m, 1H, 1-hex), 6.33 (m, 1H, 1-hex), 6.11 (s, 4H, NC₂H₂N), 2.26 (s, 24H, C₆H₂(CH₃)₃), 2.20-1.96 (m, 9H, 1-hex), 2.07 (s, 12H, C₆H₂(CH₃)₃).

2.4. Crystal Structure Determination. Data Collection and Structure Solution and Refinement. Suitable crystals of **3-Mes**, **3-SIMes**, **3-IPr**, and **4** were selected under a stereomicroscope while they were immersed in mineral oil to minimize contact with air or water. The crystal was removed from the oil using a tapered fiber, which also served to hold the crystal for data collection, and was mounted in a stream of cold nitrogen at 100(2) K. X-ray diffraction data were collected on a Bruker SMART APEX diffractometer with Mo K α (λ = 0.71073 Å) radiation. A Siemens Platform goniometer and a Charged Coupled Device (CCD) detector were used. The

data were collected using a routine to survey the reciprocal space to the extent of a full sphere to a resolution of 0.75 Å and were indexed by the APEX program.^[56] Rotation and still images showed diffractions to be sharp. Frames separated in reciprocal space were obtained and provided an orientation matrix and the initial cell parameters. Final cell parameters were obtained from the full data set. Integration of intensities and refinement of cell parameters were done using SAINT. The structures were solved by direct methods refined by iterative cycles of least-squares refinement on F^2 followed by difference Fourier synthesis using the XS software.^[57, 58] All non-hydrogen atoms were refined anisotropically. All hydrogen atoms were included in the final structure factor calculation at idealized positions and were allowed to ride on the neighboring atoms with relative isotropic displacement coefficients. Absorption corrections were based on a fitted function to the empirical transmission surface as sampled by multiple equivalent measurements.^[59] Observation during and after data collection showed that the crystal was still clear, suggesting no decomposition.

2.4.1. Crystal Structure of 3-IMes·2(C₇H₈). Red block crystals suitable for X-ray diffraction were obtained from a concentrated toluene solution left at -35 °C for 7 days. The systematic absences in the diffraction data were uniquely consistent for the space group $P2_1/c$. Two molecules of toluene are present in the unit cell.

2.4.2. Crystal Structure of 3-SIMes. Red block crystals suitable for X-ray diffraction were obtained from a concentrated toluene solution layered with hexanes. The systematic absences in the diffraction data were uniquely consistent for the space group $P2_1$. The structure was refined as an inversion twin, and the resulting Flack X parameter confirms we have the correct handedness.

2.4.3. Crystal Structure of 3-IPr·Et₂O. Red block crystals suitable for X-ray diffraction were obtained from a concentrated diethyl ether solution at -35 °C. The systematic absences in the diffraction data were uniquely consistent for the space group $P-1$. One diethyl ether molecule is present in the unit cell.

2.4.4. Crystal Structure of 4. Yellow needle crystals suitable for X-ray diffraction were obtained from a concentrated toluene solution layered with hexanes left at -35 °C for 7 days. The systematic absences in the diffraction data were uniquely consistent for the space group $P-1$. The unit cell contains 1.72 toluene molecules, which have been treated as a diffuse contribution to the overall scattering without specific atom positions by SQUEEZE/PLATON.^[60]

3. RESULTS AND DISCUSSION.

3.1. Reaction of Ni(NHC)₂ with Diphenyldiazomethane. A series of biscarbene Ni(0) complexes were synthesized using the carbenes IMes, SIMes and IPr (IMes = 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene, SIMes = 1,3-bis(2,4,6-trimethylphenyl)imidazolin-2-ylidene and IPr = 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene) (Scheme 1).^[61] The reactions of these homoleptic complexes with diazoalkanes were investigated to study the possibility of performing group transfer chemistry.

Scheme 1. NHC ligands utilized in this work.

Ni(IMes)₂ (**1-IMes**) was first prepared in 1994 by Arduengo, who took advantage of the bulky nature of IMes to prepare one of the first known two-coordinate homoleptic transition metal complexes (Scheme 2).^[55] This thermally stable, 14-electron Ni(0) complex is prepared as dark purple crystals by reaction of Ni(cod)₂ (cod = 1,5-cyclooctadiene) with 2 equivalents of the IMes ligand in tetrahydrofuran (THF). We found that this route also provides the analogous new complex Ni(SIMes)₂ (**1-SIMes**) in 92% yield (Scheme 2).

Scheme 2. Synthetic formation of **1-IMes** and **1-SIMes**.

In contrast, reaction of Ni(cod)₂ and 2 equivalents of IPr ligand forms an equilibrium between Ni(cod)₂ and Ni(IPr)₂ (**1-IPr**).^[62] Only a 30% yield of **1-IPr** could be isolated from this mixture in our hands. We therefore developed a new route to generate **1-IPr** in good yield (Scheme 3). The known complex (IPr)₂NiCl₂, **2**, was prepared as a pale pink powder in 95% yield from IPr and (PPh₃)₂NiCl₂.^[63] Further reaction of **2** with 2.1 equivalents of KC₈ affords **1-IPr** as a purple-black powder. This method provides **1-IPr** in better yield than NaH reduction starting from Ni(acac)₂^[64] (acac = acetylacetonate) and in comparable yield to reductive elimination of ethane from (IPr)₂NiMe₂ without the necessity of isolating the sensitive Ni-alkyl intermediate.^[65]

Scheme 3. Synthetic formation of **1-IPr**.

Treatment of $\text{Ni}(\text{NHC})_2$ (**1-Mes**, **1-SIMes**, and **1-Pr**) with 1 equivalent of diphenyldiazomethane at $-35\text{ }^\circ\text{C}$ followed by stirring at room temperature for 1 hour in diethyl ether affords $(\text{NHC})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ (**3-IMes**, **3-SIMes**, and **3-IPr**). Removal of the ether under vacuum followed by washing with cold hexanes affords **3-IMes**, **3-SIMes**, and **3-IPr** as red or orange-red solids in high yields (Scheme 4).

3.2. Structural Analysis of $(\text{NHC})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ Complexes. The X-ray structures of complexes **3-IMes**, **3-SIMes**, and **3-IPr** were investigated to determine how the diazoalkane ligand interacts with the Ni center. Crystals suitable for X-ray analysis of **3-IMes** (Figure 1) were obtained from a concentrated toluene solution cooled to $-35\text{ }^\circ\text{C}$ while crystals of **3-SIMes** (Figure 2) were obtained from a concentrated toluene solution layered with hexanes at room temperature. Crystals of **3-IPr** (Figure 3) were obtained from a concentrated diethyl ether solution at $-35\text{ }^\circ\text{C}$.

Scheme 4. Synthetic formation of **3-IMes**, **3-SIMes**, and **3-IPr** from their respective $\text{Ni}(\text{NHC})_2$ precursors.

Figure 1. Thermal ellipsoid plot of $(\text{IMes})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2) \cdot 2(\text{C}_7\text{H}_8)$ (**3-IMes**· $2(\text{C}_7\text{H}_8)$) drawn at 50% probability. H atoms and molecules of solvation omitted for clarity.

Figure 2. Thermal ellipsoid plot of $(\text{SIMes})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ (**3-SIMes**) drawn at 50% probability. H atoms omitted for clarity.

Figure 3. Thermal ellipsoid plot of $(\text{IPr})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)\cdot\text{Et}_2\text{O}$ (**3-IPr·Et₂O**) drawn at 50% probability. H atoms and molecules of solvation omitted for clarity.

Table 1. Selected bond lengths (Å) and angles (°) for **3**.

	3-IMes	3-SIMes	3-IPr
Ni–N(1) (Å)	1.721(2)	1.728(4)	1.748(2)
N(1)–N(2) (Å)	1.231(3)	1.208(5)	1.219(2)
N(2)–C(1) (Å)	1.349(4)	1.338(5)	1.339(3)
Ni–C(2) (Å)	1.942(3)	1.921(4)	1.984(2)
Ni–C(3) (Å)	1.907(3)	1.910(4)	1.914(2)
Ni–N(1)–N(2) (°)	161.5(2)	159.1(3)	147.2(2)
N(1)–N(2)–C(1) (°)	137.7(3)	139.1(4)	139.8(2)
C(2)–Ni–C(3) (°)	114.7(1)	123.8(2)	142.80(9)

The crystal structures of all three complexes reveal planar, three-coordinate Ni centers in which the N₂CPh₂ ligand is coordinated in an end-on κ^1 fashion to the metal. The angle between the NHC ligands [C(2)–Ni–C(3): **3-IMes** 114.7(1)° < **3-SIMes** 123.8(2)° < **3-IPr** 142.80(9)°] lie in the order consistent with the relative bulk of the three ligands as measured by percent buried volume.^[66, 67] The Ni–N(1) bond lengths of the complexes (1.721 – 1.7482 Å) fall between the experimentally obtained value for a Ni–N single bond distance of 1.881(2) Å and the Ni=N double bond distance of 1.702(2) Å as observed in (dtbpe)Ni–NH(2,6-ⁱPrC₆H₃) and (dtbpe)Ni=N(2,6-ⁱPrC₆H₃) (dtbpe = 1,2-bis(di-tert-butylphosphino)ethane), respectively.^[68] The N(1)–N(2) bond lengths for the complexes (1.208 – 1.231 Å) are closer to typical N=N double bonds (~1.24 Å) than typical N–N single bonds (~1.45 Å).^[69] The N(2)–C(1) bond lengths for the complexes (1.338 – 1.349 Å) are comparable to typical partial double bonds found in pyridine (1.352 Å).^[70] This lengthening (and weakening) of what would be expected for a C=N double bond, is consistent with the observed reduction in $\nu(\text{C}=\text{N})$ upon diazoalkane binding to a metal center, from 2032 cm⁻¹ for free N₂CPh₂^[71] to 1560 cm⁻¹ observed for each of these three complexes. The Ni–N(1)–N(2) (147.2 – 161.5°) and N(1)–N(2)–C(1) (137.7 – 139.8°) bond angles show significant deviations from linearity. Collectively, the metrical data for **3-IMes**, **3-SIMes**, and **3-IPr** are consistent with a bonding mode that is intermediate between **A**₂ and **A**₃.

The κ^1 coordination of N₂CPh₂ observed for **3-IMes**, **3-SIMes** and **3-IPr** contrasts with that observed for L₂Ni(N₂CR₂) complexes with smaller L ligands. Radius reported that when L is ⁱPr₂Im (ⁱPr₂Im = 1,3-diisopropylimidazol-2-ylidene), then N₂CPh₂ forms an η^2 -N,N complex of type **B** (Figure 1).^[27] The smaller size of ⁱPr₂Im allows the C–Ni–C angle to contract to 104.8(2)° (cf. 114.7 – 142.8° for **3**), which affords room for the side-on binding of the diazoalkane. Otsuka and Ibers have also crystallographically characterized a similar complex where Ni(CN^tBu)₂ binds diazafluorene in a type-**B** η^2 -N,N fashion.^[23] Here again the C–Ni–C angle of 100.4(2)° is significantly smaller than those seen in **3**.

3.3. Reaction and structural analysis of Ni(IMes)₂ with 1-Azidoadamantane. The observation that the (NHC)₂Ni(κ^1 -N₂CPh₂) complexes **3-IMes**, **3-SIMes**, and **3-IPr** are resistant to elimination of N₂ and thus do not form carbene complexes (vide infra) suggested that analogous azide species could exhibit similar properties. 1-Azidoadamantane (N₃Ad) was chosen because it is capable of many of the same binding modes as diphenyldiazomethane yet possesses steric bulk further from the terminal N=N moiety.

The reaction of N_3Ad with **1-IMes** in tetrahydrofuran yields a yellow solution (Scheme 5). No gas evolution was observed. From this solution, after concentration and layering with hexanes, crystals of **4** were isolated. An X-ray diffraction study of **4** revealed an η^2 -azide adduct coordinated through the terminal N atoms (Figure 4). The C–Ni–C angle in **4** of $120.1(1)^\circ$ is even larger than that seen in **3-IMes** ($114.7(1)^\circ$). The ^1H NMR spectrum reveals a diamagnetic compound that is fluxional in solution.

Scheme 5. Synthetic formation of **4** from **1-IMes**.

Figure 4. Thermal ellipsoid plot of $(\text{IMes})_2\text{Ni}(\eta^2\text{-N}_3\text{Ad})$ (**4**) drawn at 50% probability. H-atoms omitted for clarity. Select bond lengths (Å) and angles ($^\circ$): Ni(1)–N(1) = 1.926(3), Ni(1)–N(2) = 1.826(3), N(1)–N(2) = 1.234(4), N(2)–N(3) = 1.311(4), N(3)–C(1) = 1.487(5), Ni(1)–C(2) = 1.914(4), Ni(1)–C(3) = 1.918(4), C(2)–Ni(1)–C(3) = $120.1(1)$, N(1)–Ni(1)–N(2) = $38.3(1)$, N(1)–N(2)–N(3) = $139.9(3)$, N(2)–N(3)–C(1) = $114.6(3)$.

3.4. Reactivity of $(\text{NHC})_2\text{Ni}(\kappa^1\text{-N}_2\text{CPh}_2)$ Complexes. The utility of **3-IMes**, **3-SIMes** and **3-IPr** as precursors for Ni-carbene complexes was explored by studying their thermal and photochemical stability. Complexes **3-IMes**, **3-SIMes**, and **3-IPr** are remarkably robust under an inert

atmosphere in benzene up to 60 °C and do not undergo N₂ loss under thermal or photochemical conditions. At 60 °C, however, the diazoalkane ligand dissociates from the metal center to regenerate **1** cleanly.

Computational studies suggest that the formation of a carbene complex from a diazoalkane must involve the intermediacy of a κ^2 -C,N metallacycle (type **D**, Chart 1) or κ^1 -C diazoalkane intermediate (type **E**, Chart 1).^[72] Such a binding motif is unlikely in these complexes because of steric constraints imposed by the bulky NHC ligands. Space-filling models of **3-IMes** and **3-IPr** (Figure 5), where N₂CPh₂ is coordinated to the Ni center in the κ^1 -N binding mode, show considerable steric interaction between the phenyl substituents of N₂CPh₂ and the aryl substituents of the NHC ligand. This interaction would only be further exacerbated in κ^1 -C or κ^2 -C,N binding modes.

Figure 5. Space-filling models of (IMes)₂Ni(κ^1 -N₂CPh₂) (**3-IMes**, left) and (IPr)₂Ni(κ^1 -N₂CPh₂) (**3-IPr**, right).

While the (NHC)₂Ni(κ^1 -N₂CPh₂) complexes **3** do not eliminate N₂ to form stable carbenes, they do undergo stoichiometric and catalytic group transfer with alkenes to give cyclopropane products (Table 2). Treatment of a toluene solution of **3-IMes** with 1-hexene (1 equiv, 60 °C, 2 h) affords **1-IMes** and 1,1-diphenyl-2-butylcyclopropane in 90% yield, while reaction with 1-methyl-2-butene, 2-hexene, or cyclohexene provides the corresponding cyclopropane products in good yield.

Table 2. Stoichiometric group transfer reactions of (IMes)₂Ni(κ^1 -N₂CPh₂) (**3-IMes**).

$$\mathbf{3-IMes} + \begin{array}{c} R_2 \\ \diagup \quad \diagdown \\ R_1 \quad C = C \\ \diagdown \quad \diagup \\ R_3 \end{array} \xrightarrow[60^\circ\text{C}]{2\text{ hr}, -N_2} \begin{array}{c} \text{Ph} \quad \text{Ph} \\ \diagup \quad \diagdown \\ \text{C} \\ \diagdown \quad \diagup \\ R_1 \quad R_3 \end{array} + \mathbf{1-IMes}$$

Entry	Substrate	Product	Yield
1			90%
2			90%
3			86%
4			80%

The ability of $\text{Ni}(\text{NHC})_2$ to catalyze cyclopropanation was also tested (Scheme 6). Toluene solutions of 1-hexene react with N_2CPh_2 (1 equiv, 60 °C, 12 h) with 20 mol% **3-IMes** or **1-IMes** to give the corresponding cyclopropane in 90% isolated yield. There is no distinguishable difference between the catalytic rates of **3-IMes** and **1-IMes**, which is not surprising as **3-IMes** readily converts to **1-IMes** at 60 °C. The high reactivity of metal-stabilized carbenoids usually requires cyclopropanation reactions to be performed with excess olefin, by slow addition of diazoalkane, or both, to minimize the concentration of carbenes and thus avoid dimerization side products.^[9, 73-76] The ability of **1-IMes** to give high yields of cyclopropanation with a 1:1 ratio of diazoalkane to olefin is therefore particularly noteworthy.

Scheme 6. Synthetic formation of 1,1-diphenyl-2-butylcyclopropane mediated by catalytic amounts of either **1-IMes** or **3-IMes**.

3.5. *Proposed catalytic cycle for cyclopropanation.* Most transition-metal mediated cyclopropanation reactions are proposed to proceed through a metallacarbene intermediate,^[77-80] but this is not always the case. Peace has proposed diazoalkane attack on a Cu-coordinated olefin as one of the pathways by which cyclopropanes are formed with $[(\text{MeO})_3\text{P}]_2\text{CuI}$ or $\text{Cu}(\text{acac})_2$,^[81] and Maulide has proposed olefin activation for a series of Au-catalyzed intramolecular cyclopropanations where the putative metallacarbene complex is inactive for cyclopropanation.^[82] A metallacarbene intermediate is unlikely with **3-IMes** because, at the elevated temperature required for cyclopropanation, the diazoalkane ligand dissociates from the metal center rather than form a carbene (vide supra). Therefore, another catalytic pathway is probably operative.

The ability of olefins to coordinate to a metal center has significant effects on the rate and selectivity for cyclopropanations with such catalysts as CuOTf and $\text{Pd}(\text{OAc})_2$.^[83] When excess 1-hexene is added to C_6D_6 solutions of **1-IMes**, a color change occurs from purple to red and complete conversion to a new complex is observed. The ^1H NMR spectrum of this complex is consistent with an η^2 -coordination of 1-hexene to the metal with one alkene molecule for every two IMes ligands, $(\text{IMes})_2\text{Ni}(\eta^2\text{-C}_6\text{H}_{12})$ (**5**) (Scheme 7). A similar mono-alkene adduct was characterized by Radius who isolated $(^i\text{Pr}_2\text{Im})_2\text{Ni}(\text{C}_2\text{H}_4)$ from reaction of $(^i\text{Pr}_2\text{Im})_4\text{Ni}_2(\text{cod})$ and ethylene.^[84] Solutions of **5** in C_6D_6 are stable up to 80 °C with no sign of degradation by ^1H NMR. Upon removal of the solvent under vacuum, **5** completely reverts to **1-IMes**. In contrast, $(^i\text{Pr}_2\text{Im})_2\text{Ni}(\text{C}_2\text{H}_4)$, with smaller NHC ligands, is stable under vacuum.^[84]

Scheme 7. Synthetic formation of **5** upon coordination of 1-hexene to **1-IMes**.

Under cyclopropanation reaction conditions at 60 °C, it is clear that the diazoalkane adduct is not stable but that the olefin adduct is. Thus, we propose a catalytic cycle where **1-IMes**

(I) activates the olefin (II) for electrophilic attack by the diazoalkane (Scheme 8) in a similar manner to that observed by Peace.^[81] N₂ elimination from the intermediate complex (III) would likely proceed with little to no barrier, and dissociation of the cyclopropane or displacement by olefin should be facile to turn over the catalytic cycle.

Scheme 8. Proposed catalytic cycle for formation of cyclopropanation products.

Alternatively, III could eliminate N₂ to form a modified metallacyclobutane (IV).^[85] It seems unlikely a metallacyclobutane intermediate supported by two IMes ligands would be sterically achievable (*vide supra*), but it is possible to envision one IMes ligand dissociating before formation of the metallacyclobutane. After reductive elimination of the cyclopropane product and re-association of IMes, the starting complex (I) could be regenerated to complete the catalytic cycle. An argument against this mechanism is that IMes is known to react with N_2CPh_2 ,^[86] for Clyburne has shown that these reagents form the corresponding azine in 48% yield after stirring at room temperature overnight. While we did not perform this control ourselves, our elevated reaction conditions (60 °C for 12 h) would likely increase the relative azine yield. Such a reaction would decrease the amount of N_2CPh_2 available to react with our olefin, which should therefore give a poor overall yield for the cyclopropane product. Due to the high overall yield seen here for these cyclopropane products, this alternative mechanism is inconsistent with experimental results.

3.6. *Reactivity of (IMes)₂Ni(η^2 -N₃Ad) (4).* Several related Ni complexes with η^2 -bound azide ligands are known.^[87, 88] N₃Ad and N₃Mes (Mes = 2,4,6-trimethylphenyl) both bind to Ni in an η^2 fashion when Ni is supported by the chelating dtbpe ligand.^[87] This also occurs when Ni is supported by the chelating NHC ligand TMTBM (TMTBM = 3,3'-methylenebis(1-tert-butyl-4,5-dimethylimidazol-ylidene)) and reacts with N₃(2,6-ⁱPr₂C₆H₃) and N₃dmp (dmp = 2,6-dimesitylphenyl) to form (TMTBM)Ni{ η^2 -N₃(2,6-ⁱPr₂C₆H₃)} and (TMTBM)Ni(η^2 -N₃dmp), respectively.^[88] While **4** is structurally similar to these reported η^2 -azides, its reactivity is distinct. **4** is stable up to 60 °C, at which point dissociation of azide occurs to regenerate **1-IMes**; formation of the corresponding imido complex is not observed. Cooling the mixture of **1-IMes** and N₃Ad to room temperature results in the reformation of **4** along with significant decomposition. In contrast, heating (dtbpe)Ni(η^2 -N₃Ad) or (dtbpe)Ni(η^2 -N₃Mes) to 50 °C results in loss of N₂ and formation of three-coordinate imido complexes.^[87] In the case of (TMTBM)Ni{ η^2 -N₃(2,6-ⁱPr₂C₆H₃)}, N₂ extrusion could not be induced thermally, chemically, or photochemically.^[88] However, use of a bulkier azide complex (TMTBM)Ni(η^2 -N₃dmb) yielded an imido complex. This result does not appear to be general, however, as we found that reaction of **1-IMes** with N₃dmp resulted in decomposition without any apparent formation of an imido complex.^[88] Thus, only for **4** is even partially reversible azide coordination observed.

A more direct comparison between **4** and (dtbpe)Ni(η^2 -N₃Ad) is warranted because the two complexes have a similar coordination mode of the identical organoazide to a Ni(0) center but exhibit very different reactivity (vide supra). This difference is most likely due to the steric and chelating effects of the NHC versus dtbpe ligands. Pörschke's complex ((dtbpe)Ni)₂(η^2 - μ -C₆H₆) was used to synthesize (dtbpe)Ni(η^2 -N₃Ad), and during coordination of N₃Ad to the Ni center the P–Ni–P bite angle hardly changes (from 93.88(1)° to 93.75(2)°).^[87, 89] Upon loss of N₂ to form the corresponding imido complex (dtbpe)Ni=NAd, this bite angle contracts slightly to 91.69(3)°. This reactivity is due to the stability imparted by the chelating dtbpe ligand: the coordinating N₃Ad ligand is not sterically encumbered and so has the necessary space to form the proposed κ^2 -N,N metallacyclic intermediate required to induce loss of N₂.^[87] Thus, the geometry of the Ni center in the dtbpe system is independent of the identity of the coordinated azide ligand. In our current system, the C–Ni–C bite angle changes from 180° in **1-IMes** to 120.1(1)° upon coordination of N₃Ad to form **4**. If **4** were to form an imido complex, this would probably occur via a metallacyclic intermediate analogous to that proposed for formation of (dtbpe)Ni=NAd

from (dtbpe)Ni(η^2 -N₃Ad).^[87] This transformation would require further contraction of the C–Ni–C bite angle, which is unfavorable, and place the large Ad group in proximity to the mesityl groups of the NHC. This is not a problem with (dtbpe)Ni(η^2 -N₃Ad) because the ^tBu groups are pulled back from the Ni center by the chelating phosphine. Therefore, at the elevated temperatures required to induce N₂ loss and transform (dtbpe)Ni(η^2 -N₃Ad) to (dtbpe)Ni=NAd, it is more favorable for N₃Ad to dissociate from **4** to generate the stable, linear **1-IMes** precursor, before reforming **4** upon cooling.

Complex **4** was also examined for stoichiometric alkane aziridination with 1-hexene in the same manner that **3-IMes** was studied (vide supra) so as to allow for direct comparison between these systems. Upon exposure of **4** to 1-hexene and heating to 60 °C, the corresponding aziridination product was not generated. This lack of reactivity is likely due to dissociation and partial decomposition of the N₃Ad unit from the Ni center at 60 °C without loss of N₂ or formation of the coordinated imido for reasons described previously. Additionally, since 1-hexene coordinates to the Ni center in **1-IMes** to form **5**, which is stable up to 80 °C, it is likely that, at 60 °C, the alkene displaces N₃Ad without reacting. Thus, while **1-IMes** has the capacity to bind both diazoalkanes and organoazides to form coordinated Ni(0) species, only the bound diazoalkane adducts are useful for group transfer reactivity under the conditions tested here.

4. CONCLUSIONS.

In conclusion, we have prepared and characterized the complexes (IMes)₂Ni(κ^1 -N₂CPh₂) (**3-IMes**), (SIMes)₂Ni(κ^1 -N₂CPh₂) (**3-SIMes**), and (IPr)₂Ni(κ^1 -N₂CPh₂) (**3-IPr**), all of which possess a κ^1 -N end-on coordination mode to the Ni center through the terminal nitrogen atom of the diazoalkane ligand. All three complexes display Ni–N–N and N–N–C angles that significantly depart from linearity and have extensive delocalization of the π -bonds in the diazoalkane. We have also prepared and characterized the organoazide complex (IMes)₂Ni(η^2 -N₃Ad) (**4**) which possesses an η^2 coordination of the azide ligand to the Ni center. While elimination of N₂ from these complexes to create a new family of terminal Ni carbene and terminal imido complexes does not occur, **3-IMes** was successfully utilized in stoichiometric and catalytic olefin cyclopropanation reactions by producing cyclopropane products in good yields from 1:1 mixtures of diazoalkanes and olefins. **3-IMes** mediates cyclopropanation with remarkable efficiency, and so

these complexes are a promising starting point for the development of a new class of cyclopropanation catalysts.

ACKNOWLEDGEMENTS.

This paper is dedicated to the memory of Prof. Gregory L. Hillhouse who was a terrific advisor and mentor. This work was supported by National Science Foundation through grants CHE-0957816, CHE-1266281, CHE-0615274 (to G.L.H.). We thank Prof. Michael Hopkins and Prof. Richard Jordan for their comments on and assistance with the manuscript.

APPENDIX A. SUPPLEMENTARY DATA.

CIF files contain the supplementary crystallographic data for this paper. Crystallographic data tables, $^1\text{H}/^{13}\text{C}$ NMR, and mass spectra files for the structures of compounds **3-IMes**, **3-SIMes**, **3-IPr**, and **4**, as well as alternative syntheses for **1-IMes**, **1-IPr**, and **2**, can be found in the supplementary data for this article.

CORRESPONDING AUTHOR

* E-mail: nharrold@uchicago.edu

PRESENT ADDRESS

‡ Department of Chemistry, University of Wisconsin–Madison, Madison, WI 53706, USA.

NOTES

The authors declare no competing financial interest.

⊥ Deceased March 6, 2014.

REFERENCES.

- [1] D. Seyferth, *Chem. Rev.* **1955**, 55, 1155-1177.
- [2] W. R. Roper, *J. Organomet. Chem.* **1986**, 300, 167-190.
- [3] H. Werner, *J. Organomet. Chem.* **1995**, 500, 331-336.
- [4] W. A. Herrmann, *Angew. Chem. Int. Ed.* **1978**, 17, 800-812.
- [5] M. P. Doyle, *Chem. Rev.* **1986**, 86, 919-939.
- [6] D. A. Lemenovskii, M. Putala, *Russ. Chem. Rev.* **1994**, 63, 197-216.
- [7] Y. Mizobe, Y. Ishii, M. Hidai, *Coord. Chem. Rev.* **1995**, 139, 281-311.
- [8] A. Padwa, M. D. Weingarten, *Chem. Rev.* **1996**, 96, 223-270.
- [9] M. P. Doyle, D. C. Forbes, *Chem. Rev.* **1998**, 98, 911-936.
- [10] H. M. L. Davies, S. A. Panaro, *Tetrahedron* **2000**, 56, 4871-4880.

- [11] H. M. L. Davies, E. G. Antoulinakis, *J. Organomet. Chem.* **2001**, 617-618, 47-55.
- [12] R. Beckhaus, C. Santamaría, *J. Organomet. Chem.* **2001**, 617-618, 81-97.
- [13] D. Sutton, *Chem. Rev.* **1993**, 93, 995-1022.
- [14] M. Dartiguenave, M. J. Menu, E. Deydier, D. Yves, H. Siebald, *Coord. Chem. Rev.* **1998**, 178-180, 623-663.
- [15] N. P. Mankad, J. C. Peters, *Chem. Commun.* **2008**, 1061-1063.
- [16] J. J. Curley, T. Murahashi, C. C. Cummins, *Inorg. Chem.* **2009**, 48, 7181-7193.
- [17] K. Kromm, F. Hampel, J. A. Gladysz, *Organometallics* **2002**, 21, 4264-4274.
- [18] O. L. Sydora, D. S. Kuiper, P. T. Wolczanski, E. B. Lobkovsky, A. Dinescu, T. R. Cundari, *Inorg. Chem.* **2006**, 45, 2008-2021.
- [19] M. E. García, D. García-Vivó, M. A. Ruiz, P. Herson, *Organometallics* **2008**, 27, 3879-3891.
- [20] U. J. Kilgore, C. A. Sengelaub, M. Pink, A. R. Fout, D. J. Mindiola, *Angew. Chem. Int. Ed.* **2008**, 47, 3769-3772.
- [21] A. I. Nguyen, K. J. Blackmore, S. M. Carter, R. A. Zarkesh, A. F. Heyduk, *J. Am. Chem. Soc.* **2009**, 131, 3307-3316.
- [22] D. E. Herbert, N. C. Lara, T. Agapie, *Chem. Eur. J.* **2013**, 19, 16453-16460.
- [23] A. Nakamura, T. Yoshida, M. Cowie, S. Otsuka, J. A. Ibers, *J. Am. Chem. Soc.* **1977**, 99, 2108-2117.
- [24] S. Gambarotta, C. Floriani, A. Chiesi-Villa, C. Guastini, *J. Am. Chem. Soc.* **1982**, 104, 1918-1924.
- [25] J. L. Polse, A. W. Kaplan, R. A. Andersen, R. G. Bergman, *J. Am. Chem. Soc.* **1998**, 120, 6316-6328.
- [26] A. W. Kaplan, J. L. Polse, G. E. Ball, R. A. Andersen, R. G. Bergman, *J. Am. Chem. Soc.* **1998**, 120, 11649-11662.
- [27] T. Schaub, U. Radius, *Z. Anorg. Allg. Chem.* **2007**, 633, 2168-2172.
- [28] V. M. Iluc, G. L. Hillhouse, *J. Am. Chem. Soc.* **2014**, 136, 6479-6488.
- [29] For a more complete discussion of diazoalkane binding motifs, see: G. L. Hillhouse, B. L. Haymore, *J. Am. Chem. Soc.* **1982**, 104, 1537-1548.
- [30] W. A. Nugent, J. M. Mayer, *Metal-Ligand Multiple Bonds*, John Wiley & Sons, New York, **1988**.
- [31] D. E. Wigley, *Prog. Inorg. Chem.* **1994**, 42, 239-482.
- [32] S. Cenini, E. Gallo, A. Caselli, F. Ragaini, S. Fantauzzi, C. Piangiolino, *Coord. Chem. Rev.* **2006**, 250, 1234-1253.
- [33] R. Waterman, G. L. Hillhouse, *J. Am. Chem. Soc.* **2003**, 125, 13350-13351.
- [34] P. Müller, C. Fruit, *Chem. Rev.* **2003**, 103, 2905-2920.
- [35] Y. M. Badiei, A. Dinescu, X. Dai, R. M. Palomino, F. W. Heinemann, T. R. Cundari, T. H. Warren, *Angew. Chem. Int. Ed.* **2008**, 47, 9961-9964.
- [36] N. D. Harrold, R. Waterman, G. L. Hillhouse, T. R. Cundari, *J. Am. Chem. Soc.* **2009**, 131, 12872-12873.
- [37] E. R. King, E. T. Hennessy, T. A. Betley, *J. Am. Chem. Soc.* **2011**, 133, 4917-4923.
- [38] M. G. Fickes, W. M. Davis, C. C. Cummins, *J. Am. Chem. Soc.* **1995**, 117, 6384-6385.
- [39] G. Proulx, R. G. Bergman, *J. Am. Chem. Soc.* **1995**, 117, 6382-6383.
- [40] G. Proulx, R. G. Bergman, *Organometallics* **1996**, 15, 684-692.
- [41] T. A. Hanna, A. M. Baranger, R. G. Bergman, *Angew. Chem. Int. Ed.* **1996**, 35, 653-655.
- [42] G. Guillemot, E. Solari, C. Floriani, C. Rizzoli, *Organometallics* **2001**, 20, 607-615.

- [43] M. Barz, E. Herdtweck, W. R. Thiel, *Angew. Chem. Int. Ed.* **1998**, 37, 2262-2265.
- [44] H. V. R. Dias, S. A. Polach, S.-K. Goh, E. F. Archibong, D. S. Marynick, *Inorg. Chem.* **2000**, 39, 3894-3901.
- [45] G. Albertin, S. Antoniutti, D. Baldan, J. s. Castro, S. García-Fontán, *Inorg. Chem.* **2008**, 47, 742-748.
- [46] A. B. Pangborn, M. A. Giardello, R. H. Grubbs, R. K. Rosen, F. J. Timmers, *Organometallics* **1996**, 15, 1518-1520.
- [47] J. Miller, *J. Org. Chem.* **1959**, 24, 560-561.
- [48] P. A. S. Smith, C. D. Rowe, L. B. Bruner, *J. Org. Chem.* **1969**, 34, 3430-3433.
- [49] J. A. Miller, *Tetrahedron Lett.* **1975**, 16, 2959-2960.
- [50] B. C. Ranu, A. Sarkar, R. Chakraborty, *J. Org. Chem.* **1994**, 59, 4114-4116.
- [51] A. J. Arduengo III, R. Krafczyk, R. Schmutzler, H. A. Craig, J. R. Goerlich, W. J. Marshall, M. Unverzagt, *Tetrahedron* **1999**, 55, 14523-14534.
- [52] S. P. Nolan. U.S. Patent 7109348, **2006**.
- [53] V. Ritleng, E. Brenner, M. J. Chetcuti, *J. Chem. Educ.* **2008**, 85, 1646.
- [54] L. Jafarpour, E. D. Stevens, S. P. Nolan, *J. Organomet. Chem.* **2000**, 606, 49-54.
- [55] A. J. Arduengo III, S. F. Gamper, J. C. Calabrese, F. Davidson, *J. Am. Chem. Soc.* **1994**, 116, 4391-4394.
- [56] SMART; Bruker-AXS: Madison, WI, **2009**.
- [57] G. M. Sheldrick, *Acta Cryst.* **2008**, A64, 112-122.
- [58] O. V. Dolomanov, L. J. Bourhis, R. J. Gildea, J. A. K. Howard, H. Puschmann, *J. Appl. Cryst.* **2009**, 42, 339-341.
- [59] SADABS; Bruker-AXS: Madison, WI, **2012**.
- [60] <http://www.cryst.chem.uu.nl/platon/>.
- [61] T. Dröge, F. Glorius, *Angew. Chem. Int. Ed.* **2010**, 49, 6940-6952.
- [62] J. Louie, J. E. Gibby, M. V. Farnworth, T. N. Tekavec, *J. Am. Chem. Soc.* **2002**, 124, 15188-15189.
- [63] K. Matsubara, K. Ueno, Y. Shibata, *Organometallics* **2006**, 25, 3422-3427.
- [64] K. Matsubara, S. Miyazaki, Y. Koga, Y. Nibu, T. Hashimura, T. Matsumoto, *Organometallics* **2008**, 27, 6020-6024.
- [65] A. A. Danopoulos, D. Pugh, *Dalton Trans.* **2008**, 30-31.
- [66] A. C. Hillier, W. J. Sommer, B. S. Yong, J. L. Petersen, L. Cavallo, S. P. Nolan, *Organometallics* **2003**, 22, 4322-4326.
- [67] H. Clavier, S. P. Nolan, *Chem. Commun.* **2010**, 46, 841-861.
- [68] D. J. Mindiola, G. L. Hillhouse, *J. Am. Chem. Soc.* **2001**, 123, 4623-4624.
- [69] A. Hammerl, T. Klapotke, in *Encyclopedia of Inorganic Chemistry*, 2nd ed. (Ed.: R. King), Wiley & Sons, Chichester, **2005**.
- [70] F. H. Allen, O. Kennard, D. G. Watson, L. Brammer, A. G. Orpen, R. Taylor, *J. Chem. Soc., Perkin Trans. 2* **1987**, S1-S19.
- [71] M. I. Javed, M. Brewer, *Org. Synth.* **2008**, 85, 189.
- [72] R. Cohen, B. Rybtchinski, M. Gandelman, H. Rozenberg, J. M. L. Martin, D. Milstein, *J. Am. Chem. Soc.* **2003**, 125, 6532-6546.
- [73] V. K. Singh, A. DattaGupta, G. Sekar, *Synthesis* **1997**, 1997, 137-149.
- [74] X. Dai, T. H. Warren, *J. Am. Chem. Soc.* **2004**, 126, 10085-10094.
- [75] H. Pellissier, *Tetrahedron* **2008**, 64, 7041-7095.
- [76] D. Intrieri, A. Caselli, E. Gallo, *Eur. J. Inorg. Chem.* **2011**, 2011, 5071-5081.

- [77] K. H. Dötz, *Angew. Chem. Int. Ed.* **1984**, 23, 587-608.
- [78] M. P. Doyle, in *Comprehensive Organometallic Chemistry II* (Ed.: E. W. A. G. A. S. Wilkinson), Elsevier, Oxford, **1995**, pp. 387-420.
- [79] W. D. Wulff, in *Comprehensive Organometallic Chemistry II* (Ed.: E. W. A. G. A. S. Wilkinson), Elsevier, Oxford, **1995**, pp. 469-547.
- [80] D. F. Harvey, D. M. Sigano, *Chem. Rev.* **1996**, 96, 271-288.
- [81] D. S. Wulfman, R. S. McDaniel Jr, B. W. Peace, *Tetrahedron* **1976**, 32, 1241-1249.
- [82] X. Huang, S. Klimczyk, L. F. Veiros, N. Maulide, *Chem. Sci.* **2013**, 4, 1105-1110.
- [83] A. J. Anciaux, A. J. Hubert, A. F. Noels, N. Petiniot, P. Teyssie, *J. Org. Chem.* **1980**, 45, 695-702.
- [84] T. Schaub, U. Radius, *Chem. Eur. J.* **2005**, 11, 5024-5030.
- [85] C. Rodríguez-García, A. Oliva, R. M. Ortuño, V. Branchadell, *J. Am. Chem. Soc.* **2001**, 123, 6157-6163.
- [86] J. M. Hopkins, M. Bowdridge, K. N. Robertson, T. S. Cameron, H. A. Jenkins, J. A. C. Clyburne, *J. Org. Chem.* **2001**, 66, 5713-5716.
- [87] R. Waterman, G. L. Hillhouse, *J. Am. Chem. Soc.* **2008**, 130, 12628-12629.
- [88] N. D. Harrold, G. L. Hillhouse, *Chem. Sci.* **2013**, 4, 4011-4015.
- [89] I. Bach, K.-R. Pörschke, R. Goddard, C. Kopiske, C. Krüger, A. Rufinska, K. Seevogel, *Organometallics* **1996**, 15, 4959-4966.