(CAAC)CuX-Catalyzed Hydroboration of Terminal Alkynes with Pinacolborane Directed by the X-ligand

Erik A. Romero, Rodolphe Jazzar, Guy Bertrand*

Abstract: When X is a basic ligand (CAAC)CuX complexes [CAAC = cyclic (alkyl)(amino)carbene] selectively promote the (*E*)- β -hydroboration of terminal alkynes with pinacolborane instead of the dehydrogenative borylation observed with weakly nucleophilic X ligand. This methodology is applicable to a variety of terminal alkynes. Deuterium labeling experiments coupled with stoichiometric reactions give evidence towards a plausible mechanism involving a σ -mono(copper)acetylide complex.

Introduction

Due to the importance of the Suzuki-Miyaura cross-coupling reaction, organoboron esters have found wide applicability as synthons in organic chemistry [1]. Alkenyl derivatives have been prepared via transition metal-catalyzed hydroboration of the corresponding alkynes [2], with only seldom reports involving the use of inexpensive metals such as copper [3,4,5]. With this metal, Yun *et al.* [5] recently described an elegant atom efficient methodology affording selectively the (*E*)- β -hydroboration or the (*Z*)- β -hydroboration products of terminal alkynes, using 1,8-naphthalenediaminatoborane as the borane partner. There are no extensive investigations detailing the hydroboration of terminal alkynes to afford (*E*)- β -hydroboration using pinacolborane and a copper catalyst. However, Hoveyda *et al.* [3a] demonstrated that both (*E*)- β - and α -hydroboration of terminal alkynes can be obtained using bis(pinacolato)diboron (PinBBPin) as the boron partner. Note that this synthetic method requires the presence of an alkoxide, which generates a stoichiometric amount of ROBPin as waste.

Previously, we reported on the role of the X ligand in the mechanism of the (CAAC)CuX-catalyzed [CAAC = cyclic (alkyl)(amino)carbene [6,7]] azide-alkyne cycloaddition (Click reaction) [8]. We showed that basic X ligands favor the metallation step leading to the σ -mono(copper) acetylide complexes **A** [8b, 9], but disfavor the formation of the catalytically active σ,π -bis(copper)acetylide **B** (Fig. 1). Conversely, non-nucleophilic ligands favor the latter but slowly promote the former; this problem could be circumvented, however, in the presence of an external base. Using this knowledge, we recently demonstrated [10] that (CAAC)CuX complexes, featuring a weakly nucleophilic X ligand such as trifluoromethanesulfonate can selectively promote the dehydrogenative borylation of terminal alkynes with pinacol borane [11]. This result is due to the formation of **B** [8] in which the triple bond is protected, and thus cannot participate in classical hydroboration. We hypothesized, that in contrast to **B**, **A** features an unprotected and polarized C-C triple bond and should readily undergo hydroboration. Here we report that indeed a simple modification of the X ligand of (CAAC)CuX complexes allows for switching the selectivity of the reaction of pinacolborane with terminal alkynes from dehydrogenative borylation to (*E*)- β -hydroboration.

Figure 1. (CAAC)CuOTf has been shown to promote the dehydrogenative borylation of terminal alkynes due to the formation of **B**. (CAAC)CuOPh should catalyze the hydroboration of terminal alkynes via the copper acetylides **A**.

Results and discussion

We started our investigation using a stoichiometric mixture of phenylacetylene **1a** and pinacolborane in C₆D₆ at room temperature for 2 hours. We first checked that there was no reaction at room temperature in the absence of catalyst (Table 1, Entry 1). As previously reported [10], in the presence of 2.5 mol% of (CAAC)CuOTf no significant reaction occurred which stems from the difficulty of triflate (pK_a = -12) to deprotonate **1a** (Entry 2). However, in the presence of Et₃N (5 mol%), as an additive, selective formation of the dehydrogenative borylation product **3a** was observed (Entry 3). In marked contrast, with the more basic phenoxide anion (pK_a = 10), which favors the formation of **A** at the expense of the dinuclear complex **B**, the (*E*)- β -hydroboronic ester **2a** was obtained with no trace of **3a** (Entry 6). The influence of the basicity of the X ligand was confirmed, using the benzoate (pK_a = 4.2) and the acetate (pK_a = 4.8) (Entry 4-5).

A survey of solvent polarities (Entries 6-9) revealed that the reaction is in all cases very selective for the formation of **2a** with acetonitrile giving the best results (94% NMR yield). We found that decreasing the concentration from 2.3 to 1.6 mol/L had little effect on the selectivity (Entry 10), but further dilution to 1.3 mol/L significantly affects the formation of **2a** (Entry 11). Lastly, decreasing the catalyst loading below 2.5 mol% also proved detrimental to the reaction (Entries 12-13).

Table 1. Effect of the X ligand, solvent, and concentration on the β -hydroboration reaction.^a

Dipp = 2,6-Diisopropylphenyl

Entry	LCuX (mol%) X	Solvent	Conc. (M)	2a (%) ^b	3a (%) ^b
1	-	C ₆ D ₆	2.3	0	0
2	OTf (2.5)	C ₆ D ₆	2.3	1	6
3	OTf (2.5)^c	C₆D₆	2.3	1	90
4	OBz (2.5)	C ₆ D ₆	2.3	30	20
5	OAc (2.5)	C ₆ D ₆	2.3	40	10
6	OPh (2.5)	C ₆ D ₆	2.3	61	0
7	OPh (2.5)	CD ₂ Cl ₂	2.3	85	0
8	OPh (2.5)	THF-d ₈	2.3	79	0
9	OPh (2.5)	CD₃CN	2.3	94	0
10	OPh (2.5)	CD ₃ CN	1.6	91	0
11	OPh (2.5)	CD ₃ CN	1.3	77	0
12	OPh (1)	CD ₃ CN	2.3	81	0
13	OPh (0.5)	CD ₃ CN	2.3	53	0

[a] Reactions were carried out for 2 h in a test tube at RT under an argon atmosphere using 0.69 mmol of phenylacetylene and 0.71 mmol of pinacol borane. **[b]** Measured by NMR using CH₂Cl₂ or toluene as an internal standard. **[c]** 5 mol% Et₃N.

Using the optimized conditions, the scope of the hydroboration reaction was then investigated at room temperature in acetonitrile (2.3 ± 0.1 M) using 3.65 mmol of alkyne and 3.74 mmol of borane in the presence of 2.5 mol% (CAAC)CuOPh (Fig. 2). This reaction encompasses a wide range of electronically diverse terminal alkynes. In most cases, the NMR spectra of the crude reaction mixture showed excellent selectivity, and the alkenylboronic esters **2a-m** were isolated in good to excellent yields. Importantly, purification of products **2a-m** is readily achieved in air by filtration through a short plug of silica with pentane as the eluent. This protocol also allows for gram-scale syntheses, as shown with **2a**.

Figure 2. Scope of the hydroboration of terminal alkynes. **[a]** reaction time 2 h. **[b]** reaction time 4 h.

We then turned our attention to the mechanism of the reaction. We previously showed that (CAAC)CuOPh reacts with terminal alkynes to afford the copper acetylide complex **B** with elimination of phenol [8b]. Note that complexes of type **B** can be isolated, and that phenol readily reacts with pinacol borane to afford the catalytically inert PinBOPh (See SI for characterization). In accordance with recent work by Roesky *et al.*, [12] we postulate that a 1,2-addition of PinB-H across the C(sp)-C(sp) bond of **B** gives rise to a copper alkenyl boronic ester, which could not be isolated. Then, as already observed in the copper-catalyzed azide-alkyne cycloaddition reaction (CuAAC) [8], the alkyne should be acidic enough to induce a protonolysis, which affords **2** and regenerates the copper acetylide complex **B** (Fig. 3). In order to confirm this hypothesis, we performed the catalytic reaction using deuterium labelled phenylacetylene, and as expected, we observed the formation **2a-D** (90% D-incorporation). In addition, when a stoichiometric amount of copper phenylacetylide **Ba** and PinB-H were reacted with a three-fold excess of tolylacetylene, the copper tolyl acetylide **Bb** and **2a** were isolated.

-
- ⁸ a) L. Jin, D. R. Tolentino, M. Melaimi, G. Bertrand, *Sci. Adv.* 1 (2015) e1500304, b) L. Jin, E. A. Romero, M. Melaimi, G. Bertrand, *J. Am. Chem. Soc.* 137 (2015) 15696-15698.
- ⁹ Most copper acetylides are polymeric in nature with the exception of a (CAAC)CuC≡CPh⁸ and (NHC)CuC≡CPh complex: C. Nolte, P. Mayer, B. F. Straub, *Angew. Chem. Int. Ed.* 46 (2007) 2101-2103.
- ¹⁰ E. A. Romero, R. Jazzar, G. Bertrand, *Chem. Sci.* (2016) DOI: 10.1039/C6SC02668K.
- ¹¹ a) C. I. Lee, J. Zhou, O. V. Ozerov, *J. Am. Chem. Soc.* 135 (2013) 3560-3566; b) C. I. Lee, N. A. Hirscher, J. Zhou, N. Bhuvanesh, O. V. Ozerov, *Organometallics* 34 (2015) 3099-3102; c) C. I. Lee, W. C. Shih, J. Zhou, J. H. Reibenspies, O. V. Ozerov, *Angew. Chem. Int. Ed.* 54 (2015) 14003-14007; d) C. I. Lee, J. C. DeMott, C. J. Pell, A. Christopher, J. Zhou, N. Bhuvanesh, O. V. Ozerov, *Chem. Sci.* 6 (2015) 6572-6582; e) C. J. Pell, O. V. Ozerov, *Inorg. Chem. Front.* 2 (2015) 720-724.
- ¹² Y. Yang, M. Zhong, X. Ma, K. Nijesh, S. De, P. Parameswaran, H. W. Roesky, *J. Am. Chem. Soc.* 138 (2016) 2548-2551.

*Graphical abstract

Figure 3. Proposed mechanism and related experiments for the hydroboration of terminal alkynes.

In summary, we have shown that a simple replacement of a weakly coordinating X ligand by a basic anion allowed for switching the fate of the (CAAC)CuX catalysed reaction of pinacolborane with terminal alkynes. Whereas with X = OTf a dehydrogenative borylation occurs, with X = OPh (E)- β -hydroboration takes place. Deuterium labeling experiments coupled with stoichiometric reactions suggest that a σ -mono(copper)acetylide is the key catalytic species in the hydroboration. The importance of the X ligand in other copper catalyzed reactions is under active investigation.

Acknowledgements

The authors gratefully acknowledge financial support from the DOE (DE-FG02-13ER16370).

Notes and References

- 1 a) J. F. Hartwig, *Acc. Chem. Res.* 45 (2012) 864-873; b) A. J. Lennox, G. C. Lloyd-Jones, *Chem. Soc. Rev.* 43 (2014) 412-443.
- 2 For representative examples of transition metal catalyzed hydroboration with pinacol borane, see: a) S. Pereira, M. Srebnik, *Organometallics* 14 (1995) 3127-3128; b) M. Haberberger, S. Enthaler, *Chem. Asian J.* 8 (2013) 50-54; c) T. Lee, C. Baik, I. Jung, K. H. Song, S. Kim, D. Kim, S. O. Kang, J. Ko, *Organometallics* 23 (2004) 4569-4575; d) D. M. Khranov, E. L. Rosen, J. A. V. Er, P. D. Vu, V. M. Lynch, C. W. Bielawski, *Tetrahedron* 64 (2008) 6853-6862; e) C. Gunanathan, M. Hölscher, F. Pan, W. Leitner, *J. Am. Chem. Soc.* 134 (2012) 14349-14352; f) J. V. Obligation, J. M. Neely, A. N. Yazdani, I. Pappas, P. J. Chirik, *J. Am. Chem. Soc.* 137 (2015) 5855-5858; g) T. Ohmura, Y. Yamamoto, N. Miyaura, *J. Am. Chem. Soc.* 122 (2000) 4990-4991; h) Y. D. Wang, G. Kimball, A. S. Prasad, Y. Wang, *Tetrahedron Lett.* 46 (2015) 8777-8780.
- 3 a) H. Jang, A. R. Zhugralin, Y. Lee, A. H. Hoveyda, *J. Am. Chem. Soc.* 133 (2011) 7859-7871; b) Y. D. Bidal, F. Lazreg, C. S. J. Cazin, *ACS Catal.* 4 (2014) 1564-1569; c) F. Lazreg, F. Nagra, C. S. J. Cazin, *Coord. Chem. Rev.* 293-284 (2015) 48-79; d) J. Yun, *Asian J. Org. Chem.* 2 (2013) 1016-1025; e) H. R. Kim, I. G. Jung, K. Yoo, K. Jang, E. S. Lee, J. Yun, S. U. Son, *Chem. Commun.* 46 (2010) 758-760; f) J.-E. Lee, J. Kwon, J. Yun, *Chem. Commun.* (2008) 733-734.
- 4 Examples of bis(hydroboration) reaction have been reported : S. Lee, D. Li, and J. Yun, *Chem. Asian J.* 9 (2014), 2440-2443.
- 5 W. J. Jang, W. L. Lee, J. H. Moon, J. Y. Lee, J. Yun, *Org. Lett.* 18 (2016) 1390-1393.
- 6 For reviews on CAACs, see: a) M. Melaimi, M. Soleilhavoup, G. Bertrand, *Angew. Chem. Int. Ed.* 49 (2010) 8810-8849; b) D. Martin, M. Melaimi, M. Soleilhavoup, G. Bertrand, *Organometallics* 30 (2011) 5304-5313; c) M. Soleilhavoup, G. Bertrand, *Acc. Chem. Res.* 48 (2015) 256-266.
- 7 For the synthesis of CAACs, see: a) V. Lavallo, Y. Canac, C. Präsang, B. Donnadieu, G. Bertrand, *Angew. Chem. Int. Ed.* 44 (2005) 5705-5709; b) R. Jazzar, R. D. Dewhurst, J. B. Bourg, B. Donnadieu, Y. Canac, G. Bertrand, *Angew. Chem. Int. Ed.* 46 (2007) 2899-2902; c) R. Jazzar, J. B. Bourg, R. D. Dewhurst, B. Donnadieu, G. Bertrand, *J. Org. Chem.* 72 (2007) 3492-3499; d) X. Zeng, G. D. Frey, R. Kinjo, B. Donnadieu, G. Bertrand, *J. Am. Chem. Soc.* 131 (2009) 8690-8696.