LOWER BOUND FOR RANKS OF INVARIANT FORMS

HARM DERKSEN AND ZACH TEITLER

ABSTRACT. We give a lower bound for the Waring rank and cactus rank of forms that are invariant under an action of a connected algebraic group. We use this to improve the Ranestad–Schreyer–Shafiei lower bounds for the Waring ranks and cactus ranks of determinants of generic matrices, Pfaffians of generic skew-symmetric matrices, and determinants of generic symmetric matrices.

1. INTRODUCTION

Let $F \in \mathbb{C}[x_1, \dots, x_n]$ be a homogeneous form of degree d . The **Waring rank** of F , denoted $r(F)$, is the minimum number of terms in an expression for F as a linear combination of powers of linear forms:

$$F = \sum_{i=1}^r c_i \ell_i(x_1, \dots, x_n)^d, \quad c_i \in \mathbb{C}.$$

For example,

$$x_1 \cdots x_n = \frac{1}{2^{n-1}n!} \sum_{\substack{\epsilon \in \{\pm 1\}^n \\ \epsilon_1 = 1}} \left(\prod \epsilon_i \right) \left(\sum \epsilon_i x_i \right)^n,$$

so $r(x_1 \cdots x_n) \leq 2^{n-1}$. In fact $r(x_1 \cdots x_n) = 2^{n-1}$. This is a consequence of a general lower bound for Waring rank shown by Ranestad and Schreyer in [21]. For forms that are invariant under a group action, we improve the general Ranestad–Schreyer lower bound.

Power sum decompositions of this type and Waring ranks have been studied since the 19th century, thanks to their connections to the number-theoretic Waring problem, secant varieties in algebraic geometry, interpolation and quadrature methods, mixture models in statistics, and more. For comprehensive treatments, including history and applications, see [12, 14, 9, 8].

Unfortunately, Waring ranks are in general difficult to compute, and have been calculated for only a few families of polynomials. Here is a non-comprehensive list of some highlights. They are known for quadratic forms and binary forms (forms in $n = 2$ variables), classically. Waring ranks of plane cubics (i.e., forms of degree $d = 3$ in $n = 3$ variables) are known, see for example [9], [16, §8]. The Alexander–Hirschowitz Theorem [1] determines the rank of a general form, i.e., one in an open subset of the space of forms. Recently, Carlini, Catalisano, and Geramita found the Waring ranks of monomials and sums of pairwise coprime monomials [7]: the Waring rank of a sum of pairwise coprime monomials is the sum of the ranks of the monomials. More generally, if $x = (x_1, \dots, x_m)$ and $y = (y_1, \dots, y_n)$ are independent

Date: April 23, 2015.

2010 Mathematics Subject Classification. 15A21, 14N15.

Key words and phrases. Waring rank, invariant polynomials.

The first author was supported by NSF grant DMS 1302032.

variables and $F(x)$, $G(y)$ are forms of the same degree, it is conjectured that $r(F(x)+G(y)) = r(F)+r(G)$, see [25, 4, 5, 6] for very recent progress. One direction is obvious: concatenating power sum decompositions of F and G shows $r(F(x)+G(y)) \leq r(F)+r(G)$. For the reverse inequality, as for the recent determination of Waring ranks of monomials, lower bounds for Waring rank are key. Our main result is a lower bound for Waring rank which improves previous bounds for important examples such as determinants.

An interesting example of a form whose Waring rank is not yet known is the generic determinant,

$$\det_n = \det \begin{pmatrix} x_{1,1} & \cdots & x_{1,n} \\ \vdots & & \vdots \\ x_{n,1} & \cdots & x_{n,n} \end{pmatrix},$$

a form of degree n in n^2 variables. Because $\det_2$ is a quadratic form, its rank is known, namely $r(\det_2) = 4$, but $r(\det_n)$ is unknown for $n \geq 3$. As $\det_n$ is a sum of $n!$ terms of the form $x_{1,i_1} \cdots x_{n,i_n}$, each with rank 2^{n-1} , we have $r(\det_n) \leq 2^{n-1}n!$. (So $r(\det_3) \leq 24$.) This was recently improved to $r(\det_n) \leq \left(\frac{5}{6}\right)^{\lfloor n/3 \rfloor} 2^{n-1}n!$ [10, §8]. (So $r(\det_3) \leq 20$.)

Several lower bounds for Waring rank have been proposed. The classical lower bound via Sylvester's catalecticants gives $r(\det_n) \geq \binom{n}{\lfloor n/2 \rfloor}^2$; this gives $r(\det_3) \geq 9$, and asymptotically (via Stirling approximation) this lower bound grows as $2^n / \sqrt{\pi n/2}$. In [16] it is shown that $r(\det_n) \geq \binom{n}{\lfloor n/2 \rfloor}^2 + n^2 - (\lfloor n/2 \rfloor + 1)^2$; this gives $r(\det_3) \geq 14$, but this lower bound has the same asymptotic growth. Most recently, Shafiei [23], using the Ranestad–Schreyer lower bound [21], has shown $r(\det_n) \geq \frac{1}{2} \binom{2n}{n}$; this gives $r(\det_3) \geq 10$, and asymptotically it grows as $4^n / (2\sqrt{\pi n})$. (Shafiei also considers permanents, Pfaffians, symmetric determinants and permanents, etc., see [23, 22].)

In this paper we show that $r(\det_n) \geq \binom{2n}{n} - \binom{2n-2}{n-1}$; this gives $r(\det_3) \geq 14$ and asymptotically it is $\frac{3}{2}$ times the Ranestad–Schreyer–Shafiei bound. This is an example of the main result of this paper, a lower bound for Waring ranks of invariant forms under the action of a connected group.

For a polynomial F , let $\text{Diff}(F)$ be the vector space spanned by the partial derivatives of F of all orders. A special case of our main result is the following.

Theorem 1.1. *Let G be a connected algebraic group and let V be an irreducible representation of G . Let $F \in S^d V$ be an invariant form of degree d , that is, a form such that for every $g \in G$, ${}^g F = F$. Fix a basis $x_1, \dots, x_n$ for V , write $F = F(x_1, \dots, x_n)$, and let $F' = \partial F / \partial x_1$. Then $r(F) \geq \dim \text{Diff}(F) - \dim \text{Diff}(F')$.*

For example, let $V^* \cong \mathbb{C}^{n^2}$ be the space of $n \times n$ matrices and let $G = \text{SL}_n \times \text{SL}_n$ act on V^* by left and (inverted) right matrix multiplication. Then G is connected and V^* is an irreducible representation, as is V . Furthermore $\det_n$ is an invariant form, $\text{Diff} \det_n$ is spanned by the minors of all sizes of the matrix, and a first partial derivative of $\det_n$ is the determinant $\det_{n-1}$ of the complementary minor. One can check that $\dim \text{Diff}(\det_n) = \binom{2n}{n}$ and $\dim \text{Diff}(\det_{n-1}) = \binom{2n-2}{n-1}$. This shows that $r(\det_n) \geq \binom{2n}{n} - \binom{2n-2}{n-1}$. For more examples, and more details about determinant, see Section 4.

Our main theorem, Theorem 3.3, loosens the requirement for V to be an irreducible representation, generalizes to an invariant subspace of forms instead of a single invariant form, and actually gives a lower bound for cactus rank (defined below) instead of Waring rank. Also the statement is made coordinate-free.

Section 2 contains some background and basic lemmas. Our main results are in Section 3. We give examples in Section 4.

2. PREPARATION

We review some definitions and basic lemmas.

For a nonzero point p in a vector space W we write $[p]$ for the corresponding point in projective space $\mathbb{P}W$.

2.1. Apolarity. Let V be a $\mathbb{C}$ -vector space with basis $x_1, \dots, x_n$, so $S(V) = \bigoplus_{d=0}^{\infty} S^d V \cong \mathbb{C}[x_1, \dots, x_n]$, where $S(V)$ denotes the symmetric algebra. We introduce the **dual ring** $S(V^*) \cong \mathbb{C}[\partial_1, \dots, \partial_n]$, where $\partial_1, \dots, \partial_n$ is the dual basis for V^* . Then $S(V^*)$ acts on $S(V)$ by differentiation, where each ∂_i acts as $\partial/\partial x_i$. This is the **apolar pairing**. For each degree $d \geq 0$, $S^d V^* \otimes S^d V \rightarrow \mathbb{C}$ is a perfect pairing; for $e > d$, $S^e V^* \otimes S^d V \rightarrow 0$ and $S^d V^* \otimes S^e V \rightarrow S^{e-d} V$.

Let $F \in S(V)$ be a homogeneous form. The **apolar** or **annihilating ideal** $F^\perp \subseteq S(V^*)$ is $F^\perp = \{\Theta \in S(V^*) \mid \Theta F = 0\}$, a homogeneous ideal. The **apolar algebra** is $A^F = S(V^*)/F^\perp$. Note that $A^F \cong \text{Diff}(F)$, as vector spaces. In particular, the length $\ell(A^F) = \dim_{\mathbb{C}} A^F$ is equal to $\dim \text{Diff}(F)$.

One connection between Waring rank and apolarity is given by the following well-known lemma.

Apolarity Lemma. *Let $X \subseteq \mathbb{P}V$ be a scheme with saturated homogeneous defining ideal I_X . Let $\nu_d : \mathbb{P}V \rightarrow \mathbb{P}S^d V$ be the degree d Veronese map, and let $[F] \in \mathbb{P}S^d V$. Then $[F]$ is in the linear span of $\nu_d(X)$ if and only if $I_X \subseteq F^\perp$.*

Here the linear span of a scheme Z is the smallest reduced linear subspace containing Z as a subscheme. Equivalently it is the linear subspace defined by the degree 1 part of the ideal I_Z . Note that if $Z = \nu_d(X)$ then the degree 1 part of I_Z is the degree d part of I_X , interpreted as equations of hyperplanes in $\mathbb{P}S^d V$.

For proofs of the Apolarity Lemma see for example [12, Theorem 5.3], [20, §1.3], [24, §4.1].

A scheme X is called **apolar to F** if $I_X \subseteq F^\perp$.

Suppose $X = \{[\ell_1], \dots, [\ell_r]\} \subseteq \mathbb{P}V$ is a zero-dimensional reduced scheme. Then X is apolar to F if and only if $[F]$ is in the linear span of $\nu_d(X) = \{[\ell_1^d], \dots, [\ell_r^d]\}$, equivalently $F = c_1 \ell_1^d + \dots + c_r \ell_r^d$ for some scalars c_i . Hence the Waring rank $r(F)$ is the least length of a zero-dimensional *reduced* apolar scheme to F . This leads naturally to generalizations; we mention just two:

- (1) The **cactus rank** of F , denoted $cr(F)$, is the least length of *any* zero-dimensional apolar scheme to F . (The name ‘‘cactus rank’’ was introduced in [21].)
- (2) The **smoothable rank** of F , denoted $sr(F)$, is the least length of any zero-dimensional smoothable apolar scheme to F . Recall that a scheme is smoothable if it is a flat limit of smooth schemes. (Note that for a scheme of dimension 0, the notions reduced and smooth are the same.)

Evidently $r(F) \geq sr(F) \geq cr(F)$.

Remark 2.1. Earlier terminology in Definitions 5.1 and 5.66 of [12] is as follows. An apolar scheme is also called an **annihilating scheme**, cactus rank is also called **scheme length**, and smoothable rank is also called **smoothable scheme length**.

2.2. Lower bounds for rank. We have remarked that $r(F) \geq sr(F) \geq cr(F)$. We mention now some well-known lower bounds for rank, only so that we can make comparisons later on with the lower bound in Theorem 3.3.

The **Sylvester** lower bound for rank is:

$$cr(F) \geq \dim A_t^F,$$

the dimension of the t -th graded piece of A^F , for each $0 \leq t \leq d = \deg(F)$.

The **Landsberg–Teitler** lower bound for rank is the following. Assume that F cannot be written using fewer variables (that is, if $F \in S^d V'$ and $V' \subseteq V$ then $V' = V$). Let $\Sigma_t \subseteq V^*$ be the set of points at which F vanishes to order at least $t + 1$. It was shown in [16] that

$$r(F) \geq \dim A_t^F + \dim \Sigma_t.$$

The **Ranestad–Schreyer** lower bound for rank is the following. Let δ be an integer such that the apolar ideal $F^\perp$ is generated in degrees less than or equal to δ . In [21] it was proven that

$$cr(F) \geq \frac{1}{\delta} \ell(A^F).$$

The simplest way to give an upper bound for Waring rank of a form F is just to exhibit an explicit expression for F as a sum of powers. However Bernardi and Ranestad gave an interesting upper bound for cactus rank, as follows. Let l be a linear form and let F_l be a dehomogenization of F with respect to l . Let $\text{Diff}(F_l) \subseteq S$ be the subspace of S spanned by the partial derivatives of F_l of all orders. Then $cr(F) \leq \dim \text{Diff}(F_l)$, see [2, Theorem 1]. It was observed by Shafiei (and Pedro Marques) that $\dim \text{Diff}(F_l) \leq \dim \text{Diff}(F) = \ell(A^F)$, see the remarks following Proposition 3.8 in [23].

2.3. Simultaneous Waring rank. For a linear series $W \subseteq S^d V$, the **simultaneous Waring rank** $r(W)$ is the least r such that there exist linear forms $\ell_1, \dots, \ell_r \in V$ with $W \subseteq \text{span}\{\ell_1^d, \dots, \ell_r^d\}$. The **apolar ideal** $W^\perp \subseteq S(V^*)$ is $W^\perp = \bigcap_{F \in W} F^\perp$. There is an Apolarity Lemma for linear series: for a scheme $Z \subseteq \mathbb{P}V$ with vanishing ideal $I_Z \subseteq S(V^*)$, we have $\mathbb{P}W \subseteq \text{span}(\nu_d(Z))$ if and only if $I_Z \subseteq W^\perp$. As before, the simultaneous Waring rank $r(W)$ is the least length of a reduced zero-dimensional apolar scheme, so we define the **simultaneous smoothable rank** $sr(W)$ to be the least length of a smoothable zero-dimensional apolar scheme, and the **simultaneous cactus rank** $cr(W)$ to be the least length of a zero-dimensional apolar scheme. Evidently $r(W) \geq sr(W) \geq cr(W)$.

The **apolar algebra** is defined by $A^W = S(V^*)/W^\perp$. Let $\text{Diff}(W)$ be the vector subspace of $S(V)$ spanned by all the partial derivatives of all elements of W , of all orders. That is, $\text{Diff}(W) = \sum_{F \in W} \text{Diff}(F)$. As vector spaces, $A^W \cong \text{Diff}(W)$. In particular $\ell(A^W) = \dim \text{Diff}(W)$.

Note that if $W = \mathbb{C}F$ is spanned by a single form then $r(W) = r(F)$, $sr(W) = sr(F)$, $cr(W) = cr(F)$, and $\text{Diff}(W) = \text{Diff}(F)$. Even though our goal (and main interest) is in providing lower bounds for ranks of single forms, it turns out to be equally easy to prove the same lower bounds for simultaneous ranks of linear series; the desired bounds for single forms follow as the special case where $\dim W = 1$.

Each of the lower bounds for rank listed above has an analogue for simultaneous rank. See sections 2.2, 3.2-3, and 5.2 of [24] for a detailed discussion. Briefly, the Sylvester lower bound is

$$cr(W) \geq \dim A_t^W$$

for each $0 \leq t \leq d$, where $W \subseteq S^d V$. There is a generalization of the Landsberg–Teitler lower bound, but it is more complicated and not worth stating here; see [24, §3.2-3]. The Ranestad–Schreyer lower bound is

$$cr(W) \geq \frac{1}{\delta} \ell(A^W),$$

where δ is an integer such that $W^\perp$ is generated in degrees less than or equal to δ .

2.4. Preliminary results. Here are some easy lemmas.

Notation 2.2. For a projective variety $X \subset \mathbb{P}W$, we write $\tilde{X}$ for the affine cone over X . For a homogeneous ideal I , $\mathbb{V}(I)$ is the projective variety or scheme defined by I and $\tilde{\mathbb{V}}(I)$ is the affine variety or scheme defined by I .

If R is a graded ring and M is a graded R -module, then $M(d)$ denotes the module M with shifted grading: $M(d)_e = M_{d+e}$. A homomorphism between graded modules should preserve the grading. If I is an ideal of R and $x \in R$, then the colon ideal $(I : x)$ is defined by $(I : x) = \{f \in R \mid fx \in I\}$.

Lemma 2.3. *Let R be a graded ring, $I \subseteq R$ a homogeneous ideal, $x \in R$ a homogeneous element of degree d . Then the following is a short exact sequence of graded R -modules:*

$$0 \rightarrow (R/(I : x))(-d) \xrightarrow{x} R/I \rightarrow R/(I + (x)) \rightarrow 0.$$

The proof is an easy exercise.

The next lemma is well-known, although usually stated only for the case of a single form ($\dim W = 1$).

Lemma 2.4. *Let $W \subseteq S^d V$ and $\Theta \in S^e V^*$. Let $\Theta W = \{\Theta F \mid F \in W\} \subseteq S^{d-e} V$. Then $(\Theta W)^\perp = (W^\perp : \Theta)$.*

Proof. $\Psi \in (\Theta W)^\perp$ if and only if $\Psi(\Theta F) = 0$ for all $F \in W$, if and only if $\Psi\Theta \in W^\perp$. $\square$

Combining these:

Lemma 2.5. *Let $W \subseteq S^d V$, $\Theta \in S^e V^*$, and $W' = \Theta W$. Then we have a short exact sequence of graded $S(V^*)$ -modules,*

$$0 \rightarrow A^{W'}(-e) \xrightarrow{\Theta} A^W \rightarrow S(V^*)/(W^\perp + \Theta) \rightarrow 0.$$

In particular,

$$\ell(S(V^*)/(W^\perp + \Theta)) = \ell(A^W) - \ell(A^{W'}) = \dim \text{Diff}(W) - \dim \text{Diff}(W').$$

3. MAIN RESULTS

Before giving our main result, we state the following simpler theorem, which does not involve a group action.

Theorem 3.1. *Let $W \subseteq S^d V$ be a linear series of d -forms. There exists a Zariski open dense subset U of V^* such that for $\partial \in U$ (i.e., ∂ in general position) we have*

$$r(W) \geq sr(W) \geq cr(W) \geq \dim \text{Diff}(W) - \dim \text{Diff}(W'),$$

where $W' = \partial W = \{\partial F \mid F \in W\} \subseteq S^{d-1} V$.

Proof. Let $Z \subseteq \mathbb{P}V$ be a zero-dimensional apolar scheme to W of length r , with defining ideal $I_Z \subseteq F^\perp$. Since ∂ is general, the hyperplane $H \subseteq \mathbb{P}V$ defined by ∂ is disjoint from Z .

Now the affine scheme $\tilde{V}(I_Z) \subseteq V$ is one-dimensional and has no component contained in the hyperplane $\tilde{H} \subseteq V$. So $\tilde{V}(I_Z) \cap \tilde{H}$ is supported only at the origin and has length equal to $\ell(Z) = r$. That is, $\ell(S(V^*)/(I_Z + \partial)) = r$. Since $I_Z \subseteq W^\perp$ we have $\ell(S(V^*)/(W^\perp + \partial)) \leq r$. This shows that $r(W) \geq sr(W) \geq cr(W) \geq \ell(S(V^*)/(W^\perp + \partial))$. By Lemma 2.5, $\ell(S(V^*)/(W^\perp + \partial)) = \ell(A^W) - \ell(A^{W'}) = \dim \text{Diff}(W) - \dim \text{Diff}(W')$. $\square$

For a single form, $W = \mathbb{C}F$, the proof would be the same, just writing F and F' instead of W and W' throughout.

Remark 3.2. The proof of the Ranestad–Schreyer bound uses a dual form $\Theta \in F^\perp$ of degree δ , and general in the linear series $(F^\perp)_\delta$, so that Θ does not vanish at any point of Z . This requires δ to be large enough so that $(F^\perp)_\delta$ has no basepoints. In this setting, $F' = \Theta F = 0$.

A general dual linear form $\partial \in V^*$ also does not vanish at any point of Z . Lowering the degree from δ to 1 accounts for the improvement by a factor of δ in the above result, compared to the Ranestad–Schreyer bound. On the other hand, $F' = \partial F \neq 0$.

Now we state our main theorem. We make essentially the same argument, except that instead of a general hyperplane we will use a translation of a given hyperplane by a general group element.

Theorem 3.3. *Let G be a connected algebraic group and V a representation of G . Let $W \subseteq S^d V$ be an invariant subspace, i.e., ${}^g F \in W$ for all $F \in W$, $g \in G$. Let $\partial \in V^*$ be a nonzero element that is not contained in any proper subrepresentation. Let $W' = \partial W = \{\partial F \mid F \in W\} \subseteq S^{d-1} V$. Then*

$$r(W) \geq sr(W) \geq cr(W) \geq \dim \text{Diff}(W) - \dim \text{Diff}(W').$$

Proof. If the orbit $G\partial$ were dense in V^* , the result would follow immediately from Theorem 3.1, as ${}^g \partial \in V^*$ would be general for $g \in G$ general. But if the orbit $G\partial$ is closed, then a priori the orbit might completely miss the “general” open set of Theorem 3.1. So we have to argue more directly.

Suppose $Z \subseteq \mathbb{P}V$ is a zero-dimensional apolar scheme to W of length $\ell(Z) = r$. Let the support of Z be $\{p_1, \dots, p_t\}$. Let $H \subseteq \mathbb{P}V$ be the hyperplane defined by $\partial = 0$. We claim that for general $g \in G$, ${}^g Z$ is disjoint from H .

The condition ∂ is not contained in any subrepresentation is equivalent to requiring that the orbit $G\partial$ spans V^* . Then for each $i = 1, \dots, t$ there is a $g_i \in G$ such that ${}^{g_i} \partial$ does not vanish at the point p_i ; that is, the hyperplane ${}^{g_i} H$ does not contain p_i . Equivalently, ${}^{g_i^{-1}} p_i \notin H$. So there is a nonempty open subset $U_i \subseteq G$ such that ${}^g p_i \notin H$ for $g \in U_i$. Since G is connected (hence irreducible) the intersection $U = \bigcap U_i$ is a nonempty dense open set. This shows that for general elements $g \in G$ we have ${}^g Z \subseteq \mathbb{P}V \setminus H$.

Note that $W = {}^g W \subseteq \text{span}({}^g Z)$ for every $g \in G$. Now replace Z with ${}^g Z$ for a general $g \in G$. The rest of the proof is the same as the proof of Theorem 3.1. Explicitly, let $I_Z \subseteq W^\perp$ be the defining ideal of Z . The affine scheme $\tilde{V}(I_Z)$ is one-dimensional, has degree r , and has no component contained in $\tilde{H}$. Thus $\tilde{V}(I_Z) \cap \tilde{H}$ is supported only at the origin and has length equal to r . Hence $\ell(S(V^*)/(W^\perp + \partial)) \leq \ell(S(V^*)/(I_Z + \partial)) = r$. By Lemma 2.5, $\ell(S(V^*)/(W^\perp + \partial)) = \dim \text{Diff}(W) - \dim \text{Diff}(W')$. $\square$

Remark 3.4. If V is an irreducible representation of G , then every nonzero $\partial \in V^*$ meets the condition (of not being contained in a subrepresentation).

Recall that a form F is **semi-invariant** if for every $g \in G$ there is a nonzero scalar $\chi(g)$ such that ${}^gF = \chi(g)F$. Then F is semi-invariant if and only if its span $W = \mathbb{C}F$ is an invariant subspace. We obtain

Corollary 3.5. *Let G be a connected algebraic group and V a representation of G . Let $F \in S^dV$ be a semi-invariant form. Let $\partial \in V^*$ be a nonzero element that is not contained in any proper subrepresentation. Let $F' = \partial F$. Then*

$$r(F) \geq sr(F) \geq cr(F) \geq \dim \text{Diff}(F) - \dim \text{Diff}(F').$$

Here is a simple and crude lower bound.

Corollary 3.6. *Let $F \in S^dV$ be a nonzero homogeneous form. Fix a basis $\{x_1, \dots, x_n\}$ for V and the dual basis $\{\partial_1, \dots, \partial_n\}$ for V^* . Let $F = F_k x_1^k + F_{k-1} x_1^{k-1} + \dots + F_0$, $F_i \in \mathbb{C}[x_2, \dots, x_n]_{d-i}$. Suppose either that $\partial_1 \in V^*$ is general, or else that V is a representation of a connected group G , F is a semi-invariant of G , and ∂_1 lies in no proper subrepresentation of V^* . Then $r(F) \geq sr(F) \geq cr(F) \geq \dim \text{Diff}(F_k)$.*

Proof. Since $F_k = (1/k!) \partial_1^k F$ we have $F_k^\perp = (F^\perp : \partial_1^k)$ by Lemma 2.4. Then clearly $F^\perp \subseteq (F^\perp : \partial_1^k) = F_k^\perp$. And since $\partial_1^{k+1} \in F^\perp$, we have $(\partial_1) \subseteq F_k^\perp$ as well (or: $\partial_1 \in F_k^\perp$ since F_k does not involve x_1). So $F^\perp + (\partial_1) \subseteq F_k^\perp$, which implies $\ell(S(V^*)/(F^\perp + \partial_1)) \geq \ell(S(V^*)/F_k^\perp) = \dim \text{Diff}(F_k)$.

Finally, we have $cr(F) \geq \ell(S(V^*)/(F^\perp + \partial_1))$ in either case, ∂_1 is general or F is a semi-invariant, by the proofs of Theorems 3.1 and 3.3. $\square$

Compare this with the Bernardi–Ranestad upper bound for cactus rank in terms of dehomogenization:

$$\dim \text{Diff}(F_k) \leq cr(F) \leq \dim \text{Diff}(F_k + F_{k-1} + \dots + F_0).$$

4. EXAMPLES

Example 4.1. Let $V^* \cong \mathbb{C}^{n^2}$ be the space of $n \times n$ matrices. Let V have basis $\{x_{i,j} \mid 1 \leq i, j \leq n\}$ and V^* have the dual basis $\{\partial_{i,j}\}$. Let $\det_n$ be the generic $n \times n$ determinant:

$$\det_n = \det \begin{pmatrix} x_{1,1} & \cdots & x_{1,n} \\ \vdots & & \vdots \\ x_{n,1} & \cdots & x_{n,n} \end{pmatrix},$$

so $\det_n \in S^dV$ is a form of degree n in n^2 variables. Recall that the derivatives of $\det_n$ are spanned by minors. This shows $\dim(S(V^*)/\det_n^\perp)_t = \binom{n}{t}^2$, so

$$\ell(A^{\det_n}) = \sum_{t=0}^n \binom{n}{t}^2 = \binom{2n}{n}.$$

As mentioned in the introduction, the Sylvester bound shows that the cactus rank and border rank of $\det_n$ are bounded below by $\binom{n}{\lfloor n/2 \rfloor}^2$. Shafiei has shown that $\det_n^\perp$ is generated by quadrics, so the Ranestad–Schreyer bound gives $cr(\det_n) \geq \frac{1}{2} \binom{2n}{n}$.

Now $\det_n$ is invariant under the action of $\text{SL}_n \times \text{SL}_n$ on $\mathbb{C}^{n^2}$ by left and (inverted) right matrix multiplication. And the orbit of $\partial_{1,1}$ spans V^* ; in fact, the subgroup of permutation

matrices already takes $\partial_{1,1}$ to all the $\partial_{i,j}$, a basis for V^* . (In any case, this is an irreducible representation.) Since $\partial_{1,1} \det_n$ is the complementary $(n-1)$ -minor we have

$$\ell(S(V^*)/(\partial_{1,1} \det_n)^\perp) = \ell(A^{\det_{n-1}}) = \binom{2n-2}{n-1}.$$

Therefore by Theorem 3.3,

$$r(\det_n) \geq sr(\det_n) \geq cr(\det_n) \geq \binom{2n}{n} - \binom{2n-2}{n-1}.$$

Note that

$$\binom{2n-2}{n-1} = \frac{n^2}{(2n-1)(2n)} \binom{2n}{n} \approx \frac{1}{4} \binom{2n}{n},$$

so $\binom{2n}{n} - \binom{2n-2}{n-1} \approx \frac{3}{4} \binom{2n}{n}$. Hence this lower bound is asymptotically $\frac{3}{2}$ times the Ranestad–Schreyer–Shafiei bound.

See Table 1 for some values of this bound and a comparison to other bounds. The upper bound for Waring rank in this table is from [10, §8].

The upper bound for cactus rank in table 1 is the Bernardi–Ranestad upper bound. Let f be the dehomogenization of $\det_n$ with respect to $x_{n,n}$. Then $\dim \text{Diff}(f) = \dim \text{Diff}(\det_n) - 2$; indeed, derivatives of f are obtained by dehomogenizing the corresponding derivatives of $\det_n$, except that $\partial_{n,n} f = 0$ and $x_{n,n} \notin \text{Diff}(f)$. So $cr(\det_n) \leq \dim \text{Diff}(\det_n) - 2 = \binom{2n}{n} - 2$.

n	2	3	4	5	6	7	8
Sylvester	4	9	36	100	400	1225	4900
Landsberg–Teitler	4	14	43	116	420	1258	4939
Ranestad–Schreyer–Shafiei	3	10	35	126	462	1716	6435
Theorem 3.3	4	14	50	182	672	2508	9438
Upper bound for $cr(\det_n)$	4	18	68	250	922	3430	12868
Upper bound for $r(\det_n)$	4	20	160	1600	16000	224000	3584000

TABLE 1. Comparison of bounds for rank of determinant

Example 4.2. Let X be a generic $(2n) \times (2n)$ skew-symmetric matrix, that is, $X = (x_{i,j})$ such that $x_{i,j} = -x_{j,i}$ and $x_{i,i} = 0$. The Pfaffian of X is a polynomial of degree n in the entries of X , which we denote pf_n or $\text{pf}(X)$, with the property that $\text{pf}_n^2 = \det(X)$. For $n = 1, 2$ we have

$$\text{pf}_1 = \text{pf} \begin{pmatrix} 0 & x_{1,2} \\ -x_{1,2} & 0 \end{pmatrix} = x_{1,2},$$

and

$$\text{pf}_2 = \text{pf} \begin{pmatrix} 0 & x_{1,2} & x_{1,3} & x_{1,4} \\ -x_{1,2} & 0 & x_{2,3} & x_{2,4} \\ -x_{1,3} & -x_{2,3} & 0 & x_{3,4} \\ -x_{1,4} & -x_{2,4} & -x_{3,4} & 0 \end{pmatrix} = x_{1,2}x_{3,4} - x_{1,3}x_{2,4} + x_{1,4}x_{2,3}.$$

In general,

$$\text{pf}_n = \sum (-1)^\sigma x_{\sigma(1),\sigma(2)} \cdots x_{\sigma(2n-1),\sigma(2n)},$$

the sum over permutations $\sigma \in S_{2n}$ such that $\sigma(2i-1) < \sigma(2i)$ for all i and $\sigma(1) < \sigma(3) < \dots < \sigma(2n-1)$, equivalently over unordered partitions of $\{1, \dots, 2n\}$ into pairs. Note, there are $(2n-1)!! = (2n)!/(2^n n!)$ such partitions. There is a ‘‘Laplace expansion’’: for each j , $1 \leq j \leq n$,

$$\text{pf}(X) = \sum_{i < j} (-1)^{i+j+1} x_{i,j} \text{pf}(X^{i,j}) + \sum_{i > j} (-1)^{i+j} x_{i,j} \text{pf}(X^{i,j}),$$

where $X^{i,j}$ is the matrix obtained by deleting the i th and j th rows and columns of X . See [3, 11, 13].

Note that pf_n is invariant under the conjugation action of SO_{2n} on the space of skew-symmetric matrices. This is an irreducible representation of a connected group, so Theorem 3.3 applies.

By the Laplace expansion, $\partial_{n-1,n} \text{pf}_n$ is the Pfaffian of the $(2n-2) \times (2n-2)$ skew-symmetric matrix obtained by deleting the $(2n-1)$ -st and $(2n)$ -th rows and columns of X ; we may regard $\partial_{n-1,n} \text{pf}_n = \text{pf}_{n-1}$.

More generally, all the derivatives of pf_n are spanned by Pfaffians of even-sized principal (i.e., skew-symmetric) submatrices of X . So

$$\dim A_t^{\text{pf}_n} = \binom{2n}{2t}$$

for all $0 \leq t \leq n$. Therefore the Sylvester bound shows that the cactus rank and border rank of pf_n are bounded below by $\binom{2n}{2\lfloor n/2 \rfloor}$. And

$$\ell(A^{\text{pf}_n}) = \sum_{t=0}^n \binom{2n}{2t} = 2^{2n-1}.$$

Shafiei has shown that $\text{pf}_n^\perp$ is generated by quadrics [23, Theorem 4.11], so by the Ranestad–Schreyer bound, $cr(\text{pf}_n) \geq 2^{2n-2}$. Finally, Theorem 3.3 gives

$$r(\text{pf}_n) \geq sr(\text{pf}_n) \geq cr(\text{pf}_n) \geq \ell(A^{\text{pf}_n}) - \ell(A^{\text{pf}_{n-1}}) = 2^{2n-1} - 2^{2n-3} = 3 \cdot 2^{2n-3}.$$

This is exactly $\frac{3}{2}$ times the Ranestad–Schreyer–Shafiei bound.

Some values of these bounds are shown in Table 2. The upper bound for Waring rank in this table comes from the expression of pf_n as a sum of $(2n)!/2^n n!$ terms each of rank 2^{n-1} .

The upper bound for cactus rank is Shafiei’s loosening of the Bernardi–Ranestad upper bound: $cr(\text{pf}_n) \leq \ell(A^{\text{pf}_n}) = 2^{2n-1}$. This can be improved by considering a dehomogenization. (In particular, this loosening is the reason we get a worse bound for $cr(\text{pf}_2)$ than the bound for Waring rank $r(\text{pf}_2)$.)

n	2	3	4	5	6	7	8
Sylvester	6	15	70	210	924	3003	12870
Ranestad–Schreyer–Shafiei	4	16	64	256	1024	4096	16384
Theorem 3.3	6	24	96	384	1536	6144	24576
Upper bound for $cr(\text{pf}_n)$	8	32	128	512	2048	8192	32768
Upper bound for $r(\text{pf}_n)$	6	60	840	15120	332640	8468640	259459200

TABLE 2. Comparison of bounds for rank of Pfaffian

Example 4.3. Let sdet_n be the determinant of a generic symmetric matrix, that is a matrix $X = (x_{i,j})$ such that $x_{i,j} = x_{j,i}$. For example $\text{sdet}_1 = x_{1,1}$,

$$\text{sdet}_2 = \det \begin{pmatrix} x_{1,1} & x_{1,2} \\ x_{1,2} & x_{2,2} \end{pmatrix} = x_{1,1}x_{2,2} - x_{1,2}^2,$$

and so on.

Note that sdet_n is invariant under the action of SL_n on the space of symmetric matrices given by $(A, M) \mapsto AMA^t$. This is an irreducible representation of a connected group, so Theorem 3.3 applies.

Shafiei has shown

$$\dim A_t^{\text{sdet}_n} = \frac{1}{n+1} \binom{n+1}{t} \binom{n+1}{t-1},$$

see [22, Lemma 2.5]. Therefore the Sylvester bound shows that the cactus rank and border rank of sdet_n are bounded below by $\frac{1}{n+1} \binom{n+1}{\lceil n/2 \rceil} \binom{n+1}{\lceil n/2 \rceil - 1}$. And

$$\ell(A^{\text{sdet}_n}) = \frac{1}{n+2} \binom{2n+2}{n+1},$$

the $(n+1)$ -st Catalan number C_{n+1} , see [22, Corollary 2.6]. Shafiei has shown that sdet_n is generated by quadrics [22, Theorem 3.11]. So by the Ranestad–Schreyer bound, $cr(\text{sdet}_n) \geq \frac{1}{2(n+2)} \binom{2n+2}{n+1}$, see [22, Theorem 4.6].

Since $\partial_{n,n} \text{sdet}_n = \text{sdet}_{n-1}$, Theorem 3.3 gives

$$r(\text{sdet}_n) \geq sr(\text{sdet}_n) \geq cr(\text{sdet}_n) \geq C_{n+1} - C_n = \frac{1}{n+2} \binom{2n+2}{n+1} - \frac{1}{n+1} \binom{2n}{n}.$$

This is asymptotically $\frac{3}{2}$ times the Ranestad–Schreyer–Shafiei bound.

Some values of these bounds are shown in Table 3. The upper bound for cactus rank is Shafiei’s loosening of the Bernardi–Ranestad upper bound, $cr(\text{sdet}_n) \leq \ell(A^{\text{sdet}_n}) = \frac{1}{n+2} \binom{2n+2}{n+1}$; this can be improved by considering a dehomogenization.

n	2	3	4	5	6	7	8
Sylvester	3	6	20	50	175	490	1764
Ranestad–Schreyer–Shafiei	2.5	7	21	66	214.5	715	2431
Theorem 3.3	3	9	28	90	297	1001	3432
Upper bound for $cr(\text{sdet}_n)$	5	14	42	132	429	1430	4862

TABLE 3. Comparison of bounds for rank of symmetric determinant

Example 4.4. Let V^* be the space of $m \times n$ matrices, $m \leq n$. Fix the standard basis $\{x_{i,j} \mid 1 \leq i \leq m, 1 \leq j \leq n\}$ for V and the dual basis $\{\partial_{i,j}\}$ for V^* , so $x_{i,j}$ applied to a matrix is its (i,j) entry. Let X be the generic $m \times n$ matrix $X = (x_{i,j})$. Fix an integer d , $1 \leq d \leq m \leq n$, and let W_d be the linear series spanned by the d -minors of X . Then $\text{Diff}(W_d)$ is spanned by all the t -minors of X for $0 \leq t \leq d$, so

$$\dim \text{Diff}(W_d) = \sum_{t=0}^d \binom{m}{t} \binom{n}{t} = \sum_{t=0}^d \binom{m}{m-t} \binom{n}{t}.$$

In particular if $d = m \leq n$ (the case of maximal minors) then $\dim \text{Diff}(W_m) = \binom{m+n}{m}$.

Shafiei's results imply that $W_d^\perp$ is generated by quadrics. The Ranestad–Schreyer lower bound thus gives

$$cr(W_d) \geq \frac{1}{2} \sum_{t=0}^d \binom{m}{t} \binom{n}{t},$$

in particular if $d = m \leq n$ then $cr(W_m) \geq \frac{1}{2} \binom{m+n}{n}$.

Now, W_d is invariant under the action of $\text{SL}_m \times \text{SL}_n$ on V^* by left and (inverted) right matrix multiplication. Note that $W' = \partial_{1,1} W_d$ is exactly the linear series of $(d-1)$ -minors of the $(m-1) \times (n-1)$ matrix obtained by deleting the first row and column of X . So

$$\dim \text{Diff}(W') = \sum_{t=0}^{d-1} \binom{m-1}{t} \binom{n-1}{t},$$

in particular if $d = m \leq n$ then $\dim \text{Diff}(W') = \binom{m+n-2}{m-1}$. Thus

$$cr(W_d) \geq \sum_{t=0}^d \left(\binom{m}{t} \binom{n}{t} - \binom{m-1}{t-1} \binom{n-1}{t-1} \right)$$

in general (with the understanding $\binom{m-1}{-1} = \binom{n-1}{-1} = 0$), and for maximal minors, if $d = m \leq n$ then

$$cr(W_m) \geq \binom{m+n}{m} - \binom{m+n-2}{m-1}.$$

Example 4.5. Let $V_1^* \cong \mathbb{C}^{pq}$ be the space of $p \times q$ matrices, $V_2^* \cong \mathbb{C}^{qr}$ be the space of $q \times r$ matrices, and $V_3^* \cong \mathbb{C}^{rp}$ be the space of $r \times p$ matrices. We denote by $\{x_{i,j}\}_{1 \leq i \leq p, 1 \leq j \leq q}$ the standard basis of V_1 , so $x_{i,j}$ applied to a matrix is its (i, j) entry. Similarly the standard basis of V_2 is denoted $\{y_{i,j}\}_{1 \leq i \leq q, 1 \leq j \leq r}$ and the standard basis of V_3 is denoted $\{z_{i,j}\}_{1 \leq i \leq r, 1 \leq j \leq p}$. Consider the tensor

$$T_{p,q,r} = \sum_{i,j,k} x_{i,j} y_{j,k} z_{k,i} \in \mathbb{C}^{pq} \otimes \mathbb{C}^{qr} \otimes \mathbb{C}^{rp}.$$

This represents the *matrix multiplication* map $V_1 \otimes V_2 \rightarrow V_3$. We will give a lower bound for the tensor rank of $T_{p,q,r}$.

Set $V = V_1 \oplus V_2$. Let $W \subseteq S^2 V$ be the space spanned by all $\sum_{j=1}^q x_{i,j} y_{j,k}$ with $1 \leq i \leq p$ and $1 \leq k \leq r$. Note that these are the entries of the $p \times r$ matrix obtained by multiplying the matrices $(x_{i,j})_{1 \leq i \leq p, 1 \leq j \leq q}$ and $(y_{i,j})_{1 \leq i \leq q, 1 \leq j \leq r}$.

The tensor rank $tr(T_{p,q,r})$ of the tensor $T_{p,q,r}$ is the smallest number m for which there exist pure tensors $f_i(x, y) = g_i(x)h_i(y) \in V_1 \otimes V_2$, $i = 1, 2, \dots, m$ such that W is contained in the span of $f_1(x, y), \dots, f_m(x, y)$ [14, Thm. 3.1.1.1]. We can write $f_i(x, y) = \frac{1}{4}((g_i(x) + h_i(y))^2 - (g_i(x) - h_i(y))^2)$, so $tr(T_{p,q,r}) \geq \frac{1}{2}r(W)$. On the other hand, if W is contained in the space spanned by $u_1(x, y)^2, \dots, u_m(x, y)^2$ where $u_1, \dots, u_m$ are linear, then we can write $u_i(x) = v_i(x) + w_i(y)$ and W is contained in the span of $v_i(x)w_i(y)$, $i = 1, 2, \dots, m$ because W consists of forms that are bilinear in x and y . This shows that $tr(T_{p,q,r}) \leq r(W)$. We conclude that

$$\frac{1}{2}r(W) \leq tr(T_{p,q,r}) \leq r(W).$$

There is a natural action of $G = \text{GL}_p \times \text{GL}_q \times \text{GL}_r$ on V . The space W is invariant under the action of G . Let $\partial = \frac{\partial}{\partial x_{1,1}} + \frac{\partial}{\partial y_{1,1}} \in V^*$. The space $V^* \cong V_1^* \oplus V_2^*$ is the sum of

two non-isomorphic irreducible representations of G . The only proper invariant subspaces of V^* are V_1^* and V_2^* and ∂ does not lie in either of them. The space $\text{Diff}(W)$ is spanned by W , all $x_{i,j}$, all $y_{i,j}$, and $\mathbb{C}$. The dimension of W is pr while the span of all $x_{i,j}$ and all $y_{i,j}$ has dimension $pq + qr$. So the dimension of $\text{Diff}(W)$ is $pq + qr + pr + 1$. The space W' is spanned by all $y_{1,k}$ for $k > 1$, $x_{i,1}$ for $i > 1$ and $x_{1,1} + y_{1,1}$. The dimension of W' is $(p-1) + (r-1) + 1 = p + r - 1$, and the dimension of $\text{Diff}(W')$ is $p + r$. We have

$$\begin{aligned} r(W) &\geq sr(W) \geq cr(W) \geq \dim \text{Diff}(W) - \dim \text{Diff}(W') = \\ &= (pq + qr + pr + 1) - (p + r) = pq + qr + pr - p - r + 1. \end{aligned}$$

It follows that

$$\text{tr}(T_{p,q,r}) \geq \frac{1}{2}(pq + qr + pr - p - r + 1).$$

If $p = q = r = n$ then we get $r(W) \geq 3n^2 - 2n + 1$. It follows that

$$\text{tr}(T_{n,n,n}) \geq \frac{3}{2}n^2 - n + \frac{1}{2}.$$

This is a nontrivial lower bound for the tensor rank of $T_{n,n,n}$, but there are better bounds known: Recently, Massarenti and Raviolo ([18, 19, 17]) improved a lower bound of Landsberg ([15]). Both lower bounds grow asymptotically as $3n^2 - o(n^2)$. From these bounds we also get a lower bound for $r(W)$, namely $r(W) \geq 3n^2 - o(n^2)$, which is slightly worse than our bound.

REFERENCES

1. J. Alexander and A. Hirschowitz, *Polynomial interpolation in several variables*, J. Algebraic Geom. **4** (1995), no. 2, 201–222. MR 1311347 (96f:14065)
2. Alessandra Bernardi and Kristian Ranestad, *On the cactus rank of cubic forms*, J. Symbolic Comput. **50** (2013), 291–297. MR 2996880
3. David A. Buchsbaum and David Eisenbud, *Algebra structures for finite free resolutions, and some structure theorems for ideals of codimension 3*, Amer. J. Math. **99** (1977), no. 3, 447–485. MR 0453723 (56 #11983)
4. Enrico Carlini, Maria Virginia Catalisano, and Luca Chiantini, *Progress on the symmetric Strassen conjecture*, [arXiv:1405.3721](#) [math.AG], May 2014.
5. Enrico Carlini, Maria Virginia Catalisano, Luca Chiantini, Anthony V. Geramita, and Youngho Woo, *Symmetric Tensors: Rank and Strassen’s Conjecture*, [arXiv:1412.2975](#) [math.AC], Dec 2014.
6. Enrico Carlini, Maria Virginia Catalisano, Luca Chiantini, Anthony V. Geramita, and Youngho Woo, *e-computable forms and the Strassen conjecture*, [arXiv:1502.01107](#) [math.AC], Feb 2015.
7. Enrico Carlini, Maria Virginia Catalisano, and Anthony V. Geramita, *The solution to the Waring problem for monomials and the sum of coprime monomials*, J. Algebra **370** (2012), 5–14.
8. Pierre Comon, Gene Golub, Lek-Heng Lim, and Bernard Mourrain, *Symmetric tensors and symmetric tensor rank*, SIAM J. Matrix Anal. Appl. **30** (2008), no. 3, 1254–1279. MR 2447451 (2009i:15039)
9. Pierre Comon and Bernard Mourrain, *Decomposition of quantics in sums of powers of linear forms*, Signal Processing **53** (1996), no. 2–3, 93–107.
10. Harm Derksen, *On the nuclear norm and the singular value decomposition of tensors*, [arXiv:1308.3860](#) [math.OC], Aug 2013.
11. P. Heymans, *Pfaffians and skew-symmetric matrices*, Proc. London Math. Soc. (3) **19** (1969), 730–768. MR 0257105 (41 #1759)
12. Anthony Iarrobino and Vassil Kanev, *Power sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics, vol. 1721, Springer-Verlag, Berlin, 1999, Appendix C by Iarrobino and Steven L. Kleiman. MR 1735271 (2001d:14056)
13. Tadeusz Józefiak and Piotr Pragacz, *Ideals generated by Pfaffians*, J. Algebra **61** (1979), no. 1, 189–198. MR 554859 (81e:13005)

14. J. M. Landsberg, *Tensors: geometry and applications*, Graduate Studies in Mathematics, vol. 128, American Mathematical Society, Providence, RI, 2012. MR 2865915
15. ———, *New lower bounds for the rank of matrix multiplication*, SIAM J. Comput. **43** (2014), no. 1, 144–149. MR 3162411
16. J.M. Landsberg and Zach Teitler, *On the ranks and border ranks of symmetric tensors*, Found. Comp. Math. **10** (2010), no. 3, 339–366.
17. Alex Massarenti and Emanuele Raviolo, *On the rank of $n \times n$ matrix multiplication*, [arXiv:1211.6320](#) [cs.CC].
18. ———, *The rank of $n \times n$ matrix multiplication is at least $3n^2 - 2\sqrt{2}n^{\frac{3}{2}} - 3n$* , Linear Algebra Appl. **438** (2013), no. 11, 4500–4509. MR 3034546
19. ———, *Corrigendum to “The rank of $n \times n$ matrix multiplication is at least $3n^2 - 2\sqrt{2}n^{\frac{3}{2}} - 3n$ ”* [*Linear Algebra Appl.* 438 (11) (2013) 4500–4509] [[mr3034546](#)], Linear Algebra Appl. **445** (2014), 369–371. MR 3151280
20. Kristian Ranestad and Frank-Olaf Schreyer, *Varieties of sums of powers*, J. Reine Angew. Math. **525** (2000), 147–181. MR 1780430 (2001m:14009)
21. ———, *On the rank of a symmetric form*, J. Algebra **346** (2011), 340–342. MR 2842085
22. Masoumeh Sepideh Shafiei, *Apolarity for determinants and permanents of generic symmetric matrices*, [arXiv:1303.1860](#) [math.AC], Mar 2013.
23. ———, *Apolarity for determinants and permanents of generic matrices*, To appear in Journal of Commutative Algebra, 2014.
24. Zach Teitler, *Geometric lower bounds for generalized ranks*, [arXiv:1406.5145](#) [math.AG], Jun 2014.
25. Youngho Woo, *Some cases on Strassen additive conjecture*, [arXiv:1406.2213](#) [math.AG], Jun 2014.

HARM DERKSEN, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET,
ANN ARBOR, MI 48109–1043
E-mail address: hderksen@umich.edu

ZACH TEITLER, BOISE STATE UNIVERSITY, DEPARTMENT OF MATHEMATICS, 1910 UNIVERSITY DRIVE,
BOISE, ID 83725–1555
E-mail address: zzeitler@boisestate.edu