Dynamic carbonate sedimentation on the Northern Line Islands Ridge, Palmyra Basin

May 2016 draft

5 Mitchell Lyle (Corresponding Author)
College of Earth Ocean and Atmospheric Sciences, Oregon State University
mlyle@coas.oregonstate.edu

Robert Pockalny
10 Graduate School of Oceanography, University of Rhode Island
rpockalny@uri.edu

Pratigya Polissar
Lamont-Doherty Earth Observatory, Columbia University
15 Polissar@ldeo.columbia.edu

Jean Lynch-Stieglitz
School of Earth and Atmospheric Science, Georgia Institute of Technology
Jean@eas.gatech.edu

20 Samantha Bova
Department of Earth Environmental and Planetary Sciences, Brown University
Samantha_Bova@Brown.edu

25 Ann G. Dunlea
Department of Earth and Environment, Boston University
adunlea@bu.edu

Heather Ford
30 Lamont-Doherty Earth Observatory, Columbia University
hford@ldeo.columbia.edu

Jennifer Hertzberg
Department of Oceanography, Texas A&M University, now at
35 Marine Sciences Program, University of Connecticut
jennifer.hertzberg@uconn.edu

Steven Hovan
Department of Geoscience, Indiana University of Pennsylvania
40 hovan@iup.edu

Allison Jacobel
Lamont-Doherty Earth Observatory, Columbia University

jacobel@ldeo.columbia.edu

45

Christina King Wertman
Graduate School of Oceanography, University of Rhode Island
c_king@gso.uri.edu

50

Ashley E. Maloney
University of Washington, School of Oceanography
ashjames@uw.edu

55

Richard W. Murray
Department of Earth and Environment, Boston University
rickm@bu.edu

60

Julia Shackford Wilson
Department of Oceanography, Texas A&M University
juliakwilson.13@gmail.com

65

Katherine Wejnert
Fernback Science Center, Atlanta Georgia
katherine_e_wejnert@dekalbschoolsga.org

70

Ruifang C. Xie
Dept of Biogeochemistry, Max Planck Institute for Chemistry, Mainz Germany
ruifang.xie@mpic.de

75

80 **1.0 ABSTRACT:**

The Line Islands Ridge (LIR), located south of the Hawaiian Islands between 7°N and 1° S, is one of the few large central Pacific regions shallower than the regional carbonate compensation depth. Thick sequences of carbonate sediments have

accumulated around the LIR despite it being located in the sediment-starved central
85 tropical Pacific. The LIR is an important source of carbonates to the surrounding
region and deposition around the LIR has expanded the equatorial Pacific carbonate
sediment tongue by about 5% of its total area. Furthermore, sediments on the ridge
are potentially important paleoceanographic archives. A recent survey at the crest
of the LIR finds evidence for high current activity, significant erosion, but overall net
90 sediment deposition. Currents are strong enough to form sediment waves and lee
drifts in the Palmyra Basin, at the northern terminus of the LIR. Sediments along the
LIR are pelagic foraminiferal sands that are easily eroded and flow out into the
surrounding abyssal plain in active submarine channel systems. As channels
migrate, pelagic sediments fill in the abandoned channel arms. Despite significant
95 sediment losses from the top of the ridge, 1.3 km of sediment has accumulated in the
upper Palmyra Basin over basement formed 68 to 85 million years ago (Ma). Late
Neogene erosion may be more extensive than earlier erosion cycles, in response to
reduced sediment production as the Palmyra Basin exited the high productivity
equatorial latitudes. Sediments with good stratigraphic order needed for
100 paleoceanographic study are limited in this dynamic sedimentary environment, but
can be found with proper survey.

KEYWORDS

Central Pacific, carbonate sediments, Line Islands Ridge, seismic stratigraphy,

105 carbon cycle

2.0 REGIONAL SETTING:

The Line Islands Ridge (LIR, **Figure 1**) is one of the few large areas in the central Pacific that stands above the regional carbonate compensation depth of ~4800 m (Berger et al., 1976). Elevated topography associated with the LIR extends ~1000 km southeast from 7°N to ~1°S. The highest peaks of the LIR form the Line Island chain, from Kingman Reef to Kiritimati (Christmas) Island (**Figure 1**). The width of the LIR above the 3000 m depth contour varies from ~200 km in the north to < 100 km at its southern end.

Sediments on the LIR should prove to be an important archive to study Pleistocene glacial-interglacial changes in surface ocean conditions (Lynch-Stieglitz et al, 2015) as well as the Cenozoic evolution of the Pacific Ocean and circumpolar deep-water flow in the abyssal Pacific with additional scientific drilling. The high topography of the LIR provides a platform sufficiently shallow for carbonate records to exist on Cretaceous-age crust. The LIR also spans across the central Pacific equatorial region to provide transects needed to study the paleoceanographic evolution of the central Pacific.

Seismic stratigraphy around the LIR can potentially be used to understand changes in deep Pacific flow through changes in regional sediment deposition and formation of hiatus intervals. Circumpolar deep water (CPDW) now flows northward in the western Pacific, and splits to flow around the Hawaiian Islands (Talley, 2007). CPDW flowing east is blocked by the LIR, but CPDW eventually flows around it to the north between the LIR and Hawaii. A branch of bottom water

returning from the eastern Pacific flows westward along the equator to the south of
130 the LIR (Johnson and Toole, 1993).

2.1 Origin of LIR basement topography

Originally the LIR was thought to be a hotspot trace because the ridge
extends in the general direction of Pacific Plate motion (Morgan, 1972). However
dating of basalts along the chain failed to identify an age progression with distance
135 (Schlanger et al, 1984). Much of the northern Ridge was built during two major
periods of late Cretaceous volcanism, 81-86 Ma and 68-73 Ma (Davis et al., 2002) on
ocean crust that was formed at about 120 Ma. Davis et al (2002) proposed that LIR
volcanism was associated with a broad upwarping of mantle in the South Pacific
Superswell region (Larson, 1991, Adam and Bonneville, 2005). A more recent
140 interpretation (Pockalny et al., 2015) suggests the LIR may result from two small
overlapping hotspot traces, one associated with the Crough Hotspot and one with a
possible hotspot near Pukapuka Ridge.

LIR topography is also different from that of Hawaii, where eruptions have
been restricted to relatively few volcanic centers (**Figure 1**) so that large but
145 separate volcanic edifices are formed. Instead, the LIR is composed of multiple
smaller volcanoes, spread over 1-2° of longitude across the general LIR trend. In the
region between 5° and 7° N, the LIR bathymetry swells to form the largest region of
shallow bathymetry in the chain (**Figure 2**). The high topography consists of two
bounding basaltic ridges to the southwest and northeast with a basin in the center
150 filled with about 1.3 km of sediment. We refer to the northeastern ridge as

Schlanger Ridge, the southwestern ridge as Kingman Ridge, and the basin between them as Palmyra Basin.

Kingman Ridge links 3 seamounts with relatively deep saddles between them. Two of the three seamounts also reach the sea surface to form Kingman Reef and Palmyra Atoll. Schlanger Ridge, in contrast, is more massive and connected, but the crest of Schlanger Ridge is about 1000 m below the sea surface. To the south, Palmyra Basin abuts against another southeast trending basement ridge that we call Langseth Ridge. Langseth Ridge is the southeast terminus of Palmyra basin (Figure 2). The seamount that rises to form Kingman Reef has been dated to 70 Ma by ^{40}Ar - ^{39}Ar total fusion dating, (Davis et al, 2002), while a basalt from a dredge from Schlanger Ridge was dated at 76 Ma, also using the ^{40}Ar - ^{39}Ar total fusion method (Dredge 123D; Schlanger et al, 1984).

2.2 Sediments surrounding the LIR

The LIR is apparently a source of sediments to the surrounding sea floor (Figure 3; Divins, 2003; updated in Whittaker et al, 2013). The sediment deposits around the LIR are all much thicker than typical pelagic deposits on Cretaceous seafloor in the Pacific. The sediment apron surrounding the LIR is a major pelagic sediment accumulation within the tropical Pacific comparable to the thickness of sediments at the crest of the eastern equatorial Pacific sediment bulge (~600 m at its crest; Hays et al, 1969, van Andel et al, 1975; Mitchell and Lyle, 2005).

Typical sediments to the north of the Clipperton Fracture Zone (FZ) are thin, ~150-250 m (van Andel et al, 1975; Lyle and Wilson, 2006; Pälike et al, 2010) because the ocean floor was formed in the early Cenozoic or late Cretaceous when

the CCD was shallow (Arthur et al., 1985; Pälike et al, 2012). Consequently,

175 carbonates typically were dissolved before burial on the Cretaceous and Paleogene
flanks of the mid-ocean ridge, leaving only clays and radiolarian ooze. Sediments to
the south of the Clipperton FZ are thick in the position of the early Miocene equator
in the eastern Pacific (van Andel et al, 1975) but thin to the west. The region
surrounding the LIR south of Clipperton FZ (which now cuts across the LIR at about
180 2°N) were south of 4°S, i.e. in a low sedimentation regime, during the major high
sedimentation interval of the Oligocene and early Miocene. Consequently there is a
relatively thin sediment pile when compared to sediments to the east in the
equatorial Pacific sediment bulge. The eastern Pacific plate trajectory brought
sediments in the equatorial bulge through the equatorial high productivity region
185 when sediment production was highest, during the Oligocene and early Miocene.

Three DSDP sites have been drilled in the vicinity of the LIR that show how
sediments were deposited near the LIR. Site 165 is to the northwest (DSDP Leg 17
Shipboard scientific party, 1973, 490 m sediment), Site 315 is to the east (DSDP Leg
33 Shipboard scientific party, 1976, 996 m sediment), and to the south is Site 316
190 (DSDP Leg 33 Shipboard scientific party, 1976; 837 m sediment). Additional
sediment thickness data come from analog seismic reflection records acquired in
the 1970's (e.g., Normark and Spiess, 1976; Orwig, 1981) and from those by the
Glomar Challenger during the DSDP drilling expeditions themselves (Raff, 1973;
Schlanger and Winterer, 1976). Apparently, neither the thickness of sediment found
195 at Site 316 nor in nearby seismic profiles (Schlanger and Winterer, 1976) was
honored in the Divens/Whittaker grid shown in Figure 3. A ramp of thick sediments

is found from the base of the LIR to the position of Site 316, averaging a thickness around 1 sec of two-way travel time (TWTT) or around 800m.

The regional sediment thickness map in Figure 3 depicts the LIR crest as essentially sediment free, reflecting the lack of older seismic reflection and drilling data there. However, a Gloria side scan sonar survey in 1991, which also gathered seismic reflection data, found thick sediments within Palmyra Basin at the northwest end of the LIR (Paskevich et al, 2011; Barth et al, 2013, Figure 2).

Shallow water carbonates form only a small part of the excess sediment around the LIR. They were only found in the late Cretaceous basal volcanoclastic sediments of Sites 165 and 316, and from the middle Cretaceous at Site 315 (Jackson and Schlanger, 1976; Schlanger et al, 1984). The rest of the sediments are pelagic. Pelagic carbonate turbidites flowing from the LIR to the surrounding apron were critical to build the thick sediment column at Site 315 (DSDP Leg 33 Shipboard Scientific Party, 1976). Orwig (1981) has also traced turbidites sourced from the LIR that traveled to the northwest from Palmyra Basin toward Site 165, as well as to the southwest of the LIR. Because Site 165 is now well below the Neogene CCD (5040 m vs 4700 m CCD) there has been little sediment deposition since the early Miocene. Significant amounts of sediment have been generated on the LIR but then shed to the surrounding sea floor. Cretaceous turbidites sourced just after the LIR formed carried both volcanoclastics and the shallow water carbonates from the LIR.

Sediment sequences around the LIR are rarely complete but instead have significant time gaps. Hiatus periods are not regionally consistent however (Jackson and Schlanger, 1976), but apparently reflect the strength and temporal variability of

abyssal current flow around the LIR. Site 165 is missing Neogene sediments, for example, while Site 315 is missing part of the lower Miocene. The southern drillsite, Site 316, is missing much of the Oligocene. Neogene sediment sections are highly variable in thickness around the LIR, based on interpretations of seismic reflection profiles in the vicinity of the LIR (Raff, 1973, Schlanger and Winterer, 1976).

In this paper, we examine sediment deposition in the Palmyra Basin, a shallow north-facing basin at around 7°N on the LIR (Figure 2). Despite the large losses of sediment from LIR to the surrounding terrain there has been net sediment accumulation within Palmyra Basin, with average sedimentation rates since 75 Ma similar to Pleistocene sedimentation rates in the region. Assuming 75 Ma crust, the average sedimentation rate over the Palmyra Basin center is 1.7 cm/kyr compared to sedimentation rates for MIS 1-3 of 1.5 to 1.7 cm/kyr for cores nearby (cores 34, 35, and 37BB; Lynch-Stieglitz et al, 2015). Seismic horizons within Palmyra Basin are neither clean nor continuous. Instead, individual horizons can be traced for relatively short distances even though these horizons make up discernible sediment packets. The discontinuous seismic horizons are a result of downslope transport and a high-energy current regime.

3. THE MGL1208 LINE ISLANDS RIDGE SURVEY:

The MGL1208 coring, multibeam sonar, and seismic reflection survey of the LIR was the first major coring expedition to the Line Islands region in the last 30 years. Cores were collected along the LIR in order to study the Pleistocene paleoceanography of the central Pacific, including studying glacial-interglacial changes of the intertropical convergence zone (ITCZ), equatorial primary

productivity, and the El Niño-Southern Oscillation (Lynch-Stieglitz et al, 2015).

Multibeam bathymetry using the Kongsberg EM122 swath mapping system was

245 collected along the entire ship track. Multichannel high-resolution seismic reflection profiles discussed in this paper were collected near 6°N (MGL1208 Survey area E) and also south of Kiritimati (Figure 1). Digital 3.5 kHz chirp subbottom profiler data were collected over the complete survey area.

Most of MGL1208 was devoted to locating coring sites and recovering a new
250 set of sediment cores. More than 20 new piston cores, 'Big Bertha' gravity cores (6 m gravity cores using a piston core weight stand), and 2m gravity cores were collected along the LIR, with promising coring sites located by multibeam surveys and digital chirp data. The cores are described in Lynch-Stieglitz et al (2015). Pleistocene-Holocene sediments in the 2000-3500 m depth range cored on MGL1208 are all
255 pelagic foraminiferal sands and silts with no shallow water components (Lynch-Stieglitz et al, 2015). Typical sedimentation rates range from ~1cm/kyr in the north to 3cm/kyr near the equator.

The data in this paper are from the Palmyra Basin and ridges surrounding it, between 5 and 7°N near Palmyra Island (**Figures 1 and 2**). Palmyra Basin extends
260 ~100 km in a SE-NW direction, from about 5°40'N to about 6°45'N, Seismic line 1 (**Figure 2, Figure 4**) was shot near the southeastern crest of Palmyra basin, where the sea floor depth ranges between 2000 and 2400 m below sea level. Seismic line 2 (Figure 2, Figure 5) crosses the foot of the northern seamount on Langseth Ridge, roughly from west to east.

265 **4. METHODS**

Seismic reflection data for the Line Islands were collected on the R/V Langseth using the onboard Syntrak 960-24 MCS digital data acquisition system with a Digishot source controller (<http://www.ldeo.columbia.edu/research/office-of-marine-operations/langseth/scientific-instrumentation>). The seismic source for
270 the high-resolution multichannel seismic profiles was two 45 cubic inch (c.i.) generator-105 c.i. injector GI air guns. The seismic reflection data were collected with a 900 m, 60-channel streamer having 12.5 m group spacing. Shots were spaced at 25 m and data were recorded for 8 seconds and digitized at 0.5 ms intervals.

The 60-channel data were stacked after NMO correction and noise reduced
275 via spectral shaping, with a 30 Hz lower cutoff, 100% pass from 60Hz to 450 Hz and then ramping down to a high cutoff at 900 Hz. The data were then migrated assuming 1500 m/s acoustic velocity. Three seismic reflection lines (19 hours of survey) were collected in the vicinity of Palmyra Basin, and another 33 hours were collected where LIR crossed the equator. Seismic data are archived at the University
280 of Texas Academic Seismic portal of the Marine Geoscience Data System (<http://www.ig.utexas.edu/sdc/cruise.php?cruiseIn=mgl1208>).

Estimates of depth in the seismic profiles use a seismic velocity of 1500 to 1600 m/s in unconsolidated sediments (Hamilton et al, 1982). Total thickness of sediments in Palmyra basin were estimated by comparison to TWTT to basement at
285 Site 315 versus depth drilled, plus shipboard measurements of seismic velocity in the lower sediments to extrapolate below the bottom depth of Site 315 km (Site 315, DSDP Leg 33 scientific party, 1976).

Swathmap bathymetry was collected using the Kongsberg EM122 multibeam system on the R/V Langseth. The data were ping-edited to remove outliers using MB-System software (available from <http://www.ldeo.columbia.edu/res/pi/MB-System/>) and plotted with the open-source GMT mapping software (<http://gmt.soest.hawaii.edu/>). Additional swathmap bathymetry data come from the US interagency Extended Continental Shelf Program (ECS; Barth et al, 2013; Gardner and Calder, 2010). The ECS Program data are found along the SW side of Palmyra Basin along Kingman Ridge and to the west of the LIR.

5. SURFACE FEATURES: EROSION AND CURRENT ACTIVITY

The most prominent sedimentary features around the crest of the LIR indicate an energetic sedimentary environment with significant sediment movement. Submarine channels are prominent along the edges of LIR, e.g., as found around the rim of Palmyra Basin (Figure 2; Barth et al, 2013), while sediment waves are prominent at the saddle between Kingman and Langseth Ridges (Figures 4,6, 7).

In most of Figure 2, the apparently smooth topography is an artifact of satellite ocean bathymetry (Smith and Sandwell, 1997). Satellite topography is adequate to find large-scale bathymetric features, but cannot resolve bathymetric features < 10km in spacing, even as large as abyssal hill topography (Goff et al, 2004). Multibeam mapping is needed to reach bathymetric resolution sufficient to study the sedimentation and its relationship to the LIR seamounts and ridges, as around the rim of Palmyra basin (Figure 2).

The sediments on the LIR are being channeled to the surrounding fan systems via a network of anastomosing small channels that join main channels

within each sub-basin. They drain to abyssal plains that form the apron surrounding the LIR. A well-imaged large channel system can be seen on Figure 2 at 5.8°N, 161.5°W, just south of Palmyra Atoll. A large channel system has been poorly imaged by bathymetry but runs almost due south to north from the upper to the lower Palmyra Basin and carries sediments to the abyssal plain around 6°40'N, 162°30'W. The channel system has been better imaged by the sidescan GLORIA survey of the US EEZ around Kingman Reef and Palmyra Island in 1991 (Paskevich et al, 2011, Barth et al, 2013, Jones et al, 2013).

Channels and sub-channels have little sinuosity because they are (1) formed on steep topography, (2) LIR sediments are coarse-grained, and (3) the sediments have little cohesion (Clark et al, 1992). Sediments collected by coring on MGL1208 are typically foraminiferal sands, with about 50% sand fraction (>64μ ; Lynch-Stieglitz et al, 2015). The remainder is nannofossil carbonate.

The 3 cores attempted in the upper Palmyra Basin in 1800-2100 m of water collected only traces of foraminiferal sand in the core catchers either because the cores did not penetrate deeply enough to retain the sediments or the sediments drained out. The carbonates at the top of the basin are apparently winnowed, much richer in coarse fraction, and are thus too sandy and poorly cohesive to be cored with the methods we used. Sediments with similar composition on the sides of the basaltic seamounts and ridges appear to slough into channels easily, often leaving avalanche scars behind.

The upper Palmyra Basin, where the seismic reflection profiles were acquired (Figure 2, **Figure 4**, MGL1208 Seismic line 1; Figure 5: Seismic Line 2) is

divided into a NE channeled regime along the base of Schlanger Ridge and a SW
335 regime with smoother sediments. The NE channels drain from higher topography
and are perched with respect to the SW part of the seismic line, where smoother
sediments and the basin depocenter are located. A seamount to the south of the
main axis of Schlanger Ridge (Figure 2) helps to enclose the NE subbasin and
steepen the basement topography. In contrast, the slope above the SW end of
340 Seismic Line 1 is relatively shallow.

Smooth sediments as well as sediment waves are also found in the gap
between Kingman and Langseth Ridges (**Figures 6 and 7**). The smoother sediment
fill extends about halfway across the upper Palmyra Basin, from the southern edge
to the basin depocenter. Further downslope to the NW, the upper reaches of large
345 channels that drain northward disrupt smooth sediment deposition . These
channels are large enough that they can be traced even in the low-resolution
satellite bathymetry in Figure 2. To the NE, as Seismic Line 1 passes across the crest
of a saddle across Schlanger Ridge, smooth sediments are also deposited. A lee drift
of sediments is found to the NE of a small basalt cone in the gap (Figure 4),
350 indicating recent current flow primarily from south to north.

Sediment waves are found at the gap between Kingman and Langseth Ridges
(Figure 2, **Figure 6**). They form north of a step upward in sea floor level possibly
buttressed by basement topography (**Figure 7**). Bottom currents through Kingman-
Langseth Gap cause a series of sand waves that have ~1.3 km wavelength and wave
355 heights of 10-20 m (**Figure 6**).

The sediment waves have developed over long periods of time, since similar waves can be found to depths of ~ 0.2 seconds two way travel time (TWTT), ~ 150 -
160 m deep in the sediment column as inferred by the undulating sediment horizons (**Figure 7**; the right hand side of Seismic Line 1). The wavelength in Figure
360 7 is ~ 2 km rather than 1.3 km because the seismic line cuts the wave field obliquely. Given typical Pleistocene LIR sedimentation rates of 1-2 cm/kyr north of 5°N (Lynch-Stieglitz et al, 2015), sediment waves have been forming since ~ 8 -16 Ma

Large abyssal sediment waves are thought to occur primarily in fine-grained muds at current speeds of 0.05 to 0.25 m/s (Flood, 1988; Stow et al, 2009). The
365 foraminiferal sand waves we document here are apparently too coarse for these types of sediment waves—sand waves typically require higher near-bottom current flows of ≥ 0.5 m/s, and significant sediment supply (Stow et al, 2009). However, hollow whole foraminiferal tests are significantly less dense than a solid calcite or quartz sphere of the same size. Because their effective density is ~ 1.5 g/cm³ vs 2.7
370 g/cm³ for a solid calcite sphere, they can be eroded very easily (Miller and Komar, 1977). Current speeds in the range of 0.15-0.2 m/s are sufficient to erode 100 to 300 μm size tests from foraminiferal sands. Moderate currents will also preferentially transport whole tests and leave behind test fragments because the fragments are smaller and denser (Miller and Komar, 1977). Moderate speed
375 bottom currents sweeping through the gap between Langseth and Kingman Ridges into Palmyra Basin could preferentially sort whole tests and deposit them in sediment waves found in Figure 6. We did not core the seamount gap to be able to study the differences in sediment composition there, however.

Abyssal current velocities over the LIR can be gleaned from the HYCOM data
380 assimilation ocean circulation model reanalysis (Hybrid Coordinate Ocean Model;
<http://hycom.org>) which has a $1/12^\circ$ (9 km) horizontal scale and a variable vertical
scale. Below 2 km, model ocean layers are 500 m thick. For the 6-year period from
18 Sept 2008 to 11 Sept 2014, daily average u and v velocity data from every 2nd
day were extracted from the HYCOM reanalysis and then averaged to make the
385 current velocities shown in Figure 6. HYCOM model current speeds are high in the
vicinity of the Palmyra Basin sediment waves, averaging between 0.2-0.25 m/s,
sufficiently vigorous to form the observed waves (Figure 6).

The HYCOM model may also bias bottom current velocities low. Comparison
of HYCOM model eddy kinetic energy to measurements at a moored array in the Gulf
390 of Mexico suggested that HYCOM may underestimate abyssal current velocities by
~40% (Rosburg et al, 2012). Also, the HYCOM model does not resolve tidal currents,
so actual near-bottom velocities may be even larger when the tidal cycle reinforces
the mean current. Finally, current speeds should be locally higher at flow
constrictions like that formed by the gap between Kingman and Schlanger Ridges
395 than the average in a typical $9 \text{ km} \times 9 \text{ km}$ HYCOM box.

The sediments to the south of the gap are smooth and the topography is
significantly deeper than upstream, forming a smooth platform at ~2340 m water
depth, with a step up to the 2250 m water depth of the Palmyra basin fill. The gap
has been adequately surveyed by multibeam to confirm that the platform does not
400 have sediment channels typically found elsewhere near the LIR crest. Such

smoothing suggests transport by the currents flowing through the gap to fill in channels and smooth the sediment surface.

Sediments only about 250 m thick are imaged at the SW end of Seismic Line 1 (**Figure 7**) versus ~1.3 km of sediment at the depocenter (see Section 6.3 for thickness estimate). However, Seismic Line 1 ends near the base of a seamount rather than in the gap between Kingman Ridge and Langseth Ridge. Within the gap there is about 400 m of sediment (0.5 s TWTT), as found at the western end of Seismic Line 2 (**Figure 5**). The thick sediments and smooth topography within the Kingman gap indicates net deposition and the current indicators suggest net transport from the south.

The currents that produce the sand waves have varied through time. Occasional hiatuses can be found at the end of Line 1 (**Figure 7**) where a surface was formed and there was apparently little sediment deposition for some time. A prominent seismic horizon within the sand waves is marked on Figure 7 roughly at 150 ms TWTT (~120 mbsf) and marks the upper surface of a series of sediment waves. The lower surface of this sediment packet has cut into previously deposited sediments and is found at a depth about 75 m below the hiatus surface. These sediments mark a relict surface from an interval of higher current speeds through the gap.

6. SEISMIC HORIZONS AND PALMYRA BASIN CHANNELS

Although basaltic basement forms the high topography both to the north and south of Palmyra basin, the high topography between the two basement ridges results from sediment fill (**Figures 4, 5, 8**). Despite surface erosion, there has been

net sediment deposition since the late Cretaceous, with more than 1 km of sediment
425 deposited in the axis of the central basin. Even with strong currents, the gaps in both
Kingman and Schlanger Ridges are sediment covered. Within the saddle on
Schlanger Ridge at the NE end of Line 1 (6° 20' N; Figure 4) flat topography formed
by sediments has gradually expanded southward into the basin, perhaps by
relatively high supply of sediments from the rest of Schlanger Ridge and by tidal
430 currents moving sediment to the ridge flanks.

Figure 4 is a comparison between the surface bathymetric features from the
multibeam survey with the seismic reflection profile along Seismic Line 1. Major
channel systems are found on both sides of Schlanger Ridge; the northern channel is
outside of Palmyra Basin and drains to the east, while the channel within Palmyra
435 Basin flows to the southwest and is perched relative to the rest of Palmyra Basin.
Both channels are cutting down into older sediments. Significant sediment erosion
(~500m) has occurred on either side of Schlanger Ridge within the channels, but
200-600 m of older sediments still remain (Figure 4, Figure 8). The interchannel
highs do not have interpretable seismic stratigraphy. The poor seismic continuity
440 within the channels may result from repeated episodes of filling followed by channel
cutting.

A second system of small wide channels occupies the upper Palmyra basin to
the south of the sedimentary ridge that divides the central channel system from the
more deeply cut channels in the NE (**Figure 4**). There is net sediment deposition
445 near the basin axis—older deeper channels are buried under younger sediment fill.
Unlike the channels near Schlanger Ridge that originate on basalt seamounts and

relatively steep topography, there is a gentle slope upwards to the crest of Palmyra Basin 20 km to the east and 100 m shallower in depth than the SW end of Seismic line 1 (Figure 2). The lower net slopes leads to less erosion.

450 However, erosion of surface sediments has occurred in the middle of Seismic Line 1, as shown by truncation of sediment layers, near the lowest point of Line 1. Further southwest, the sediment waves discussed in the last section are slowly burying high basement topography near Kingman Ridge.

 This broad channel near the center of Seismic Line 1 flows westward within
455 the Palmyra Basin to eventually join with a large channel flowing northward away from Kingman Ridge (Figure 2) and the combined system then flows north and downslope out of the Palmyra Basin. The channels carry sediment from the ridges down the length of Palmyra Basin and as far north as DSDP Site 165 (Orwig, 1981), at least through the early Miocene. Further survey and analysis is needed to make a
460 quantitative estimate of sediment loss from Palmyra Basin to the north, however.

6.1 Continuity of LIR Seismic Horizons: comparison to Cocos Ridge

 There is poor continuity along seismic horizons from the LIR (**Figure 9, top**). Sediment layering on the LIR contrasts with the classic pelagic drape found on the north flank of the Cocos ridge in the eastern equatorial Pacific (**Figure 9, bottom**;
465 Line 2 from the MV1014 Cocos Ridge survey, Liao and Lyle, 2014). Both seismic lines are played back to the same scale, and both were shot with paired 45-105 cubic inch GI guns. Seismic horizons found in the Cocos Ridge profile are easy to trace along the entire line despite significant changes in topography, sediment

thickness, and occasional erosion. Erosional areas on the Cocos Ridge are easy to
470 identify by breaks in seismic horizon continuity.

In contrast, the seismic horizons in the Palmyra Basin are not nearly as
continuous. Seismic horizons are prominent for a few km, and then often fade away.
Reflection amplitudes often abruptly change strength, as at the boundaries of
channels. The changes in amplitude make the seismic record appear blotchy. We
475 hypothesize that the poor continuity is due to the dynamic nature of sedimentation
in the Palmyra Basin and that changes in seismic amplitude reflect along-strike
changes in sediment physical properties (grain size, porosity, bulk density) caused
by current sorting and the resulting variability in local deposition environments.
Sediments may not only be sorted before being deposited by the high currents, but
480 they can also be reworked as currents or density flows move them from their place
of initial deposition to where they are finally buried.

The difference in seismic response of sediments on Cocos Ridge with respect
to those in the Palmyra Basin results from multiple factors: (1) higher biogenic
silica content of the biogenic carbonates on Cocos Ridge, (2) lower current speeds,
485 and (3) lower gradient from the sediment sources to the depocenter. Cocos Ridge
sediments are biogenic carbonates but also have 10-60% diatom bio-silica (ODP Leg
202 Scientific Party, 2003). In contrast, LIR sediments are essentially 100%
carbonate. Cyclic deposition of carbonate-rich versus carbonate-poor sediments
produces and the strong density contrast between these two sediment types
490 produces good acoustic impedance changes between layers and strong, easily
traceable seismic horizons typical of the Cocos Ridge and the eastern Pacific in

general (Liao and Lyle, 2014, Mayer et al., 1986). The diatoms also form a sediment framework to help bind together the carbonates and keep them from eroding as easily as pure carbonates.

Both mean and maximum current speeds are lower at Cocos Ridge as compared to the LIR and concentrated around the ridge tops (Figure 10, as compared to Figure 6; note the different color schemes). Maximum near-bottom current speeds at the LIR are 0.2-0.25 m/s, with spots that exceed 0.4 m/s. Current speeds are in the range of 0.15 to 0.20 m/s along the Cocos Ridge crest. These are sufficient to winnow sediment and preferentially erode fines from the top of the ridge (Marcantonio et al, 2014; Liao and Lyle, 2014) but weaker maximum velocities (~0.1 m/s) downslope allow relatively even pelagic sedimentation of the resuspended sediment near the Cocos Ridge depocenter (near Site 1241).

Finally, the gradient from the top of Cocos Ridge to the depocenter is relatively shallow so that channeling is not as favored. The Cocos Ridge drops only 300 m over 50 km from the top of the ridge to Site 1241, so that there is less gradient to allow development of channeling. The Cocos Ridge, at least near Site 1241, appears to be a system where the downhill sediment flux is dominated by resuspension and downhill transport, while the LIR sediments are redeposited via channeled flows.

6.2 Modern sediment deposition and channeling from seamounts at the LIR

Paradoxically, where LIR sediments are not disturbed or moved, sediment packets can be correlated over long distances. Pleistocene cores were recovered on

MGL1208 spanning 800 km along the LIR that all have a common bulk density
515 stratigraphy also replicated by oxygen isotope stratigraphy (Lynch-Stieglitz et al,
2015). However, in order to collect undisturbed Pleistocene sediment sections, the
MGL1208 scientific party had to search for small areas of flat topography where
sediments had not sloughed off LIR slopes. These areas were typically found on
interfluvies between active channel systems and at water depths ~3,000 m. Careful
520 surveying was essential for finding suitable areas to core.

The high energy at the sea floor along the LIR and easy erodibility of the
sediment can also result in significant local hiatuses and sediment redeposition at
any particular location. Within the MGL1208 core set, some cores had
stratigraphically complete sections for up to 450,000 kyr, but others recovered only
525 a few meters of undisturbed sediments before a hiatus was reached.

Seismic Line 2 (**Figure 5**; see Figure 4 for line location) follows along the
base of the Langseth Ridge and gives insights into how sediment forms channels as
they drain sediments from basement high topography. The beginning (western end)
of Line 2 is at the gap between Kingman and Langseth Ridges where there is smooth
530 topography and thick sediments. There is also a strong reflection shallow in the
section that may be the same hiatus noted in Figure 7. The seismic horizons have
little continuity, like Seismic line 1, also showing the dynamic depositional
environment.

As Seismic Line 2 crosses over the toe of the seamount to the south of the gap
535 (Figures 2 and 4), deposition of sediment in between minor basement ridges is
imaged. Oldest sediments were preferentially deposited in basement valleys

between spurs from the seamount with some sediment drape. Channels originating from Langseth Ridge now cut into the youngest sediment column (Figure 5) where much of the sediment surface may be relict. Surprisingly, the channels are only partly contained by the basalt ridges. Many basalt spurs were first buried by sediments and then have been exhumed later. The largest channels have developed on the eastern end of Seismic Line 2, as the slope starts to fall away to the southeast, into the next basin south of Palmyra Basin along the LIR. It is clear from the multibeam bathymetry that the channels originate on Langseth Ridge, and likely result from focused downflow of foraminiferal sands. However, the ubiquity of channels points to a high production and deposition of sediments on the LIR basalt ridges, that feed significant volumes of sediment into the basins via the channel systems.

6.3 Sediment Deposition in Palmyra Basin

Figure 8 is an interpretation of Seismic Line 1. Basalt basement in the center of Palmyra Basin is now at a depth of about 3500 mbsl, or ~2 km below the flanking ridge heights, and ~2 km above the average sea floor surrounding the LIR. Even if the ridges were near sea level when they first formed, Palmyra Basin was still relatively deep although uplifted from typical sea floor..

There is no ground truth for a seismic stratigraphy of the basin, but it is likely that the lower (brown) unit is primarily Cretaceous and early Paleogene in age, given that the LIR formed around 70-85 Ma. At Site 315, the Paleocene and Eocene are marked by low sedimentation rates (DSDP Leg 33 Scientific Party, 1976), so we

assume that there is only a thin Paleogene section. We have picked basement as the
560 hard lower reflections. However, this may also include some cherty sections near
basement difficult to distinguish from basalt. Cherts have been found in the
Paleocene-Oligocene sections of Sites 315 and 316.

Also based on Site 315, we propose that most of the deposition in the basin
occurred in the late Oligocene-early Miocene, when Palmyra Basin was carried
565 across the equatorial zone by plate tectonics (~18 Ma equator crossing using a fixed
hotspot model, Engebretson et al., 1985 with Gripp and Gordon, 1990 for 0-5 Ma). A
similar age of equator crossing (18 Ma) is achieved using the hybrid Torsvik et al
(2008) paleomagnetic absolute reference frame plate model.

Assuming that the brown unit in Figure 8 is probably late Cretaceous-
570 Paleogene, we propose that the blue unit is Oligocene-middle Miocene, and the light
green sediment unit is possibly middle-late Miocene. Based on typical
sedimentation rates, the youngest pink unit is probably < 5 Ma and represents the
Plio-Pleistocene.

Thick sediments are found in the center of Palmyra Basin, with a total TWTT
575 of 1.4 s. If acoustic velocities within Palmyra Basin sediments are similar to those at
Site 315, there is 1.3-1.4 km of sediment at the deepest point. From the Site 315
seismic-drilling comparison, 1.1 s TWTT is equivalent to 1 km of sediment over
basement (Site 315, DSDP Leg 33 scientific party, 1976), and using an estimated
sediment velocity of 2.3 km/s in the lower Site 315 sediments, the addition 0.3
580 seconds should be equivalent to about 350 additional meters of sediment.

While there is no seismic cross-line from MGL1208 down the axis of Palmyra Basin, it is clear from the bathymetry and confirmed by the GLORIA seismic reflection data reported by Barth et al (2013) that the entire Palmyra Basin is a thick sediment pile. The northern LIR topography thus combines basaltic volcanism that created the topographic framework surrounding Palmyra Basin, and sedimentation that has since filled it. The LIR edifice ends at about 7.5°N where the volcanic topography that makes up Kingman and Schlanger Ridges ends and where there is a gap to the next major seamounts to the north. Excess sediment flows out in channels extending to the north of the LIR.

Initial sediment deposition in the Cretaceous was biased towards the NE (Schlanger Ridge) side of Palmyra Basin (Figure 8). Significant deposition occurred in the NE end of Line 1 during the brown time interval, with less deposition to the southwest. The upper boundary of the brown unit appears to be truncated at both ends of the seismic line, in the NE and SW. Because Seismic Line 1 crosses a subbasin near Schlanger Ridge with seamounts surrounding it on 3 sides, sediment deposition in the NE was enhanced by downslope sediment transport from the Schlanger Ridge.

As the basin matured, the axis of sedimentation migrated to the southwest and sediments began building up at the southwest end of the profile. Little sediment was deposited at the SW end of Seismic Line 1 during the brown interval. In contrast, the blue sediment unit (Oligocene-lower Miocene?) is thickest in the center of the basin. The upper boundary appears conformable with the overlying

sediments. Both the brown and blue units thin significantly as Kingman Ridge is approached, in the SW part of Figure 8.

605 We could not trace the light green unit into the channeled subbasin in the NE, partly because the seismic line crosses channels from the base of a seamount offset to the south of the main Schlanger Ridge that helps to form the subbasin (see map in Figure 2).

 The light green unit thickness is more uniform over the rest of the basin, but
610 the thickest sediments have shifted even further SW than those of the blue unit. The pink Plio-Pleistocene unit is continuously deposited to the SW but is severely cut by channels in the NE half of Seismic Line 1. We could not confidently trace it into the NE subbasin.

 There is a deposit of material with high acoustic reflectivity found on both
615 flanks of Schlanger Ridge that may be the remnants of an early channel system or perhaps volcanoclastics and shallow carbonates associated with LIR formation. Volcanoclastics are found in the basal sediments of Sites 165, 315, and 316 around the LIR (DSDP Leg 17 Scientists, 1973; DSDP Leg 33 Scientists, 1976) and shallow water carbonates are found in the basal sediments of Site 315.

620 Buried channels are marked by highly reflective seismic packets that are the response from sides of channels or meanders within the channel system (Figure 8). They are prominent in the NE cutting into the boundary between the brown and blue horizons. To the northeast of the Palmyra Basin depocenter we find evidence of a persistent channel formed low in the sediment sequence. We have outlined a
625 channel, shown by dotted lines in the brown sequence, with additional channel

levees being found in the lower blue sequence. A similar channel cuts into the upper sediment column, suggesting that the position of this large channel is partly controlled by basement topography. Despite these channel systems, there has been a net filling of Palmyra Basin.

630 There may be net loss of the shallowest sediments within Palmyra Basin in the modern system, indicated by downcutting channels. Near Schlanger Ridge there is major downcutting into the older sediment sequences. In the central part of the seismic profile, however, there is both downcutting and filling suggesting that loss of sediment from the upper Palmyra Basin has been episodic and is largely balanced
635 by fresh sediment production upslope.

7. DISCUSSION AND CONCLUSIONS

Large-scale regions of elevated topography like the LIR are important for carbonate deposition within the Pacific Ocean. During the late Mesozoic and early Cenozoic, the Pacific CCD was often shallower than the average depth of the mid-
640 ocean ridge system, and carbonate was only deposited on anomalously shallow topography like the LIR or large igneous provinces like the Ontong Java Plateau, Manihiki Plateau, or Shatsky Rise (Neal et al, 1997). The lack of a large deep sediment carbonate reservoir in the equatorial Pacific probably helped to amplify swings in the CCD in the Paleogene (Pälike et al, 2012).

645 Scientific drilling and seismic reflection profiles crossing the LIR give evidence that the high topography has been responsible for collecting a sediment pile that is much thicker than the average in the surrounding tropical Pacific, especially for the Paleogene. The LIR and the sediment aprons around it extend

carbonate deposition about 5° to the west of where the intersection of the CCD and
650 average mid-ocean ridge depth should have ended carbonate burial. The LIR and its
flank deposits make up roughly 5% of the total depositional area of equatorial
Pacific carbonate zone. Since the sediment pile is mostly carbonates, the
preservation of this sediment is important to the Cenozoic carbon cycle. The
presence of large areas of mid-depth carbonates (~2000-3000 m) on topographic
655 highs like the LIR helps to buffer changes in carbonate chemistry of Pacific waters
during large atmospheric CO₂ transients like the fossil fuel transient today, or
perhaps the transient associated with emplacement of the Columbia River Basalts at
the early-middle Miocene boundary.

7.1 LIR sedimentation versus the Maldives Ridge

660 LIR carbonate sedimentation is different from sedimentation at other pelagic
tropical ridges like the Maldives or Hawaii, because the relatively small volume of
volcanism built the LIR framework in contrast to voluminous volcanism at the
Maldives Ridge or massive individual volcanic centers as in the Hawaiian Island
chain. .

665 The LIR has a distinct pattern of volcanism that is later buried beneath
sediments. There were large numbers of small seamounts and basalt ridges that
formed the LIR, perhaps by crossing small hotspot trails (Pockalny et al, 2015) or by
mantle upwarping (Davis et al, 2002). This volcanism formed a relatively wide
region of elevated ocean floor with sub parallel seamount ridges, culminating in the
670 Palmyra Basin in the north.

What is pertinent to the sedimentation is that no large near-sea level basalt platform was formed. Instead, a series of small islands formed that have since sunk to become atolls. Because the ocean crust is about 50 million years older than the seamount volcanism, subsidence from crustal cooling was relatively minor after volcanism stopped. Shallow carbonate production was limited by seamount area that reached into the near-surface ocean. Around the Palmyra Basin, only Kingman Reef and Palmyra Island maintain reefs along the southwestern Kingman Ridge. The seismic evidence suggests that the northeastern Schlanger Ridge may have been only briefly emergent. In between the two bounding ridges, the Palmyra Basin was ~2 km below the Cretaceous ocean surface. Only 3 other atolls besides Kingman Reef and Palmyra Island (Teraina, Tabuaeran, and Kiritimati) form islands along the LIR. Each of these occupies the summit of a volcanic edifice, separated by 100 km or more from each other. The lack of large numbers of atolls is evidence that most of the LIR was never above sea level.

Further south, the matrix of seamounts that form the LIR has formed many small basins along the ridge. While a few seamounts have joined to form linear chains, many are separated by 50 km or so, and form subbasins along the ridge. Several subbasins that were crossed during the MGL1208 survey also show development of major channel systems draining sediment from the shallower regions.

In contrast, The Maldives Ridge—a hotspot ridge in the tropical Indian Ocean—exemplifies sedimentation along a ridge with voluminous volcanism. Since the Maldives Ridge was at or near the Paleocene ridge crest when formed at the

Reunion hot spot (57 Ma, Duncan and Hargraves, 1990) it was also subject to much

695 larger subsidence as well (Coffin, 1993) to provide accommodation for a thick
shallow carbonate sequence. The Maldives Ridge consists of a massive volcanic
edifice that reached sea level and then slowly subsided (Aubert, 1994; Aubert and
Droxler, 1996, Betzler et al, 2015). The central region was further downfaulted and
subsided more rapidly than the two flanks of the ridge (Aubert, 1994), The high
700 volume of volcanism produced a very large edifice on which sediment then
accumulated. The shallow but downfaulted central part of the ridge provided a
shallow sediment catchment as well.

The Maldives platform is as wide as the entire Palmyra basin on the northern
LIR (Figure 2) but mostly is less than 500 m deep. Apparently much of the edifice
705 reached sea level and subsequently subsided more slowly than reefs grew, allowing
reef and shallow carbonate sediments to deposit nearly as rapidly as the ridge
subsided. The more rapid subsidence of the center forms a deeper Maldives Inner
Sea flanked by a series of atolls along each ridge flank (Aubert, 1994). The basalt
crust of the Maldives Ridge is now topped by 2-3 km of shallow-water carbonates
710 (Aubert and Droxler, 1996). The differing volume of volcanism that formed each
ridge along with differing subsidence history caused the differences in
sedimentation between the Maldives and the LIR. While the Maldives formed a large
shallow platform that allowed both reef formation and a catchment for shallow
carbonates, volcanism at the LIR formed small isolated islands. Nevertheless the LIR
715 uplift formed mid-depth topography above the Pacific CCD where carbonates could
accumulate and then be channeled to its apron. Thus the Maldives became a large

shallow-water carbonate platform, while the LIR became primarily a reservoir for pelagic carbonates.

7.2 Sediment deposition in the Palmyra Basin

720 In the absence of drilling in Palmyra Basin, we can only infer the ages of different sediment units based upon drilling on the LIR apron. However, it is clear from Site 315 that highest deposition rates regionally occurred in the late Oligocene-early Miocene, associated not only with the time when Palmyra Basin was crossing the equator but also with the high sedimentation rates and deep CCDs found
725 throughout the Pacific at that time (Lyle, 2003; Moore et al, 2004; Pälike et al, 2012).

 The accumulation of sediment within Palmyra Basin is somewhat faster than sediment burial in the LIR apron, but not hugely so. We estimate that there is 1.3 km of sediment within Palmyra basin, as compared to 800-1000 m drilled on either side of the LIR at Sites 315 and 316 (DSDP Leg 33 Scientific Party, 1973). While the
730 oldest sediments from the DSDP sites contain shallow water carbonates, most of the sediment pile appears to be pelagic in origin. The higher sediment accumulation in Palmyra Basin probably results from the interaction of currents with the ridges surrounding it to preferentially deposit sediment from the tops of the basalt ridge into Palmyra basin.

735 Since the area of the ridges surrounding Palmyra Basin make up only ~20-25% of the total basin area, shedding of sediment from the ridges alone could not provide all of the excess sediment regionally. Instead, early development of shallow topography by volcanism along the LIR may have initially provided a relatively shallow sea floor, upon which carbonates could be deposited with minimal

dissolution. The large region of turbidite deposition on the flanks of the LIR indicates that a significant fraction of sediment first was deposited on the LIR and then flowed outward from the high topography.

One of the remaining mysteries of LIR sedimentation is how the LIR appears to have generated more pelagic carbonate production regionally than one would normally expect in the pelagic realm. From studies of Pleistocene cores taken on the MGL1208 cruise, the average sediment deposition in Palmyra Basin is close to the level that could be supplied by surface water production. The Pleistocene sedimentation rates (1-2 cm/ky; Lynch-Stieglitz et al., 2015) are similar to the long-term average needed to fill the basin. The amount of shallow bare ridge tops should be sufficient to supply some sediment to the surrounding apron, but it is unclear how the LIR could support voluminous turbidite deposition in the region around it. If there were an 'island effect' of elevated primary productivity along the LIR near islands (e.g., Andrade et al., 2013) it was probably not large. The few islands along the LIR are small and probably have small influence on the regional productivity. Developing a sedimentary balance depends upon better estimates of sediment loss to the LIR apron.

Sediments from Kingman Ridge on the SW side of Palmyra Basin probably are an important sediment source for Palmyra Basin even though there is also significant loss from Kingman Ridge outward to the LIR apron. Because of the prevailing abyssal currents traveling from the SW to the NE, the southern Kingman Ridge should be the most important source of horizontally transported sediment to Palmyra Basin. Sediment waves indicate that the SW to NE current directions

prevailed over the late Neogene at least. Current activity has shaped the Palmyra basin sediment fill since the basin was first formed. Early in the basin's history, there were high acoustic impedance (high current velocity?) deposits along Schlanger Ridge flanks. Nevertheless, sediment deposition appears to have been biased toward Schlanger Ridge during the Cretaceous and Paleogene (the brown unit in Figure 8), perhaps being sourced by the more massive ridge on the NE of Palmyra basin.

Flow of sediments downhill and relatively high current velocity has caused pelagic sediment drape to be poorly developed over Palmyra Basin, as indicated by the mediocre continuity of seismic horizons. Instead of relatively quiet pelagic sediment rain to the sea floor, there is abundant evidence of current sorting that deposited different size classes of sediment in different locations, and there has probably also always been at least some level of downslope transport.

The relatively large amount of channeling in the younger sediments, especially near Schlanger Ridge, may reflect a slowdown in sedimentation since the Miocene, perhaps because of plate tectonic movement away from the equatorial high productivity zone. At Site 315 the highest sedimentation rates broadly matched when Site 315 (now at 4°10' N, Figure 4) was in the equatorial zone. However, sedimentation rates at Site 315 radically increased after 27 Ma, when Site 315 was positioned at 4°S rather than near the equator. Clearly other factors were in play to increase the sedimentation since the middle Oligocene. It appears that a major factor in the change in sedimentation was the increase of equatorial Pacific

785 carbonate deposition in the middle-late Oligocene and broadly deeper CCD (van
Andel et al, 1975; Pälke et al, 2012).

7.3 The LIR and regional carbonate sedimentation

. DSDP drilling around the LIR has shown that there are thick carbonate
sediment deposits around the LIR, with turbidites apparently sourced from the LIR.

790 However, the general elevation of the crust surrounding the LIR must also have
contributed to the regional preservation of carbonates. Turbidites are common in
the drilled sediments, but a significant but unquantified amount of the sediment
column is pelagic in origin. Spot coring (a core every 100 m or so in the upper
sediments) during drilling of these early DSDP sites makes it difficult to assess the

795 Neogene contribution of the LIR to the surrounding apron. However, since
carbonate deposition has continued since the Cretaceous, initial uplift during the
late Cretaceous, 40 to 50 million years after the local ocean crust originally formed,
and initial volcanoclastic sedimentation was sufficient to build a sediment column
shallow enough to maintain carbonate burial (DSDP Leg 17 Party, 1973; DSDP Leg
800 33 Scientific Party, 1976). Now Sites 315 and 316 are still above the CCD despite
being on roughly 120 Ma ocean crust because they both have accumulated nearly 1
km of sediment. In contrast, the deeper and more distal Site 165 stopped
accumulating turbidites in the early Miocene, as it dropped below the CCD.

Today, the ubiquitous presence of channels from the LIR to the apron gives
805 evidence that the LIR itself is still contributing large volumes of sediment to the
surrounding region. In Palmyra Basin we find that these channels are both large
and active. Moderate levels of anomalous volcanism long after the ocean crust was

formed combined with relatively unstable sediments accumulating along seamounts at the the LIR have provided both large-scale loss of surface sediments (Lynch-
810 Stieglitz et al, 2015) but also have built significant accumulations of carbonate sediments at and around the LIR.

8. ACKNOWLEDGEMENTS

NSF grants OCE-1157486 (M.L.), OCE-1158994 (R.P.), OCE-1159053 (J.L.-S.), and OCE-1158886 (P.J.P.) funded the MGL1208 cruise responsible for collecting the
815 cores, multibeam and seismic reflection data. The captain and crew of the R/V Marcus Langseth and the LDEO Marine Office are gratefully acknowledged for the success of MGL1208. Funding for the analysis and publication of this paper came from USGS Grant G13AC00351 to Pockalny in support of the US Extended Continental Shelf project.

820 Abyssal current data were extracted from the HYCOM reanalysis. Funding to develop HYCOM is from the National Ocean Partnership Program and the Office of Naval Research. Data assimilative products using HYCOM are funded by the U.S. Navy. Computer time was made available by the DoD High Performance Computing Modernization Program. The output is publicly available at <http://hycom.org>.

9. FIGURE CAPTIONS

Figure 1: Location of Line Islands Ridge with respect to the Hawaiian Islands. White dashed lines mark Pacific crustal ages in millions of years. Crustal ages are from Müller et al, 2008. The lack of offsets at fracture zones represents errors

830 *in the global data set. Red line and stars follow the tectonic backtrack of*
Palmyra Basin, on the northern Line Islands Ridge based on the fixed hotspot
Pacific Plate model of Engebretson et al, 1985, spliced to Gripp and Gordon
plate motions for 0-5 Ma . The location of Figure 2 is shown by the box and is
the location of MGL1208 survey area E.

835 *Figure 2: Bathymetric map of the Northern Line Island Ridge showing satellite*
bathymetry and multibeam mapping from MGL1208 and from Gardner and
Caulder (2010) The map also identifies Palmyra basin, Kingman Ridge,
Schlanger Ridge, and Langseth Ridge. Seismic lines 1 and 2 from the MGL1208
E area seismic survey are dashed, while the remainder of the MGL1208
840 *trackline is shown as a fine solid line. The box outlines the bathymetry shown in*
Figure 4.

Figure 3: Regional sediment thickness around the LIR from Divins (2003) and
Whittaker et al (2013). The black line outlines the 3000 m bathymetric contour
on the LIR. DSDP Site 165 is to the NW of LIR; Site 315 is to the east; and Site
845 *316 is to the south. Drilled sediment thicknesses are shown. The crest of the LIR*
is identified to be without sediment in this regional map, reflecting the lack of
data.

Figure 4: Bathymetry to the southeast of Seismic line 1, with the seismic reflection
profile displayed below. See Figure 2 for the orientation of the figure. The
850 *tracklines for seismic lines 1 and 2 are also shown as black lines, and include*
the trackline as the seismic gear was recovered after Seismic line 2 was
completed. The white line shows the ship's track after the seismic experiment.

Open circles mark the hourly position of the seismic source, while the black circles mark the position at 12-hour intervals marked GMT month/day in 2012.

855 *Red arrows mark the start and end of Seismic lines 1 and 2. The Seismic Line 2 profile is in Figure 5. A well-developed perched channel system is imaged in the northeast, next to Schlanger Ridge, where a seamount to the south of the main ridge forms a subbasin. A wide shallow channel system runs downslope from the Palmyra Basin depocenter. The location of sand waves and the seismic*
860 *profiles shown in figure 7 are shown on the right side of Seismic line 1.*

Figure 5: Seismic Line 2 from the MGL1208 Site E Survey, crossing the gap between Kingman and Langseth Ridges (west end) and the foot of Langseth Ridge. See Figure 2 and Figure 4 for the line position. Basal sediments tend to drape over topography, although they accumulated faster between basement ridges.

865 *Younger channels now erode into the older basin fill.*

Figure 6: (A) Bathymetric expression of sediment waves near the gap between Kingman and Langseth Ridges, and (B) maximum abyssal current vectors over LIR from the HYCOM model (Hybrid Coordinate Ocean Model; <http://hycom.org>), a data assimilating ocean circulation model with 9 km grid
870 *spacing. The 3000 m contour is highlighted to outline the Northern LIR. Black dots are locations of cores taken on MGL1208. Current vectors through the Kingman/Langseth Ridge gap are 0.2 m/s, flowing to the NE through the region where sediment waves were found.*

Figure 7: Seismic expression of sediment waves near the gap between Kingman and
875 *Langseth Ridges from Seismic Line 1. Top, digital subbottom profiler record at*

the same length scale as the bottom MCS record but with higher vertical exaggeration. The profile locations are shown on Figure 4. A prominent horizon may mark a hiatus period, while lower down in the sediment column sediment waves have eroded into older sediment.

880 *Figure 8: Seismic interpretation of sediment deposition within Palmyra Basin by interpreting horizons based upon the sediment column drilled at Site 315. Brown marks a sediment packet that is likely Cretaceous-Paleogene in age, blue is probably Oligocene-middle Miocene, and light green is probably late Miocene. Pink sediments are interpreted to be Pliocene-Pleistocene. There are*

885 *about 1.3 to 1.4 km of sediment at the depocenter along Seismic Line 1 based on seismic velocities at Site 315. Figure 9: Continuity of seismic horizons in Palmyra Basin (seismic line 1, top profile) compared to Cocos Ridge MV1014 Seismic Line 2 (bottom profile, Liao and Lyle, 2014; profile location is shown in Figure 10). Cocos Ridge forms the northern boundary of the Panama Basin in*

890 *the eastern Pacific. The basement age along MV1014-Line 2 is ~11.5 Ma, as compared to ~80 Ma along LIR Seismic Line 1. Both seismic lines are plotted at the same scale, and both used the same GI seismic source. Seismic horizons are much less continuous at LIR than at Cocos Ridge, evidence of higher current activity at the LIR.*

895 *Figure 10: Maximum current velocities from HYCOM above the Cocos Ridge. Magenta Line marks the position of MV1014 Seismic Line 2 from Liao and Lyle (2014) shown in Figure 9. Black dot is the location of ODP Site 1241 (ODP Leg 202 Scientific Party, 2003) Maximum current velocities range from 0.05 to 0.15*

m/s, or about 0.1 m/s slower than current velocities across the upper Palmyra

900 *Basin along LIR Seismic Line 1.*

10. REFERENCES

Adam, C. and A. Bonneville, 2005, Extent of the South Pacific Superswell. *Journal of Geophysical Research*, B110, B09408, 14pp, doi:10.1029/2004JB003465.

905

Arthur, M. A., W. E. Dean, and S. O. Schlanger (1985), Variations in the global carbon cycle during the Cretaceous related to climate, volcanism, and changes in atmospheric CO₂, in *AGU Geophysical Monograph 32*, pp 504-529, American Geophysical Union.

910

Aubert, O., 1994. Origin and stratigraphic evolution of the Maldives (central Indian Ocean, Rice University, Houston Texas.

915

Aubert, O., Droxler, A.W., 1996. Seismic stratigraphy and depositional signatures of the Maldivian carbonate system (Indian Ocean). *Marine and Petroleum Geology* 13 (5), 503-536.

920

Barth, Ginger A, Eakins, Barry W., Scheirer, Daniel S, Wong Florence L., and Jones, Meghan R., 2013; A fresh look at the geologic evolution of Kingman Reef and Palmyra Atoll, U.S. Line Islands. American Geophysical Union, Fall Meeting 2013, abstract #OS13B-1698,

925

Berger, W.H., Adelseck, C.G.J., and Mayer, L.A., 1976, Distribution of carbonate in surface sediments of the Pacific Ocean: *Journal of Geophysical Research*, 81, 2617-2627.

930

Betzler, C., Kroon, D., Reijmer, J.J.G., 2000. Synchronicity of major late Neogene sea level fluctuations and paleoceanographically controlled changes as recorded by two carbonate platforms. *Paleoceanography* 15 (6), 722-730.

Clark, J.D., Kenyon, N.H., Pickering, K.T., 1992. Quantitative analysis of the geometry of submarine channels: implications for the classification of submarine fans. *Geology* 20, 633-636.

- 935 Coffin, M. F. (1993). Emplacement and subsidence of Indian Ocean plateaus and submarine ridges. *Synthesis of results from scientific drilling in the Indian Ocean*, AGU Geophys. Monograph 70, 115-125.
- 940 Davis, A.S., L.B. Gray, D.A. Clague, J.R. Hein, 2002. The Line Islands revisited: New $^{40}\text{Ar}/^{39}\text{Ar}$ geochronologic evidence for episodes of volcanism due to lithospheric extension. *Geochemistry Geophysics Geosystems*, 3, 28 pp. , doi: 10.1029/2001GC000190
- 945 Divins, D.L., *Total Sediment Thickness of the World's Oceans & Marginal Seas*, NOAA National Geophysical Data Center, Boulder, CO, 2003.
www.ngdc.noaa.gov/mgg/sedthick/sedthick.html
- DSDP Leg 17 Shipboard Scientific Party, 1973. Site 165. Init Repts, DSDP, 17, 47-102.
- 950 DSDP Leg 33 Shipboard Scientific Party, 1976, Site 315. Init Repts, DSDP, 33, 37-104.
- DSDP Leg 33 Shipboard Scientific Party, 1976, Site 316. Init Repts, DSDP, 33, 105-159.
- 955 Duncan, R. A., and R. B. Hargraves, 1990. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of basement rocks from the Mascarene Plateau, the Chagos Bank, and the Maldives Ridge, in *Proc IODP, Sci Res.*, 115, edited, pp. 43-51, IODP, College Station TX.
- 960 Engebretson, D.C., Cox, A., Gordon, R.G., 1985. Relative Motions Between Oceanic and Continental Plates in the Pacific Basin. Geological Society of America Special Paper 206, 59 pp
- Flood, R.D., 1988. A lee wave model for deep-sea mudwave activity. *Deep Sea Research*, 35 (6), 973-983.
- 965 Gardner, J.V., and B.R. Calder, 2010. U.S. Law of the Sea Cruise to Map the Southern Flank of the Kingman Reef-Palmyra Atoll section of the Line Islands, Equatorial Pacific Ocean, University of New Hampshire, Center for Coastal and Ocean Mapping/Joint Hydrographic Center, Durham, 81 pp.
- 970 Goff, J.A., Smith, W.H.F., Marks, K.M., 2004. The contributions of abyssal hill morphology and noise to altimetric gravity fabric. *Oceanography* 17 (1), 24-37.
- 975 Gripp, A.E., Gordon, R.G., 1990. Current plate velocities relative to the hotspots incorporating the NUVEL-1 global plate motion model. *Geophysical Research Letters* 17 (8), 1109-1112.

- 980 Hamilton, E. L., R. T. Bachman, W. H. Berger, T. C. Johnson, and L. A. Mayer (1982),
Acoustic and related properties of calcareous deep-sea sediments, *Journal of
Sedimentary Petrology*, 52(3), 733-753.
- 985 Hays, J.D., Saito, T., Opdyke, N.D., Burckle, L.H., 1969. Pliocene-Pleistocene sediments
of the equatorial Pacific: Their paleomagnetic, biostratigraphic, and climatic
record. *Geological Society of America Bulletin* 80 (August), 1481-1514.
- Jackson, E.D., and S.O. Schlanger, 1976. Regional syntheses, Line Islands Chain,
Tuamotu Island Chain, and Manihiki Plateau, central Pacific Ocean. *Init Repts,
DSDP*, 33, 915-927. .
- 990 Johnson, G.C., Toole, J.M., 1993. flow of deep and bottom waters in the Pacific at
10°N. *Deep Sea Research I*, 40(2), 371-394.
- Jones, M.R., Eakins, B., and Barth, G., 2013, Seafloor geology of the U.S. Line Islands
Region (abs.): American Geophysical Union, 2013 Fall Meeting, OS13B-1699.
- 995 Larson, R.L., 1991, Latest pulse of Earth: Evidence for a mid-Cretaceous superplume.
Geology, 19, 547-550.
- 1000 Liao, Y., Lyle, M., 2014. Late Miocene to Pleistocene sedimentation and sediment
transport on the Cocos Ridge, eastern tropical Pacific Ocean. *Marine Geology*
355, 1-14.
- 1005 Lyle, M., Wilson, P.A., 2006. Leg 199 synthesis: evolution of the equatorial Pacific in
the early Cenozoic. In: Wilson, P.A., Lyle, M., Firth, J.V. (Eds.), *Proceedings of
the ODP, Scientific Results*, 199. Ocean Drilling Program, College Station TX,
pp. 1-39.
- 1010 Lynch-Stieglitz, J., et al. (2015), Glacial-interglacial changes in central tropical Pacific
surface seawater property gradients, *Paleoceanography*, 30, 423-438,
doi:10.1002/2014PA002746.
- 1015 Marcantonio, F., Lyle, M., Ibrahim, R., 2014. Particle sorting during sediment
redistribution processes and the effect on ²³⁰Th-normalized mass
accumulation rates. *Geophysical Research Letters* 41, 5547-5554.
- Mayer, L.A., Shipley, T.H., Winterer, E.L., 1986. Equatorial Pacific seismic reflectors
as indicators of global oceanographic events. *Science* 233, 761-764.

- 1020 Miller, M.C., Komar, P.D., 1977. The development of sediment threshold curves for unusual environments (Mars) and for inadequately studied materials (foram sands). *Sedimentology* 24, 709-721.
- 1025 Mitchell, N., Lyle, M., 2005. Patchy deposits of Cenozoic pelagic sediments in the central Pacific. *Geology* 33, 49-52.
- Morgan, W.J., 1972. Deep mantle convection plumes and plate motions. *American Association of Petroleum Geologists Bulletin*, 56, 203-213.
- 1030 Müller, R.D., Sdrolias, M., Gaina, C., Roest, W.R., 2008. Age, spreading rates, and spreading asymmetry of the world's ocean crust. *Geochemistry, Geophysics, Geosystems* 9 (4), 19 pp.
- 1035 Neal, C. R., Mahoney, J. J., Kroenke, L. W., Duncan, R. A., & Petterson, M. G. (1997). The Ontong Java Plateau. *Large Igneous Provinces: Continental, Oceanic, and Planetary Flood Volcanism*, 183-216.
- 1040 Normark, W.R., Spiess, F.N., 1976. Erosion on the Line Islands archipelagic apron: effect of small-scale topographic relief. *Geological Society of America Bulletin* 87, 286-296.
- ODP Leg 202 Scientific party, 2003. Site 1241, Proceedings of the Ocean Drilling Program, Initial Reports, doi:10.2973/odp.proc.ir.202.112.2003
- 1045 Orwig, T.L., 1981. Channeled turbidites in the eastern central Pacific Basin. *Marine Geology* 39, 33-57.
- 1050 Pälike, H., Lyle, M., Nishi, H., Raffi, I., Ridgwell, A., Gamage, K., Klaus, A., and the Exp 320/321 Shipboard scientific party, 2012. A Cenozoic record of the equatorial Pacific carbonate compensation depth. *Nature* 488, 609-614.
- 1055 Paskevich, V.F., Wong, F.L., O'Malley, J.J., Stevenson, A.J., and Gutmacher, C.E., 2011, GLORIA sidescan-sonar imagery for parts of the U.S. Exclusive Economic Zone and adjacent areas: U.S. Geological Survey Open-File Report 2010-1332, <http://pubs.usgs.gov/of/2010/1332/>
- Pockalny, R., Scheirer, D., Barth, G., Eakins, B., 2015. A two-phase volcanic model for the origin of Line Islands Ridge. AGU Fall meeting.

- 1060 Raff, A.D., 1973. Underway Data. Initial Reports, Deep Sea Drilling Project 17, 337-364.
- Rosburg, K. C., K. A. Donohue, E. Chassignet, 2012. Performance Evaluation of HYCOM Gulf of Mexico. Ocean Sciences Meeting, Salt Lake City, Utah
- 1065 Schlanger, S.O., Winterer, E.L, 1976. Underway geophysical data: navigation, bathymetry, magnetics, and seismic profiles. . Init Repts, DSDP, 33, 655-693.
- 1070 Schlanger, S.O., Garcia, M.O., Keating, B.H., Naughton, J.J., Sager, W.W., Haggerty, J.A., Philpotts, J.A., 1984. Geology and geochronology of the Line Islands. Journal of Geophysical Research, Earth 89 (B13), 11,261-11,272.
- Smith, W.H.F., Sandwell, D.T., 1997. Global seafloor topography from satellite altimetry and ship depth soundings. Science 277, 1957-1962.
- 1075 Stow, D.A.V., Hernandez-Molina, F.J., Llave, E., Sayago-Gil, M., Diaz del Rio, V., Branson, A., 2009. Bedform-velocity matrix: the estimation of bottom current velocity from bedform observations. Geology 37 (4), 327-330.
- 1080 Talley, L.D. 2007. Hydrographic Atlas of the World Ocean Circulation Experiment (WOCE): Volume 2: Pacific Ocean. International WOCE Project Office, Southampton, UK. www-pord.ucsd.edu/whp_atlas/Pacific_index.html
- 1085 Torsvik, T.H., Müller, R.D., Van Der Voo, R., Steinberger, B., Gaina, C., 2008. Global plate motion frames: toward a unified model. Reviews of Geophysics 46, 44 pp.
- van Andel, T.H., Heath, G.R., Moore, T.C., 1975. Cenozoic History and Paleooceanography of the central equatorial Pacific Ocean, GSA Memoir 143.
- 1090 Geological Society of America.
- Whittaker, Joanne, Alexey Goncharov, Simon Williams, R. Dietmar Müller, German Leitchenkov (2013) Global sediment thickness dataset updated for the Australian-Antarctic Southern Ocean, Geochemistry, Geophysics, Geosystems. DOI: 10.1002/ggge.20181
- 1095

Figure 1: Location map, central Pacific

Figure 2: Northern LIR map

Lyle et al, Figure 2

Figure 3. regional sediment thickness map

Figure 4: Palmyra Basin multibeam plus seismic

Figure 5: Seismic line 2

Figure 6: LIR sedwaves, map and currents

Figure 7: LIR sedwaves seismic profiles

Figure 8: Interpretation of seismic line 1

Figure 9: comparison of LIR and Cocos R seismic profiles

Figure 10: currents on Cocos Ridge

HYCOM Max. Velocity (m/s)