Detection and Risk Assessment of Diarrheagenic *E. coli* in Recreational Beaches of Brazil

Vanessa F.V. Rodrigues^{1,2}, Irma N. G. Rivera¹, Keah-Ying Lim², Sunny C. Jiang^{2,*}

¹Biomedical Science Institute, Microbiology Department, University of São Paulo, Estado de São Paulo, SP, 05508-000, Brazil

²Department of Civil and Environmental Engineering, University of California, Irvine, CA 92697-2175, U.S.A.

Running title: Toxic *E. coli* in Brazilian Beaches

*Corresponding author: 844E Engineering Tower, University of California, Irvine, CA 92617; Email: sjiang@uci.edu; Tel: 949-824-5527

Abstract

Marine beaches are important recreational and economic resources in Brazil, but the beaches' water quality is negatively impacted by the discharge of domestic sewage effluent. The occurrence of diarrheagenic *E. coli* among the *E. coli* isolated from three Brazilian marine beaches was investigated. Multiplex and single step PCR were used to screen 99 *E. coli* isolates for ten target toxin genes. Six toxin genes, *stx1*, *eae*, *estp*, *esth*, *astA*, and *bfpA*, were identified in 1% to 35% of the isolates. A quantitative microbial risk assessment (QMRA) of human exposure to diarrheagenic *E. coli* during marine recreation was carried out. The results indicated that the diarrheagenic *E. coli* risk is well below the U.S. EPA's recommended daily recreational risk benchmark. However, the overall recreational health risk due to all pathogens in the water could be much higher and exceeded the U.S. EPA's benchmark.

Keywords: diarrheagenic *E. coli*, recreational health, quantitative microbial risk assessment, Brazilian beaches

Graphical abstract

- What is the recreational health risk posed by the presence of diarrheagenic *E. coli* in the water of Brazilian beaches?
- Is the recreational water quality criteria of Brazil adequate for the health protection of bathers?

Introduction

Marine recreational waters are often affected by both point source pollution from sewage outfalls and industrial discharges, and non-point source pollution from diffused sewage discharges, stormwater runoff, resuspension of contaminated sediments, human bather's shedding, and animal fecal inputs (**Hamilton et al., 2010; King, 2013; Rivera et al., 2008; Rivera and Martins, 1996; U.S. EPA**). In Brazil, marine beaches are important recreational and economic resources, but the beaches' water quality is negatively impacted by agriculture discharges and effluent of partially treated or untreated domestic sewage. According to Trata Brazil Institute, only 62.3% of human sewage generated in Brazil in 2013 was treated (**Instituto Trata Brasil, 2014**). The scenario is the worst in coastal regions — in Baixada Santista region (São Paulo state), only 60% of human sewage was collected by sewer systems with the remaining 40% released directly to the environment as diffused source of discharge. Of the 60% sewage collected through sewer systems, only 10% was fully treated using conventional wastewater treatment process, whereas the remaining was partially treated through gross removal of solids and chlorination (**Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2005**). Discharge of untreated or partially treated human sewage to beaches is usually elevated during peak recreational season due to the influx of tourist population that generates excess sewage, which overwhelms the sewer collection system resulting in direct discharge of raw sewage to the beaches as diffused sources. The excess sewage collected through the sewer system is only partially treated because of the limited treatment capacity of the local wastewater treatment plants (**Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2005**).

The untreated or improperly treated sewage contains toxic chemicals, pathogens, and high concentrations of organic and inorganic nutrients, which can have adverse health effects on recreational bathers through inhalation of aerosols, accidental ingestion of water, and wound infection (**Abdelzaher et al., 2010; Abessa et al., 2005; Braga et al., 2000; Westrell et al., 2004; World Health Organization, 1999**). Gastrointestinal illnesses (GI) are the most commonly reported and documented recreational water diseases, which are caused by enteric viruses, bacteria, and protozoa that are commonly associated with human sewage (**Centers for Disease**

Control and Prevention; Fleisher et al., 2010; Turbow et al., 2008). The high nutrient concentration associated with sewage discharge can also trigger the proliferation of native marine opportunistic pathogens such as *Vibrio spp.*, which causes both GI and wound infections **(Dickinson et al., 2013).** Eyes, ear, and skin infections among recreational bathers are also well documented, but their infective agents have not been well identified and quantified in recreational waters impacted by both point source **(Haile et al., 1999; Wade et al., 2010)** and non-point source pollution **(Colford et al., 2012; Colford Jr et al., 2007; Fleisher et al., 2010).**

In order to protect the health of beachgoers in Brazil, the environmental sanitation agency of São Paulo State performs routine monitoring of the microbial water quality of beaches for fecal indicator bacteria (thereafter abbreviated as “FIB” that include enterococci, thermotolerant coliform, and *E. coli*). The current water quality criteria for recreational waters are based in Conama Resolution n° 274/2000, which classifies beaches into two categories of use: “Proper” or “Improper”. For a beach to be considered “Proper” for safe recreational activities, the water samples must contain less than 100 enterococci, 1000 thermotolerant coliform, and 800 *E. coli* per 100 mL in 80% of the five samples collected from the same sampling point for five consecutive weeks **(Conselho nacional do meio ambiente (CONAMA), 2008).** However, it is unclear whether these water quality criteria are protective of the health of bathers.

The level of FIB in recreational water is traditionally used to indicate human sewage contamination of the water; a higher FIB level generally corresponds to a higher recreational health risk. This is, however, a gross generalization because a specific type of FIB (e.g. enterococci, thermotolerant coliform, or *E. coli*) may be more suitable to indicate the recreational risk in a specific type of water (e.g. fresh or marine). Putting this into context, the U.S. EPA set different thresholds of FIB level based on the type of recreational waters; *E. coli* are only used for indicating the health risk of recreational freshwater whereas enterococci are used for both marine and freshwater. Furthermore, these thresholds are established based on a large-scale epidemiological study in the U.S.—the National Epidemiological and Environmental Assessment of Recreational (NEEAR) water studies, wherein the threshold level of each FIB corresponds to a range of NEEAR-gastrointestinal illness (NGI) rates of 32 or 36 illness cases per 1000 bathers

(the definition of the **(U.S. EPA, 2012)**). These illness rates also serve as a measure for the level of protection provided to assure the safety of people recreating in the waters. In comparison, there is no clear rationale for the selection of the FIB standard set by the Conama Resolution n° 274/2000. For example, the threshold level of 800 CFU/100 mL *E. coli* concentration set in Conama Resolution n° 274/2000 (regardless of the type of recreational water) is significantly higher than the 126 CFU/ 100 mL *E. coli* threshold (geometric mean of samples collected over thirty days and no sample exceeds 410 CFU/100 mL) set by U.S. EPA for sewage-impacted freshwater. Again, the 100 CFU/100mL enterococcus threshold of Conama Resolution is also more than twice as high as the U.S. EPA threshold of 35 CFU/100mL. Ultimately, this leads to the important question regarding the adequacy of Conama Resolution n° 274/2000's water quality criteria for recreational health protection due to the unknown health risk that is associated with these criteria.

In the absence of a costly epidemiological study, recreational health risks of sewage-impacted beaches can be estimated through understanding the occurrence of pathogen, the level of human exposure, and the pathogen dose-response relationship, following the Quantitative Microbial Risk Assessment (QMRA) framework. Towards assessing the recreational health risks of Brazilian beaches, we investigated the fraction of diarrheagenic *E. coli* among the total *E. coli* population isolated from water samples of three Brazilian coastal regions impacted by different levels of human sewage contamination in order to understand the recreational risk specific to this group of pathogen.

Diarrheagenic *E. coli* is a well-known etiological agent of human diarrheal diseases (**Kaper and Hacker, 1999; Nataro and Kaper, 1998; Russo and Johnson, 2000**), which represents a small portion of the diverse serotypes and strains of *E. coli* found in human and other warm-blooded animal guts. Diarrheagenic *E. coli* are grouped into six categories: enteropathogenic (EPEC), enterotoxigenic (ETEC), enteroinvasive (EIEC), enterohaemorrhagic (EHEC), enteroagregative (EAEC), and *E. coli* with diffuse adherence (DAEC) (**Nataro and Kaper, 1998**). The pathogenicity of *E. coli* in the diarrheagenic categories is usually assessed through searching for virulence-associated genes, such as *estp*, *esth* and *elt* in ETEC; *bfpA* and *eae* in EPEC; *stx1*, *stx2* and *eae* in

EHEC; *invE* in EIEC, and *aggR* and *astA* in EAEC (**Fujioka et al., 2013**). In spite of the significant burden of diarrheagenic *E. coli* to human health, its presence/abundance has not been investigated extensively in recreational beach waters due to the technical challenges for detecting toxin genes. Here we report the detection of EPEC, ETEC, EIEC, EHEC and EAEC in three coastal regions in the São Paulo state using PCR methods. A QMRA model was then used to estimate the health risks associated with exposure to diarrheagenic *E. coli* during recreational activities at those beaches. Finally, the total recreational illness risk is then inferred through analyzing the contribution of diarrheagenic *E. coli* to the total risk of recreating in sewage-impacted water.

The role of *E. coli* as an indicator of marine recreational water health risk has been investigated in several previous studies. For example, Bonamano, et al. (**2015**) modeled the dispersion of viable and total *E. coli* cells in an artificial semi-enclosed bathing area of an Italian beach. The study showed that a high concentration of *E. coli* was found during the summer tourist season, which was attributed to the overflow of sewage treatment plants. However, the limitations of *E. coli* as an indicator for marine recreational beach safety is also well known, which is further highlighted by the U.S. EPA recommendation of using enterococcus as the indicator due the sensitivity of *E. coli* to osmotic stress (**U.S. EPA, 2012**). A primary goal of this study is to improve the current understanding of *E. coli* in marine waters through evaluation of the contribution of diarrheagenic *E. coli* to recreational health risks. The results of this study provide scientific ground that can be used for supporting the further development of the recreational water quality criteria set by the Conama Resolution n° 274/2000.

Materials and Methods

Study sites.

The water samples were collected between 2005 and 2007 from three coastal regions in the São Paulo State, namely, Baixada Santista (BS), São Sebastião Channel (SS), and Ubatuba (U) (Figure 1). Each region was impacted by different levels of anthropogenic pollution from both sewage, industrial outfalls, and diffused discharge of untreated raw sewage and agriculture runoff (**Burbano-Rosero et al., 2011**). Sewage generated in these coastal regions is either discharged into the coastal environment directly untreated, or is collected by sewer systems and sent to treatment plants. Of the sewage collected, a portion is treated using conventional biological wastewater treatment process and discharged directly to surface waters; the remaining portion is partially treated through the removal of gross solids followed by chlorination. This partially treated sewage is released into the ocean through submarine outfalls that are situated 8-24 meters below the water surface (**Companhia de Tecnologia de Saneamento Ambiental (CETESB)**). Little is known of the impact of these minimally treated sewage outfalls to recreational beaches. Neither the transport pattern of sewage plumes nor the microbiological quality of the surrounding waters was monitored. These are in dramatic contrast to the highly regulated submarine outfalls in the U.S.

Santos/ São Vicente Bay, in Baixada Santista, is the entry point of two main urban waterways to the Atlantic Ocean. Santos and São Vicente Channels collect diffused urban runoff and sewage effluents from two of the most populated coastal cities in São Paulo State — São Vicente and Santos with a combined population of ~743,000 inhabitants in 2005 (**Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2014**). Baixada Santista coast is also the home of four submarine sewage outfalls as shown in Figure 1. Samples for this study were collected offshore from the mouth of São Vicente (BS1) channel, a preserved area (BS2), and near the mouth of Santos channel (BS3).

Figure 1: Recreational beaches (Ubatuba, São Sebastião Channel and Baixada Santista) in coastal region São Paulo state investigated in this study

São Sebastião channel is 25 km in length and flows between the city of São Sebastião and the Ilhabela island. The two cities were reported to have ~73,000 and ~25,000 inhabitants in 2005, respectively. The region is famous for its beaches, which makes it a popular tourist destination, especially for people from the state of São Paulo. Four outfalls were identified in the region— three submarine outfalls releasing partially treated sewage, and one releasing industrial sewage (Figure 1). Samples were collected offshore near the Marine Biology Center – CEBIMar- USP in front Praia do Segredo (SS1) and near the Baraqueçaba beach (SS2).

The last region included in this study, Ubatuba, is a small tourist city with ~79,000 inhabitants. Approximately 33% of the sewage generated in the region is collected and treated by sewage treatment plants. The region is also not known to have any submarine outfalls. Samples were collected from two sites offshore near Saco da Ribeira, at a small dock for tourist boats (UB1 and UB2).

All water samples were collected onboard a small boat using sterile sampling equipment following standard methods for the examination of water and wastewater (**WEF and APHA, 2005**, and also see (**Burbano-Rosero et al., 2011**) for additional details). The sampling sites were located between the shoreline and sewage outfalls to capture the impact from both diffused onshore surface discharge and the submarine discharge of partially treated sewage. Samples were taken monthly during the summer recreational peak season between February and May 2006 (February to April 2006 for Ubatuba) and between January and March 2007. All samples were transported in ice coolers and processed within 6 hours of collection (**WEF and APHA, 2005**).

***E. coli* quantification and isolation.**

The *E. coli* concentration (CFU/ 100 mL) was determined by membrane filtration technique on mFC agar (Difco). The blue colonies were isolated and counted as *E. coli* based on biochemical reactions. All *E. coli* isolates from the three regions were deposited in the collection of Microbial Ecology Laboratory of Biomedical Science Institute (ICB) at University of São Paulo, Brazil (USP). Detailed information for *E. coli* isolation is described in (**Burbano-Rosero et al., 2011**).

Genomic DNA extraction and PCR amplification of virulence-associated genes of diarrheagenic E. coli.

The *E. coli* isolates stored frozen in glycerol were transferred to Luria-Bertani (LB) broth (Difco) for recovery, then individual colonies were re-isolated on MacConkey and Eosin Methylene Blue Agar plates (Difco). The biochemical identification of each isolate was re-confirmed by IMViC (Indole, Methyl red, Voges-Proskauer, and in Citrate test) following the American Public Health Association's (APHA) recommendation (**WEF and APHA, 2005**). Isolates that matched *E. coli* profile were incubated overnight at 37°C in LB broth and were used for the genomic DNA extraction using the *Wizard Genomic DNA Purification* (Promega) kit following the manufacturer's instructions. The DNA quality, concentration, and purity were evaluated through electrophoresis on 1.0% agarose gel in 1x TAE buffer (40 mM Tris acetate, 2 mM Na₂EDTA·2H₂O [pH 8.0]) and

spectrometrically (NanoDrop 1000 Thermo Scientific). Reconfirmed *E. coli* isolates were utilized for genetic identification of virulence genes, wherein 30 isolates were collected from Baixada Santista, 49 from São Sebastião Channel, and 20 from Ubatuba.

The PCR oligonucleotide primers used in this study are listed in Table 1 and the reaction conditions were adapted from a protocol by Fujioka et al. **(Fujioka et al., 2013)**. Two PCR assays were performed to verify the occurrence of target virulence genes. The first PCR was a multiplex-PCR targeting genes *stx1*, *stx2*, *eae*, *bfpA*, *aggR*, *elt*, *invE*, and *astA*. The reaction was performed using the HotStart PCR kit by PreMix Bioneer (Korea) containing primers (20µM stock solution concentration) and DNA template (100 ng/µL) as following: 0.1 µL each of *stx2*F and *sxt2*R; 0.2 µL each of *stx1*F, *sxt1*R, *elt*F, *elt*R, *invE1*, *invE2*; 0.25 µL each of *bfpA*F and *bfpA*R; 0.3 µL each of *astA*F and *astA*R; 0.35 µL each of *eae*F, *eae*R, *aggR1* and *aggR2*, 5 µL of the DNA template, and the final volume was adjusted to 20 µL with nuclease free water (Sigma). The mixture was preheated at 94 °C for 5 min, and then amplified for 35 cycles using the program: denaturation at 94 °C for 1 min, annealing at 55 °C for 1 min, extension at 72 °C for 1 min with a final extension step at 72 °C for 10 min. The PCR products were separated by electrophoresis (100 V, 10 min, after 90 V, 1 h 20 min) in a 1.5% agarose gel (Invitrogen) and stained by ethidium bromide (0.5 µg/mL) and photographed in the photosystem Quantity One (Bio-Rad). The sample was scored as positive for the specific gene when the expected size amplicons were observed. All the positive reactions from multiplex PCR were re-confirmed by single step PCR.

In the second PCR assay, individual amplifications for the *estp* and *esth* genes were carried out. The PCR cycles include the initial denaturation at 94 °C for 5 min, followed by 35 cycles of 95 °C, 15 sec; 55 °C, 30 sec; 68 °C, 10 sec and the final extension at 68 °C for 3 min. In all reactions, reference *E. coli* strains were used as positive controls.

Table 1. Primers used for PCR amplification of the virulence genes for five categories of diarrheagenic *E. coli*

Target gene	Primer name	Primer sequence (5' to 3')	T _m (°C)	Amplicon size (bp)	Reference
<i>stx1</i>	<i>stx 1 F</i>	AGTTAATGTGGTGGCGAAGG	60	347	(Fujioka et al., 2013)
	<i>stx 1 R</i>	CACCAGACAATGTAACCGC	58		
<i>stx2</i>	<i>stx 2-F</i>	TTCGGTATCCTATTCCCGG	58	589	(Fujioka et al., 2013)
	<i>stx 2-R</i>	CGTCATCGTATACACAGGAG	60		
<i>eae</i>	<i>eae-F</i>	CCCGAATTCGGCACAAGCATAAGC	74	881	(Oswald et al., 2000)
	<i>eae-R</i>	CCCGGATCCGTCTCGCCAGTATTCG	82		
<i>bfpA</i>	<i>bfpA-F</i>	AATGGTGCTTGCGCTTGCTGC	66	324	(Gunzberg et al., 1995)
	<i>bfpA-R</i>	GCCGCTTTATCCAACCTGGTA	64		
<i>aggR</i>	<i>aggR-1</i>	GTATACACAAAAGAAGGAAGC	58	254	(Ratchtrachenchai et al., 2012)
	<i>aggR-2</i>	ACAGAATCGTCAGCATCAGC	60		
<i>elt</i>	<i>elt-F</i>	AACGTTCCGGAGGTCTTATG	60	511	(Fujioka et al., 2013)
	<i>elt-R</i>	CAACCTTGTGGTGCATGATG	60		
<i>esth</i>	<i>esth-F</i>	TTCACCTTTCCCTCAGGATG	60	172	(Fujioka et al., 2013)
	<i>esth-R</i>	ATAGCACCCGGTACAAGCAG	62		
<i>estp</i>	<i>estp-F</i>	ACTGAATCACTTGACTCTTCA	58	120	(Fujioka et al., 2013)
	<i>estp-R</i>	TCACAGCAGTAAAATGTGTTGT	60		
<i>invE</i>	<i>invE-1</i>	GCAGGAGCAGATCTTGAAG	58	208	(Fujioka et al., 2013)
	<i>invE-2</i>	GAAAGGCACGAGTGACTTTC	60		
<i>astA</i>	<i>astA-F</i>	CCATCAACACAGTATATCCG	58	101	(Fujioka et al., 2013)
	<i>astA-R</i>	ACGGCTTTGTAGTCCTTCCA	60		

Quantitative microbial risk assessment.

The health risks associated with exposure to diarrheagenic *E. coli* recreating in contaminated beaches were assessed using the QMRA framework as outlined by National Research Council (**National Research Council, 1983**). QMRA was carried out using the Monte Carlo technique to account for uncertainties and variability associated with risk estimates. All the computations were performed using MATLAB R2012a (The Mathworks Inc., Natick, MA).

Hazard identification.

The hazard of human exposure to *E. coli* harboring virulence-associated genes is well-documented (**Koczura et al., 2015**). The percentage of *E. coli* isolates from each region exhibiting virulence-associated genes was used to estimate the percentage of diarrheagenic *E. coli* in the total *E. coli*. For this purpose, the PCR results for the presence/absence of virulence genes in each isolate were input into a binary matrix containing 1's and 0's to represent "positive" and "negative" (amplifications and no amplifications), respectively (see APPENDIX). It has been noted that an isolate can harbor more than one virulence gene. In determining the number of virulence strains among *E. coli* isolates, the isolate with more than one virulence genes was only counted once. The concentration of diarrheagenic *E. coli* was estimated using

$Conc_{d\ E. coli} = \% \text{ virulence} \times Conc_{E. coli}$, where $Conc_{d\ E. coli}$ is the estimated concentration of diarrheagenic *E. coli* (CFU/100mL), $\% \text{ virulence}$ is the percentage of isolates that are identified to contain at least one virulence gene (%), and $Conc_{E. coli}$ is the concentration of total *E. coli* at the study site (CFU/100mL).

Since the concentration of *E. coli* at each study site varies with time, the probability of encountering a specific concentration of *E. coli* is determined using a triangular distribution, in which the lower bound, peak, and upper bound of the distribution were characterized by the minimum, arithmetic mean, and maximum *E. coli* values from each site, respectively (**Haas, 1996; Haas et al., 1999**).

Exposure assessment.

The level of human exposure to diarrheagenic *E. coli* during water recreation is dependent on gender, age group, and the types of recreational activities (**U.S. EPA, 2012**), which results from the different volume of water ingested during a marine swimming event. Based on a previous exposure study (**Schets et al., 2011**), the volume of water ingested by each gender and age group during a recreational event can be characterized as a gamma distribution, wherein the volume of water ingested (mL) during a swimming event is $G(r=0.45, \lambda=60)$ for men, $G(r=0.51, \lambda=35)$ for women, and $G(r=0.58, \lambda=55)$ for children. Assuming independency between pathogen concentration and water volume ingested, the pathogen exposure dose is then calculated as the joint probability of water ingested during marine swimming event and concentration of pathogenic *E. coli* as: $D_{ingested} = Vol_{ingested} \times Conc_{d\ E. coli}/100$, where $D_{ingested}$ is the estimated dose of diarrheagenic *E. coli* ingested (CFU), $Vol_{ingested}$ is the volume of water ingested by a human subject (mL), and $Conc_{d\ E. coli}$ is the estimated concentration of diarrheagenic *E. coli* (CFU/100mL).

Dose-response assessment.

The probability of illness per swimming event was estimated using the Beta-Poisson model for virulent *E. coli* O157:H7, $P_{ill} = 1 - \left(1 + \frac{D_{ingested}}{\beta} \right)^{-\alpha}$, where P_{ill} is the probability of illness due to a swimming event (**Haas et al., 2000**). The best-fit estimates for the model, $\alpha = 0.49$ and $\beta = 1.89 \times 10^5$, are used (**Haas et al., 2000**).

Simulation of the worst-case scenario.

For simulating the worst case scenario that represents the maximum human health risk from exposure to diarrheagenic *E. coli* during water recreation, the maximum allowed concentration of *E. coli* (800 CFU/ 100 mL) based on the Conama Resolution n° 204/2000 was used as the worst-case scenario benchmark. This benchmark is multiplied by the mean value for the percentage of virulence-associated genes of diarrheagenic *E. coli* encountered at all study sites to estimate the maximum

diarrheagenic *E. coli* concentration, which is then used in the QMRA model for estimating the maximum health risk associated with diarrheagenic *E. coli*.

Results

Concentration of *E. coli*.

The minimum value for *E. coli* concentration was assumed to be 0.99 CFU/100 mL at certain sampling dates where the presence of *E. coli* was below the quantifiable limit (≤ 1 CFU/100mL). The minimum (min), arithmetic mean (mean), and maximum (max) concentration (CFU/ 100 mL) of *E. coli* from the three coastal regions were: 0.99, 154, and 912 at Baixada Santista; and 0.99, 7.84, and 86 at São Sebastião Channel; while all values are ≤ 1 CFU/100mL at Ubatuba.

Detection of virulence-associated genes of diarrheagenic *E. coli*.

The *stx2*, *aggR*, *elt*, and *invE* genes were not detected in all *E. coli* isolates examined. Among the Ubatuba samples (n=20), virulence-associated genes were detected 20% of the time, wherein *astA* and *bfpA* genes were detected in 15% and 5% of the isolates, respectively. For samples from Baixada Santista (n=30), 30% of the isolates contained *astA* and 3% contained *esth* genes. At São Sebastião channel (n=49), 47% of the *E. coli* isolates were positive for *astA*, 4% positive for *eae* and 2% isolates also contained *stx1* or *estp* (2%, each). The summary of virulence-associated genes detected among isolates is shown in Table 2.

Table 2: Number of isolates showing virulence-associated genes of diarrheagenic *E. coli*, frequency (%) of detection by PCR among isolates, and the total *E. coli* concentration from Ubatuba, Baixada Santista, and São Sebastião Channel, São Paulo state, Brazil.

Sampling sites	# of isolates	Genes						Total <i>E. coli</i> concentration (CFU/100mL)			
		<i>stx1</i>	<i>eae</i>	<i>estp</i>	<i>esth</i>	<i>astA</i>	<i>bfpA</i>	<i>Min</i>	<i>Max</i>	<i>A.Mean</i>	<i>G.Mean</i>
São Sebastião Channel	49	1 (2%)	2 (4%)	1 (2%)	0	23 (47%)	0	1.0	86.0	7.8	1.7
Baixada Santista	30	0	0	0	1 (3%)	9 (30%)	0	1.0	910.0	154.8	18.5
Ubatuba	20	0	0	0	0	3 (15%)	1 (5%)	1.0	1.0	1.0	1.0
Total	99	1	2	1	1	35	1				

Risk Characterization.

Ubatuba region was not considered in the risk assessment because of its low concentration of total *E. coli* (<1 CFU/100mL) and low detection frequency for virulence-associated genes of diarrheagenic *E. coli*. A comparison of the recreational risks at Baixada Santista and São Sebastião Channel showed that the diarrheagenic *E. coli* illness risk was over an order-of-magnitude higher at Baixada Santista than São Sebastião Channel. The 95th percentile illness risk is 2.86×10^{-4} for men, 1.90×10^{-4} for women, and 3.43×10^{-4} for children recreating at Baixada Santista; 4.29×10^{-5} to men, 2.83×10^{-5} for women, and 4.90×10^{-5} for children at São Sebastião Channel. The worst-case scenario (95th percentile) of the risk is 8×10^{-4} for men, 5.42×10^{-4} for women and 8.73×10^{-4} for children. The boxplot of illness risk (Figure 2) indicated the illness risk due to exposure to diarrheagenic *E. coli* at both beaches are below the 32—36 NGI per 1000 ($3.2 - 3.6 \times 10^{-3}$) bathers set by the U.S. EPA for the recreational health risk (U.S. EPA, 2012).

Figure 2: Box-and-whisker plot showing the illness risks among swimmers from different sex/age groups recreating at São Sebastião Channel and Baixada Santista. The illness risks that correspond to the *E. coli* criterion set by Conama Resolution n°274/2000 for each sex/age group are also plotted for comparison. Each box represents the lower, median, and upper quartile (e.g. 25th, 50th, and 75th percentile values) of the distribution, where the whiskers extend $1.5 \times (75^{\text{th}} \text{ percentile value} - 25^{\text{th}} \text{ percentile value})$ from each end of the box. Markers graphed outside of each whisker are considered as outliers.

Discussion

Diarrheogenic *E. coli* in marine waters.

E. coli is used as a water quality indicator in many parts of the world to indicate fecal pollution of water and to guide bathing water quality management. Studies from around the globe have continued to learn about the many different factors that can affect the behaviors of *E. coli* in the water, and thus, its implications in risk management. For example, an investigation of *E. coli* pollution in a Baltic sea lagoon (**Schippmann et al. 2013**) showed that *E. coli* concentration only gives an incomplete picture of sources and risks of pollutants, which does not support the decision making for bathing water quality management. In the contrast, a modeling study conducted in Hong Kong indicated that the fate and transport of *E. coli* from a sewage outfall near eight marine beaches were successfully predicated by spatial-temporal decay of *E. coli* in marine water (**Chan et al. 2013**). In another study, *E. coli* concentration was found between 0-90 CFU/100mL along Teluk Kemang beach in Malaysia and the concentration was significantly correlated with temperature and oxidation reduction potential of the water (**Praveena et al. 2013**). So far, limited studies have investigated the occurrence of diarrheogenic *E. coli* in recreational marine waters (**Albertini; Almeida et al., 2012; Hamilton et al., 2010**) due to the technical challenges in genetic identification of diarrheogenic genes and the low occurrence of diarrheogenic *E. coli* among the total *E. coli*.

E. coli populations are highly diverse in phenotypes, ecological niches and survival characteristics (**Selander and Levin, 1980**). Some studies suggest that diarrheogenic *E. coli* strains were derived from commensal strains by acquisition of virulence-associated genes localized in chromosomes or mobile genetic elements (**Finlay and Falkow, 1997**). Diarrheogenic *E. coli* are categorized to EPEC, ETEC, EIEC, EAEC, EHEC and DAEC groups (**Nataro and Kaper, 1998; World Health Organization, 1999**). However, some authors questioned the importance of DAEC as a diarrheogenic category. Only the virulent gene markers from the first five groups of diarrheogenic *E.*

coli were screened among *E. coli* isolates from marine coastal waters in this study to grasp the level of human exposure to this group of pathogen.

The disease outbreaks associated with diarrheagenic *E. coli* during freshwater recreations are frequently reported in the U.S. and European countries due to the existence of a comprehensive surveillance system (**Centers for Disease Control and Prevention; Rangel et al., 2005; Verma et al., 2007**). Children are among the most vulnerable population to the exposure of diarrheagenic *E. coli* because of their higher rate of water ingestion during water recreation and lower rate of immunity (**Wade et al., 2008**). For this reason, a risk assessment of diarrheagenic *E. coli* in recreational waters is important to understand the public health risks in sewage-impacted water.

The *eae* gene was detected in two of the 49 isolates (4%) from São Sebastião Channel, one of them was also positive for *estp* gene, which identifies the isolate as atypical EPEC or ETEC. Similar detection frequency (3.6%) of *eae* gene was reported among 24,493 *E. coli* isolated from marine recreational waters at Avalon Bay, California (**Hamilton et al., 2010**), a beach location that has documented sewage contamination (**Boehm et al., 2003**).

The co-occurrence of *bfpA* and *astA* genes was identified in one of 20 isolates at Ubatuba, which characterizes the isolate as a potential typical EPEC or EAEC. The low frequency of *bfpA* gene detection in samples from marine waters also agrees with the results of Hamilton et al. (**Hamilton et al., 2010**), who evaluated the occurrence of typical EPEC by searching for EAF plasmid and found only eight positive among 24,493 isolates (<1%). Additional testing may be necessary to confirm the EPEC characterization of the isolate.

The *esth* gene, which represents the variant ETEC isolated from humans (**Nataro and Kaper, 1998**), was found in one of 30 isolates (3%) from Baixada Santista. This detection in the offshore location of the ocean could be due to the release of minimally treated human sewage (**Companhia de Tecnologia de Saneamento Ambiental**

(CETESB), 2006) through submarine outfall from the high volume of sewage during the summer vacation season.

The *stx1* gene that identifies EHEC was found in one of 49 isolates (2%) from São Sebastião Channel. The detection of *stx1* is rare among environmental *E. coli* isolates. It was not found in the Avalon study **(Hamilton et al., 2010)** where nearly 25,000 *E. coli* isolates were screened, but it was found in a transecting study along a freshwater to estuarine water salinity gradient using a culture enrichment study **(Masters et al., 2011)**. The detection of this virulent human pathogen in recreational water of São Sebastião Channel provides clear evidence for heavy human sewage contamination in the area, especially taking into the consideration that water samples were taken offshore. The concentration of *E. coli* and associated virulence genes are possibly even higher at these beaches due to the lower dilution effect from the ocean water compounded with the influence from both submarine sewage outfalls and diffused non-point source of raw human sewage discharge since 40% of the raw sewage was not collected. This result raises health concerns from exposure to fresh human sewage during water recreation at these beaches.

The São Sebastião Channel is considered a mesotrophic marine area due to the release of human waste from four outfalls, including three outfalls (Araçá, Saco da Capela and Cigarras) for domestic sewage discharge and one (Tebar) for industrial discharge **(Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2007)**. The heavy influence of human sewage in the region is reflected in this study by the relatively higher frequency of detection and greater diversity of virulence-associated genes of diarrheagenic *E. coli* (*stx1*, *eae*, *estp* and *astA*).

The heat-stable enterotoxin-1 *astA* gene is the most frequently detected among *E. coli* isolates from all three regions (up to 47%). The role that atypical EPEC strains harboring the *astA* gene play in causing human disease is unclear. The high frequency detection of this gene reflects its prevalence among different categories of diarrheagenic *E. coli*. Previous studies showed 20.7 to 41% ETEC harbor *astA* gene,

and it was identified in 88% EHEC, 22% EPEC and 13% DAEC (**Ménard and Dubreuil, 2002; Savarino et al., 1993**). The gene was also present in 16.6% of *E. coli* strains without other known virulence genes (**de Sousa and Dubreuil, 2001**).

Health risk associated with diarrheagenic E. coli.

The presence of diarrheagenic *E. coli* in sewage-polluted beach water is well justified for health concerns. However, our QMRA results showed that the presence of diarrheagenic *E. coli* in the waters of Baixada Santista and São Sebastião were not significant health threats to beachgoers (specifically bathers) recreating in the regions. This observation holds even for a hypothetical beach location with *E. coli* concentration that reaches the benchmark value (800 CFU/100 mL) of Conama Resolution n° 274/2000. The predicted illness risks for the beachgoers were at least one-order of magnitude lower than U.S. EPA's recommended daily recreational benchmark risk of 32 or 36 NGI per 1000 bathers.

It is important to recognize that these results are not indicative of the overall safety of beachgoers as diarrheagenic *E. coli* only represents one of the many pathogens present in the waters. In fact, previous QMRA studies assessing the human health risks associated with recreating in waters affected by human fecal inputs often accounted for a wide range of pathogens with enteric viruses attributing to a major fraction of the recreational risks (**Soller et al., 2010a; Soller et al., 2010b**). Based on assumptions of pathogen concentration measured in human sewage or projections from the patterns of waterborne diseases in the U.S., Soller et al., (**2010a**) estimated that the risk attributed to diarrheagenic *E. coli* in sewage-polluted water ranged between 10^{-5} and 10^{-6} , which accounts for 0.04% of the total waterborne disease risk. Interestingly, these previous diarrheagenic *E. coli* risk estimates are within the same order-of-magnitude of our QMRA results, which are based on the real diarrheagenic *E. coli* data collected from Brazilian recreational beaches. This agreement provides a degree of confidence for converting the risk of diarrheagenic *E. coli* to the total risk.

Using a reverse approach employed by Soller et al. (2010a), a simple calculation to convert the median illness risk of diarrheagenic *E. coli* to total illness risks can be made by dividing the range of diarrheagenic *E. coli* median risk value by 0.04%

($\frac{2.7 \times 10^{-6}}{0.0004}$ and $\frac{4 \times 10^{-5}}{0.0004}$). The results indicate a median total illness risk range of 7 to 100

illness cases per 1000 bathers across the Baixada Santista and São Sebastião regions. The high-end value of this estimate will exceed the U.S. E.P.A. benchmark by nearly three folds.

The main caveat of the above extrapolation is that all *E. coli* studies used by Soller et al. (2010a,b) were taken place in freshwater environment, which may not be applicable to marine beaches. *E. coli* is known to decay faster in marine water than in freshwater due to its low tolerance to osmotic shock (**Gourmelon et al., 1997**). For the same reason, *E. coli* is not recommended by the U.S. EPA for marine beach water quality monitoring; enterococci is used instead, which has been shown to be more resistant to osmotic shock. The absence of *E. coli* in marine water does not necessarily indicate the absence of enteric viruses and protozoa, which have a better survival rate than *E. coli*. Based on this knowledge, the risk specific to diarrheagenic *E. coli* represents an even smaller portion of the total recreational risk in marine beaches than that in freshwater beaches.

QMRA, although a powerful tool for estimating recreational health risk at beaches in the absence of a large scale epidemiological study, is limited by the understanding of all the disease-causing agents in the recreational water. There are hundreds of different viruses, bacteria and protozoa associated with discharge of sewage to recreational waters that are potentially pathogenic to human. It is impossible to identify and quantify all or even a reliable fraction of them. The study presented here uses diarrheagenic *E. coli* as a case study to illustrate health risk specific to a group of pathogens. This approach provides insight to interpretation of the results between indicators and pathogens and to provoke new thinking in recreational health risk management. Regardless of the level of the risk attributed by this specific pathogen, the detection of

virulent *E. coli* strains in marine water is a clear indication of sewage contamination and the likely presence of other human pathogens.

Environmental and economic impacts

In addition to the human health concern of detecting virulent *E. coli* of human sewage origin in marine waters, the results also alerts the possible environmental damages due to the discharge of nutrients and other pollutants carried by partially treated and untreated sewage. These damages may include eutrophication, reduced biodiversity, and imbalance in the number of indigenous organisms (**Epstein et al., 1993**). According to Tommasi (**Tommasi, 1979**), the urban sewage is the main culprit of ecosystem degradation in coastal waters in Brazil due to inefficient treatment plants and inadequate sanitation infrastructure. The results of this study confirmed the significant sewage influence in marine environment, which guides the environmental management decisions for recreational health, marine biodiversity, and socio-economic practices.

Acknowledgements

This research was partially supported by a grant from CNPq and FAPESP. The travel support for V. F.V. Rodrigues (formerly Da Silva) was provided by Santander Bank fellowship. Graduate Fellowship support to K.Y. Lim was provided by the U.S. National Science Foundation Partnerships for International Research and Education (OISE-1243543).

This manuscript is dedicated to Dr. Irma N. G. Rivera who had devoted her life to research until the last minute.

References

- Abdelzaher, A.M., Wright, M.E., Ortega, C., Solo-Gabriele, H.M., Miller, G., Elmir, S., Newman, X., Shih, P., Bonilla, J.A., Bonilla, T.D., 2010. Presence of pathogens and indicator microbes at a non-point source subtropical recreational marine beach. *Applied and environmental microbiology* 76, 724-732.
- Abessa, D.M., Carr, R.S., Rachid, B.R., Sousa, E.C., Hortelani, M.A., Sarkis, J.E., 2005. Influence of a Brazilian sewage outfall on the toxicity and contamination of adjacent sediments. *Marine Pollution Bulletin* 50, 875-885.
- Albertini, L.S., Ecologia, fatores associados à virulência e diversidade de *Escherichia coli* isolados de amostras de água de lastro, água de regiões portuárias e moluscos bivalves no Brasil. Universidade de São Paulo.
- Almeida, N.C., Barros, A.L.R., Arouche, S.P., Ferro, T.A.F., Moraes, F.H.R., Neto, V.M., Figueiredo, P.d.M.S., 2012. Detecção de enteropatógenos e teste de susceptibilidade a agentes sanitizantes de cepas diarreio gênicas de *Escherichia coli* isoladas das praias de São Luis-Maranhão. *Revista de Patologia Tropical* 41.
- Boehm, A.B., Fuhrman, J.A., Mrše, R.D., Grant, S.B., 2003. Tiered approach for identification of a human fecal pollution source at a recreational beach: case study at Avalon Bay, Catalina Island, California. *Environmental science & technology* 37, 673-680.
- Bonamano, S., Madonia, A., Borsellino, C., Stefani, C., Caruso, G., De Pasquale, F., Piermattei, V., Zappala, G., Marcelli, M., 2015. Modeling the dispersion of viable and total *Escherichia coli* cells in the artificial semi-enclosed bathing area of Santa Marinella (Latium, Italy). *Marine Pollution Bulletin* 95, 141-154.
- Braga, E.S., Bonetti, C.V., Burone, L., Bonetti Filho, J., 2000. Eutrophication and bacterial pollution caused by industrial and domestic wastes at the Baixada Santista Estuarine System—Brazil. *Marine Pollution Bulletin* 40, 165-173.
- Burbano-Rosero, E., Ueda-Ito, M., Kisielius, J., Nagasse-Sugahara, T., Almeida, B., Souza, C., Markman, C., Martins, G., Albertini, L., Rivera, I., 2011. Diversity of somatic coliphages in coastal regions with different levels of anthropogenic activity in Sao Paulo State, Brazil. *Applied and environmental microbiology* 77, 4208-4216.
- Centers for Disease Control and Prevention, Healthy Swimming/Recreational Water: Recreational Water Illnesses (RWIs). Available at: <http://www.cdc.gov/healthywater/swimming/rwi/> (Accessed on June 10, 2015)
- Chan, S.N., Thoe, W., Lee, J.H.W., 2013. Real-time forecasting of Hong Kong beach water quality by 3D deterministic model. *Water Research* 47, 1631-1647.
- Colford, J.M., Schiff, K.C., Griffith, J.F., Yau, V., Arnold, B.F., Wright, C.C., Gruber, J.S., Wade, T.J., Burns, S., Hayes, J., 2012. Using rapid indicators for *Enterococcus* to assess the risk of illness after exposure to urban runoff contaminated marine water. *Water Research* 46, 2176-2186.
- Colford Jr, J.M., Wade, T.J., Schiff, K.C., Wright, C.C., Griffith, J.G., Sandhu, S.K., Burns, S., Hayes, J., Sobsey, M., Lovelace, G., 2007. Water quality indicators and the risk of illness in non-point source impacted recreational waters. *Epidemiology* 18, 27-35.
- Companhia de Tecnologia de Saneamento Ambiental (CETESB), Emissários submarinos. Available at: <http://water.epa.gov/polwaste/nps/index.cfm> (Accessed on May 28, 2015)

- Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2005. Relatório de qualidade daa águas litorâneas do Estado de São Paulo–Balneabilidade das Praias–2005. 254.
- Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2006. Relatório de qualidade daa águas litorâneas do Estado de São Paulo–Balneabilidade das Praias–2006. 365.
- Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2007. Relatório de qualidade daa águas litorâneas do Estado de São Paulo–Balneabilidade das Praias–2007. 294.
- Companhia de Tecnologia de Saneamento Ambiental (CETESB), 2014. Relatório de qualidade daa águas litorâneas do Estado de São Paulo–Balneabilidade das Praias–2014. 224.
- Conselho nacional do meio ambiente (CONAMA), 2008. Resolução no. 274. 29 November 2000.
- de Sousa, C.P., Dubreuil, J.D., 2001. Distribution and expression of the *astA* gene (EAST1 toxin) in *Escherichia coli* and *Salmonella*. *International journal of medical microbiology* 291, 15-20.
- Dickinson, G., Lim, K.-Y., Jiang, S.C., 2013. Quantitative microbial risk assessment of pathogenic *Vibrios* in marine recreational waters of Southern California. *Applied and environmental microbiology* 79, 294-302.
- Epstein, P.R., Ford, T.E., Colwell, R.R., 1993. Marine ecosystems. *The Lancet* 342, 1216-1219.
- Finlay, B.B., Falkow, S., 1997. Common themes in microbial pathogenicity revisited. *Microbiology and Molecular Biology Reviews* 61, 136-169.
- Fleisher, J.M., Fleming, L.E., Solo-Gabriele, H.M., Kish, J.K., Sinigalliano, C.D., Plano, L., Elmir, S.M., Wang, J.D., Withum, K., Shibata, T., Gidley, M.L., Abdelzaher, A., He, G., Ortega, C., Zhu, X., Wright, M., Hollenbeck, J., Backer, L.C., 2010. The BEACHES Study: health effects and exposures from non-point source microbial contaminants in subtropical recreational marine waters. *International Journal of Epidemiology* 39, 1291-1298.
- Fujioka, M., Otomo, Y., Ahsan, C.R., 2013. A novel single-step multiplex polymerase chain reaction assay for the detection of diarrheagenic *Escherichia coli*. *Journal of microbiological methods* 92, 289-292.
- Gourmelon, M., Pommepuy, M., Touati, D., Cormier, M., 1997. Survival of *Escherichia coli* exposed to visible light in seawater: analysis of *rpoS*-dependent effects. *Canadian journal of microbiology* 43, 1036-1043.
- Gunzburg, S.T., Tornieporth, N.G., Riley, L.W., 1995. Identification of Enteropathogenic *Escherichia-Coli* by Pcr-Based Detection of the Bundle-Forming Pilus Gene. *J Clin Microbiol* 33, 1375-1377.
- Haas, C.N., 1996. How to average microbial densities to characterize risk. *Water Research* 30, 1036-1038.
- Haas, C.N., Rose, J.B., Gerba, C.P., 1999. Quantitative microbial risk assessment. John Wiley & Sons.
- Haas, C.N., Thayyar-Madabusi, A., Rose, J.B., Gerba, C.P., 2000. Development of a dose-response relationship for *Escherichia coli* O157: H7. *International journal of food microbiology* 56, 153-159.
- Haile, R.W., Witte, J.S., Gold, M., Cressey, R., McGee, C., Millikan, R.C., Glasser, A., Harawa, N., Ervin, C., Harmon, P., 1999. The health effects of swimming in ocean water contaminated by storm drain runoff. *Epidemiology* 10, 355-363.

- Hamilton, M.J., Hadi, A.Z., Griffith, J.F., Ishii, S., Sadowsky, M.J., 2010. Large scale analysis of virulence genes in *Escherichia coli* strains isolated from Avalon Bay, CA. *Water Research* 44, 5463-5473.
- Instituto Trata Brasil, 2014. Ranking do Saneamento. Available at: <http://www.tratabrasil.org.br/datafiles/estudos/ranking/relatorio-completo-2014.pdf> (Accessed on May 28, 2015)
- Kaper, J.B., Hacker, J.r., 1999. Pathogenicity islands and other mobile virulence elements. ASM Press.
- King, L.J., 2013. Combating the Triple Threat: The Need for a One Health Approach. *Microbiology Spectrum* 1, 1-9.
- Koczura, R., Krysiak, N., Taraszewska, A., Mokracka, J., 2015. Coliform bacteria isolated from recreational lakes carry class 1 and class 2 integrons and virulence-associated genes. *Journal of applied microbiology* 119, 594-603.
- Masters, N., Wiegand, A., Ahmed, W., Katouli, M., 2011. *Escherichia coli* virulence genes profile of surface waters as an indicator of water quality. *Water Research* 45, 6321-6333.
- Ménard, L.-P., Dubreuil, J.D., 2002. Enteraggregative *Escherichia coli* heat-stable enterotoxin 1 (EAST1): a new toxin with an old twist. *Critical reviews in microbiology* 28, 43-60.
- Nataro, J.P., Kaper, J.B., 1998. Diarrheagenic *escherichia coli*. *Clinical microbiology reviews* 11, 142-201.
- National Research Council, 1983. Risk Assessment in the Federal Government: Managing the Process. The National Academies Press.
- Rangel, J.M., Sparling, P.H., Crowe, C., Griffin, P.M., Swerdlow, D.L., 2005. Epidemiology of *Escherichia coli* O157: H7 outbreaks, United States, 1982–2002. *Emerg Infect Dis.* 11, 603-609.
- Ratchtrachenchai, O.-A., Subpasu, S., Ito, K., 2012. Investigation on enteraggregative *Escherichia coli* infection by multiplex PCR. *Bulletin of the Department of Medical Sciences* 39, 211-220.
- Rivera, I., Paula, C., Souza, C., Melo, I., Azevedo, J., 2008. *Microbiologia Aquática Marinha. Microbiologia Ambiental*, 610-627.
- Rivera, I.N., Martins, M.T., 1996. Bactérias enteropatogênicas no ambiente aquático. *Rev. ciênc. farm* 17, 115-136.
- Russo, T.A., Johnson, J.R., 2000. Proposal for a new inclusive designation for extraintestinal pathogenic isolates of *Escherichia coli*: ExPEC. *Journal of Infectious Diseases* 181, 1753-1754.
- Savarino, S.J., Fasano, A., Watson, J., Martin, B.M., Levine, M.M., Guandalini, S., Guerry, P., 1993. Enteraggregative *Escherichia coli* heat-stable enterotoxin 1 represents another subfamily of *E. coli* heat-stable toxin. *Proceedings of the National Academy of Sciences* 90, 3093-3097.
- Schets, F.M., Schijven, J.F., de Roda Husman, A.M., 2011. Exposure assessment for swimmers in bathing waters and swimming pools. *Water Research* 45, 2392-2400.
- Selander, R.K., Levin, B.R., 1980. Genetic diversity and structure in *Escherichia coli* populations. *Science* 210, 545-547.
- Soller, J.A., Bartrand, T., Ashbolt, N.J., Ravenscroft, J., Wade, T.J., 2010a. Estimating the primary etiologic agents in recreational freshwaters impacted by human sources of faecal contamination. *Water Research* 44, 4736-4747.

Soller, J.A., Schoen, M.E., Bartrand, T., Ravenscroft, J.E., Ashbolt, N.J., 2010b. Estimated human health risks from exposure to recreational waters impacted by human and non-human sources of faecal contamination. *Water Research* 44, 4674-4691.

Tommasi, L., 1979. Considerações ecológicas sobre o sistema estuarino de Santos (SP). Honorary Professor Thesis. Universidade de São Paulo, Instituto Oceanográfico 2.

Turbow, D.J., Kent, E.E., Jiang, S.C., 2008. Web-based investigation of water associated illness in marine bathers. *Environmental research* 106, 101-109.

U.S. EPA, Polluted runoff: Nonpoint source pollution. Available at: <http://water.epa.gov/polwaste/nps/index.cfm> (Accessed on February 25, 2015)

U.S. EPA, 2012. Recreational Water Quality Criteria, EPA 820-F-12-058. Office of Water, Washington, DC, United States Environmental Protection Agency.

Verma, A., Bolton, F., Fiefield, D., Lamb, P., Woloschin, E., Smith, N., McCann, R., 2007. An outbreak of *E. coli* O157 associated with a swimming pool: an unusual vehicle of transmission. *Epidemiology and infection* 135, 989-992.

Wade, T.J., Calderon, R.L., Brenner, K.P., Sams, E., Beach, M., Haugland, R., Wymer, L., Dufour, A.P., 2008. High sensitivity of children to swimming-associated gastrointestinal illness: results using a rapid assay of recreational water quality. *Epidemiology* 19, 375-383.

Wade, T.J., Sams, E., Brenner, K.P., Haugland, R., Chern, E., Beach, M., Wymer, L., Rankin, C.C., Love, D., Li, Q., 2010. Rapidly measured indicators of recreational water quality and swimming-associated illness at marine beaches: a prospective cohort study. *Environmental Health* 9, 1-14.

WEF and APHA, 2005. Standard methods for the examination of water and wastewater. American Public Health Association (APHA): Washington, DC, USA.

Westrell, T., Schönning, C., Stenström, T.-A., Ashbolt, N., 2004. QMRA (quantitative microbial risk assessment) and HACCP (hazard analysis and critical control points) for management of pathogens in wastewater and sewage sludge treatment and reuse. *Water Science & Technology*, 23-30.

World Health Organization, 1999. Health-based monitoring of recreational waters: The feasibility of a new approach (The'Annapolis protocol'), Protection of the Human Environment Water, Sanitation and Health Series. World Health Organization (WHO). Sustainable Development and Healthy Environments.

• What is the recreational health risk posed by the presence of diarrheagenic *E. coli* in the water of Brazilian beaches?

• Is the recreational water quality criteria of Brazil adequate for the health protection of bathers?