Mesozoic echinoid diversity in Portugal: investigating fossil record quality and environmental constraints on a regional scale

Bruno Claro Pereira^{a,b,c*}; Michael J. Benton^b, Marcello Ruta^d, Octávio Mateus^{b, c}

a. School of Earth Sciences, University of Bristol, Bristol, BS8 1RJ, UK

b. GeoBioTec, Dept. Earth Sciences, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Portugal

c. Museu da Lourinhã, Lourinhã, Portugal

d. School of Life Sciences, Faculty of Science, University of Lincoln, LN6 7TS, UK

Abstract: Several analyses of diversity through geological time use global, synoptic databases, and this practice often makes it difficult to distinguish true signals in changing diversity from regional-scale sampling and/or geological artefacts. Here we investigate how echinoid diversity changed through the Mesozoic of the Lusitanian basin in Portugal based on a comprehensive, revised database, and seek to distinguish biological signal from geological or environmental constraints. The observed diversity pattern is far from having a defined trend, showing many fluctuations that appear to be linked with gaps in the geological record. This study revealed that, independently of the method used, whether correlation tests or model fitting, the diversity signal is not completely explained by the studied sampling proxies. Among the different proxies, marine facies variation in combination with outcrop area best explains the palaeodiversity curve.

Keywords: echinoid; diversity; sampling proxies; fossil record; Portugal

* Corresponding author. Tel: +44 (0) 117 9545400; fax: +44 (0) 117 9545420

E-mail address: glbmcp@bristol.ac.uk (B. C. Pereira)

1. Introduction

A long-standing debate in current palaeobiological research concerns the significance of global diversity patterns retrieved from raw counts of fossils through time (e.g., Raup, 1972, 1976; Sepkoski et al., 1981; Smith, 2007; Benton et al., 2011; Hannisdal and Peters, 2011; McGowan and Smith, 2011; Benton et al., 2013; Brocklehurst and Fröbisch, 2014; Dunhill et al., 2014), and numerous quantitative studies of the quality of the fossil record allow researchers to agree that fossil diversity is biased by many factors.

Many studies have shown a strong correlation between global fossil diversity and numbers of fossil-bearing formations. This strong correlation has been interpreted in three distinct ways: (1) bias or ‘megabias’; covariation between palaeodiversity and formation time series was formerly widely seen as key evidence for intrinsic deficiencies in the fossil and rock record (i.e., Peters and Foote, 2001, 2002; Wang and Dodson, 2006; Fröbisch, 2008; Barrett et al., 2009; Butler et al., 2009; Benson et al., 2010; Benson and Butler, 2011; Butler et al., 2011; Upchurch et al., 2011); (2) common cause; covariation between palaeodiversity and such geological signals as formation count or rock volume, could result from a single driver such as sea level change, prompting some researchers to invoke a common cause model (Peters, 1995; Hannisdal and Peters, 2011); (3) redundancy; palaeodiversity and fossil formation counts could be intimately linked because both metrics accumulate in tandem as a result of the ways in which palaeontologists assemble data through research time (Benton et al., 2011, 2013). In contrast to that strong relationship at global scale, studies performed at regional or local scale may find weak (Benton et al., 2004; Mander and Twitchett, 2008; Benton, 2012; Dunhill et al., 2012, 2013, 2014), or strong (Crampton et al., 2003; Smith and Benson, 2013) relationships between the two. If there truly are megabiases that can be measured by rock record proxies, fine-scale investigation should reveal stronger correlations between alpha and beta diversity and the proxies than time series recorded at global scale,

because statistical noise ought to be minimized, and thus evidence ought to emerge that one phenomenon drives the other.

Echinoderms, and especially echinoids, have a long collecting record and a wide geographical distribution. Their high-magnesium calcite endoskeletons, either associated or disarticulated, can be found in different marine facies, and are widely represented in the fossil record. This might seem surprising because such mineralogy is unstable, but the recrystallization of high-magnesium to low-magnesium calcite during diagenesis does not result in loss or significant transformation of the ossicle structure. Echinodermata are characterized by their calcitic endoskeleton, which is relatively stable during diagenesis. The endoskeleton composition excludes the possibility of different preservation rates among different echinoderm groups. It is assumed that echinoid preservation potential is uniform, because the endoskeleton composition appears to be constant through time. This leads to good preservation of echinoids and easy collection from marine outcrops (Smith, 2007b).

Despite the high potential for preservation of echinoids, taphonomic bias, such as was noted by Smith (2007b), cannot be ruled out. After the death of an echinoderm, the soft tissues decompose in a few days. Without these tissues, the endoskeleton disaggregates and the ossicles are easily dissociated, which are often small and invisible to the naked eye. This might mean that species with larger macroscopic ossicles would be recognized in the field, while smaller ossicles would remain unnoticed.

Here, we choose to study a regionally restricted, but well sampled series of echinoid faunas from Portugal in order to control for certain biases. For example, Jackson and Johnson (2001) highlighted the geographical bias and the poor sampling of high-diversity, low-latitude marine faunas in global datasets. Further, the geological record of the Northern hemisphere is better studied than the Southern. Here, this geographical bias (Smith, 2007a) is avoided by focusing on one restricted area. Also, sampling efficiency issues (Smith, 2007a)

are minimized in this study, because the database does not arise from compilation of the efforts of many investigators each with different enthusiasms and collecting modes, but is based mostly on geological survey collections, which derive from comprehensive sampling by a limited number of collectors, trained to a particular search style, and covering the landscape equally and efficiently.

In the past, diversity analyses have been performed at different taxonomic levels (for example, species, Smith and Benson, 2013; genus, Dunhill et al. 2014a, b; or family level, Jepson and Penney, 2007). The choice of which level to use seems arbitrary and greatly dependent on practical questions relating to the information that the authors could obtain. It is not axiomatic which level is more representative of the true diversity signal. Generally, the aim is to document an accurate species-level signal in order to understand macroevolution, so a family-level diversity plot would be regarded as likely to generalize the data too much to be useful. Palaeontologists have often preferred to compile their palaeodiversity data at generic, rather than species level, for two reasons (e.g. Foote and Sepkoski, 1999; Smith, 2007b; Mayhew et al., 2012; Smith et al., 2012; Dunhill et al., 2014a, b): species-level compendia may show greater volatility, because new species are established more frequently than new genera; and species boundaries are often more debated than generic boundaries in the primary taxonomic literature.

Robeck et al. (2000) demonstrated that a finely subdivided taxonomic grouping provides a more precise sampled diversity analysis than coarser subdivisions. They concluded that the use of finer taxonomic divisions leads to a more precise measurement of the number of species recorded from the rock record than coarser divisions. The sampled diversity does not correspond to the real biological diversity, since is not corrected for sampling problems.

Echinoids clade can be subdivided into two major clades, the regulars and the irregulars. These two groups evolved at different times in the echinoid tree, with irregulars evolving from regular echinoids (Kroh and Smith, 2010). The evolution from one to the other allowed echinoids to occupy both soft sediment and higher energy habitats. In addition, the evolution and radiation from regulars to irregulars occurred during the time frame under study. Therefore, it is possible that the two groups have different diversity patterns in relation to environmental changes, making it worthwhile to investigate the diversity patterns of both.

Smith and Benson (2013) compared echinoid diversity in the Cretaceous of the U.K. against several rock and environmental proxies. Their results retrieved a strong association between the palaeodiversity signal and a combination of outcrop area with marine facies. It would, therefore, be interesting to see whether their key conclusions are applicable to other regional studies of the same group, and the Portuguese echinoid record provides impetus for the present study.

The aim of this study is explore the nature of the Portuguese echinoid fossil record, by investigating the palaeodiversity signal and comparing it with the coeval, and already documented echinoderm record from the U.K. We will explore similarities and differences between the two records; similarities could represent true biological signals, or reflect the preservation conditions that influenced both areas. The marine Mesozoic succession in Portugal is ideal as the basis for a regional-scale case study on echinoderm fossil diversity because it has a well sampled Mesozoic fossil record, with more than 251 taxa represented around a thousand specimens in total, and it lies at the convergence of two oceanic realms (Tethys to the South and the North Atlantic to the West).

2. Materials and Methods

2.1. *Diversity curves*

The diversity dataset comprises echinoid occurrences in the Portuguese fossil record from specimens in public collections. Prior to this analysis, all these occurrences were subjected to thorough alpha-taxonomic revision by B.C.P., based on examination of type specimens and original materials throughout, down to species level whenever possible (Appendix A). Because the dataset results from a taxonomic revision performed by a single person, and the revision has been done recently, this standardizes the species identification criteria and reduces the risk of taxonomic bias, as identified by Smith (2007a).

The analysed specimens comprise tests (complete and their fragments) and spine fragments (articulated and disarticulated). The vast majority of these specimens (around 95% of the observed fossils) are at the Museu Geológico (Lisbon, Portugal), and were almost entirely collected during the geological surveys of the country, comprising specimens from the Lusitanian and Algarve Basins. In addition, other Portuguese museums that have relevant Mesozoic echinoderm specimens, including the Museu Mineralógico e Geológico (Coimbra University, Portugal) and the Museu da Lourinhã (Lourinhã, Portugal) were also surveyed for specimens. The vast majority of the analysed taxa were collected during geological surveys of the Portuguese territory. This type of campaign should ensure equal sampling effort, through the different formations and localities, analogous to other such case studies based on geological survey collections and records from New Zealand (Crampton et al., 2003) and the United Kingdom (Dunhill et al., 2012, 2013, 2014a, b). The argument is that geological survey palaeontologists collect every fossil they can find, even those of very poor quality, and will search every square kilometre of their mapping areas, thus ensuring a geographically evenly distributed sampling effort. In more usual situations, palaeontologists generally do not follow such a search model, but go straight to known, rich localities, and so over-collect where fossils are rich (the so-called “bonanza effect”; Raup, 1977), and under-collect where fossils have not yet been found. Similarly to this study, the diversity data used by Dunhill et

al. (2012, 2013; 2014a, b) was mainly based on information gathered by the British Geological Survey, with few additions from other sources. The correlation tests between diversity and sampling metrics on those studies could support either bonanza or bias effects. But, Dunhill et al. (2014b, Fig. 4) information transfer statistical tests indicate a redundancy between the two metrics.

The specimens analysed in this work came from the Lusitanian (West of the Iberian Peninsula) and Algarve basins (Southwest of the Iberian Peninsula), corresponding to subsiding basins beside two passive continental margins (Fig. 1). The two basins are infilled with Mesozoic and Cenozoic sediments and their formation is linked with the opening of the Atlantic Ocean to the west, and the Tethys Ocean to the south. The Portuguese Mesozoic echinoid fauna comprises 251 species, after the mentioned taxonomic revision. This taxonomic revision resulted in the addition of 5 new taxa. From the 251 Portuguese taxa verified by the taxonomical revision, 197 species possessed tests and were found on the Lusitanian basin.

Many of those species were described for the first time in Portugal. In the majority of cases, the observed fossils belong to old collections (late 19th to early 20th century), making it difficult to establish a precise age distribution in many cases. The fossil ages provided by old publications often require revision, providing ages at stage level and, therefore, poor age range precision. For the diversity analysis and when a more restricted range was required, it was considered that the taxa in question would occupy the entire stage/substage interval. In total, 243 species were collected in the Lusitanian basin, 13 species in the Algarve basin, and 5 species appear in both basins. These numbers demonstrate a quite significant difference in diversity signal from both basins, with only about five per cent of species in the Algarve Basin.

The differences in the numbers of species between the two basins are significant. The Algarve basin has been historically less sampled than the Lusitanian basin, with many of the geological surveys and subsequent works focusing on the former basin. This fact can explain the difference between the palaeodiversity numbers of the two basins, in association with possible longer hiatus periods experienced by the Algarve basin. For the sake of providing results that are as accurate as possible and minimizing possible collecting bias, we will consider only diversity data from the Lusitanian basin.

The Mesozoic sediments in the Lusitanian basin range in age from Induan (Early Triassic) to Maastrichtian (Late Cretaceous), while the echinoid occurrences span from Sinemurian (Early Jurassic) to Maastrichtian and provide the most complete echinoderm fossil record in Portugal. The specimen ages were revised, when possible, with the aid of the latest local stratigraphic information and long-term, independent dating of the relevant marine Portuguese formations (Dinis et al., 2008; Kullberg et al., 2013). The stage and substage time intervals used in the dataset follow the boundaries in Gradstein et al. (2012), except in some additional subsequent revisions to substage boundaries, which are cited separately below.

In the past, it was common practice to erect echinoid species based exclusively on isolated spine morphology. However, recent studies noted the potential for high morphological variability of this character in individual specimens, especially in cidaroids. Depending on its test position, the spine can be long or short, but can also possess different overall shape and ornamentation. This means that a single echinoid species may possess different spine morphotypes, and this has resulted in the establishment of multiple synonymous species, many based on isolated spines. On the other hand, some echinoid groups show low spine plasticity, in which the same spine morphotype may be present in different species. Therefore, species identified only on spine morphology are not reliable and

are excluded in the current study, to avoid counting the same species twice. By focusing only on the Lusitanian basin, and excluding taxa identified solely from spine morphology, the final dataset used for the analyses consists of 195 echinoid species.

2.1.1. *The “Lazarus effect”*

Whereas taxonomic level does not much affect the palaeodiversity curves for sampled-in-range data, the effect is quite different for range-through plots. In these cases, missing data, especially “Lazarus taxa” (i.e. disappearance and reappearance of taxon in the fossil record), are incorporated, and the relative frequency and sizes of gaps can vary depending on completeness of documentation and taxonomic level (Fara, 2001). Such corrections can reduce some gaps in the fossil record, for example, gaps resulting from basin discontinuities, like those that will be noted in our data. To correct for this effect, the generic dataset was checked for possible gaps in each genus range. Whenever this occurred, those gaps were filled. Both the corrected and uncorrected datasets were assessed for correlation and time series models.

2.1.2. *Collecting effort*

A collecting effort curve, or discovery curve, has often been used as a means of assessing the rate of species description and the trustworthiness of the dataset. The objective is to investigate whether the number of species within defined parameters (taxonomic, geographic, temporal) is reaching saturation, or if many species still remain to be discovered from accessible outcrops. When the level of saturation is approached, the plot of species accumulation against research time overall generally is a sigmoid curve, i.e. initial low rates of discoveries, followed by a rapid increase in the number of species added, and then a new slow rate. This new slow rate tends towards an asymptote, which represents a hypothetical

maximum record of species for the defined sampling universe (Benton et al., 2011), and is limited by possible bias in the rock record.

The collecting effort curve for the Lusitanian basin echinoids was built through the compilation of the number of species against the year they were described based on numerous publications (Forbes, in Sharpe, 1850; Lorient, in Choffat, 1880; Choffat, 1880; Lorient, 1884; Lorient, 1887-1888; Lorient, 1890-1891; Lorient, 1900; Lorient, 1905; Lambert, 1913; Lambert, 1915 to 1916; Rey, 1972; Vadet and Willie, 2002). The new species that resulted from B.C.P.'s taxonomic revision are also included in this curve. The collecting curve includes only those species that are currently regarded as valid, not the sum total of all species as described.

2.2. *Sampling proxies*

2.2.1. *Formations*

Previous authors have used a variety of parameters as sampling proxies to explain the diversity signal. Fossiliferous formation numbers has been widely used as a sampling proxy (Peters and Foote, 2001, 2002; Fröbisch, 2008; Barrett et al., 2009; Butler et al., 2009, 2011; Benson et al., 2010; Benson and Butler, 2011; Mannion et al., 2011; Upchurch et al., 2011; Benson and Mannion, 2012; Lloyd, 2012; Benson and Upchurch, 2013). This metric has been proposed as a good sampling proxy since it reflects multiple sampling controls, including rock area, thickness, lithological variation, and some aspects of human effort (Peters and Foote, 2001). Therefore, it should also provide a summary of amount of record (Raup, 1976), research effort (Sheehan, 1977) and lithological variability.

However, others argue that fossiliferous formation count is a poor sampling proxy (Crampton et al., 2003; Wignall and Benton, 1999; Peters, 2006; Smith 2007; Peters and Heim, 2010; Benton et al., 2011, 2013; Dunhill, 2011, 2012; Dunhill et al., 2014b). These

authors present five reasons for their critique: 1) definitions of formations are arbitrary and they vary significantly with stratigraphic age, geography and environment; 2) formations tend not to correlate with rock volume, area of exposure or collecting effort; 3) they are largely dependent on facies heterogeneity; 4) they are partially redundant with palaeodiversity, both measures reflecting rock heterogeneity; 5) their volume varies at least over eight orders of magnitude.

We compiled two formations datasets for the Mesozoic of the Lusitanian Basin (based on Dinis et al., 2008; Kullberg et al., 2013), one including all formations, and the other the purely marine formations (Appendix A). The inclusion of terrestrial and transitional facies may seem counterintuitive, as echinoids are exclusively marine animals, but many of the Portuguese Mesozoic formations contain both marine and continental facies, and we considered it best to include all possible settings for echinoids. In any case, we offer results based both on the full formations dataset, and the strictly marine formations dataset.

2.2.2. *Facies heterogeneity*

Habitat heterogeneity has been suggested as another factor influencing the diversity signal obtained from the fossil record. Each type of habitat results in the production and accumulation of different types of sediments, and the organisms living there are adapted to those conditions. Theoretically, a greater diversity of habitats may imply a higher number of species, leading to a higher diversity overall. Crampton et al. (2006) suggested that the number of formations might be a good estimator of this rock facies heterogeneity.

In this study, we use variation in sedimentary facies as an indicator of numbers of habitats. To calculate this indicator, the description of each formation was examined (described in Dinis et al. 2008; Kullberg et al. 2013) to establish which facies were more

dominant. The facies terms used were: limestones and dolomites; marls; sandstones; mudstones; conglomerates and evaporites.

A subdataset with just marine facies was also created, excluding conglomerates, evaporites and parts of the sandy facies. This subdataset is similar to the facies division used by Smith and Benson (2013). All the truly marine formations yield macrofossils, implying that the marine facies curve also corresponds to a fossiliferous facies curve. The sum of the number of each facies through time is used as an indicator of facies variability and, thus, an indicator of habitat variability.

2.2.3. *Outcrop area*

Another proxy used in this study is the outcrop area, or map area, corresponding roughly to the area available for retrieving fossils. As noted before (Dunhill, 2011, 2012), there can be a substantial difference between areas of outcrop and exposure: exposure is strictly the area of rock available for study at the surface, whereas outcrop always represents a larger area on the map, including areas of non-exposure where the rocks are covered by soil, water, and human developments (roads, cities). In our case, we could not obtain data on exposure, but at least the outcrop areas are assessed within a limited geographic area in which the degree of concealment (i.e. the difference between outcrop and exposure) for each formation is roughly equivalent.

Previous studies have tested the relationship between diversity and outcrop area, with some favouring (Raup, 1972; Raup, 1976; Peters and Foote, 2001, 2002; Smith, 2001; Crampton et al., 2003; Smith 2007a; Smith and McGowan, 2007; Smith et al., 2012; Smith and Benson, 2013) and others discarding this relationship (Benton et al., 2011; Dunhill, 2011, 2012; Dunhill et al., 2012, 2013, 2014). One explanation for the covariation of palaeodiversity and outcrop areas was that outcrop area is a metric of sampling, and rises and

falls in both time series could simply record collecting opportunity rather than any real diversity signal (Smith 2001, 2007a; Crampton et al., 2003). Alternatively, the covarying signals might reflect the common cause model (Peters, 2005; Hannisdal and Peters, 2011), a kind of species-area effect driven by sea level change, especially in the case of shallow marine formations, as here.

To test for covariation of palaeodiversity and outcrop area signals, the method of Smith and McGowan (2007) and Smith and Benson (2013) was used here. Their method consists in using the distribution grid of geological maps, together with the information from the map memoirs, to retrieve a count of how many grid units contain each fossiliferous formation. This method provides an even distribution grid through the area of study and provides even estimates of the formation area. These areas were then organized in sequence by time bins. Smith and McGowan (2007) tested this method at different map scales and concluded that the curve obtained had the same pattern, regardless of which map scale was used, just changing the amplitude of the peaks.

The Portuguese Geological Survey (Laboratório Nacional de Energia e Geologia, LNEG) produced geological maps at the scales 1:50000, 1:200000, 1:500000 and 1:1000000. Unfortunately, both the Lusitanian and the Algarve basins are not fully mapped at the 1:50000 and 1:200000 scales. For this reason, geological maps at 1:500000 and 1:1000000 were used, but subdividing these maps with the grid of maps of lower scale. This could be achieved by using an on-line tool called Geoportal (<http://geoportal.lneg.pt/geoportal/mapas/index.html>), provided by LNEG, where the geological maps can be interactively visualized at different scales and grids superimposed from other scales. At the usable scales mentioned above, the website provides coarse information, not providing information for each individual formation. Instead, it provides groups of units and their age ranges.

At the scale 1:1000000, the website provides groups of units, corresponding to coarse stratigraphic packages, and their age ranges at series level (Appendix A). On the other hand, maps at the scale 1:500000 give a little more information. At this scale, the units comprise a mixture of formal and informal stratigraphic units, some of them correlated with established formations, again with a range at series level. To improve the precision of the age range of each unit, the unit's descriptions were compared with the works of Dinis et al. (2008) and Kullberg et al. (2013) and, where possible, a more precise age range for each group of sedimentary packages was attributed. For this study, only data from sedimentary units were extracted, excluding the volcanic deposits from the end of the Cretaceous, which lack of macrofossils. With the units established and their age ranges constrained, the distribution of each formation at both scales was extracted. For each map scale in use, the grid of the 1:50000 geological maps was superimposed and the number of grid units in which each sedimentary group appeared was counted (Appendix A). At the scales used and the information provided by the used tool, it was not possible determine the area for each individual formation, despite all efforts.

2.2.4. *Palaeotemperature*

Echinoids can provide important information about biotic and climatic evolution. Therefore, it is reasonable to compare the evolution of this group with climate parameters, like temperature. It has been suggested that temperature controls echinoid diversity, with cooling periods decreasing their global diversity (McKinney et al., 1992). Carbon and oxygen stable isotopes have been used as proxies for palaeoclimate signal (e.g. Voigt et al., 2003; Carpentier et al., 2006; Martin-Garin et al. 2010; Suan et al., 2010; Price et al., 2013), with both isotopic signatures providing information regarding warming or cooling periods. For this study, the isotope dataset was compiled from different publications (Saito and Donk,

1974; Price and Sellwood, 1994; Picard et al., 1998; Veizer et al., 1999; Voigt, 2000; Cresta et al. 2001; Wierzbowski 2002; Voigt et al., 2003, 2004; Carpentier et al., 2006; Fursich et al., 2005; Aguirre-Urreta et al., 2008; Price and Teece, 2010; Suan et al., 2010; Huber et al., 2011; Lazar et al., 2011; Alberti et al., 2012; Armendáriz et al., 2013; Price et al., 2013), comprising a total of 1341 samples from different localities with a worldwide distribution, some of which are Portuguese localities. The isotopic signatures used were extracted from Mesozoic brachiopods, oysters and foraminifera, with no evidence for diagenetic alteration (Appendix A). The compilation of the dataset follows the criteria used by Price et al. (2013) and, therefore, information on wood, belemnite and bulk rock isotopes was discarded. The same authors preferred not to adjust the $\delta^{18}\text{O}$ values for secular variation, due to inconsistencies in the use of the ‘carbonate clumped isotope’ method, and preferred to use this proxy as an indicator of temperature changes, in which more negative isotope ratios indicate seawater warming. Bearing this in mind, a compilation of isotope ratios is here used as a proxy for temperature changes.

2.2.5. *Sea level*

Another factor normally stated to influence diversity is sea level change. This proxy has long been recognized as a potential driver of diversity changes, especially linking sea level falls with extinctions (Chamberlin, 1909). Fluctuations in global sea level are a consequence of changes in the ocean water or oceanic basin volumes, linked with tectonic events and temperature changes. Later, Newell (1967) suggested a correlation between periods of loss of epicontinental seas during sea-level lowstands, and major palaeontological turnovers.

Further, with sea level rise, flooding of cratons occurs, increasing the area of shallow marine shelf available for organisms to occupy. This increase of shallow waters also

increases the area for marine sediment deposition, resulting in an increase in the rock record. On the other hand, regressions reduce the habitat area and the volume and surface area of previously deposited sediments, due to erosion. This coupling of shelf rock and fossil records can be interpreted either as evidence for a ‘common cause’ model in which both rock volume and fossil diversity rise and fall in concert because both are driven by sea level rises and falls (Peters, 2005). Alternatively, the very changes in sea level can be seen as controlling fossil preservation by erosion during transgressions (Smith, 2007b). McGowan and Smith (2008) found from cross-correlation analyses that there is some evidence for intercontinental correlations, apparently linked to long-term sea-level fluctuations, at first and, occasionally, second-order cycles.

In the absence of a comprehensive local sea level curve, the Portuguese Mesozoic echinoid palaeodiversity was compared with two global sea level curves (Haq et al., 1987; Miller et al., 2005) and a western European sea level curve (Graciansky et al., 1998). The raw data for the global curves was provided by Miller et al. (2005). The raw data for the Graciansky et al. (1998) sea level curve could not be obtained.

Because the raw data for this last curve are not provided, the Grabit tool in the Matlab program (Doke, 2007) was used to extract point values from the figure provided by Graciansky et al. (1998, Appendix Chart 1) and extracted as many points as needed to replicate the original curve. The output of the program is a file with the x and y coordinates of each point (see Appendix A).

2.3. Generating the diversity, formations, facies and outcrop curves

The diversity, formations, facies and outcrop values through time were calculated and plotted in R v.2.14.1 (R Development Team, 2011) using a script created by B.C.P. (see Appendix B), using part of Graeme Lloyd’s ‘rt_diversity_ranges.r’ script. This script divides

the time frame into equal time bins and searches through the dataset for taxa that occur within each time bin. For our analysis we used a time bin bandwidth of 1 Myr.

Currently, there are two commonly used methods to address the issue of plotting diversity through time, the use of time bins or the use of arbitrary bins, like stages or substages. The first method is highly dependent on the precision of the dataset, but tries to be a faithful representation of reality. The second method overestimates diversity by stating, for example, that a certain species that could have lived for 5 million years, actually spanned just 2 million years. Also, the use of stages, or substages, implies extra mathematical corrections, for addressing the uneven time distribution of those categories. With these differences in mind, the time bin projection was chosen, to try to obtain the highest precision possible.

2.4. *Pairwising the datasets*

The oxygen isotope, carbon isotope and sea level datasets did not possess the same time bin distribution as our diversity dataset. The statistical methods that we used can only be performed if all datasets are distributed in the same time bins (paired). For the isotopic datasets, the function ‘ksmooth’ in R was used to create a smooth line based on the full set of irregularly spaced isotopic values. For this curve, we used a bandwidth of 10 Myr.

Then, functions based on the isotopic and sea level curves were created, through the ‘splinefun’ function in R, using the method ‘monoH.FC’. After the different functions had been created, the values for each diversity time bin midpoint were extracted, and these were used for time series and correlation comparisons with the diversity values for each time bin (Appendix B).

2.5. *Generalized differencing*

The palaeodiversity signal and tested sampling metrics were initially considered as raw values and then subjected to generalized differencing. This manipulation of the data aims to remove general background patterns ('detrending'), by identifying any trends in the data, through a linear model, and the differences from it. This data transformation was carried out in R, using Graeme Lloyd's script 'functions_2.r' (Lloyd, 2008) together with an automated R code, written by B.C.P. (Appendix B). This data manipulation was previously used by, for example, Benson and Butler (2011), Benton et al. (2013) and Smith and Benson (2013).

2.6. *Pairwise correlation analyses*

Spearman rank correlation tests were performed to explore covariation between the palaeodiversity data and the different sampling and environmental proxies. The strength of such correlations indicates the extent to which each variable might explain part of the diversity signal variance.

Then, to circumvent the possibility of redundancy between sampling proxies, correlation tests between the proxies were performed. If the correlation was high between two proxies, that would imply that the proxies were based on the same signal and, in consequence, one of them should be excluded. All correlation tests were also performed in the program R, using a script developed by B.C.P. (Appendix B). This script uses the R function 'cor.test' to perform the correlation analyses, with the Spearman correlation method, and also corrects the results for false discovery rate. For this correction, the more strict Bonferroni method was used.

2.7. *Generalized least squares model fitting*

Correlation tests can only compare two variables at a time, for example, palaeodiversity with one sampling proxy or environmental driver. However, a modelling

technique can investigate the explanatory power of each variable independently and the possibility of combinations. Such modelling approaches were previously used to assess palaeontological data (e.g. Marx and Uhen, 2010; Benson and Butler, 2011; Butler et al., 2011; Benson and Mannion, 2012; Benton et al., 2011, 2013; Smith and Benson, 2013). This study uses the generalized least squares (gls) model, based on the assumption that the values of the response variable are dependent, and common to ecological data across space and time. This method is commonly used to model correlated observations, which was assumed in the case of our data and proxies.

For this method, the 'gls' function in the 'nlme' library (Pinheiro et al., 2004) in the R program was used, on the raw data measures. To use this function, an R script was written by B.C.P. (Appendix B). Initially the routine was run with all putative explanatory variables together and, then variables were gradually removed and different combinations were tested. For the initial analyses, tests to determine which best autoregressive order values (p) in corARMA were ideal for our study had to be performed. These tests used the diversity data at species level, $p = 1, 2, 3, 4, 7, 8$ and 20 , in a model that considered several proxies under study (total formations, marine formations, oxygen isotopes, carbon isotopes and a global sea level). They were performed separately for diversity based on species, genera and genera corrected for Lazarus effect curves, and used the sea level curve based on the Haq et al. (1987) data.

As expected, some of the variables explain the variance in the response variable better than others, as measured by Akaike Information Criterion (AIC) values. According to Burnham and Anderson (2002), the Akaike Information Criterion corrected (AICc) values should be used instead of the AIC, when the number of observations divided by the number of estimated parameters is lower than 40. For our study, this fraction is definitively higher than 40. Nevertheless, the AICc values were also extracted, using the function 'AICc.gls' in

the R library ‘AICcmodavg’ (Mazerolle, 2012). Lower AIC or AICc values can be interpreted as indicating better models.

3. Results

3.1. *The collecting curve*

The collecting curve of the Lusitanian basin (Fig. 2) shows that the rate of discovery of echinoids appears to have reached a steady state. Accumulation of new taxa shows significant growth during the late 1890s and early 1900s, especially by the work of Perceval de Loriol (1828 to 1908). After this author, few articles were published, but these have added only a few new species and genera. This pattern of the curve suggests it is unlikely that significant numbers of new echinoid species will be found in the future, and that there can be some confidence in the dataset and conclusions extracted from it.

3.2. *Palaeodiversity through time*

The comparison of the diversity curves at species, genus and family level shows that the pattern of the echinoid diversity curve is the same, independent of taxonomic level (Fig. 3A). The curves differ in the amplitude of peaks and in a slight smoothing of the curve at higher taxonomic levels. Plots of species through time changed relatively little with taxonomic revision, whereas the generic plots changed significantly. Changes in the generic assignments of many of the species arise because the old taxonomic assignments in the late 19th century have not subsequently been revised.

In the Lusitanian basin, the number of taxa varied significantly during the Mesozoic (Fig. 3A). The palaeodiversity curve is far from having a general trend, instead being characterized by peaks of diversification and drops of apparent extinctions. Many of these drops in diversity coincide with well-established basin discontinuities (Fig. 3B), and seem to

indicate some sort of cause and effect. The time with most species is the Late Jurassic, standing out quite clearly on the diversity curve, and is preceded by the Callovian to Oxfordian hiatus (Kullberg et al. 2013). Also, the existence of marine formations is not indicative of the presence of echinoids. For example, during a great part of the Pliensbachian and part of the Toarcian there are no records of echinoids, even though a considerable number of marine formations existed during those ages (Fig. 3C).

3.2.1. *Regular vs irregular echinoids*

The palaeodiversity behaviour of echinoids was analysed, as a whole, and as different subclades. Regular and irregular echinoids show rather different palaeodiversity histories through the Mesozoic (Fig. 4A). These changes in the diversity of both groups might be related to changes in sedimentation type in the basin. The Jurassic of the Lusitanian basin is dominated by carbonates and mudstones, associated with a calm and tropical environment, while the Cretaceous was much more siliciclastic, with accumulation of mudstones, sandstones and marls. This change in the type of sedimentation could have led to environmental adaptation of the echinoids, inducing the diversification of endobenthonic forms, which are especially characteristic of the irregular echinoids.

3.2.2. *Corrected palaeodiversity curves*

The imprecision of echinoid classifications based on isolated spines was previously mentioned. Inclusion or exclusion of species based on spines has significant impact on the sampled diversity values. Removal of spine-based species significantly decreases the diversity of several times, especially during the Late Jurassic and Early Cretaceous. The higher impact of this subtraction on Jurassic than on Cretaceous diversity can be explained by the fact that regular echinoids were more abundant than irregular echinoids during the

Jurassic. Regular echinoids have larger and more conspicuous spines, such as the cidaroids, while irregular echinoids have smaller and undifferentiated spines. Nevertheless, the overall diversity pattern remains the same with or without these species (Fig. 4B). Correlation tests (Appendix A) confirm the similarity of the two curves, whether assessed on raw data ($\rho=0.964$, $p<0.001$) or on data after generalized differencing ($\rho=0.832$, $p<0.001$). Therefore, it is preferable to use the dataset without the doubtful taxa, bearing in mind that this curve represents a minimum estimate of diversity.

When range data are calculated by filling gaps, to avoid the Lazarus effect at the species level, there are few changes in the palaeodiversity curve, with the ranges of only three species changed. On the other hand, at the level of genera, there are significant changes (Fig. 4C), presumably because there were many changes to total ranges when Lazarus gaps were filled. In particular, many drops are removed or significantly attenuated, that had previously been demonstrated to be associated with major basin discontinuities. Nevertheless, important and long temporal decreases in diversity are still well marked at the end of the Kimmeridgian and Hauterivian, even after correction. Although the global disappearance of a taxon cannot be undone, the Lazarus effect can be circumvented. The local extinction of a certain taxon can precede recolonization from other habitats.

In our analyses, corrections for the Lazarus effect artificially attenuated many decreases in diversity. However, these corrections did not always produce improvements in correlation, slightly increasing it in some cases, but decreasing it in others (Appendix A). This raises the question of when it is appropriate to correct for Lazarus gaps. Generally, it makes sense to perform such corrections because the objective is to establish the best possible estimate of the overall palaeodiversity. The fact that the uncorrected palaeodiversity time series correlates better with the rock metrics than the corrected palaeodiversity signal

confirms that some at least of the Lazarus gaps arise from inadequacies of sampling, as might be expected.

3.3. *Sampling variables*

3.3.1. *Number of formations*

In a few sections of our diversity plot, drops in diversity seem to be associated with decreases in the number of formations (Fig. 3C). For example, the Hauterivian diversity decrease is linked to a decrease in the number of both marine and total number of formations. The decrease of both diversity and number of formations is associated with the D13 basin discontinuity, which occurred in the Barremian, and corresponds to a regression maximum, which led to the general emersion of the Lusitanian Basin (Kullberg et al. 2013).

There is low to moderate correlation between palaeodiversity and total number of formations ($\rho \approx 0.4$ and $p < 0.001$, for both raw data and after generalized differencing), and with number of marine formations (Table 1; Fig. 3C; Appendix A). For both raw data and after generalized differencing, the correlation between diversity and formations is slightly higher at lower taxonomic levels, even after applying the corrections against the Lazarus effect. After generalized differencing of the data, the correlation values between palaeodiversity and number of formations slightly increase, suggesting moderate values of correlation. On the other hand, the number of marine formations shows the same behaviour, but with considerably lower values (Table 1 and Appendix A). The application of generalized differencing slightly improves this correlation.

3.3.2. *Facies heterogeneity*

As for the diversity signal, the number of all facies and marine facies varies through time. The curve for number of facies per time bin does not have a specific trend, showing peaks and troughs (Fig. 5A). The same pattern is also observed in the marine facies curve.

The Spearman's rho values obtained for comparisons of time series of facies or marine facies and diversity indicate little to moderate correlation (Table 1 and Appendix A), at the levels of species, genera and corrected genera. The correlation values obtained for a simple correlation between diversity and facies were moderate ($\rho \approx 0.6$ and $p < 0.001$), while the same correlation, after generalized differencing, gave systematically lower rho values ($\rho \approx 0.35$ and $p < 0.001$). Further, the correlation between diversity and just marine facies retrieved systematically slightly lower values, for both raw data and after generalized differencing.

3.3.3. *Outcrop area*

The amount of sedimentary rock corresponding to the different time bins is not evenly distributed in the Lusitanian basin: the Cretaceous time bins show generally higher outcrop areas than the Jurassic time bins (Fig. 5B). Correlations between diversity and outcrop area change markedly according to the map scale. The outcrop dataset based on the maps at 1:500000 systematically yielded better correlation values than those based on the 1:1000000 scale (Table 1 and Appendix A). The correlations between diversity and outcrop area based on geological maps at the scale 1:500000 yielded high (raw data, $\rho \approx 0.7$ and $p < 0.001$) and moderate (generalized differenced, $\rho \approx 0.4$ and $p < 0.001$) rho values.

3.3.4. *Diversity and sampling proxies*

Different correlation test results were obtained from the comparison between diversity and the rock record proxies. The correlation between formations and marine formations,

formations and marine formations with facies and marine facies retrieved high rho values, for the raw data and moderate values, after generalized differencing (Table 1 and Appendix A). This shows that these proxies share a common trend and should be tested separately, implying that substantial redundancy exists between some pairs of these proxies. Also, the correlation tests between facies and marine facies with outcrop area (1:50000) were associated with moderate rho values (around 0.30), for both raw data and after generalized differencing. Although a moderate correlation, these values appear to suggest a common factor influencing both proxies, and the results extracted from them should be interpreted carefully.

The rock proxies that showed consistently highest correlation values with diversity (at all taxonomic levels tested) were facies variation, both total number and just marine facies (Table 1 and Appendix A). This moderate correlation remained similar after applying generalized differencing to the data. This implies that, as with the previously mentioned studies, facies heterogeneity might have some control over the biological signal obtained from the geological record, or that the sampling across facies is influencing the diversity signal. Sedimentary facies are highly dependent on environmental factors, which also influence diversity. Higher numbers of habitats produce a higher numbers of facies and, also, hold potential for higher rates of diversification. These higher rates of diversification are linked to the potential for new habitats to be occupied by different species. It was surprising that the total number of facies had a slightly higher correlation with diversity, than did marine facies alone. The reason for this could be that by using just marine facies variation, we are not encompassing the full range of habitats in a particular time period. For example, by focusing on just marine facies, transitional environments that, at some point, could have held echinoids are excluded. Therefore, by just focusing on marine facies we are excluding the full range of habitats.

On the other hand, the correlation values between palaeodiversity and the number of formations and marine formations is considerably less significant, for both raw data and after applying generalized differencing (Table 1 and Appendix A). The results of the Spearman correlation tests are significantly lower than those obtained for the other rock proxies, with the correlation slightly increasing after generalized differencing, at all taxonomic levels tested (Table 1 and Appendix A). The expansion or contraction of marine habitats related with sea level changes would lead to variation in the number of marine formations, due to changes in sedimentation style. The correlation values obtained seem to support this idea.

The raw correlation between diversity and outcrop area yielded different correlation values, according to the map scale used. While for maps at 1:1000000, the correlation values were weak, for the scale 1:500000, the correlation was significantly stronger (Table 1 and Appendix A). On the other hand, after applying generalized differencing, both scales were associated with quite low correlation values. These low correlation values appear to indicate that the diversity signal obtained by fossil counting is not biased by an accessibility factor, but on outcrop alone. If that bias existed, time periods with larger outcrop areas would have yielded more fossils and vice versa, and the correlation between diversity and this proxy would be high. Nevertheless, the influence of sampling area cannot be entirely excluded.

3.4. *Environmental proxies*

3.4.1. *Temperature driver*

From the observation of the two isotopic ratios, it seems that there were no significant long-term temperature changes (longer than 10 Myr, which is the bandwidth for projecting the data), during most of the Mesozoic, since both oxygen and carbon isotopic signatures do not show major fluctuations (Figs. 5C and 6A). Isotopic values indicate that global temperature was significantly warmer than today, since the oxygen isotopic ratios are more

negative and the carbon isotopic ratios are higher than the present day. For example, Myers et al. (2012a, 2012b) show a surface palaeotemperature for the upper Kimmeridgian/lower Tithonian Lourinhã Formation between 27°C and 34°C (average 31°C).

There is very little correlation between palaeodiversity and both isotopic curves (Figs. 5C and 6A; Table 1 and Appendix A), especially after applying generalized differencing ($\rho \approx 0.10$ and normally $p > 0.003$), suggesting that temperature had little influence on echinoid palaeodiversity. It is unclear whether this is an artefact of imprecise data, or an indication that the temperature of seawater does not affect echinoid diversity.

3.4.2. *Sea level driver*

The three sea level curves used in this study (Haq et al., 1987; Graciansky et al., 1998; Miller et al., 2005) show overall trends of increase of sea level through the Mesozoic, but with different orders of magnitude (Fig. 5C), and agree on an apparent overall trend for a stabilization of sea level during the Late Cretaceous. This overall trend of increase in sea level is punctuated by small drops, which vary in magnitude according to the model chosen and do not always coincide between models.

Correlation tests between palaeodiversity and the sea level curves (Table 1 and Appendix A) produced rather low Spearman's ρ values, both for the raw ($\rho \approx 0.2$ and $p < 0.001$) and generalized differenced data ($\rho < 0.1$ and $p < 0.001$). The results vary slightly according to the taxonomic level chosen. For the raw data, the highest correlation values were obtained between species and Miller's curve. Regardless of the taxonomic level chosen, Miller's curve yielded the highest values, while Graciansky's resulted consistently in lowest values. Looking at the generalized differenced correlations, the results were not always statistically significant, yielding significantly lower ρ values than the raw data

comparisons. Also, the results vary according to the taxonomic level chosen, decreasing with higher taxonomic levels.

3.4.3. *Diversity and the environmental proxies*

The low correlation values obtained between diversity and the sea level could result from the fact that none of the curves used truly correspond to the sea level variations, in the Lusitanian Basin. In this basin, sea level was controlled by global/regional sea level changes, but also by local tectonic events, that would have played an important role in determining the area of continental flooding. Also, this low correlation appears to go against the “common-cause” hypothesis. For this hypothesis to be true, we ought to obtain a concrete correlation between sea level and diversity, which is not the case in this study.

As previously enunciated, temperature changes had little influence on palaeodiversity, going against the commonly enunciated that echinoids are a good proxy for inferring palaeotemperature (e.g. McKinney et al., 1992). This result is also opposite to the findings of Mayhew et al. (2008, 2012). These authors demonstrated that temperature correlated with different measures of palaeodiversity at global scale, although in different ways (with highs in palaeodiversity either correlated with lows or highs in palaeotemperature). But, it must be borne in mind that that study was made across the entire Phanerozoic and, by consequence, used a broader and less precise data. Again, the regional-scale records ought to show this correlation even more convincingly, if the relation between temperature and diversity exists. Perhaps the analyses by Mayhew et al. (2008, 2012) provide evidence for correlations because of summing effects of randomly varying regional-scale data sets, and there is in fact no evidence that palaeotemperature drives global palaeodiversity. To sum up, it appears that both environmental proxies analysed had very little influence on the palaeodiversity.

3.5. *Model fitting analysis*

Before the model analyses were performed, the possibility of convergence between proxies was investigated. The use of two variables with the same behaviour would bias the results of the model towards that signal. To circumvent this, correlation tests were undertaken between the different variables. Please, see Appendix A for the results of such tests. When the correlation results between two variables were high, these variables were used in separate models.

From tests performed to investigate the best autoregressive order value (p), the value of p that retrieved the lowest AIC value was 1 (Appendix A). The remaining orders not only resulted in lower AIC values, but also substantially increased the time required for computation. In consequence, all subsequent tests used 1 as the value of p . For tests using $p = 2$ and 3, the program reported an error.

For the diversity data at species level and using all meaningful proxies (marine formations, marine facies, outcrop at 1:1000000 map scale, oxygen and carbon isotopes, Graciansky et al. 1998 sea level curve), it is clear that the only proxies with a meaningful p -value (values of 0) are facies heterogeneity and outcrop area. All other proxies show rather higher p -values (above 0.05), suggesting a lower correlation of these proxies with palaeodiversity. After this initial test, other possible models were assessed in different combinations, by removing and adding proxies. Constant to all models were that the marine facies counts and outcrop areas (at the scale 1:500000) showed highly significant correlations with palaeodiversity ($p < 0.001$). The coefficient values between palaeodiversity and these two proxies are positive (Appendix A), in combination and isolated. This indicates that, when outcrop area and marine facies heterogeneity increase, the number of species retrieved also increases.

The model fitting analyses show a similar result to the correlation tests. In all the models calculated, the isotopic and sea level variations appear to have had very little influence on the diversity signal, with the coefficient values varying according to the proxy. Nevertheless, it is possible to infer that the increase of temperature might have allowed an increase in diversity. The carbon isotopic signature systematically yielded positive coefficients while the oxygen isotopes were constantly negative. Although the two proxies have opposite signals with respect to diversity, they both point to the same behaviour. With the temperature rise, the carbon isotopes tend to be more positive and the oxygen more negative. So, and even if the models indicate that temperature had very little influence on diversity, the rises in temperature seem to be associated with a small rise in diversity.

While the relation between temperature and diversity seems to have some trend, the model fitting analyses for the sea level curves have no particular direction. The coefficient values for sea level were low and with mixed signals, positive in some cases and negative in others (Appendix A). Also, the coefficient values are relatively low, which presumably relates to the uncertainty of the signal.

Different models were also run with the objective of understanding which proxy explained more of the palaeodiversity time series, whether marine facies heterogeneity or outcrop area (at the scale 1:500000). In sequence, the model fits for single proxies with lowest AIC values (Appendix A) are marine facies, then all facies, and then outcrop area. All the other models with the proxies under study resulted in higher AIC values and higher p-values (Appendix A). The same models were calculated with diversity based on genera and genera corrected for gaps in the record. At corrected generic level, the AIC values are smaller than for the same models with diversity at species level (3855.082, tested with marine facies and outcrop area 1:1000000). Other combinations of models at different taxic levels and without corrected palaeodiversity data provided higher AIC values, and so are not included

here. The AICc values were also obtained for several of the models but the differences between the AIC and AICc values are minimal and, therefore, neglected here.

The model fitting analyses (gls) allowed better understanding of the relationship between the rock proxies and palaeodiversity. Facies and marine facies are the proxies that yield the best scores, with highly significant p-values of 0 (Appendix A). The different models show different directionalities between diversity and proxies, according to the proxy in question. The coefficients are systematically positive for facies, marine facies and outcrop at the scale 1:50000. On the other hand, outcrop at the scale 1:1000000 retrieved negative values, while formations and marine formations produced positive and negative coefficient values (Appendix A).

The model that yielded the lowest AIC value included marine facies and outcrop area, at the scale 1:50000 (AIC=5511.596), with both yielding p-values of 0. The coefficient value obtained from this model was higher for the marine facies count than for the outcrop area. The results for the other models are provided in Appendix A.

After the model fitting analyses, the two proxies used in the model with the best results were combined and correlated with palaeodiversity. The combination of marine facies and outcrop area resulted in moderate correlation values with the diversity signal, for the pairwise correlation tests (for the raw data and after generalized differencing) and model fitting. The two proxies combined resulted in better correlation values than for the two proxies isolated, and implies that both proxies in conjunction conditioned the diversity signal. Smith and Benson (2013) obtained a similar result for the Cretaceous echinoids of the U.K. However, and the correlation values obtained by those authors were significantly higher. Possible reasons for this will be addressed below.

The correlation between diversity and the combination of marine facies and outcrop area is significant. Nevertheless, interpretations of these correlations need to be cautious. The

correlation tests between marine facies and outcrop area demonstrate the existence of some relationship between the two proxies, with rho values of 0.286 (raw data) and 0.353 (after generalized differencing), and suggest a common factor influencing both proxies.

3.6. *Diversity, marine facies and outcrop area*

The Spearman rank correlation tests (Table 1 and Appendix A) show variable rho values and a tendency to low p-values. Since the gls results indicate that marine facies and outcrop area (at the scale 1:500000) were the proxies that best explained the diversity signal, these two proxies were further tested with the Spearman correlation test. This allows comparison with the results of Smith and Benson (2013). In both cases, in Portugal and in the UK, these two proxies yielded the lowest AIC values of all tested models, suggesting that facies heterogeneity and outcrop area together best explain the palaeodiversity signal.

With this result, we decided to further test the correlation between diversity and these two variables combined, by multiplying the marine facies signal with the outcrop curve. The same procedure was used by Smith and Benson (2013), and provides a proportion of the outcrop area of marine facies per time bin. The result was a stronger correlation (raw data) between diversity and the two proxies combined, but with variable rho values. The rho values for the raw data are systematically equal to, or lower than, those obtained for the marine facies and outcrop area isolated, implying that the combined proxies provide the highest explanatory power (Table 1 and Appendix A). On the other hand, the correlation values after generalized differencing yielded significantly better results than those from the two proxies isolated, with consistently low rho values. This suggests that overall trends were obscuring some aspects of the finer-scale, bin-by-bin correlations.

4. Discussion

4.1. *Palaeodiversity interpretations*

In the Lusitanian basin, there are no records of echinoids in the Triassic, because sedimentation was predominantly terrestrial. The first marine incursion happened in the early Hettangian, while true and stable marine conditions only became established during the Sinemurian, with the development of a carbonate ramp (Kullberg et al. 2013). The first echinoids appear in the Lusitanian basin only during the early Sinemurian (Fig. 3B), cidaroid spine fragments from São Pedro de Moel, Marinha Grande (Coimbra Formation). These are synchronous with the first true marine incursions into the Lusitanian basin. Later, rocks from the middle Sinemurian yield the first identifiable echinoid remains.

During the Pliensbachian and most of the Toarcian, no echinoids were found in the Lusitanian basin. At the beginning of the Pliensbachian, there was a rapid deepening of the basin, with an inundation maximum in the *Margaritatus* zone (upper Pliensbachian), according to Kullberg et al. (2013). By the late Toarcian and onwards, the number of echinoid taxa slowly increases, with a few drops, until the end of the Callovian.

The Callovian to Oxfordian transition is marked by a stratigraphic hiatus at the basin scale, which lasted until the middle Oxfordian, producing a hiatus in the fossil record. This episode corresponds to an early event of tectonic inversion, restricted to the Lusitanian Basin (Kullberg et al. 2013). Echinoid diversity peaked during the Late Jurassic, to about twice the level seen elsewhere in the remaining time series.

A significant drop marks the Jurassic to Cretaceous transition in echinoid diversity. This absence can be partially explained by the absence of marine sediments during the early Berriasian (Fig. 3C).

After this echinoid diversity drop, diversity recovers to levels not as high as the Jurassic record. This new diversification event coincides with a great increase in the number of irregular echinoids (Fig. 4A). During the Aptian, there was a significant decrease in

echinoid diversity, which lasted some time and is contemporaneous with the regional so-called “Aptian crisis” (Kullberg et al., 2013). After a progressive sea level rise initiated in the early Aptian, the Lusitanian basin received a rapid and high-energy influx of clastic material, transported by fluvial systems and eroding the lower layers, according to Rey (1999). This freshwater input would have had a major impact on the marine ecosystem, especially on stenohaline animals such as echinoderms. This environmental change may explain the significant decrease in echinoid diversity during great part of that stage.

During the Cenomanian, an extensive carbonate platform was established (Dinis et al., 2008), with, once again, conditions for the proliferation and diversification of echinoderms, including echinoids (Fig. 3A). By the end of the middle of the Late Cretaceous, the Lusitanian basin could no longer accommodate sediments and the sea retreated. The Coniacian to Campanian is marked by a shift in the sedimentation depocentre to offshore. Only small areas of the northern part of basin could support restricted shallow marine waters (Dinis et al., 2008), not ideal environmental conditions for echinoids, while the southern region suffered erosion. This restriction in the habitat could explain the poor diversity of echinoids for most of the Late Cretaceous.

After this narrative, it is clear that the evolution of the Lusitanian basin had a direct impact on the diversity signal obtained from the fossil record. As Smith (2007b) indicated, tectonic drivers change the basin’s area and volume, which influences the areal extent of continental flooding and, therefore, directly the diversity of marine faunas, according to a kind of species-area effect. Also, these tectonic events imply changes in basin sedimentation. As an example of this, Figure 3B demonstrates a good correlation between many of the diversity drops with major basin discontinuities.

4.2. *Comparing the Portuguese with the U.K. diversity*

Smith and Benson (2013) found that the British Cretaceous echinoids also occupied a passive margin linked to the opening of the North Atlantic. The number of species in their study is slightly larger than the dataset here, and it is focused on the Cretaceous. Their dataset is divided into stages, which divides the data into uneven time bins and provides a coarser timescale than here.

In the U.K., echinoid diversity during most of the Early Cretaceous is relatively low, with fewer than 10 species per stage. During the Aptian, echinoid diversity greatly increases, a trend that continues until the Cenomanian. After this period of increase, diversity decreases to about half the level that it reached during the Cenomanian. The numbers of species remained more or less stable until the Maastrichtian, when diversity decreased significantly.

The comparison between Portuguese and U.K. echinoid diversity shows a few similarities and a few differences (Fig. 6C). While the Early Cretaceous record of the U.K. is rather poor in echinoids, with fewer than 10 species through most of this time period, the Portuguese record is rather diverse, with about four times more species. This changes during the Aptian, and from this stage onwards, U.K. echinoid diversity is significantly larger than the Portuguese. Similarly in both areas, the Late Cretaceous starts with high diversity, decreasing afterwards. In order to be able to correlate the two echinoid datasets, we transformed our dataset to the stage level and performed the correlation tests applied for comparing our diversity signal with the proxies.

Echinoid diversity from the U.K. Cretaceous revealed a strong correlation between the diversity signal and the grouping of outcrop area and marine facies, for both raw data and for first differenced data. As previously mentioned, the same proxies used in combination and correlated with diversity yielded lower values, for both the raw and generalized differenced data. The difference between the correlation values obtained for the two studies could result from differences in the resolution of the datasets used. As mentioned before, Smith and

Benson (2013) organized their data by stages and used the stage midpoint to plot the data, while our data is organized by substages and 1 Myr time bins. Also, our outcrop dataset is based on maps with a larger scale than Smith and Benson's (2013) study.

In order to understand how similar the Portuguese and U.K. faunas were, we decided to correlate the two diversity signals. If the two diversity signals were a reflection of the geological evolution of the different basins, in a sense, biased by the rock record, the correlation between the two diversity datasets should be low. On the other hand, if biological factors were strongly imprinted on the diversity signals, the correlations between the data from the two basins should be high. We converted our initial dataset to the frame of Smith and Benson's (2013) dataset and performed the same correlation tests between the two datasets as were done for the Portuguese data alone. The tests were done on both the raw data and after applying generalized differencing. The rho values obtained were low for the raw data, and moderate after generalized differencing, but with high p-values (Table 1 and Appendix A). The high p-values provide very little confidence to state that the two datasets are statistically different.

4.3. *Common-cause vs Megabias hypothesis*

Two hypotheses seek to explain the apparent correlation between diversity and some sampling proxies, the common-cause and the megabias. Smith and Benson (2013) interpreted the strong correlation between the combined proxies and palaeodiversity as an indicator that the rock record strongly influences the diversity signal. They consider that the rock record is highly conditioned by periods of deposition and erosion, with erosion events obliterating information, thus the diversity signal is a direct reflection of Geology. We obtained similar results to Smith and Benson (2013), although with a less strong correlation between diversity

and the combined proxies. Therefore, we could assume that the rock record, and its availability, controls the diversity signal and, therefore, supports the megabiases hypothesis.

But, these results can also be interpreted as in favour of the common-cause hypothesis. The higher correlation obtained with diversity and the two proxies combined, in contrast to values for each proxy alone, could be a consequence of the moderate correlation values retrieved between marine facies and outcrop area. This moderate relationship could indicate a common factor influencing both proxies, which would be influencing diversity as well. The results of a strong correlation between diversity and facies area availability can also suggest a diversity control through the area of continental flooding, or area of habitat. For this to be true, a stronger correlation between diversity and sea level would be expected, something that this study did not reveal. Also, the generally poor correlation results (after generalized differencing) could argue against the idea that the correlation of fossil diversity and formation count provides evidence for megabiases, and that other factors may have had a more important role in controlling echinoid diversity. If true, local diversity signals should be intrinsically related with the number of formations at smaller scales. If this is not the case, as we suggest, then it seems that there may be some kind of summing effect, where numerous uncorrelated signals, when lumped together, produce an apparently reasonable correlation (Benton, 2012).

Other factors that might have influenced the diversity of this group have yet to be tested, such as predator-prey pressure, nutrient availability, or tectonics. Tectonics induces changes in the basin volume, in the area of continental flooding and, in consequence, influences the number of formations (Smith, 2007b). In sum, tectonics controls the overall amount and heterogeneity of the geological record that is preserved, as well as the quantity of fossil material that reaches us. As noted before, there is evidence for a relationship between the decrease in palaeodiversity and basin discontinuities (Fig. 3B), with diversity apparently

decreasing close to these discontinuities. This fact to be true and not an artefact of the method used seems to support the common-cause hypothesis. Discontinuities in the rock record at basin scale result from tectonic uplift and/or sea level changes. Such events would ultimately cause the erosion of the rock record and shrinking and loss of habitats would precede it. This change in habitat distribution would influence the overall diversity prior to the discontinuity. Although clearly an important driver, tectonics is hard to quantify for use in correlation tests.

The very low correlation values are in line with the conclusions of other studies at regional and local scales (for example, Dunhill et al., 2012, 2013, 2014a, b) and indicate that none of the factors assessed is, on its own, truly driving Portuguese echinoid palaeodiversity. Many previous studies have identified strong correlations between global diversity and the number of fossil-bearing formations (for example, Benson and Butler, 2011; Smith and Benson, 2013). On the other hand, our local study obtained similar results to other regional and local studies, where this correlation is poor. It would be good to identify definite cause-effect linkages, and these ought to be clear at local and regional scales, where exact matching between rises and falls in the time series under comparison ought to be evident.

5. Conclusions

Echinoid diversity during the Mesozoic in the Lusitanian basin is far from having a general trend, differing from previous global studies in other taxonomic groups. Many of the variations in diversity seem to be linked to basinal discontinuities and, therefore, with the evolution of the basin.

Comparison between the plots based on species, genera, and families suggests that diversity patterns are independent of taxonomic level. The different levels show similar diversity patterns, differing only in the magnitude of the changes.

The two different testing methods used to investigate the influence of the different sampling proxies on the diversity patterns obtained similar results. Both Spearman and gls tests indicated poor to moderate correlation between many of the analysed proxies and the diversity patterns. Among the different proxies tested, marine facies variation together with outcrop area showed highest correlation with the palaeodiversity data. Furthermore, both correlation tests and linear model fitting agree that the temperature and sea level changes appear to have had very little influence on echinoid diversity.

Acknowledgments

We thank G. Price for providing some of the isotopic data used in this study, Graeme Lloyd for providing some of the R scripts, and Andrew Smith, for his comments on this manuscript. BCP thanks Pablo Palacios, for his many discussions regarding many of the setbacks experienced with the R script, and all people's comments on this work provided during the last EEC meeting. This work was funded by FCT doctoral training grant SFRH/BD/68891/2010 to BCP. We also thank the Finn Surlyk, and the referees of the MS for many very helpful comments.

Appendix. Supplementary data

Supplementary data to this article can be found online at

References

- Aguirre-Urreta, M.B., Price, G.D., Ruffell, A.H., Lazo, D.G., Kalin, R.M., Ogle, N., Rawson, P.F., 2008. Southern hemisphere Early Cretaceous (Valanginian-Early Barremian) carbon and oxygen isotope curves from the Neuquén Basin, Argentina. *Cretaceous Research* 29, 87–99. doi:10.1016/j.cretres.2007.04.002.

963 Alberti, M., Fürsich, F.T., Pandey, D.K., 2012. The Oxfordian stable isotope record ($\delta^{18}\text{O}$,
 964 $\delta^{13}\text{C}$) of belemnites, brachiopods, and oysters from the Kachchh Basin (western India)
 965 and its potential for palaeoecologic, palaeoclimatic, and palaeogeographic
 966 reconstructions. *Palaeogeography, Palaeoclimatology, Palaeoecology* 344-345, 49–68.
 967 doi:10.1016/j.palaeo.2012.05.018.

968 Alroy, J., 2010. Geographical, environmental and intrinsic biotic controls on Phanerozoic
 969 marine diversification. *Palaeontology* 53: 6, 1211–1235, doi: 10.1111/j.1475-
 970 4983.2010.01011.x

971 Armendáriz, M., Rosales, I., Bádenas, B., Piñuela, L., Aurell, M. and García-Ramos, J.C.,
 972 2013. An approach to estimate lower Jurassic seawater oxygen-isotope composition
 973 using $\delta^{18}\text{O}$ and Mg/Ca ratios of belemnite calcites (Early Pliensbachian, northern
 974 Spain). *Terra Nova* 25, 439–445. doi:10.1111/ter.12054.

975 Barrett, P.M., McGowan, A.J., Page, V., 2009. Dinosaur diversity and the rock record.
 976 *Proceedings of the Royal Society B* 276, 2667-2674.

977 Benson, R.B.J., Butler, R.J., 2011. Uncovering the diversification history of marine tetrapods:
 978 ecology influences the effect of geological sampling biases. *Geological Society Special*
 979 *Publications* (London) 358, 191-208.

980 Benson, R.B.J., Butler, R.J., Lindgren, J., Smith, A.S., 2010. Mesozoic marine tetrapod
 981 diversity: mass extinctions and temporal heterogeneity in geological megabiases
 982 affecting vertebrates. *Proceedings of the Royal Society B* 277, 829-834.

983 Benson, R.B.J., Mannion, P.D., 2012. Multivariate models are essential for understanding
 984 vertebrate diversification in deep time. *Biology Letters* 8, 127-130.

985 Benson, R.B.J., Upchurch, P., 2013. Diversity trends in the establishment of terrestrial
 986 vertebrate ecosystems: interactions between spatial and temporal sampling bases.
 987 *Geology* 41, 43-46. doi:10.1130/G33543.1

988 Benton, M.J., 2012. No gap in the middle Permian record of terrestrial vertebrates. *Geology*
989 40, 339-342.

990 Benton, M.J., Dunhill, A.M., Lloyd, G.T., Marx, F.G., 2011. Assessing the quality of the
991 fossil record: insights from vertebrates. In: McGowan, A.J., Smith, A.B. (eds.)
992 Comparing the Geological and Fossil Records: Implications for Biodiversity Studies.
993 Geological Society Special Publications (London) 358, 63-94.

994 Benton, M.J.; Ruta, M.; Dunhill, A.M., Sakamoto, M., 2013. The first half of tetrapod
995 evolution, sampling proxies, and fossil record quality. *Palaeogeography,*
996 *Palaeoclimatology, Palaeoecology* 372, 18-41.

997 Benton, M.J., Tverdokhlebov, V.P., Surkov, M.V., 2004. Ecosystem re-modelling among
998 vertebrates at the Permian-Triassic boundary in Russia. *Nature* 432, 97-100.

999 Brocklehurst, N., Fröbisch, J., 2014. Current and historical perspectives on the completeness
1000 of the fossil record of pelycosaurian-grade synapsids. *Palaeogeography,*
1001 *Palaeoclimatology, Palaeoecology* 399, 114–126.

1002 Burnham, K.P.; Anderson, D.R., 2002. *Model Selection and Multimodel Inference: A*
1003 *Practical Information-Theoretic Approach*. 2nd edition, Springer-Verlag, 488 pp.

1004 Butler, R.J., Barrett, P.M., Nowbath, S., Upchurch, P., 2009. Estimating the effects of the
1005 rock record on pterosaur diversity patterns: implications for hypotheses of bird/pterosaur
1006 competitive replacement. *Paleobiology* 35, 432-446.

1007 Butler, R.J., Benson, R.B.J., Carrano, M.T., Mannion, P.D., Upchurch, P., 2011. Sea level,
1008 dinosaur diversity and sampling biases: investigating the 'common cause' hypothesis in
1009 the terrestrial realm. *Proceedings of the Royal Society of London B* 278, 1107-1113.

1010 Carpentier, C., Martin-Garin, B., Lathuiliere, B., Ferry, S., 2006. Correlation of reefal
1011 Oxfordian episodes and climatic implications in the eastern Paris Basin (France). *Terra*
1012 *Nova* 18, 191-200.

1013 Chamberlin, T.S., 1909. Diastrophism as the ultimate basis of correlation. *Journal of Geology*
1014 17, 689-693.

1015 Choffat, P., 1880. *Étude Stratigraphique et Paléontologique des Terrains Jurassiques du*
1016 *Portugal. Lisbon.*

1017 Crampton, J.S., Beu, A.G., Cooper, R.A., Jones, C.A., Marshall, B., Maxwell, P.A., 2003.
1018 Estimating the rock volume bias in paleobiodiversity studies. *Science* 301, 358-360.

1019 Crampton, J.S., Foote, M., Beau, A.G., Cooper, R.A., Matcham, I., Jones, C.M., Maxwell,
1020 P.A., Marshall, B.A., 2006. Second-order sequence stratigraphic controls on the quality
1021 of the fossil record at an active margin: New Zealand Eocene to recent shelf molluscs.
1022 *Palaios* 21, 86-105.

1023 Cresta, S., Goy, A., Ureta, S., Arias, C., Barrón, E., Bernad, J., Canales, M. L., García-Joral,
1024 F., García-Romero, E., Gialanella, P. R., Gómez, J. J., González, J. A., 2001. The Global
1025 Boundary Stratotype Section and Point (GSSP) of the Toarcian-Aalenian Boundary
1026 (lower-middle Jurassic). *Episodes* 24, 166-175.

1027 Dera, G., Brigaud, B., Monna, F., Laffont, R., Pucéat, E., Deconinck, J.-F., Pellenard, P.,
1028 Joachimski, M.M., Durlet, C., 2011. Climatic ups and downs in a disturbed Jurassic
1029 world. *Geology* 39: 3, 215-218.

1030 Dinis, J.L., Rey, J., Cunha, P.P., Callapez, P., Pena dos Reis, R., 2008. Stratigraphy and
1031 allogenic controls of the western Portugal Cretaceous: an updated synthesis. *Cretaceous*
1032 *Research* 29, p. 772-780.

1033 Doke, J., 2007. Grabit.m, The MathWorks MatLab Central Website.
1034 (<http://www.mathworks.com/matlabcentral/fileexchange/7173-grabit>). Accessed in
1035 August 2014.

1036 Dunhill, A.M., 2011. Using remote and a GIS to quantify rock exposure area in England and
 1037 Wales: implications for palaeodiversity studies. *Geology* 39, 11-114. doi:
 1038 10.1130/G31503.1
 1039 Dunhill, A.M., 2012. Problems with using rock outcrop area as a paleontological sampling
 1040 proxy: rock outcrop and exposure area compared with coastal proximity, topography,
 1041 land use, and lithology. *Paleobiology* 38, 126-143.
 1042 Dunhill, A.M., Benton, M.J., Newell, A.J., Twitchett, R.J., 2013. Completeness of the fossil
 1043 record and the validity of sampling proxies: a case study from the Triassic of England
 1044 and Wales. *Journal of the Geological Society (London)* 170, 291-300.
 1045 Dunhill, A.M., Benton, M.J., Twitchett, R.J., Newell, A.J., 2012. Completeness of the fossil
 1046 record and the validity of sampling proxies at outcrop level. *Palaeontology* 55, 1155-
 1047 1175.
 1048 Dunhill, A.M., Benton, M.J., Twitchett, R. J., Newell, A.J., 2014a. Testing the fossil record:
 1049 sampling proxies and scaling in the British Triassic-Jurassic. *Palaeogeography,*
 1050 *Palaeoclimatology, Palaeoecology* 404, 1-11. doi: 10.1016/j.palaeo.2014.03.026
 1051 Dunhill, A.M., Hannisdal, B., Benton, M.J., 2014b. Disentangling rock record bias and
 1052 common-cause from redundancy in the British fossil record. *Nature Communications* 5:
 1053 4818. doi:10.1038/ncomms5818
 1054 Fara, E., 2001. What are Lazarus taxa?. *Geological Journal* 36, 291–303. doi:
 1055 10.1002/gj.879.abs.
 1056 Foote, M., Sepkoski, J.J.Jr., 1999. Absolute measures of the completeness of the fossil
 1057 record. *Nature* 398, 415-417.
 1058 Forbes, E., in Sharpe, D., 1850. Description of fossil Echinidae from Portugal. *Quarterly*
 1059 *Journal of the Geological Society London* 6, 195-199. doi:
 1060 10.1144/GSL.JGS.1850.006.01-02.18

1061 Fröbisch, J., 2008. Global taxonomic diversity of Anomodonts (Tetrapoda, Therapsida) and
 1062 the terrestrial rock record across the Permian-Triassic boundary. PLOS ONE 3, e3733.

1063 Fursich, F.T., Singh, I.B., Joachimski, M., Krumm, S., Schfirt, M., Schlirf, S., 2005.
 1064 Palaeoclimate reconstructions of the middle Jurassic of Kachchh (western India): an
 1065 integrated approach based on palaeoecological, oxygen isotopic, and clay mineralogical
 1066 data. Palaeogeography, Palaeoclimatology, Palaeoecology 217, 289-309.

1067 Graciansky, P.-C. de, Hardenbol, J., Jacquin, T., Vail, P. R., eds., 1998, Mesozoic and
 1068 Cenozoic sequence stratigraphy of European basins, Society for Sedimentary Geology
 1069 (SEPM) Special Publication 60, 786 p.

1070 Gradstein, F.M, Ogg, J.G., Schmitz, M.D., Ogg, G.M., 2012. The Geologic Time Scale 2012.
 1071 Elsevier, 1176. doi: 10.1016/B978-0-444-59425-9.00004-4.

1072 Hannisdal, B.; Peters, S.E., 2011. Phanerozoic Earth System Evolution and Marine
 1073 Biodiversity. Science 334: 6059, 1121-1124. doi: 10.1126/science.1210695

1074 Haq, B.U., Hardenbol, J., Vail, P.R., 1987. Chronology of fluctuating sea levels since the
 1075 Triassic. Science 235, 1156-1166. doi:10.1126/science.235.4793.1156

1076 Huber, B.T., MacLeod, K.G., Gröcke, D.R., Kucera, M., 2011. Paleotemperature and
 1077 paleosalinity inferences and chemostratigraphy across the Aptian/Albian boundary in the
 1078 subtropical North Atlantic. Paleoceanography 26, PA422. doi:10.1029/2011PA002178

1079 Jackson, J.B.C., Johnson, K.G., 2001. Measuring past biodiversity. Science, 293, 2401-2403.
 1080 doi:10.1126/science.1063789

1081 Jepson, J.E., Penney, D., 2007. Neuropteran (Insecta) palaeodiversity with predictions for the
 1082 Cretaceous fauna of the Wealden. Palaeogeography, Palaeoclimatology, Palaeoecology
 1083 248, 109-118. doi:10.1016/j.palaeo.2006.11.015

1084 Kroh, A., Smith, A.B., 2010. The phylogeny and classification of post-Palaeozoic echinoids.
 1085 Journal of Systematic Palaeontology 8:2, 147-212.

- 1086 Kullberg, J.C., Rocha, R.B., Soares, A.F., Rey, J., Terrinha, P., Azerêdo, A.C., Callapez, P.,
 1087 Duarte, L.V., Kullberg, M.C., Martins, L., Miranda, R., Alves, C., Mata, J., Madeira, J.,
 1088 Mateus, O., Moreira, M., Nogueira, C.R., 2013. III.3. A Bacia Lusitaniana: Estratigrafia,
 1089 Paleogeografia e Tectónica. In: Dias, R., Araújo, A., Terrinha, P., Kullberg, J.C. (eds.),
 1090 Geologia de Portugal, II – Geologia Meso-cenozóica de Portugal, 196-347.
- 1091 Lambert, J., 1913. Échinides Calloviens du plateau de Césareda (Portugal). Comunicações
 1092 dos Serviços Geológicos de Portugal IX, 69-76.
- 1093 Lambert, J., 1915-1916. Note sur quelques échinides de la grande oolithe (Bathonien) et du
 1094 Callovien du massif de Porto-de-Moz (Portugal). Comunicações Serviços Geológicos
 1095 du Portugal XI, 85-96.
- 1096 Lazar, I., Panaiotu, C.E., Grigore, D., Sandy, M.R., Peckmann, J., 2011. An unusual
 1097 brachiopod assemblage in a Late Jurassic (Kimmeridgian) stromatactis mud-mound of
 1098 the Eastern Carpathians (Haghimas Mountains), Romania. *Facies* 57, 627-647.
- 1099 Lloyd, G., 2008. Dating phylogenetic trees of fossil taxa.
 1100 <http://www.graemetlloyd.com/methdpf.html>
- 1101 Lorient, P., 1884. Notes pour servir a l'étude des échinodermes I. *Recueil Zoologique Suisse* 1,
 1102 605-643.
- 1103 Lorient, P., 1887-1888. *Recueil d'Études Paléontologiques sur la Faune Crétacique du*
 1104 *Portugal*, vol. II, Description des Echinides. Commission des Travaux Géologiques du
 1105 Portugal, 1 and 2.
- 1106 Lorient, P., 1890-1891. Description de la Faune Jurassique du Portugal, Embranchement des
 1107 Échinodermes. Commission des Travaux Géologiques du Portugal.
- 1108 Lorient, P., 1900. Notes pour servir a l'etude des Echinodermes VIII. *Revue Suisse de*
 1109 *Zoologie*, vol. 8, 55-96.

1110 Lorient, P., 1905. Notes pour Servir à L'étude Des Échinodermes, Genève, second series III,
1111 119-146.

1112 Mander, L., Twitchett, R. J., 2008. Quality of the Triassic-Jurassic bivalve fossil record in
1113 Northwest Europe. *Palaeontology* 51, 1213-1223.

1114 Martin-Garin, B., Lathuiliere, B., Geister, J., Ramseyer, K., 2010. Oxygen Isotopes and
1115 Climatic Control of Oxfordian coral reefs (Jurassic, Tethys). *Palaios* 25, 721-729.

1116 Marx, F. G., Uhen, M. D., 2010. Climate, critters, and cetaceans: Cenozoic drivers of the
1117 evolution of modern whales. *Science* 327, 993-996.

1118 Mayhew, P.J., Jenkins, G.B., Benton, T.G. 2008. A long-term association between global
1119 temperature and biodiversity, origination and extinction in the fossil record. *Proceedings*
1120 *of the Royal Society B* 275, 47-53.

1121 Mayhew, P.J., Bell, M. A., Benton, T.G., McGowan, A. J., 2012. Biodiversity racks
1122 temperature over time. *PNAS* 109: 38, 15141-15145.

1123 Mazerolle, M.J., 2012. Package ‘‘AICcmodavg. Model selection and multimodel inference
1124 based on (Q)AIC (c) version 1.24. R Foundation for Statistical Computing, Vienna,
1125 Austria.

1126 McGowan, M.L., Smith, A.B., 2008. Are global Phanerozoic marine diversity curves truly
1127 global? A study of the relationship between regional rock records and global
1128 Phanerozoic marine diversity. *Paleobiology* 34; 1, 80-103.

1129 McGowan, M. L., Smith, A. B., (eds.) 2011. Comparing the Geological and Fossil Records:
1130 Implications for Biodiversity Studies. Geological Society, London.

1131 McKinney, M.L., McNamara, K.J., Carter, B.D., Donovan, S.K., 1992. Evolution of
1132 Paleogene echinoids: a global and regional view. In: Prothero, D.R., Berggren, W.A.
1133 (eds.). *Eocene-Oligocene Climatic and Biotic Evolution*. Princeton University Press,
1134 349-367. ISBN: 9780691604954

1135 Miller, K.G., Kominz, M.A., Browning, J.V.; Wright, J.D.; Mountain, G.S., Katz, M.E.,
 1136 Sugarman, P.J., Cramer, B.S., Christie-Blick, N., Pekar, S.F., 2005. The Phanerozoic
 1137 Record of Global Sea-Level Change. *Science* 310, 1293-1298.
 1138 Myers, T.S., Tabor N.J., Jacobs L.L., Mateus O., 2012a. Palaeoclimate of the Late Jurassic of
 1139 Portugal: Comparison with the Western United States. *Sedimentology* 59:6, 1695–1717.
 1140 Myers, T.S., Tabor N.J., Jacobs L.L., Mateus O., 2012b. Estimating soil pCO₂ using paleosol
 1141 carbonates: implications for the relationship between primary productivity and faunal
 1142 richness in ancient terrestrial ecosystems. *Paleobiology* 38, 585–604.
 1143 Newell, N.D., 1967. Paleontological gaps in geochronology. *Journal of Paleontology* 36, 592-
 1144 610.
 1145 Peters, S.E., Foote, M., 2001. Biodiversity in the Phanerozoic: a reinterpretation.
 1146 *Paleobiology* 27,583-601.
 1147 Peters, S.E., Foote, M., 2002. Determinants of extinction in the fossil record. *Nature* 416,
 1148 420-424.
 1149 Picard, S., Garcia, J.P., Lecuyer, C., Sheppard, S.M.F., Cappetta, H., Emig, C.C., 1998. $\delta^{18}\text{O}$
 1150 values of coexisting brachiopods and fish: Temperature differences and estimates of
 1151 paleo-water depths. *Geology* 26, 975-978.
 1152 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., 2004. NLME: linear and nonlinear mixed
 1153 effects models. R package version 3.1-53.
 1154 ([http://www.stats.bris.ac.uk/R/src/contrib/2.0.1-patched/Recommended/nlme_3.1-](http://www.stats.bris.ac.uk/R/src/contrib/2.0.1-patched/Recommended/nlme_3.1-56.tar.gz)
 1155 [56.tar.gz](http://www.stats.bris.ac.uk/R/src/contrib/2.0.1-patched/Recommended/nlme_3.1-56.tar.gz))
 1156 Price, G., Sellwood, B., 1994. Paleotemperatures indicated by Upper Jurassic (Kimmeridgian
 1157 Tithonian) fossils from Mallorca determined by oxygen-isotope composition.
 1158 *Palaeogeography, Palaeoclimatology, Palaeoecology* 110: 1-2, 1-10.

1159 Price, G.D., Teece, C., 2010. Reconstruction of Jurassic (Bathonian) palaeosalinity using
 1160 stable isotopes and faunal associations. *Journal of the Geological Society* 167: 6, 1199–
 1161 1208.

1162 Price, G.D.; Twitchett, R.J.; Wheeley, J.R. , Buono, G. (2013). Isotopic evidence for long
 1163 term warmth in the Mesozoic. *Science Reports* 3: 1438. doi:10.1038/srep01438

1164 Raup, D.M., 1972. Taxonomic diversity during the Phanerozoic. *Science* 177, 1065-1071.

1165 Raup, D.M., 1976. Species diversity in the Phanerozoic: an interpretation. *Paleobiology* 2,
 1166 289–297.

1167 Rey, J., 1972. Recherches géologiques sur le Crétacé inférieur de l'Estremadura (Portugal).
 1168 *Memória dos Serviços Geológicos de Portugal*, new series, 21.

1169 Robeck, H.E., Maley, C.C., Donoghue, M.J. (2000). Taxonomy and temporal diversity
 1170 patterns. *Paleobiology* 26, 171–187.

1171 Saito, T., Van Donk, J., 1974. Oxygen and carbon isotope measurements of Late Cretaceous
 1172 and Early Tertiary foraminifera. *Micropaleontology* 20, 152–177.

1173 Sepkoski J.J., Jr, Bambach, R.K., Raup, D.M., Valentine, J.W., 1981. Phanerozoic marine
 1174 diversity and the fossil record. *Nature* 293, 435-437.

1175 Smith, A.B., 2001. Large-scale heterogeneity of the fossil record: implications for
 1176 Phanerozoic biodiversity studies. *Philosophical Transactions of the Royal Society of*
 1177 *London B* 356, 351–367.

1178 Smith, A.B., 2007a. Marine diversity through the Phanerozoic: problems and prospects.
 1179 *Journal of the Geological Society* 164, 731–745.

1180 Smith, A.B., 2007b. Intrinsic versus extrinsic biases in the fossil record: Contrasting the
 1181 fossil record of echinoids in the Triassic and early Jurassic using sampling data,
 1182 phylogenetic analysis and molecular clocks. *Paleobiology* 33, 310–323.

1183 Smith, A.B., Benson, R.B.J., 2013. Marine diversity in the geological record and its
 1184 relationship to surviving bedrock area, lithofacies diversity, and original marine shelf
 1185 area. *Geology* 41, 171–174. doi:10.1130/G33773.1.

1186 Smith, A.B., Lloyd, G.T., McGowan, A.J., 2012. Phanerozoic marine diversity: rock record
 1187 modelling provides an independent test of large-scale trends. *Proceedings of the Royal*
 1188 *Society B* 279, 4489–4495.

1189 Smith, A.B., McGowan, A.J., 2005. Cyclicity in the fossil record mirrors rock outcrop area.
 1190 *Biology Letters* 1, 443–445.

1191 Suan, G., Mattioli, E., Pittet, B., Lécuyer, C., Suchéras-Marx, B., Duarte, L.V., Philippe, M.,
 1192 Reggiani, L., Martineau, F., 2010. Secular environmental precursors to Early Toarcian
 1193 (Jurassic) extreme climate changes. *Earth and Planetary Science Letters* 290: 3–4, 448–
 1194 458. ISSN 0012-821X, <http://dx.doi.org/10.1016/j.epsl.2009.12.047>

1195 Upchurch, P., Mannion, P., Benson, R., Butler, R., Carrano, M., 2011. Geological and
 1196 anthropogenic controls on the sampling of the terrestrial fossil record: a case study from
 1197 the Dinosauria. *Geological Society (London) Special Publications* 358, 209–240.

1198 Vadet, A., Willie, E., 2002. Quelques oursins du Lusitanien du Portugal. *Échinidés nouveaux*
 1199 *peu ou mal connus. Annales de la Société D'Histoire Naturelle du Boulonnais* II: 3, 8–18.

1200 Veizer, J., Bruckschen, P., Diener, A., Azmy, K., Ebner, S., Buhl, D., Ala, D., Bruhn, F.,
 1201 Strauss, H., Korte, C., Carden, G.A.F., Jasper, T., Godderis, Y., Pawellek, F., Podlaha,
 1202 O.G., 1999. $^{87}\text{Sr}/^{86}\text{Sr}$, $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ evolution of Phanerozoic seawater. *Chemical*
 1203 *Geology* 161, 59–88.

1204 Voigt, S., 2000. Stable oxygen and carbon isotopes from brachiopods of southern England
 1205 and northwestern Germany: estimation of upper Turonian palaeotemperatures.
 1206 *Geological Magazine* 137, 687–703.

1207 Voigt, S., Wilmsen, M., Mortimore, R. N., Voigt, T., 2003. Cenomanian palaeotemperatures
1208 derived from the oxygen isotopic composition of brachiopods and belemnites: evaluation
1209 of Cretaceous palaeotemperature proxies. *International Journal of Earth Sciences* 92: 2,
1210 285-299.

1211 Voigt, S., Gale, A.S., Flögel, S., 2004. Midlatitude shelf seas in the Cenomanian-Turonian
1212 greenhouse world: Temperature evolution and North Atlantic circulation.
1213 *Paleoceanography* 19, PA4020. doi: 10.1029/2004PA001015.

1214 Wall, P.D., Ivany, L.C., Wilkinson, B.H., 2009. Revisiting Raup: exploring the influence of
1215 outcrop area on diversity in light of modern sample-standardization techniques.
1216 *Paleobiology* 35, 146-167.

1217 Wall, P.D., Ivany, L.C., Wilkinson, B.H., 2011. Impact of outcrop area on estimates of
1218 Phanerozoic terrestrial biodiversity trends. *Geological Society (London) Special*
1219 *Publications* 358, 53-62.

1220 Wang, S.C., Dodson, P., 2006. Estimating the diversity of dinosaurs. *PNAS* 103, 13601-
1221 13605.

1222 Wierzbowski, H., 2002. Detailed oxygen and carbon isotope stratigraphy of the Oxfordian in
1223 Central Poland. *International Journal of Earth Sciences* 91, 304-314.

Table 1

Correlation coefficients (Spearman's rho, ρ) and p-values between the palaeodiversity and the different proxies, at species level. The palaeodiversity signal refers to the diversity of the Lusitanian Basin, excluding taxa identification based on spine morphology.

Comparison	Spearman test		Generalized differences + Spearman test	
	ρ	p-value	P	p-value
Diversity ^ Total number of formations	0.3478908	< 2.2e-16	0.4750611	< 2.2e-16
Diversity ^ Number of marine formations	0.1424233	1.773e-07	0.2783111	< 2.2e-16
Diversity ^ All Facies	0.6349632	< 2.2e-16	0.337582	3.004e-06
Diversity ^ Marine Facies	0.5643569	< 2.2e-16	0.3516584	1.086e-06
Diversity ^ Outcrop (1:1000000)	0.4052215	9.622e-09	-0.05812138	0.4316
Diversity ^ Outcrop (1:500000)	0.6965468	< 2.2e-16	0.3668568	3.429e-07
Diversity ^ δ^{18} Oxygen	0.1746603	1.36e-10	-0.006837237	0.8031
Diversity ^ δ^{13} Carbon	0.2074738	1.991e-14	-0.04681501	0.08765
Diversity ^ Sea level (Haq et al., 1987)	0.2958423	< 2.2e-16	-0.04797601	0.08006
Diversity ^ Sea level (Miller et al., 2005)	0.3293531	< 2.2e-16	0.02278169	0.4061
Diversity ^ Sea level (Graciansky et al., 1998)	0.3005157	< 2.2e-16	-0.05214827	0.05707
Diversity ^ Outcrop (1:500000) x Marine Facies	0.7253658	< 2.2e-16	0.4631989	4.165e-11
Lusitanian ^ U.K. species diversity	0.2952271	0.3515	0.4363636	0.1825

Table 2

Comparison of generalized-differenced time series models between palaeodiversity and the proxies. The palaeodiversity signal refers to the diversity of the Lusitanian Basin, excluding taxa based only on spines.

Proxies	AIC	Coefficients	p-values
All Facies	5546.096	1.001621	0
Marine Facies	5526.639	1.266207	0
Outcrop (1:50000)	5628.395	0.178438	0
Outcrop (1:50000) + Marine Facies	5511.596	0.104395	0
		1.15307	0

Figure legends

Fig. 1. Map with the location of the areas under study. The Lusitanian and Algarve Basins are highlighted.

Fig. 2. Cumulative collectors curve of echinoid species descriptions through historical time, for the Lusitanian basin. Taxa based on spines morphology excluded.

Fig. 3. (A) Echinoid species (solid line), genera (dashed line) and family (dashed dotted line) diversity, after removing species based on spine morphology, in the Mesozoic of the Lusitanian Basin. (B) Echinoid species diversity, after excluding species based on spine morphology (solid line) and major basin discontinuities (dark grey rectangular zones), after Kullberg *et al.* (2013), in the Mesozoic of the Lusitanian Basin. (C) Echinoid species diversity (solid line), excluding species based on spine morphology, total number of formations (dashed line) and number of marine formations (dashed and dotted line) for the Mesozoic of Lusitanian Basin. The grey rectangles represent the stages.

Fig. 4. (A) Regular (solid line) and Irregular (dashed line) echinoid species diversity, in the Lusitanian Basin, after excluding taxa identified based on spine morphology. (B) Number of echinoid species described for the studied time period (solid line) and number of echinoid species, when excluding the species described/identified on spine morphology (dashed line), for the Portuguese Mesozoic. (C) Echinoid diversity at genera (solid line) and corrected genera for the Lazarus effect (dashed line) curves, in the Mesozoic of the Lusitanian Basin. The grey rectangles represent the stages.

Fig. 5. (A) Echinoid species diversity, after removing species based on spine morphology (solid line), with the facies (dashed line) and marine facies (dashed dotted line) variation for the Lusitanian Basin. (B) Echinoid species diversity without species based on spine morphology (solid line), with the outcrop area based on the geological maps at the scale 1:1000000 (dashed line) and 1:500000 (dashed dotted line). (C) Echinoid species diversity, after removing species based on spine morphology (solid line), in the Mesozoic of the Lusitanian Basin, isotopic values of δ^{18} Oxygen of brachiopods, oysters and foraminifera (grey dots) and the smooth curve generated with the Kernel regression method, with a bandwidth of 5 Myr (dashed dotted line).

Fig.6. (A) Echinoid species diversity, after removing species based on spine morphology (solid line), in the Mesozoic of the Lusitanian Basin, isotopic values of δ^{13} Carbon of brachiopods, oysters and foraminifera (grey dots) and the smooth curve generated with the Kernel regression method, with a bandwidth of 5 Myr (dashed dotted line). (B) Echinoid species diversity in the Mesozoic of the Lusitanian Basin (solid black line) and global sea level curves based on Haq et al., 1987, Graciansky et al. (1998) and Miller et al., 2005 (light grey line). (C) Comparison between the Portuguese (solid line) and the U.K. (dashed line) Cretaceous echinoid diversity.

Fig. 1.

1284

1285 **Fig. 2.**

Fig. 3.

1288

1289 **Fig. 4.**

1290

1291 **Fig. 5.**

Fig. 6.

Bruno Miguel Claro Pereira
Department of Earth Sciences
Wills Memorial Building
Queens Road
Bristol
BS8 1RJ
United Kingdom
T: +44 (0)117 331 5133
E-mail: glbmcp@bristol.ac.uk

2 February 2015

Dear Finn Surlyk (editor of *Palaeo* 3),

I hereby resubmit the manuscript entitled "**Mesozoic echinoid diversity in Portugal: investigating fossil record quality and environmental constraints on a regional scale**", for your reconsideration for publication in *Palaeo*-3. The manuscript was previously entitled "**Echinoid diversity in Portugal, sampling proxies and fossil record quality: a regional-scale study**". The title has been changed following the comment from reviewer 1.

We would like to thank all the comments that you and the reviewers provided to the submitted manuscript. All the points raised were fair and we tried to accommodate those suggestions, in the new version. Please, report to the response document, provided as an attached file of the resubmission, for a more detailed response to each comment made.

Yours Sincerely,

A handwritten signature in black ink, appearing to read "Bruno Pereira", with a large, sweeping flourish underneath.

Bruno Pereira

Figure 1
[Click here to download high resolution image](#)

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6