Precession-driven Changes in Iceland-Scotland Overflow Water Penetration and Bottom Water Circulation on Gardar Drift Since ~ 200 ka

A. C. Elmore^{1*}, J.D. Wright², T. Chalk³

¹ Department of Geography, Durham University, Durham, England, DH1 3LE

² Department of Earth and Planetary Sciences, Rutgers University, New Brunswick, NJ
08854

³ Department of Physical Oceanography, Woods Hole Oceanographic Institution, 266 Woods
Hole Road, Woods Hole, MA 02543

* Corresponding author's address aurora.elmore@durham.ac.uk

1. Abstract

Benthic foraminiferal stable isotopic records from a transect of sediment cores south of the Iceland-Scotland Ridge reveal that the penetration depth of Iceland-Scotland Overflow Water (ISOW) varied on orbital timescales with precessional pacing over the past ~ 200 kyr. Similar, higher benthic foraminiferal $\delta^{13}\text{C}$ values (~ 1.0 ‰) were recorded at all transect sites downstream of the Iceland-Scotland Ridge during interglacial periods (Marine Isotope Chrons 5 and 1), indicating a deeply penetrating ISOW. During glacial periods (Marine Isotope Chrons 6, 4, and 2), benthic foraminiferal $\delta^{13}\text{C}$ values from the deeper (2700-3300 m), southern sites within this transect were significantly lower (~ 0.5 ‰) than values from the northern (shallower) portion of the transect (~ 1.0 ‰), reflecting a shoaling of ISOW and greater influence of glacial Southern Component Water (SCW) in the deep Northeast Atlantic. Particularly during intermediate climate states, ISOW strength is driven by precessional cycles, superimposed on the large-scale glacial-interglacial ISOW variability.

26 Millennial-scale variability in the penetration of ISOW, likely caused by high-frequency
27 Heinrich and Dansgaard-Oeschger Events, is most pronounced during intermediate climate
28 states.

29

30 **2. Introduction**

31 Variability in the formation of deep Northern Component Water (NCW; analogous
32 to modern North Atlantic Deep Water; NADW) is linked to regional and global climate
33 changes on orbital time-scales (e.g., Broecker and Denton, 1989; Raymo et al., 1989). Causal
34 mechanisms for variations in NCW formation are thought to be associated with incoming
35 solar radiation (insolation), fresh-water input (Elmore and Wright, 2011; Elmore et al., 2015),
36 and/or sea ice extent (Broecker and Denton, 1989; Rind et al., 2001; Dokken et al., 2013).
37 During the late Pleistocene, changes in insolation in the high northern latitudes were the
38 product of the combined influences of variations in the Earth's orbital parameters of
39 eccentricity (~ 400 kyr and 100 kyr), obliquity (~ 40 kyr) cycles, and precession (~ 23 kyr)
40 (Laskar et al., 1993). Since past variations in NCW production are tied to climatic conditions
41 (including surface water temperature and wind strength), studies have linked particular NCW
42 circulation patterns to different climate states, notably the glacial and interglacial end-
43 members (e.g., Broecker et al., 1985; Imbrie et al., 1993; Liseicki et al., 2008). However,
44 insolation changes are not simply sinusoidal because the total insolation received is derived
45 from distinct orbital periodicities (Laskar et al., 1993); and therefore, the deep-water
46 circulation states may be unique for each glacial and interglacial period, as well as for
47 intermediate climate states. Thus, there is significant scope for variations of NCW production
48 to change both on glacial-interglacial timescales and with the precise orbitals associated,
49 however these NCW variations remain unresolved.

50 Modern NADW is produced by the interplay between five intermediate to deep
51 water-mass components, Iceland-Scotland Overflow Water (ISOW; ~ 5 Sv), Denmark Strait
52 Overflow Water (DSOW; ~ 5 Sv), Antarctic Bottom Water (AABW; ~ 1 Sv), Labrador Sea
53 Water (~ 1 Sv), and Mediterranean Outflow Water (~ 1 Sv) (Worthington, 1976). Kuijpers et
54 al. (1998) suggested that ISOW is the most important water mass for forming NADW;
55 however, uncertainty remains as to the historical flux, strength, and penetration depth, related
56 to the density of ISOW, particularly during intermediate climate states. While variability in
57 each of the components of NADW is not fully understood, it has been proposed that changes
58 in these components have contributed to the orbital-scale climate variability (Hillaire-Marcel
59 et al., 1994, Raymo et al., 2004; Millo et al., 2006). Iceland-Scotland Overflow Water forms
60 by convection in the Norwegian and Greenland Seas and the Arctic Ocean (Worthington,
61 1976; Mauritzen, 1996) and penetrates to >4 km depth (Dickson et al., 1990). The total
62 contribution of ISOW is related to surface temperature and salinity in the Nordic Seas
63 (Duplessy et al., 1988a), as well as to the volume of surface inflow (Worthington, 1976),
64 amount of sea ice cover (Prins et al., 2002; Raymo et al., 2004), wind forcing (Kohl et al.,
65 2007) and regional tectonics (Wright and Miller, 1996). Sea-level and sill depth assert a
66 control on ISOW contribution because they regulate the sensitive cross sectional volume
67 above the Iceland-Scotland-Faeroe Ridge, through-which overflow water can pass between
68 the open North Atlantic and the Nordic Seas (Millo et al., 2006). Thus, researchers suggested
69 that ISOW strength and penetration depth has varied on time-scales of millions of years
70 (Wright and Miller, 1996), tens of thousands of years (i.e. orbital scale; Duplessy et al.,
71 1988a; Raymo et al., 2004), thousands of years (i.e. millennial; Oppo et al., 1995; Dokken
72 and Hald, 1996; McManus et al., 1999), hundreds of years (Bianchi and McCave, 1999;
73 Moffa-Sanchez et al., 2015), years (Turrell et al., 1999), or seasons (Hatun et al., 2004).

74 Iceland-Scotland Overflow Water formation was likely vigorous and deeply
 75 penetrating during interglacial Marine Isotope Chron (MIC) 1 and 5e (Duplessy et al.,
 76 1988a,b; Kissel et al., 1997) and weaker and/or less deeply penetrating during the Last
 77 Glacial Maximum (LGM; MIC 2; Duplessy et al., 1998a; Kissell et al., 1997; Yu et al., 2008;
 78 Sarnthein et al., 2007); however, the history of ISOW is unclear for other glacial and
 79 intermediate climate states. For instance, Kuijpers et al. (1998) argued for enhanced
 80 formation of ISOW during MIC 6, while Kissel et al. (1997) and Rasmussen et al. (2003)
 81 considered it minimal. Kissel et al. (1997) presumed ISOW was vigorous during MIC 5d,
 82 while Kuijpers et al. (1998) and Rasmussen et al. (2003) suggest it was minimal.
 83 Additionally, several studies suggest ISOW was vigorous during MIC 3 (Duplessy et al.,
 84 1998b; Kissel et al., 1997; Kuijpers et al., 1998), while others suggest a variable ISOW due to
 85 higher-frequency Heinrich Events and Dansgaard/Oeschger Cycles (Rasmussen et al., 1996a;
 86 1996b; Kissel et al., 1999; Prins et al., 2001; 2002; Rasmussen and Thomsen, 2009). Because
 87 ISOW is a large contributor to NCW (Schmitz and McCartney, 1993; Kissel et al., 1997;
 88 Hansen and Osterhus, 2000), variations in ISOW strength have been proposed to exert a large
 89 control over NCW circulation patterns (Kuijpers et al., 1998).

90 Large-scale deepwater circulation patterns in the North Atlantic have been
 91 reconstructed using geochemical records from benthic foraminiferal Cd/Ca (Boyle and
 92 Keigwin, 1987), Zn/Ca (Marchitto et al., 2002), ϵ_{Nd} (Roberts et al., 2011; Yu et al., 2008;
 93 Crockett et al., 2011), and especially $\delta^{13}C$ (e.g., Boyle and Keigwin, 1987; Oppo and
 94 Fairbanks, 1987; Duplessy et al., 1988a; Sarnthein et al., 1994; Raymo et al., 1990; Oppo and
 95 Lehman, 1995; Oppo et al., 1995; Flower et al., 2000; Curry and Oppo, 2005; Olsen and
 96 Ninnemann, 2010). Modern NADW is nutrient depleted, and thus, has high $\delta^{13}C$ values (1 –
 97 1.5 ‰; Kroopnick et al., 1972; Kroopnick, 1985); modern NADW includes reintrained North
 98 Atlantic waters as well as contributions of ISOW, DSOW, and LSW, which have $\delta^{13}C$ values

of ~ 0.8 ‰, ~ 0.8 ‰ (Dokken et al., 2013; Bauch et al., 2001), and 0.8-1.2 ‰ (Winsor et al., 2012), respectively. In contrast, the nutrient enriched Southern Ocean is vertically well mixed due to the strong West Wind Drift, and thus has an isotopic composition of $\delta^{13}\text{C} \sim -0.4$ ‰ (Kroopnick, 1985). Deep Pacific Ocean $\delta^{13}\text{C}$ values are ~ 0.0 ‰, reflecting the long residence time of these waters (Kroopnick, 1985). As such, measured $\delta^{13}\text{C}$ values of biogenic calcite in the deep Southern Ocean reflect the relative inputs of deepwater masses from the Pacific, Indian, and Atlantic Oceans (Oppo and Fairbanks, 1987; Charles and Fairbanks, 1992; Lynch-Stieglitz, 2007).

During glacial-interglacial transitions of the late Pleistocene, benthic foraminiferal $\delta^{13}\text{C}$ studies, supported by other geochemical studies, determined that the large-scale patterns of deepwater circulation vary between glacial and interglacial climate states (Boyle and Keigwin, 1987; Oppo and Fairbanks, 1987; Duplessy et al., 1988a; Oppo and Lehman, 1995; Roberts et al., 2011; Gebbie, 2014). Cross sections of benthic foraminiferal $\delta^{13}\text{C}$ values of the Atlantic during the LGM (GEOSECS; Curry et al., 1988; Oppo and Horowitz, 2000; Venz and Hodell, 2002; Raymo et al., 2004; Curry and Oppo, 2005) show that a shoaling of NCW currents allowed for the northward intrusion of Antarctic Bottom Water (AABW) into the northern North Atlantic, up to 60°N . Studies have therefore suggested that the core of NCW shoaled by as much as ~ 2 km during the LGM (Sarthein et al., 1994; Curry et al., 1988; Duplessy et al., 1988a; Oppo and Lehman, 1995; Gebbie, 2014), coincident with a weakened and/or shallower ISOW (Duplessy et al., 1988b; Kissel et al., 1997; Yu et al., 2008).

The objective of this study was to determine variability in deep-water circulation patterns in the eastern North Atlantic over the past 200 kyr on orbital and millennial time-scales. Herein, new $\delta^{13}\text{C}$ records from sediment cores 11JPC and 3GGC, from the southern end of Gardar Drift, are presented for the past 200 ka (Figure 1a). New records from these two cores are combined with published records from seven other cores with a variety of

depths and sufficient data within the interval of interest to form our “Gardar Drift Transect” (Table 1; Figure 1). This transect provides a unique opportunity to track variations in the influence of ISOW. Transect cores, combined with cores from the South Atlantic (Ocean Drilling Program; ODP Site 1090) and mid- latitude North Atlantic (Deep Sea Drilling Program; DSDP Site 607), allow for the examination of southern sourced waters in the eastern northern North Atlantic in a similar method to the Atlantic cross sections (e.g., Curry and Oppo, 2005), but on a finer scale (Figure 1b). To examine the depth to which ISOW penetrated, benthic foraminiferal $\delta^{13}\text{C}$, a semi-conservative water mass tracer, was compared along a depth transect south of the Iceland-Scotland Ridge; with the occurrence of low $\delta^{13}\text{C}$ water on southern Gardar Drift (here taken to be southern sourced) being indicative of a shoaled ISOW.

3. Methods

3.1 Regional Setting

Gardar and Bjorn Drifts are key deposits for studying North Atlantic paleoceanography since they lie within the flow of modern-day ISOW (Figure 1a; e.g., Oppo and Lehman, 1995; Bianchi and McCave, 1999). As ISOW flows south along the Reykjanes Ridge, contourite drift deposits form along each edge of the current, with Bjorn Drift to the north and Gardar Drift to the South (Figure 1a; Davies and Loughton, 1972; Bianchi and McCave, 1999). Contourite drifts, like Gardar and Bjorn, are found throughout the North Atlantic where large quantities of sediments are ‘plastered’ against already existing bathymetric features (Hollister et al., 1978). These contourite drift deposits provide an excellent location for paleoceanographic reconstructions due to the high sedimentation rates (Figure 1a; e.g., McCave et al., 1980; McCave and Tucholke, 1986; Channell et al., 1997; Hall et al., 1998; Bianchi and McCave, 1999; Faugers et al., 1999; Praetorius et al., 2008;

Thornalley et al., 2010; 2011a; Elmore and Wright, 2011; Elmore et al., 2015). To utilize the paleoceanographic information in this archive, the study examined jumbo piston core, 11JPC (2707 m), and giant gravity core, 3GGC (3305 m; Table 1; Figure 1). Both cores were collected by the *R/V Knorr* from southern Gardar Drift on cruise 166, leg 14 in 2002.

3.2 Samples and Sedimentological Data Collection

To generate estimates of weight percent of carbonate (wt. % CaCO_3) and weight percent of coarse fraction (wt. % CF), we sampled cores 11JPC and 3GGC at ~ 5 cm intervals for the entire length of both cores (23.60 m for 11JPC and 1.84 m for 3GGC). Samples were split approximately in half and each half was dried overnight in a 50 °C oven. One half of each sample was weighed (Dry Wt. _{initial}) and combined with ~ 40 ml of 1.0 M acetic acid in a 50 ml centrifuge tube for 24 hours. Each sample was agitated during the process to ensure complete removal of carbonate from the sample. The vials were centrifuged for 1 minute to separate the sample from the acid, which was then decanted. Samples were reprocessed in acetic acid, and then rinsed with 40 ml of deionized water three times, centrifuging for one minute before each decanting. Samples were then oven-dried and weighed again (Dry Wt. _{-CaCO₃}). Assuming negligible opal content since biogenic silica was not observed during microscopic examination, the weight percent calcium carbonate (Wt. % CaCO_3) was then determined using the following equation:

$$[1] \quad \text{Wt. \% CaCO}_3 = (1 - \text{Dry Wt.}_{-\text{CaCO}_3} / \text{Dry Wt.}_{\text{initial}}) * 100$$

Weight percent coarse fraction (Wt. % CF), was determined by weighing the other half of the dried sample before (wt. _{unwashed}) and after (wt. _{washed}) being washed through a 63

µm sieve. Dry samples were soaked in a dilute calgon solution to prevent flocculation during washing. Wt. % CF was then calculated using the following equation:

$$[2] \quad \text{Wt. \% CF} = (\text{wt.}_{\text{washed}} / \text{wt.}_{\text{unwashed}}) * 100$$

3.3 Stable Isotopic Analysis

In order to assess the paleoceanographic changes over Gardar Drift since 200 ka, stable isotopic analyses of $\delta^{18}\text{O}$ were performed on benthic foraminifera to establish age control (e.g., Lisiecki and Raymo, 2005); and the simultaneous analysis of $\delta^{13}\text{C}$ was measured since epifaunal benthic foraminifera faithfully record changes in the $\delta^{13}\text{C}$ value of the bottom waters in which they live (e.g., Curry and Oppo, 2005). For this study, only *P. wuellerstorfi* tests were chosen for benthic foraminiferal analysis since some *Cibicidoides* taxa (e.g., *C. robertsoniensis*) do not record equilibrium values and may be up to 1 ‰ lower in $\delta^{13}\text{C}$ values in this region (Elmore, 2009). Planktic foraminifera were also analyzed for stable isotopic concentration to ensure that changes in surface productivity were not a controlling factor on benthic $\delta^{13}\text{C}$ values (Mackensen et al., 1993).

For stable isotopic analyses, ~ 15 tests of the planktic foraminifera, *Globigerina bulloides* and ~ 5 tests of benthic foraminifera, *Planulina wuellerstorfi*, were handpicked under a binocular microscope from the 250 – 350 µm size fraction of the washed samples. The samples were analyzed using a Micromass Optima Mass Spectrometer equipped with an automated Multiprep at the Rutgers University Stable Isotope Laboratory. Samples were reacted in phosphoric acid for 15 minutes at 90 °C. Measured values are reported using standard δ -notation and are compared to Vienna PeeDee Belemnite using an internal lab standard that is routinely calibrated with NBS-19 (1.95 ‰ $\delta^{13}\text{C}$, -2.20 ‰ $\delta^{18}\text{O}$; Coplen et al., 1983). The internal lab standard is offset from NBS-19 by 0.1 ‰ for $\delta^{13}\text{C}$ and 0.04 ‰ for

198 $\delta^{18}\text{O}$. The 1- σ precision of standards during analysis for this project was typically 0.05 ‰ for
 199 $\delta^{13}\text{C}$ and 0.09 ‰ for ‰ $\delta^{18}\text{O}$.

200 Differences between bottom water $\delta^{13}\text{C}$ values are the result of the waters' source
 201 region ($\delta^{13}\text{C}$ values are biologically-fractionated during primary productivity and undergo
 202 temperature-dependent fractionation during air-sea gas exchange; Lynch-Stieglitz et al.,
 203 1995); $\delta^{13}\text{C}$ values are also controlled by water mass ageing and productivity fluctuations
 204 above the waters' flow path. Additionally, benthic foraminiferal $\delta^{13}\text{C}$ values can be affected
 205 by organic carbon settling to the deep sea above the site, which is controlled by primary
 206 productivity and remineralization (Mackensen et al., 1993). Thus, benthic foraminiferal $\delta^{13}\text{C}$
 207 records at a particular site may also contain a localized signal relating to changing organic
 208 carbon flux and/or water mass remineralization properties. For southern Gardar Drift Sites for
 209 which data is presented here, there are no significant similarities between the planktic and
 210 benthic $\delta^{13}\text{C}$ records within each core (Figure 2, Figure 3), indicating that surface processes
 211 are not dominating the benthic foraminiferal $\delta^{13}\text{C}$ signal. Similar $\delta^{13}\text{C}$ values among all
 212 Gardar Drift sites (Figure 5c), especially during interglacials, provides additional confidence
 213 in the use of $\delta^{13}\text{C}$ as a reliable water mass tracer on Gardar Drift (Praetorius et al., 2008;
 214 Thornalley et al., 2010; 2011a; Elmore and Wright, 2011; Elmore et al., 2015). Thus, for the
 215 remainder of this manuscript our Gardar Drift benthic foraminiferal $\delta^{13}\text{C}$ data has been
 216 confidently used as a paleo-watermass tracer.

217

218 **3.4 Age Models**

219 The age model for the top 333 cm of 11JPC was constrained by 15 AMS ^{14}C ages
 220 (Table 2), which were all previously published in Elmore & Wright (2011) and Elmore et al.
 221 (2015). As previously published, the AMS samples were comprised of 4 - 6 mg of planktic
 222 foraminifera *Globogenerina bulloides* that were selected using a binocular microscope, ultra

sonificated in deionized water, and analyzed at the Keck Center for Accelerator Mass Spectrometry at the University of California, Irvine (Elmore and Wright, 2011). These published radiocarbon ages were then converted to calendar ages according to the Fairbanks0805 calibration, and a standard 400-year reservoir correction was applied (Fairbanks et al., 2005). Since the publication of these AMS ^{14}C dates, temporally evolving reservoir age corrections have been proposed for studies of cores from this region (Thornalley et al., 2011b; Stern and Lisiecki, 2013), application of constant correction/adjustment does not change the conclusions of this study because 1) the majority of our study interval (210-40 ka) is earlier than the utility of AMS ^{14}C dating (40 ka-present), and 2) the standard reservoir correction was also consistently applied to any AMS ^{14}C dates used in the previously published age models. We are not endeavoring to look at the high frequency variations among these cores and thus any relatively minor offsets in age models will not significantly affect the conclusions in this study. The published age model from Elmore and Wright (2011) revealed a ~ 18 kyr hiatus, constrained by 3 AMS ^{14}C dates from 227-282 cm (Table 2). Based on an abrupt color change in the core, the top of this hiatus was located at 222 cm (see supplement of Elmore and Wright, 2011). This section of sediment is interpreted to represent a mass transport event and thus data from this interval were not interpreted as part of this study. The previously published AMS ^{14}C date of ~ 33.81 ka at 333 cm is considered to be below the mass transport event because it conforms to the linear sedimentation rate defined by the remainder of chrono-stratigraphic tie points below the event (Figure 2E).

Age model information below 333 cm is not based on radiocarbon and is being presented here for the first time. The Lachamp Event (40 ka) was identified at 430 cm (H. Evans, personal communication; Table 2). Additional chrono-stratigraphic tie points were determined by comparing measured foraminiferal $\delta^{18}\text{O}$ values to a stacked, benthic

foraminiferal $\delta^{18}\text{O}$ record by Lisiecki and Raymo (2005; Table 2). The error associated with the Lisiecki and Raymo (2005) stack is reported to be ~ 2 kyr, though relative errors will be significantly lower at individual sites tied to the LR04 chronology. In the case of 11JPC, the benthic foraminiferal $\delta^{18}\text{O}_{P. wuellerstorfi}$ record shows a very strong agreement to the LR04 stack as well as to the other records in the region (Figure 4) and the age model shows nearly linear sedimentation rates through the interval of study (Figure 2E).

An age model for 3GGC was established based entirely on visual correlation among similar records (i.e., wiggle matching) rather than AMS ^{14}C dating (see Table 3), thus cores 3GGC and 11JPC are on a common age model. The record of benthic $\delta^{18}\text{O}$ from 3GGC (Figure 3a) was first visually compared to the Lisiecki and Raymo (2005) LR04 stack. A further, fine-tuning of the age model was completed by visual comparison of benthic and planktic $\delta^{18}\text{O}$, benthic $\delta^{13}\text{C}$, % CF, and % CaCO_3 from GGC 3 to the proximal core, 11JPC (Figure 3; Table 3). Since there are no available AMS ^{14}C dates, the error associated with this age model is proportionate to the error reported for the LR04 benthic stack of ~ 2 kyr.

As with the previously published age model from Elmore et al. (2015) and Elmore and Wright (2011) that constrains the ages for the top of 11JPC, all previously published benthic foraminiferal $\delta^{13}\text{C}$ records used in the construction of the Gardar Transect were kept on their originally published chronologies (Table 1). The selected cores each have substantial resolution to be used for comparison, with each published measured sample representing between 0.4 and 1.3 kyrs of sedimentation (Table 1). Thus these cores are ideally geographically located and capable of comprising a depth transect to trace watermass changes on Gardar Drift (Figure 1). To support the use of each chronology remaining as it was published, there is a remarkable similarity in trends and values of the $\delta^{18}\text{O}$ records from all sites investigated here, when the original benthic foraminiferal $\delta^{18}\text{O}$ data is compared (Figure 4). Since benthic foraminiferal $\delta^{18}\text{O}$ is largely controlled by extra-regional processes

(including continental ice volume), the similarity in records is a strong indication that the age models for these sites are comparable, and relatively minor offsets will not affect the conclusions of this study.

3.5 Transect Construction

To evaluate ISOW changes in the Late Pleistocene, we compiled the benthic foraminiferal $\delta^{13}\text{C}$ data from all available core locations with high-resolution data over the time period of interest (Table 1). The range in length (1600 km), range in water depth (1600 m), and confined geographic area of the sites within the Gardar Drift transect allows for the detection of ISOW penetration depth by monitoring variations in benthic foraminiferal $\delta^{13}\text{C}$ values (Figure 1). Figure 5C shows our new $\delta^{13}\text{C}_{P. wuellerstorfi}$ data, as well as all previously published (i.e., raw) data from the cores used to form the Gardar transect, on their original age models. To assess this variability in $\delta^{13}\text{C}$ gradients on Gardar Drift, data from each of the sites in the Gardar transect were treated in two ways. First, benthic foraminiferal $\delta^{13}\text{C}$ data from all sites in the Gardar region from 0-200 ka (Figure 5C) were smoothed using a Gaussian fit with a span of 2 kyrs. This provided an average benthic foraminiferal $\delta^{13}\text{C}$ value. Uncertainties for this average were calculated by using a Monte Carlo simulation, where the external reproducibility of the data was taken into account. This was performed by perturbing the $\delta^{13}\text{C}$ data over a large sample number ($n=1,000$), and summing the resultant records before smoothing (as above) over a 2-kyr window. All sites were weighted equally (where they contained data within the sampling window; i.e. gaps of > 2 kyr are omitted) to provide an unbiased final compilation. The standard deviation from the mean was then calculated along the timeseries (Figure 5E).

Secondly, Gardar Drift core sites were divided in to North and South regions (Figure 1) to examine watermass changes along the drift crest. For the Northern Gardar Transect, a

composite record was generated for northern sites, ODP site 984, ODP site 983, EW9302 JPC8, and V29-202. The weighted mean for Northern Gardar and 95% confidence intervals are presented along with the standard deviations from the mean, calculated as above (Figure 5F). A composite record for the sites on Southern Gardar, 11JPC, 3GGC, and Neap 18k, was determined using the same method (Figure 1). The difference between the North Gardar transect benthic foraminiferal $\delta^{13}\text{C}$ record and the South Gardar transect $\delta^{13}\text{C}$ records was then determined, utilizing the full uncertainty characterized by the composites (Figure 5G). All Monte Carlo statistics were carried out in the R statistical analysis program (R Development Core Team, 2010).

Time-series analyses were completed on the north-south difference and standard deviation of the benthic foraminiferal $\delta^{13}\text{C}$ transect data using the AnalySeries 2.0.4.2 program (Paillard et al., 1996) using a span of 0.01 to identify recurring periodicities (Figure 6). Spectral analyses were completed using the Blackman-Tukey method in the same software (Blackman and Tukey, 1958).

The $\delta^{13}\text{C}$ value of the Southern Component Water (SCW) end member was determined by using ODP Site 1090 from the Atlantic sector of the Southern Ocean (Table 1). As an additional monitor of the intrusion of SCW into the North Atlantic, benthic foraminiferal $\delta^{13}\text{C}$ data was included from ODP Site 607, located below the subtropical North Atlantic (Ruddiman et al., 1989; Raymo et al., 1989). Core locations, water depths, and age model information for each site are provided in Table 1.

4. Results

4.1 Core KN166-14 11JPC

Marine Isotope Chronozones (MIC) 1 through 9a were defined in core 11JPC using the planktic and benthic oxygen isotope records, which show the typical saw-toothed pattern

that characterized the late Pleistocene benthic foraminiferal oxygen isotope curves (Lisiecki and Raymo, 2005; Figure 4). Over this interval, the $\delta^{18}\text{O}$ values range from 0.8 to 3.8 ‰ for *G. bulloides*, and from 2.5 to 4.5 ‰ for *P. wuellerstorfi* (Figure 2). Sharp decreases in $\delta^{18}\text{O}$ are recorded at transitions from glacial to interglacial chronozones (MIC 8 – 7.5, MIC 6 – 5e; Figure 2).

Benthic foraminiferal $\delta^{13}\text{C}$ records from core 11JPC record orbital-scale variability, with values from -0.3 to 1.5 ‰ for *P. wuellerstorfi* (Figure 2). The lowest values in benthic $\delta^{13}\text{C}$ are recorded during MIC 8, 6, 4, and 3 (Figure 2). Values of benthic $\delta^{13}\text{C}$ are highly variable during MIC 3, likely due to the high frequency Heinrich or Dansgaard/Oeschger Events (Figure 2).

Through the interval of study, the record of Wt. % CF varies on glacial-interglacial timescales with values ranging from 0 – 35 % for core 11JPC (Figure 2). Low Wt. % CF (< 15 %) values are recorded during interglacial MICs 7.5, 7.3-7.1, 5e-5c, 5a, and 1 (Figure 2). The highest Wt. % CF values are recorded from 1700 to 1550 cm, corresponding to MIC 6 (Figure 2). The observed variability in Wt. % CF values indicates either: an increase in fine particles during interglacials, an increase of large particles during glacials, or both (Figure 2); winnowing or dissolution could also effect the Wt. % CF record, however foraminifera appear visually well preserved. Increased ice rafted detritus (IRD) abundances during glacial periods are a well-documented feature of North Atlantic sediment records (Ruddiman, 1977; Bond and Lotti, 1995; McManus et al., 1999; Venz et al., 1999; Andrews, 2000), and have been documented during the cold Younger Dryas from this sediment core (Elmore and Wright, 2011); this suggests that higher glacial Wt. % CF is mainly due to a glacial increase in IRD (Figure 2). The MIC 3 section is characterized by Wt. % CF values that are highly variable and range from 0 to 25 %; this is likely caused by increased IRD during Heinrich Events (Bond and Lotti, 1995; Figure 2).

Weight percent carbonate also varies on glacial-interglacial time-scales in core 11JPC (Figure 2), as has been shown in other North Atlantic cores (Ruddiman et al., 1987; Ortiz et al., 1999). Low Wt. % CaCO_3 values (20 to 30 %) are recorded in the sections from 2200 to 2100, 1900 to 1800, 1600 to 1500 and 500 to 300 cm (Figure 2). These low % CaCO_3 values are coincident with elevated % CF suggesting that the record of % CaCO_3 is determined mainly by the dilution of carbonate by larger IRD during glacial periods (Figure 2). Highest values of % CaCO_3 are recorded in 11JPC sediments from 2100 to 2050, 1500 to 1450, 1275 to 1225, 950 to 800, and 150 to 0 cm (Figure 2). These zones of high % CaCO_3 correspond to interglacial periods with high productivity (Figure 2).

4.2 Core KN166-14 3GGC

According to the age model described above, core 3GGC can be subdivided into chronozones of the upper MIC 3, Last Glacial Maximum (LGM), Bolling/Allerod, Younger Dryas, and Holocene (Figure 3). Oxygen isotope values for *G. bulloides* and *P. wuellerstorfi* are highest (~ 3.5 and 4.5 ‰, respectively) in the LGM section (~ 25 to 15 ka; Figure 3). Termination 1 is recorded by a decrease in the $\delta^{18}\text{O}$ values in all species, marking the Bolling/Allerod section. Increasing $\delta^{18}\text{O}$ values reflect the cooler conditions during the Younger Dryas section (~ 13.1 to 11.5 ka; Figure 3). The lowest $\delta^{18}\text{O}$ values in 3GGC represent the Holocene section (~ 11.5 to 0 ka; Figure 3).

Benthic and planktic $\delta^{13}\text{C}$ values decrease in the upper MIC3 and LGM sections for *G. bulloides* and *P. wuellerstorfi* (from 0.0 to -1.0 and from 1.2 to 0.4 ‰, respectively); values of each species are similar in these sections (Figure 3). *P. wuellerstorfi* $\delta^{13}\text{C}$ values increase in the Bolling/Allerod, Younger Dryas, and lower Holocene to values of 1.0 and 0.5 ‰, respectively (Figure 3). Unlike *P. wuellerstorfi*, *G. bulloides* $\delta^{13}\text{C}$ values decrease in the Bolling/Allerod, Younger Dryas, and lower Holocene to a value of -1.0 (Figure 3).

The record of Wt. % CF is variable for core 3GGC, with values ranging from 0 to 35 % (Figure 3). Low Wt. % CF (< 10 %) is observed in the upper MIC 3 and throughout MIC 1 (Figure 3). Higher Wt. % CF (> 20 %) is observed during the LGM, likely due to an increase in IRD (Figure 3; McManus et al., 1999).

4.3 Transect Results

A comparison of benthic foraminiferal $\delta^{13}\text{C}$ records from sites downstream of the Iceland-Scotland Ridge reveals a large degree of variability, indicating changes in watermasses (Figure 5c). The sites in the southern portion of the transect show greater variability than those in the northern part (Figure 5c; Figure 5f). This is reflective of the greater influence of SCW, with low $\delta^{13}\text{C}$ values, recorded from sites that are more distal from the Iceland-Scotland Ridge (Figure 5d). Lower variability in $\delta^{13}\text{C}$ values in the northern sites, or sites proximal to the ridge, indicates that they were almost always bathed by the same watermass with unchanging composition, which is likely ISOW (Figure 5c). The difference in benthic foraminiferal $\delta^{13}\text{C}$ values between all transect sites is minimal (< 0.5 ‰) during interglacials, MIC 5 and the Holocene, indicating that all transect sites were bathed by a similar water mass during interglacial periods (Figure 5c, 5d). The benthic foraminiferal $\delta^{13}\text{C}$ values of the northern transect compilation and southern transect compilation are also similar during interglacial periods, also indicating that all of the transect sites were bathed by a similar watermass, likely ISOW (Figure 5f).

The standard deviation from the ensemble mean of all benthic foraminiferal $\delta^{13}\text{C}$ data is less than 0.25 ‰ during interglacial periods (therefore within analytical reproducibility), providing statistical representations of the small gradient in benthic foraminiferal $\delta^{13}\text{C}$ values within the transect sites (Figure 5e). The difference between the northern and southern transect composite benthic foraminiferal $\delta^{13}\text{C}$ records is also lower during interglacials (< 0.2

398 ‰) indicating a small gradient in benthic foraminiferal $\delta^{13}\text{C}$ (Figure 5f, 5g). The small
 399 increase in standard deviation values in the earliest portion MIC 5e is likely due to
 400 differences within the age models for individual records over the abrupt transition from MIC
 401 6 to 5e (Figure 5e). Since the northernmost cores used in this comparison (ODP site 984 and
 402 ODP site 983) lie northward of the maximum possible extent of AABW (Curry and Oppo,
 403 2005), they should always record the $\delta^{13}\text{C}$ signature of a northern source and can be
 404 approximated as the northern end member (Raymo et al., 1989). The low variability in the
 405 benthic foraminiferal $\delta^{13}\text{C}$ records during the interglacial periods thus indicates that all sites
 406 are recording a northern sourced watermass during interglacial periods (Figure 5e).
 407 Additionally, while benthic foraminiferal $\delta^{13}\text{C}$ values at DSDP site 607 are very similar to
 408 southern Gardar values during interglacial periods, benthic foraminiferal $\delta^{13}\text{C}$ values from
 409 the South Atlantic site (ODP site 1090) are significantly lower, suggesting that southern
 410 sourced waters were not prevalent in the northern North Atlantic during these periods (Figure
 411 5f). Thus, a small benthic foraminiferal $\delta^{13}\text{C}$ gradient during MIC 5e and the Holocene
 412 indicates that a deeply penetrating ISOW bathed all sites on Gardar Drift (Figure 5g).

413 The gradient in benthic foraminiferal $\delta^{13}\text{C}$ values between the transect sites is large
 414 (up to ~ 1.5 ‰) during glacial periods MIC 6, 4, and 2; this suggests that different water
 415 masses bathed the north and south regions of this transect during each of these glacial periods
 416 (Figure 5f; Figure 5g). The standard deviation (an indication of data spread) of the benthic
 417 foraminiferal $\delta^{13}\text{C}$ data shows higher values during MIC 6, 4 and the LGM (> 0.25 ‰)
 418 relative to the spread in MIC 5e and 1; this shows that the southern portion of the transect is
 419 bathed by a different watermass than the northern portion. The difference between the
 420 northern and southern transect composite benthic foraminiferal $\delta^{13}\text{C}$ records are both also
 421 larger during these glacial periods, suggesting two distinct water masses (Figure 5g). During
 422 MIC 6, 4 and the LGM, benthic foraminiferal $\delta^{13}\text{C}$ values from southern Gardar Drift are

more similar to values at DSDP 607 and ODP 1090 than the benthic foraminiferal $\delta^{13}\text{C}$ values at northernmost transect sites, indicating that SCW waters invaded the North Atlantic, reaching the southern flanks of Gardar Drift (Figure 5f; Figure 5g). Because of the location of this transect, the intrusion of SCW onto southern Gardar Drift is evidence for the shoaling of ISOW during glacial periods.

The warmer substages of MIC 5, 5e, 5c, and 5a, are characterized by low ($< 0.5 \text{ ‰}$) benthic foraminiferal $\delta^{13}\text{C}$ variability on Gardar Drift (Figure 5a; Figure 5b). The standard deviation of the benthic foraminiferal $\delta^{13}\text{C}$ during MIC 5 is generally $< 0.25 \text{ ‰}$ (Figure 5e). The difference between the northern and southern composite benthic foraminiferal $\delta^{13}\text{C}$ records is also lower during MIC 5e, 5c, 5a (Figure 5f, 5g). The gradient is slightly larger during the cooler substages of MIC 5, MIC 5d and 5b ($\sim 0.5 \text{ ‰}$; Figure 5g). Increased difference between the northern and southern composite benthic foraminiferal $\delta^{13}\text{C}$ records and increased standard deviation of benthic foraminiferal $\delta^{13}\text{C}$ values during MIC 5d and 5b signify a greater range in benthic foraminiferal $\delta^{13}\text{C}$ values within the transect (Figure 5g). This could suggest that ISOW was shallower during MIC 5d and 5b than during the other sub-stages of MIC 5 (Figure 5f; Figure 5g). However, MIC 5d and 5b are characterized by ISOW that was only slightly shoaled.

Some higher frequency variability is superimposed on the longer-term trends during MIC 4 and 3 in core 11JPC records of % CF, *G. bulloides* $\delta^{18}\text{O}$, *P. wuellerstorfi* $\delta^{13}\text{C}$, and % CaCO_3 (Figure 2). Percent coarse fraction decreased from 70 to 60 ka and reached minimum values at ~ 58 ka, followed by a general increase from 58 to 35 ka (Figure 2C). Percent Carbonate and *P. wuellerstorfi* $\delta^{13}\text{C}$ records increased from 70 to 60 ka and reached maximum values at ~ 58 ka, followed by a general decrease in value from 58 to 35 ka (Figure 2d, 2b). The period from 70 to 60 ka was characterized by a general decrease in the standard deviation of the benthic foraminiferal $\delta^{13}\text{C}$ values in the transect, followed by an increase in

standard deviation from 58 to 35 ka, suggesting a shoaling of ISOW (Figure 5g). *Globigerina bulloides* $\delta^{18}\text{O}$ values became increasingly variable during the period from 58 to 35 ka, suggesting changing surface water conditions that are attributed to increased freshwater inputs (Figure 2a). Together, these records indicate an overarching change in circulation and climate whereby a general decrease in ice rafting and freshwater input leads to deepening of ISOW from 70 to 60 ka, and a general increase in ice rafting is linked to a shoaling ISOW from 58 to 35 ka (Figure 2, 5).

Spectral analysis of the standard deviation of the benthic foraminiferal $\delta^{13}\text{C}$ values for the transect sites, which indicates the size of the gradient between northern and southern transect sites, reveals a strong precessional signal (Figure 6). The precessional control on the north-south gradient indicates that the depth of ISOW penetration is paced by this orbital forcing (Figure 6).

5. Discussion- Orbital Scale Variability in ISOW Penetration Depth

Variability in NCW formation is often linked to changes in insolation (Figure 5b), known as Milankovich Cycles (Hays, 1976; Imbrie and Imbrie, 1980; Imbrie et al., 1993; Raymo, 1997; Raymo et al., 2004; Lisiecki and Raymo, 2005). Between 3 and 1 Ma, the 41-kyr obliquity cycles dominated benthic foraminiferal $\delta^{13}\text{C}$ records (Raymo et al., 1989; 1997; Shackleton et al. 1990; Lisiecki and Raymo, 2005). Following the Mid-Pleistocene Climate Transition, the 100-kyr eccentricity cycle dominated ice volume trends through the late Pleistocene (Hays, 1976; Raymo et al., 2004; Lisiecki et al., 2008). Northern Component Water strength has been shown to follow these orbital-scale climate changes (e.g., Raymo et al., 2004). The depth of ISOW penetration also follows orbital-scale climate cycles with a strong 100-kyr periodicity (Figure 5g, 6; Duplessy et al., 1988a; Kissel et al., 1997).

In addition to the longer eccentricity and obliquity cycles, the shorter ~ 23 -kyr

precession cycle is also visible in climate records (Figure 6), most notably through the sub-stages of MIC 5 (Raymo et al., 2004), as well as in the Mediterranean sapropels (Hilgen, 1991); however, precession is considered to be a weak driver of large-scale ice volume changes (Lisiecki et al., 2008). Precession-driven peaks in high-latitude northern hemisphere insolation can be seen at ~ 50 and ~ 60 ka (Figure 5b; Laskar et al., 2004); however, previous studies of NCW strength do not show corresponding changes during MIC 3 (Raymo et al., 2004). Similarly, insolation peaks at ~ 150 and ~ 170 ka caused by the precessional forcing occur during MIC 6, which is generally presumed to be a prolonged glacial period of decreased NCW formation (Raymo et al., 2004).

Our benthic foraminiferal $\delta^{13}\text{C}$ records (Figure 5d; Figure 5f) indicate that ISOW penetrated to deeper water depths during interglacial periods, consistent with modern observations (Worthington, 1976) and other paleoceanographic reconstructions (Duplessy et al., 1988a; 1988b; Kissel et al., 1997). During glacial periods, ISOW shoaled according to our records (Figure 5f; Figure 5g), consistent with proxy evidence for the intrusion of AABW into the northern North Atlantic (Boyle and Keigwin, 1987; Curry et al., 1988; Duplessy et al., 1988; McManus et al., 1999; Marchitto et al., 2002; Curry and Oppo, 2005). This suggests that the general patterns of ISOW variability are driven by the 100-kyr eccentricity forcing. However, the variability in the benthic foraminiferal $\delta^{13}\text{C}$ values from the sites within the Gardar Drift transect also indicates that ISOW depth penetration varied in concert with precessional cycles (Figure 5g; Figure 6). These cycles are easily seen during the precession minima at ~ 185 and ~ 145 ka, during MIC 6, which are both associated with increased benthic foraminiferal $\delta^{13}\text{C}$ gradients from the standard deviation data, indicating shallower ISOW due to decreased insolation (Figure 5g). These results are in opposition to those of Kuijpers et al. (1998) who suggested that ISOW formation was enhanced during MIC 6. Additionally, the precessionally driven insolation minima at ~ 70 and ~ 25 ka, and

slight minima at ~ 45 ka, have corresponding maxima in benthic foraminiferal $\delta^{13}\text{C}$ gradients (Figure 5g; Figure 6).

The mechanism that links precession cycles and ISOW strength must not be strictly temperature dependent since neither ice core (Alley, 2004) nor paleoceanographic temperature records show a pronounced precessional signal (Raymo et al., 2004; Liseicki et al., 2009). However, we propose that, high northern latitude summer insolation minima, driven mainly by precession cycles, led to increased low-elevation glaciers and sea-ice, which in turn provided the fresh meltwater necessary to impede convection in the Nordic Seas, and thus hindered the formation of ISOW. Additionally, the precessional forcing strongly affects low-latitude climate variations (McIntyre and Molino, 1996) and thus precessionally-driven changes in the tropical carbon cycle could be driving the precessional variability seen in the Gardar Drift benthic foraminiferal $\delta^{13}\text{C}$ records.

A strong precessional control on ISOW strength (Figure 6) is especially interesting considering that ISOW is a large contributor to NCW (Worthington, 1976), and precession is not considered to exert a strong control on NCW production strength (Raymo et al., 2004; Lisiecki et al., 2008). This apparent difference may be due to the fact that the Gardar Drift transect is in a confined, gateway location that is sensitive to even subtle changes in NCW. One possible explanation for the apparent disagreement between ISOW variations and NCW variations is that many NCW strength records do not have the temporal resolution to decipher precession-scale variability. Another possible explanation is that other NCW contributors compensate for decreased ISOW production during some precession minima; such orbital-scale variability has been seen in records of Denmark Straits Overflow Water (e.g., Fagel et al., 2004; Milla et al., 2006) and Labrador Sea Water (Hillaire-Marcel et al., 1994).

Our new benthic foraminiferal $\delta^{13}\text{C}$ data examines glacial-interglacial changes in the depth penetration of ISOW along the Gardar Drift, with interglacial ISOW penetrating more

deeply than ISOW during glacial periods (Figure 5). This may suggest a changing influence of ISOW to NADW during large climate transitions in the Late Pleistocene. Although $\delta^{13}\text{C}$ data is sometimes ambiguous, our carbon isotopic data provide an important constraint for a change in ISOW penetration depth since $\delta^{13}\text{C}$ is the only proxy that can currently provide the required data density to examine ISOW depth in 3 dimensions. Despite the limitations of $\delta^{13}\text{C}$, we have identified that ISOW was shallower during glacial intervals; since ISOW is a large contributor to NADW, this agrees with reconstructions that glacial NADW was shallower, weaker, or non-existent (e.g., Curry and Oppo, 2005; Thornalley et al., 2011a; Stern and Lisiecki, 2013; Flower et al., 2000; Lynch-Stieglitz et al., 2007). Indeed, there is a strong agreement between our Southern Gardar $\delta^{13}\text{C}$ record and the intermediate depth North Atlantic record from Lisiecki et al. (2008), suggesting that our records may have a wider importance to the North Atlantic (Figure 5D). Our data also shows SCW infringement on Southern Gardar Drift, which is in agreement with numerous reconstructions from the North Atlantic that demonstrate southern sourced water infringement using a variety of proxies from Pa/Th to Cd/Ca and neodymium isotopes (e.g., Marchitto et al., 2002; Wilson et al., 2014; Alvarez Zarikian et al., 2009; McManus et al., 1999). While our $\delta^{13}\text{C}$ data can only identify similar watermasses, and thus is not a direct link to ISOW volume or flow strength, sedimentological proxies in the region have suggested that glacial ISOW was weaker according to sediment magnetic properties (Kissel et al., 1999) or stronger according to grain-size and Sr/Nd isotopes. Both stronger and weaker ISOW strength can be reconciled with the shallower ISOW that we report.

6. Conclusions

Benthic foraminiferal $\delta^{13}\text{C}$ records from a Gardar Drift transect reveal that the depth of ISOW penetration varied on orbital time-scales. Similar benthic foraminiferal $\delta^{13}\text{C}$ values

were recorded along our transect during MIC 5e and 1, indicating that ISOW was deeply penetrating during interglacial periods (Figure 5d; Figure 5f). During MIC 6, 4, and 2, ISOW shoaled significantly, allowing the intrusion of SCW into the southern region of Gardar Drift (Figure 5g; Figure 6). This relationship is consistent with previous studies linking NCW variation with the 100-kyr climate cycles of the late Pleistocene (e.g., Hays, 1976; Raymo et al., 1990; 2004). However, the Gardar Drift transect reveals a strong imprint of the precession component of insolation on ISOW production, particularly during intermediate climate states, such as MIC 3, and during the glacial states of MIC 6 and 4 (Figure 5d; Figure 6). The expression of precession in MIC 5 is strong in the climate signal (*c.f.*, benthic foraminiferal $\delta^{18}\text{O}$ changes from MIC 5a to 5e) but muted in the $\delta^{13}\text{C}$ gradient on Gardar Drift. This suggests that, during the substages of MIC 5: 1) ISOW production remained deeply penetrating, such that the mixing zone between NCW and SCW lay south of Gardar, and therefore Gardar Drift was insensitive to precession-driven changes; or 2) the smaller ice sheets in the northern hemisphere did not contribute substantial meltwater to the Norwegian Sea, and therefore could not affect ISOW depth through either flux or density variations. On shorter time-scales, high-frequency changes in ISOW penetration depth are also linked to freshwater inputs, likely due to Heinrich or Dansgaard-Oeschger Events.

Acknowledgements

We thank J. McManus, K. Miller, Y. Rosenthal, G. Mountain, and D. Thornalley for comments on earlier versions of this manuscript, K. Allen, L. Neitzke, S. Henderson, and R. Mortlock for helpful discussions, H. Evans and J. Channell for magnetostratigraphy, and J. Southon for AMS ^{14}C analytical assistance. The manuscript was significantly improved by comments from the Editor and two anonymous reviewers. This research was supported by National Science Foundation grant OCE-0095219 to J.D. Wright.

573 **References**

- 574 Alley, R.B.. 2004. GISP2 ice core temperature and accumulation data. *IGBP PAGES/World*
 575 *Data Center for Paleoclimatology Data Contribution Series*, 13.
 576
- 577 Alvarez Zarikian, C.A., A.Y. Stepanova, and Jens Grutzner. 2009. Glacial-interglacial
 578 variability in deep sea ostracod assemblage composition at IODP Site U1314 in the
 579 subpolar North Atlantic. *Marine Geology*. 258 (1-4); 69-87.
 580
- 581 Andrews, J.T.. 2000. Icebergs and iceberg rafted detritus (IRD) in the North Atlantic: facts
 582 and assumptions. *Oceanography*. 13 (3); 100-108.
 583
- 584 Bauch, H.A., Erlenkeuser, H., Spielhagen, R.F., Struck, U., Matthiessen, J., Thiede, J.,
 585 Heinemeier, J., 2001. A multiproxy reconstruction of the evolution of deep and
 586 surface waters in the subarctic Nordic seas over the last 30,000 yr. *Quat. Sci. Rev.* 20,
 587 659–678.
 588
- 589 Bianchi, G.G. and N. McCave, 1999. Holocene periodicity in North Atlantic climate and
 590 deep-ocean flow south of Iceland. *Nature*. 397; 515-517.
 591
- 592 Blackman, R. B. and J. W. Tukey. 1958. The Measurement of Power Spectra From the Point
 593 of View of Communication Engineering. Dover Publications, New York. 190 pp.
 594
- 595 Bond, G.C. and R. Lotti. 1995. Iceberg Discharges into the North Atlantic on Millennial
 596 Time Scales During the Last Glaciation. *Science*. 267; 1005-1010
 597
- 598 Boyle, E.A. and L.D. Keigwin, 1987. North Atlantic thermohaline circulation during the past
 599 20,000 years linked to high-latitude surface temperature. *Nature*. 330; 35-40.
 600
- 601 Broecker, W.S., Peteet, D.M., and D. Rind. 1985. Does the ocean-atmosphere system have
 602 more than one stable mode of operation? *Nature*. 315; 21-26.
 603
- 604 Broecker, W.S. and G.H. Denton. 1989. The role of ocean-atmosphere reorganizations in
 605 glacial cycles. *Geochimica et Cosmochimica Acta*. 53; 2465-2501.
 606
- 607 Chapman, M.R. and N.J. Shackleton. 1999. Global ice-volume fluctuations, North Atlantic
 608 ice-rafting events, and deep-ocean circulation changes between 130 and 70 ka.
 609 *Geology*. 27 (9); 795-798.
 610
- 611 Charles, C.D. and R.G. Fairbanks. 1992. Evidence from Southern Ocean sediments for the
 612 effect of North Atlantic deep-water flux on climate. *Nature*. 355; 416-419.
 613
- 614 Channell, J.E.T., Hodell, D.A., and B. Lehman. 1997. Relative geomagnetic paleointensity
 615 and $\delta^{18}\text{O}$ at ODP Site 983 (Gardar Drift, North Atlantic) since 350ka. *Earth and*
 616 *Planetary Science Letters*. 153; 103-118.
 617
- 618 Coplen, T.B., Kendall, C., and J. Hopple. 1983. Comparison of stable isotope reference
 619 samples. *Nature*. 302; 236-238.
 620

- Crocket, K.C., D. Vance, M. Gutjahr, G.L. Foster, and D.A. Richards. 2011. Persistent Nordic deep-water overflow to the glacial North Atlantic. *Geology*. 39; 515-518.
- Curry, W.B., Duplessy, J.C., Labeyrie, L.D., and N.J. Shackleton. 1988. Changes in the distribution of $\delta^{13}\text{C}$ of deep water ΣCO_2 between the last glaciation and the Holocene. *Paleoceanography*. 3(3); 317-341.
- Curry, W.B. and D.W. Oppo. 2005. Glacial water mass geometry and the distribution of $\delta^{13}\text{C}$ and ΣCO_2 in the western Atlantic Ocean. *Paleoceanography*. 20; 1-12.
- Davies, T.A. and A.S. Loughton. 1972. Sedimentary processes in the North Atlantic. In: Loughton, A.S., Berggren, W.A. et al. (Eds.), Initial Reports of Deep Sea Drilling Project, Vol. 12. U.S. Government Printing Office, Washington, D.C., pp. 905-934.
- Dickson, R.R., Gmitrowicz, E.M., and A.J. Watson. 1990. Deep water renewal in the northern North Atlantic. *Nature*. 344; 848-850.
- Dokken, T. and M. Hald. 1996. Rapid climatic shifts during isotopes stages 2-4 in the Polar North Atlantic. *Geology*. 24 (7); 599-602.
- Dokken, T. M., K. H. Nisancioglu, C. Li, D. S. Battisti, and C. Kissel (2013), Dansgaard-Oeschger cycles: Interactions between ocean and sea ice intrinsic to the Nordic seas, *Paleoceanography*, 28, 491-502, doi:10.1002/palo.20042.
- Duplessy, J.C., Shackleton, N.J., Fairbanks, R.G., Labeyrie, L., Oppo, D.W., and N. Kallel. 1988a. Deepwater source variations during the last climatic cycle and their impact on the global deep-water circulation. *Paleoceanography*. 3(3); 342-360.
- Duplessy, J.C., Labeyrie, L., and P.L. Blanc. 1988b. Norwegian Sea Deep Water Variations Over the Last Climatic Cycle: Paleo-oceanographical Implications. *Long and Short Term Variability of Climate*. 16; 86-116.
- Elmore, A.C. and J.D. Wright. 2011. North Atlantic Deep Water and climate variability during the Younger Dryas cold period. *Geology*. 39 (2); 107-110.
- Elmore, A.C., Wright, J.D., and J. Southon, 2015. Continued meltwater influence on North Atlantic Deep Water instabilities during the early Holocene. *Marine Geology*. 360; 17-24.
- Fagel, N., Innocent, C., Gariépy, C. and C. Hillaire-Marcel. 2002. Sources of Labrador Sea sediments since the last glacial maximum inferred from Nd-Pb isotopes. *Geochimica et Cosmochimica Acta*. 66 (14); 2569-2581.
- Fairbanks, R.G., Mortlock, R.A., Chiu, T.-C., Cao, L., Kaplan, A., Guilderson, T.P., Fairbanks, T.W., Bloom, A.L., Grootes, P.M., and M.-J. Nadeau. 2005. Radiocarbon calibration curve spanning 0 to 50,000 years BP based on paired ^{230}Th , $^{234}\text{U}/^{238}\text{U}$ and ^{14}C dates on pristine corals. *Quaternary Science Reviews*. 24; 1781-1796.
- Faugers, J.C., Stow, D.A.C., Imbert, P., and A. Viana. 1999. Seismic features diagnostic of contourite drifts. *Marine Geology*. 162; 1-38.

- Flower, B.P., Oppo, D.W., McManus, J.F., Venz, K.A., Hodell, D.A., and J.L. Cullen. 2000. North Atlantic intermediate to deep water circulation and chemical stratification during the past 1 Myr. *Paleoceanography*. 15; 388-403.
- Gebbie, G. 2014. How much did Glacial North Atlantic Water shoal? *Paleoceanography*. 29 (3); 190-209.
- Hall, I.R., McCave, I.N., Chapman, M.R., and N.J. Shackleton. 1998. Coherent deep flow variation in the Iceland and American basins during the last interglacial. *Earth and Planetary Science Letters*. 164; 15-21.
- Hansen, B. and S. Osterhus. 2000. North Atlantic-Nordic Seas exchange. *Progress in Oceanography*. 45 (2); 109-208.
- Hatun, H., Sando, A.B., Drang, H., and M. Bentsen. 2004. Seasonal to Decadal Temperature Variations in the Faroe-Shetland Inflow Water. *Geophysical Monograph- American Geophysical Union*.
- Hays, J.D., Imbrie, J., and N.J. Shackleton. 1976. Variations in the Earth's Orbit: Pacemaker of the Ice Ages. *Science*. 194 (4270); 1121-1134.
- Hilgen, F.J.. 1991. Astronomical calibration of Gauss to Matuyama sapropels in the Mediterranean and implication for the Geomagnetic Polarity Time Scale. *Earth and Planetary Science Letters*. 104; 226-244.
- Hillaire-Marcel, C., A. de Vernal, M. Lucotte, and A. Mucci. 1994. The Labrador Sea during the late Quaternary. *Canadian Journal of Earth Sciences*. 31; 1-4.
- Hollister, C.D., Flood, R.D., and I.N. McCave. 1978. Plastering and decorating in the North Atlantic. *Oceanus*. 21 (1); 5-13.
- Imbrie, J. and J.Z. Imbrie. 1980. Modeling the Climatic Response to Orbital Variations. *Science*. 207 (4434); 943-953.
- Imbrie, J., A., Berger, E.A. Boyle, S.C. Clemens, A. Duffy, W.R. Howard, G. Kulka, J. Kitzbach, D.G. Martinon, A. McIntyre, A.C. Mix, B. Molino, J.J. Morley, L.C. Peterson, N.G. Pisias, W.L. Prell, M.E. Raymo, N.J. Shackleton, and J. R. Toggweiler. 1993. On the structure and origin of major glaciation cycles 2. The 100,000-year cycle. *Paleoceanography*. 8 (6); 699-736.
- Jansen, E., Raymo, M.E., Blum, P., et al., 1996. Site 983. In *Proceedings of Ocean Drilling Program, Initial Reports*. 162; 139-167.
- Kissel, C., Laj, C., Lehman, B., Labyrie, L., and V. Bout-Roumazeilles. 1997. Changes in the strength of the Iceland-Scotland Overflow Water in the last 200,000 years: Evidence from magnetic anisotropy analysis of core SU90-33. *Earth and Planetary Science Letters*. 152; 25-36.

- Kissel, C., Laj, C., Labeyrie, L., Dokken, T., Voelker, A., and D. Blamart. 1999. Rapid climatic variations during marine isotopic stage 3: magnetic analysis of sediments from Nordic Seas and North Atlantic. *Earth and Planetary Science Letters*. 171; 489-502.
- Kleiven, H.F., Jansen, E., Curry, W.B., Hodell, D.A., and K. Venz. 2003. Atlantic Ocean thermohaline circulation changes on orbital to sub-orbital timescales during the mid-Pleistocene. *Paleoceanography*, 18 (1); 1-13.
- Kohl, A., D. Stammer, and B. Cornuelle. 2007. Interannual to Decadal Changes in the ECCO Global Synthesis. *Journal of Physical Oceanography*. 27; 313-337.
- Kroopnick, P., R.F. Weiss, and H. Craig. 1972. Total CO₂, ¹³C, and dissolved oxygen-¹⁸O at GEOSECS II in the North Atlantic. *Earth and Planetary Science Letters*. 16; 103-110.
- Kroopnick, P.. 1985. The Distribution of ¹³C in the Atlantic Ocean. *Earth and Planetary Science Letters*. 49; 469-484.
- Kuijpers, A., Troelstra, S.R., Wisse, M., Nielsen, S.H., and T.C.E. van Weering. 1998. Norwegian Sea overflow variability and NE Atlantic surface hydrography during the past 150,000 years. *Marine Geology*. 152; 75-99.
- Laskar, J., Joutel, F., and F. Boudin. 1993. Orbital, precessional, and insolation quantities for the Earth from -20 Myr to +10 Myr. *Astronomy and Astrophysics*. 270; 522-533.
- Lisiecki, L.E. and M.E. Raymo. 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$ records. *Paleoceanography*. 20 (1003); 1-17.
- Lisiecki, L.E., Raymo, M.E., and W.B. Curry. 2008. Atlantic overturning responses to Late Pleistocene climate forcings. *Nature*. 456 (6); 85-88.
- Lynch-Stieglitz, J., Adkins, J.F., Curry, W.B., Dokken, T., Hall, I.R., Herguera, J.C., Hirschi, J.J.M., Ivanova, E.V., Kissell, C., Marchal, O., Marchitto, T.M., McCave, I.N., McManus, J.F., Mulitza, S., Ninnemann, U., Peeters, F., Yu, E.-F., and R. Zahn. 2007. Atlantic Meridional Overturning Circulation During the Last Glacial Maximum. *Science*. 316; 66-69.
- Mackensen, A., H.-W. Hubberten, T. Bickert, G. Fischer, and D.K. Futterer. 1993. The $\delta^{13}\text{C}$ in benthic foraminiferal tests of *Fontbotia wuellerstorfi* (Schwager) relative to the $\delta^{13}\text{C}$ of dissolved inorganic carbon in Southern Ocean deep water: Implications for glacial ocean circulation models. *Paleoceanography*. 8(5); 587-610.
- Marchitto, T.M., Oppo, D.W., and W.B. Curry. 2002. Paired benthic foraminiferal Cd/Ca and Zn/Ca evidence for a greatly increased presence of Southern Ocean Water in the glacial North Atlantic. *Paleoceanography*. 17 (3); 1-16.
- McCave, I.N, Lonsdale P.F., Hollister C.D., and W.D. Gardner. 1980. Sediment Transport over the Hatton and Gardar Contourite Drifts. *Journal of Sedimentary Petrology*. 50 (4); 1049-1062.

- McCave, I.N. and B.E. Tucholke. 1986. Deep current-controlled sedimentation in the western North Atlantic. In Vogt, P.R. and B.E. Tucholke (Eds.), *The Geology of North America, Vol. M, The Western North Atlantic Region*. Geological Society of America, Boulder, CO; 451-468.
- Mauritzen, C. 1996. Production of dense overflow waters feeding the North Atlantic across the Greenland-Scotland Ridge. Part 1: Evidence for a revised circulation scheme. *Deep Sea Research Part 1: Oceanographic Research Papers*. 4 (6); 769-806.
- McIntyre, A. and B. Molino. 1996. Forcing of Atlantic Equatorial and Subpolar Millennial Cycles by Precession. *Science*. 274; 1867-1870.
- McIntyre, K., Ravelo, A.C., and M.L. Delaney. 1999. North Atlantic Intermediate Waters in the late Pliocene to early Pleistocene. *Paleoceanography*. 14; 324-335.
- McManus, J.F., Oppo, D.W., and J.L. Cullen. 1999. A 0.5-Million-Year Record of Millennial-Scale Climate Variability in the North Atlantic. *Science*. 283; 971-975.
- Millo, C., Sarnthein, M., Voelker, A., and H. Erlenkeuser. 2006. Variability of the Denmark Strait Overflow during the Last Glacial Maximum. *Boreas*. 1; 50-60.
- Moffa-Sanchez, P., I.R. Hall, D.J.R. Thornalley, S. Barker, and C. Stewart. 2015. Changes in the strength of the Nordic Seas Overflows over the past 3000 years. *Quaternary Science Reviews*. 123; 134-143.
- Olsen, A. and U. Ninnemann. 2010. Large $\delta^{13}\text{C}$ Gradients in the Preindustrial North Atlantic Revealed. *Science*. 330 (6004); 658-659.
- Oppo, D.W. and R.G. Fairbanks. 1987. Variability in the deep and intermediate water circulation of the Atlantic Ocean during the past 25,000 years; Northern Hemisphere modulation of the Southern Ocean. *Earth and Planetary Science Letters*. 86 (1); 1-15.
- Oppo, D.W. and S.J. Lehman. 1995. Suborbital timescale variability of North Atlantic Deep Water during the past 200,000 years. *Paleoceanography*. 10(5); 901-910.
- Oppo, D.W., Raymo, M.E., Lohmann, G.P. Mix, A.C., Wright, J.D., and W.L. Prell. 1995. A $\delta^{13}\text{C}$ record of Upper North Atlantic Deep Water during the past 2.6 million years. *Paleoceanography*. 10 (3); 373-394.
- Oppo, D.W., Horowitz, M., and S.J. Lehman. 1997. Marine Core Evidence for Reduced Deep Water Production During Termination II Followed By a Relatively Stable Substage 5e (Eemian). *Paleoceanography*. 12; 51-63.
- Oppo, D.W., Keigwin, L.D., McManus, J.F., and J.L. Cullen. 2001. Persistent suborbital climate variability in marine isotope stage 5 and Termination II. *Paleoceanography*. 16(3); 280.
- Ortiz, J.D., Mix, A., Harris, S., O'Connell, S., 1999. Diffuse spectral reflectance as a proxy for percent carbonate content in North Atlantic sediments. *Paleoceanography* 14,

- 820 171-186.
- 821
- 822 Paillard, D., Labeyrie, L., and P. Yiou. 1996. Macintosh program performs time-series
823 analysis. *Eos Trans. AGU.* 77; 379.
- 824
- 825 Prins, M.A., Troelstra, S.R., Kruk, R.W., van der Borg, K., de Jong, A.F.M., and G.J. Weltje.
826 2001. The Late Quaternary Sediment Record of the Reykjanes Ridge, North Atlantic.
827 *Radiocarbon.* 43 (2B); 939-947.
- 828
- 829 Prins., M.A., Bouwer, L.M., Beets, C.J., Troelstra, S.R., and G.J. Weltje. 2002. Ocean
830 circulation and iceberg discharge in the glacial North Atlantic: Inferences from
831 unmixing of sediment size distribution. *Geology.* 30 (6); 555-558.
- 832
- 833 R Development Core Team, 2010, R: A language and environment for statistical computing:
834 Vienna, Austria, R Foundation for Statistical Computing.
- 835
- 836 Rasmussen, T.L Thomson, E., van Weering, T.C.E., and L. Labeyrie. 1996a. Rapid changes
837 in surface and deep water conditions at the Faeroe Margin during the last 58,000
838 years. *Paleoceanography.* 11(6); 757-771.
- 839
- 840 Rasmussen, T.L Thomson, E., Labeyrie, L., and T.C.E. van Weering. 1996b. Circulation
841 changes in the Faeroe-Shetland Channel correlating with cold events during the last
842 glacial period (58-10 ka). *Geology.* 24; 937-940.
- 843
- 844 Rasmussen, T.L., E. Thomsen, A. Kuijpers, and S. Wastegard. 2013. Late warming and early
845 cooling of the sea surface in the Nordic seas during MIS 5e (Eemian Interglacial).
846 *Quaternary Science Reviews.* 22 (8-9); 809-821.
- 847
- 848 Rasmussen, T.L. and E. Thomsen. 2009. Ventilation changes in intermediate water on
849 millennial time scales in the SE Nordic seas, 65-11 kyr BP. *Geophysical Research*
850 *Letters.* 36l; 1-5.
- 851
- 852 Raymo, M.E., Ruddiman, W.F., Backman, J., Clement, B.M., and D.G. Martinson. 1989.
853 Late Pliocene variation in Northern Hemisphere ice sheets and North Atlantic deep
854 water circulation. *Paleoceanography.* 4; 413-446.
- 855
- 856 Raymo, M.E., Ruddiman, W.F., Backman, J., Clemet, B.M., and D.G. Martinson. 1990.
857 Evolution of Atlantic-Pacific $\delta^{13}\text{C}$ gradients over the last 2.5 my. *Earth and Planetary*
858 *Science Letters.* 97; 353-368.
- 859
- 860 Raymo, M.E.. 1997. The timing of major climate terminations. *Paleoceanography.* 12; 577-
861 585.
- 862
- 863 Raymo, M.E., Oppo, D.W., Flower, B.P., Flower, Hodell, D.A., McManus, J.F., Venz, K.A.,
864 Kleiven, K.F., and K. McIntyre. 2004. Stability of North Atlantic Water Masses in
865 Face of Pronounced Climate Variability During the Pleistocene. *Paleoceanography.*
866 19; 1-13.
- 867

- Revel, M., Cremer, M., Grousset, F.E., and L. Labeyrie. 1996. Grain-size and Sr-Nd isotopes as tracer of paleo-bottom current strength, Northeast Atlantic Ocean. *Marine Geology*. 131; 233-249.
- Rind, D., Dumeny, P., Russell, G.L., Sheth, S., Collins, D., Schmidt, G.A., and J. Teller. 2001. Effects of glacial meltwater in the GISS Coupled Atmosphere-Ocean Model: Part 1: North Atlantic Deep Water response. *Journal Geophysical Research Letters*. 106; 27335-27354.
- Roberts, N.L., A.M. Piotrowski, J.F. McManus, and L.D. Keigwin. 2010. Synchronous deglacial overturning and water mass source changes. *Science*. 327; 75-78.
- Ruddiman, W.F.. 1977. Late Quaternary deposition of ice-rafted sand in the subpolar North Atlantic (lat 40 to 65N). *Geological Society of America Bulletin*. 88; 1813-1827.
- Ruddiman, W.F., McIntyre, A., Raymo, M.E., 1987. Paleoenvironmental results from North Atlantic sites 607 and 609. Initial. Rep. DSDP 94, 855-878.
- Ruddiman, W.F., Raymo, M.E., Martinson, D.G., Clement, B.M., and J. Backman, 1989. Mid-Pleistocene evolution of Northern Hemisphere climate, *Paleoceanography*, 4; 353-412.
- Sarnthein, M., Winn, K., Jung, S.J.A., Duplessy, J.-C., Labeyrie, L., Erlenkeuser, H., and G. Ganssen. 1994. Changes in east Atlantic deepwater circulation over the last 30,000 years: Eight time slice reconstructions. *Paleoceanography*. 9 (2); 209-267.
- Sarnthein, M., Grootes, P.M., Kennett, J. P., and M.-J. Nadeau. 2007. ¹⁴C reservoir ages show deglacial changes in ocean currents and carbon cycle. In *Ocean Circulation: Mechanisms and Impacts – Past and Future Changes of Meridional Overturning* (eds. A. Schmittner, J.C.H. Chiang and S.R. Hemming), American Geophysical Union, Washington, D.C.. doi:10.1029/173GM13.
- Schmitz, W.J. and M.S. McCartney. 1993. On the North Atlantic Circulation. *Reviews of Geophysics*. 31 (1); 29-49.
- Shipboard Scientific Party. 1999. Site 1090. In: Gersonde, R., Hodell, D.A., Blum, P., et al. (Eds.), *Proc. ODP Init. Rep.* 177; 1-101.
- Stern, J.V. and L.E. Lisiecki. 2013. North Atlantic circulation and reservoir age changes over the past 41,000 years. *Geophysical Research Letters*. 40 (14); 3693-3697.
- Thornalley, D.J.R., McCave, I.N., and H. Elderfield. 2010. Freshwater input and abrupt deglacial climate change in the North Atlantic. *Paleoceanography*. 25 (1).
- Thornalley, D.J.R., Barker, S., Broecker, W.S., Elderfield, H., and I.N. McCave. 2011a. The Deglacial Evolution of North Atlantic Deep Convection. *Science*. 331 (6014); 202-205.
- Thornalley, D.J.R., McCave, I.N., and H. Elderfield. 2011b. Tephra in deglacial ocean sediments south of Iceland: stratigraphy, geochemistry and oceanic reservoir ages.

- 918 *Journal of Quaternary Science*. 26 (2); 190-198.
- 919
- 920 Turrell, W.R., Slessor, G., Adams, R.D, Payne, R., and P.A. Gillibrand. 1999. Decadal
- 921 variability in the composition of Faroe Shetland Channel bottom Water. *Deep-Sea*
- 922 *Research I*. 46; 1-25.
- 923
- 924 Venz, K.A. and D.A. Hodell. 2002. New evidence for changes in Plio-Pleistocene deep
- 925 circulation from Southern Ocean ODP Leg 177 Site 1090. *Paleogeography,*
- 926 *Paleoclimatology, Paleoecology*. 182; 197-220.
- 927
- 928 Wilson, D.J., K.C. Crocket, T. van de Flierdt, L. Robinson, and J.F. Adkins. 2014. Dynamic
- 929 intermediate ocean circulation in the North Atlantic during Heinrich Stadial 1: A
- 930 radiocarbon and neodymium isotope perspective. *Paleoceanography*. 29 (11); 1072-
- 931 1093.
- 932
- 933 Winsor, K., A.E. Carlson, G.P. Klinkhammer, J.S. Stoner, and R.G. Hatfield. 2012. Evolution
- 934 of the northeast Labrador Sea during the last interglaciation. *Geochemistry,*
- 935 *Geophysics, Geosystems*. 13 (11). 1-17.
- 936
- 937 Worthington, L.V.. 1976. On the North Atlantic Circulation. Johns Hopkins University Press;
- 938 Baltimore, MD.
- 939
- 940 Wright, J.D. and K.G. Miller. 1996. Control of North Atlantic Deep Water circulation by the
- 941 Greenland-Scotland Ridge. *Paleoceanography*. 11(2); 157-170.
- 942
- 943 Yu, J., H. Elderfield, and A.M. Piotrowski. 2008. Seawater carbonate ion- $\delta^{13}\text{C}$ systematics
- 944 and application to glacial-interglacial North Atlantic Ocean circulation. *Earth and*
- 945 *Planetary Science Letters*.
- 946
- 947
- 948
- 949
- 950
- 951
- 952
- 953
- 954
- 955
- 956

957 **FIGURE CAPTIONS:**

958 **Figure 1a:** Bathymetric map of the North Atlantic, including bottom water currents (blue),
 959 contourite drifts (uppercase lettering), and the northern components of NADW; ISOW,
 960 DSOW, and LSW (grey arrows). Core locations from this study and previously published
 961 studies are shown, including northern Gardar Drift sites (blue) ODP 984, EW9302, V29-202,
 962 and ODP983, and southern Gardar Drift sites (red) 11JPC (this study), 3GGC (this study),
 963 and Neap 18k. The location of the cross section in Figure 1b is designated by the white line.
 964 GEOSECS stations for the $\delta^{13}\text{C}$ data presented in Figures 1c, 1d, and 1e are designated by a
 965 cross for station 24 (red), 23 (blue), and 19 (green).

966

967 **Figure 1b:** Bathymetric cross section of the southern Norwegian Sea (NS) and the region
 968 south of the Iceland-Faeroes Ridge (IFR), as shown by the white line on Figure 1a. Bold lines
 969 show idealized directions of the major modern water masses, North Atlantic Current (NAC;
 970 orange) and Iceland-Scotland Overflow Water (ISOW; navy). The location of Bjorn and
 971 Gardar Drifts are shown, as are the projected locations of all cores used in this study (blue
 972 circles for northern Gardar Drift and red circles for southern Gardar Drift). Late Holocene *P.*
 973 *wuellerstorfi* $\delta^{13}\text{C}$ values from the selected cores are shown, where available. The vertical
 974 dashed lines represent the locations of recently measured profiles from GEOSECS stations 24
 975 (red; C), 23 (blue; D), and 19 (green; E; Kroopnick et al., 1974). Coretop benthic
 976 foraminiferal $\delta^{13}\text{C}$ data from cores used in this study are included where available.

977

978 **Figure 2:** Proxy records from core 11JPC plotted versus sediment depth. **(A)** $\delta^{18}\text{O}$ *G.*
 979 *bulloides* (light blue) and $\delta^{18}\text{O}$ *P. wuellerstorfi* (red), **(B)** $\delta^{13}\text{C}$ *G. bulloides* (blue) and $\delta^{13}\text{C}$ *P.*
 980 *wuellerstorfi* (red), **(C)** % CF (orange), **(D)** % CaCO_3 (green), and **(E)** the age model with
 981 age-depth tie points are shown. Marine Isotope Chrons 1 through 8 are also shown.

982

983 **Figure 3:** Proxy records for core 3GGC plotted versus sediment depth. **(A)** $\delta^{18}\text{O}$ *G. bulloides*
 984 (light blue) and $\delta^{18}\text{O}$ *P. wuellerstorfi* (pink), **(B)** $\delta^{13}\text{C}$ *G. bulloides* (blue) and $\delta^{13}\text{C}$ *P.*
 985 *wuellerstorfi* (red), **(C)** % CF (orange), **(D)** the age model, with age-depth tie points is shown.
 986 The Holocene, Younger Dryas (YD), Bolling/Allerod (BA), Last Glacial Maximum (LGM)
 987 and MIC 3 are also indicated.

988

989 **Figure 4:** A: The LR04 benthic foraminiferal stack (light blue; Lisiecki and Raymo, 2005) is
 990 shown for reference over our interval of interest. B: All benthic foraminiferal $\delta^{18}\text{O}$ data from
 991 northern portion of the Gardar Drift transect (red), the southern portion of the transect (blue),
 992 including our 2 new cores (11JPC and 3GGC). The very good correspondence between all
 993 records shows strong confidence in the age models of all cores. The 95% confidence interval
 994 of all data is shown in grey, which includes uncertainty propagated by a Monte Carlo
 995 ($n=1,000$) simulation.

996

997 **Figure 5** A: The LR04 benthic foraminiferal stack (light blue; Lisiecki and Raymo, 2005) is
 998 shown for reference over our interval of interest. Light grey vertical bars indicate glacial
 999 intervals of LGM, MIS4, and MIS6. B: Insolation at 65°N for the last 200 kyr (grey; Laskar
 1000 et al. 2004). C: All benthic foraminiferal $\delta^{13}\text{C}$ data from the northern portion of the Gardar
 1001 transect (red symbols; ODP Site 984, EW9302 JPC8, ODP Site 983, and V29-202) and the
 1002 southern portion of the transect (blue symbols; 11JPC, 3GGC, and NEAP18k) are shown
 1003 with age, including the 95% confidence interval of all Gardar Drift data (grey swath). D:
 1004 Average of all Gardar *P. wuellerstorfi* $\delta^{13}\text{C}$ values interpolated in 2-kyr intervals (black line)
 1005 plotted with ± 2 standard deviation (purple swath) is shown versus age from 0 to 200 ka.
 1006 The mid-latitude North Atlantic DSDP site 607 (black line) and South Atlantic ODP site

1007 1090 (green line) are shown for comparison. The mid-depth Atlantic composite record from
 1008 sites ODP925, ODP927, ODP928, and GeoB1214 from Lisiecki et al. (2008) is also shown
 1009 (dashed line) E: Standard deviation of all Gardar transect $\delta^{13}\text{C}$ values interpolated in 2-kyr
 1010 intervals (orange). F: The average of $\delta^{13}\text{C}$ *P. wuellerstorfi* records from the northern transect
 1011 sites (red line), compared with the southern transect sites (blue line), both shown with
 1012 associated confidence intervals (swaths). The mid-latitude North Atlantic (DSDP site 607;
 1013 black line) and South Atlantic (ODP site 1090; green line) are again shown for comparison.
 1014 G: The difference between $\delta^{13}\text{C}$ values of the smoothed northern and southern transects from
 1015 panel F (black line with pink swath indicating the confidence intervals). Note: the northern
 1016 quartet of sites and southern trio are given equal weighting.

1017

1018 **Figure 6:** Blackman-Tukey spectral analysis of the standard deviation benthic foraminiferal
 1019 $\delta^{13}\text{C}$ values for the Gardar Drift transect from 0 to 200 ka (orange line) and of the difference
 1020 between the smoothed Northern and Southern Gardar transects (purple line) determined using
 1021 AnalySeries. Power can be seen in the obliquity (41 kyr) and precession (23 kyr) bands.

Figure 1a

Figure 1b

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Table 1: Locations and water depths of cores used in this study.

Core	Lat	Lon	Water Depth (m)	Age Model Reference	Interval in compilation	Average age resolution (kyr per sample)
NORTHERN GARDAR DRIFT SITES:						
ODP site 984	61.43	-24.08	1660	Raymo et al., 2004	0-210 ka	1.30
EW9302 JPC8	61.00	-25.00	1915	Oppo et al., 2001; Oppo et al., 1997	57-135 ka	0.54
ODP site 983	60.40	-23.63	1995	Jansen et al., 1996; McIntyre et al., 1999; Raymo et al., 1998; Kleiven et al., 2003; Raymo et al., 2004;	0-210 ka	0.81
V29-202	61.00	-21.00	2658	Oppo & Lehman, 1995	0-160 ka	0.62
SOUTHERN GARDAR DRIFT SITES:						
KN166-14 11JPC	56.24	-27.65	2707	This Study	0-13 ka & 33-210 ka	0.92
KN166-14 3GGC	55.52	-26.53	3305	This Study	0-27 ka	0.40
Neap 18k	52.77	-30.35	3275	Chapman & Shackleton, 1999	55-125 ka	0.65
ATLANTIC SITES: (Not included in the compilation):						
DSDP site 607	41.00	-32.96	3427	Raymo et al., 2004		
ODP site 1090	42.91	-8.90	3702	Venz & Hodell, 2002		

Table 2: Age model for core KN166-14 11JPC was previously published for 0-33.81 ka (Elmore and Wright, 2011) and the interval from 33.81 to the bottom of the core is presented here for the first time. Tie points were determined by AMS ^{14}C (white boxes), magneto-stratigraphy (orange box) or chrono-stratigraphic comparison to a stacked benthic foraminiferal $\delta^{18}\text{O}$ record by Liseicki and Raymo (2005; blue boxes). AMS ^{14}C ages in red were not included in the age model (Elmore and Wright, 2011).

Depth (cm)	Age (ka)
0	0.67
45	2.87
105	6.05
127	7.06
143	8.38
145	6.99
147	8.58
149	8.86
153	8.88
163	9.65
183	11.45
198	11.99
213	13.34
220	13.62
227	34.07
250	33.71
282	31.91
333	33.81
430	40.00
626	62.70
779.7	74.50
1003	88.00
1260	103.00
1394	110.00
1518	129.30
1558	138.20
1636	174.50
1795	201.00
1929	226.00
2083	254.00
2290	295.00

Table 3: Age model for core KN166-14 3GGC was determined based on chrono-stratigraphic comparison to a stacked benthic foraminiferal $\delta^{18}\text{O}$ record by Liseicki and Raymo (2005).

Depth (cm)	Age (ka)
0	0
82	11.5
94	13.3
102	15
184	30